



1 **Supplementary material**



1


2 Figure S1: Global maps with annual average reference potential evaporation (PET; m
 3 day⁻¹). On top annual average CRU Penman-Monteith reference PET. In the right
 4 column reference PET obtained with the Penman-Monteith (PM), the standard
 5 Hargreaves (HGorig) and Blaney-Cridde (BCorig) and in the right column reference
 6 PET obtained with Priestley-Taylor (PT), Hargreaves with increased multiplication
 7 factor (HGrecal) and the re-calibrated Blaney-Cridde equation (BCrecal) are
 8 displayed.



1
2
3
4
5
6


Figure S2a: Global maps with seasonal average daily actual evapotranspiration (m day⁻¹). From left to right Penman-Monteith (PM), Hargreaves with increased multiplication factor (HGrecal) and re-calibrated Blaney-Criddle (BCrecal) and from top to bottom the DJF, MAM, JJA and SON seasons.

1



2


3 Figure S2b: Global maps with seasonal average daily actual evapotranspiration (m
4 day⁻¹). From left to right Priestley-Taylor (PT), the original Hargreaves equation
5 (HGorig) and the original Blaney-Criddle equation (BCorig) and from top to bottom
6 the DJF, MAM, JJA and SON seasons.



1

2 Figure S3a: Global maps with seasonal average daily cell specific runoff (m day⁻¹).
 3 From left to right Penman-Monteith (PM), Hargreaves with increased multiplication
 4 factor (HGrecal) and re-calibrated Blaney-Criddle (BCrecal) and from top to bottom
 5 the DJF, MAM, JJA and SON seasons.

1



2

3 Figure S3b: Global maps with seasonal average daily cell specific runoff (m day⁻¹).
4 From left to right Priestley-Taylor (PT), the original Hargreaves equation (HGorig)
5 and the original Blaney-Criddle equation (BCorig) and from top to bottom the DJF,
6 MAM, JJA and SON seasons.

7