



Technical note: Multiple wavelet coherence for untangling scale-specific and localized multivariate relationships in geosciences

Wei Hu^{1,3} and Bing Cheng Si^{2,3}

¹New Zealand Institute for Plant & Food Research Limited, Private Bag 4704, Christchurch 8140, New Zealand

²College of Hydraulic and Architectural Engineering, Northwest A&F University, Yangling 712100, China

³University of Saskatchewan, Department of Soil Science, Saskatoon, SK S7N 5A8, Canada

Correspondence to: Wei Hu (wei.hu@plantandfood.co.nz) and Bing Cheng Si (bing.si@usask.ca)

Received: 4 April 2016 – Published in Hydrol. Earth Syst. Sci. Discuss.: 27 April 2016

Revised: 8 July 2016 – Accepted: 13 July 2016 – Published: 8 August 2016

Abstract. The scale-specific and localized bivariate relationships in geosciences can be revealed using bivariate wavelet coherence. The objective of this study was to develop a multiple wavelet coherence method for examining scale-specific and localized multivariate relationships. Stationary and non-stationary artificial data sets, generated with the response variable as the summation of five predictor variables (cosine waves) with different scales, were used to test the new method. Comparisons were also conducted using existing multivariate methods, including multiple spectral coherence and multivariate empirical mode decomposition (MEMD). Results show that multiple spectral coherence is unable to identify localized multivariate relationships, and underestimates the scale-specific multivariate relationships for non-stationary processes. The MEMD method was able to separate all variables into components at the same set of scales, revealing scale-specific relationships when combined with multiple correlation coefficients, but has the same weakness as multiple spectral coherence. However, multiple wavelet coherences are able to identify scale-specific and localized multivariate relationships, as they are close to 1 at multiple scales and locations corresponding to those of predictor variables. Therefore, multiple wavelet coherence outperforms other common multivariate methods. Multiple wavelet coherence was applied to a real data set and revealed the optimal combination of factors for explaining temporal variation of free water evaporation at the Changwu site in China at multiple scale-location domains. Matlab codes for multiple wavelet coherence were developed and are provided in the Supplement.

1 Introduction

Geoscience data such as topography, climate, and ocean waves usually present cyclic patterns, with high-frequency (small-scale) processes being superimposed on low-frequency (large-scale) processes (Si, 2008). More often than not, geoscience data are transient, consisting of a variety of frequency regimes that may be localized in space or time (Torrence and Compo, 1998; Si and Zeleke, 2005; Graf et al., 2014). The transient characteristics exist widely in non-stationary processes, but also sometimes occur in stationary processes (Feldstein, 2000). The wavelet method is a common tool for detecting multi-scale and localized features of transient processes in geosciences. Bivariate wavelet coherence has been widely used for untangling scale-specific and localized relationships for transient processes in areas including geophysics (Lakshmi et al., 2004; Müller et al., 2008), hydrology (Labat et al., 2005; Das and Mohanty, 2008; Tang and Piechota, 2009; Carey et al., 2013; Graf et al., 2014), soil science (Si and Zeleke, 2005; Biswas and Si, 2011), meteorology (Torrence and Compo, 1998), and ecology (Polansky et al., 2010). This method, however, is limited to two variables. Processes in geosciences are usually complex and may be affected by more than two environmental factors. A method is needed for analysing multivariate (>2 variables) and localized relationships at multiple scales.

Several methods have been used for characterizing multivariate relationships. For example, multiple spectral coherence (MSC) has been used to explore the scale-specific relationships between soil-saturated hydraulic conductivity (K_s)

and multiple soil physical properties (Koopmans, 1974; Si, 2008), but requires a stationary data series, which is rare in geosciences. Multivariate empirical mode decomposition (MEMD), a data-driven method, decomposes each variable into different components (intrinsic mode functions (IMFs)) with each IMF corresponding to a common scale inherent in multiple variables (Rehman and Mandic, 2010). The MEMD method is meritorious due to its ability to deal with both transient and nonlinear systems. The combination of the squared multiple correlation coefficient and MEMD (MCC_{memd}) has been used to explore the multivariate control of soil water content and K_s at multiple scales (Hu and Si, 2013; She et al., 2013, 2015; Hu et al., 2014). However, the sum of variances from different components did not typically equal the total variance of the original series, which may produce misleading MCC_{memd} results. In addition, multivariate relationships in geosciences are most likely to change with time or space due to the transient nature of the processes involved. However, localized multivariate relationships are not available using any of the existing multivariate methods. Therefore, extending the wavelet coherence from two variables to multiple variables is required.

An attempt to extend wavelet coherence from two to three variables has been made by Mihanović et al. (2009). Their method was also applied later in the marine sciences (Ng and Chan, 2012a, b). Limitations arise when using the trivariate wavelet coherence: first, only two predictor variables are considered; second, the two predictor variables must be orthogonal. Otherwise, extremely high or low (spurious) coherence ($\gg 1$ or < 0) may be produced. This spuriousness is inconsistent with the definition of coherence, which may limit the application of this method in geosciences where environmental variables are usually cross correlated. Therefore, a robust method for calculating MWC, which produces coherence in the closed interval of $[0, 1]$, is needed.

The objective of this paper is to develop an MWC that applies to cases where there are multiple environmental variables, of which may be cross correlated. This method is first tested with artificial data sets to demonstrate its advantages over existing multivariate methods. The superiority of the new method over others can be assessed by determining whether the known major features of the artificial data are demonstrated by these methods. The new method is then applied to a temporal series of evaporation (E) from free water surface and meteorological factors at the Changwu site in Shaanxi, China.

2 Theory

Bivariate wavelet coherence can be understood as the traditional correlation coefficient localized in the scale-location domain (Grinsted et al., 2004). Just as correlation coefficients can be extended from two variables to multiple (> 2) variables, wavelet coherence between two variables may also be

extended to multiple variables. Similar to bivariate wavelet coherence, MWC is based on a series of auto- and cross-wavelet power spectra, at different scales and spatial (or temporal) locations, for the response variable and all predictor variables.

Following Koopman (1974), a matrix representation of the smoothed auto- and cross-wavelet power spectra for multiple predictor variables X ($X = \{X_1, X_2, \dots, X_q\}$) can be written as

$$\overset{\leftrightarrow}{W}^{X,X}(s, \tau) = \begin{bmatrix} \overset{\leftrightarrow}{W}^{X_1, X_1}(s, \tau) & \overset{\leftrightarrow}{W}^{X_1, X_2}(s, \tau) & \dots & \overset{\leftrightarrow}{W}^{X_1, X_q}(s, \tau) \\ \overset{\leftrightarrow}{W}^{X_2, X_1}(s, \tau) & \overset{\leftrightarrow}{W}^{X_2, X_2}(s, \tau) & \dots & \overset{\leftrightarrow}{W}^{X_2, X_q}(s, \tau) \\ \vdots & \vdots & \ddots & \vdots \\ \overset{\leftrightarrow}{W}^{X_q, X_1}(s, \tau) & \overset{\leftrightarrow}{W}^{X_q, X_2}(s, \tau) & \dots & \overset{\leftrightarrow}{W}^{X_q, X_q}(s, \tau) \end{bmatrix}, \quad (1)$$

where $\overset{\leftrightarrow}{W}^{X_i, X_j}(s, \tau)$ is the smoothed auto-wavelet power spectra (when $i = j$) or cross-wavelet power spectra (when $i \neq j$) at scale s and spatial (or temporal) location τ , respectively. For the detailed calculation of smoothed auto- and cross-wavelet power spectra, see Sect. S1 in the Supplement.

The matrix of smoothed cross-wavelet power spectra between response variable Y and multiple predictor variables X can be defined as

$$\overset{\leftrightarrow}{W}^{Y,X}(s, \tau) = \left[\overset{\leftrightarrow}{W}^{Y, X_1}(s, \tau) \overset{\leftrightarrow}{W}^{Y, X_2}(s, \tau) \dots \overset{\leftrightarrow}{W}^{Y, X_q}(s, \tau) \right], \quad (2)$$

where $\overset{\leftrightarrow}{W}^{Y, X_i}(s, \tau)$ is the smoothed cross-wavelet power spectra between Y and X_i at scale s and spatial (or temporal) location τ .

The smoothed wavelet power spectrum of response variable Y is $\overset{\leftrightarrow}{W}^{Y,Y}(s, \tau)$. Following Koopmans (1974), the MWC at scale s and location τ , $\rho_m^2(s, \tau)$, can be written as

$$\rho_m^2(s, \tau) = \frac{\overset{\leftrightarrow}{W}^{Y,X}(s, \tau) \overset{\leftrightarrow}{W}^{X,X}(s, \tau)^{-1} \overline{\overset{\leftrightarrow}{W}^{Y,X}(s, \tau)}}{\overset{\leftrightarrow}{W}^{Y,Y}(s, \tau)}. \quad (3)$$

where $\overline{\overset{\leftrightarrow}{W}^{Y,X}(s, \tau)}$ is a complex conjugate of $\overset{\leftrightarrow}{W}^{Y,X}(s, \tau)$.

When only one predictor variable (e.g. X_1) is included in X , Eq. (3) is the equation for bivariate wavelet coherence, $\rho_b^2(s, \tau)$, which can be expressed as (Torrence and Webster, 1999; Grinsted et al., 2004):

$$\rho_b^2(s, \tau) = \frac{\overset{\leftrightarrow}{W}^{Y, X_1}(s, \tau) \overline{\overset{\leftrightarrow}{W}^{Y, X_1}(s, \tau)}}{\overset{\leftrightarrow}{W}^{X_1, X_1}(s, \tau) \overset{\leftrightarrow}{W}^{Y, Y}(s, \tau)}, \quad (4)$$

Therefore, bivariate wavelet coherence is consistent with multiple wavelet coherence if only one predictor variable is included. In addition, the wavelet phase between a response variable (Y) and a predictor variable (X_1) is

$$\phi(s, \tau) = \tan^{-1} \left(\text{Im} \left(W^{Y, X_1}(s, \tau) \right) / \text{Re} \left(W^{Y, X_1}(s, \tau) \right) \right), \quad (5)$$




Figure 1. (a) Stationary and (b) non-stationary series of response variables (y for stationary and z for non-stationary case) encompassing five cosine waves (y_1 to y_5 for stationary and z_1 to z_5 for non-stationary case) with different dimensionless scales.

where Im and Re denote the imaginary and real part of $W^{Y,X_1}(s, \tau)$, respectively. Note that the phase information between a response variable Y and multiple predictor variables X cannot be obtained.

Multiple wavelet coherence at the 95% confidence level is calculated using the Monte Carlo method (Grinsted et al., 2004). Surrogate spatial series (i.e. red noise) of all variables are generated with a Monte Carlo simulation based on their first-order autocorrelation coefficient (AR1). The MWC at each scale and location is calculated using the simulated spatial series. This is repeated an adequate number of times (e.g. 1000) (Grinsted et al., 2004). At each scale, MWCs at all locations outside the cones of influence, from all simulations are ranked in ascending order. The value at the 95th percentile represents the 95% confidence level for the MWC at that scale. The Matlab codes and user manual document for calculating MWC and significance level are provided in the Supplement (Sect. S2–S4 in the Supplement).

3 Data and analysis

3.1 Artificial data for method test

The method is tested using a stationary and non-stationary artificial data set, generated following Yan and Gao (2007). The response variable (y for the stationary case and z for the non-stationary case) encompasses five cosine waves (y_1 to y_5 for the stationary case and z_1 to z_5 for the non-stationary case), with different dimensionless scales (Fig. 1). For the stationary case, $y_1 = \cos(2\pi x / 4)$, $y_2 = \cos(2\pi x / 8)$, $y_3 = \cos(2\pi x / 16)$, $y_4 = \cos(2\pi x / 32)$, and $y_5 = \cos(2\pi x / 64)$, where $x = 0, 1, 2, \dots, 255$. There is one regular cycle every 4, 8, 16, 32, and 64 locations, representing dimensionless scales of 4, 8, 16, 32, and 64 for y_1 , y_2 , y_3 , y_4 , and y_5 , respectively (Fig. 1a). The regular cycles make each predictor and response series stationary. For the non-stationary case, $z_1 = \cos(500\pi(x/1000)^{0.5})$, $z_2 = \cos(250\pi(x/1000)^{0.5})$, $z_3 = \cos(125\pi(x/1000)^{0.5})$, $z_4 = \cos(62.5\pi(x/1000)^{0.5})$, and $z_5 = \cos(31.25\pi(x/1000)^{0.5})$, where $x = 0, 1, 2, \dots, 255$. The equation containing the square root of the location term results in the gradual change in frequency (scale), with the greatest dimensionless scales of 4, 8, 16, 32, and 64 at the right hand side for z_1 , z_2 , z_3 , z_4 , and z_5 , respectively (Fig. 1b). The average scales for these predictor variables are 3, 5, 9, 17, and 32, respectively. The location-varying scales make each predictor and response variable non-stationary.

For both the stationary and non-stationary series, the variance of the response variable is 2.5. The predictor variables, each with a variance of 0.5, are orthogonal to each other, and contribute equally to the total variance of the response variable. The cosine-like artificial data sets mimic many time series such as seismic signals, turbulence, air temperature, precipitation, hydrologic fluxes, and the El Niño–Southern Oscillation. They also mimic geoscientific spatial series such as ocean waves, seafloor bathymetry, land surface topography, and soil water content along a hummocky landscape. Therefore, they are representative of a geoscience data series and are suitable for testing the new method.

Multiple wavelet coherence between the response variable y (or z) and two (y_2 and y_4 , or z_2 and z_4) or three (y_2 , y_3 , and y_4 , or z_2 , z_3 , and z_4) predictor variables were calculated. The advantage of the artificial data is that the known scale and localized features for all variables, and the known relationships between the response and each predictor variable, are exact. By definition, the coherence is 1 at scales corresponding to those of the included predictor variables, and 0 at other scales.

To demonstrate the advantages of MWC in dealing with abrupt changes (a type of transient and localized feature), the second half of the original series of y_2 (or z_2) or y_4 (or z_4) is replaced by 0, and MWC between the response variable and new set of predictor variables is calculated. We anticipate that

the coherence changes from 1 to 0 at the location where the new predictor variable becomes 0.

Predictor variables may not be as regular as those shown in Fig. 1, and may also be cross correlated to one another. For these reasons, zero-mean white noises with standard deviations of 0.3, 1, and 4 are added to the predictor variables of y_2 (or z_2) and y_4 (or z_4). The resulting noised series have correlation coefficients of 0.9, 0.5, and 0.1, respectively, with their original predictor variable. Therefore, we will refer to them as weakly, moderately, and highly noised series, respectively. Multiple wavelet coherences between the response variable and different predictor variables (original and noised series) are calculated to demonstrate the performance of MWC when noised or correlated predictor variables are involved. Only the non-stationary case will be demonstrated, because the performances of MWC for stationary and non-stationary cases are similar.

The MWC is compared to the MSC (Koopmans, 1974; Si, 2008) and MCC_{memd} (Hu and Si, 2013), which are widely used for spatial or temporal series analysis in different disciplines. The advantages of the new method over these two methods will be demonstrated mainly in terms of relationships between response and predictor variables at various scales of the response variable. The MSC is calculated based on the calculated auto- and cross-wavelet power spectra, using an equation similar to Eq. (3). The detailed introduction of this method can be found in Si (2008). For the calculation of MCC_{memd} , a set of response and predictor variables form a multivariate data series for MEMD. The MEMD is a data-driven method and has the ability to align common scales present within multivariate data. Please refer to Rehman and Mandic (2010) and Hu and Si (2013) for the MEMD analysis, and the website (<http://www.commsp.ee.ic.ac.uk/~mandic/research/emd.htm>) for the related Matlab codes. The original series of response and predictor variables can be decomposed by the MEMD, into different components (IMFs) with varying scales. For IMFs at the same scale, multiple stepwise regressions are conducted between response and predictor variables, and the multiple correlation coefficients for each scale-specific IMF are calculated.

3.2 Real data for application

Daily evaporation (E) from free water surfaces in an E601 evaporation pan (pan diameter of 61.8 cm), and other meteorological factors (i.e. relative humidity, mean temperature, sun hours, and wind speed) were collected from 1 January 1979 to 31 December 2013 at the Changwu site in Shaanxi, China. All these meteorological data are obtained from the China Meteorological Administration. The Changwu site is a transition area between semi-arid and sub-humid climates, where agricultural productivity is mainly limited by water. Monthly averages of all variables were used in this study, because we are mainly interested in seasonal and inter-annual variability.




Figure 2. Multiple wavelet coherence (a) between response variable y and predictor variables y_2 and y_4 ; (b) between response y and predictors y_2 , y_3 , and y_4 ; (c) between response z and predictors z_2 and z_4 ; and (d) between response z and predictors z_2 , z_3 , and z_4 . The artificial data series (y) encompasses five cosine waves (y_1 , y_2 , y_3 , y_4 , and y_5) with different scales for the stationary case, and the artificial data series (z) encompasses five cosine waves (z_1 , z_2 , z_3 , z_4 , and z_5) with different scales for the non-stationary case. The predictor variables, connected by a hyphen, are shown in the top right corner of each subplot. Thin solid lines demarcate the cones of influence, and thick solid lines show the 95 % confidence levels.

4 Results and discussion

4.1 MWC with orthogonal predictor variables

For the stationary data, there are two narrow, horizontal bands (red colour) representing an MWC value of around 1, at the respective scales of 8 and 32 for all locations (Fig. 2a). These two bands also correspond to the scales of 8 and 32, respectively, for the two predictor variables. When an additional predictor variable with the scale of 16 is introduced, a wide band appears from 6 to 40 (Fig. 2b), signifying that the MWC equals approximately 1 at all locations, at the scales of 8, 16, and 32. As anticipated, when all five predictor variables with scales ranging from 4 to 64 are included, coherence values of close to 1 are found in the whole scale-location domain (data not shown).

The application of MWC to the non-stationary data sets shows that the scales with significant MWC values gradually increase as distance increases (Fig. 2c and d). This increase in the scales is due to the non-stationarity of the variables (Fig. 1b). For example, when predictor variables of z_2 and z_4 are included, scales of the two bands corresponding to MWC around 1 increase from 4 to 8 and from 8 to 32, respectively (Fig. 2c). Furthermore, as expected, for only one




Figure 3. Multiple wavelet coherence (a) between y and y_2h_0 and y_4 ; (b) between y and y_2 and y_4h_0 ; (c) between z and z_2h_0 and z_4 ; and (d) between z and z_2 and z_4h_0 . The variables y_2h_0 (or z_2h_0) and y_4h_0 (or z_4h_0) refer to the new series of y_2 (or z_2) and y_4 (or z_4), in which the second half is replaced by 0.

predictor variable (stationary and non-stationary), MWC reduces to bivariate wavelet coherence; there is only one band of coherence around 1, which corresponds to the scale of that predictor variable (data not shown). Note that the significant MWC values for both stationary and non-stationary cases are not exactly 1 at all scales or locations, due to the smoothing effect along both scales and locations. However, the mean MWC values of the significant bands are very high (i.e. 0.94–1.00), and the MWC values at the centre of the significant band are 1, which corresponds to the exact scale of a predictor variable.

When the point values in the second half of the data series of a predictor variable are replaced by 0, the MWC values in that half of the data series are almost 0 at scales corresponding to that predictor variable (Fig. 3). For the stationary case, when the point values in the second half of the data series of predictor variable y_2 (or y_4) are replaced by 0, the MWC values are around 1 at the scale of 8 (or 32) in the first half of the transect, and 0 in the second half (Fig. 3a and b). Similar results are also found for the non-stationary case (Fig. 3c and d). This is expected because the constant series of 0 is not correlated to the response variables at any scale. Much like bivariate wavelet coherence, the MWC method is able to detect abrupt changes in the data series, and has the advantages of dealing with localized multivariate relationships.

4.2 MWC with noised and correlated predictor variables

When z_2 and a noised series derived from z_2 are included as predictor variables, there is only one band of coherence




Figure 4. Multiple wavelet coherence of an artificial data series (z) encompassing five cosine waves ($z_1, z_2, z_3, z_4,$ and z_5) with different scales and (a) z_2 and noised z_2 , (b) z_2 and noised z_4 , and (c) $z_2, z_4,$ and noised z_4 for the non-stationary case. The variables z_2wn (z_4wn), z_2mn (z_4mn), and z_2sn (z_4sn) indicate weakly, moderately, and strongly noised z_2 (z_4) series, respectively. Weakly, moderately, and strongly noised series are correlated with original series, with correlation coefficients of 0.9, 0.5, and 0.1, respectively.

close to 1 at scales corresponding to z_2 , irrespective of the correlation between z_2 and a noised series of z_2 (Fig. 4a). When z_2 and a noised series of z_4 are included as predictor variables, the coherence depends on the degree of the noise (Fig. 4b). For weakly noised series, there are two bands of coherence of around 1, corresponding to the scales of z_2 and z_4 , respectively. The percentage area of significant coherence (PASC) is 23 %, which equals that of when z_2 and z_4 are included. With the increasing magnitude of noise, the coherence and corresponding PASC at the scales corresponding to z_4 decrease. When z_2 and a strongly noised series of z_4 are considered, the band of coherence around 1, at scales corresponding to z_4 , disappears.

The inclusion of a third noised z_4 variable substantially increases the area with high coherence (in red) as compared to the case when only z_2 and z_4 are included (Fig. 4c). This indicates that MWC will increase as the number of predictor variables increases, with the highest coherence less or equal to 1, irrespective of the number of predictor variables. However, the area of significant coherence may not necessarily increase because of the simultaneously increased statistical significance threshold (Ng and Chan, 2012a). In fact, the PASC values for three predictor variables (19–20 %) are lower than those of the two predictor variables (23 %). This indicates that, in this case, two predictor variables are better than three

in terms of explaining the variations of the response variable. This occurs because the variance of the response variable that is explained by the noised variable is already accounted for by other variables. Therefore, only an additional variable that can independently explain a fair amount of variance could contribute significantly to explaining variations of a response variable (Fig. 4b). This may also explain why there is only one band of coherence around 1 at scales corresponding to z_2 , when z_2 and a noised series of z_2 are included (Fig. 4a). This information is helpful in choosing predictor variables for developing scale-specific predictions, especially when predictor variables are correlated.

4.3 Comparison with other multivariate methods

4.3.1 MSC

The MSC as a function of scale is shown in Fig. 5a. For the stationary case, when y_2 and y_4 are included as predictor variables, there are two plateaus centred at the scales of 8 and 28, representing a coherence of 1. As expected, when an additional predictor variable y_3 is added, the corresponding scale of 16 also shows coherence of 1. The MSC produces similar scale-specific relationships, as MWC does for a stationary data set, with exception given to the centred scale (i.e. 28) with a coherence of 1. Here, the scale with a unity MSC deviates from the expected value (i.e. 32) for predictor variable y_4 . For the non-stationary case, however, the MSC is much lower than 1 for the predictor variables of z_2 and z_4 ; an MSC of 1 is present only at the scale of 8 when an additional predictor variable z_3 is added. Obviously, the MSC underestimates the multivariate relationships, and is not suitable for non-stationary processes (Si, 2008) due to its inability to deal with localized features. The MSC at a specific scale provides the average of multivariate relationships, across all locations. Due to the change in scale of a predictor variable with location for the non-stationary case, the MSC deviates greatly from 1.

The MSC decreases at scales when the second half of the included predictor variable series are replaced by 0 for both the stationary and non-stationary series (Fig. 5b). For example, when the second half of the y_4 series in the stationary case is replaced by 0, the MSC at scales of around 32 decreases from 1 to 0.52. Although the MSC, throughout the second half of the series, can detect the decrease of coherence at the scales corresponding to the 0 values, the exact locations for the decrease cannot be identified. In fact, the coherence decreases only in the second half of the series, and does not change in the first half of the series. The location for the decrease can be easily identified by the MWC, but not by MSC. This further demonstrates the inability of the MSC to deal with localized features.




Figure 5. Multiple spectral coherence (MSC) of an artificial data series (y or z) encompassing five cosine waves (y_1 to y_5 ; or z_1 to z_5) with different scales and (a) two (y_2 and y_4 ; or z_2 and z_4) or three (y_2 , y_3 , and y_4 ; or z_2 , z_3 , and z_4) data series, and (b) two (y_2 and y_4 ; or z_2 and z_4) data series when the second half of one data series is replaced by 0. The variables y_2h_0 (or z_2h_0) and y_4h_0 (or z_4h_0) refer to the new series of y_2 (or z_2) and y_4 (or z_4) in which the second half is replaced by 0.

4.3.2 MCC_{memd}

Five intrinsic mode functions (IMFs) with non-negligible variance, are obtained for multivariate data series. While the obtained scales for the response variable y are in agreement with the true scales for the stationary case, the obtained scales (i.e. 3, 6, 11, 21, and 43) for the response variable z deviate slightly from the average scales for the non-stationary case. For the response variable, the contribution of IMFs to the total variance generally decreases (from 20 to 13% for stationary, and from 27 to 11% for non-stationary) from IMF1 to IMF5. This disagrees with the fact that each scale contributes equally (i.e. 20%) to the total variance. In addition, the sum of variances over all IMFs for each variable is less than 100% (ranging from 84 to 93%), indicating that MEMD cannot capture all the variances. For the detailed results of MEMD, see Sect. S5 in the Supplement.

The MCC_{memd} as a function of scale, is shown in Fig. 6a. For the stationary case, when predictor variables of y_2 and y_4 are included, the MCC_{memd} values are 0.98 and 0.93, respectively, at scales corresponding to those of y_2 and y_4 . When




Figure 6. Multiple correlation coefficient between multivariate empirical mode decomposition (MCC_{memd}) of an artificial series (y or z) and (a) two (y_2 and y_4 ; or z_2 and z_4) or three (y_2 , y_3 , and y_4 ; or z_2 , z_3 , and z_4) data series, and (b) two (y_2 and y_4 ; or z_2 and z_4) data series when the second half of one data series is replaced by 0. The variables y_2h_0 (or z_2h_0) and y_4h_0 (or z_4h_0) refer to the new series of y_2 (or z_2) and y_4 (or z_4) in which the second half is replaced by 0.

a predictor variable of y_3 is included, the MCC_{memd} values are 1.00, 1.00, and 0.96, respectively, at scales corresponding to those of y_2 , y_3 , and y_4 . For the non-stationary, two-predictor-variable case, the corresponding MCC_{memd} values are 0.80 and 0.85. For the non-stationary, three-predictor-variable case, the corresponding MCC_{memd} values are 0.95, 0.99, and 0.91, respectively. Therefore, the MCC_{memd} can be used to determine the scale-specific multivariate relationships. Similar to MSC, however, the MCC_{memd} underestimates the multivariate relationships, especially for the non-stationary case with less predictor variables. On the contrary, the MCC_{memd} also overestimates the multivariate relationships. For example, when considering only predictor variables corresponding to scales of 8, 16, and 32, the MCC_{memd} value for the stationary case is 0.47 at the scale of 64. This deviates much from the expected MCC_{memd} value of 0 (Fig. 6a). The possible underestimation and overestimation by the MCC_{memd} may come from the decomposition errors inherent in the MEMD algorithm (Rehman and Mandic, 2010).

Similar to MSC, the localized multivariate relationships cannot be obtained from MCC_{memd} . This can be better ex-

plained by the decrease of MCC_{memd} when half of the series of the predictor variables is replaced by 0 (Fig. 6b). Take the stationary case for example, the MCC_{memd} values at the scales corresponding to y_2 and y_4 decrease from 0.98 to 0.49, and from 0.93 to 0.62, respectively, when the second half of the y_2 and y_4 series is replaced by 0.

As explained above, the MWC has advantages in untangling localized multivariate relationships as compared to the common multivariate methods. It is important to reveal the multivariate relationships which vary with time or space, that are associated with different processes. For example, discharge usually occurs on knolls, while recharge usually occurs in neighbouring depressions (Gates et al., 2011). Therefore, the controlling factors of soil water storage may vary with the land element characteristics of a location. Local controls may be more important on knolls, while non-local controls may be more important in depressions (Grayson et al., 1997). In a temporal domain, vegetation transpiration contributes more to the evapotranspiration in the growing seasons, which may result in the changes of environmental factors explaining temporal variations of evapotranspiration in different seasons.

4.4 Application of the MWC

Each meteorological factor was significantly correlated to E , but the dominant factors explaining variations in E differed with scale. For example, the relative humidity was the dominating factor at small (2–8 months) and large (> 32 months) scales, while temperature was the dominating factor at the medium (8–32 months) scales. Overall, the relative humidity corresponded to the greatest mean MWC (0.62) and PASC value (40%) at multiple scale-location domains. For the detailed relationships between E and each factor, see Sect. S6 in the Supplement.

The MWC analysis shows that the combination of relative humidity and mean temperature produced the greatest mean MWC (0.82) and PASC (49%) among all two-factor cases. This suggested that relative humidity and mean temperature were the most appropriate factors for explaining variations in E at multiple scale-location domains (Fig. 7a). However, adding an additional factor such as sun hours, which was the best among all three-factor cases, increased the average coherence (0.91), but slightly decreased the PASC to 48% (Fig. 7b). This indicated that sun hours were not significantly different from red noise in explaining additional variation in E . Similar results were found when the wind speed was added. This occurs because most areas with significant coherence between E and sun hours or wind speed were a subset of areas with significant coherence between E and relative humidity or mean temperature (see Sect. S3 in the Supplement). Therefore, relative humidity and mean temperature were adequate for explaining the temporal variation of E at various scales at this site. This was consistent with Li et al. (2012), who indicated that relative humidity and mean




Figure 7. Multiple wavelet coherence between evaporation (E) from water surfaces and meteorological factors, such as (a) relative humidity and mean temperature and (b) relative humidity, mean temperature, and sun hours, at the Changwu site in Shaanxi, China.

temperature were the two main contributors to the temporal change of potential evapotranspiration on the Chinese Loess Plateau.

5 Conclusions

Multiple wavelet coherence was developed to determine scale-specific and localized multivariate relationships in geosciences. The new method was tested and compared with existing multivariate methods, using an artificial data set. The new method can be used to determine the proportion of the variance of a response variable that is explained by predictor variables, at a specific scale and location (spatially or temporally). As compared with bivariate wavelet coherence, more variation may be explained at multiple scale-location domains by the MWC. Including more variables is only beneficial if the variables are not strongly cross correlated, and can independently explain a fair amount of variability in a response variable. Therefore, the best combinations of variables that explain multivariate, spatial, or temporal variability at multiple scales can be determined. This is important for optimizing variables to develop scale-specific prediction.

The MSC and MCC_{memd} can determine multivariate relationships at multiple scales, but localized multivariate relationships are not available. Furthermore, both MSC and MCC_{memd} are likely to underestimate the degree of multivariate relationships for non-stationary processes. In addition, the performance of MCC_{memd} relies on the performance of MEMD, which needs further development. Application of the MWC into the real data set indicates that the combination of relative humidity and mean temperature are the optimal factors that can be used to explain temporal variations of E at the Changwu site in China.

Limitations of the new method also exist. Theoretically, any number of predictor variables can be included in the multiple wavelet analysis. However, the statistical significance threshold usually increases with the number of predictor variables (Grinsted et al., 2004; Ng and Chan, 2012a). In addition,

the inclusion of too many predictor variables may result in the statistical significance threshold at particular wavelet scales (e.g. the lowest and largest scales) to approach unity. This would restrict the availability of statistical information. Furthermore, similar to bivariate wavelet analysis, the new method also suffers from the multiple-testing problem (Maraun and Kurths, 2004; Maraun et al., 2007; Schaeffli et al., 2007; Schulte et al., 2015; Schulte, 2016). Therefore, a more robust statistical significance testing method may be beneficial to the new method.

In summary, multiple wavelet coherence has advantages over existing multivariate methods, and provides an effective vehicle for untangling complex spatial or temporal variability for multiple controlling factors at multiple scales and locations. It may also be used as a data-driven tool for modelling and predicting various processes in the area of geosciences, such as precipitation, drought, soil water dynamics, stream flow, and atmospheric circulation.

6 Data availability

The meteorological data sets can be obtained from the China Meteorological Administration.

The Supplement related to this article is available online at doi:10.5194/hess-20-3183-2016-supplement.

Acknowledgements. The Matlab codes for calculating multiple wavelet coherence are developed based on the codes provided by A. Grinsted (<http://www.glaciology.net/wavelet-coherence>) and, together with the user manual, are available in the Supplement (Sect. S2–S4). The project was funded by the National Natural Science Foundation of China (41371233), the Natural Sciences and Engineering Research Council of Canada (NSERC), Agriculture Development Fund of Saskatchewan, and the New Zealand Institute for Plant & Food Research under the *Land Use Change and Intensification* programme. We thank the two anonymous reviewers for their constructive comments.

Edited by: B. Schaeffli

Reviewed by: two anonymous referees

References

- Biswas, A. and Si, B. C.: Identifying scale specific controls of soil water storage in a hummocky landscape using wavelet coherence, *Geoderma*, 165, 50–59, doi:10.1016/j.geoderma.2011.07.002, 2011.
- Carey, S. K., Tetzlaff, D., Buttle, J., Laudon, H., McDonnell, J., McGuire, K., Seibert, J., Soulsby, C., and Shanley, J.: Use of color maps and wavelet coherence to discern seasonal

- and interannual climate influences on streamflow variability in northern catchments, *Water Resour. Res.*, 49, 6194–6207, doi:10.1002/wrcr.20469, 2013.
- Das, N. N. and Mohanty, B. P.: Temporal dynamics of PSR-based soil moisture across spatial scales in an agricultural landscape during SMEX02: A wavelet approach, *Remote Sens. Environ.*, 112, 522–534, doi:10.1016/j.rse.2007.05.007, 2008.
- Feldstein, S. B.: The timescale, power spectra, and climate noise properties of teleconnection patterns, *J. Clim.*, 13, 4430–4440, doi:10.1175/1520-0442(2000)013<4430:TTPSAC>2.0.CO;2, 2000.
- Gates, J. B., Scanlon, B. R., Mu, X. M., and Zhang, L.: Impacts of soil conservation on groundwater recharge in the semi-arid Loess Plateau, China, *Hydrogeol. J.*, 19, 865–875, 2011.
- Graf, A., Bogen, H. R., Drüe, C., Hardelauf, H., Pütz, T., Heine-mann, G., and Vereecken, H.: Spatiotemporal relations between water budget components and soil water content in a forested tributary catchment, *Water Resour. Res.*, 50, 4837–4857, doi:10.1002/2013WR014516, 2014.
- Grayson, R. B., Western, A. W., Chiew, F. H. S., and Blöschl, G.: Preferred states in spatial soil moisture patterns: local and nonlocal controls, *Water Resour. Res.*, 33, 2897–2908, doi:10.1029/97WR02174, 1997.
- Grinsted, A., Moore, J. C., and Jevrejeva, S.: Application of the cross wavelet transform and wavelet coherence to geophysical time series, *Nonlin. Processes Geophys.*, 11, 561–566, doi:10.5194/npg-11-561-2004, 2004.
- Hu, W. and Si, B. C.: Soil water prediction based on its scale-specific control using multivariate empirical mode decomposition, *Geoderma*, 193–194, 180–188, doi:10.1016/j.geoderma.2012.10.021, 2013.
- Hu, W., Biswas, A., and Si, B. C.: Application of multivariate empirical mode decomposition for revealing scale- and season-specific time stability of soil water storage, *Catena*, 113, 377–385, doi:10.1016/j.catena.2013.08.024, 2014.
- Koopmans, L. H.: *The spectral analysis of time series*, Academic Press, New York, 119–164, 1974.
- Labat, D.: Recent advances in wavelet analyses: Part I. A review of concepts, *J. Hydrol.*, 314, 275–288, doi:10.1016/j.jhydrol.2005.04.003, 2005.
- Lakshmi, V., Piechota, T., Narayan, U., and Tang, C. L.: Soil moisture as an indicator of weather extremes, *Geophys. Res. Lett.*, 31, L11401, doi:10.1029/2004GL019930, 2004.
- Li, Z., Zheng, F. L., and Liu, W. Z.: Spatiotemporal characteristics of reference evapotranspiration during 1961–2009 and its projected changes during 2011–2099 on the Loess Plateau of China, *Agric. For. Meteorol.*, 154–155, 147–155, doi:10.1016/j.agrformet.2011.10.019, 2012.
- Maraun, D. and Kurths, J.: Cross wavelet analysis: significance testing and pitfalls, *Nonlin. Processes Geophys.*, 11, 505–514, doi:10.5194/npg-11-505-2004, 2004.
- Maraun, D., Kurths, J., and Holschneider, M.: Nonstationary Gaussian processes in wavelet domain: synthesis, estimation, and significance testing, *Phys. Rev. E*, 75, 016707, doi:10.1103/PhysRevE.75.016707, 2007.
- Mihanović, H., Qrlić, M., and Pasrić, Z.: Diurnal thermocline oscillations driven by tidal flow around an island in the Middle Adriatic, *J. Marine Syst.*, 78, S157–S168, doi:10.1016/j.jmarsys.2009.01.021, 2009.
- Müller, W. A., Frankignoul, C., and Chouaib, N.: Observed decadal tropical Pacific–North Atlantic teleconnections, *Geophys. Res. Lett.*, 35, L24810, doi:10.1029/2008GL035901, 2008.
- Ng, E. K. W. and Chan, J. C. L.: Geophysical applications of partial wavelet coherence and multiple wavelet coherence, *J. Atmos. Ocean. Tech.*, 29, 1845–1853, doi:10.1175/JTECH-D-12-00056.1, 2012a.
- Ng, E. K. W. and Chan, J. C. L.: Interannual variations of tropical cyclone activity over the north Indian Ocean, *Int. J. Climatol.*, 32, 819–830, doi:10.1002/joc.2304, 2012b.
- Polansky, L., Wittemyer, G., Cross, P. C., Tambling, C. J., and Getz, W. M.: From moonlight to movement and synchronized randomness: Fourier and wavelet analyses of animal location time series data, *Ecology*, 91, 1506–1518, doi:10.1890/08-2159.1, 2010.
- Rehman, N. and Mandic, D. P.: Multivariate empirical mode decomposition, *Proc. R. Soc. A*, 466, 1291–1302, doi:10.1098/rspa.2009.0502, 2010.
- Schaeffli, B., Maraun, D., and Holschneider, M.: What drives high flow events in the Swiss Alps? Recent developments in wavelet spectral analysis and their application to hydrology, *Adv. Water Resour.*, 30, 2511–2525, doi:10.1016/j.advwatres.2007.06.004, 2007.
- Schulte, J. A.: Cumulative areawise testing in wavelet analysis and its application to geophysical time series, *Nonlin. Processes Geophys.*, 23, 45–57, doi:10.5194/npg-23-45-2016, 2016.
- Schulte, J. A., Duffy, C., and Najjar, R. G.: Geometric and topological approaches to significance testing in wavelet analysis, *Nonlin. Processes Geophys.*, 22, 139–156, doi:10.5194/npg-22-139-2015, 2015.
- She, D. L., Liu, D. D., Peng, S. Z., and Shao, M. A.: Multiscale influences of soil properties on soil water content distribution in a watershed on the Chinese Loess Plateau, *Soil Sci.*, 178, 530–539, doi:10.1097/SS.0000000000000021, 2013.
- She, D. L., Zheng, J. X., Shao, M. A., Timm, L. C., and Xia, Y. Q.: Multivariate empirical mode decomposition derived multi-scale spatial relationships between saturated hydraulic conductivity and basic soil properties, *Clean-Soil Air Water*, 43, 910–918, doi:10.1002/clen.201400143, 2015.
- Si, B. C.: Spatial scaling analyses of soil physical properties: A review of spectral and wavelet methods, *Vadose Zone J.*, 7, 547–562, doi:10.2136/vzj2007.0040, 2008.
- Si, B. C. and Zeleke, T. B.: Wavelet coherence analysis to relate saturated hydraulic properties to soil physical properties, *Water Resour. Res.*, 41, W11424, doi:10.1029/2005WR004118, 2005.
- Tang, C. L. and Piechota, T. C.: Spatial and temporal soil moisture and drought variability in the Upper Colorado River Basin, *J. Hydrol.*, 379, 122–135, doi:10.1016/j.jhydrol.2009.09.052, 2009.
- Torrence, C. and Compo, G. P.: A practical guide to wavelet analysis, *Bull. Am. Meteorol. Soc.*, 79, 61–78, doi:10.1175/1520-0477(1998)079<0061:apgtwa>2.0.CO;2, 1998.
- Torrence, C. and Webster, P. J.: Interdecadal changes in the ENSO-monsoon system, *J. Clim.*, 12, 2679–2690, doi:10.1175/1520-0442(1999)012<2679:ICITEM>2.0.CO;2, 1999.
- Yan, R. and Gao, R. X.: A tour of the Hilbert–Huang transform: an empirical tool for signal analysis, *IEEE Instrum. Meas. Mag.*, 10, 40–45, doi:10.1109/MIM.2007.4343566, 2007.