



A systematic assessment of drought termination in the United Kingdom

Simon Parry^{1,2}, Robert L. Wilby², Christel Prudhomme^{1,2}, and Paul J. Wood²

¹Centre for Ecology & Hydrology, Wallingford, Oxfordshire, OX10 8BB, UK

²Department of Geography, Loughborough University, Loughborough, Leicestershire, LE11 3TU, UK

Correspondence to: Simon Parry (spar@ceh.ac.uk)

Received: 20 November 2015 – Published in Hydrol. Earth Syst. Sci. Discuss.: 15 January 2016

Revised: 19 August 2016 – Accepted: 5 September 2016 – Published: 20 October 2016

Abstract. Drought termination can be associated with dramatic transitions from drought to flooding. Greater attention may be given to these newsworthy and memorable events, but drought terminations that proceed gradually also pose challenges for water resource managers. This paper defines drought termination as a distinctive phase of the event. Using observed river flow records for 52 UK catchments, a more systematic and objective approach for detecting drought terminations is demonstrated. The parameters of the approach are informed by a sensitivity analysis that ensures a focus on terminations of multi-season to multi-year droughts. The resulting inventory of 467 drought terminations provides an unprecedented historical perspective on this phenomenon in the UK. Nationally and regionally coherent drought termination events are identifiable, although their characteristics vary both between and within major episodes. Contrasting drought termination events in 1995–1998 and 2009–2012 are examined in greater depth. The data are also used to assess potential linkages between metrics of drought termination and catchment properties. The duration of drought termination is moderately negatively correlated with elevation ($r_s = -0.47$) and catchment average rainfall ($r_s = -0.42$), suggesting that wetter catchments in upland areas of the UK tend to experience shorter drought terminations. More urbanized catchments tend to have gradual drought terminations (contrary to expectations of flashy hydrological response in such areas), although this may also reflect the type of catchments typical of lowland England. Significant correlations are found between the duration of the drought development phase and both the duration ($r_s = -0.29$) and rate ($r_s = 0.28$) of drought termination. This suggests that prolonged drought development phases tend to be followed by shorter and more

abrupt drought terminations. The inventory helps to place individual events within a long-term context. The drought termination phase in 2009–2012 was, at the time, regarded as exceptional in terms of magnitude and spatial footprint, but the Thames river flow record identifies several comparable events before 1930. The chronology could, in due course, provide a basis for exploring the complex drivers, long-term variability, and impacts of drought termination events.

1 Introduction

Drought termination, generally defined as the end point of a drought, has been neglected in research literature relative to drought onset. Studies which address this phenomenon have focused on extreme transitions at the end of a drought (e.g. Yang et al., 2012; Ning et al., 2013), but there has been a lack of attention devoted to assessing the full range of drought termination types and characteristics. Whilst abrupt drought terminations may result in more destructive and newsworthy impacts (e.g. Webster et al., 2011; Lavers and Villarini, 2013; Parry et al., 2013), gradual drought terminations are problematic for water resource managers who must reconcile public relations with continued water restrictions during wet weather.

Some studies systematically identify and characterize droughts themselves (e.g. Hisdal et al., 2001; Pfister et al., 2006; Marsh et al., 2007; Fleig et al., 2011; Li et al., 2013), but these have generally not considered the drought termination phase. A limited historical perspective can be gained from studies of drought termination on an event basis, including those based on hydrometeorological (e.g. Kienzie,




Figure 1. Locations of the 52 study catchments, colour-coded by region. The regions are abbreviated in Figs. 4, 5, and 6 as follows: Western Scotland: WS; Eastern Scotland: ES; Northern Ireland: NI; North-west England: NWE; North-east England: NEE; North & Central Wales: NCW; Midlands: MID; South-west UK: SWUK; Anglian: ANG; Southern England: SE. Inset: the constituent countries of the UK.

2006; Marengo et al., 2008), remotely sensed (e.g. Wang et al., 2013; Chew and Small, 2014), or experimental catchment data (e.g. Miller et al., 1997; Lange and Hansler, 2012). Even considering several events (e.g. Eltahir and Yeh, 1999; Shukla et al., 2011) is too limited a sample to generalize or move beyond qualitative descriptions (e.g. Parry et al., 2013). A systematic assessment would enable a more robust analysis of the spatial and temporal variability of drought termination. Moreover, the importance of the end of a drought has already been recognized as a criterion in a hydrological drought typology and a basis for differentiating drought types (Van Loon and Van Lanen, 2012; Van Loon et al., 2015).

Studies that systematically identify the end of droughts in the historical record (e.g. Mo, 2011; Kam et al., 2013; Maxwell et al., 2013; Patterson et al., 2013) have typically considered drought termination to be instantaneous. There are two notable exceptions: Bonsal et al. (2011) sub-

divided drought into six stages, one of which is the concept of drought termination as a phase considered herein, and Nkemdirim and Weber (1999) expressed the concept of a rate of drought termination using Palmer Drought Severity Index units over time.

Preliminary steps have been taken to identify and characterize the spatial signature of a single drought termination for 15 catchments in the UK (Parry et al., 2016), and to apply the same assessment technique in a temporal analysis of drought terminations in a single catchment for the period 1883–2013 (Parry et al., 2015). The approach adopted in these studies differs from others (e.g. Kam et al., 2013; Patterson et al., 2013) by considering drought termination to be a period of a drought event with its own start, end and duration between these points.

By combining these spatial (Parry et al., 2016) and temporal approaches (Parry et al., 2015), the aim of this study is to derive chronologies of drought termination for 52 UK catchments. These data are subsequently used to assess the historical variability of drought termination and to explore the link between drought termination metrics and catchment properties. A sensitivity analysis of the drought termination metrics to methodological parameters is included; the selection of parameters that results from this analysis is also informed by the focus of this study on the termination of multi-season to multi-year droughts. It is anticipated that a better understanding of the physical processes driving drought termination will lead to improved water resources management and forecasting during these problematic episodes in the future.

2 Data

Catchments were selected on the basis of their area and record length, favouring larger catchments with longer records in order to maximize the spatial and temporal coverage of the chronologies. This selection was supplemented by additional catchments to improve representation of the diversity of hydrogeological conditions in the UK. The resulting 52 catchments (Fig. 1; Table A1) account for more than 40 % of the gauged area of the UK whilst capturing some of the longest river flow records. Nearly half (21 of 52) of the catchments are classified as near-natural, and these are predominantly located in northern and western areas of the UK. To the south and east and for the larger catchments, flows may be affected by anthropogenic influences (such as abstractions and return flows) which can mask changes associated with drought termination (Ning et al., 2013). A naturalized river flow series is used for the Thames; no other naturalized series are available for the study catchments. River flow data were obtained from the UK National River Flow Archive (NRFA). Start dates range between January 1883 and June 1982, but all series extend to September 2013. Time series of monthly mean river flows were derived for each catchment for every month in which at least 90 % of the daily




Figure 2. A conceptualization of drought termination definition and metrics. The three parameters are as follows: D is the number of months of below-average flows required for the drought development phase to begin; R is the number of months of intermittent above-average flows permitted within D ; and T is the number of consecutive months of above-average flows required for the end of the drought termination phase. t_{sd} is the time of the start of drought development, t_{ed} is the time of the end of drought development, t_{st} is the time of the start of drought termination, and t_{et} is the time of the end of drought termination. The grey horizontal line represents an anomaly of zero, below which flows are below-average and above which flows are above-average.

data were available. Metadata on catchment area, median elevation, Standard-period Average Annual Rainfall for 1961–1990 (hereafter SAAR6190; Spackman, 1993), Base Flow Index (hereafter BFI; Gustard et al., 1992), and urban extent (Marsh and Hannaford, 2008) were also obtained for each catchment from the NRFA (Table A1).

3 Methodology

3.1 Defining drought termination

Drought termination is defined here as a phase of a drought, rather than an instantaneous point in time. The threshold level method (Zelenhasić and Salvai, 1987) has been applied on a monthly time step, and drought events are subdivided at the point of the maximum negative flow anomaly (Bravar and Kavvas, 1991) into two phases: drought development and drought termination (Fig. 2). Drought termination is characterized by its duration (e.g. Bonsal et al., 2011), rate of change (e.g. Correia et al., 1987; Nkemdirim and Weber, 1999), and seasonality (e.g. Mo, 2011).

For each catchment, monthly mean flow data were converted into a percentage anomaly of the monthly long-term average (LTA), calculated from a 1971–2000 reference period (Eq. 1).

$$Z_{anom,t} = 100((Z_{obs,t}/Z_{LTA,m}) - 1), \quad (1)$$

where t is the time step index, m is the month of the time step, $Z_{anom,t}$ is the percentage anomaly at t , $Z_{obs,t}$ is the observed value at t , and $Z_{LTA,m}$ is the LTA at m . Where river flow records commence after 1971 (13 of the 52 catchments; Table A1), the monthly LTA is an average of all available monthly mean flows within the 1971–2000 timeframe. Of these 13 catchments, only five sets of monthly LTAs are derived from less than 24 years of available data, and all catchments have at least 19 years in the 1971–2000 period.

The start of a drought development phase (t_{sd} where “s” is start and “d” is development; Fig. 2) is the first month of D consecutive months (pre-defined by the user) for which $Z_{anom,t}$ is negative. R months within the D -month duration are permitted to be above average, to account for minor wet interludes during the development of the drought. Once a drought has been initiated, the end of the drought termination phase (t_{et} where “e” is end and “t” is termination; Fig. 2) is the last month of T consecutive months for which $Z_{anom,t}$ is greater than $Z_{LTA,m}$. The termination magnitude (TM; Fig. 2) is $Z_{anom,t}$ at t_{et} .

The end of the drought development phase (t_{ed} ; Fig. 2) is the month with the largest negative $Z_{anom,t}$ value (defining the drought magnitude, DM; Fig. 2) between t_{sd} and t_{et} . The start of the drought termination phase (t_{st} ; Fig. 2) is the next month after t_{ed} .

The conceptual diagram in Fig. 2 illustrates the two phases of drought and some of the associated drought termination




Figure 3. Demonstrations of the sensitivity of drought termination metrics to parameter selection for the Scottish Dee and Itchen catchments. D , R , and T are the three parameters of the methodology: $D7$ – $D36$ are 7- to 36-month durations over which Z_{anom} is negative; $R1$ – $R3$ are the number of months (1, 2, or 3) within the D -month duration for which Z_{anom} is permitted to be positive; $T2$ – $T6$ are the number of consecutive months (2–6) for which Z_{anom} is positive. The metrics are as follows: “Ev dec $^{-1}$ ”: number of events per decade; TDD: total drought duration (drought development duration and drought termination duration taken together); DTR: drought termination rate. The bold box on each response surface shows the combination of parameters used to derive the drought termination chronologies in this study.

metrics. The drought termination duration (DTD; Fig. 2) is the number of months between t_{st} and t_{et} . The drought termination rate (DTR; Fig. 2) is the difference between the drought magnitude and the termination magnitude, divided by the drought termination duration. The drought termination seasonality is a code relating to the seasons through which drought termination occurs. For example, if the start of drought termination is in autumn and the end of drought termination is in the next winter, the drought termination seasonality would be “Aut–Win”. Because seasonality is assessed on the entire drought termination period rather than its beginning or end, when drought termination durations span four or more seasons they are considered not to have a seasonality.

3.2 Parameter selection

At the outset, expert judgement was used to select parameters which identified well-known hydrological droughts in the historical record. A drought chronology for the UK (Marsh et al., 2007) identified an average of two events per decade over the last 50 years. Experimentation with different parameter sets suggested that a moderately high value for D is required to ensure a focus on multi-season and multi-year droughts. The value of R must balance between identifying unrealistically large numbers of events or none at all. The hydrological variability of many catchments in the UK requires the value of T to be greater than one, to account for wet interludes during droughts. Combining these findings with prior expert knowledge on drought occurrence in the UK, the following parameters were identified as appropriate for the aims of this study: $D = 10$; $R = 1$; $T = 2$.

Once the parameters had been selected, response surfaces (e.g. Fig. 3) were used to provide quantitative support for this decision. At first glance across a range of catchment sizes, characteristics, and hydroclimatic settings, the parameters above generally satisfy the approximate events per decade criteria outlined above. Two contrasting catchments were selected to illustrate typical patterns of sensitivity in the response surfaces. The Scottish Dee (Eastern Scotland; Fig. 3, left) is a relatively wet upland catchment with impermeable geology and a flashy hydrological response, whilst the Itchen (Southern England; Fig. 3, right) is a relatively dry lowland catchment with permeable geology and a buffered hydrological response. The identified combination of parameters ($D = 10$; $R = 1$; $T = 2$) is indicated by bold boxes on the response surfaces in Fig. 3.

The response surfaces illustrate how the number of drought events identified varies with parameter selection. Fewer events were identified with increasing D (moving from left to right in Fig. 3, top left and top right) due to stricter criteria for drought initiation. Conversely, increasing R (for a given D and T , moving from bottom to top in Fig. 3, top left and top right) detected more events because this relaxed the initiation criteria (ratio between D and R) to allow more intermittent months above the average flow threshold. As T increased (for a given D and R , moving from bottom to top in Fig. 3, top left and top right), the number of identified events decreased as the threshold for completion of drought termination became more stringent. These patterns were consistent across a range of catchment sizes, characteristics and hydroclimatic settings.

Although the number of identified events was the primary verification provided by the response surfaces, variations in the average characteristics of the resulting events were also explored. For total drought duration (TDD), increasing T for the Scottish Dee (moving from bottom to top in Fig. 3, middle left) caused identified droughts to lengthen considerably and resulted in merging of previously distinct events into unrealistically long periods (e.g. exceeding 120 months, or




Figure 4. A chronology of drought termination for all 52 study catchments. Red bars indicate drought development, blue bars indicate drought termination, white bars indicate no drought development or drought termination, and grey bars signify periods before gauged river flow records began. On the x axis, a decade (e.g. 1990–2000) is comprised of 120 monthly time steps and there are 1569 monthly time steps along the entire x axis (January 1883 to September 2013, inclusive).

10 years). The Itchen did not exhibit this behaviour (Fig. 3, middle right), suggesting that individual drought events were typically separated by long spells (greater than 6 months) with above threshold flows such that merging was less likely. This was consistent with the lower variability of river flows in groundwater influenced catchments like the Itchen. Similar contrasts between the two catchments were also apparent for drought termination rate (DTR; Fig. 3, bottom left and bottom right), in part because duration is a component of the DTR calculation. Higher values of T caused more merging of events in responsive catchments such as the Scottish Dee, increasing TDD (and DTD) and thereby reducing DTR. Although not directly comparable due to the different nature of the indicators used, this finding is consistent with a previous study of two catchments with contrasting river flow regimes in which less stringent criteria for drought identification increased the duration of droughts in the more responsive catchment to a far greater degree (Tallaksen et al., 1997). This suggests more stringent criteria are required for more responsive catchments.

In general, drought termination metrics showed greater sensitivity to parameter values in more responsive catchments (less responsive catchments were insensitive). Severe initiation criteria (high D and low R) and larger values of T are not appropriate for responsive catchments because these combinations are physically implausible, resulting in

the merging of events into unrealistic durations with corresponding effects on derived drought termination metrics.

These key findings of the sensitivity analysis verified the initial decision on parameter selection. Values of $D = 10$, $R = 1$, and $T = 2$ do not over- or under-represent drought occurrence for catchments of different size, geology, or average rainfall, whilst primarily identifying severe multi-year and multi-season events that form the focus of this study. For these reasons the same parameter values were applied to all 52 catchments in this study, and enabled a comparison of drought termination characteristics across catchments without the influence of variations in parameter selection.

3.3 Correlation analysis

Potential relationships between drought termination characteristics and catchment properties were explored through a correlation analysis. Since the majority of drought termination characteristics are not normally distributed, and to limit the influence of outliers, the Spearman rank correlation test (Spearman, 1944) was applied to the inventory of drought development and drought termination characteristics and catchment metadata. Correlation analysis was performed using all 52 catchments, as well as on a subset of catchments with at least 10 drought termination events. By omitting catchments with only a few identified events, a subset of catchments is retained for which catchment average drought termination

characteristics are more robust against the potential variability exhibited by individual atypical events.

4 Results

4.1 Spatio-temporal variability of drought termination

Drought termination chronologies for all 52 catchments, approximately ordered from the north-west (top) to the south-east (bottom) of the UK, are presented in Fig. 4. This allows visual inspection of the spatial coherence of drought events over a common data period beginning in the early 1970s. At a national scale, droughts have been relatively infrequent, occurring only in 1975–1977 and 1995–1998. Regional droughts affected southern and eastern areas in 1988–1993, 2004–2007, and 2009–2012. Drought-poor periods are also evident, the longest of which was the decade following the 1975–1977 event during which there were few prolonged droughts at either regional or national scales.

Prior to 1970, a lack of river flow data before gauged records commenced (particularly in northern and western areas of the UK; Table A1) limits the assessment of the spatial coherence of drought phases, but events in 1962–1964 and 1959 are identifiable in longer records in South-west UK, Anglian, Southern England and the Midlands. Persistent drought conditions (with intermittent drought terminations) within the 1890–1910 “Long Drought” (Marsh et al., 2007) are observed in the Thames river flow record from 1883.

Drought terminations show considerable spatio-temporal variability. For example, the 1988–1993 event had a notably uneven temporal evolution, with the transition to drought termination occurring early in the drought followed by a long drought termination phase for catchments in South-west UK and Anglian, whereas shorter drought terminations were apparent in the rest of the country. Fewer droughts have occurred in northern and western areas of the UK than in southern and eastern areas, while drought terminations tend to occur over longer time periods in the south. However, it is important to note the wide range of variability in drought termination characteristics exhibited within individual catchments. Two drought termination events are singled out for more detailed analysis: 1995–1998, the most widespread event since the 1970s; and 2009–2012, reported as unprecedented in the historical record (Parry et al., 2013).

4.2 Event analysis: 1995–1998

Drought in 1995–1998 affected all but one of the study catchments (Fig. 5; left), offering the best opportunity to analyse the spatial variability of drought termination within a single, severe event. The overall duration of drought was up to 3 years in the south and east in the UK but generally shorter in the north. There were two distinct patterns of drought termination. In the north and west, the drought termination phase began within 6 months of the start of drought development,

and long drought termination phases (three or more seasons) followed in 13 catchments. In contrast, drought termination started almost 2 years later in 25 catchments, mainly in the south and east. The transition to drought termination was generally spatially coherent across North & Central Wales, Midlands, South-west UK and Southern England, with the exceptions of the Conwy (NCW), Tywi (SWUK) and Great Ouse (SE).

Drought termination durations were generally longer (by 6 to 9 months) for catchments in the Southern England and Anglian regions (Fig. 5; top right). Conventionally referred to as the 1995–1997 drought in the literature (e.g. Marsh et al., 2013; Spraggs et al., 2015), it was the second half of 1998 before catchments in parts of lowland England (e.g. the Warwickshire Avon, Colne, Thames, Itchen and Dorset Avon) had completed the drought termination phase. The drought termination rate displayed a west–east divide in 1995–1998, particularly apparent for Wales, southern and eastern England, and the Midlands (Fig. 5; middle right). Whilst much of Wales and south-west England exhibited drought termination rates of 16–32 % per month, this decreased to less than 8 % per month across large areas of south-east England. Further north, the pattern was more mixed. Two-season drought terminations (Fig. 5; bottom right) generally were confined to the far northern parts of Scotland and England. Three-season drought terminations started in the autumn in Scotland and in the winter in Wales, south-west England and the Midlands. Long drought terminations (more than 8 months across four or more seasons) in many catchments in Western Scotland, Northern Ireland, North-west England, North-east England, Anglian, and Southern England prevented an assessment of drought termination seasonality.

4.3 Event analysis: 2009–2012

In contrast to the 1995–1998 event, the 2009–2012 drought was regional, primarily affecting North & Central Wales, South-west UK, Anglian, Southern England and the Midlands. The temporal sequencing of drought termination was also more regionally variable than in 1995–1998. Drought terminations began much sooner (early summer 2010) in North-west England, and had ended whilst drought continued to develop further south (Fig. 6; left). Drought terminations started in South-west UK up to a year before those in Anglian and the Midlands. In Anglian, Southern England and the Midlands, drought termination began in winter 2011/12 or spring 2012 and ended in late spring or early summer 2012. The end of the drought termination phase was much more spatially coherent in 2009–2012 than in 1995–1998.

Drought termination durations in 2009–2012 were generally 6 months or less (Fig. 6; top right), much shorter than those for 1995–1998. There was a gradient in drought termination duration from north-east to south-west across the affected catchments. The shortest durations (1–3 months) occurred across southern and eastern England and the Mid-




Figure 5. The 1995–1998 drought termination: chronologies of drought development and drought termination (a); drought termination duration (b); drought termination rate (c); drought termination seasonality (d).

lands, but lasted longer (10–18 months) for catchments in the south-west of England and Wales. The highest drought termination rates (more than 32 % per month) occurred in the largest catchments, whilst lower values (less than 16 % per month) were restricted to smaller catchments in Northern Ireland, North-east England and the far south of England (Fig. 6; middle right). Drought termination rates in 2009–

2012 showed a similar gradient to drought termination duration. There was more uniformity in drought termination rate across the drought-affected area for 2009–2012 than in 1995–1998, and drought terminations were generally more abrupt in 2009–2012.

There was greater seasonality for the 2009–2012 drought (Fig. 6; bottom right) than for the 1995–1998 event because




Figure 6. The 2009–2012 drought termination: chronologies of drought development and drought termination (a); drought termination duration (b); drought termination rate (c); drought termination seasonality (d).

drought terminations were generally shorter and started at different times. Catchments in southern and eastern England, the Midlands, and north Wales experienced drought terminations in spring and/or summer. Drought terminations in the winter months were uncommon for the 2009–2012 event. Winter drought terminations were restricted to the Warwick-

shire Avon (Midlands) and smaller catchments in the Anglian and Southern England regions.

4.4 Drought termination and catchment properties

The above analysis offers a qualitative assessment of the impact of catchment type on drought termination charac-

Table 1. Spearman correlations between drought termination characteristics and both catchment properties and drought development characteristics. Correlations are presented for individual events (rows $n = 467$) and for catchment mean drought characteristics (rows $n = 52$). Asterisks (*) denote statistical significance at the 95 % confidence level. Drought termination characteristics are defined as DTD: drought termination duration; and DTR: drought termination rate. Drought development characteristics are defined as DDD: drought development duration; and DM: drought magnitude. Catchment properties are denoted as SAAR6190: Standard-period Average Annual Rainfall for 1961–1990; and BFI: Base Flow Index.

	n	Catchment properties					Drought development characteristics	
		Area	Median elevation	SAAR 6190	BFI	Urban extent	DDD	DM
DTD	467	−0.02	−0.14*	−0.12*	0.06	0.13*	−0.29*	−0.18*
DTD	52	−0.24	−0.47*	−0.42*	0.19	0.34*	0.05	−0.02
DTR	467	0.02	0.12*	0.12*	−0.19*	−0.14*	0.28*	−0.05
DTR	52	0.13	0.19	0.15	−0.18	−0.35*	0.01	−0.17

teristics. Longer drought termination durations occurred in groundwater influenced catchments of southern and eastern England (e.g. the Stringside in Anglian and the Itchen and Dorset Avon in Southern England) during both 1995–1998 and 2009–2012, although this link does not apply for all identified drought termination events in the historical record. However, the synchronicity of the end of drought termination in spring 2012 (Fig. 6; left), when compared to the incoherent end of drought termination in 1995–1998 (Fig. 5; left), suggests that catchment properties are less influential during abrupt drought terminations than during gradual events.

Spearman correlations (r_s) between drought characteristics (magnitude, termination duration, and termination rate) and five catchment properties (catchment area, median elevation, SAAR6190, BFI, and urban extent) were calculated from the inventory of events. Correlations were assessed for individual drought events ($n = 467$) as well as for catchment averaged values ($n = 52$; Table 1) and were considered statistically significant where $p < 0.05$.

The highest r_s (−0.47; $p < 0.001$) was found for catchment average drought termination duration and median elevation, suggesting that upland catchments tend to experience shorter drought terminations. A similar correlation value was found between SAAR6190 ($r_s = -0.42$; $p = 0.002$) and drought termination duration, most likely due to the association between elevation and rainfall ($r_s = 0.71$; $p < 0.001$). Drought termination rate and urban extent are negatively correlated ($r_s = -0.35$; $p = 0.012$). This association may be influenced by a groundwater signal that is generally stronger in the more urbanized south and east of the UK, although r_s values for BFI and drought termination rate are small (−0.18; $p = 0.193$).

Spearman correlations were also derived for a subset of the study catchments, with 17 out of the 52 meeting the criteria of at least 10 identified drought termination events (Table A1). A statistically insignificant correlation was found between catchment average drought termination rate and BFI

($r_s = -0.42$; $p = 0.093$). This is consistent with the expectation of faster drought termination rates (i.e. more abrupt drought endings) in lower BFI (i.e. more responsive) catchments. For this subset of catchments, relationships between drought termination duration and both elevation and rainfall again corresponded to the highest values of r_s , but the linkage between urban extent and drought termination duration ($r_s = 0.44$; $p = 0.076$) was comparable.

For correlations between the properties of the drought development phase and drought termination characteristics, significant relationships were detected for drought development duration with both drought termination duration ($r_s = -0.29$; $p < 0.001$) and drought termination rate ($r_s = 0.28$; $p < 0.001$). This implies that sustained periods of drought development tend to be succeeded by shorter and more abrupt drought terminations. Relationships with catchment average drought development characteristics are not statistically significant, but assessments with the larger individual event dataset found that most associations (e.g. between drought magnitude and drought termination duration, or between drought development duration and drought termination rate) are significant.

5 Discussion

This study has systematically discretized drought terminations in historical river flow records for the UK for the first time. The detection method identified 467 drought events across 52 study catchments, providing a comprehensive inventory for further analysis of the historical variability of drought termination. Two aspects are explored here: a preliminary assessment of linkages between drought termination characteristics and catchment properties, including features of the preceding drought development phase (informed by the correlation analysis above); and a re-appraisal of drought termination characteristics in 2009–2012 within a broader hydrological context. In addition, this section also

corroborates the inventory of drought events and their terminations against existing work in the research literature, and considers the influence of the data and methodology on the results.

5.1 Drought termination characteristics and catchment properties

The spatio-temporal variability in drought termination within individual events (Figs. 4, 5, and 6) reflects the amount and timing of rainfall as well as its modulation by local catchment properties. This supports the findings of earlier studies that show hydrological drought termination to be more spatially variable than drought development, owing to the heterogeneity of catchment characteristics (e.g. Nkemdirim and Weber, 1999; Bell et al., 2013; DeChant and Moradkhani, 2015). However, the balance between the importance of rainfall distribution (in space and time) and catchment properties varies. In responsive catchments rainfall receipt will largely determine drought termination, whilst characteristics of the catchment may have more influence in those that are less responsive.

Some of the strongest correlations were found between drought termination duration and both elevation and catchment average rainfall (SAAR6190). This is likely to be because catchments in wetter upland areas of the UK are typically impermeable and responsive to rainfall, translating to shorter drought terminations. The correlations between urban extent and both drought termination duration and drought termination rate imply that drought terminations tend to be longer and more gradual in catchments with larger urban areas. This contradicts the expectation that typically impermeable urban areas may exhibit more abrupt drought terminations. The more urbanized catchments of the UK are generally in the south-east with more permeable geology, and it may be that lower responsiveness to rainfall negates the impact of the urban extent. Note also that the urban extent data are based on satellite imagery from 1998 to 2000 and, therefore, do not reflect the changing proportion of a catchment as built area outside of this short period. Further research could be undertaken to assess the impact of increasing urbanized area on changes in drought termination characteristics within certain study catchments under increasing development pressure (e.g. the Great Ouse in Southern England).

The BFI is widely regarded as a proxy for groundwater influence in the UK. However, water storage in lakes and seasonal snow cover can also be locally important, with BFI values of 0.43–0.60 for the Spey, Deveron, Scottish Dee and Naver in northern Scotland despite negligible groundwater influence. Whilst these impermeable catchments typically respond rapidly to rainfall, catchments with similar BFI values in areas of groundwater influence further south are less responsive. BFI is often considered to reflect catchment responsiveness, but the presence of lakes and/or snow cover in some responsive catchments of the north and west of the UK

mean that elevation is a better indicator of the spatial variability of responsiveness in the UK than BFI. This may explain why correlations between drought termination characteristics and elevation are stronger than those with BFI. By excluding catchments in Scotland that exhibit mismatches between BFI and responsiveness (through the use of the subset of 17 catchments with at least 10 events), the correlation analysis found a stronger association between drought termination rate and BFI. This linkage, as well as the qualitative observation of longer drought terminations in groundwater influenced catchments, is consistent with previous studies that report longer duration drought termination in soil moisture and groundwater levels (e.g. Eltahir and Yeh, 1999; Thomas et al., 2014).

Stronger relationships identified in the larger dataset between drought development and drought termination characteristics suggest that catchment averaging of metrics prior to correlation analysis may smooth out unique associations, resulting in information loss and obscuring some signals. A weak negative (but statistically significant) correlation was found between drought magnitude and drought termination duration, contrary to the pattern reported for two multi-year droughts in the US (Nkemdirim and Weber, 1999). The most important linkages were between drought development duration and both drought termination duration and drought termination rate.

5.2 Validating the chronologies of drought and drought termination

The rarity of national-scale droughts over the instrumental period (i.e. 1970s onwards) – limited to events in the mid-1970s and mid/late 1990s – corroborates previous work on regional drought in Europe (Hannaford et al., 2011). The locus of the 1988–1993 drought in the south-east of the UK confirms the chronology of Marsh et al. (2007). Time series of regional drought (Hannaford et al., 2011) identify a number of minor periods of river flow deficiency in the decade following the 1975–1977 event, but such episodes were not prolonged or severe enough to be detected in this study. However, the 1962–1964 drought was identifiable here despite the limited spatial coverage of river flow data. This event has been cited as an important multi-year drought at both UK and European scales (Parry et al., 2012). Similarly, Marsh et al. (2007) identify both the 1959 event and the 1890–1910 “Long Drought” when cataloguing major droughts in the UK. Whilst the use of standardized indicators (e.g. Hannaford et al., 2011) identifies the same amount of time under deficit conditions in each region, it is clear that stream-flow deficiencies are fewer but more prolonged in southern and eastern areas of the UK, confirming the results presented herein.

Validating the drought termination phases in Fig. 4 is less straightforward because of the relative lack of focus in the literature on the end of a drought relative to its other char-

Table 2. Catchments for which the drought termination rate during the 2009–2012 event was the largest of any previous event in the historical record.

Catchment	Number of drought events	Drought termination rate (% per month)		Year of drought termination ranking 2nd by drought termination rate
		2009–2012	Rank 2 event	
Severn	17	90.6	26.5	1997
Derwent	7	62.3	42.6	1976
Trent	11	56.3	28.0	1959/60
Warwickshire Avon	20	49.6	33.7	1963
Thames	35	38.1	37.2	1929/30
Teme	8	33.6	29.6	1975/76
Sydling Water	10	30.8	25.5	1974
Itchen	9	21.1	12.5	1963
Carron	3	18.2	11.9	2001

acteristics. Some of the longest drought termination durations correspond to the 1988–1993 drought, particularly for the Witham in the Anglian region, reflecting previous findings that the recovery from this drought was generally prolonged and particularly so in groundwater influenced catchments (Marsh et al., 1994). Conversely, the abrupt nature of drought terminations corresponding to the 1975–1977 event, evident in the chronologies presented herein, has been widely reported in the literature (e.g. Doornkamp et al., 1980; Rodda and Marsh, 2011).

5.3 Drought termination rate for 2009–2012 in a historical context

The rate of drought termination in 2009–2012 was particularly abrupt – more so than any other event identified in the post-1970 common data period. Almost a third (9 out of 31) of the drought-affected catchments in 2009–2012 registered new maxima for drought termination rate (Table 2). For the Severn, the drought termination in 2009–2012 was almost 4 times more abrupt than any other event since records began in 1929. This ranks amongst the top five most abrupt drought terminations for any event in any of the 52 study catchments ($n = 467$), although lagging substantially behind the most abrupt drought termination in this same dataset, the Whiteadder Water (Eastern Scotland) in 2004–2007, which was a third larger than the second ranked event. Drought magnitudes in 2009–2012 were not exceptional, but it was the differences between drought magnitudes and termination magnitudes over such short drought termination durations that were particularly noteworthy in establishing new maximum drought termination rates. This suggests that exceptional rainfall totals accumulated over short durations (assessed as greater than a 100-year return period; Bell et al., 2013) were more important than the severity of the preceding drought.

Research conducted in the immediate aftermath of the 2009–2012 event suggested that the drought termination was unprecedented in the historical record (Parry et al., 2013; Marsh et al., 2013). However, the assessment of the rarity of such abrupt transitions was based on ratios between average river flows over arbitrarily defined periods (May–July and the preceding December–March; Marsh et al., 2007). The more systematic approach adopted here allows an objective re-appraisal of the historical context across all timeframes. Although the drought termination event in 2009–2012 remains the most abrupt on record for the Thames (Table 2), there were three other comparably abrupt drought terminations between 1883 and 1930. This suggests that the rarity of the 2009–2012 drought termination may have been overstated (in the specific case of the Thames).

The drought termination phases in 2009–2012 and 2004–2007 were the most abrupt on record for 17 and 13 % of the 52 catchments, respectively; no other event registered new maxima in more than 10 % of the catchments, although this is difficult to assess consistently prior to 1970 due to limitations in data availability. These recent severe multi-year droughts featured consecutive dry winters (Wilby et al., 2015), supporting the view that long droughts result in more abrupt drought termination phases. However, the possibility that drought termination rates are becoming more abrupt warrants further exploration.

The wide variation in drought termination rates both between and within catchments suggests that different drought termination mechanisms are at work. Drought termination reflects a complex interplay of the specific hydroclimatic conditions with local catchment properties, even for groundwater influenced permeable catchments (in which the rainfall signal is substantially modulated by geology). Groundwater drought termination has been observed to be much slower than drought development in the western US (Bravar and Kavvas, 1991). Whether this applies to individual events in groundwater influenced catchments in this study would

depend on the extent to which deficits have propagated to groundwater. The artificial depletion of groundwater aquifers in Southern England may also have impacted drought termination characteristics in some catchments (e.g. the Itchen). The approach adopted in this study could be extended to groundwater level records as a further line of research. Similar variability in drought terminations was reported by Bonsal et al. (2011), and was attributed by Kam et al. (2013) to differences in rainfall intensity determined by the synoptic conditions (e.g. tropical cyclones).

5.4 Drought termination seasonality for 2009–2012 in a historical context

The drought termination in 2009–2012 occurred through the spring and early summer, an unusual but not unprecedented event. Only 8 of the 467 drought terminations occurred entirely in spring or in summer. Five of these eight relate to the 2009–2012 event (the Severn, Trent, Derwent and Witham in spring, and the Colne in summer). With the exception of the Severn, the drought termination in 2009–2012 is the only single-season event in the historical record for each catchment. Drought terminations across both spring and summer are similarly rare. Of the 13 events (out of 467) with spring–summer drought termination seasonality, five occurred in 2009–2012 (the Yscir, Exe, Thames, Itchen and Sydling Water; Fig. 6, bottom right). Of the remaining eight events, no other drought termination is represented by more than two catchments. For the Thames, the only previous example of a drought termination entirely within the spring and summer was in 1888. Other studies have also found that it is unlikely that multi-season droughts will terminate in two seasons or less (Karl et al., 1987).

Rather than simply the wettest season, it is the season with the greatest potential for large positive rainfall anomalies that is most likely to facilitate drought termination (Karl et al., 1987; Mo, 2011). In the UK these two factors coincide; hence, winter provides the greatest likelihood for drought termination (Van Loon et al., 2014). The larger evaporative demand in summer reduces the effectiveness of all but the most extreme rainfall, explaining the tendency for drought terminations in the winter half-year. Of the 467 drought terminations, single season events were more common in autumn (eight) and winter (eight) than in spring (five) and particularly summer (three).

At regional scales, variation in drought termination seasonality is likely to be determined by catchment properties, such as storage causing lagged responses. For catchments in Scotland, the influence of snow may also influence drought termination. Where seasonal snowpacks exist, winter drought terminations may be delayed until the snowmelt season (Van Loon et al., 2014). However, the large variability of drought termination characteristics and the moderate to weak correlations with catchment properties imply that a range of physical processes are involved. At national or

continental scales, larger-scale drivers such as El Niño and La Niña events in the Pacific (e.g. Tomasella et al., 2011; Marengo and Espinoza, 2016), switches in Atlantic temperatures (Wilby, 2001; Folland et al., 2015), and tropical cyclones (e.g. Kam et al., 2013; Patterson et al., 2013) have been shown to be a factor in drought termination events. Further research is required to assess the extent to which changes in these and other synoptic drivers might be influencing the seasonality of drought terminations in the UK. For instance, Matthews et al. (2016) report relatively low frequencies of summer cyclones in the period 1961–1990 but a marked resurgence in counts since the 1990s.

5.5 Impact of methodology and data on results

Although the detection procedure utilized herein applied consistent rules, the parameter values used to define a drought and its phases can influence the resulting chronology. This is illustrated by the sensitivity analysis (Fig. 3) and has been reported by other studies (e.g. Patterson et al., 2013). Drought termination phases following shorter drought developments, for example driven by summer heatwaves, would not be well represented by the parameter settings used in this study. This is because the parameters which determine the initiation of drought development (D and R) require below-average river flows for at least 9 of 10 consecutive months, a timeframe which is too prolonged to adequately characterize typical single season drought events. In addition, events in the more hydrologically responsive north and west of the UK might be less well represented because droughts in these wetter regions are typically shorter than multi-season in duration. However, the spatial variability in the number of identified droughts is consistent with the levels of service set by regional water companies, with drought-induced water restrictions expected more frequently in the south-east of the UK than in the north. Nevertheless, there is a need to more comprehensively assess the sensitivity of derived chronologies of drought termination to the choice of detection parameters.

The monthly time step used in this study may also be limiting. Drought termination can occur rapidly, perhaps within a few days in some instances of intense cyclonic activity. Under these circumstances, monthly data may obscure accurate definitions of the end of drought termination or underestimate the drought termination rate. In addition, the use of a monthly average flow threshold is higher than those usually applied in threshold-based studies. Low flow thresholds such as Q_{70} (Hisdal et al., 2001) and Q_{80} (e.g. Tallaksen et al., 2009) have been widely used in the literature, and threshold levels between Q_{70} and Q_{90} are generally considered appropriate (Fleig et al., 2006). The use of an average flow threshold would be expected to increase the overall duration of drought (as illustrated by Tallaksen et al., 1997) as well as the drought development and drought termination phases. However, applying a lower threshold would sub-divide well-

known multi-year drought events (e.g. 1995–1998 and 2009–2012 from this study) into a number of more severe episodes, each with their own drought termination. In order to focus on multi-season to multi-year droughts, a higher threshold is required. A previous study that applied thresholds between Q_{50} and Q_{90} found that a higher threshold level identified more multi-year droughts (Tallaksen et al., 1997). It is acknowledged that the suitability of different thresholds is specific to individual perceptions or applications.

The approach utilized in this study focuses on the status of river flows, which can increase substantially over relatively short timescales and replenish water supplies rapidly without having to account for a deficit that has accumulated during the drought development phase. However, it is acknowledged that deficit volume approaches (in which the accumulated volume of water “lost” during drought development is recovered) may be important for studies which focus on the overall water balance.

The potential influence of abstractions from surface and groundwater sources during drought development may artificially extend the duration of the drought termination phase. The catchments used in this study include some of the largest in the UK in order to maximize spatial coverage, and few of these could be described as near-natural. Abstractions to meet higher water demand during drought development, particularly during heatwave conditions, combine with lower natural recharge. Drought-terminating rainfall must account for this “anthropogenic deficit” in addition to the natural hydrological deficit. There is a regional bias in the anthropogenic influence on river flows, with more impacted catchments in the south and east of the UK and more near-natural catchments in the north and west. Whilst this spatial pattern also reflects the number of droughts identified, the selection of parameters that favour major multi-season droughts is probably more influential. The use of monthly mean river flows may also dilute the impact of artificial influences on individual days.

6 Conclusions

For the first time, terminations of multi-season to multi-year droughts in the UK have been systematically identified and characterized. This study detected 467 events in 52 catchments covering a range of geographical settings, and provides chronologies of both drought development and drought termination phases. This information provides a new perspective on the historical variability of drought termination in the UK that is potentially useful for water resource managers and researchers in a range of fields including ecology, geomorphology and water quality. It is hoped that characterizing 467 drought termination events will underpin further research into any emerging trends and provide the basis for the development of a drought termination typology. It should be noted that the chronology of drought termina-

tion presented herein has been derived using parameters that were informed by a sensitivity analysis and ensuring a focus on multi-season to multi-year droughts in the UK. For other applications across a range of locations and/or considering alternative definitions of droughts, it is recognized that alternative parameters may be required.

Investigations into the link between drought termination characteristics and catchment properties or drought development characteristics would be strengthened by a larger sample of events. Stronger correlations were found for catchment average drought termination metrics when using the subset of catchments with at least 10 identified events, although this subset is biased towards catchments with longer records predominantly in southern and eastern areas of the UK. The BFI is not an adequate predictor of the responsiveness of a catchment. Further exploration of potential linkages between drought termination characteristics and catchment properties should seek to use variables which are more closely related to river flow responsiveness than BFI (e.g. a flashiness index; Baker et al., 2007). The use of potential associations between drought termination characteristics and those of the preceding drought development phase by water resource managers is constrained by weak to moderate correlations and requires further research before useful conclusions can be drawn. Ideally, coupled land–atmosphere model experiments would be performed to explore possible links between drought duration or magnitude and terminating rainfall mechanisms.

The identification and characterization of 467 drought terminations have provided a comprehensive historical context within which to place the notable 2009–2012 event. This illustrates the variability of drought termination characteristics in the UK, re-assessing the conclusion (based on a subset of newsworthy examples) that droughts tend to terminate abruptly. The long-term context could be improved further through the use of river flow reconstructions (e.g. Jones and Lister, 1998; Jones et al., 2006) to “fill in the grey space” in Fig. 4, which represents the best historical perspective provided by available observed data. The method used in this study has the flexibility to produce similarly comprehensive chronologies of drought termination in groundwater level records, water quality metrics or ecological indices, to trace the propagation of drought termination throughout the river system and hydrological cycle. Drought termination in river flows and groundwater levels may not synchronize even within the same catchment due to lagged response times. Hence, even when a drought terminates abruptly with severe river flooding, (contrary to public expectations) water restrictions may not be removed until groundwater levels respond. The complexities associated with this propagation of drought termination require further research.

7 Data availability

The river flow data used in this study can be accessed from the National River Flow Archive (<http://nrfa.ceh.ac.uk/>), the UK's focal point for and principle repository of hydrometric data. All of the data for the 52 catchments used herein are freely available to download.

Appendix A

Table A1. Metadata for the 52 study catchments. The subset of 17 catchments referred to in Sects. 4.4 and 5.1 is indicated with asterisks (*).

Region	Catchment	Record length (years)	Area (km ²)	Median elevation (m)	SAAR6190 (mm)	BFI	Urban extent (%)
W Scotland	Naver	37	477	187	1384	0.43	0.0
W Scotland	Carron	35	138	342	2620	0.26	0.0
W Scotland	Nevis	32	69	518	2912	0.27	0.1
W Scotland	Clyde	51	1903	252	1129	0.46	3.0
W Scotland	Ayr	38	574	212	1214	0.30	0.6
W Scotland	Cree	51	368	212	1760	0.28	0.2
W Scotland	Nith	37	477	288	1460	0.39	0.2
E Scotland	Findhorn	56	782	408	1064	0.40	0.0
E Scotland	Spey*	62	2861	420	1120	0.60	0.1
E Scotland	Deveron*	54	955	209	928	0.57	0.2
E Scotland	Scottish Dee*	85	1370	508	1109	0.53	0.1
E Scotland	Tay	62	4587	395	1425	0.65	0.2
E Scotland	Forth	33	1036	180	1752	0.41	0.0
E Scotland	Whiteadder Water	45	503	230	813	0.51	0.2
E Scotland	Tweed	52	4390	255	955	0.52	0.3
N Ireland	Mourne	32	1844	153	1288	0.39	0.3
N Ireland	Faughan	38	273	173	1219	0.47	0.4
N Ireland	Lagan	42	492	95	916	0.43	3.2
NW England	Eden	47	2287	210	1183	0.49	0.8
NW England	Kent	46	209	205	1732	0.41	1.8
NW England	Ribble	54	1145	198	1353	0.34	3.7
NE England	South Tyne	52	751	333	1148	0.34	0.2
NE England	Tees	58	818	370	1141	0.34	0.4
NE England	Ure	56	915	264	1118	0.39	0.8
NE England	Derwent	41	1586	102	765	0.70	0.8
N&C Wales	Conwy	50	345	328	2055	0.28	0.1
N&C Wales	Welsh Dee	77	1013	347	1369	0.54	0.4
N&C Wales	Severn*	93	4325	127	913	0.53	2.0
N&C Wales	Teme	44	1480	191	818	0.55	0.7
N&C Wales	Wye*	78	4010	199	1011	0.54	0.7
Midlands	Trent*	56	7486	118	761	0.64	10.5
Midlands	Warwickshire Avon*	78	2210	96	654	0.51	4.9
SW UK	Tywi	56	1090	220	1534	0.47	0.2
SW UK	Yscir	42	63	361	1299	0.46	0.0
SW UK	Tone	53	202	120	966	0.60	1.6
SW UK	Torridge*	54	663	146	1186	0.38	0.4
SW UK	Exe*	58	601	235	1248	0.50	0.6
SW UK	Dart	56	248	347	1765	0.52	0.7
SW UK	Warleggan	45	25	232	1442	0.70	0.2
SW UK	Sydling Water*	45	12	190	1032	0.88	0.5
Anglian	Lud	46	55	89	699	0.90	2.2
Anglian	Witham*	55	298	91	614	0.69	3.5
Anglian	Bedford Ouse*	81	1460	101	636	0.53	3.5
Anglian	Stringsides	49	99	20	629	0.84	0.7
Anglian	Wensum	45	398	57	684	0.75	1.3
Anglian	Colne*	55	238	68	566	0.52	2.2
S England	Thames*	131	9948	100	706	0.63	6.6
S England	Great Stour*	50	345	75	747	0.70	3.2
S England	Bull	36	41	58	820	0.37	0.9
S England	Itchen	56	360	107	833	0.96	2.9
S England	Dorset Avon*	49	324	129	745	0.91	1.3
S England	Stour*	41	1073	83	861	0.64	2.0

Author contributions. Simon Parry devised the approach, selected the catchments, and coordinated the writing of the paper. Robert L. Wilby provided input on the structure and content of the paper and the impetus for the correlation analysis. Christel Prudhomme and Paul J. Wood provided feedback on the different paper structures and content. All authors contributed to the manuscript writing and commented on the analyses.

Acknowledgements. This research was funded through the Learning & Development programme at the Centre for Ecology & Hydrology (CEH), as well as the Natural Environment Research Council's (NERC) "Analysis of historic drought and water scarcity in the UK" (NERC grant ref.: NE/L01016X/1) and "Improving predictions of drought to inform user decisions (IMPETUS)" (NERC grant ref.: NE/L010267/1) projects. River flow data and catchment metadata were provided by the UK National River Flow Archive at CEH. The manuscript was improved following valuable feedback provided by two reviewers, Henny van Lanen and an anonymous reviewer, and by journal editor Lena Tallaksen. The authors would also like to thank Katie Muchan, Filip Kral, Lucy Barker, and Shaun Harrigan (all CEH) for their assistance with river flow data and metadata, spatial data, graphics, and statistics, respectively.

Edited by: L. M. Tallaksen

Reviewed by: Henny van Lanen and one anonymous referee

References

- Baker, D. B., Richards, R. P., Loftus, T. T., and Kramer, J. W.: A new flashiness index: characteristics and applications to midwestern rivers and streams, *J. Am. Water Resour. As.*, 40, 503–522, 2007.
- Bell, V. A., Davies, H. N., Kay, A. L., Marsh, T. J., Brookshaw, A., and Jenkins, A.: Developing a large-scale water-balance approach to seasonal forecasting: application to the 2012 drought in Britain, *Hydrol. Process.*, 27, 3003–3012, 2013.
- Bonsal, B. R., Wheaton, E. E., Meinert, A., and Siemens, E.: Characterizing the Surface Features of the 1999–2005 Canadian Prairie Drought in Relation to Previous Severe Twentieth Century Events, *Atmos.-Ocean*, 49, 320–338, 2011.
- Bravar, L. and Kavvas, M. L.: On the physics of droughts, II. Analysis and simulation of the interaction of atmospheric and hydrologic processes during droughts, *J. Hydrol.*, 129, 299–330, 1991.
- Chew, C. C. and Small, E. E.: Terrestrial water storage response to the 2012 drought estimated from GPS vertical position anomalies, *Geophys. Res. Lett.*, 41, 6145–6151, 2014.
- Correia, F. N., Santos, M. A., and Rodrigues, R. R.: Engineering Risk In Regional Drought Studies, in: *Engineering Reliability and Risk in Water Resources*, Proceedings of the NATO Advanced Study Institute on "Engineering Reliability and Risk in Water Resources", edited by: Duckstein, L. and Plate, E. J., Tucson, Arizona, USA, 19 May–2 June 1985, Martinus Nijhoff Publishers, Dordrecht, 1987.
- DeChant, C. M. and Moradkhani, H.: Analyzing the sensitivity of drought recovery forecasts to land surface initial conditions, *J. Hydrol.*, 526, 89–100, 2015.
- Doornkamp, J. C., Gregory, K. J., and Burn, A. S.: *Atlas of Drought in Britain 1975–76*, Institute of British Geographers, London, 1980.
- Eltahir, E. A. B. and Yeh, P. J.-F.: On the asymmetric response of aquifer water level to floods and droughts in Illinois, *Water Resour. Res.*, 35, 1199–1217, 1999.
- Fleig, A. K., Tallaksen, L. M., Hisdal, H., and Demuth, S.: A global evaluation of streamflow drought characteristics, *Hydrol. Earth Syst. Sci.*, 10, 535–552, doi:10.5194/hess-10-535-2006, 2006.
- Fleig, A. K., Tallaksen, L. M., Hisdal, H., and Hannah, D. M.: Regional hydrological drought in north-western Europe: linking a new Regional Drought Area Index with weather types, *Hydrol. Process.*, 25, 1163–1179, 2011.
- Folland, C. K., Hannaford, J., Bloomfield, J. P., Kendon, M., Svensson, C., Marchant, B. P., Prior, J., and Wallace, E.: Multi-annual droughts in the English Lowlands: a review of their characteristics and climate drivers in the winter half-year, *Hydrol. Earth Syst. Sci.*, 19, 2353–2375, doi:10.5194/hess-19-2353-2015, 2015.
- Gustard, A., Bullock, A., and Dixon, J. M.: *Low Flow Estimation in the United Kingdom*, Institute of Hydrology Report No. 108, 1992.
- Hannaford, J., Lloyd-Hughes, B., Keef, C., Parry, S., and Prudhomme, C.: Examining the large-scale spatial coherence of European drought using regional indicators of precipitation and streamflow deficit, *Hydrol. Process.*, 25, 1146–1162, 2011.
- Hisdal, H., Stahl, K., Tallaksen, L. M., and Demuth, S.: Have streamflow droughts in Europe become more severe or frequent?, *Int. J. Climatol.*, 21, 317–333, 2001.
- Jones, P. D. and Lister, D. H.: Riverflow reconstructions for 15 catchments over England and Wales and an assessment of hydrologic drought since 1865, *Int. J. Climatol.*, 18, 999–1013, 1998.
- Jones, P. D., Lister, D. H., Wilby, R. L., and Kostopoulou, E.: Extended riverflow reconstructions for England and Wales, 1865–2002, *International J. Climatol.*, 26, 219–231, 2006.
- Kam, J., Sheffield, J., Yuan, X., and Wood, E. F.: The Influence of Atlantic Tropical Cyclones on Drought over the Eastern United States, *J. Climate*, 26, 3067–3086, 2013.
- Karl, T., Quinlan, F., and Ezell, D. S.: Drought Termination and Amelioration: Its Climatological Probability, *J. Climate Appl. Meteorol.*, 26, 1198–1209, 1987.
- Kienzie, S. W.: The Use of the Recession Index as an Indicator for Streamflow Recovery After a Multi-Year Drought, *Water Resour. Manage.*, 20, 991–1006, 2006.
- Lange, J. and Haensler, A.: Runoff generation following a prolonged dry period, *J. Hydrol.*, 464–465, 157–164, 2012.
- Lavers, D. A. and Villarini, G.: Were global numerical weather prediction systems capable of forecasting the extreme Colorado rainfall of 9–16 September 2013?, *Geophys. Res. Lett.*, 40, 1–6, 2013.
- Li, S., Xiong, L., Dong, L., and Zhang, J.: Effects of the Three Gorges Reservoir on the hydrological droughts at the downstream Yichang station during 2003–2011, *Hydrol. Process.*, 27, 3981–3993, 2013.
- Marengo, J. A. and Espinoza, J. C.: Extreme seasonal droughts and floods in Amazonia: causes, trends and impacts, *Int. J. Climatol.*, 36, 1033–1050, doi:10.1002/joc.4420, 2016.
- Marengo, J. A., Nobre, C. A., Tomasella, J., Oyama, M. D., de Oliveira, G. S., de Oliveira, R., Camargo, H., Alves, L. M., and

- Brown, I. F.: The Drought of Amazonia in 2005, *J. Climate*, 21, 495–516, 2008.
- Marsh, T. J. and Hannaford, J. (Eds.): UK Hydrometric Register. Hydrological data UK series, NERC Centre for Ecology & Hydrology, Wallingford, 2008.
- Marsh, T. J., Monkhouse, R. A., Arnell, N. W., Lees, M. L., and Reynard, N. S.: The 1988–92 drought, NERC Institute of Hydrology, Wallingford, 1994.
- Marsh, T., Cole, G., and Wilby, R.: Major droughts in England and Wales, 1800–2006, *Weather*, 62, 87–93, 2007.
- Marsh, T. J., Parry, S., Kendon, M. C., and Hannaford, J.: The 2010–12 drought and subsequent extensive flooding, NERC Centre for Ecology & Hydrology, Wallingford, 2013.
- Matthews, T., Murphy, C., Wilby, R. L., and Harrigan, S.: A cyclone climatology of the British-Irish Isles 1871–2012, *Int. J. Climatol.*, 36, 1299–1312, doi:10.1002/joc.4425, 2016.
- Maxwell, J. T., Ortegen, J. T., Knapp, P. A., and Soule, P. T.: Tropical Cyclones and Drought Amelioration in the Gulf and South-eastern Coastal United States, *J. Climate*, 26, 8440–8452, 2013.
- Miller, J. D., Gaskin, G. J., and Anderson, H. A.: From Drought to Flood: Catchment Responses Revealed using Novel Soil Water Probes, *Hydrol. Process.*, 11, 533–541, 1997.
- Mo, K. C.: Drought onset and recovery over the United States, *J. Geophys. Res.*, 116, D20106, doi:10.1029/2011JD016168, 2011.
- Ning, N. S. P., Gawne, B., Cook, R. A., and Nielsen, D. L.: Zooplankton dynamics in response to the transition from drought to flooding in four Murray-Darling basin rivers affected by differing levels of flow regulation, *Hydrobiologia*, 702, 45–62, 2013.
- Nkemdirim, L. and Weber, L.: Comparison between the Droughts of the 1930s and the 1980s in the Southern Prairies of Canada, *J. Climate*, 12, 2434–2450, 1999.
- Parry, S., Hannaford, J., Lloyd-Hughes, B., and Prudhomme, C.: Multi-year droughts in Europe: analysis of development and causes, *Hydrol. Res.*, 43, 689–706, 2012.
- Parry, S., Marsh, T., and Kendon, M.: 2012: from drought to floods in England and Wales, *Weather*, 68, 268–274, 2013.
- Parry, S., Prudhomme, C., Wilby, R. L., and Wood, P. J.: Chronology of drought termination for long records in the Thames catchment, in: *Drought: research and science-policy interfacing*, edited by: Andreu, J., Solera, A., Paredes-Arquiola, J., Haro-Montegudo, D., and Van Lanen, H. A. J., Leiden, CRC Press, Balkema, 165–170, 2015.
- Parry, S., Prudhomme, C., Wilby, R. L., and Wood, P. J.: Drought termination: Concept and characterisation, *Prog. Phys. Geogr.*, doi:10.1177/0309133316652801, 2016.
- Patterson, L. A., Lutz, B. D., and Doyle, M. W.: Characterization of Drought in the South Atlantic, United States, *J. Am. Water Resour. Assoc.*, 49, 1385–1397, 2013.
- Pfister, C., Weingartner, R., and Luterbacher, J.: Hydrological winter droughts over the last 450 years in the Upper Rhine basin: a methodological approach, *Hydrol. Sci. J.*, 51, 966–985, 2006.
- Rodda, J. C. and Marsh, T. J.: The 1975–76 Drought – a contemporary and retrospective review. NERC Centre for Ecology & Hydrology, Wallingford, 2011.
- Shukla, S., Steinemann, A. C., and Lettenmaier, D. P.: Drought Monitoring for Washington State: Indicators and Applications, *J. Hydrometeorol.*, 12, 66–83, 2011.
- Spackman, E.: Calculation and Mapping of Rainfall Averages for 1961–90, University of Salford, Manchester, 1993.
- Spearman, C. E.: “General intelligence”, objectively determined and measured, *Am. J. Psychol.*, 15, 201–293, 1994.
- Spraggs, G., Peaver, L., Jones, P., and Ede, P.: Re-construction of historic drought in the Anglian Region (UK) over the period 1798–2010 and the implications for water resources and drought management, *J. Hydrol.*, 526, 231–252, 2015.
- Tallaksen, L. M., Madsen, H., and Clausen, B.: On the definition and modelling of streamflow drought duration and deficit volume, *Hydrol. Sci. J.*, 42, 15–33, 1997.
- Tallaksen, L. M., Hisdal, H., and Van Lanen, H. A. J.: Space-time modelling of catchment scale drought characteristics, *J. Hydrol.*, 375, 363–372, 2009.
- Thomas, A. C., Reager, J. T., Famiglietti, J. S., and Rodell, M.: A GRACE-based water storage deficit approach for hydrological drought characterization, *Geophys. Res. Lett.*, 41, 1537–1545, 2014.
- Tomasella, J., Borma, L. S., Marengo, J. A., Rodriguez, D. A., Cuartas, L. A., Nobre, C. A., and Prado, M. C. R.: The droughts of 1996–1997 and 2004–2005 in Amazonia: hydrological response in the river main-stem, *Hydrol. Process.*, 25, 1228–1242, 2011.
- Van Loon, A. F. and Van Lanen, H. A. J.: A process-based typology of hydrological drought, *Hydrol. Earth Syst. Sci.*, 16, 1915–1946, doi:10.5194/hess-16-1915-2012, 2012.
- Van Loon, A. F., Tiedeman, E., Wanders, N., Van Lanen, H. A. J., Teuling, A. J., and Uijlenhoet, R.: How climate seasonality modifies drought duration and deficit, *J. Geophys. Res.-Atmos.*, 119, 4640–4656, 2014.
- Van Loon, A. F., Ploum, S. W., Parajka, J., Fleig, A. K., Garnier, E., Laaha, G., and Van Lanen, H. A. J.: Hydrological drought types in cold climates: quantitative analysis of causing factors and qualitative survey of impacts, *Hydrol. Earth Syst. Sci.*, 19, 1993–2016, doi:10.5194/hess-19-1993-2015, 2015.
- Wang, H., Jia, L., Steffen, H., Wu, P., Jiang, L., Hsu, H., Xiang, L., Wang, Z., and Hu, B.: Increased water storage in North America and Scandinavia from GRACE gravity data, *Nat. Geosci.*, 6, 38–42, 2013.
- Webster, P. J., Toma, V. E., and Kim, H.-M.: Were the 2010 Pakistan floods predictable?, *Geophys. Res. Lett.*, 38, L04806, doi:10.1029/2010GL046346, 2011.
- Wilby, R. L.: Downscaling summer rainfall in the UK from North Atlantic ocean temperatures, *Hydrol. Earth Syst. Sci.*, 5, 245–257, doi:10.5194/hess-5-245-2001, 2001.
- Wilby, R. L., Prudhomme, C., Parry, S., and Muchan, K. G. L.: Persistence of hydrometeorological droughts in the United Kingdom: A regional analysis of multi-season rainfall and river flow anomalies, *Journal of Extreme Events*, 2, 1550006, doi:10.1142/S2345737615500062, 2015.
- Yang, S., Wu, B., Zhang, R., and Zhou, S.: Relationship Between an Abrupt Drought-Flood Transition over Mid-Low Reaches of the Yangtze River in 2011 and the Intra-seasonal Oscillation over Mid-High Latitudes, *Ac. Meteorol. Sin.*, 27, 129–143, 2012.
- Zelenhasi, E. and Salvai, A.: A Method of Streamflow Drought Analysis, *Water Resour. Res.*, 23, 156–168, 1987.