

Supplement of Hydrol. Earth Syst. Sci., 22, 3493–3513, 2018
<https://doi.org/10.5194/hess-22-3493-2018-supplement>
© Author(s) 2018. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

The temporally varying roles of rainfall, snowmelt and soil moisture for debris flow initiation in a snow-dominated system

Karin Mostbauer et al.

Correspondence to: Karin Mostbauer (karin.mostbauer@students.boku.ac.at)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Model equations

Table S1. Water balance equations and constitutive functions (abbreviations see Table 1 and sect. 2.2, 3.1.1, 3.2).

Reservoir	Water balance equation	Constitutive functions
Snow reservoir	$\Delta S_{snow}/\Delta t = P_s - M$	<p>100 m elevation zones:</p> $P = P_l + P_s$ $P_s = \begin{cases} P & T_{mean} \geq T_{temp} \\ 0 & T_{mean} < T_{temp} \end{cases}$ $M = \begin{cases} \min \begin{cases} melt_f \cdot (T_{mean} - T_{temp}) \\ S_{snow} \cdot t^{-1} + M_{glacier} \end{cases} & T_{mean} \geq T_{temp} \\ 0 & T_{mean} < T_{temp} \end{cases}$ $M_{glacier} = (melt_f \cdot (T_{mean} - T_{temp}) - S_{snow} \cdot t^{-1}) \cdot \text{glacier extent}$ <p>[assumption: $S_{glacier}$ is unlimited]</p>
Unsaturated reservoir	$\Delta S_u/\Delta t = P_l + M - Q_{uf} - Q_{up} - E_a - Q_{us}$	$Q_{uf} = (1-D) \cdot C_r \cdot (P_l + M); \quad C_r = 1/[1 + e^{(-S_u/S_{u,max}+0.5)/\beta}]$ $Q_{up} = D \cdot C_r \cdot (P_l + M); \quad C_r = 1/[1 + e^{(-S_u/S_{u,max}+0.5)/\beta}]$ $E_a = E_p \cdot \min \begin{cases} S_u/S_{u,max} \cdot 1/L_p \\ 1 \end{cases}$ $Q_{us} = S_u / S_{u,max} \cdot P_{max}$
Fast reservoir	$\Delta S_f/\Delta t = Q_{uf} - Q_f$	$Q_f = S_f \cdot (1 - e^{-K_f \cdot t}) \cdot t^{-1}$
Slow reservoir	$\Delta S_s/\Delta t = Q_{up} + Q_{us} - Q_s$	$Q_s = S_s \cdot (1 - e^{-K_s \cdot t}) \cdot t^{-1}$

Boxplots






Figure S1. Examples of ill-performing system variables (a-b) and ill-performing absolute x-axis definitions for storage elements (c-d): both the debris flow event days' boxplots (blue) and the threshold values' boxplots (grey) show wide variations within the behavioral models i.e. the classification of the debris flows' system variables (high moisture, low moisture, etc.) would – to a greater extent – depend on which model parameter set one considers as "correct" (which cannot be determined due to model equifinality).


Timeseries plots




(d) 1964






(e) 1965




(f) 1966







(j) 1982



(k) 1985




(l) 1987




(m) 1989

21




(n) 1998

22
a-d



(o) 2003

23






Figure S2. Observed daily stream flow Q_{obs} (black solid line), daily mean temperature T_{mean} at mean elevation (grey solid line) and maximum temperature T_{max} at mean elevation (black solid line) as well as, based on observed precipitation data, modelled daily rainfall P_r (dark blue downward columns for 5th percentile, incl. grey downward columns for 95th percentile), daily snowfall P_s (light blue downward columns for 5th percentile, incl. grey downward columns for 95th percentile) and daily snowmelt M (dark blue upward columns for 5th percentile, incl. grey upward columns for 95th percentile), modelled stream flow (dark blue line for the median and the grey shaded area for the 5/95th percentiles of all behavioral model solutions) and modelled relative soil moisture (solid blue line for the median and the grey shaded area for the 5/95th percentiles) for all years with debris flow occurrence.