

Input parameter	Description	Values (using ^{18}O as a tracer)	Parametric sources
h	Relatively humidity	0.63	Measured by the humidity meter
T ($^{\circ}\text{C}$)	Water temperature	15.66	Monitored with divers
δ_{surface} (^{18}O) ‰	Surface water isotopic compositions	-12.45	Average value of surface inflow samples
δ_{gw} (^{18}O) ‰	Groundwater isotopic compositions	-11.97	Average value of porewater samples
δ_{L} (^{18}O) ‰	Lake water isotopic compositions	-12.54	Average value of Ximen Co Lake water samples
F_{gw} (mm d^{-1})	LGD rates	14.18	Calculated based on the ^{222}Rn mass balance model
ε^* (^{18}O) ‰	Effective equilibrium isotopic enrichment factor	10.12	Eqs. (S13)–(S14)
C_{k} (^{18}O) ‰	Kinetic constant for ^{18}O	14.2	Constants based on the evaporating experiment
ε_{k} (^{18}O) ‰	Kinetic enrichment factor	5.2	From Eq. (S15)
ε (^{18}O) ‰	Total isotopic enrichment factor	15.33	The sum of ε^* and C_{k}
α^* (^{18}O) ‰	Effective isotopic equilibrium factor	1.01	$\alpha^* = 1 + \varepsilon^*$
δ_{a} (^{18}O) ‰	Isotopic composition of ambient air	-23.12	Estimated with δ_{in} and δ_{a}
δ_{in} (^{18}O) ‰	Isotopic composition of surface inflow water	-13.41	Average value of surface inflow water
δ_{E} (^{18}O) ‰	Isotopic compositions of evaporating vapour	-35.1	From Eq. (S12)
$[\text{Cl}^-]_{\text{in}}$ (mgL^{-1})	Chloride concentrations in surface inflow water	0.91	Filed data
$[\text{Cl}^-]_{\text{L}}$ (mgL^{-1})	Chloride concentrations in lake water	1.02	Filed data
$[\text{Cl}^-]_{\text{gw}}$ (mgL^{-1})	Chloride concentrations in groundwater	1.48	Filed data