



# Groundwater–glacier meltwater interaction in proglacial aquifers

Brighid É. Ó Dochartaigh<sup>1</sup>, Alan M. MacDonald<sup>1</sup>, Andrew R. Black<sup>2</sup>, Jez Everest<sup>1</sup>, Paul Wilson<sup>3</sup>, W. George Darling<sup>4</sup>, Lee Jones<sup>5</sup>, and Mike Raines<sup>5</sup>

<sup>1</sup>British Geological Survey, Lyell Centre, Research Avenue South, Edinburgh EH14 4AP, UK

<sup>2</sup>School of Social Sciences, University of Dundee, Dundee DD1 4HN, UK

<sup>3</sup>Geological Survey of Northern Ireland, Dundonald House, Upper Newtownards Road, Belfast BT4 3SB, UK

<sup>4</sup>British Geological Survey, Maclean Building, Wallingford OX10 8BB, UK

<sup>5</sup>British Geological Survey, Environmental Science Centre, Keyworth NG12 5GG, UK

**Correspondence:** Alan M. MacDonald (amm@bgs.ac.uk)

Received: 15 March 2019 – Discussion started: 24 April 2019

Revised: 30 July 2019 – Accepted: 22 September 2019 – Published: 5 November 2019

**Abstract.** Groundwater plays a significant role in glacial hydrology and can buffer changes to the timing and magnitude of flows in meltwater rivers. However, proglacial aquifer characteristics or groundwater dynamics in glacial catchments are rarely studied directly. We provide direct evidence of proglacial groundwater storage, and quantify multi-year groundwater–meltwater dynamics, through detailed aquifer characterisation and intensive high-resolution monitoring of the proglacial system of a rapidly retreating glacier, Virkisjökull, in south-eastern Iceland. Proglacial unconsolidated glaciofluvial sediments comprise a highly permeable aquifer ( $25\text{--}40 \text{ m d}^{-1}$ ) in which groundwater flow in the shallowest 20–40 m of the aquifer is equivalent to 4.5 % (2.6 %–5.8 %) of mean river flow, and 9.7 % (5.8 %–12.3 %) of winter flow. Estimated annual groundwater flow through the entire aquifer thickness is 10 % (4 %–22 %) the magnitude of annual river flow. Groundwater in the aquifer is actively recharged by glacier meltwater and local precipitation, both rainfall and snowmelt, and strongly influenced by individual precipitation events. Local precipitation represents the highest proportion of recharge across the aquifer. However, significant glacial meltwater influence on groundwater within the aquifer occurs in a 50–500 m river zone within which there are complex groundwater–river exchanges. Stable isotopes, groundwater dynamics and temperature data demonstrate active recharge from river losses, especially in the summer melt season, with more than 25 % and often > 50 % of groundwater in the near-river aquifer zone sourced from glacier meltwater. Proglacial aquifers such as these are common globally, and future changes in glacier coverage and pre-

cipitation are likely to increase the significance of groundwater storage within them. The scale of proglacial groundwater flow and storage has important implications for measuring meltwater flux, for predicting future river flows, and for providing strategic water supplies in de-glaciating catchments.

## 1 Introduction

A major challenge in modern hydrology is predicting changes in freshwater flows and storage resulting from glacier retreat in response to climate change (Jiménez Cisneros et al., 2014). Most glaciers worldwide have been in retreat since the mid-19th century, with the loss of global glacier ice accelerating during the 21st century (Zemp et al., 2015). This change has the potential to affect over 1 billion people who live in catchments where glacier melt contributes to river flow (Kundzewicz et al., 2008). Glacial retreat is expected to increase meltwater river flows until the mid-late 21st century (Jiménez Cisneros et al., 2014; Lutz et al., 2014). Longer term, as glacier ice loss continues, meltwater flows will decrease (Jiménez Cisneros et al., 2014). This lessening of the role of glaciers in regulating flows will change the nature of glacier-fed rivers and the importance of other water sources in glacier catchments: rainfall, snowmelt and groundwater. Predicted impacts include changes to the frequency and magnitude of flooding (Jiménez Cisneros et al., 2014); hydroelectric power production (Laghari, 2013); drinking water and irrigation (Kundzewicz et al., 2008);

ecosystem functioning of catchments (Brown et al., 2007); and groundwater recharge (Taylor et al., 2013).

The role of groundwater storage in the hydrology of de-glaciating catchments is recognised, but to date there has been little direct hydrogeological investigation of groundwater–meltwater interactions (Levy et al., 2015; Vincent et al., 2019) with calls for more (Heckmann et al., 2016; Vincent et al., 2019). Indirect studies, inferred from river flow, indicate groundwater in Himalayan glacial catchments may be a significant source of delayed discharge to rivers (Andermann et al., 2012). Modelling of Himalayan catchments suggests that increased glacial melt this century will increase groundwater recharge from glacier runoff and the groundwater baseflow component in river flow (Immerzeel et al., 2013). Glacier meltwater rivers in Alaska can potentially lose half their annual flow to groundwater (Liljedahl et al., 2017). Groundwater can comprise 15 %–75 % of winter river flows in glacial catchments in the European Alps, Canadian Rockies, Peruvian Andes and Iceland (Malard et al., 1999; Hood et al., 2006; Bury et al., 2011; McKenzie et al., 2014; Baraer et al., 2015; MacDonald et al., 2016). Direct experimental studies of groundwater in glacial environments are rare (Vincent et al., 2019): e.g. subglacial groundwater behaviour (Sigurðsson, 1990; Boulton et al., 2001, 2007a, b); groundwater flow in relict rock glaciers (Winkler et al., 2016; Harrington et al., 2018); and the behaviour of shallow (< 3 m) groundwater in glacial outwash plains in Iceland (Robinson et al., 2008, 2009a, b). The latter Icelandic studies demonstrated meltwater recharge to proglacial aquifers and linked retreating glaciers with declining groundwater levels (Levy et al., 2015).

In this study, we directly investigate the 3-D aquifer properties of a proglacial floodplain (referred to here as a *sandur*) of the Virkisjökull glacier in south-eastern Iceland, to 15 m depth, using geophysics, drilling and hydraulic conductivity testing; and provide continuous time series data for groundwater, river stage/flow and precipitation over 3 years, with campaign sampling for stable isotopes. We explore the relationships between groundwater, glacier meltwater flows and precipitation, revealing seasonal and spatial hydrological patterns.


Iceland provides an ideal observatory for studying groundwater in de-glaciating catchments. Ice melt from glaciers, which cover ~ 11 % of Iceland, provides an estimated third of total river runoff (Björnson and Pálsson, 2008), but glacier retreat across Iceland (Sigurðsson et al., 2007) is forecast to produce significant changes in glacial catchment hydrology (Aðalgeirsdóttir et al., 2011). The British Geological Survey (BGS), in collaboration with Veðurstofa Íslands (the Icelandic Meteorological Office), has studied the Virkisjökull catchment since 2009, monitoring rapid glacier retreat (Bradwell et al., 2013), retreat mechanisms (Phillips et al., 2013, 2014), and researching glacial meltwater hydrology (MacDonald et al., 2016; Flett et al., 2017; Mackay et al., 2018).

## Study site

Virkisjökull is an outlet glacier of the Vatnajökull ice cap in south-eastern Iceland (Fig. 1), within the Virkisá River basin, which has a catchment area of ~ 32.5 km<sup>2</sup> to the confluence with the Svinafellsá River (MacDonald et al., 2016). Virkisjökull drains ice steeply south-westwards from an elevation of > 1800 m a.s.l. on the ice cap summit to < 150 m a.s.l. at its terminus, with an average gradient of approximately 0.25. It has a high mass balance gradient, with net annual accumulation of more than 7 m w.e. a<sup>-1</sup> (metres of water equivalent per annum) at the ice cap summit (Guðmundsson, 2000) and net annual ice melt of more than 8 m w.e. a<sup>-1</sup> in the main ablation zone (Flett, 2016). The equilibrium line altitude on Virkisjökull is approximately 1150 m a.s.l. (MacDonald et al., 2016). The glacier has been retreating since 1990 (Hannisdóttir et al., 2015), with a marked acceleration in retreat rates since 2005 (Bradwell et al., 2013), during which time the glacier terminus has retreated by ~ 1 km and there has been extensive surface lowering.

The Virkisá River emerges from a small, shallow proglacial lake that has formed during the recent rapid deglaciation, and flows initially for 1 km over bedrock, flanked by moraines, and then for 4 km across the Virkisjökull sandur to the Svinafellsá River (Fig. 1). The river drains glacial meltwater and virtually all precipitation falling on Virkisjökull glacier, adjacent hillslopes and proglacial moraines. It occupies a single channel across the upper sandur, separating into a number of distinct channels across the lower sandur (Fig. 1). The mean summer river flow over 3 years of continuous monitoring (2011–2014) ranged from 5.3 to 7.9 m<sup>3</sup> s<sup>-1</sup>; and significant river flow occurred in winter (mean 1.6–2.4 m<sup>3</sup> s<sup>-1</sup>). Isotopic studies (MacDonald et al., 2016), validated by numerical modelling (Mackay et al., 2018), demonstrate that summer river flows are governed by glacier ice melt, and that winter flows are a combination of glacier meltwater, local precipitation and groundwater flow. The Virkisjökull sandur falls from 100 to 50 m a.s.l. with a surface gradient of 0.017 (Fig. 1). Over much of the sandur where river channels are actively migrating, there is little vegetation cover and no soil development. In more stable areas thin soils and more developed vegetation cover occur (Fig. 1). The groundwater catchment on the sandur associated with outflow from Virkisjökull has been estimated by using the surface water catchment identified from lidar and dGPS (Fig. 1).

The proglacial area has a maritime climate with cool summers (mean summer air temperature 8–12 °C) and mild winters (1 °C). Air temperature in the Virkisá basin is controlled mainly by altitude, with an average annual lapse rate of  $-5^{\circ}\text{C km}^{-1}$  (Flett, 2016; Mackay et al., 2018). Mean annual precipitation south-west of the Vatnajökull ice cap, including the Virkisjökull sandur, is ~ 1800 mm; precipitation on the eastern side of the ice cap averages 3000 mm a<sup>-1</sup>, and can exceed 7000 mm a<sup>-1</sup> on the ice cap summit (Guðmundsson,



**Figure 1.** Virkisjökull study catchment. **(a)** Study area on Virkisjökull sandur, south-eastern Iceland, encompassing 6 km<sup>2</sup> groundwater catchment originating at a proglacial lake outlet. Hillshade model generated from LiDAR DEM ©Veðurstofa Íslands, 2010. **(b)** Piezometer M1 on the upper sandur near the catchment edge, showing established sandur vegetation and, in the middle distance, the area of moraines. **(c)** Virkísá River on the lower sandur in the summer melt season showing braided channels and an active, unvegetated sandur surface. Piezometers in the sandur are in three transects: Upper (U1–U2), Middle (M1–M3), and Lower (L1–L3).

2000). The proglacial area receives  $\sim 150$  precipitation days per year, estimated from interpretation of 3 years of daily photographs (MacDonald et al., 2016), which also show that snow cover, even in winter, rarely lasts for more than a week before melting. Potential evapotranspiration over the sandur was estimated at  $\sim 450 \text{ mm a}^{-1}$  by Einarsson (1972) and actual evapotranspiration at  $100\text{--}414 \text{ mm a}^{-1}$  by Jónsdóttir (2008).

## 2 Methodology

### 2.1 Aquifer characterisation

Eight boreholes were drilled into the sandur to 9–15 m depth during July and August 2012, in three transects approximately perpendicular to the river along a 3 km longitudinal reach in the upper, middle, and lower study catchment (Fig. 1a). Sediment samples collected during drilling were lithologically logged. The boreholes were installed as piezometers in September 2012, with 88 mm diameter uPVC plain casing to at least 5–12 m depth and a 3–6 m length of 0.5 mm slotted well screen below this (Table S1 in the Supplement). A further two boreholes were drilled into volcanic bedrock, to 5.5 and 13.75 m depth, between the glacier terminus and the upper edge of the sandur (Fig. 1a). Three methods were used to establish the physical aquifer properties of the sandur: (1) infiltration tests to 0.15 m depth at 20 locations, using a Guelph permeameter, and saturated hydraulic conductivity calculated by the Laplace method (Reynolds et al., 1983) (Table S2); (2) particle size analysis on 42 sandur sediment samples to 0.5 m depth, at 22 locations, and hydraulic conductivity estimated using a modified Hazen formula suitable for heterogeneous glacial deposits (MacDonald et al., 2012; Williams et al., 2019) (Table S3); and (3) constant rate pumping tests of between 3.5 and 6 h in each sandur piezometer, at rates of  $0.5\text{--}1.8 \text{ L s}^{-1}$ , and transmissivity estimated by the Jacob time-drawdown and Theis recovery methods corrected for unconfined conditions (Kruseman and de Ridder, 1994).

To measure aquifer thickness and depth to bedrock, two Tromino® passive seismic surveys were undertaken transversely across the Virkisjökull sandur, and a third longitudinally down the Svinafellsandur aquifer 4.5 km to the west, using a single broad-band three-component seismometer with one vertical and two horizontal components. Measurements were recorded for 15 min at 50–100 m lateral intervals and data processed to derive depth to bedrock assuming typical shear wave velocities of  $400\text{--}600 \text{ m s}^{-1}$  for Icelandic glacial sands and gravels (Bessason and Kaynia, 2002; Castellaro et al., 2005). These data were interpreted with a previous seismic reflection survey in the area to infer sediment thickness and potential layering (Guðmundsson et al., 2002).

### 2.2 Groundwater, surface hydrology, and precipitation monitoring

Monitoring of groundwater levels and temperature in sandur piezometers, at 15 min intervals, was undertaken from August 2012 to May 2015 (34 months) using In-Situ Inc. Rugged Troll 100 non-vented pressure transducers at 7–8.4 m depth (Ó Dochartaigh et al., 2019). Two In-Situ Rugged Barometer Trolls measured air pressure and temperature. River stage and discharge data are available for August 2012–May 2015 from an automatic stream gauge at Virkisá bridge (Fig. 1) with two water-level sensors, checked using daily photographs and continuous flow measurements from a radar mounted beneath a bridge (MacDonald et al., 2016). From April 2013 to March 2015 river stage and temperature were additionally monitored continuously every 15 min adjacent to piezometer U1 by an In-Situ Inc. Rugged Troll 100 pressure transducer (Fig. 1). Rainfall data and temperature for the proglacial area were measured at the closest of the three Automatic Weather Stations installed by BGS in the catchment (AWS1; 156 m a.s.l.). These weather stations were not equipped to measure snowfall, but daily photographs enabled periods of snowfall to be estimated. Long-term weather data from the Fagurhólsmýri weather station operated by the Icelandic Meteorological Office (IMO) approximately 12 km south of the study site, and national scale gridded products (Nawri et al., 2017), were used to check the plausibility of weather data measured on site.

Hierarchical cluster analysis of groundwater-level data was carried out on the entire dataset. Data were treated using the Standardized Groundwater level Index (Bloomfield and Marchant, 2013), which indicated the optimal number of clusters is four. Groundwater flow was estimated assuming a mean aquifer width of 1 km, an aquifer thickness at the river gauge from the passive seismic interpretation, an average measured groundwater-level gradient of 0.016, and hydraulic conductivity from the median of all measured values ( $n = 64$ ). Uncertainty was calculated from the interquartile range of measured K and uncertainty in aquifer thickness interpretation. The hydraulic conductivity of the deeper, unmeasured, sandur aquifer layer was estimated using the formula of MacDonald et al. (2012) taking into account a change in sediment state from very loose to loose and firm, which is likely to over-estimate the reduction in pore space due to loading (Schmidt and McDonald, 1979). The total volume of groundwater stored in the aquifer was estimated using a conservative estimate of the average aquifer porosity of 15 % (Parrieux and Nicoud, 1990).

### 2.3 Groundwater isotopic sampling and analysis

Physico-chemical analysis and modelling were based on samples of groundwater from piezometers and springs collected during three summer campaigns in September 2012, 2013, and 2014 and three winter (pre-melt) campaigns in

January 2013, April 2013, and May 2014 (MacDonald et al., 2019). Groundwater sampling from piezometers was carried out after purging by low-flow pumping until stable readings were obtained for field-measured parameters. Field measurements of specific electrical conductance (SEC), temperature and bicarbonate alkalinity by titration pH (Table S4), and dissolved oxygen and redox potential (Eh) were made at the time of sampling. Samples for stable isotopes  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  were collected unfiltered in glass or Nalgene™ polyethylene bottles and analysed at BGS laboratories by isotope ratio measurement on a VG-Micromass Optima mass spectrometer. Data are quoted in permil (‰) with respect to Vienna Standard Mean Ocean Water (VSMOW) (IAEA/WMO, 2019); measurement precision was  $\pm 0.1\text{ ‰}$  for  $\delta^{18}\text{O}$  and  $\pm 1.0\text{ ‰}$  for  $\delta^2\text{H}$ . Local precipitation stable isotope composition and a local meteoric line were estimated from International Atomic Energy Agency station data for Reykjavík (IAEA/WMO, 2019), supported by estimates for southeastern Iceland (Arnason, 1977) and for southern Iceland (Sveinbjörnsdóttir et al., 1995), as described in MacDonald et al. (2016). The isotopic composition of the Virkisá River as it enters the sandur was established by sampling campaigns from September 2011 to December 2014 (MacDonald et al., 2016).

The high topographic gradient of the catchment, with large climatic differences between the upland glacial accumulation area ( $> 1800\text{ m a.s.l.}$ ) and the lowland temperate proglacial area ( $< 150\text{ m a.s.l.}$ ), results in two easily distinguished isotopic compositions: (1) glacier meltwater; and (2) precipitation across the proglacial area. A binary mixing model for  $\delta^2\text{H}$  was applied to investigate the relative contributions of local precipitation and of river water (which is dominated by glacier melt) to sandur groundwater, based on a two-component mixing equation. The end members applied for  $\delta^2\text{H}$  composition were  $-76.1\text{ ‰}$  for river water (Table S4) and  $-58.5\text{ ‰}$  for average annual local precipitation (MacDonald et al., 2016). The fraction of local precipitation in sandur groundwater (FGW) was calculated using the formula  $\text{FGW} = (\delta^2\text{H}_R - \delta^2\text{H}_P)/(\delta^2\text{H}_{GW} - \delta^2\text{H}_R)$ , where  $\delta^2\text{H}_R$  is the composition of river water;  $\delta^2\text{H}_P$  is the composition of local precipitation; and  $\delta^2\text{H}_{GW}$  is the composition of sampled groundwater. Since most river recharge to the aquifer occurs during the summer months when river flow is high and dominated by glacier melt, the impact of the small evolution in stable isotope composition down-river observed in winter due to groundwater baseflow (MacDonald et al., 2016) is insignificant, particularly when compared to the large difference between river flow and local precipitation isotopic composition.

### 3 Results


#### 3.1 Sandur structure and aquifer properties

The groundwater study catchment covers  $6\text{ km}^2$  and encompasses the sandur, adjacent hillslopes and moraines, and river outflow from the proglacial lake (Fig. 1). Geophysical evidence from the passive seismic and previous seismic reflection survey indicates that (Figs. 2, S1) depth to bedrock increases from approximately  $60\text{--}100\text{ m}$  in the upper sandur to  $100\text{--}150\text{ m}$  in the lower sandur. The shallow aquifer material comprises loosely consolidated, moderately to poorly sorted, dominantly medium- to coarse-grained glaciofluvial sand, gravel, and cobbles (Fig. 2). All the sediment is of volcanic origin and has been transported and deposited by the Virkisá River. The deeper deposits are not exposed, but nearby seismic interpretation confirms that the material is generally uniform to  $> 50\text{ m}$ , reflecting the similar sediment derivation and deposition mechanisms (Guðmundsson et al., 2002). Although not directly observed in the seismic data, there is a possibility that at greater depth ( $> 50\text{ m}$ ) there exist more consolidated Pleistocene aged sediments which have been compacted by ice loading during earlier glaciations (Guðmundsson et al., 2002). Observations of bedrock from nearby exposures and two boreholes drilled in bedrock reveal relatively massive and poorly fractured volcanic rock.

The sandur aquifer is highly permeable to at least  $15\text{ m}$  depth, with a median hydraulic conductivity of  $35\text{ m d}^{-1}$  (IQ range  $25\text{--}40\text{ m d}^{-1}$ ) (Fig. 2a, Tables S2, S3). Transmissivity of the upper  $15\text{ m}$  is  $100\text{--}2500\text{ m}^2\text{ d}^{-1}$  with a median value of  $600\text{ m}^2\text{ d}^{-1}$ , consistent with hydraulic conductivity measurements (Table S1). The permeability of the deeper sandur aquifer was not directly measured. However, given the grain size distribution is the same as the shallow aquifer, and assuming the worst case of compaction due to burial and ice loading (Schmidt and McDonald, 1979), median hydraulic conductivity may have reduced to  $15\text{ m d}^{-1}$  or at a worst case  $6\text{ m d}^{-1}$  (MacDonald et al., 2012). By contrast, the underlying bedrock has low transmissivity, less than the lower limit from the experimental methods employed (transmissivity  $< 0.25\text{ m}^2\text{ d}^{-1}$ ). The sandur aquifer is unconfined. Depth to groundwater ranges from 0 to  $4.4\text{ m}$  below ground level and maximum measured seasonal groundwater-level fluctuations are  $1.0\text{--}3.6\text{ m}$ . From  $1\text{ km}$  down-sandur from its upper edge, there is extensive groundwater discharge at the ground surface via perennial and ephemeral springs (Fig. 2). A conservative estimate of the volume of groundwater stored in the full thickness of the aquifer is  $51 \pm 15\text{ million m}^3$ , approximately  $1\text{--}2\%$  of estimated ice volume in the glacier (Mackay et al., 2018).

#### 3.2 Groundwater dynamics

Groundwater-level elevation falls from the upper to lower sandur, with a gradient of  $0.018$  across the upper and  $0.013$



**Figure 2.** Geometry, geology, and hydrogeology of the sandur aquifer. **(a)** Hydraulic conductivity and summer groundwater-level contours. Other legend as Fig. 1. Hillshade model generated from LiDAR DEM © Veðurstofa Íslands, 2010; **(b)** schematic cross section of the hydrogeology, showing the locations of the piezometer transects, spring discharge area, and indicative groundwater flow lines; **(c)** geological section through the river bank showing heterogeneous glaciofluvial deposition; **(d)** perennial groundwater-fed stream on the lower sandur, associated with extensive growth of mosses and other aquatic vegetation; **(e)** groundwater discharge to the otherwise inactive river channel on the lower sandur, flowing to the active channel in the distance.

across the lower sandur (Fig. 2). In the upper sandur, closest to the glacier, groundwater levels adjacent to the glacial meltwater channel are on average 1 m below river stage for most or all of the year (Fig. 3a, b), leading to a strong piezometric gradient away from the river to groundwater. Across the middle sandur, groundwater levels close to the river vary from

0.5 m below to 0.5 m above adjacent river stage, leading to complex river–groundwater interactions. Here, piezometric gradients are generally from river to aquifer in the summer melt season, when river flows are highest, and from aquifer to river in winter, driven by high winter precipitation and associated recharge. From 2 km down-sandur, groundwater levels

are above the adjacent river stage for much of the year, creating a piezometric gradient that drives visible groundwater discharge through seeps and springs to the river (Fig. 2d) and ephemeral and perennial springs (Fig. 2e).

Hierarchical cluster analysis of groundwater-level data indicates two patterns of groundwater-level fluctuation (Fig. 3c): one driven primarily by local precipitation; and the second driven partly by precipitation but also strongly influenced by river stage, especially in summer (Fig. 3d). Groundwater levels showing the first pattern (in piezometers U2, M1, M2, M3, and L3) fluctuate dominantly in response to individual precipitation events and longer-term precipitation patterns. The magnitude of groundwater-level fluctuations typically increases with distance from the river. Rainfall is higher than its long-term average throughout most of the winter and lower in summer, and this is generally reflected in the groundwater-level fluctuations (Fig. 3d). Groundwater levels showing the second pattern (in piezometers U1, L1, and L2) fluctuate in response to river stage as well as local precipitation, at seasonal (Fig. 3d) and also at event timescales (Fig. 3a). River stage is typically higher than its long-term average during peak summer melt, and groundwater levels in this group also remain close to or higher than their long-term average throughout the summer (Fig. 3d). The strongest response to river stage at a seasonal timescale is in piezometer L1, where groundwater levels during the 2013 summer melt season remained consistently higher than throughout the three winters from 2012 to 2014 (Fig. 3d). The strongest response at an event timescale is in piezometer U1, where groundwater levels show consistent diurnal fluctuations during the summer melt season that coincide with diurnal melt-controlled fluctuations in the river stage (Fig. 3a).


Piezometers U1 and U2 (20 and 90 m from the river, respectively) illustrate the relative impacts of summer glacier meltwater flows and large winter precipitation events on groundwater level–river stage gradients (Fig. 3). In summer, low precipitation and large glacier meltwater flows cause groundwater levels in U1 to rise above U2, creating a piezometric gradient away from the river (Fig. 3a). During individual winter rain storms, groundwater levels in U2 rise higher than in U1, creating a piezometric gradient towards the river (Fig. 3b) and driving baseflow to the river further downstream in the middle sandur.

Mean estimated annual groundwater flow through the shallow part of the aquifer calculated using Darcy's equation (20–40 m thick) is  $0.19 \text{ m}^3 \text{ s}^{-1}$  (IQ range  $0.093\text{--}0.30 \text{ m}^3 \text{ s}^{-1}$ ), equivalent to 4.5 % (2.7 %–5.8 %) of mean annual river flow and 9.7 % (5.8 %–12 %) of mean winter river flow. The relatively small seasonal variation in groundwater levels means there is no significant seasonal variation in estimated groundwater flow across the aquifer. Overall groundwater flow through the total depth of the sandur aquifer is estimated as  $0.42 \text{ m}^3 \text{ s}^{-1}$  ( $0.12\text{--}1.1 \text{ m}^3 \text{ s}^{-1}$ ), equivalent to 9.8 % (3.6 %–22 %) of mean annual river flow and 21 % (7.7 %–46 %) of mean winter river flow.

### 3.3 Stable isotopes and temperature

Stable isotope composition ( $\delta^{2}\text{H}$  and  $\delta^{18}\text{O}$ ) in groundwater from piezometers and springs was compared to that of glacier meltwater and local rainfall (Fig. 4, Table S4). Previous studies have demonstrated that glacial meltwater and local rainfall on the proglacial area are easily distinguished using  $\delta^{2}\text{H}$  and  $\delta^{18}\text{O}$  due to the high elevation of the accumulation area (MacDonald et al., 2016). Across individual piezometers, springs, and the river, variability between sampling campaigns was much less than variability between sites (Table S4). In particular, the river samples (taken as the river enters the sandur and therefore largely glacier meltwater) exhibited little seasonal variability  $-76.1 \pm 2.6 \delta^{2}\text{H}$  ( $n = 19$ ). Therefore, mean values from across the campaigns were used for analysis. Groundwater stable isotope compositions vary considerably across the sandur, spanning the range of compositions expected from glacier meltwater and local precipitation (Fig. 4). Piezometers (U1, L1, L2) identified from their hydrographs as most influenced by the river have isotopic compositions similar to river water, while piezometers whose hydrographs are influenced more by precipitation have a much wider range of isotopic composition, with U2 and M3 similar to local rainfall, and M2, M1, and L3 a mixture between local rainfall and river water. The springs showed a wide variety of compositions.

A binary mixing model developed for  $\delta^{2}\text{H}$  indicates the relative proportion of precipitation and glacier meltwater in groundwater (Fig. 4b, c) and demonstrates a clear relationship with distance from the meltwater river. Within a zone extending up to 50 m from the river in the upper sandur, 130 m in the central sandur, and 500 m in the lower sandur, groundwater in piezometers generally comprises more than 50 % glacier meltwater. Shallower groundwater from springs within this river zone is more influenced by local precipitation, but still comprises more than 25 % glacier meltwater. Beyond this zone, groundwater from both piezometers and springs consistently comprises less than 25 % river water (Fig. 4c). Since the binary mixing model uses glacier meltwater as its endpoint, it is likely to be conservative in the proportion of river–groundwater interactions as it does not account for evolution of the river water stable isotope composition downstream due to groundwater baseflow. Selected hydrochemical tracers and water temperature also help distinguish these two zones (Table S4). Specific electrical conductance (SEC) and bicarbonate ( $\text{HCO}_3^-$ ) are significantly lower in those piezometers strongly influenced by the river than those where precipitation influence is dominant (Fig. S1). River water temperature is relatively constant year-round at an average of  $1.7^\circ\text{C}$ , and mean annual groundwater temperature is lowest in piezometers close to the river and highest in those furthest from the river (Table S4).




**Figure 3.** Groundwater levels, river stage, and precipitation. **(a)** Groundwater levels in the upper sandur during a 14 d dry period in summer (for the legend, see panel **d**). **(b)** Groundwater levels in the upper sandur during a 14 d rainy period in winter (for the legend, see panel **d**). Piezometer U1 (solid) is 20 m from the river; piezometer U2 (dashed) is 90 m from the river. **(c)** Dendrogram obtained by hierarchical cluster analysis of groundwater-level data from sandur piezometers (piezometer locations in Fig. 1). The highest-level break shows two clusters representing piezometers where groundwater is influenced dominantly by local precipitation (U2, M1, M2, M3, L3) and piezometers where groundwater is influenced by the meltwater river (U1, L1, L2). Boxes show the four optimal sub-clusters indicated by data standardisation. **(d)** Monthly running mean of river stage (m), hourly precipitation (mm), and groundwater level (m), as a variation from the long-term average (LTA = 0).

## 4 Discussion

This study in Iceland shows that proglacial floodplains can form thick, highly permeable aquifers. By directly quantifying aquifer parameters and groundwater–glacier meltwater interaction, we have provided evidence of the significance of groundwater in proglacial hydrology. This has important implications for measuring glacial meltwater flux, for predicting future river flows and ecological impacts, and for water supplies in de-glaciating catchments. Similar thick proglacial glaciofluvial aquifers with high permeability and storage occur in other active glacial environments, e.g. elsewhere in Iceland (Robinson et al., 2008), the European Alps (Parrié and Nicoud, 1990), and the Peruvian Andes (McKenzie et al., 2014), and with rapid deglaciation occurring globally proglacial aquifers are developing in many other locations, increasing the importance of characterising groundwater (Vincent et al., 2019).

### 4.1 Groundwater flow

Our study shows that significant water can flow through a glacierised catchment as groundwater, despite groundwater representing only a small proportion of the volume of water stored in glacial ice in the catchment. Reliable measurements of glacier meltwater are important for calibrating cryospheric-hydrological models (Bliss et al., 2014; Lutz et al., 2014; Mackay et al., 2018). The estimated volume of groundwater flow through the shallowest 20–40 m of the Virkisjökull proglacial aquifer is significant,  $0.19 \text{ m}^3 \text{ s}^{-1}$ , equivalent to approximately 4.5 % of mean annual river flow or 9.7 % of mean winter river flow, with estimates of 9.8 % and 21 %, respectively, if flow through the full thickness of the aquifer is considered. Other studies in Iceland have proposed that a similarly large proportion of meltwater ( $0.5\text{--}1 \text{ m w.e. a}^{-1}$ ) can flow through the groundwater system, either from sub-glacial or proglacial recharge (Sigurdsson, 1990; Hemmings et al., 2016); meltwater river losses to groundwater of up to 50 % have also been reported (Liljedahl



**Figure 4.** Stable isotope composition of waters and results of the binary mixing model of  $\delta^2\text{H}$  in groundwater. (a) Stable isotope composition of sandur groundwater in piezometers and springs. Individual piezometers labelled; piezometer locations in Fig. 1. Also shown are ranges in stable isotope composition of precipitation and river water (MacDonald et al., 2016). Plotted on the Local Meteoric Water Line (LMWL) calculated for Reykjavík. (b) Plot of the mean proportion of groundwater recharged from the river using binary mixing of the model of  $\delta^2\text{H}$  by perpendicular distance from the river and down-sandur. (c) Map of the mean proportion of groundwater estimated to be recharged from the river using a binary mixing model for  $\delta^2\text{H}$ . Hillshade model generated from LiDAR DEM © Veðurstofa Íslands, 2010.

et al., 2017). Measuring river flow in catchments with active glaciers is notoriously difficult, given the harsh conditions, actively changing river beds, and wide ranges in flows and sediment load. Measurements are therefore subject to high uncertainty. Here, we demonstrate that groundwater adds another source of uncertainty. Measurements of river flows that rely solely on river stage in the proglacial area are likely to underestimate total annual meltwater flows, with much higher relative errors at low flows. Similar potential underestimation in glacier melt estimations due to groundwater flow have recently been reported in South America (Saberi et al., 2019).

#### 4.2 Meltwater–groundwater interaction

Groundwater–glacier meltwater interactions are controlled by relative differences in water levels between the river and the proglacial aquifer and vary both spatially, down the catchment, and seasonally. There is year-round active recharge of

river water to the aquifer in the upper catchment, complex interaction in the middle of the sandur, and extensive groundwater baseflow to the river and springs across the lower catchment. Distinct patterns of groundwater–glacier meltwater dynamics are observed in groundwater-level fluctuations and in groundwater stable isotopic composition, temperature, and chemistry. In a zone extending up to 50–500 m from the river, the influence of the river on groundwater overshadows that of local precipitation. Here, recharge of glacier meltwater to the aquifer from river losses has a significant impact on the physical, chemical, and stable isotopic characteristics of groundwater in the proglacial aquifer. The aquifer provides additional water storage and groundwater discharges back to the river further downstream through a large number of springs and seeps (Figs. 1 and 4). This is consistent with other studies in glacier-dominated catchments, which inferred groundwater baseflow to rivers of 15 %–75 % (Malard et al., 1999; Hood et al., 2006; Bury et al., 2011; McKenzie et al., 2014; MacDonald et al., 2016).

However, away from the river the aquifer is recharged dominantly from local precipitation. Active precipitation recharge to the aquifer is evident from groundwater stable isotopic composition and groundwater-level response to precipitation and reflects high annual precipitation (rainfall and snow), high aquifer permeability, and low evapotranspiration linked to limited soil development and vegetation cover. Recharge is likely to occur not only from direct precipitation on the sandur surface, but also from ephemeral streams draining from hillslopes and groundwater seepage from surrounding moraines. Groundwater discharge via springs and baseflow in the lower catchment supports surface water flows and local ecosystems and comprises groundwater derived mainly from local precipitation (Fig. 4).

Looking forward, as the glacier continues to melt, the proglacial aquifer will continue to have a buffering effect on river flow. High river flows will recharge the aquifer, whether caused by glacier ice melt, snowmelt or winter storms, as occurs in relic glacial outwash aquifers now in now-temperate areas (e.g. MacDonald et al., 2014), and will sustain springs, baseflow, and surface ecosystems further down the catchment. Local precipitation falling on the aquifer is likely to continue to be a major source of aquifer recharge and contribution to river baseflow in addition to the groundwater discharging from other glacial deposits emerging within the landscape (MacDonald et al., 2016). In upland areas in northern Europe where glaciofluvial deposits from past glaciations are present, detailed studies have demonstrated that groundwater often comprises more than 50 % of flow to river headwaters (Soulsby et al., 2005; Blumstock et al., 2015; Scheliga et al., 2017). Therefore, as glaciers continue to melt, groundwater baseflow is likely to become an increasingly important proportion of river flow in de-glaciating catchments.

#### 4.3 Proglacial aquifers as strategic water resources

This study has demonstrated that the Virkisjökull sandur is a highly productive aquifer with regular recharge. Similar thick proglacial glaciofluvial aquifers occur throughout the world and are increasing in extent as glaciers recede, and are likely to also have the potential to sustain high-quality reliable water supplies. In formerly glaciated areas, these aquifers are often targeted for public water supply (e.g. Ó Dochartaigh et al., 2015) because of their ability to sustain high-yielding boreholes, their connectivity with rivers that provides additional recharge, and the generally high chemical quality of the groundwater compared to surface water. If projected glacier losses and increased precipitation in glacierised catchments are realised (Jiménez Cisneros et al., 2014), proglacial aquifers, recharged by local precipitation, represent a potentially significant store of high-quality water in regions around the world that currently rely on glacier melt for water supply.

#### 5 Conclusions

Three years of investigations of groundwater and glacier meltwater at Virkisjökull, south-eastern Iceland, have enabled the aquifer parameters of the proglacial floodplain to be reliably characterised and seasonal groundwater–glacier meltwater dynamics to be quantified. The key findings from the research are the following.

1. Direct measurements of aquifer characteristics show consistently high permeability ( $35 \text{ m d}^{-1}$ ,  $n = 64$ , IQR  $25\text{--}40 \text{ m d}^{-1}$ ) and volume of groundwater storage ( $50 \pm 15$  million  $\text{m}^3$ ). The proglacial floodplain therefore forms a highly productive aquifer.
2. Significant water flows as groundwater through the shallowest  $20\text{--}40 \text{ m}$  of the proglacial floodplain ( $0.19 \text{ m}^3 \text{ s}^{-1}$ ; IQ range  $0.09\text{--}0.30 \text{ m}^3 \text{ s}^{-1}$ ), equivalent to 4.5 % of mean annual meltwater flow and 9.7 % of mean winter flow. If the full thickness of the aquifer is considered, then groundwater flows of  $0.42 \text{ m}^3 \text{ s}^{-1}$  (IQ range  $0.12\text{--}1.1 \text{ m}^3 \text{ s}^{-1}$ ) – equivalent to 9.8 % (3.6 %–22 %) of mean annual river flow and 21 % (7.7 %–46 %) of mean winter river flow – are possible.
3. Groundwater is recharged both from the glacial meltwater river and local precipitation falling on the aquifer or draining from nearby hillslopes. Glacier meltwater is particularly important in a zone from 50 to 500 m from the river, where glacier meltwater comprises > 25 % and often > 50 % of recharge.
4. There are complex but consistent river–groundwater interactions: in the upper sandur, closest to the glacier, the river loses to groundwater much of the year; in the middle sandur the river loses to the groundwater in the summer melt and gains from groundwater in the winter low flows; in the lower sandur groundwater provides baseflow to the river through springs and baseflow seeps.

Proglacial aquifers are common worldwide and increasing in extent with deglaciation. These findings, therefore, have wider implications for measuring glacier meltwater flux, for predicting future river flows, and for water supplies in de-glaciating catchments. Effectively understanding and characterising groundwater flows and storage in catchments with glaciers, and incorporating this in hydrological models, will strengthen our ability to predict and manage the hydrological and environmental impacts of accelerating glacier retreat.

*Data availability.* Water chemistry and groundwater-level data are available freely from the National Geoscience Data Centre (<https://doi.org/10.5285/3c28c1e9-d19d-431c-8c30-e9a014447d7b>, Ó Dochartaigh et al., 2019, and <https://doi.org/10.5285/14da9c02-c5ec-4019-8e5c-06c744d8be9d>, MacDonald et al., 2019). River stage and precipitation are available from the authors on request.

*Supplement.* The supplement related to this article is available online at: <https://doi.org/10.5194/hess-23-4527-2019-supplement>.

*Author contributions.* BEOD managed the field sampling campaign and installation of piezometers and wrote early drafts of the manuscript. AMM oversaw the research and analysis and wrote the final draft of the paper. PW, MR, BEOD, JE, ARB, AMM, and LJ undertook fieldwork and analysis of individual components of the research and WGD provided interpretation of the stable isotopes. All commented on final draft of the manuscript.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This research is published with the permission of the Executive Director of the British Geological Survey (NERC). We thank Vatnajökuls Þjóðgarður for permission to install monitoring equipment; Vatnsborun ehf for borehole drilling and installations; Icelandair for assistance with equipment transport; and Veðurstofa Íslands and the people of Svinafell for research support. Tim Heaton at the NERC Stable Isotope Facility undertook stable isotope analysis; James Sorensen carried out cluster analysis; and Crai Woodward at BGS assisted with diagrams (all BGS).

*Financial support.* This research has been supported by the BGS-NERC Earth Hazards and Observatories Directorate.

*Review statement.* This paper was edited by Wouter Buytaert and reviewed by Aude Vincent and one anonymous referee.

## References

- Aðalgeirs Óttir, G., Guðmundsson, S., Björnsson, H., Pálsson, F., Jóhannesson, T., Hannesdóttir, H., Sigurðsson, S. P., and Berthier, E.: Modelling the 20th and 21st century evolution of Hoffellsjökull glacier, SE-Vatnajökull, Iceland, *The Cryosphere*, 5, 961–975, <https://doi.org/10.5194/tc-5-961-2011>, 2011.
- Andermann, C., Longuevergne, L., Bonnet, S., Crave, A., Davy, P., and Gloaguen, R.: Impact of transient groundwater storage on the discharge of Himalayan rivers, *Nat. Geosci.* 5, 127–132, <https://doi.org/10.1038/NGEO1356>, 2012.
- Arnason, B.: Hydrothermal systems in Iceland traced by deuterium, *Geothermics*, 5, 125–151, [https://doi.org/10.1016/0375-6505\(77\)90015-3](https://doi.org/10.1016/0375-6505(77)90015-3), 1977.
- Baraer, M., Mckenzie, J., Mark, B. G., Gordon, R., Bury, J., Condom, T., Gomez, J., Knox, S., and Fortner, S. K.: Contribution of groundwater to the outflow from ungauged glaciated catchments: A multi-site study in the tropical Cordillera Blanca, Peru, *Hydrol. Process.*, 29, 2561–2581, <https://doi.org/10.1002/hyp.10386>, 2015.
- Bessason, B. and Kaynia, A. M.: Site amplification in lava rock on soft sediments, *Soil Dyn. Earthq. Eng.*, 22, 525–540, [https://doi.org/10.1016/S0267-7261\(02\)00035-0](https://doi.org/10.1016/S0267-7261(02)00035-0), 2002.
- Björnsson, H. and Pálsson, F.: Icelandic glaciers, *Jökull*, 58, 365–386, 2008.
- Bliss, A., Hock, R., and Radić, V.: Global response of glacier runoff to twenty-first century climate change, *J. Geophys. Res.-Earth*, 119, 717–730, <https://doi.org/10.1002/2013JF002931>, 2014.
- Bloomfield, J. P. and Marchant, B. P.: Analysis of groundwater drought building on the standardised precipitation index approach, *Hydrol. Earth Syst. Sci.*, 17, 4769–4787, <https://doi.org/10.5194/hess-17-4769-2013>, 2013.
- Blumstock, M., Tetzlaff, D., Malcolm, I. A., Nuetzmann, G., and Soulsby, C.: Baseflow dynamics: Multi-tracer surveys to assess variable groundwater contributions to montane streams under low flows, *J. Hydrol.*, 527, 1021–1033, <https://doi.org/10.1016/j.jhydrol.2015.05.019>, 2015.
- Boulton, G. S., Dobbie, K. E., and Zatsepин, S.: Sediment deformation beneath glaciers and its coupling to the subglacial hydraulic system, *Quatern. Int.*, 86, 3–28, [https://doi.org/10.1016/S1040-6182\(01\)00048-9](https://doi.org/10.1016/S1040-6182(01)00048-9), 2001.
- Boulton, G. S., Lunn, R., Vidstrand, P., and Zatsepин, S.: Subglacial drainage by groundwater-channel coupling, and the origin of esker systems: Part 1 – glaciological observations, *Quaternary Sci. Rev.*, 26, 1067–1090, <https://doi.org/10.1016/j.quascirev.2007.01.007>, 2007a.
- Boulton, G. S., Lunn, R., Vidstrand, P., and Zatsepин, S.: Subglacial drainage by groundwater-channel coupling, and the origin of esker systems: Part II – theory and simulation of a modern system, *Quaternary Sci. Rev.*, 26, 1091–1105, <https://doi.org/10.1016/j.quascirev.2007.01.006>, 2007b.
- Bradwell, T., Sigurðsson, O., and Everest, J.: Recent, very rapid retreat of a temperate glacier in SE Iceland, *Boreas*, 42, 959–973, <https://doi.org/10.1111/bor.12014>, 2013.
- Brown, L. E., Milner, A. M., and Hannah, D. M.: Groundwater influence on alpine stream ecosystems, *Freshwater Biol.*, 52, 878–890, <https://doi.org/10.1111/j.1365-2427.2007.01739.x>, 2007.
- Bury, J. T., Mark, B. G., McKenzie, J. M., French, A., Baraer, M., In Huh, K., Luyo, M. A. Z., and López, R. J. G.: Glacier recession and human vulnerability in the Yanamarey watershed of the Cordillera Blanca, Peru, *Climatic Change*, 105, 179–206, <https://doi.org/10.1007/s10584-010-9870-1>, 2011.
- Castellar, S., Mulargia, F., and Bianconi, L.: Passive Seismic Stratigraphy: a new efficient, fast and economic technique, *Geologia Tecnica e Ambientale*, 3, 76–102, 2005.
- Einarsson, M. Á.: Evaporation and potential evapotranspiration in Iceland, *Vedurstofa Íslands/The Icelandic Meteorological Office*, Reykjavík, Iceland, 1972.
- Flett, V., Maurice, L., Finlayson, A., Black, A. R., and MacDonald, A. M.: Meltwater flow through a rapidly deglaciating glacier and foreland catchment system: Virkisjökull, SE Iceland, *Hydrol. Res.*, 48, 1666–1681, <https://doi.org/10.2166/nh.2017.205>, 2017.
- Flett, V. T.: Glacier retreat and projected river regime changes in the hydrologically highly-coupled Virkisjökull catchment, SE Iceland, PhD, University of Dundee, Dundee, UK, 2016.
- Guðmundsson, M. T.: Mass balance and precipitation on the summit plateau of Öræfajökull, SE-Iceland, *Jökull*, 48, 49–54, 2000.

- Guðmundsson, M. T., Bonnel, A., and Gunnarsson, K.: Seismic soundings of sediment thickness on Skeiðarársandur, SE Iceland, *Jökull*, 51, 53–64, 2002.
- Hannesdóttir, H., Björnsson, H., Pálsson, F., Aðalgeirsþóttir, G., and Guðmundsson, S.: Changes in the southeast Vatnajökull ice cap, Iceland, between ~1890 and 2010, *The Cryosphere*, 9, 565–585, <https://doi.org/10.5194/tc-9-565-2015>, 2015.
- Harrington, J. S., Mozil, A., Hayashi, M., and Bentley, L. R.: Groundwater flow and storage processes in an inactive rock glacier, *Hydrolog. Process.*, 32, 3070–3088, <https://doi.org/10.1002/hyp.13248>, 2018.
- Heckmann, T., McColl, S., and Morche, D.: Retreating ice: research in pro-glacial areas matters, *Earth Surf. Proc. Land.*, 41, 271–276, <https://doi.org/10.1002/esp.3858>, 2016.
- Hemmins, B., Whitaker, F., Gottsmann, J., and Hawes, M. C.: Non-eruptive ice melt driven by internal heat at glaciated stratovolcanoes, *J. Volcanol. Geoth. Res.*, 327, 385–397, <https://doi.org/10.1016/j.jvolgeores.2016.09.004>, 2016.
- Hood, J. L., Roy, J. W., and Hayashi, M.: Importance of groundwater in the water balance of an alpine headwater lake, *Geophys. Res. Lett.*, 33, L13405, <https://doi.org/10.1029/2006GL026611>, 2006.
- IAEA/WMO: Global Network of Isotopes in Precipitation (GNIP) Water Isotope System for Data Analysis, Visualization, and Electronic Retrieval (WISER). International Atomic Energy Authority/World Meteorological Organisation, available at: [http://www-naeb.iaea.org/napc/ih/IHS\\_resources\\_gnip.html](http://www-naeb.iaea.org/napc/ih/IHS_resources_gnip.html), last access: 14 April 2019.
- Immerzeel, W. W., Pellicciotti, F., and Bierkens, M. F. P.: Rising river flows throughout the twenty-first century in two Himalayan glacierized watersheds, *Nat. Geosci.*, 6, 742–745, <https://doi.org/10.1038/ngeo1896>, 2013.
- Jiménez Cisneros, B. E., Oki, T., Arnell, N. W., Benito, G., Cogley, J. G., Döll, P., Jiang, T., and Mwakalila, S. S.: Freshwater resources, in: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Field, C. B., Barros, V. R., Dokken, D. J., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., Genova, R. C., Girma, B., Kissel, E. S., Levy, A. N., MacCracken, S., Mastrandrea, P. R., and White, L. L., Cambridge University Press, Cambridge, UK, 229–269, 2014.
- Jónsdóttir, J. F.: A runoff map based on numerically simulated precipitation and a projection of future runoff in Iceland, *Hydrolog. Sci. J.*, 53, 100–111, <https://doi.org/10.1623/hysj.53.1.100>, 2008.
- Kruseman, G. P. and de Ridder, N. A.: Analysis and Evaluation of Pumping Test Data (2nd ed.), Publication 47, Intern. Inst. for Land Reclamation and Improvement, Wageningen, the Netherlands, 370 pp., 1994.
- Kundzewicz, Z. W., Mata, L. J., Arnell, N. W., Döll, P., Jiminez, B., Miller, K., Oki, T., Sen, Z., and Shiklomanov, I.: The implications of projected climate change for freshwater resources and their management, *Hydrolog. Sci. J.*, 53, 3–10, <https://doi.org/10.1623/hysj.53.1.3>, 2008.
- Laghari, J.: Climate change: Melting glaciers bring energy uncertainty, *Nature*, 502, 617–618, <https://doi.org/10.1038/502617a>, 2013.
- Levy, A., Robinson, Z., Krause, S., Waller, R., and Weatherill, J.: Long-term variability of proglacial groundwater-fed hydrological systems in an area of glacier retreat, Skeiðarársandur, Iceland, *Earth Surf. Proc. Land.*, 40, 981–994, <https://doi.org/10.1002/esp.3696>, 2015.
- Liljedahl, A. K., Gadeke, A., O’Neil, S., Gatesman, T. A., and Douglas, T. A.: Glacierized headwater streams as aquifer recharge corridors, subarctic Alaska, *Geophys. Res. Lett.*, 44, 6876–6885, <https://doi.org/10.1002/2017GL073834>, 2017.
- Lutz, A. F., Immerzeel, W. W., Shrestha, A. B., and Bierkens, M. F. P.: Consistent increase in High Asia’s runoff due to increasing glacier melt and precipitation, *Nat. Clim. Change.*, 4, 587–592, <https://doi.org/10.1038/NCLIMATE2237>, 2014.
- MacDonald, A. M., Maurice, L., Dobbs, M. R., Reeves, H. J., and Auton, C. A.: Relating in situ hydraulic conductivity, particle size and relative density of superficial deposits in a heterogeneous catchment, *J. Hydrol.*, 434–435, 130–141, <https://doi.org/10.1016/j.jhydrol.2012.01.018>, 2012.
- MacDonald, A. M., Lapworth, D. J., Hughes, A. G., Auton, C. A., Maurice, L., Finlayson, A., and Goody, D. C.: Groundwater, flooding and hydrological functioning in the Findhorn floodplain, Scotland, *Hydrolog. Res.*, 45, 755–773, <https://doi.org/10.2166/nh.2014.185>, 2014.
- MacDonald, A. M., Black, A. R., Ó Dochartaigh, B. É., Everest, J., Darling, W. G., Flett, V., and Peach, D. W.: Using stable isotopes and continuous meltwater river monitoring to investigate the hydrology of a rapidly retreating Icelandic outlet glacier, *Ann. Glaciol.*, 57, 151–158, <https://doi.org/10.1017/aog.2016.22>, 2016.
- MacDonald, A. M., Ó Dochartaigh, B. É., and Fallas, H. C.: Water chemistry and stable isotope data, Virkisjökull Glacier Observatory, 2011–2018, British Geological Survey, Dataset, <https://doi.org/10.5285/14da9c02-c5ec-4019-8e5c-06c744d8be9d>, 2019.
- Mackay, J. D., Barrand, N. E., Hannah, D. M., Krause, S., Jackson, C. R., Everest, J., and Aðalgeirsþóttir, G.: Glacio-hydrological melt and run-off modelling: application of a limits of acceptability framework for model comparison and selection, *The Cryosphere*, 12, 2175–2210, <https://doi.org/10.5194/tc-12-2175-2018>, 2018.
- Malard, F., Tockner, K., and Ward, J. V.: Shifting Dominance of Subcatchment Water Sources and Flow Paths in a Glacial Floodplain, Val Roseg, Switzerland, *Arct. Antarct. Alp. Res.*, 31, 135–150, <https://doi.org/10.2307/1552602> 1999.
- McKenzie, J. M., Gordon, R. P., Baraer, M., Lautz, L. K., Mark, B. G., Chavez, D., and Aubry-Wake, C.: Hydrogeology in glaciated high-elevation Andean watersheds – results from the Cordillera Blanca, Peru, Conference Paper 102-13, Geological Society of America Annual Meeting, 19–22 October 2014, Vancouver, British Columbia, Canada, 2014.
- Nawri, N., Pálmasen, B., Petersen, G. N., Björnsson, H., and Þorsteinsson, S.: The ICRA atmospheric reanalysis project for Iceland, Tech. rep., Icelandic Meteorological Office, Reykjavík, Iceland, 2017.
- Ó Dochartaigh, B. É., MacDonald, A. M., Fitzsimons, V., and Ward, R.: Scotland’s aquifers and groundwater bodies. British Geological Survey Open Report OR/15/028, 63 pp., available at: <http://nora.nerc.ac.uk/id/eprint/511413/> (last access: April 2019), 2015.

- Ó Dochartaigh, B. É., MacDonald, A. M., Wilson, P., and Everest, J.: Groundwater Monitoring Data, Virkisjökull Glacier Observatory, 2012–2018, British Geological Survey, Dataset, <https://doi.org/10.5285/3c28c1e9-d19d-431c-8c30-e9a014447d7b>, 2019.
- Parrioux, A. and Nicoud, G. F.: Hydrological behaviour of glacial deposits in mountainous areas, IAHS Publication, 190, 291–312, 1990.
- Phillips, E., Finlayson, A., and Jones, L.: Fracturing, block faulting, and moulin development associated with progressive collapse and retreat of a maritime glacier: Falljökull, SE Iceland, *J. Geophys. Res.–Earth*, 118, 1–17, <https://doi.org/10.1002/jgrf.20116>, 2013.
- Phillips, E., Finlayson, A., Bradwell, T., Everest, J., and Jones, L.: Structural evolution triggers a dynamic reduction in active glacier length during rapid retreat: evidence from Falljökull, SE Iceland, *J. Geophys. Res.–Earth*, 119, 2194–2208, <https://doi.org/10.1002/2014JF003165>, 2014.
- Reynolds, W. D., Elrick, D. E., and Topp, G. C.: A re-examination of the constant head well permeameter method for measuring saturated hydraulic conductivity above the water table, *Soil Sci.*, 136, 250–268, 1983.
- Robinson, Z. P., Fairchild, I. F., and Russell, A. J.: Hydrogeological implications of glacial landscape evolution at Skeiðarársandur, SE Iceland, *Geomorphology*, 97, 218–236, <https://doi.org/10.1016/j.geomorph.2007.02.044>, 2008.
- Robinson, Z. P., Fairchild, I. F., and Arrowsmith, C.: Stable isotope tracers of shallow groundwater recharge dynamics and mixing within an Icelandic sandur, Skeiðarársandur, IAHS Publication, 326, 119–125, 2009a.
- Robinson, Z. P., Fairchild, I., and Spiro, B.: The sulphur isotope and hydrochemical characteristics of Skeiðarársandur, Iceland: identification of solute sources and implications for weathering processes, *Hydrol. Process.*, 23, 2212–2224, <https://doi.org/10.1002/hyp.7368>, 2009b.
- Saberi, L., McLaughlin, R. T., Ng, G.-H. C., La Frenierre, J., Wickert, A. D., Baraer, M., Zhi, W., Li, L., and Mark, B. G.: Multi-scale temporal variability in meltwater contributions in a tropical glacierized watershed, *Hydrol. Earth Syst. Sci.*, 23, 405–425, <https://doi.org/10.5194/hess-23-405-2019>, 2019.
- Scheliga, B., Tetzlaff, D., Nuetzmann, G., and Soulsby, C.: Groundwater isoscapes in a montane headwater catchment show dominance of well-mixed storage, *Hydrol. Proc.*, 31, 3504–3519, <https://doi.org/10.1002/hyp.11271>, 2017.
- Schmidt, V. and McDonald, D. A.: The role of secondary porosity in the course of sandstone diagenesis, in: Aspects of diagenesis, edited by: Scholle, P. A., and Schlüter, P. R., Society of Economic Paleontologists and Mineralogists Special Publication, 26, 175–207, 1979.
- Sigurðsson, F.: Groundwater from glacial areas in Iceland, *Jökull*, 40, 119–145, 1990.
- Sigurðsson, O., Jonsson, T., and Jóhannesson, T.: Relation between glacier-termini variations and summer temperatures in Iceland since 1930, *Ann. Glaciol.*, 42, 395–401, <https://doi.org/10.3189/172756407782871611>, 2007.
- Soulsby, C., Malcolm, I. A., Youngson, A. F., Tetzlaff, D., Gibbins, C. N., and Hannah, D. M.: Groundwater–surface water interactions in upland Scottish rivers: hydrological, hydrochemical and ecological implications, *Scot. J. Geol.*, 41, 39–49, <https://doi.org/10.1144/sjg41010039>, 2005.
- Sveinbjörnsdóttir, Á. E., Johnsen, S. J., and Arnórsson, S.: The use of stable isotopes of oxygen and hydrogen in geothermal studies in Iceland. Proceedings of the World Geothermal Congress, 18–31 May 1995, Florence, Italy, 1043–1048, 1995.
- Taylor, R. G., Scanlon, B., Döll, P., Rodell, M., van Beek, R., Wada, Y., Longuevergne, L., Leblanc, M., Famiglietti, J. S., Edmunds, M., Konikow, L., Green, T. R., Chen, J., Taniguchi, M., Bierkens, M. F. P., MacDonald, A., Fan, Y., Maxwell, R. M., Yechieli, Y., Gurdak, J. J., Allen, D. M., Shamsuddoha, M., Hiscock, K., Yeh, P. J.-F., Holman, I., and Treidel, H.: Ground water and climate change, *Nat. Clim. Change*, 3, 322–329, <https://doi.org/10.1038/nclimate1744>, 2013.
- Vincent, A., Violette, S., and Aðalgeirsdóttir, G.: Groundwater in catchments headed by temperate glaciers: A review, *Earth-Sci. Rev.*, 188, 59–76, <https://doi.org/10.1016/j.earscirev.2018.10.017>, 2019.
- Williams, J. D. O., Dobbs, M. R., Kingdon, A., Lark, R. M., Williamson, J. P., MacDonald, A. M., and Ó Dochartaigh, B. É.: Stochastic modelling of hydraulic conductivity derived from geotechnical data: an example applied to central Glasgow, *Earth Environ. Sci. Trans. R. Soc. Edinb.*, 108, 141–154, <https://doi.org/10.1017/S1755691018000312>, 2019.
- Winkler, G., Wagner, T., Pauritsch, M., Birk, S., Kellerer-Pirklbauer, A., Benischke, R., Leis, B., Morawetz, R., Schreilechner, G., and Hergarten, S.: Identification and assessment of groundwater flow and storage components of the relict Schöneben Rock Glacier, Niedere Tauern Range, Eastern Alps (Austria), *Hydrogeol. J.*, 24, 937–953, <https://doi.org/10.1007/s10040-015-1348-9>, 2016.
- Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S. U., Hoelzle, M., Paul, F., Haeberli, W., Denzinger, F., Ahlström, A. P., Anderson, B., Bajracharya, S., Baroni, C., Braun, L. N., Cáceres, B. E., Casassa, G., Cobos, G., Dávila, L. R., Delgado Granados, H., Demuth, M. N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Hagen, J. O., Holmlund, P., Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnina, V. V., Portocarrero, C. A., Prinz, R., Sangewar, C. V., Severskiy, I., Sigurðsson, O., Soruco, A., Usualiev, R., and Vincent, C.: Historically unprecedented global glacier decline in the early 21st century, *J. Glaciol.*, 61, 745–762, <https://doi.org/10.3189/2015JoG15J017>, 2015.