

Supplement of Hydrol. Earth Syst. Sci., 24, 4777–4791, 2020
<https://doi.org/10.5194/hess-24-4777-2020-supplement>
© Author(s) 2020. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Socio-hydrological data assimilation: analyzing human–flood interactions by model–data integration

Yohei Sawada and Risa Hanazaki

Correspondence to: Yohei Sawada (yohei.sawada@sogo.t.u-tokyo.ac.jp)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Contents of this file

Figures S1 to S10

Introduction

This supporting information provides the figures which support the findings shown in the main article.




Figure S1. Timeseries of (a) high water level $W(t)$, (b) the flood protection level (or levee height) $H(t)$, (c) the distance of the center of mass of the human settlement from the river $D(t)$, (d) the size of the human settlement $G(t)$, (e) the intensity of flooding events $F(t)$, and (f) the social awareness of the flood risk $M(t)$ simulated by the data assimilation experiment in which the observations of F , G , D , H , and M are assimilated into the model every 50 years with 5000 ensembles in the experiment 1 (see section 3.1). The time step is annual. Grey, red, and black lines are the ensemble members, their mean, and the synthetic truth, respectively.




Figure S2. The ratio of RMSEs of the no data assimilation experiment (NoDA) to those of the data assimilation experiments in which the observation error is set to 10%, 15%, 20%, and 25% of the synthetic true values in the experiment 1 (see section 3.1.1). Blue, orange, gray, and yellow bars are RMSEs of the size of the human settlement $G(t)$, the center of mass of the human settlement from the river $D(t)$, the flood protection level (or levee height) $H(t)$, and the social awareness of the flood risk $M(t)$.




Figure S3. The ratio of RMSEs of the no data assimilation experiment (NoDA) to those of the data assimilation experiments in which all of observations (F, G, D, H, and M) are assimilated with 20, 50, 100, 250, 1000, and 5000 ensembles in the experiment 1 (see section 3.1.1). Blue, orange, gray, and yellow bars are RMSEs of the size of the human settlement $G(t)$, the center of mass of the human settlement from the river $D(t)$, the flood protection level (or levee height) $H(t)$, and the social awareness of the flood risk $M(t)$.




Figure S4. Timeseries of (a) high water level $W(t)$, (b) the flood protection level (or levee height) $H(t)$, (c) the distance of the center of mass of the human settlement from the river $D(t)$, (d) the size of the human settlement $G(t)$, (e) the intensity of flooding events $F(t)$, and (f) the social awareness of the flood risk $M(t)$ simulated by the data assimilation experiment in which the observations of F , G , D , H , and M are assimilated into the model every 50 years with 5000 ensembles in the experiment 2 (see section 3.1.2). The time step is annual. Grey, red, and black lines are the ensemble members, their mean, and the synthetic truth, respectively.




Figure S5. Timeseries of (a) the cost of levee raising γ_E , (b) the rate by which new properties can be built φ_P , (c) the rate of decay of levees κ_T , (d) memory loss rate μ_S estimated by the data assimilation of all observations (F, G, D, H, and M) with 5000 ensembles every 50 years in the experiment 2 (see section 3.1.2). The time step is annual. Grey, red, and black lines are the ensemble members, their mean, and the synthetic truth, respectively.




Figure S6. The ratio of RMSEs of the no data assimilation experiment (NoDA) to those of the data assimilation experiments in which the observation error is set to 10%, 15%, 20%, and 25% of the synthetic true values in the experiment 2 (see section 3.1.2). (a) Blue, orange, gray, and yellow bars are RMSEs of the size of the human settlement $G(t)$, the center of mass of the human settlement from the river $D(t)$, the flood protection level (or levee height) $H(t)$, and the social awareness of the flood risk $M(t)$. (b) Blue, orange, gray, and yellow bars are RMSEs of the cost of levee raising γ_E , the rate by which new properties can be built φ_P , the rate of decay of levees κ_T , memory loss rate μ_S .




Figure S7. The ratio of RMSEs of the no data assimilation experiment (NoDA) to those of the data assimilation experiments in which all of observations (F, G, D, H, and M) are assimilated with 20, 50, 100, 250, 1000, and 5000 ensembles in the experiment 2 (see section 3.1.2). (a) Blue, orange, gray, and yellow bars are RMSEs of the size of the human settlement $G(t)$, the center of mass of the human settlement from the river $D(t)$, the flood protection level (or levee height) $H(t)$, and the social awareness of the flood risk $M(t)$. (b) Blue, orange, gray, and yellow bars are RMSEs of the cost of levee raising γ_E , the rate by which new properties can be built φ_P , the rate of decay of levees κ_T , memory loss rate μ_S




Figure S8. Timeseries of (a) the cost of levee raising γ_E , (b) the rate by which new properties can be built φ_P , (c) the rate of decay of levees κ_T , (d) memory loss rate μ_S estimated by the data assimilation of observations of G and H with 5000 ensembles in the real-world experiment in the city of Rome. The timestep is annual. Grey and red lines are the ensemble members and their mean, respectively.




Figure S9. Same as Figure S8 but only real data of G are assimilated.




Figure S10. Same as Figure S8 but only real data of H are assimilated.