

## *Corrigendum to* "In-stream *Escherichia coli* modeling using high-temporal-resolution data with deep learning and process-based models" published in Hydrol. Earth Syst. Sci., 25, 6185–6202, 2021

Ather Abbas<sup>1</sup>, Sangsoo Baek<sup>1</sup>, Norbert Silvera<sup>2</sup>, Bounsamay Soulileuth<sup>3</sup>, Yakov Pachepsky<sup>4</sup>, Olivier Ribolzi<sup>5</sup>, Laurie Boithias<sup>5</sup>, and Kyung Hwa Cho<sup>1</sup>

<sup>1</sup>School of Urban and Environmental Engineering, Ulsan National Institute of Science and Technology, Ulsan 689-798, Republic of Korea

<sup>2</sup>Institute of Ecology and Environmental Sciences of Paris (iEES-Paris), Sorbonne Université,

Univ. Paris Est Creteil, IRD, CNRS, INRA, Paris, France

<sup>3</sup>IRD, IEES-Paris UMR 242, c/o National Agriculture and Forestry Research Institute, Vientiane, Lao PDR <sup>4</sup>Environmental Microbial and Food Safety Laboratory, USDA-ARS, Beltsville, MD, USA <sup>5</sup>Géosciences Environnement Toulouse, Université de Toulouse, CNRS, IRD, UPS, Toulouse, France

Correspondence: Kyung Hwa Cho (khcho@unist.ac.kr) and Laurie Boithias (laurie.boithias@get.omp.eu)

Published: 18 March 2022

The authors regret that the published version of the above article contained an error in the acknowledgements and financial support section. The correct funding statements are as follows.

Acknowledgements. This work was supported by the Korea Environment Industry and Technology Institute (KEITI) through the Aquatic Ecosystem Conservation Research Program, funded by the Korea Ministry of Environment (MOE) (2020003050001). The authors sincerely thank the Lao Department of Agricultural Land Management (DALaM) for its support, including granting the permission for field access, and the M-TROPICS Critical Zone Observatory (https://mtropics.obs-mip.fr, last access: 16 March 2022), which belongs to the French Research Infrastructure OZCAR (http://www.ozcar-ri.org/, last access: 16 March 2022), for data access. The authors also thank Campus France (PHC STAR 41510WH) for their financial support. *Financial support.* This work was supported by the KEITI through the Aquatic Ecosystem Conservation Research Program, funded by the MOE (2020003050001).

The authors would like to apologize for any inconvenience.