



Improving streamflow simulation through machine learning-powered data integration and its potential for forecasting in the Western U.S.

Yuan Yang¹, Ming Pan¹, Dapeng Feng², Mu Xiao¹, Taylor Dixon¹, Robert Hartman³, Chaopeng Shen⁴, Yalan Song⁴, Agniv Sengupta¹, Luca Delle Monache¹, and F. Martin Ralph¹

¹Center for Western Weather and Water Extremes, Scripps Institution of Oceanography, University of California San Diego, CA, USA

²Department of Earth System Science, Stanford University, Stanford, CA, USA

³Robert K. Hartman Consulting Services, Roseville, CA, USA

⁴Civil and Environmental Engineering, Pennsylvania State University, PA, USA

Correspondence: Yuan Yang (yuanyangthu@gmail.com)

Received: 10 April 2025 – Discussion started: 30 April 2025

Revised: 16 August 2025 – Accepted: 4 September 2025 – Published: 21 October 2025

Abstract. Accurate streamflow forecasts are crucial but remain challenging for the arid Western United States (U.S.). Recently, machine learning methods such as long short-term memory (LSTM) have exhibited high accuracy in streamflow simulation and strong abilities to integrate observations to enhance performance. This study evaluated an LSTM-based data integration approach that incorporates streamflow (Q) and snow water equivalent (SWE) observations to improve streamflow estimations across different lag times (1–10 d, 1–6 months) and timescales (daily and monthly) over hundreds of basins in the Western U.S. Integrating Q at the daily scale provided the greatest improvements, increasing the median Kling-Gupta Efficiency (KGE) of 646 basins from 0.80 to 0.96 when integrating 1 d lagged Q , and remaining at 0.89 even with a 10 d lag. Integrating Q at the monthly scale also enhanced streamflow estimations, though to a lesser extent than at the daily scale, with the median KGE rising from 0.80 to 0.86 when integrating 1-month lagged streamflow. The next most notable improvement resulted from integrating SWE at the monthly scale, where the median KGE improved to 0.86 when integrating 1-month lagged SWE. Furthermore, SWE integration showed greater benefits at the monthly scale in snow-dominated basins during snowmelt season, which was beneficial for spring-summer flow estimations. However, integrating SWE at the daily scale did not show improvements. These results highlight the potential of this LSTM-based data integration approach for both short-

term and long-term streamflow forecasting due to its performance, automation and efficiency.

1 Introduction

Accurate, reliable, and easily implementable hydrological forecasts are crucial for Western United States (U.S.), a region characterized by arid conditions and high water demand (Baker et al., 2021; Fleming et al., 2021; Hunt et al., 2022; Pierce et al., 2008). Short-term forecasts aid in flood risk mitigation, while long-term forecasts facilitate water allocation, reservoir operations, hydropower generation, and drought resilience (Broxton et al., 2023; Yaseen et al., 2015). However, this region's complex topography, including deserts, mountains, valleys, and coastal areas, along with its localized climate dynamics, such as atmospheric rivers, monsoons, and seasonal snowpack, pose significant challenges for accurate streamflow forecasting (Zeng et al., 2018).

Operational agencies employ various streamflow forecast practices, tailored to their specific needs and regional characteristics. The U.S. Department of Agriculture Natural Resources Conservation Service (NRCS) utilizes principal component regression (PCR), a statistical model to predict streamflow based on selected predictors (Garen, 1992; Perkins et al., 2009). The National Weather Service (NWS)

River Forecast Centers (RFC) developed the Hydrological Ensemble Forecast System (HEFS), which uses the Sacramento Soil Moisture Accounting (SAC-SMA) and SNOW-17 models to generate streamflow forecasts across different timescales (Brown et al., 2014; Demargne et al., 2014). While historically successful, these techniques have become less skillful due to regional climate change and other technical limitations, necessitating potential upgrades or replacements (Fleming and Goodbody, 2019). For instance, the recently developed National Water Model is intended to serve as the basis for the future U.S. streamflow forecasting system (Cosgrove et al., 2024). Additionally, these models require extensive manual expertise for domain-specific implementation, such as subjective predictor selection, careful empirical regression identification, and labor-intensive parameter calibration (Fleming et al., 2021). Moreover, they struggle to ingest new observations to enhance streamflow forecasts without substantial structural modifications, such as recalibrating regressions or integrating data assimilation techniques (Franz et al., 2014; Gichamo and Tarboton, 2019). For example, the California-Nevada RFC (CNRFC) employs a “forecasters-in-the-loop” approach, where forecasters manually adjust predictions as new information becomes available, leveraging their prior experience to enhance forecast accuracy.

With the ever-increasing data availability and large advancements in computing technologies, machine learning (ML) models have emerged as promising alternatives to alleviate these limitations. ML models can automatically extract useful information from complex datasets and generate accurate estimation without requiring extensive knowledge of the underlying physical systems (LeCun et al., 2015; Prasad et al., 2017; Schmidhuber, 2015; Shen, 2018; Shen et al., 2023), thereby reducing the need for manual interventions. Moreover, ML models can easily absorb new datasets during training (Shen, 2018), scale efficiently to multiple catchments (Feng et al., 2020; Kratzert et al., 2018), and extrapolate proficiently to ungauged basins (Feng et al., 2021; Kratzert et al., 2019a). Therefore, a surge in applying ML models for streamflow forecasting has been observed in recent years (Fleming et al., 2021; Nearing et al., 2024). For example, the multi-model machine learning metasystem (M^4) is currently being developed as the next-generation operational forecasting system in NRCS (Fleming and Goodbody, 2019). Among the various ML models, one increasingly popular model is the Long Short-Term Memory (LSTM) network, a specifically designed version of recurrent neural network (RNN) for long-term sequential datasets (Greff et al., 2016; Hochreiter and Schmidhuber, 1997). With its unique structure of memory cells and gating mechanisms, LSTM effectively manages the flow of information over long sequences, enabling the retention of relevant input data while discarding less important information. A growing body of research has demonstrated LSTM’s seemingly incomparable performance in streamflow estimation at both daily and monthly scales (Ayana et al.,

2023; Cheng et al., 2020; Clark et al., 2024; Dalkilic et al., 2023; Feng et al., 2020, 2021; Frame et al., 2022; Gauch et al., 2021; Kratzert et al., 2019a; Lees et al., 2021; Nearing et al., 2024).

Incorporating observations is important to improve streamflow estimation, as it helps adjust model states to better represent actual hydrological conditions (Sabzipour et al., 2023). In the context of LSTM-based models, this can be achieved through methods such as data assimilation (DA) or data integration (DI, Feng et al., 2020; Song et al., 2024), the latter also referred to as “autoregression” in Nearing et al. (2022). Similar to traditional DA in hydrological models, DA in LSTM-based models computes the difference between simulations and observations, and propagates it backward into the model to update the model’s internal states. This process relies on inverse procedures, such as variational optimization, and ensemble-based conditional probability estimation, which are not only computationally intensive but also highly sensitive to parameters related to error distributions, regularization coefficients, and resampling procedures (Bannister, 2017; Nearing et al., 2018; Snyder et al., 2008). In contrast, DI directly incorporates observations as inputs and lets LSTM autonomously learn how to optimally utilize this information to enhance estimation. A comparative analysis by Nearing et al. (2022) demonstrated that DI is more accurate and computationally efficient than DA, making it a preferable approach for improving LSTM-based streamflow estimation.

Several studies have demonstrated that directly integrating streamflow observations into the LSTM inputs can significantly improve daily streamflow estimation but only at one or several gauges (Khoshkalam et al., 2023; Le et al., 2019; Sabzipour et al., 2023). Feng et al. (2020), Mangukiya et al. (2023) and Nearing et al. (2022) extended this analysis to large-scale datasets, yet their findings remained constrained to the daily timescale. On the other hand, snow is the primary source of water in the Western U.S., contributing approximately 53 % of the total streamflow (Li et al., 2017). Despite its critical role, few studies have investigated the impact of integrating snow observations into LSTM on streamflow estimation. One exception is Thapa et al. (2020), which showed that incorporating snow cover area as an input improved monthly streamflow estimation, though this analysis was limited to only one gauge. Furthermore, different hydrological variables exhibit varying persistence within the water cycle. Snow, for example, has a longer memory effect since it acts as a natural reservoir that stores water during winter and gradually releases water throughout the spring and summer snowmelt season. However, a gap remains in the literature regarding the comprehensive evaluation of how different observations, such as streamflow (Q) and snow water equivalent (SWE), affect streamflow estimation across multiple timescales.

Motivated by the demonstrated performance of LSTM, this study evaluated a flexible LSTM-based data integra-

tion approach that incorporates different observations (Q and SWE) to improve streamflow simulations across multiple timescales and hundreds of basins in the Western U.S. In this study, retrospective simulations were conducted using observed meteorological forcings, rather than weather forecasts. Given that accurate simulations form the foundation of reliable streamflow forecasting, the demonstrated performance of this data integration approach in retrospective simulations underscores its potential value for forecasting applications. The findings of this study provide critical insights into (1) the effectiveness of LSTM-based data integration for improving streamflow forecasting in the Western U.S. and (2) the different influence of Q and SWE observations on forecast performance across varying timescales.

2 Methods

2.1 Data

We selected a total of 646 basins (all dots in Fig. 1a) in the Western U.S. from the U.S. Geological Survey (USGS) Geospatial Attributes of Gages for Evaluating Streamflow II (GAGEII; Falcone, 2011; Falcone et al., 2010) database for model training. Basin selection was based on several criteria, including boundary accuracy, basin area, data length, reservoir influences, and visual inspection (Appendix A). To further investigate the effect of integrating SWE data, we identified a subset of 429 snow-dominated basins (blue dots in Fig. 1a) from the selected 646 basins (Appendix A), while the remaining basins (orange) are classified as rain-dominated.

We utilized five forcing variables from CW3E 1 km 1-hourly Meteorological Forcing on NWM Grid (CW3E-Forcing, Pan, 2025) dataset and monthly leaf area index (LAI) climatology (no interannual change) from PROBA-V (Fuster et al., 2020) (Table E1). CW3E-Forcing is generated using an elevation-based downscaling and merging procedure to ingest a series of inputs from different sources with different temporal/spatial resolutions, domains, periods of coverage, and lag times. Key features of this forcing dataset include its long-term record (spanning from 1979 to the present), high resolution (1 km, 1 h), and national-scale coverage across the conterminous United States. Here, we utilized the aggregated daily retrospective data from 1983 to 2022. Note that in this study, we performed retrospective experiments to show the effectiveness of the DI-LSTM approach, therefore, no forecasted forcings were used.

To inform LSTM about basin rainfall-runoff behaviors, we calculated the top 10 sensitive basin attributes according to Kratzert et al. (2019b), including climate, topography, and soil attributes (Table E1) as additional inputs to train the models. These attributes were static and appended to the forcing data as input for LSTM.

The daily streamflow data, used both as the training target as well as the input of streamflow integration experiments,

were obtained directly from the USGS Water Information System.

For SWE, we used the daily 4 km gridded SWE data from the University of Arizona dataset (Broxton et al., 2016; Zeng et al., 2018). This dataset is derived through ordinary Kriging interpolation of SWE values from the Snow Telemetry (SNOTEL) sites and further enhanced by incorporating snow depth measurements from thousands of NWS Cooperative Observer Program (COOP) stations (Dawson et al., 2017).

All gridded data were spatially averaged to the basin scale from their original resolutions. All dynamic datasets were aggregated to both daily and monthly timescales to conduct experiments at these two temporal resolutions.

2.2 Modeling

Due to the great potential of LSTM in hydrological modeling, we adopted the LSTM model to investigate the effects of data integration. Additional LSTM details are in Appendix B.

Overall, we trained two types of LSTM models to assess the potential of leveraging lagged observations to improve streamflow estimation (Fig. 1b). The first type is a standard LSTM model that does not perform data integration (DI) and does not use any historical Q or SWE observations. It serves as a valuable benchmark for the comparison against DI-LSTM model. The inputs consist solely of forcings and basin attributes at the current time step and can be expressed as:

$$\mathbf{I}^t = [\mathbf{x}_0^t, \mathbf{A}], \quad (1)$$

Where t is the current time step, $\mathbf{I}^t$ represents the raw input to the model (before data pre-processing), $\mathbf{x}_0^t$ stands for dynamic forcings, and $\mathbf{A}$ represents static basin attributes.

The second type of model is DI-LSTM, which refers to the incorporation of lagged observations (y) into the model (Fig. 1c). The inputs of DI-LSTM can be expressed as:

$$\mathbf{I}^t = [\mathbf{x}_0^t, \mathbf{A}, y^{t-N}], \quad (2)$$

where N is the lag time step, and y^{t-N} is N -step lagged Q or SWE directly from observations. In other words, we fed a N -step-lagged variable y , and let DI-LSTM decide how to use it to dynamically update both cell and hidden states, as well as the LSTM weights, thereby minimizing the accumulation of compounding errors and achieving a better estimation. The only difference between DI-LSTM model and the standard LSTM is whether lagged observations are incorporated in the inputs. Compared with the complex DA techniques used in conceptual or process-based models, this LSTM-powered DI method is relatively straightforward. Its higher computational efficiency and lower development costs make it a promising candidate for operational implementation.

2.3 Experiments

In this study, we evaluated our DI algorithm with two variables: lagged Q and SWE. Given that the effects of DI are




Figure 1. (a) Study basins: blue dots stand for snow-dominated basins, orange dots stand for rain-dominated basins. (b) models: LSTM vs. DI-LSTM model. (c) DI-LSTM with data integration of N -step lagged observations.

expected to vary across different timescales, we tested the algorithm at both daily and monthly scales across all selected basins. For the daily scale, lag times ranged from 1 to 10 d were considered, aligning with the focus of short-term operational forecasts, which typically target lead times within 10 d due to rapidly increasing uncertainty beyond this range. For the monthly scale, 1- to 6-month lags were chosen to reflect typical forecasting horizons used in broader water resource planning and management. In the following text, we used $\text{DI}(Q-N)$ or $\text{DI}(\text{SWE}-N)$ to denote the integration with Q or SWE from N time steps ago. Additionally, to assess whether integrating SWE has a more pronounced effect in snow-dominated basins, we conducted an additional set of LSTM and $\text{DI}(\text{SWE}-N)$ experiments specifically for the 429 snow-dominated basins. In total, 52 experiments were conducted in this study. A summary of these experiments is provided in Table 1.

For each experiment, training data from all selected basins during the 1983–2002 period was used to train LSTM and DI-LSTM models, enabling the network to learn a general understanding of the rainfall-runoff process. The inputs included six meteorological features and 10 static basin attributes (Table E1). The loss function was the Root-Mean-Squared Error (RMSE). Standard pre-processing techniques, including normalization and standardization, were applied to ensure compatibility across different input types and to facilitate effective parameter optimization (See Appendix C

for details). Lagged observations were directly appended to the original LSTM inputs and underwent the same preprocessing procedures. Hyperparameters, such as the number of hidden/cell states and the length of the input sequence, were determined separately for daily and monthly scales. For the daily scale, hyperparameter combinations were inherited from our previous studies (Feng et al., 2020; Song et al., 2024; Yang et al., 2025). For the monthly scale, hyperparameters were determined through a simple grid search across a predefined range of values (Table E2). Final selections were based on analysis of training and validation RMSE learning curves, with the chosen settings minimizing validation RMSE while avoiding overfitting. A fast and flexible LSTM framework from the open-source hydroDL repository (Fang et al., 2021) was implemented.

Missing values are common in streamflow data, yet a naive LSTM cannot operate if any of its inputs are missing. To address this limitation in $\text{DI}(Q)$ experiments, we initially trained the standard LSTM model by filling in missing data with the mean of the training period and subsequently replaced the missing lagged streamflow data with the corresponding LSTM-modeled streamflow data at the same lag time. To prevent missing target (streamflow) values from influencing the model training, for all experiments, the loss function calculation excluded simulations where the corresponding streamflow observations were missing.

Table 1. Experiments.

Time Scale	Lag Time (N)	DI Observations	Training Basins	Experiment Name
Daily	1–10 d	Q	All	Daily DI(Q - N)
		SWE	All & snow-dominated*	Daily DI(SWE- N)
Monthly	1–6 months	Q	All	Monthly DI(Q - N)
		SWE	All & snow-dominated*	Monthly DI(SWE- N)

* Only used in Sect. 4.2

To account for stochasticity in the neural network training and to provide more reliable results (Fig. E1), we performed an ensemble of six randomly seeded trainings, and the mean of all six model simulations was used for the model evaluation.

2.4 Evaluation

We evaluated the ensemble mean simulations from two types of models, LSTM and DI-LSTM, for 2003–2022, independent from the training period. The differences between the two kinds of simulations showed the effect of integrating lagged observations. Metrics adopted to evaluate model performance included the modified Kling-Gupta Efficiency (KGE, Kling et al., 2012) and its three component metrics: correlation coefficient (CC, for temporal coherence), relative variability (RV, for bias in variability), and relative bias (RB, for bias in magnitude). The equations of the four metrics are shown in Table E3. We also calculated the percent bias of the top 2 % peak flow range (FHV) and the percent bias of the bottom 30 % low flow range (FLV) to highlight the performance of the model for peak flows and baseflow, respectively.

3 Results

3.1 The effectiveness of DI(Q) at the daily scale

The daily baseline LSTM without any DI already showed a very promising simulation, with a median KGE of 0.80, a median CC of 0.92, a median RV of 0.94, and a median RB of -10.34% during the test period (Table 2, Fig. 2). Better performance can be seen over more humid regions, while only 12 basins show negative KGE values (Fig. 3), these basins are located in hyper-arid regions with predominantly zero streamflow throughout the evaluation period (e.g., gauge c in Fig. E2). This result, consistent with previous studies, such as Feng et al. (2024), Kratzert et al. (2019b) and Nearing et al. (2024), highlights the ability of a large-scale LSTM model to learn hydrologic behaviors across diverse basins without strong prior structural assumptions.


Overwhelming benefits were observed from integrating lagged streamflow, consistent with previous studies in CONUS (Feng et al., 2020; Nearing et al., 2022), India (Man-

gukiya et al., 2023) and Canada (Khoshkalam et al., 2023; Sabzipour et al., 2023). Compared to the baseline LSTM, all DI(Q) experiments exhibited significantly improved median values (Table 2, $p \leq 0.05$, Kolmogorov-Smirnov test, Eghbali, 1979; Smirnov, 1948) as well as substantially reduced variability across all metrics (Fig. 2). After integrating the 1 d lagged streamflow, the median KGE, CC, RV and RB improved to 0.96, 0.98, 0.96 and 1.24% , respectively, approaching nearly perfect values. Negative KGE values were observed in only three basins, all located in hyper-arid regions with mean daily streamflow below $1 \text{ m}^3 \text{ s}^{-1}$. Integrating lagged daily streamflow also improved the relative bias of both low and high flows. Although the median FLV remained largely unchanged, which was already close to zero, the variability of FLV was largely reduced, indicating consistently low values across basins. The underestimation of high flows was significantly reduced, with median values shifting closer to zero and a narrower range of variability. The compaction of FHV was less pronounced than that of FLV. Peak flows often occur over shorter timescales (e.g., during storm events lasting less than 1 d), and thus their predictability relies more on immediate forcings than on accumulated hydrologic memory. As a result, the integration of lagged streamflow was less effective in improving high flow estimates than low flow estimates. Nevertheless, the benefits of DI(Q) were still noticeable with FHV, demonstrating the role of antecedent conditions in influencing flooding.

Spatially, ubiquitous and heterogeneous benefits from daily DI(Q - N) can be observed over the whole Western U.S. Taking DI(Q -1) as an example, most gauges experienced a boost of 0.1–0.3 in KGE, and about 83 % of basins had a KGE larger than 0.9 (Fig. 3). The largest improvements were found in the Rocky Mountains and Sierra Nevada Ranges, where KGE values were boosted from < 0.6 to 0.9–1. For instance, gauges a and b (Fig. E2), located in this mountainous region, illustrate cases where DI(Q -1) substantially improved streamflow simulations. At gauge a , both underestimation and overestimation were notably reduced, resulting in a high KGE of 0.965. At gauge b , DI(Q -1) effectively corrected the pronounced underestimation of baseflow, yielding strong overall performance. Improvements were also observed in the northern region. At gauge d in the Pacific Northwest (Fig. E2), DI(Q -1) reduced peak flow overestima-

Table 2. Median KGE of LSTM, DI(Q) and DI(SWE) experiments.

	Daily Scale		Monthly Scale		Monthly scale, April–July	
	Q	SWE	Q	SWE	Q	SWE
LSTM	0.80	0.80	0.80	0.80	0.76	0.76
$N = 1$ (day/month)	0.96	0.80	0.86	0.82	0.81	0.79
$N = 2$ (day/month)	0.95	0.80	0.85	0.82	0.79	0.78
$N = 3$ (day/month)	0.94	0.81	0.85	0.82	0.80	0.78
$N = 4$ (day/month)	0.93	0.80	0.84	0.81	0.78	0.76
$N = 5$ (day/month)	0.92	0.80	0.84	0.81	0.78	0.76
$N = 6$ (day/month)	0.92	0.80	0.83	0.80	0.78	0.76
$N = 7$ (day/month)	0.91	0.81	–	–	–	–
$N = 8$ (day/month)	0.90	0.81	–	–	–	–
$N = 9$ (day/month)	0.90	0.81	–	–	–	–
$N = 10$ (day/month)	0.89	0.80	–	–	–	–

**Figure 2.** Performance of LSTM (black) and DI(Q - N) ($N = 1$ – 10) experiments (red) at the daily scale. The “B” on the x-axis stands for baseline LSTM, and N stands for DI(Q - N) experiment. The black horizontal line stands for the median value of the baseline LSTM. The grey horizontal line shows perfect value for RV, RB, FLV, and FHV. The boxplots display the median, 25th/75th percentiles, the lowest datum above $Q1 - 1.5 \cdot (Q3 - Q1)$ (lower whisker), and the highest datum below $Q3 + 1.5 \cdot (Q3 - Q1)$ (upper whisker).

tion and increased the KGE to 0.947. The spatial pattern of improvements shows a positive correlation with the streamflow autocorrelation, with the strongest benefits in regions with high streamflow autocorrelation (Fig. E3). In several southern basins, utilizing lagged streamflow observations did not improve simulations. For example, DI(Q -1) did not improve the simulation at gauge c in the southwest (Fig. E2), which exhibited no baseflow and 1 d flash peaks. One possible explanation is that these are highly arid basins with low streamflow autocorrelation and flash floods (Li et al., 2022; Mangukiya and Sharma, 2025; Saharia et al., 2017). The sudden sharp streamflow peaks in these basins typically persist for less than one day and have little relationship with the previous day’s streamflow, limiting the effectiveness of lagged streamflow observations.

In general, more recent observations typically contribute more to predictive improvements (Cheng et al., 2020; Sabzipour et al., 2023). The benefits of daily DI(Q) gradually decayed as N increased, with a corresponding widening of metric variability (Fig. 2). This gradual decay of DI(Q) benefits, to a certain extent, reflects the memory length of hydrological processes (Feng et al., 2020; Sabzipour et al., 2023). However, even in the DI(Q -10) experiment, the median KGE, CC, RV and RB remained at 0.89, 0.95, 0.95 and -3.00% , respectively, still outperforming the baseline LSTM. This demonstrates that integrating streamflow from 10 d ago remains valuable for daily streamflow simulations. If implemented in a forecasting mode, the results suggest that near real-time streamflow observations could be leveraged to enhance short range streamflow forecast across these basins




Figure 3. Comparison of KGE spatial patterns over the Western U.S. for experiments at the daily scale (left), monthly scale (middle) and monthly scale but only evaluation for April to July (right). From top to bottom: (a–c) LSTM, (d–f) $DI(Q-1)$, (g–i) $\Delta KGE = KGE_{DI(Q-1)} - KGE_{LSTM}$. $N1/N2$ on (g)–(i) stands for the number of basins where $DI(Q-1)/LSTM$ performs better, respectively.

in the Western U.S., relative to models without such observations.

3.2 The effectiveness of $DI(Q)$ at the monthly scale

At the monthly scale, the baseline LSTM simulated streamflow well, achieving a median KGE of 0.80, quite similar to the daily-scale results. This consistency in performance across temporal resolutions aligns with findings from Yao et al. (2023), indicating that the standard LSTM is largely unaffected by changes in temporal resolution. Integrating lagged streamflow observations from 1 to 6 months ago also significantly improved model performance, yielding higher median values and reduced variability across all metrics. Monthly $DI(Q-1)$ achieved a median KGE of 0.86 (Fig. 4a) and enhanced simulations in about 76 % of basins (Fig. 3). For example, $DI(Q-1)$ largely reduced the underestimation in the baseflow and overestimation in the peak flow, leading to much higher KGE values for gauges *a*, *b* and *d* in Fig. E2. However, its effectiveness remained limited in hyper-arid regions, such as at gauge *c* (Fig. E2), where overall simulation accuracy did not improve. $DI(Q-6)$ still exhibited a higher median KGE (0.83) and a smaller spread, showing the ad-

vantage of integrating monthly streamflow. However, the improvements at the monthly scale were less pronounced than those at the daily scale. This was expected since the monthly streamflow autocorrelation is usually weaker (Fig. E3), and lagged streamflow provides reduced predictive value. Effective water management in the Western U.S. depends heavily on spring–summer (April–July) streamflow volume forecasts, commonly referred to as seasonal Water Supply Forecasts (WSFs). To assess model performance during this critical period, we evaluated streamflow from April to July. When evaluated specifically for the April–July period, LSTM performed slightly worse than the full-year analysis, with a median KGE of 0.76, but with a similar spatter pattern (Fig. 3). As in the full-year results, several arid basins in the southern region exhibited very low KGE values, highlighting the need for further research to improve simulations in arid environments. However, integrating lagged monthly streamflow significantly contributes to better performance, with higher median KGE values for monthly $DI(Q-1)$ and monthly $DI(Q-6)$ (0.81 and 0.78, respectively) as well as reduced variability (Fig. 4c). The improvements for the April–July flow exhibited a spatial pattern similar to those observed for year-round flow, albeit with reduced magnitude. This dif-

ference in magnitude is likely attributable to loss functions in monthly DI(Q) experiments being optimized for year-round flow rather than specifically tailored to the April–July period.

3.3 The effectiveness of DI(SWE) at the daily scale

In contrast to daily DI(Q), integrating lagged SWE data at the daily scale did not improve streamflow simulations in terms of KGE (Fig. 5). This outcome aligns with expectations, as snow-related processes typically have a longer memory effect. Moreover, temperature, one of the model inputs, partially reflects snow dynamics, which the LSTM can effectively leverage through its memory states to estimate streamflow. However, significant improvements were still observed in CC and RV, indicating that DI(SWE) can enhance temporal dynamics and reduce variability biases. The overestimation was reduced, particularly during low-flow conditions, while underestimation worsened, leading to poorer RB medians. This increased underestimation may stem from the prevalence of seasonal snowpack in most basins, where abundant days with zero SWE values could introduce bias when integrated into the model. Additionally, the quality of the SWE dataset itself likely plays a role. Further investigation, such as utilizing SWE data from Airborne Snow Observatory (ASO, Painter et al., 2016) or snow course, is needed to better understand the underestimation issue.

Spatially, most improvements were observed in the Rocky Mountains (Fig. 6), where deeper snowpack usually exists and flow is dominated by snow. To further investigate whether the effect of integrating lagged SWE varies across different snowpacks, we evaluated model performance separately over rain-dominated basins (orange dots in Fig. 1a) and snow-dominated basins (blue dots in Fig. 1b). Figure 7a and e present the KGE values of the LSTM model, while Fig. 7b and f show the KGE differences between DI(SWE) and LSTM at the daily scale for both types of basins. The baseline LSTM performed better in snow-dominated basins, with a higher median KGE of 0.80 (compared to 0.77 for rain-dominated basins) and smaller variability (Fig. 7). In terms of KGE differences, snow-dominated basins showed no obvious improvement, with a median Δ KGE of zero, while more rain-dominated basins exhibited negative Δ KGE after integrating lagged SWE. These rain-dominated basins are mainly located on the west side of the Cascade Mountains, the eastern slope of the Rocky Mountains, and the Southwest, where snowmelt is less dominant and rainfall contributes significantly to streamflow. Consequently, utilizing lagged SWE data did not show an impact on streamflow; instead, adding more zero SWE values into the LSTM model led to increased underestimation, ultimately degrading performance. To illustrate the effect of daily DI(SWE) in different hydrologic regimes, we highlight two representative gauges from snow- and rain-dominated basins. Gauge *a*, located in Yellowstone National Park (Fig. E4), sits at a high

elevation (2355 m) and receives substantial winter snowfall, which serves as a primary contributor to streamflow. Integrating daily SWE data at this site helped reduced the underestimation of peak flows. In contrast, gauge *b*, situated in California's Central Coast region (Fig. E4), experiences minimal snowfall and is predominantly influenced by seasonal rainfall. As a result, incorporating near-zero SWE data did not improve simulation performance at this site.

Considering the delayed effect of snow processes on streamflow generation, we further investigated the effect of integrating SWE from different seasons (accumulation and snowmelt) on streamflow. The snow accumulation and snowmelt season are defined individually for each basin and each water year following the methodology of Trujillo and Molotch (2014) (Appendix D). Here we focused exclusively on snow-dominated basins, as minimal improvements were observed in rain-dominated basins (Fig. 7). Figure 8 shows the metric differences between DI(SWE) and LSTM during accumulation and snowmelt seasons over snow-dominated basins. The percentage of basins with positive Δ CC increased from 53 %–61 % during accumulation season to 73 %–77 % during snowmelt season. Notably, the median values of Δ CC during snowmelt season exceeded even the 75th percentiles of accumulation season (Fig. 8b), indicating stronger performance gains in temporal dynamics. More improvements were also observed in RV during snowmelt season, with more basins showing RV values closer to ideal value 1 (negative $|RV-1|$) and larger negative median $\Delta|RV-1|$ (Fig. 8c). However, larger $\Delta|RB|$ were also observed during the snowmelt season. As a result, when considering the comprehensive metric, KGE, snowmelt season demonstrated only a slight improvement in median Δ KGE compared to the accumulation season.

3.4 The effectiveness of DI(SWE) at the monthly scale

Due to the long memory of snow processes in the hydrological cycle, integrating lagged SWE at the monthly scale provided benefits to streamflow simulation, as evidenced by slightly higher median KGE values as well as smaller spreads (Fig. 4). For instance, integrating lagged SWE from one month ago led to improved KGE in about 65 % of basins (Fig. 6), with the median KGE increasing from 0.80 to 0.82. A similar spatial pattern of improvements, with slightly higher magnitude as indicated by the darker blue dots in Fig. 6i, was also observed when evaluating spring–summer (April–July) streamflow.

The benefits of DI(SWE) at the monthly scale gradually declined as N increased, reflecting the decreasing persistence of snow in the hydrological cycle and its diminishing predictive value over longer lag periods (Fig. 4). However, DI(SWE-6) still showed some improvements, with slightly higher 25th and 75th percentiles and smaller interquartiles, despite an almost unchanged median. This suggests that integrating SWE data from six months ago remains informa-




Figure 4. KGE boxplots for DI(Q -N) (left) and DI(SWE-N) (right) at monthly scale (top) and monthly scale but only evaluation from April to July (bottom).




Figure 5. Performance of LSTM (black) and DI(SWE-N) ($N = 1$ –10) experiments (green) at the daily scale. The “B” on the x -axis stands for baseline LSTM, and N stands for the DI(SWE-N) experiment. The black horizontal line stands for the median value of the baseline LSTM. The grey horizontal line shows perfect value for RV, RB, FLV, and FHV.

tive for streamflow simulation. Therefore, if implemented in a forecasting mode, the findings suggest that near real-time SWE observations have the potential to enhance long-term monthly streamflow forecasts, relative to models without such observations.

The benefits of DI(SWE) at the monthly scale were more pronounced in snow-dominated basins compared to rain-dominated basins (Fig. 7c and g). For example, as shown in Fig. E4, the snow-dominated gauge *a* exhibited substantial improvement in peak flow simulation, while the hydropograph at the rain-dominated gauge *b* showed little to

change. This improvement difference became even more evident when evaluating streamflow from April to July, the primary snowmelt season (Fig. 7d and i), further emphasizing the greater impact of DI(SWE) in snow-dominated basins during snowmelt season.




Figure 6. Comparison of KGE spatial patterns over the Western U.S. at the daily scale (left), monthly scale (middle) and monthly scale but only evaluation for April to July (right). From top to bottom: (a–c) LSTM, (d–f) DI(SWE-1), (g–i) $\Delta KGE = KGE_{DI(SWE-1)} - KGE_{LSTM}$. $N1/N2$ on (g)–(i) stands for the number of basins where DI(SWE-1)/LSTM performs better, respectively.




Figure 7. Comparison of KGE over rain-dominated basins (top) and snow-dominated basins (bottom). (a, e) black boxplots stand for KGE of baseline LSTM, and D , $M1$, $M2$ on the x-axis stand for the results of daily scale, monthly scale and monthly scale but evaluation only for April to July. (b–d, f–h) colored boxplots stand for KGE difference between DI(SWE-N) and LSTM ($\Delta KGE = KGE_{DI(SWE-N)} - KGE_{LSTM}$) at the daily, monthly, and monthly scale but evaluation only for April to July. The grey horizontal lines are zero.




Figure 8. Metric differences between DI(SWE-N) and LSTM over snow accumulation and snowmelt seasons (difference in KGE, CC, $|RV-1|$, and $|RB|$) over snow-dominated basins. $\Delta|RV-1|$ is used since the ideal value of RV is 1. Only median and interquartile range (25th–75th) are shown here. N stands for DI(SWE-N) experiment. The grey horizontal lines show zero.

4 Discussions

4.1 Comparison of integrating different observations at different timescales

Figure 9 summarizes the median KGE values for all experiments at different timescales over all basins, as shown in Figs. 2, 4, and 5, separately. The benefits of different integration experiments can be roughly ranked as follows:

daily DI(Q) > monthly DI(Q) > monthly DI(SWE)
> daily DI(SWE)

Consistent patterns were also observed specifically over snow-dominated basins, as shown in Fig. E5. It is counterintuitive that even over snow-dominated basins at the monthly scale and during April–July period, integrating lagged streamflow observations provided greater improvements than integrating SWE, despite snow being a key predictor of spring–summer flow in the snow-dominated Western U.S. (Fleming et al., 2024; Koster et al., 2010; Shukla and Lettenmaier, 2011; Wood et al., 2016). This outcome is likely attributable to the inherent characteristics of the LSTM architecture. Due to its memory-based structure, the LSTM is well-suited for capturing long-term dependencies and cumulative processes. As a result, it can effectively learn the snow-related dynamics implicitly from historical meteorological forcings (e.g., precipitation and temperature) and streamflow responses, without requiring explicit SWE input (Feng et al., 2020; Jiang et al., 2022; Modi et al., 2025). For

example, the model may internally infer snowpack accumulation when precipitation coincides with subfreezing temperatures and simulate melt-driven streamflow increases when temperature rise. Consequently, because the model already captures key snow dynamics internally, the integration of external SWE observations provides less incremental value than integrating direct streamflow observations.

In the monthly-scale analysis, DI(Q) yielded slightly greater improvements when evaluated over the entire year, whereas DI(SWE) showed a marginally larger enhancement in spring–summer flow estimates when integrating lagged SWE from 1–3 months prior.

4.2 Comparison of DI(SWE) between snow-dominated basins and all basins

From the above analysis, we found that DI(SWE) experiments showed greater improvements when evaluated over snow-dominated basins. To further explore this, we conducted the same DI(SWE) experiments exclusively trained over snow-dominated basins to determine if additional gains could be achieved. As expected, training the models (both LSTM and DI(SWE)) over a more homogeneous group of basins provided higher performance (Fig. E6). Figure 10 shows the median ΔKGE between DI(SWE) and the corresponding baseline LSTM over all basins and snow-dominated basins. Similar to daily DI(SWE) trained over all basins, daily DI(SWE) trained exclusively over snow-dominated basins did not enhance streamflow estimation and even slightly degraded performance. However, at the




Figure 9. Median KGE values of all experiments at the daily scale (left), monthly scale (middle) and monthly scale but only evaluation for April to July (right) over all basins. N on the x -axis stands for DI(Q - N) or DI(SWE- N) experiment.

monthly scale, DI(SWE) improved streamflow estimations for both the whole year flow and April–July flow. This improvement became more pronounced for the April–July period, reinforcing the finding that integrating SWE has a larger effect on streamflow estimation over snow-dominated basins during snowmelt season.

4.3 Potential operational forecast applications and limitations

ML is gaining popularity in hydrology research and operational communities. This trend is driven by several key factors, including its easy implementation without substantial development and operational costs, strong model performance, ability to handle complex prediction tasks, and flexible model structure to adapt new datasets as additional predictors during training. Moreover, ML enables automated and objective modeling, minimizing the need for extensive manual interventions and subjective decision-making (Fleming et al., 2021, 2024; Modi et al., 2025).

This study evaluated the performance of an LSTM-powered data integration model that integrates lagged Q and SWE observations across various lag times at both daily and monthly scales. The pronounced improvements observed in the retrospective experiments highlight its potential for forecasting applications. In forecasting mode, recent observations can be incorporated into the LSTM model to dynamically update hydrological conditions, reducing the initialization errors compared to models that rely solely on forecasted forcings. In this framework, the “lag time” in retrospective simulations corresponds to the “lead time” in forecasting mode. In other words, integrating recent Q or SWE data into the LSTM model could enhance streamflow forecasts in the Western U.S. at both short lead times (daily scale) and extended lead times (monthly scale), relative to the baseline LSTM model without such integration. Given its demonstrated effectiveness, flexibility, and automation, this data integration framework holds promises for real-time hydrolog-

ical forecasting, offering valuable applications in water resource management.

Despite much promise, the DI-LSTM approach would have certain limitations when applied to operational streamflow forecasting. First, the improvements demonstrated in this study may be less pronounced in real-world forecasting applications. Here, retrospective simulations were used, leveraging observed meteorological forcings to evaluate the effectiveness of DI-LSTM for streamflow simulations, thereby providing an upper bound on potential performance. However, operational forecast systems rely on predicted forcings, which inherently contain significant uncertainties that impact streamflow forecasts. Additionally, the accuracy of weather forecasts is expected to decay with increasing lead time, further diminishing the DI-LSTM predictive skill for longer lead time. Therefore, further research is necessary to assess the performance of DI-LSTM in an operational setting using actual forecasted meteorological inputs. Moreover, collaboration with the meteorological community is essential to improving the accuracy of forcing predictions. Second, this study provides deterministic streamflow estimation with limited uncertainty analysis. Uncertainty is inherent in all aspects of hydrological modeling, and its estimation is critical for actionable hydrological forecasts (Fang et al., 2020; Klotz et al., 2022). To address uncertainty due to random initial weights and biases, this study employed six repeated runs with different random seeds. However, uncertainties related to model inputs and observational data for model training were not explicitly considered. Recent studies have introduced various methods to quantify uncertainty in ML-based models for different uncertainty sources, such as Markov Chain Monte Carlo, variational inference, Monte Carlo dropout, Mixture density networks and ensemble techniques (Abdar et al., 2021). Future work should further explore uncertainty quantification to enhance forecast reliability and underpin decision-making in water resources management.




Figure 10. Median Δ KGE between DI(SWE) and LSTM over all basins (green) and snow-dominated basins (blue). From left to right are results at the daily scale, monthly scale and monthly scale but evaluation for April to July. N on the x -axis stands for DI(SWE- N) experiment.

5 Conclusions

Based on LSTM, we evaluated a flexible data integration approach (DI-LSTM) incorporating different observations, e.g., Q and SWE, across multiple lag times at both daily and monthly scales over hundreds of basins in the Western U.S. By comparing DI-LSTM with the baseline LSTM, we assessed the impact of integrating lagged observations on streamflow estimations. The key findings in the Western U.S. are summarized as follows:

1. The baseline LSTM without integrating any lagged observations already showed strong predictive capability in the Western U.S., achieving a median KGE of 0.80 at both daily and monthly scales.
2. Integrating Q at the daily scale yielded the most substantial improvements, with significantly improved median values and reduced spread across all performance metrics. The median KGE across 646 basins increased to 0.96 with the integration of 1 d lagged streamflow and remained at 0.89 even with a 10 d lag. Integrating Q at the monthly scale also improved streamflow estimations, though to a lesser extent, with the median KGE increasing from 0.80 to 0.86 when integrating streamflow from 1 month ago.
3. Integrating lagged SWE at the monthly scale led to better accuracy, whereas its integration at the daily scale did not improve streamflow estimations. This finding reflects the long-term memory of snow processes in the hydrological cycle, which extends beyond short timescales.
4. The benefits of integrating SWE were more pronounced in snow-dominated basins during the snowmelt season, highlighting its value for improving spring-summer flow estimations.
5. Overall, the benefits of integrating different observations at different timescales for streamflow estimations can be roughly ranked as follows: daily DI(Q) > monthly DI(Q) > monthly DI(SWE) > daily DI(SWE).

Due to its strong predictive performance, automation without the need for extensive domain-specific customization, and flexibility to ingest additional observations, the DI-LSTM approach demonstrates large potential for short-term (e.g., 1–10 d) and long-term (1–6 months) operational streamflow forecasts in the Western U.S. However, further studies, such as using real forecasted forcing data, are needed to assess its performance under realistic forecasting conditions.

Appendix A: Training basin selection and snow-dominated basin selection

We performed a screening to identify suitable training basins in the Western US by implementing the following procedure:

1. *Basin area:* Only basins within the range of 50–5000 km² were selected. Basins smaller than 50 km² were discarded due to probable artificial boundaries. The maximum area threshold was applied since channel routing effects become apparent at the daily scale in larger basins (Gericke and Smithers, 2014).
2. *Data length:* only basins with at least 10 year data during the training period (1983–2002) were selected to ensure sufficient data for training.
3. *Reservoir influences:* To minimize the effect of river regulation by dams or reservoirs, only basins with degree of regulation (DOR) no greater than 0.1 were selected (Ouyang et al., 2021). The DOR is defined as the ratio of total reservoir capacity within a basin to the mean annual cumulative discharge, with total reservoir capacity data sourced from GAGEII.
4. *Visual inspection:* Since some data are collected manually, they may contain errors in reported discharge values. We excluded basins with potentially erroneous discharge records, such as those with an unreasonably high magnitude far exceeding precipitation or with abrupt, dramatic differences between time intervals.

For most basins in the Western U.S., streamflow during the April–July period (spring to early summer) is primar-

ily driven by snowmelt or contemporaneous rainfall. In this region, 1 April is widely used as the transition point from snow accumulation season and snowmelt (Musselman et al., 2021). The maximum SWE between October and April is commonly used as an indicator of the total snow available for melt-driven streamflow (Musselman et al., 2021; Mote et al., 2018).

To quantify the relative contributions of snowmelt and rainfall to streamflow, we calculated two correlation indices: (1) the correlation between maximum SWE (October to April) and total streamflow volume (April–July), denoted as $\text{Corr}(\text{maxSWE}, Q_{\text{tot}})$, and (2) the correlation between total rainfall (April–July) and total streamflow volume for the same period, denoted as $\text{Corr}(P_{\text{tot}}, Q_{\text{tot}})$. Based on these indices, snow-dominated basins were identified using the following two criteria:

1. $\text{Corr}(\text{maxSWE}, Q_{\text{tot}}) > \text{Corr}(P_{\text{tot}}, Q_{\text{tot}})$
2. $\text{Corr}(\text{maxSWE}, Q_{\text{tot}}) > 0.1$,

Criterion 1 ensures that snow has a greater influence than rainfall on streamflow, while criterion 2 excludes basins with negligible snow influence, thereby retaining only those basins where snowmelt meaningfully contributes to streamflow.

Appendix B: LSTM model

LSTM introduces “memory cells” and “gates” to keep and filter information. Cell states allow information to be stored over long time periods, which is desirable for modeling processes such as snow accumulation and snowmelt. The input, forget and output gates control the flow of information, controlling what to let in, what to forget, and what to output from the system, respectively. These gates are all trained automatically and simultaneously, using input data to predict the target variable. The forward propagation equations of the LSTM model are described by the following equations:

$$\text{Input transformation: } \mathbf{x}^t = \text{ReLU}(\mathbf{W}_I \mathbf{I}^t + \mathbf{b}_I), \quad (\text{B1})$$

$$\text{Input node: } \mathbf{g}^t = \tanh(\text{D}(\mathbf{W}_{gx} \mathbf{x}^t) + \text{D}(\mathbf{W}_{gh} \mathbf{h}^{t-1}) + \mathbf{b}_g), \quad (\text{B2})$$

$$\text{Input gate: } \mathbf{i}^t = \sigma(\text{D}(\mathbf{W}_{ix} \mathbf{x}^t) + \text{D}(\mathbf{W}_{ih} \mathbf{h}^{t-1}) + \mathbf{b}_i), \quad (\text{B3})$$

$$\text{Forget gate: } \mathbf{f}^t = \sigma(\text{D}(\mathbf{W}_{fx} \mathbf{x}^t) + \text{D}(\mathbf{W}_{fh} \mathbf{h}^{t-1}) + \mathbf{b}_f), \quad (\text{B4})$$

$$\text{Output gate: } \mathbf{o}^t = \sigma(\text{D}(\mathbf{W}_{ox} \mathbf{x}^t) + \text{D}(\mathbf{W}_{oh} \mathbf{h}^{t-1}) + \mathbf{b}_o), \quad (\text{B5})$$

$$\text{Cell state: } \mathbf{s}^t = \mathbf{g}^t \odot \mathbf{i}^t + \mathbf{s}^{t-1} \odot \mathbf{f}^t, \quad (\text{B6})$$

$$\text{Hidden state: } \mathbf{h}^t = \tanh(\mathbf{s}^t) \odot \mathbf{o}^t, \quad (\text{B7})$$

$$\text{Output: } \mathbf{y}^t = \mathbf{W}_{hy} \mathbf{h}^t + \mathbf{b}_y \quad (\text{B8})$$

where $\mathbf{I}^t$ represents the raw input to the model, $\mathbf{x}^t$ represents the input vector to the LSTM cell. ReLU is the rectified linear unit, σ is the sigmoidal function, $\odot$ is the element-wise

multiplication operator, D is the dropout operator. $\mathbf{W}$ and $\mathbf{b}$ with different subscripts represent the gate-specific network weights and bias parameters, respectively. $\mathbf{g}^t$ is the output of the input node, $\mathbf{i}^t$, $\mathbf{f}^t$, $\mathbf{o}^t$ are the input, forget, and output gates, respectively; $\mathbf{h}^t$ represents the hidden states, $\mathbf{s}^t$ represents the memory cell states and $\mathbf{y}^t$ represents the predicted output.

Appendix C: Data pre-processing for LSTM and DI-LSTM

During the iterations of the training process, basins from the entire dataset were randomly sampled to form a mini-batch each time to calculate the loss function. This batching method typically assumes that model errors are identically distributed among basins within the same mini-batch. Without data preprocessing or normalization, the loss function would inherently pay more attention to wetter and larger basins compared to drier or smaller basins. To prevent this imbalance, we applied standard pre-processing techniques, including normalization and standardization, following Feng et al. (2020).

First, we normalized the daily discharge by basin area and mean daily precipitation to obtain a dimensionless discharge value as the target variable.

Then we transformed the distributions of daily discharge and precipitation as close to Gaussian as possible, since these two typically have Gamma distributions, using the equation:

$$v^* = \log_{10}(\sqrt{v} + 0.1) \quad (\text{C1})$$

where v and v^* are the variables before and after transformation, respectively. 0.1 is added inside the log to avoid making the log of zero. Transforming the data to a Gaussian distribution enhances the stability and efficiency of gradient-based optimization methods in LSTM. Additionally, it reduces the impact of extreme peak values during model training, improving the model's representation of low-flow conditions.

Finally, standardization was applied to all input features (forcings, static basin attributes, and lagged observations), as well as the output (discharge) by subtracting the mean value and then dividing by the standard deviation of training-period data.

Appendix D: Snow season definition

The snow accumulation and snowmelt season are defined individually for each basin and each water year (1 October to 30 September) following the methodology of Trujillo and Molotch (2014). For each water year each basin, the date of peak annual SWE is identified. The snow season is then defined as the continuous period during which SWE remains greater than zero and includes the peak SWE. This snow season is subsequently divided into two parts: the accumu-




Figure D1. Snow season definitions. Peak SWE is the highest snow water equivalent (SWE) value in a water year.

lation season, which occurs before the peak SWE date, and the snowmelt season, which follows it (Fig. D1).

Note that the seasonal analysis in this study focuses exclusively on the main SWE curve, i.e., the continuous SWE curve associated with the peak SWE. In basin-years with intermittent snow, there may be several snow accumulation and melt cycles prior to and/or after the main SWE curve which are not accounted for in this analysis.

Appendix E




Figure E1. Performance comparison between the ensemble mean and individual random seed simulations across different experiments at the daily scale: (a) LSTM, (b) $DI(Q-1)$, and (c) $DI(SWE-1)$. “meanflow” refers to the ensemble mean derived from six simulations, while “seed 1” through “seed 6” represent the results from individual random seeds.




Figure E2. Time series plots for selected basins to illustrate the benefits of DI(Q) across different flow regimes. Numbers in the legends represent KGE values of the simulations. (a1–d1) time series comparisons for the daily experiments, (a2–d2) time series comparisons for the monthly experiments. (e) the locations of the corresponding basins.




Figure E3. Spatial distribution of (a) 1 d-lag and (b) 1-month-lag autocorrelation function of streamflow (ACF(1)).




Figure E4. Time series plots for selected basins to illustrate the benefits of DI(SWE) across snow- and rain-dominated basins. Numbers in the legends represent KGE values of the simulations. (a1–b1) time series comparisons for the daily experiments, (a2–b2) time series comparison for the monthly experiments. (c) the locations of the corresponding basins.




Figure E5. Median KGE values of all experiments at the daily scale (left), monthly scale (middle) and monthly scale but only evaluation for April to July (right) over snow-dominated basins. N on the x -axis stands for DI(Q - N) or DI(SWE- N) experiment.




Figure E6. Median KGE values of DI(SWE- N) over all basins (green) and snow-dominated basins (blue). From left to right are results at the daily scale, monthly scale and monthly scale but only evaluation for April to July. N on the x -axis stands for DI(SWE- N) experiments.

Table E1. Summary of the forcing data and attribute variables used in this study.

	Variable	Data Source	Units
Forcing	Daily precipitation	CW3E-Forcing (Pan, 2025)	mm d ⁻¹
	Daily maximum temperature	CW3E-Forcing (Pan, 2025)	°C
	Daily minimum temperature	CW3E-Forcing (Pan, 2025)	°C
	Daily mean surface downwelling shortwave	CW3E-Forcing (Pan, 2025)	W m ⁻²
	Daily mean 10 m wind	CW3E-Forcing (Pan, 2025)	m s ⁻¹
	Monthly LAI climatology	PROBA-V LAI (Fuster et al., 2020)	–
Attributes	Mean daily precipitation	CW3E-Forcing (Pan, 2025)	mm d ⁻¹
	High precipitation duration – the average duration of high precipitation events (number of consecutive days ≥ 5 times mean daily precipitation)	CW3E-Forcing (Pan, 2025)	days
	Fraction of precipitation falling as snow (i.e., on days colder than 0 °C)	CW3E-Forcing (Pan, 2025)	–
	Aridity – P/PET, where PET is estimated by the Hargreaves (1994) method	CW3E-Forcing (Pan, 2025)	–
	Frozen days – days colder than 0 °C	CW3E-Forcing (Pan, 2025)	days
	Area	basin boundary file	km ²
	Mean elevation	GMTED (Amatulli et al., 2018a)	m above sea level
	Mean slope	GMTED (Amatulli et al., 2018a)	°
	Geological permeability	GLHYMPS V2 (Huscroft et al., 2018a)	m ²
	Soil sand content	SoilGrids (Hengl et al., 2017)	%

Table E2. Hyperparameters for the LSTM or DI-LSTM model.

Hyperparameter	Daily Scale	Monthly Scale	
	Best value	Grid search	Best value
Length of training instances	365	12, 24, 36, 48	48
Mini-batching size	100	50, 100, 150, 200	50
LSTM dropout rate	0.5	0, 0.2, 0.5	0.5
LSTM hidden size	256	128, 256	256
Number of training epochs	300	[100, 600]	300
Number of stacked LSTM layer	1	1	1

Table E3. The definition of KGE and its three component metrics.

Metric	Equation	Perfect Value
CC	$CC = \frac{\text{cov}(Q_o, Q_m)}{\sigma_{Q_o} \cdot \sigma_{Q_m}}$	1
RV	$RV = \frac{\sigma_{Q_m} / \mu_{Q_m}}{\sigma_{Q_o} / \mu_{Q_o}}$	1
RB	$RB = \frac{\sum_{i=1}^N Q_{m,i} - \sum_{i=1}^N Q_{o,i}}{\sum_{i=1}^N Q_{o,i}} \times 100$	0
KGE	$KGE = 1 - \sqrt{(CC - 1)^2 + RB^2 + (RV - 1)^2}$	1

Note, Q_o , Q_m represent streamflow observations and simulations, respectively. cov, σ and μ represent covariance, standard deviation and mean, respectively.

Code and data availability. The source codes for LSTM-based rainfall-runoff simulations are from hydroDL, which is available at: <https://doi.org/10.5281/zenodo.5015120> (Fang et al., 2021).

CW3E-Forcing is available at: <https://www.reachhydro.org/home/records/1-km-conus-forcing> (last access: 10 October 2025) (<https://doi.org/10.5281/zenodo.14714512>, Pan, 2025). The PROBA-V LAI is available at: <https://land.copernicus.eu/global/products/lai> (last access: 10 October 2025). Elevation data from GMTED is available at: <https://doi.org/10.1594/PANGAEA.867115> (Amatulli et al., 2018b). Geological permeability from GLHYMPS V2 is available at: <https://doi.org/10.5683/SP2/TTJNU> (Huscroft et al., 2018b). Soil sand content data from SoilGrids is available at: <https://soilgrids.org/> (last access: 10 October 2025).

The daily streamflow data from USGS is available at: <https://waterdata.usgs.gov/nwis> (last access: 10 October 2025). UA SWE dataset: https://climate.arizona.edu/data/UA_SWE/DailyData_4km/ (last access: 10 October 2025). The reservoir storage information is from GAGEII attributes: <https://pubs.usgs.gov/publication/70046617> (last access: 10 October 2025).

Author contributions. YY: initial idea, modeling, analysis, visualization, and writing. MP: initial idea, conceptualization and project administration. YY took the lead in the preparation of the manuscript, but all the authors contributed.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Also, please note that this paper has not received English language copy-editing. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. The authors thank the developer of datasets used in this research for their efforts in creating and sharing valuable resources. Part of the analysis was conducted using Delta, managed by National Center for Supercomputing Applications at University of Illinois Urbana-Champaign.

Financial support. This research has been supported by the Cooperative Institute for Research to Operations in Hydrology (CIROH) with funding under award NA22NWS4320003 from the NOAA Cooperative Institute Program. The statements, findings, conclusions, and recommendations are those of the author(s) and do not necessarily reflect the opinions of NOAA.

Review statement. This paper was edited by Xing Yuan and reviewed by two anonymous referees.

References

- Abdar, M., Pourpanah, F., Hussain, S., Rezazadegan, D., Liu, L., Ghavamzadeh, M., Fieguth, P., Cao, X., Khosravi, A., Acharya, U. R., Makarek, V., and Nahavandi, S.: A review of uncertainty quantification in deep learning: Techniques, applications and challenges, *Inf. Fusion*, 76, 243–297, <https://doi.org/10.1016/j.inffus.2021.05.008>, 2021.
- Amatulli, G., Domisch, S., Tuanmu, M.-N., Parmentier, B., Ranipeta, A., Malczyk, J., and Jetz, W.: A suite of global, cross-scale topographic variables for environmental and biodiversity modeling, *Sci. Data*, 5, 180040, <https://doi.org/10.1038/sdata.2018.40>, 2018a.
- Amatulli, G., Domisch, S., Tuanmu, M.-N., Parmentier, B., Ranipeta, A., Malczyk, J., and Jetz, W.: A suite of global, cross-scale topographic variables for environmental and biodiversity modeling, links to files in GeoTIFF format, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.867115>, 2018b.
- Ayana, Ö., Kanbak, D. F., Kaya Keleş, M., and Turhan, E.: Monthly streamflow prediction and performance comparison of machine learning and deep learning methods, *Acta Geophys.*, 71, 2905–2922, <https://doi.org/10.1007/s11600-023-01023-6>, 2023.
- Baker, S. A., Rajagopalan, B., and Wood, A. W.: Enhancing Ensemble Seasonal Streamflow Forecasts in the Upper Colorado River Basin Using Multi-Model Climate Forecasts, *J. Am. Water Resour. Assoc.*, 57, 906–922, <https://doi.org/10.1111/1752-1688.12960>, 2021.
- Bannister, R. N.: A review of operational methods of variational and ensemble-variational data assimilation, *Q. J. R. Meteorol. Soc.*, 143, 607–633, <https://doi.org/10.1002/qj.2982>, 2017.
- Brown, J. D., Wu, L., He, M., Regonda, S., Lee, H., and Seo, D.-J.: Verification of temperature, precipitation, and streamflow forecasts from the NOAA/NWS Hydrologic Ensemble Forecast Service (HEFS): 1. Experimental design and forcing verification, *J. Hydrol.*, 519, 2869–2889, <https://doi.org/10.1016/j.jhydrol.2014.05.028>, 2014.
- Broxton, P. D., Dawson, N., and Zeng, X.: Linking snowfall and snow accumulation to generate spatial maps of SWE and snow depth, *Earth Space Sci.*, 3, 246–256, <https://doi.org/10.1002/2016EA000174>, 2016.

- Broxton, P. D., Van Leeuwen, W. J. D., Svoma, B. M., Walter, J., and Biederman, J. A.: Subseasonal to seasonal streamflow forecasting in a semiarid watershed, *J. Am. Water Resour. Assoc.*, 59, 1493–1510, <https://doi.org/10.1111/1752-1688.13147>, 2023.
- Cheng, M., Fang, F., Kinouchi, T., Navon, I. M., and Pain, C. C.: Long lead-time daily and monthly streamflow forecasting using machine learning methods, *J. Hydrol.*, 590, 125376, <https://doi.org/10.1016/j.jhydrol.2020.125376>, 2020.
- Clark, S. R., Lerat, J., Perraud, J.-M., and Fitch, P.: Deep learning for monthly rainfall–runoff modelling: a large-sample comparison with conceptual models across Australia, *Hydrol. Earth Syst. Sci.*, 28, 1191–1213, <https://doi.org/10.5194/hess-28-1191-2024>, 2024.
- Cosgrove, B., Gochis, D., Flowers, T., Dugger, A., Ogden, F., Graziano, T., Clark, E., Cabell, R., Casiday, N., Cui, Z., Eicher, K., Fall, G., Feng, X., Fitzgerald, K., Frazier, N., George, C., Gibbs, R., Hernandez, L., Johnson, D., Jones, R., Karsten, L., Ke-feleg, H., Kitzmiller, D., Lee, H., Liu, Y., Mashriqui, H., Mat-tern, D., McCluskey, A., McCreight, J. L., McDaniel, R., Midek-isa, A., Newman, A., Pan, L., Pham, C., RafieeiNasab, A., Ras-mussen, R., Read, L., Rezaeianzadeh, M., Salas, F., Sang, D., Sampson, K., Schneider, T., Shi, Q., Sood, G., Wood, A., Wu, W., Yates, D., Yu, W., and Zhang, Y.: NOAA's National Water Model: Advancing operational hydrology through continental-scale modeling, *J. Am. Water Resour. Assoc.*, 60, 247–272, <https://doi.org/10.1111/1752-1688.13184>, 2024.
- Dalkilic, H. Y., Kumar, D., Samui, P., Dixon, B., Yesilyurt, S. N., and Katipoğlu, O. M.: Application of deep learning approaches to predict monthly stream flows, *Environ. Monit. Assess.*, 195, 705, <https://doi.org/10.1007/s10661-023-11331-5>, 2023.
- Dawson, N., Broxton, P., and Zeng, X.: A New Snow Density Parameterization for Land Data Initialization, *J. Hydrometeorol.*, 18, 197–207, <https://doi.org/10.1175/JHM-D-16-0166.1>, 2017.
- Demargne, J., Wu, L., Regonda, S. K., Brown, J. D., Lee, H., He, M., Seo, D.-J., Hartman, R., Herr, H. D., Fresch, M., Schaake, J., and Zhu, Y.: The Science of NOAA's Operational Hydrologic Ensemble Forecast Service, *Bull. Am. Meteorol. Soc.*, 95, 79–98, <https://doi.org/10.1175/BAMS-D-12-00081.1>, 2014.
- Eghbali, H. J.: K-S Test for Detecting Changes from Land-sat Imagery Data, *IEEE Trans. Syst. Man Cybern.*, 9, 17–23, <https://doi.org/10.1109/TSMC.1979.4310069>, 1979.
- Falcone, J. A.: GAGES-II: Geospatial Attributes of Gages for Evaluating Streamflow, U.S. Geological Survey, <https://doi.org/10.5066/P96CPHOT>, 2011.
- Falcone, J. A., Carlisle, D. M., Wolock, D. M., and Meador, M. R.: GAGES: A stream gage database for evaluating natural and altered flow conditions in the conterminous United States, *Ecol-ogy*, 91, 621–621, <https://doi.org/10.1890/09-0889.1>, 2010.
- Fang, K., Kifer, D., Lawson, K., and Shen, C.: Evaluating the Potential and Challenges of an Uncertainty Quantification Method for Long Short-Term Memory Models for Soil Moisture Predictions, *Water Resour. Res.*, 56, e2020WR028095, <https://doi.org/10.1029/2020WR028095>, 2020.
- Fang, K., Shen, C., and Feng, D.: mhpi/hydroDL: MHPI-hydroDL, Zenodo [code], <https://doi.org/10.5281/zenodo.5015120>, 2021.
- Feng, D., Fang, K., and Shen, C.: Enhancing streamflow forecast and extracting insights using long-short term memory networks with data integration at continental scales, *Water Resour. Res.*, 56, e2019WR026793, <https://doi.org/10.1029/2019WR026793>, 2020.
- Feng, D., Lawson, K., and Shen, C.: Mitigating Prediction Error of Deep Learning Streamflow Models in Large Data-Sparse Regions With Ensemble Modeling and Soft Data, *Geophys. Res. Lett.*, 48, e2021GL092999, <https://doi.org/10.1029/2021GL092999>, 2021.
- Feng, D., Beck, H., de Bruijn, J., Sahu, R. K., Satoh, Y., Wada, Y., Liu, J., Pan, M., Lawson, K., and Shen, C.: Deep dive into hydrologic simulations at global scale: harnessing the power of deep learning and physics-informed differentiable models (δ HBV-globe1.0-hydroDL), *Geosci. Model Dev.*, 17, 7181–7198, <https://doi.org/10.5194/gmd-17-7181-2024>, 2024.
- Fleming, S. W. and Goodbody, A. G.: A Machine Learning Metasystem for Robust Probabilistic Nonlinear Regression-Based Forecasting of Seasonal Water Availability in the US West, *IEEE Access*, 7, 119943–119964, <https://doi.org/10.1109/ACCESS.2019.2936989>, 2019.
- Fleming, S. W., Garen, D. C., Goodbody, A. G., McCarthy, C. S., and Landers, L. C.: Assessing the new Natural Resources Conservation Service water supply forecast model for the American West: A challenging test of explainable, automated, ensemble artificial intelligence, *J. Hydrol.*, 602, 126782, <https://doi.org/10.1016/j.jhydrol.2021.126782>, 2021.
- Fleming, S. W., Rittger, K., Oaida Taghialatela, C. M., and Graczyk, I.: Leveraging Next-Generation Satellite Remote Sensing-Based Snow Data to Improve Seasonal Water Supply Predictions in a Practical Machine Learning-Driven River Forecast System, *Water Resour. Res.*, 60, e2023WR035785, <https://doi.org/10.1029/2023WR035785>, 2024.
- Frame, J. M., Kratzert, F., Klotz, D., Gauch, M., Shalev, G., Gilon, O., Qualls, L. M., Gupta, H. V., and Nearing, G. S.: Deep learning rainfall–runoff predictions of extreme events, *Hydrol. Earth Syst. Sci.*, 26, 3377–3392, <https://doi.org/10.5194/hess-26-3377-2022>, 2022.
- Franz, K. J., Hogue, T. S., Barik, M., and He, M.: Assessment of SWE data assimilation for ensemble streamflow predictions, *J. Hydrol.*, 519, 2737–2746, <https://doi.org/10.1016/j.jhydrol.2014.07.008>, 2014.
- Fuster, B., Sánchez-Zapero, J., Camacho, F., García-Santos, V., Verger, A., Lacaze, R., Weiss, M., Baret, F., and Smets, B.: Quality Assessment of PROBA-V LAI, fAPAR and fCOVER Collection 300 m Products of Copernicus Global Land Service, *Remote Sens.*, 12, 1017, <https://doi.org/10.3390/rs12061017>, 2020.
- Garen, D. C.: Improved Techniques in Regression-Based Streamflow Volume Forecasting, *J. Water Resour. Plan. Manag.*, 118, 654–670, [https://doi.org/10.1061/\(ASCE\)0733-9496\(1992\)118:6\(654\)](https://doi.org/10.1061/(ASCE)0733-9496(1992)118:6(654)), 1992.
- Gauch, M., Mai, J., and Lin, J.: The proper care and feeding of CAMELS: How limited training data affects streamflow prediction, *Environ. Model. Softw.*, 135, 104926, <https://doi.org/10.1016/j.envsoft.2020.104926>, 2021.
- Gericke, O. J. and Smithers, J. C.: Review of methods used to estimate catchment response time for the purpose of peak discharge estimation, *Hydrol. Sci. J.*, 59, 1935–1971, <https://doi.org/10.1080/02626667.2013.866712>, 2014.
- Gichamo, T. Z. and Tarboton, D. G.: Ensemble Streamflow Forecasting Using an Energy Balance Snowmelt Model Coupled to a Distributed Hydrologic Model with Assimilation of Snow and

- Streamflow Observations, *Water Resour. Res.*, 55, 10813–10838, <https://doi.org/10.1029/2019WR025472>, 2019.
- Greff, K., Srivastava, R. K., Koutník, J., Steunebrink, B. R., and Schmidhuber, J.: LSTM: A search space odyssey, *IEEE Trans. Neural Netw. Learn. Syst.*, 28, 2222–2232, <https://doi.org/10.1109/TNNLS.2016.2582924>, 2016.
- Hargreaves, G. H.: Defining and Using Reference Evapotranspiration, *J. Irrig. Drain. Eng.*, 120, 1132–1139, [https://doi.org/10.1061/\(ASCE\)0733-9437\(1994\)120:6\(1132\)](https://doi.org/10.1061/(ASCE)0733-9437(1994)120:6(1132)), 1994.
- Hengl, T., Mendes de Jesus, J., Heuvelink, G. B. M., Ruiperez Gonzalez, M., Kilibarda, M., Blagotić, A., Shangguan, W., Wright, M. N., Geng, X., Bauer-Marschallinger, B., Guevara, M. A., Vargas, R., MacMillan, R. A., Batjes, N. H., Leenaars, J. G. B., Ribeiro, E., Wheeler, I., Mantel, S., and Kempen, B.: SoilGrids250m: Global gridded soil information based on machine learning, *PloS One*, 12, e0169748, <https://doi.org/10.1371/journal.pone.0169748>, 2017.
- Hochreiter, S. and Schmidhuber, J.: Long short-term memory, *Neural Comput.*, 9, 1735–1780, <https://doi.org/10.1162/neco.1997.9.8.1735>, 1997.
- Hunt, K. M. R., Matthews, G. R., Pappenberger, F., and Prudhomme, C.: Using a long short-term memory (LSTM) neural network to boost river streamflow forecasts over the western United States, *Hydrol. Earth Syst. Sci.*, 26, 5449–5472, <https://doi.org/10.5194/hess-26-5449-2022>, 2022.
- Huscroft, J., Gleeson, T., Hartmann, J., and Börker, J.: Compiling and Mapping Global Permeability of the Unconsolidated and Consolidated Earth: GLobal HYdrogeology MaPS 2.0 (GLHYMPS 2.0), *Geophys. Res. Lett.*, 45, 1897–1904, <https://doi.org/10.1002/2017GL075860>, 2018a.
- Huscroft, J., Gleeson, T., Hartmann, J. and Börker, J.: Compiling and mapping global permeability of the unconsolidated and consolidated Earth: GLobal HYdrogeology MaPS 2.0 (GLHYMPS 2.0), *Borealis*, V1 [data set], <https://doi.org/10.5683/SP2/TTJNIU>, 2018b.
- Jiang, S., Zheng, Y., Wang, C., and Babovic, V.: Uncovering Flooding Mechanisms Across the Contiguous United States Through Interpretive Deep Learning on Representative Catchments, *Water Resour. Res.*, 58, e2021WR030185, <https://doi.org/10.1029/2021WR030185>, 2022.
- Khoshkalam, Y., Rousseau, A. N., Rahmani, F., Shen, C., and Abbasnezhadi, K.: Applying transfer learning techniques to enhance the accuracy of streamflow prediction produced by long Short-term memory networks with data integration, *J. Hydrol.*, 622, 129682, <https://doi.org/10.1016/j.jhydrol.2023.129682>, 2023.
- Kling, H., Fuchs, M., and Paulin, M.: Runoff conditions in the upper Danube basin under an ensemble of climate change scenarios, *J. Hydrol.*, 424–425, 264–277, <https://doi.org/10.1016/j.jhydrol.2012.01.011>, 2012.
- Klotz, D., Kratzert, F., Gauch, M., Keefe Sampson, A., Brandstetter, J., Klambauer, G., Hochreiter, S., and Nearing, G.: Uncertainty estimation with deep learning for rainfall–runoff modeling, *Hydrol. Earth Syst. Sci.*, 26, 1673–1693, <https://doi.org/10.5194/hess-26-1673-2022>, 2022.
- Koster, R. D., Mahanama, S. P. P., Livneh, B., Lettenmaier, D. P., and Reichle, R. H.: Skill in streamflow forecasts derived from large-scale estimates of soil moisture and snow, *Nat. Geosci.*, 3, 613–616, <https://doi.org/10.1038/ngeo944>, 2010.
- Kratzert, F., Klotz, D., Brenner, C., Schulz, K., and Herrnegger, M.: Rainfall–runoff modelling using Long Short-Term Memory (LSTM) networks, *Hydrol. Earth Syst. Sci.*, 22, 6005–6022, <https://doi.org/10.5194/hess-22-6005-2018>, 2018.
- Kratzert, F., Klotz, D., Herrnegger, M., Sampson, A. K., Hochreiter, S., and Nearing, G. S.: Toward Improved Predictions in Ungauged Basins: Exploiting the Power of Machine Learning, *Water Resour. Res.*, 55, 11344–11354, <https://doi.org/10.1029/2019WR026065>, 2019a.
- Kratzert, F., Klotz, D., Shalev, G., Klambauer, G., Hochreiter, S., and Nearing, G.: Towards learning universal, regional, and local hydrological behaviors via machine learning applied to large-sample datasets, *Hydrol. Earth Syst. Sci.*, 23, 5089–5110, <https://doi.org/10.5194/hess-23-5089-2019>, 2019b.
- Le, X.-H., Ho, H. V., Lee, G., and Jung, S.: Application of Long Short-Term Memory (LSTM) Neural Network for Flood Forecasting, *Water*, 11, 1387, <https://doi.org/10.3390/w11071387>, 2019.
- LeCun, Y., Bengio, Y., and Hinton, G.: Deep learning, *Nature*, 521, 436–444, <https://doi.org/10.1038/nature14539>, 2015.
- Lees, T., Buechel, M., Anderson, B., Slater, L., Reece, S., Coxon, G., and Dadson, S. J.: Benchmarking data-driven rainfall–runoff models in Great Britain: a comparison of long short-term memory (LSTM)-based models with four lumped conceptual models, *Hydrol. Earth Syst. Sci.*, 25, 5517–5534, <https://doi.org/10.5194/hess-25-5517-2021>, 2021.
- Li, D., Wrzesien, M. L., Durand, M., Adam, J., and Lettenmaier, D. P.: How much runoff originates as snow in the western United States, and how will that change in the future?, *Geophys. Res. Lett.*, 44, 6163–6172, <https://doi.org/10.1002/2017GL073551>, 2017.
- Li, Z., Gao, S., Chen, M., Gourley, J. J., Liu, C., Prein, A. F., and Hong, Y.: The conterminous United States are projected to become more prone to flash floods in a high-end emissions scenario, *Commun. Earth Environ.*, 3, 1–9, <https://doi.org/10.1038/s43247-022-00409-6>, 2022.
- Mangukiy, N. K. and Sharma, A.: Deep Learning-Based Approach for Enhancing Streamflow Prediction in Watersheds With Aggregated and Intermittent Observations, *Water Resour. Res.*, 61, e2024WR037331, <https://doi.org/10.1029/2024WR037331>, 2025.
- Mangukiy, N. K., Sharma, A., and Shen, C.: How to enhance hydrological predictions in hydrologically distinct watersheds of the Indian subcontinent?, *Hydrol. Process.*, 37, e14936, <https://doi.org/10.1002/hyp.14936>, 2023.
- Modi, P., Jennings, K., Kasprzyk, J., Small, E., Wobus, C., and Livneh, B.: Using Deep Learning in Ensemble Streamflow Forecasting: Exploring the Predictive Value of Explicit Snowpack Information, *J. Adv. Model. Earth Syst.*, 17, e2024MS004582, <https://doi.org/10.1029/2024MS004582>, 2025.
- Mote, P. W., Li, S., Lettenmaier, D. P., Xiao, M., and Engel, R.: Dramatic declines in snowpack in the western US, *Npj Clim. Atmospheric Sci.*, 1, 2, <https://doi.org/10.1038/s41612-018-0012-1>, 2018.
- Musselman, K. N., Addor, N., Vano, J. A., and Molotch, N. P.: Winter melt trends portend widespread declines in snow water resources, *Nat. Clim. Change*, 11, 418–424, <https://doi.org/10.1038/s41558-021-01014-9>, 2021.

- Nearing, G., Yatheendradas, S., Crow, W., Zhan, X., Liu, J., and Chen, F.: The Efficiency of Data Assimilation, *Water Resour. Res.*, 54, 6374–6392, <https://doi.org/10.1029/2017WR020991>, 2018.
- Nearing, G. S., Klotz, D., Frame, J. M., Gauch, M., Gilon, O., Kratzert, F., Sampson, A. K., Shalev, G., and Nevo, S.: Technical note: Data assimilation and autoregression for using near-real-time streamflow observations in long short-term memory networks, *Hydrol. Earth Syst. Sci.*, 26, 5493–5513, <https://doi.org/10.5194/hess-26-5493-2022>, 2022.
- Nearing, G., Cohen, D., Dube, V., Gauch, M., Gilon, O., Harrigan, S., Hassidim, A., Klotz, D., Kratzert, F., Metzger, A., Nevo, S., Pappenberger, F., Prudhomme, C., Shalev, G., Shenzen, S., Tekalign, T. Y., Weitzner, D., and Matias, Y.: Global prediction of extreme floods in ungauged watersheds, *Nature*, 627, 559–563, <https://doi.org/10.1038/s41586-024-07145-1>, 2024.
- Ouyang, W., Lawson, K., Feng, D., Ye, L., Zhang, C., and Shen, C.: Continental-scale streamflow modeling of basins with reservoirs: Towards a coherent deep-learning-based strategy, *J. Hydrol.*, 599, 126455, <https://doi.org/10.1016/j.jhydrol.2021.126455>, 2021.
- Painter, T. H., Berisford, D. F., Boardman, J. W., Bormann, K. J., Deems, J. S., Gehrke, F., Hedrick, A., Joyce, M., Laidlaw, R., Marks, D., Mattmann, C., McGurk, B., Ramirez, P., Richardson, M., Skiles, S. M., Seidel, F. C., and Winstral, A.: The Airborne Snow Observatory: Fusion of scanning lidar, imaging spectrometer, and physically-based modeling for mapping snow water equivalent and snow albedo, *Remote Sens. Environ.*, 184, 139–152, <https://doi.org/10.1016/j.rse.2016.06.018>, 2016.
- Pan, M.: CW3E 1-km 1-hourly meteorological forcing on NWM grid. Center for Western Weather and Water Extremes (CW3E), Zenodo [data set], <https://doi.org/10.5281/zenodo.14714512>, 2025.
- Perkins, T. R., Pagano, T. C., and Garen, D. C.: Innovative operational seasonal water supply forecasting technologies, *J. Soil Water Conserv.*, 64, 15A–17A, <https://doi.org/10.2489/jswc.64.1.15A>, 2009.
- Pierce, D. W., Barnett, T. P., Hidalgo, H. G., Das, T., Bonfils, C., Santer, B. D., Bala, G., Dettlinger, M. D., Cayan, D. R., Mirin, A., Wood, A. W., and Nozawa, T.: Attribution of Declining Western U.S. Snowpack to Human Effects, *J. Clim.*, 21, 6425–6444, <https://doi.org/10.1175/2008JCLI2405.1>, 2008.
- Prasad, R., Deo, R. C., Li, Y., and Maraseni, T.: Input selection and performance optimization of ANN-based streamflow forecasts in the drought-prone Murray Darling Basin region using IIS and MODWT algorithm, *Atmospheric Res.*, 197, 42–63, <https://doi.org/10.1016/j.atmosres.2017.06.014>, 2017.
- Sabzipour, B., Arsenault, R., Troin, M., Martel, J.-L., Brissette, F., Brunet, F., and Mai, J.: Comparing a long short-term memory (LSTM) neural network with a physically-based hydrological model for streamflow forecasting over a Canadian catchment, *J. Hydrol.*, 627, 130380, <https://doi.org/10.1016/j.jhydrol.2023.130380>, 2023.
- Saharia, M., Kirstetter, P.-E., Vergara, H., Gourley, J. J., Hong, Y., and Giroud, M.: Mapping Flash Flood Severity in the United States, *J. Hydrometeorol.*, 18, 397–411, <https://doi.org/10.1175/JHM-D-16-0082.1>, 2017.
- Schmidhuber, J.: Deep learning in neural networks: An overview, *Neural Netw.*, 61, 85–117, <https://doi.org/10.1016/j.neunet.2014.09.003>, 2015.
- Shen, C.: A Transdisciplinary Review of Deep Learning Research and Its Relevance for Water Resources Scientists, *Water Resour. Res.*, 54, 8558–8593, <https://doi.org/10.1029/2018WR022643>, 2018.
- Shen, C., Applying, A. P., Gentine, P., Bandai, T., Gupta, H., Tarkovsky, A., Baity-Jesi, M., Fenicia, F., Kifer, D., Li, L., Liu, X., Ren, W., Zheng, Y., Harman, C. J., Clark, M., Farthing, M., Feng, D., Kumar, P., Aboelyazeed, D., Rahmani, F., Song, Y., Beck, H. E., Bindas, T., Dwivedi, D., Fang, K., Höge, M., Rackauckas, C., Mohanty, B., Roy, T., Xu, C., and Lawson, K.: Differentiable modelling to unify machine learning and physical models for geosciences, *Nat. Rev. Earth Environ.*, 1–16, <https://doi.org/10.1038/s43017-023-00450-9>, 2023.
- Shukla, S. and Lettenmaier, D. P.: Seasonal hydrologic prediction in the United States: understanding the role of initial hydrologic conditions and seasonal climate forecast skill, *Hydrol. Earth Syst. Sci.*, 15, 3529–3538, <https://doi.org/10.5194/hess-15-3529-2011>, 2011.
- Smirnov, N.: Table for Estimating the Goodness of Fit of Empirical Distributions, *Ann. Math. Stat.*, 19, 279–281, <https://doi.org/10.1214/aoms/1177730256>, 1948.
- Snyder, C., Bengtsson, T., Bickel, P., and Anderson, J.: Obstacles to High-Dimensional Particle Filtering, *Mon. Weather Rev.*, 136, 4629–4640, <https://doi.org/10.1175/2008MWR2529.1>, 2008.
- Song, Y., Tsai, W.-P., Gluck, J., Rhoades, A., Zarzycki, C., McCrary, R., Lawson, K., and Shen, C.: LSTM-Based Data Integration to Improve Snow Water Equivalent Prediction and Diagnose Error Sources, *J. Hydrometeorol.*, 25, 223–237, <https://doi.org/10.1175/JHM-D-22-0220.1>, 2024.
- Thapa, S., Zhao, Z., Li, B., Lu, L., Fu, D., Shi, X., Tang, B., and Qi, H.: Snowmelt-Driven Streamflow Prediction Using Machine Learning Techniques (LSTM, NARX, GPR, and SVR), *Water*, 12, 1734, <https://doi.org/10.3390/w12061734>, 2020.
- Trujillo, E. and Molotch, N. P.: Snowpack regimes of the Western United States, *Water Resour. Res.*, 50, 5611–5623, <https://doi.org/10.1002/2013WR014753>, 2014.
- Wood, A. W., Hopson, T., Newman, A., Brekke, L., Arnold, J., and Clark, M.: Quantifying Streamflow Forecast Skill Elasticity to Initial Condition and Climate Prediction Skill, *J. Hydrometeorol.*, 17, 651–668, <https://doi.org/10.1175/JHM-D-14-0213.1>, 2016.
- Yao, Y., Zhao, Y., Li, X., Feng, D., Shen, C., Liu, C., Kuang, X., and Zheng, C.: Can transfer learning improve hydrological predictions in the alpine regions?, *J. Hydrol.*, 625, 130038, <https://doi.org/10.1016/j.jhydrol.2023.130038>, 2023.
- Yang, Y., Feng, D., Beck, H. E., Hu, W., Abbas, A., Sengupta, A., Delle Monache, L., Hartman, R., Lin, P., Shen, C., and Pan, M.: Global Daily Discharge Estimation Based on Grid Long Short-Term Memory (LSTM) Model and River Routing, *Water Resour. Res.*, 61, e2024WR039764, <https://doi.org/10.1029/2024WR039764>, 2025.
- Yaseen, Z. M., El-shafie, A., Jaafar, O., Afan, H. A., and Sayl, K. N.: Artificial intelligence based models for streamflow forecasting: 2000–2015, *J. Hydrol.*, 530, 829–844, <https://doi.org/10.1016/j.jhydrol.2015.10.038>, 2015.

Zeng, X., Broxton, P., and Dawson, N.: Snowpack Change From 1982 to 2016 Over Conterminous United States, *Geophys. Res. Lett.*, 45, 12940–12947, <https://doi.org/10.1029/2018GL079621>, 2018.