



# How montane forests shape snow cover dynamics across the central European Alps

Vincent Haagmans<sup>1,2</sup>, Giulia Mazzotti<sup>1,3,4</sup>, Clare Webster<sup>1,5,6</sup>, and Tobias Jonas<sup>1</sup>

<sup>1</sup>WSL Institute for Snow and Avalanche Research SLF, Davos, Switzerland

<sup>2</sup>Department of Civil, Environmental and Geomatic Engineering, ETH Zürich, Zürich, Switzerland

<sup>3</sup>Université Grenoble Alpes, INRAE, CNRS, IRD, Grenoble INP, IGE, Grenoble, France

<sup>4</sup>Université Grenoble Alpes, Université de Toulouse, Météo-France, CNRS, CNRM, Centre d'Études de la Neige, St. Martin d'Hères, France

<sup>5</sup>Department of Geosciences, University of Oslo, Oslo, Norway

<sup>6</sup>Department of Geography, University of Zurich, Zürich, Switzerland

**Correspondence:** Vincent Haagmans (vincent.haagmans@slf.ch)

Received: 7 August 2025 – Discussion started: 6 October 2025

Revised: 5 January 2026 – Accepted: 12 January 2026 – Published: 30 March 2026

**Abstract.** A substantial fraction of seasonal snow is stored in mid-latitude montane forests, serving as an essential temporal water storage. Across vast areas, snow cover dynamics result from processes controlled by forest structure, topography, climate, and weather variability. As data availability has limited our ability to disentangle how these four key controls interact across landscapes within complex topography, most forest snow studies have focused on only one or two of these controls. In this study, we employed the process-based FSM2oshd forest snow model framework for an in-depth analysis of the current state of forest snow water resources across the central European Alps. Over the 8 years analysed, forest snow accounted for 20%–30% of total snow storage in midwinter. In the various effects of existing forest cover on snow, pronounced differences were found depending on elevation, aspect, region, and year. While the presence of forest usually led to a decrease in peak SWE, it decelerated snowmelt, often leading to a later snow disappearance date, particularly on south-facing slopes. However, variability between years and regions was strong enough to shift or even reverse such trends, with snow-scarce years accentuating relative differences in the effects of forests on snow cover. With forest disturbances projected to increase and snow storage to further decline, enhanced complexity of snow cover dynamics in montane forests is to be expected. This places greater emphasis on understanding how the effects of key controls such as forest structure, topography, and weather interact.

## 1 Introduction

Forest and seasonal snow co-occur on about 20% of the surface area of mid-latitude mountain ranges (Deschamps-Berger et al., 2025), which serve as essential temporal water storage for downstream populated regions (Immerzeel et al., 2020; Viviroli et al., 2007). In the European Alps, for example, snowmelt is the primary seasonal runoff signal (Beniston, 2012) and accounts for up to 40% of annual runoff in Switzerland (BAFU, 2021). Therefore, understanding how forest impacts snow cover across mountain ranges is important for water resource and forest resource management (Dickerson-Lange et al., 2021, 2023), streamflow forecasting (Pomeroy et al., 2012; Sun et al., 2018; Troendle, 1983), as well as ecological processes (Cooper et al., 2020; Sanmiguel-Vallelado et al., 2021).

During the snow accumulation period, interception of snow by forest canopies (Hedstrom and Pomeroy, 1998; Roth and Nolin, 2019) decreases below-canopy snow depth (Dharmadasa et al., 2023; Mazzotti et al., 2019a; Moeser et al., 2015) and snow water equivalent (Harestad and Bunnell, 1981; López-Moreno and Latron, 2008; López-Moreno and Stähli, 2008; Stähli and Gustafsson, 2006). As winter progresses, snow ablation is influenced by the presence of forest canopies, which mask the forest floor from shortwave radiation and enhance longwave radiation (Essery et al., 2008; Link et al., 2004; Lundquist et al., 2013; Mazzotti et al.,

2019b; Sicart et al., 2004). This usually results in less radiative energy reaching the forest floor compared to open areas (Link et al., 2004; Strasser et al., 2011). Reduced wind speeds in forests also limit the magnitude of turbulent fluxes (Conway et al., 2018; Price, 1988; Roth and Nolin, 2017). These processes occur wherever forests and snow coexist, but their relative magnitudes and net effects on forest snow resources vary depending on forest structure, topography, weather, and climate (Lundquist et al., 2013; Roth and Nolin, 2017; Safa et al., 2021).

Forest structure exerts first-order control on the strength of most forest-snow processes at the local site scale, whereas climate exerts first-order control between regions (e.g., Broxton et al., 2020). Forest structure is highly heterogeneous, and the spatial scales at which different forest-snow processes act strongly vary. This results in spatiotemporal patterns of forest-snow processes that are highly complex at scales from the individual tree crown (meter; Mazzotti et al., 2019a) to the site/plot scale (meter-hectare; Koutantou et al., 2022) and beyond (meter-valley; Mazzotti et al., 2023). Furthermore, local climatic conditions influence the likelihood that different processes dominate the spatial variability of snow cover within a specific region. For example, denser canopies will intercept more snow than sparse canopies (Moeser et al., 2015). At the same time, the efficiency of canopy interception depends strongly on, but is not limited to, air temperature and wind speed (Cebulski and Pomeroy, 2025; Katsushima et al., 2023; Lundquist et al., 2021). Next, melting processes are primarily driven by incoming shortwave radiation to the snow surface, which is higher in the open and within sparser canopies (Mazzotti et al., 2019b). The effects of shortwave radiation on canopy heating and emitted longwave radiation are also more pronounced in sparse canopies exposed to direct shortwave radiation (Haagmans et al., 2025; Webster et al., 2017). The strengths of these radiative differences are subsequently more pronounced in mid-latitude compared to higher-latitude environments (Seyednasrollah and Kumar, 2019).

Apart from prevailing climatic conditions, variability in meteorological conditions from one winter to the next also affects the evolution of (forest) snow cover patterns. This interannual variability affects snow accumulation and ablation processes differently in forests compared to open areas, with the most substantial differences in snow storage between forested and open sites typically observed during warm and wet winters (López-Moreno and Stähli, 2008). In contrast, these differences are usually smaller during cold and dry winters (Strasser et al., 2011). Meanwhile, the variability of forest snow cover increases during prolonged dry spells (López-Moreno and Latron, 2008) and cold, snow-scarce winters (López-Moreno and Stähli, 2008). Interannual variability in meteorological conditions has also been key to explaining snow accumulation and ablation dynamics across opposing forested slopes of a subalpine valley (Mazzotti et al., 2023).

Finally, topographic controls, particularly elevation and aspect, are additional key factors contributing to the complex differential patterns of snow storage between forests and open areas. At low and mid-elevations, snow can be deeper and persist longer in the open compared to adjacent forest sites. In contrast, forests at higher elevations can maintain snow longer than the adjacent open site (Roth and Nolin, 2017). Topographic aspect has also been observed to exert strong control over differences in snow retention between open and forested sites. Forests on sun-exposed slopes have been found to reduce incoming radiation to the snow surface and delay snowmelt timing relative to adjacent open sites, whereas forests on the opposing shaded slope increase radiation and advance the timing of snowmelt relative to the open site (Ellis et al., 2011). Moreover, consistently deeper snow depths were observed on a forested, shaded slope compared to the opposing sun-exposed forested slope (Koutantou et al., 2022). A detailed modeling study of these opposing sunlit and shaded slopes showed that forest structure, year-to-year weather variability, and topography interact in intricate ways to form complex accumulation and ablation patterns of forest snow (Mazzotti et al., 2023). This is particularly relevant to consider for regions with transitional climates (Dickerson-Lange et al., 2023), such as the European Alps (Bozzoli et al., 2024).

Many forest snow studies have investigated only one or two of the key controls in isolation (i.e., climate, forest structure, year-to-year weather variability, and/or topography) as scale or data availability limited the ability to disentangle how the four key controls interact across landscapes with complex topography (e.g., Ellis et al., 2011; Roth and Nolin, 2017; Seyednasrollah and Kumar, 2019). Especially in the European Alps, studies have been limited to small study sites and did not extend beyond the scale of small Alpine valleys (e.g., Geissler et al., 2023; Mazzotti et al., 2023). The spatiotemporal effects of montane forests on snow water resources, resulting from the combined effects of forest structure, topography, and year-to-year weather variability across different climate zones in Europe, are thus still largely unknown. As snow storage in the European Alps declines (Marty et al., 2017; Matiu et al., 2021), it is increasingly important to understand the different controls on snow water storage across the European Alpine range to predict, manage, and understand the consequences of these declines. Even more so, since past land-use-driven human interferences, as well as natural disturbances such as bark beetles, wildfires, and windthrow, continue to influence forest structure in the European Alps strongly (Bebi et al., 2017).

Past studies of combined controls on forest snow storage have mainly been limited by the lack of spatiotemporally continuous forest snow observations in complex landscapes over the spatial extents required to thoroughly investigate interactions between climate, topography, interannual weather patterns, and forest structure. While airborne laser scanning (ALS) has been employed for repetitive mapping of montane

forest snow cover in complex terrain (e.g., Broxton et al., 2015; Currier et al., 2019; Kostadinov et al., 2019; Mazzotti et al., 2019a; Safa et al., 2021), continuous mapping across entire mountain ranges has yet to be achieved. Where comprehensive observational datasets are missing, advanced process-based models can complement our understanding of forest snow processes (Harpold et al., 2020; Mazzotti et al., 2023). These models have the advantage of providing consistent gapless multi-variable datasets that are impossible to observe over entire mountain ranges.

In this study, we use output from the FSM2oshd model, which is run by the national operational snow hydrological service (OSHD) across Switzerland and adjacent river basins, covering the central European Alps (Mott et al., 2023). FSM2oshd is a mass- and energy-balance-based forest snow model specifically developed to operate over large spatial scales while maintaining process representation and sub-grid parameterization for snow-canopy-atmosphere interactions. A point-based version of the model (FSM2) was used in Mazzotti et al. (2023) at 2 m spatial resolution during six winters to explore interactions between forest structure, topography, and weather in a small Swiss alpine catchment. Upscaling approaches for running FSM2 as the gridded FSM2oshd have been validated by Mazzotti et al. (2021), making it a suitable model choice for this study's purpose. Model output for eight consecutive hydrological years (2017–2024) is used to answer the following research questions (RQ):

**RQ1:** *What is the overall impact of forests on snow storage in the central European Alps?*

**RQ2:** *How do topography and climate across the study region affect forest impacts on seasonal snow dynamics?*

**RQ3:** *How do forest impacts on seasonal snow dynamics vary between years?*

First, the FSM2oshd model framework's capability to accurately represent forest snow cover dynamics at its 250 m resolution is demonstrated across six focus regions. Subsequently, the FSM2oshd output is analyzed and discussed with respect to the above research questions RQ1–RQ3, yielding new insights into forest snow dynamics and its driving factors across large spatiotemporal scales. To the best of our knowledge, such an in-depth analysis of the current state of forest snow water resources in the central European Alps has not been attempted to date.

## 2 Methods

### 2.1 Study area

The study area is situated in central Europe, slightly extending across the Swiss national borders to include the tributaries to all lakes and rivers in Switzerland (Fig. 1). It spans


nearly 58 000 km<sup>2</sup>, with elevations ranging from 184–4806 m above sea level (m a.s.l.). 48.5 % of the area is above 1000 m, and 19.4 % is above 2000 m. The climate on the north side of the Alps is strongly influenced by its proximity to the Atlantic Ocean. In contrast, the Mediterranean Sea primarily influences the south side of the Alps (MeteoSwiss, 2025a).

Across the study area, 41 % is either forested or directly affected by nearby forest presence (e.g., shading). Therein, six focus regions, each approximately 1200 km<sup>2</sup> in size, were defined to analyze forest snow cover dynamics across different climatological conditions and forest types (Fig. 1). These are grouped as follows: the non-Alpine Jura region and the northern Alpine flank (Interlaken, Schwyz regions), which are generally exposed to weather systems originating from the Atlantic, and the more central Alpine regions (Rhône bend, Gotthard, Davos regions) that are exposed to weather systems originating from the Atlantic and the Mediterranean. Characteristics of each focus region are summarized in Table 1.

### 2.2 The Swiss operational snow-hydrological model framework

Based on Mott et al. (2023), we provide a brief overview of the OSHD model framework, focusing on the FSM2oshd snowpack model and its forest snow routines. FSM2oshd provides physics-based simulations of snow and melt distribution at 250 m resolution over the entire study area. It is forced by output from the 1 km numerical weather prediction system COSMO, run by MeteoSwiss (2025b), which is downscaled to 250 m. Moreover, assimilation of data from 444 snow monitoring stations in the study area ensures consistency with available observations of snow height and snow water equivalent (SWE) using methods detailed in Magnusson et al. (2017, 2014) and Cluzet et al. (2024).

The snow cover's mass and energy balances are solved using separate model instances for open, forested, and glacierized areas, while snow is not accumulated on large water bodies that never freeze. Accordingly, grid cells are divided into four corresponding land cover fractions (cf. Fig. 2, Mott et al., 2023), and grid cell-level snow cover properties are computed as weighted averages of these different instances. A forest mask is used to partition grid cells into open areas and forest-covered fractions. Note that open areas influenced by the presence of adjacent forest, i.e., within 20 m distance of forest edges, are included in the forest fraction (see Fig. 2.2–2.3). All key processes through which the forest canopy affects mass and energy exchange between the atmosphere and the sub-canopy snowpack are incorporated in FSM2oshd simulations: canopy interception of snowfall is followed by either sublimation into the atmosphere or unloading onto the ground. The canopy also regulates the transmission of shortwave radiation while enhancing longwave radiation and attenuating wind. Accurate representation of forest snow processes at the 250 m model resolution is



**Figure 1.** Map of the study area with canopy cover fraction ( $f_{veg}$ ) and six focus regions defined to analyze forest snow cover across different climatological conditions and forest types.

**Table 1.** Topographical, land cover, and meteorological characteristics of the six focus regions as outlined in Fig. 1. Meteorological characteristics are based on the model forcings evaluated over the study period and grid cells with forest cover. Topographic characteristics are derived from land cover datasets used for our simulations; see Sect. 2.2 for further information.

	Jura (A)	Interlaken (B)	Schwyz (C)	Rhône bend (D)	Gotthard (E)	Davos (F)
Elevation range (m a.s.l.)	430–1660 m	556–4007 m	418–3049 m	347–3236 m	408–3416 m	430–3309 m
Area below 2000 m	100 %	84.1 %	88.9 %	71.8 %	35 %	43.4 %
Forest cover fraction	21.8 %	45.2 %	20.7 %	26.2 %	29 %	34.4 %
Mean leaf area index (LAI)	2.46	1.75	2.28	2.05	1.71	2.65
Mean canopy cover fraction of forested terrain ( $f_{veg}$ )	0.53	0.41	0.47	0.46	0.42	0.53
Mean $T_{annual}$ in forested terrain	8.33 °C	8.03 °C	7.99 °C	8.16 °C	5.9 °C	5.63 °C
Mean $T_{DJF}$ in forested terrain	0.69 °C	0.83 °C	0.70 °C	0.79 °C	−1.22 °C	−1.53 °C
Mean annual precipitation in forested terrain	1263 mm	1293 mm	1666 mm	1042 mm	1427 mm	1115 mm

achieved by using canopy structure descriptors (e.g., canopy cover fraction  $f_{veg}$ ) and time-varying transmissivities computed at high spatial resolution, which are subsequently averaged over the forest fraction of a grid cell (Mazzotti et al., 2021). For example, shortwave transmission is calculated at a 10 m spacing to account for the effect of heterogeneous canopy structure; the results are then upscaled to the model grid resolution, as described in Mazzotti et al. (2021) and

Webster et al. (2023). Note that the model instance representing forest snow only runs for grid cells that have forest cover, while simulations for open (i.e., non-forested) conditions are calculated for every grid cell. This approach enables a direct assessment of the effect of forest cover relative to open conditions, even in grid cells fully covered by forest. For open areas, the snow cover fraction (SCF) is defined using the algorithm by Helbig et al. (2021). The SCF of forested areas

is parameterized from snow depth using a hyperbolic tangent model (Essery, 2015).

### 2.3 Evaluating modeled forest snow cover dynamics using PlanetScope RGB imagery

The FSM2 model has been extensively validated, particularly with respect to individual forest snow processes (Mazzotti et al., 2020a), hyper-resolution simulations (Mazzotti et al., 2020b, 2023), and its performance in upscaled simulations from 1–50 m (Mazzotti et al., 2021). In this study, we also demonstrate the capabilities of FSM2 at the 250 m resolution (i.e., FSM2oshd) to accurately model forest snow cover across the range of climates and forests in the study area.

Over the European Alps, snow cover retrievals derived from high-resolution (1–5 m) optical satellite imagery are the only distributed and regularly available information for evaluating modeled forest snow across large areas and at scales where forest snow processes vary. The PlanetScope constellation acquires such satellite imagery, providing near-daily coverage of the six focus regions at approximately 3 m resolution (Frazier and Hemingway, 2021). Because PlanetScope acquisitions are collected near the edge of very low Earth orbit at small off-nadir view angles, the resulting images enable direct peering into complex forested terrain, including small forest gaps (Pflug et al., 2024). This study utilized PlanetScope acquisitions, available as image composites, to assess modeled forest snow cover in the focus regions. The composites were selected based on the following criteria: (i) no or minimal cloud cover existed over the evaluated region, (ii) no intercepted snow was visible in the canopy to avoid possible misclassification of snow in the canopy for snow on the ground, and (iii) alignment with three specific evaluation goals related to snow cover dynamics:

- Interannual evaluation: one region, one PlanetScope composite per year acquired around 1 April, over six years.
- Seasonal evaluation: one region, six PlanetScope composites within one year.
- Interregional evaluation: six regions, one PlanetScope composite for approximately the same date (around 1 April)

The data product used in this study is “ortho\_visual”, which is post-processed explicitly for visual analyses (Planet Labs Inc., 2023). Further details are provided in Table 2.

The procedure for evaluating modeled snow cover in FSM2oshd consisted of six steps, the initial four of which are visualized in Fig. 2.

#### Step 1: Prepare a PlanetScope composite for evaluation

For a specific evaluation goal, we selected a suitable PlanetScope composite, ensuring satisfactory quality

based on visual inspection. The corresponding composite was downloaded, and the 250 m FSM2oshd model grid overlaid on it (Fig. 2.1).

#### Step 2: Assign land cover types

Based on land cover information available within the FSM2oshd framework, for each model grid cell in a focus region, we determined whether open areas and/or forests are present. When open areas are present (Fig. 2.2, gray area, inverse of forest mask), the grid cell was assigned to the open land cover mask. If forest cover is present (Fig. 2.2, green area, forest mask), the grid cell was assigned either to the sparse or the dense forest land cover mask based on canopy cover fraction as a proxy of forest cover density (sparse:  $f_{veg} < 0.5$ , dense:  $f_{veg} > 0.5$ , see Fig. 1). Note that this procedure led to double assignments wherever open land cover and either sparse forest or dense forest co-occur in a grid cell (cf. Fig. 2, Mott et al., 2023).

#### Step 3: Estimate snow cover fraction (SCF) for individual grid cells

To the best of our knowledge, no algorithm is currently available to reliably estimate within-forest SCF in PlanetScope retrievals (e.g., in small forest discontinuities). We therefore conducted a manual assessment of SCF. Based on the observable snow cover fraction in a grid cell, a SCF estimate was made by assigning a corresponding SCF class: bare (0), very little (0.05), patchy (0.25), about half (0.5), mostly continuous (0.75), almost continuous (0.95), and continuous (1); see Fig. 2.3. Note that the SCF of grid cells was assessed separately for the different land cover types, and only for that part of a grid cell assigned to the respective land cover type.

Using seven distinct SCF classes was the best trade-off between covering the full range (0–1) of SCFs (i.e., information content) and reliable manual assignment. On average, per PlanetScope composite, SCF was assessed for 630 grid cells for each of the three land cover types, ensuring representation of the full SCF range (0–1) and varying elevations and aspects; see Fig. 2.4 (upper panel). In total, 16 composites and  $\sim 30\,000$  individual grid cells were evaluated (Table 2).

#### Step 4: Derive SCF profiles from the observed SCF of individual grid cells

Estimated SCFs of individual grid cells were binned into 100 m elevation bands for each land cover type to provide a mean SCF value per elevation band while also accounting for aspect (i.e., north-facing ( $> 270^\circ - \leq 90^\circ$ ) or south-facing ( $> 90^\circ - \leq 270^\circ$ )). A PlanetScope composite hence yielded six SCF profiles (see Fig. 2.4, lower panel).

**Table 2.** An overview of the analyzed PlanetScope composites for the focus regions and respective evaluation goals. The number of evaluated grid cells per land cover type is provided for each PlanetScope composite of a focus region. The acquisition time of the images in the composites is between 9 and 11 UTC.

Focus region	Date	Evaluation goal	No. evaluated grid cells with open area land cover	No. evaluated grid cells with sparse forest land cover	No. evaluated grid cells with dense forest land cover
Rhône bend	1 April 2020	Interregional	773	748	625
Interlaken	1 April 2020	Interregional	748	1030	613
Gotthard	3 April 2020	Interregional	564	595	373
Schwyz	1 April 2020	Interregional	743	616	505
Jura	10 February 2022	Interregional	636	819	670
Davos	29 March 2019	Interannual	529	648	576
Davos	7 February 2020	Seasonal	270	268	272
Davos	8 March 2020	Seasonal	454	536	503
Davos	19 March 2020	Seasonal	489	525	515
Davos	1 April 2020	Seasonal, interregional, interannual	652	510	503
Davos	13 April 2020	Seasonal	652	490	578
Davos	23 April 2020	Seasonal	789	503	511
Davos	1 April 2021	Interannual	1008	887	897
Davos	28 March 2022	Interannual	693	678	708
Davos	6 April 2023	Interannual	854	826	789
Davos	6 April 2024	Interannual	772	734	596

**Step 5:** Derive SCF profiles from the modeled SCF of individual grid cells

Here, step 4 was repeated using model data to yield equivalent profiles of modeled SCF for those grid cells assessed in step 3.

**Step 6:** Compare observed SCF to modeled SCF

Finally, the SCF profiles enabled the evaluation of differences between modeled and observed SCF dynamics, separately for each land cover type and by aspect.

## 2.4 Metrics for analyzing modeled forest impact on snow cover dynamics

To isolate the impact of forest on snow cover dynamics in each grid cell and compare it with non-forested conditions, our approach leveraged the different model instances used in the FSM2oshd framework to represent different land cover types (cf. Sect. 2.2). Several metrics were computed to characterize the snow season for each of these two model outputs, as well as for the grid-cell-averaged product accounting for all land cover types. Metrics were calculated over all grid cells and eight hydrological years (HY) from 2017–2024, each starting on 1 September.


First, a qualitative, categorical characterization of the snow season across a region is provided based on snow cover types. Here, we assessed these for all three model outputs and across the whole study area using a simplified classification solely relying on seasonal snow cover duration, adapted from

Sturm et al. (1995), Sturm and Liston (2021), and López-Moreno et al. (2024):

- *Ephemeral snow*: a snow cover that persists for more than one but less than 60 consecutive days.
- *Marginal snow*: a snow cover that persists during 60–120 consecutive days.
- *Seasonal snow*: a snow cover that persists for at least 120 consecutive days but has disappeared by 31 August (end of HY).
- *Perennial snow*: snow cover accumulated over the season persists until 31 August, i.e., through the hydrological year and beyond.

Second, we relied on quantitative descriptors of snow accumulation and melt for further in-depth analyses. These follow Mazzotti et al. (2023) and include:

- *Peak SWE*: the maximum value of snow water equivalent on the ground.
- *Snow disappearance date (SDD)*: the date of the first occurrence of SWE < 10 mm after peak SWE, also referred to as melt-out.
- *Center time of snowmelt runoff (CT)*: the date at which 50 % of the total annual snowmelt runoff (at the grid-cell level) has been released (Stewart et al., 2004).



**Figure 2.** The initial four (of six) steps in the workflow for validating modeled (forest) snow cover fraction (SCF) using PlanetScope RGB imagery. The PlanetScope composite captures the Davos region.

Unless otherwise specified, the analyses in the results section assessed the impact of forest cover in terms of the difference between the model instances representing forested and open land cover types.

### 3 Results

#### 3.1 Evaluation of modeled forest snow cover based on PlanetScope imagery

Figure 3 shows an evaluation of modeled snow cover fraction during spring for six consecutive years in the Davos region. Aspect-dependent SCF differences are apparent. Approximating the snow line by  $SCF = 0.5$ , we found an elevation difference between the snow lines on north- and south-facing aspects of approximately 400 m during this time of the snow season. This elevation difference was similar between PlanetScope acquisitions from different years. However, the absolute snow line elevation varied from year to year, despite observations made at the same time of year. For exam-

ple, in 2019 and 2021, the snow line was 400 m lower than in 2023 and 2024. Remarkably, differences in SCF profiles between the three land cover types (open area, sparse forest, and dense forest) typically remained within  $\sim 50$  m, with some notable exceptions between forested and open areas in 2022 and 2023. All the above findings from observations were generally well represented by the FSM2oshd model, except that in some cases the model overestimated differences between SCF profiles of the three land cover classes, particularly on the north-facing slopes in 2020 and 2024.

Figure A1 illustrates the progression of the snow line with elevation throughout winter and spring of 2020 in the Davos region, representing the seasonal evaluation. Both modeled and observed SCF profiles showed a progressive increase of  $\sim 500$  m in snow line elevation between 8 March and 23 April, while the spread between aspects gradually increased. FSM2oshd could even represent specific features, such as the observed substantial SCF difference between sparse and dense forests on north-facing slopes on 19 March, or the drop in SCF at high elevation in south-facing slopes on 13 and 23 April. Considerable discrepancies were present

only in the 7 February comparison, when the modeled forest SCF was underestimated shortly after a minor snowfall two days before the PlanetScope scene acquisition, due to melt starting too early in the model.

The observation-based SCF profiles (left side of Fig. A2) showed similar patterns across the six focus regions, and differences were generally minor compared to the variability between aspects and years (cf. Fig. 3), the Jura region, however, being an exception. Furthermore, differences in SCF profiles between aspects were notably smaller in the Rhône bend and Jura regions than in the other regions. FSM2oshd provided an excellent match to SCF observations for the Rhône bend, Gotthard, and Davos regions, but a less close match for the Interlaken and Schwyz regions. Lower accuracy in these two cases could also be attributed to a minor snowfall that occurred two days prior to the acquisition of the PlanetScope imagery. This snowfall of a few centimeters was sufficient to make the snow cover appear nearly uniform in the observations, but FSM2oshd did not accurately represent this. The Jura region stands out in that the profiles of observed SCF on south-facing aspects were captured very well by FSM2oshd, whereas modeled SCF on north-facing aspects at elevations of 700–1100 m was significantly lower than the observed values. This again resulted from a prior snowfall event, from which the resulting snow cover persisted longer on the steep, shaded north-facing slopes of the Jura than was modeled.

Mean absolute error (MAE) statistics further confirmed the excellent match between observations and model (see Table A1). The MAEs, averaging at 0.16 SCF, were low and remarkably consistent across PlanetScope composites and land cover types. Slightly larger MAEs ( $\sim 0.25$ ) were only determined for the evaluations of the Interlaken and Schwyz regions on 1 April 2020. Note that this assessment was based on the evaluation of many grid cells for each land cover class within a focus region. On average across all PlanetScope composites, 4% of all grid cells classified as “open area,” 13% of all grid cells with a “sparse forest,” and 15% of all grid cells with a “dense forest” land cover type were evaluated. We thus concluded that FSM2oshd is well capable of representing observed SCF variability between years, aspects, and throughout a winter season, even if in 3 of the 16 composites, some shortcomings of the model’s response to recent snowfall were notable. Therefore, we consider the model results to provide a highly accurate representation of snow cover evolution in both open and forested environments, which constitutes the basis for the subsequent analyses.

### 3.2 Overall impact of forests on seasonal snow cover

Figure 4 summarizes the impact of forests on snow water resources across the study area and for the 2017–2024 hydrological years. The maximum amount of water stored as snow during a hydrological year averaged at 126 mm and ranged


from 87–192 mm (Fig. 4A), corresponding to a total volume of water between 4.8 and 10.6 km<sup>3</sup>. Forested areas had an average peak SWE storage of 56 mm, which during the study period ranged from 28–108 mm, corresponding to a total volume of water between 0.7 and 2.6 km<sup>3</sup> (Fig. 4B). Forest peak SWE typically occurred 1–2 months earlier than total peak SWE.

To quantify the relevance of forest snow water storage, we computed the fraction of SWE stored in forests relative to total SWE (Fig. 4C). This fraction reached an absolute maximum of 0.34 in early winter of December 2021. On average, it peaked at 0.23 in early winter. It then remained relatively stable for about two months before decreasing as total SWE storage continued to increase, whilst forest SWE storage remained approximately constant. The period during which forests stored snow was significantly shorter than the overall SWE storage period in the study area. This difference resulted from the abundance of snow at elevations well above the tree line.

Alternatively, comparing actual snow water storage to a model scenario without forests provided an estimate of snow water storage sensitivity to forest cover in absolute terms (Fig. 4D). If forest cover was unaccounted for over the entire model domain, total SWE storage would, at its peak, increase by only 5 mm on average, which is 3.7% of the current mean total peak SWE (cf. Fig. 4A and D). The increase of the maximum SWE storage, in this scenario, reached 9 mm during HY2021 and occurred 11 d later than the total peak SWE during that year (Fig. 4A). This indicates the potential for prolonged overall storage without trees, as can also be observed by comparing the tails of the SWE curves for forests (Fig. 4B) and no forests (Fig. 4D), respectively.

The impacts of forests on snow cover dynamics outlined above mean that the snow cover type at a given location can also differ depending on whether forest cover is present or not. Using the snow cover type classification detailed in Sect. 2.4, we present a qualitative assessment of forest impacts on seasonal snow in Fig. 5. Over 95% of the land area in the study domain was snow-covered during the study period, for which the dominant snow cover types were ephemeral (50.5%) and seasonal (34.5%), as illustrated in Fig. 5A. Ephemeral snow cover was primarily found in the lowlands below 600 m, while seasonal snow cover was typically present in most of the Alps above 1000 m. Marginal snow cover (14%) occurred mainly above 750 m in the pre-Alps and high alpine valleys, acting as a transition zone between ephemeral and seasonal snow cover. Perennial snow cover (1.5%) closely aligned with glacier cover and was found only above 2500 m. While elevation was the primary factor controlling snow cover type, it also depended on aspect (Fig. 5C); for example, seasonal snow tended to occur approximately 200 m lower on north-facing aspects than on south-facing aspects.

Comparing the snow cover types determined for the model outputs of the forested and open land cover types revealed




**Figure 3.** Comparison of observed (PlanetScope) and modeled (FSM2oshd) snow cover fraction (SCF) for the Davos focus region and the 2019–2024 period. SCF profiles are shown separately for three land cover types (i.e., dense forest, sparse forest, and open area) on north- or south-facing slopes.

where and how forests, in addition to topographic factors, influenced this classification. Transitions between snow cover types due to the presence of forest are mapped out in Fig. 5B. Of all grid cells with forest cover and seasonal snow, 8.5 % had a snow cover duration that was extended by forests: 6 % were classified as marginal in the presence of forest and as ephemeral in its absence, while 2.5 % were seasonal with forest and marginal without. The opposite, i.e., a shortening of the snow cover season, was found in 4.3 % of these grid cells, with 2.5 % classified as ephemeral with forest and marginal

without, and 1.8 % as marginal with and seasonal without forest, respectively.

These transitions between snow cover types occurred across large areas in the Jura and the Pre-Alps. Here, the snow cover type was identified as marginal in the presence of forest but as ephemeral in its absence. The same kind of transition was also found across the Alps on some south-facing slopes above 1000 m. Alpine south-facing aspects above 1500 m more often featured seasonal snow cover when forest was present, but marginal snow cover in the



**Figure 4.** Snow storage evolution during eight hydrological years (HY) for each day of the year (DoY) in terms of snow water equivalent (SWE), summed across the whole study area (A) and for the forested part of the study domain only (B). Ratio of SWE stored in forests to total SWE storage (C), and storage difference if there were no forests (D).

open. On north-facing aspects above 1500 m, snow cover type was commonly marginal in the presence of forest but seasonal in the open. Below this elevation on north-facing aspects, snow cover types were ephemeral and marginal in the presence and absence of forest, respectively. However, considerable variability in snow cover type divergences existed between hydrological years, as visualized in Fig. 5D.

### 3.3 Spatial differences in forest impacts on snow accumulation and ablation dynamics across the study area


To explore how the impacts of forests on snow cover vary spatially across complex topography and climatic gradients, we analyzed how snow cover dynamics descriptors introduced in Sect. 2.4 varied in terms of elevation and aspect across the six focus regions introduced in Sect. 2.1.

Figure 6 presents the mean difference in peak SWE, i.e.,  $\Delta\text{Peak SWE}$ , computed as the peak SWE in the open land cover type minus the peak SWE in the forested land cover types. Note that the same sign convention was applied to all subsequent analyses. In the Alps, peak SWE in open areas was almost everywhere higher than in forested terrain. Notably, a tendency toward greater deficits in forests above  $\sim 1300$  m existed. In contrast, at lower elevations and away from the Alps, peak SWE was slightly higher in forests. Moreover, the three more central Alpine focus regions exhibited aspect dependency, where peak SWE showed a remarkably higher deficit in forests on north-facing slopes (D–F, Fig. 6). Such aspect-dependent patterns were not observed

in the two focus regions on the northern flanks of the Alps (B and C, Fig. 6). In the Jura Mountains (A), the highest elevations exhibited a distinct pattern. Here, forested areas stored slightly higher peak SWE compared to nearby open areas.

Figure 7 presents the mean difference in snow disappearance day between forested and open areas ( $\Delta\text{SDD}$ ). Across the Alps, whether snow persisted longer in forests or open areas was generally dictated by elevation and aspect. Within the five Alpine focus regions (B–F, Fig. 7), snow persisted longer in open areas above  $\sim 1700$ – $1850$  m for 1–5 weeks, regardless of aspect. Below this elevation, snow remained longer in forests on south-facing slopes for a few days to two weeks. Oppositely, on north-facing slopes, snow persisted longer in open areas for a few days to two weeks. Within the (non-Alpine) Jura region, snow persisted longer in the forests for 1–4 weeks, regardless of aspect (A, Fig. 7). These differences in SDD were most pronounced at elevations between 1100 and 1550 m.

Much before the snow cover disappears, the snowpack already generates runoff from melting. The timing of snow accumulation relative to melting determines when precipitation, which is temporally stored as snow, becomes available for runoff generation. To contrast this timing between forested and nearby open areas, we analyzed the date on which 50 % of the snowmelt volume is released (CT) in Fig. 8. We found that, on average, whether CT occurred first in forested or nearby open areas strongly depended on the region, elevation, and aspect. Away from the tree line ( $< 1850$  m), the focus regions centered in the Alps (D–F, Fig. 8) exhibited a clear aspect dependence on whether CT



**Figure 5.** Dominant snow cover types across the study area during the 2017–2024 hydrological years (A), for which (C) captures the distribution across elevation and aspect. Transitions between snow cover types due to the presence of forest cover are shown in (B), while (D) captures its variability across aspects and elevations for each year. The four percentage rates shown for each hydrological year indicate the ratio of grid cells that transitioned between snow cover types due to the presence of forest cover.

occurred earlier in forests or in nearby open areas. Here, compared to open areas, forests on south-facing slopes delayed CT by at least a few days and up to two weeks. In contrast, forests on north-facing slopes advanced CT by up to two weeks, a pattern that was also observed more generally above 1850 m across all regions, regardless of aspect. Within the two focus regions on the northern Alpine flank (B and C, Fig. 8), forests (< 1850 m) in all aspects delayed CT by at least a few days and up to two weeks, while these delays were more pronounced on south-facing slopes. Significantly longer delays of CT by forests of 2–3 weeks were observed in the (non-Alpine) Jura region (A, Fig. 8).


### 3.4 Temporal variations in forest impacts on snow accumulation and melt dynamics

In this Section, we examined variations in forest snow cover dynamics over the eight hydrological years. The differences

in peak SWE, SDD, and CT between forested and nearby open areas in the Davos region were analyzed. The winter conditions for each hydrological year were categorized as either snow-scarce ( $\leq -20\%$  mean peak SWE), average, or snow-rich ( $\geq +20\%$  mean peak SWE).

The difference in peak SWE storage between open areas and forests varied considerably between years and with elevation and aspect (Fig. 9A). Generally, peak SWE storage was higher in open areas, with this difference being more pronounced on north-facing slopes. Snow storage differences consistently increased with elevation up to the tree line. In snow-rich years, differences in peak SWE storage were amplified, while they were minimized during snow-scarce years. Notably, during the 2020 hydrological year, forests on south-facing slopes at lower elevations had higher peak SWE storage.

The day of snow disappearance also exhibited considerable variability across elevations and aspects between years



**Figure 6.** Mean difference in peak SWE between open and forested areas ( $\Delta\text{Peak SWE}$ ) during the 2017–2024 hydrological years. For each of the six focus regions, a polar plot is provided (A–F). These show the average  $\Delta\text{Peak SWE}$  for eight 150 m elevation bands (950–2150 m) across eight aspect classes, where the labels refer to the directions in which slopes face.

(Fig. 9B). In many locations, snow persisted longer in forests in one year but disappeared earlier in another. This variability in  $\Delta\text{SDD}$  could be attributed to winter conditions: during snow-rich winters, snow generally persisted longer in the open, whereas during average and snow-scarce winters, aspect-dependent patterns emerged with snow persisting longer in south-facing forests. Near the tree line ( $> 1850$  m),  $\Delta\text{SDD}$  was less pronounced, and during most winters, snow persisted longer in the open.


Differential patterns of the center time of snowmelt runoff ( $\Delta\text{CT}$ ) showed additional variability that did not necessarily follow patterns of  $\Delta\text{Peak SWE}$  and  $\Delta\text{SDD}$  linked to winter conditions (Fig. 9C). During snow-rich winters, CT occurred later in forests on south-facing slopes for one year (HY2018), while occurring later in the open during another

year (HY2021). Conversely, during snow-scarce and average winters, patterns of  $\Delta\text{SDD}$  and  $\Delta\text{CT}$  showed greater conformity; for example, snow persisting longer in forests generally co-occurred with CT occurring later in forests.

## 4 Discussion

### 4.1 The role of accumulation versus ablation for differential snow cover dynamics

The analysis presented in Sect. 3.3 reveals considerable differences between open and forested terrain in both peak SWE (Fig. 6) and SDD (Fig. 7). The question arises as to whether these differences occur primarily due to accumulation or ablation processes. On south-facing slopes in the Alps, snow




**Figure 7.** Mean difference in snow disappearance day between open and forested areas ( $\Delta$ SDD) during the 2017–2024 hydrological years. For each of the six focus regions, a polar plot is provided (A–F). These show the average  $\Delta$ SDD for eight 150 m elevation bands (950–2150 m) across eight aspect classes, where the labels refer to the directions in which slopes face.

typically persists longer in forests than in the open (B–F, Fig. 7), even though peak SWE is on average lower in these forests (B–F, Fig. 6). Hence, snow persistence in south-facing slopes appears to be mainly a result of slower ablation rates in forests in comparison to the open. However, on north-facing slopes, snow persists longer in the open where peak SWE storage is also higher. Here, it is likely the difference in accumulation that drives the overall effect of forests on snow persistence.

However, spatiotemporal differences in accumulation and ablation, as well as their variations between forested and open terrain, are more intricate than the above description of the overall effect suggests. In Fig. 10, we show that conditions averaged over large scales are driven by the range of grid-cell-to-grid-cell variability, rather than representing

conditions prevailing in any majority of grid cells. To this end, we present an evaluation of the contrasting conditions between south- and north-facing grid cells from only one region (Davos) and one elevation band (1400–1550 m) for HY2022. There is considerable variability in the amount of snow that builds up during the accumulation period in the open (peak  $SWE_{open}$ ), which is primarily due to local differences in concurrent melt, modulated by terrain shading and variable microclimatic conditions. Spatial variability in snowfall is less influential in this regional evaluation (data not shown).


On south-facing slopes, the overall (average) effect of forest amounts to a positive  $\Delta$ Peak SWE (Fig. 10D, dashed line) and a negative  $\Delta$ SDD (Fig. 10E, dashed line) consistent with results shown in Figs. 6F and 7F. However, grid cells that




**Figure 8.** Mean difference in the date on which 50 % of the snowmelt volume is released between open and forested areas ( $\Delta$ CT) during the 2017–2024 hydrological years. For each of the six focus regions, a polar plot is provided (A–F). Polar plots show the  $\Delta$ CT for eight 150 m elevation bands (950–2150 m) across eight aspect classes, where the labels refer to the directions in which slopes face.

contribute to the negative  $\Delta$ SDD are predominantly those that feature a low peak  $SWE_{\text{open}}$  (Fig. 10E, F2, and F3). In contrast, grid cells that contribute to the positive  $\Delta$ Peak SWE are predominantly those that feature a high peak  $SWE_{\text{open}}$  (Fig. 10D and F1). So, despite the overall effect, there may not be a single grid cell that features both a positive  $\Delta$ Peak SWE and a negative  $\Delta$ SDD at the same location. On north-facing slopes, there is less potential for pronounced ablation during accumulation, which is why grid cells with peak  $SWE_{\text{open}}$  values below 100 mm are largely missing. This promotes higher  $\Delta$ Peak SWE (Fig. 10A) and essentially inhibits negative  $\Delta$ SDD values (Fig. 10B and C1–C3), resulting in an overall much higher  $\Delta$ Peak SWE and a positive  $\Delta$ SDD (cf. Figs. 6F and 7F).

These considerations can also be extrapolated to explain differences in  $\Delta$ SDD between years and regions. For instance, HY2020 was a snow-scarce year with low accumulation (i.e., peak  $SWE_{\text{open}}$ ), particularly in the Interlaken and Schwyz regions. This resulted in many grid cells with very negative  $\Delta$ SDD values, even on north-facing slopes, leading to a reduced aspect dependence in both regions (Fig. 7B and C). On the other hand, snow-rich and/or cold years that lead to pronounced  $\Delta$ Peak SWE (Fig. 9A, 2018/2019/2021; Davos region) provide less potential for many grid cells with very negative  $\Delta$ SDD values, hence showing overall neutral or slightly positive  $\Delta$ SDD even on south-facing slopes (cf. Fig. 9B). Finally, in the Jura region, conditions are generally different, with, on average, half the amount of peak SWE compared to all the other regions. This favors conditions sim-



**Figure 9.** Interannual variability of forest snow cover dynamics in the Davos region for eight hydrological years (HY), characterized by differential peak SWE (A), differential snow disappearance day (B), and differential center time of snowmelt runoff (C) between forested and open areas. Winter conditions are categorized as either snow-scarce ( $\leq -20\%$  mean peak SWE), average, or snow-rich ( $\geq +20\%$  mean peak SWE). Polar plots show the respective variable for eight elevation bands, each spanning 150 m (950–2150 m) across eight aspect classes, representing the directions in which slopes face.



**Figure 10.** An evaluation of the contrasting conditions in terms of  $\Delta\text{Peak SWE}$  and  $\Delta\text{SDD}$  relative to values of  $\text{peak SWE}_{\text{open}}$  for all grid cells of the 1400–1550 m elevation band on north-facing (A, B) and south-facing (D, E) aspects for HY2022. Additionally, SWE time series of forested and open fractions are presented for three selected grid cells on north-facing aspects (C1–C3) and south-facing aspects (F1–F3), see colored markers in A, B, D, E, and corresponding line plots in C1–C3 and F1–F3.

ilar to those in panels F2 and F3 of Fig. 10, with negative  $\Delta\text{SDD}$  irrespective of elevation and aspect. In addition, the topography of the Jura Mountains differs from that of the Alps as the highest elevations are plateau-like, i.e., relatively flat.

The combined effects of forest cover, topography, and interannual weather variability on forest snow dynamics, as discussed above, provide further context for past observations by Koutantou et al. (2022), who found significant differences in small-scale snow height distributions between two opposing forested north- and south-facing slopes within the study area. Subsequently, Mazzotti et al. (2023) were able to link these differences to ablation from early-season insolation on south-facing slopes in the absence of topographic shading. By considering additional topography-driven processes and including detailed canopy structure information, both aforementioned studies advanced the conceptual framework of Lundquist et al. (2013), who focused on the com-

parison of forested versus non-forested conditions at the site scale. Here, our results demonstrate that the combined effects of these processes, initially investigated at the scale of individual trees, also affect snow distribution dynamics at much coarser spatial scales and over large extents. The relevance of both forest cover and topography on snow cover dynamics beyond the valley scale has also been noted by Broxton et al. (2020), who found SWE differences to increase between areas with more or less forest cover as winter progressed, influenced by terrain shading or the lack thereof on opposing slopes.

#### 4.2 Forest structure effects on snow storage

Distinct aspect- and elevation-dependent patterns of  $\Delta\text{Peak SWE}$  are apparent in Figs. 6 and 9A. Not surprisingly, these patterns generally match those of leaf area index (LAI, cf. Fig. A3), confirming that canopy snow processes have a con-

siderable impact on snow accumulation on the ground. For example, in the Rhône bend, Gotthard, and Davos regions, LAI is consistently higher on north-facing aspects within the 1100–1700 m range, correlating with increased  $\Delta$ Peak SWE (Fig. 6), regardless of how snow-rich a winter is (Fig. 9A). In the Interlaken and Schwyz regions, there is little aspect dependency of LAI, which explains the absence of corresponding  $\Delta$ Peak SWE patterns (B and C, Fig. 6). Above 1700 m across the study domain, LAI rapidly decreases, explaining why, even though cumulative annual snowfall further increases with elevation, differential peak SWE storage does not. These interrelations demonstrate the role of snow interception, which is known to scale with LAI (Hedstrom and Pomeroy, 1998). The longer and more snow is trapped in the canopy, the more intercepted snow will sublimate back into the atmosphere instead of contributing to snow accumulation on the ground (Lundquist et al., 2021; Pomeroy et al., 1998). Hence, aside from ablation processes, peak SWE should, in first order, decrease with increasing LAI, explaining the correlation between the patterns in Figs. 6 and A3. However, ablation can make a difference. Areas above 1250 m in the Jura region provide an example in which peak SWE is consistently higher within forests (Fig. 6A) due to considerably more snowmelt during the accumulation period in the open, equivalent to the examples shown in Fig. 10F2 and F3. This phenomenon has also been reported during winter storms with rain on snow, in which ablation in forests was limited by lower wind speeds that reduced the magnitude of turbulent exchanges at the snow surface (Marks et al., 1998). Alternatively, slightly higher SWE accumulation in forests could also result from differences in sublimation (Gelfan et al., 2004).

Current forest structures across the Alps resulted from past land-use-driven human interferences, as well as natural disturbances such as bark beetles, wildfires, and windthrow (Bebi et al., 2017). Coniferous forests are prevalent, and this biome is where future changes in forest structure are likely to be most pronounced (Seidl et al., 2017), in turn affecting forest snow dynamics. For example, beetle infestations can reduce LAI and thus interception, thereby increasing snow accumulation on the ground (Boon, 2012). In addition, forests are unevenly altered by wildfires, creating patches of varying burn severity (Koshkin et al., 2022), where an increase in overstory mortality is associated with increased sub-canopy snow accumulation (Maxwell and St Clair, 2019). Hence, especially in forests with dense canopies, local disturbances could lead to an overall increase in peak SWE storage. In the study domain, this applies specifically to north-facing slopes in the central Alps (Rhône bend, Gotthard, and Davos regions, cf. Fig. A3). However, topography also influences how forest disturbances affect snow cover, as both terrain and canopy affect the amount of sunlight reaching the snowpack (Rinehart et al., 2008). On north-facing slopes, sparser forests often increase snow storage due to limited winter insolation and thus ablation (Broxton et al., 2020; Harpold

et al., 2020). In contrast, south-facing slopes receive more sunlight, complicating the effects of disturbances. Denser forests can reduce snow storage via interception and increased longwave radiation (Broxton et al., 2020), while canopy removal may also accelerate ablation and reduce peak SWE (Mazzotti et al., 2023). With forest disturbances projected to increase (Seidl et al., 2017), understanding their effects on snow storage in complex terrain is crucial, though it falls beyond the scope of this study. Currently, most of such research originates from North America, driven by concerns related to water and fire management (e.g., Dickerson-Lange et al., 2023, 2021; Lewis et al., 2023; Moeser et al., 2020; Pomeroy et al., 2012). However, given the sensitivity of snow process interactions to forest structure, climate, and topography, corresponding studies from other regions with different environmental conditions are equally vital.

### 4.3 The hydrological relevance of forest snow

Forest snow can account for more than 30 % of total snow storage, yet surprisingly small differences result when comparing simulations with and without forest. With no forest cover at all, total peak snow storage would only increase by a few percent (Fig. 4). This may seem surprising, considering in how many ways forests influence snow processes. However, although approximately 40 % of the study area is forest-covered or affected by its presence, more than 50 % of this area is located at elevations below 1000 m. This considerably constrains the effect of forests on snow storage at the national level in absolute terms in two ways: at those elevations, unloading of intercepted snow usually happens within a few days, leaving little time for respective sublimation losses. Moreover, most of the total mass is accumulated higher up in the mountains, leaving differences between forested and open areas at lower elevations only a small share. In addition, while peak SWE is increased due to a lack of canopy interception (Fig. 6), pronounced ablation in the open during accumulation on south-facing slopes might offset some of this increase (cf. peak  $SWE_{open}$  in Fig. 10A and D; Sect. 4.1). Lastly, different meteorological conditions in the open (e.g., higher wind speeds) will result in increased sublimation losses from the ground (Strasser et al., 2008). Combined, these factors may explain why only a relatively small increase in total peak SWE storage would result if there were no forests in the study domain.

Even if the effect on total snow storage is relatively small, the presence of forest implies considerable changes to the spatial distribution and temporal evolution of snow cover, as is evidenced by the complex patterns seen in Figs. 6–9, and as discussed above. The increased spatial heterogeneity of forest snow cover is known to disperse the occurrence of snowmelt runoff over time. As a result, downstream rivers can be expected to have a flatter seasonal hydrograph, reducing peak flow (Pomeroy et al., 2012) and decreasing the risk of spring snowmelt flooding (Hendrick et al., 1971). Recent

modeling studies have shown that at the watershed scale in the US, increasing forest heterogeneity leads to increased low streamflows into late summer (Sun et al., 2018). The extent of these runoff changes depends on local climatic conditions and forest structure (Currier et al., 2022).

Our findings confirm spatially heterogeneous effects of forest cover on the timing of snowmelt runoff, which shifts depending on aspect and elevation (Fig. 8). Interestingly, findings in the central Alps (D–F, Fig. 8) are in line with Ellis et al. (2011) who found that forests substantially delayed snowmelt onset on south-facing slopes, whereas on north-facing slopes, snowmelt onset occurred at the same time but progressed much more quickly in forests; this in contrast to other regions where aspect dependent responses were largely missing. Not surprisingly, patterns of  $\Delta CT$  (Fig. 8) corresponded to those of  $\Delta SDD$  (Fig. 7), indicating that dependencies discussed in Sect. 4.1 also have larger-scale impacts on snowmelt runoff generation. Still, other factors, such as hillslope effects, soil characteristics, and soil frost, also affect how heterogeneous snowmelt runoff in montane forests translates into a hydrograph response (Pomeroy et al., 2012; Redding and Devito, 2011).

#### 4.4 Opportunities and limitations of a simulation-based analysis

Using model output data in this study was necessary, as no equivalent measurements exist. Even in areas that undergo periodic ALS acquisitions (Dwivedi et al., 2024; Painter et al., 2016), temporal coverage would likely not be sufficient to reveal the level of detail discussed here (e.g., Sect. 4.1). The approach of using data from physically based forest snow models to analyze effects that are difficult to observe is not new. Exploiting hyper-resolution simulations with FSM2 (2 m), Mazzotti et al. (2023) advanced the understanding of forest snow process interactions in complex terrain at the level of individual mass and energy balance components, providing the basis for this work. Other recent studies have used similar approaches, for example, to highlight the importance of fine-scale forest structure, which governs a range of relevant forest snow process interactions (Broxton et al., 2021). Throughout the past decades, modeling studies have played an important role in shaping today's understanding of both specific forest snow processes (e.g., Pomeroy et al., 1998) and how process interactions in montane forests modulate the subcanopy snow cover (e.g., Pomeroy et al., 2012; Strasser et al., 2011).

Of course, the FSM2oshd model used here, like any model, does not provide perfectly accurate simulations of the actual conditions; however, the model has previously been shown to accurately represent forest snow processes across various sites, years, climates, and spatial resolutions (Mazzotti et al., 2020a, b, 2023). In addition, data assimilation strategies ensured that the model remained consistent with daily snow depth observations from  $\sim 500$  stations

(Cluzet et al., 2024; Mott et al., 2023). Further validation of FSM2oshd simulations in this study aimed to provide additional confidence in the approaches used for upscaling the physical representations of meter-scale forest snow processes to 250 m resolution (cf., Mazzotti et al., 2021). With no ALS data available, PlanetScope satellite imagery proved an effective means of validating FSM2oshd over such a large area at the necessary spatial resolution. This demonstrated the ability of FSM2oshd to capture complex patterns of spatiotemporal snow cover dynamics (Figs. 3, A1, and A2), expressed by a persistently low MAE (Table A1), which also attests to the capabilities of the methodology used for ingesting three-dimensional forest structure data (Mott et al., 2023; Webster et al., 2023).

While FSM2oshd can represent lateral redistribution processes (Quéno et al., 2024), these process representations are effective only at higher spatial resolutions and would require grid spacing of 100 m or finer. Here, we prioritized a large spatial extent over the high resolution required to include redistribution processes. Including snow redistribution processes might have yielded additional insights, even though subalpine forests in the Alps are typically surrounded by narrow, steep terrain, which provides shelter from strong winds. Nevertheless, we acknowledge that the model setup used here may underpredict sublimation losses of snow in open terrain, particularly in wind-exposed settings at high elevation, i.e., above the tree line.


## 5 Conclusions

This work provided a comprehensive analysis of how mountain forests affect snow cover dynamics across a large area of 58 000 km<sup>2</sup> with variable climate and complex topography in the central European Alps. Data for this study were obtained from simulations using the process-based FSM2oshd forest snow model (Mazzotti et al., 2020b, a; Mott et al., 2023). FSM2oshd has undergone extensive validation, both here and in several prior studies, demonstrating its ability to accurately simulate snow cover dynamics in complex mountainous landscapes. Simulations were conducted at an hourly temporal and 250 m spatial resolution over eight hydrological years. The study area extended from the lowlands north of the Alps to the lowlands south of them, encompassing elevations between 184 and 4806 m. Meteorological forcings were obtained from the high-resolution numerical weather prediction system COSMO, run by MeteoSwiss, and were debiased and downscaled using the approaches detailed in Mott et al. (2023). Canopy data were derived from high-resolution lidar acquisitions, enabling the detailed representation of radiation transmission through the three-dimensional canopy structure using CanRad (Webster et al., 2023).


Our analysis revealed extensive spatial variability of snow cover dynamics in response to variable forest cover, topography, and climatological conditions (Figs. 6 and 7). A particu-

larly striking finding is the marked difference in how snow in north- and south-facing terrain is affected by the presence of forest cover. Effects that were previously only known from meter-resolution observations or simulations, particularly the decisive role of early-season melt (Mazzotti et al., 2023), were carried over to the coarser-scale simulations analyzed here, with potential consequences for runoff generation from snowmelt at catchment scales (Figs. 8 and 10). Variability between years was found to be strong enough to shift or even reverse typical trends with elevation, aspect, between regions, or both, where low-snow years accentuated relative differences in the effects of forest on snow cover (Fig. 9). These results highlight that extensive datasets across large spatial and temporal extents, be it model-based or observational, are necessary to fully embrace in which intricate ways key factors interact to control forest snow processes. Given the hydrological and ecological relevance of forest snow in a changing environment, our work encourages ongoing efforts to advance modeling and observational approaches applicable at these large spatiotemporal scales.

Appendix A




**Figure A1.** Comparison of observed (PlanetScope) and modeled (FSM2oshd) snow cover fraction (SCF) for the Davos focus region during winter and spring of 2020. SCF profiles are shown separately for three land cover types (i.e., dense forest, sparse forest, and open area) on north- or south-facing slopes.



**Figure A2.** Comparison of observed (PlanetScope) and modeled (FSM2oshd) snow cover fraction (SCF) for the six focus regions at the end of winter (approx. 1 April). SCF profiles are shown separately for three land cover types (i.e., dense forest, sparse forest, and open area) on north- or south-facing slopes.

**Table A1.** Performance of the FSM2oshd model to simulate snow cover fraction relative to observations expressed by the mean absolute error (MAE). The percentage of grid cells in an acquisition, assessed by land cover type, is also indicated.

Focus region	Date	Open area MAE (% grid cells)	Sparse forest MAE (% grid cells)	Dense forest MAE (% grid cells)
Rhône bend	1 April 2020	0.158 (4.8 %)	0.125 (9.4 %)	0.128 (16.5 %)
Interlaken	1 April 2020	0.216 (5.0 %)	0.229 (10.4 %)	0.219 (24.5 %)
Gotthard	3 April 2020	0.169 (3.7 %)	0.152 (11.8 %)	0.181 (22.1 %)
Schwyz	1 April 2020	0.255 (4.7 %)	0.269 (7.2 %)	0.399 (13.0 %)
Jura	10 February 2022	0.207 (4.9 %)	0.142 (9.6 %)	0.200 (9.5 %)
Davos	29 March 2019	0.165 (3.2 %)	0.118 (14.9 %)	0.132 (14.4 %)
Davos	7 February 2020	0.046 (1.6 %)	0.196 (6.2 %)	0.171 (6.8 %)
Davos	8 March 2020	0.096 (2.7 %)	0.138 (12.4 %)	0.171 (12.6 %)
Davos	19 March 2020	0.170 (3.0 %)	0.110 (12.1 %)	0.127 (12.9 %)
Davos	1 April 2020	0.136 (3.9 %)	0.102 (11.8 %)	0.138 (12.6 %)
Davos	13 April 2020	0.131 (3.9 %)	0.113 (11.3 %)	0.120 (14.5 %)
Davos	23 April 2020	0.138 (4.8 %)	0.126 (11.6 %)	0.134 (12.8 %)
Davos	1 April 2021	0.168 (6.1 %)	0.122 (20.4 %)	0.153 (22.4 %)
Davos	28 March 2022	0.168 (4.2 %)	0.122 (15.6 %)	0.141 (17.7 %)
Davos	6 April 2023	0.167 (5.2 %)	0.152 (19.0 %)	0.169 (19.7 %)
Davos	6 April 2024	0.171 (4.7 %)	0.117 (16.9 %)	0.132 (14.9 %)



**Figure A3.** Leaf area index (LAI). For each of the six focus regions, a polar plot is provided (A–F). These show the average LAI for eight 150 m elevation bands (950–2150 m) across eight aspect classes, where the labels refer to the directions in which slopes face.

*Code availability.* Code for the FSM2oshd snow model is available via Mott et al. (2026). COSMO forcing data is available from MeteoSwiss at <https://www.meteoswiss.admin.ch/services-and-publications/service/weather-and-climate-products/weather-forecasts.html>, last access: 6 August 2025.

*Data availability.* FSM2oshd model output data used in analyses of this manuscript will be made available upon request. PlanetScope satellite imagery is available at <https://www.planet.com/products/satellite-monitoring/>, last access: 6 August 2025.

*Author contributions.* VH: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing. GM: Conceptualization, Methodology, Software, Supervision, Writing – Review & Editing. CW: Methodology, Software, Writing – Review & Editing. TJ: Conceptualization, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing.

*Competing interests.* The contact author has declared that none of the authors has any competing interests.

*Disclaimer.* Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

*Acknowledgements.* We thank SLF's Operational Snow Hydrological Service for providing the simulations that served as the basis for this study. We also thank the two anonymous reviewers for their constructive and detailed comments, which improved this manuscript. GM received funding from the Swiss National Science Foundation, grants 500PN\_202741 and 5R5PN\_225378. Generative AI tools, including Grammarly, ChatGPT, were used to assist in improving the clarity, grammar, and language of this manuscript. These tools were not used to generate original scientific content (e.g., figures), perform data analysis, or draw conclusions. All intellectual content, interpretations, and final decisions reflect the authors' work.

*Financial support.* This research has been partly supported to GM by the Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung (grant-nos.: 500PN\_202741 and 5R5PN\_225378).

*Review statement.* This paper was edited by Christa Kelleher and reviewed by two anonymous referees.

## References

- Bebi, P., Seidl, R., Motta, R., Fuhr, M., Firm, D., Krumm, F., Conedera, M., Ginzler, C., Wohlgenuth, T., and Kulakowski, D.: Changes of forest cover and disturbance regimes in the mountain forests of the Alps, *For. Ecol. Manag.*, 388, 43–56, <https://doi.org/10.1016/j.foreco.2016.10.028>, 2017.
- Beniston, M.: Is snow in the Alps receding or disappearing?, *WIREs Clim. Change*, 3, 349–358, <https://doi.org/10.1002/wcc.179>, 2012.
- Boon, S.: Snow accumulation following forest disturbance, *Ecohydrology*, 5, 279–285, <https://doi.org/10.1002/eco.212>, 2012.
- Bozzoli, M., Crespi, A., Matiu, M., Majone, B., Giovannini, L., Zardi, D., Brugnara, Y., Bozzo, A., Berro, D. C., Mercalli, L., and Bertoldi, G.: Long-term snowfall trends and variability in the Alps, *Int. J. Climatol.*, 44, 4571–4591, <https://doi.org/10.1002/joc.8597>, 2024.
- Broxton, P. D., Harpold, A. A., Biederman, J. A., Troch, P. A., Molotch, N. P., and Brooks, P. D.: Quantifying the effects of vegetation structure on snow accumulation and ablation in mixed-conifer forests, *Ecohydrology*, 8, 1073–1094, <https://doi.org/10.1002/eco.1565>, 2015.
- Broxton, P. D., van Leeuwen, W. J. D., and Biederman, J. A.: Forest cover and topography regulate the thin, ephemeral snowpacks of the semiarid Southwest United States, *Ecohydrology*, 13, e2202, <https://doi.org/10.1002/eco.2202>, 2020.
- Broxton, P. D., Moeser, C. D., and Harpold, A.: Accounting for Fine-Scale Forest Structure is Necessary to Model Snowpack Mass and Energy Budgets in Montane Forests, *Water Resour. Res.*, 57, e2021WR029716, <https://doi.org/10.1029/2021WR029716>, 2021.
- Cebulski, A. C. and Pomeroy, J. W.: Theoretical Underpinnings of Snow Interception and Canopy Snow Ablation Parameterisations, *WIREs Water*, 12, e70010, <https://doi.org/10.1002/wat2.70010>, 2025.
- Cluzet, B., Magnusson, J., Quéno, L., Mazzotti, G., Mott, R., and Jonas, T.: Exploring how Sentinel-1 wet-snow maps can inform fully distributed physically based snowpack models, *The Cryosphere*, 18, 5753–5767, <https://doi.org/10.5194/tc-18-5753-2024>, 2024.
- Conway, J. P., Pomeroy, J. W., Helgason, W. D., and Kinari, N. J.: Challenges in Modeling Turbulent Heat Fluxes to Snowpacks in Forest Clearings, *J. Hydrometeorol.*, 19, 1599–1616, <https://doi.org/10.1175/JHM-D-18-0050.1>, 2018.
- Cooper, A. E., Kirchner, J. W., Wolf, S., Lombardozi, D. L., Sullivan, B. W., Tyler, S. W., and Harpold, A. A.: Snowmelt causes different limitations on transpiration in a Sierra Nevada conifer forest, *Agr. Forest Meteorol.*, 291, 108089, <https://doi.org/10.1016/j.agrformet.2020.108089>, 2020.
- Currier, W. R., Pflug, J., Mazzotti, G., Jonas, T., Deems, J. S., Bormann, K. J., Painter, T. H., Hiemstra, C. A., Gelvin, A., Uhlmann, Z., Spaete, L., Glenn, N. F., and Lundquist, J. D.: Comparing Aerial Lidar Observations With Terrestrial Lidar and Snow-Probe Transects From NASA's 2017

- SnowEx Campaign, *Water Resour. Res.*, 55, 6285–6294, <https://doi.org/10.1029/2018WR024533>, 2019.
- Currier, W. R., Sun, N., Wigmosta, M., Cristea, N., and Lundquist, J. D.: The impact of forest-controlled snow variability on late-season streamflow varies by climatic region and forest structure, *Hydrol. Process.*, 36, e14614, <https://doi.org/10.1002/hyp.14614>, 2022.
- Deschamps-Berger, C., López-Moreno, J., Gascoin, S., Mazzotti, G., and Boone, A.: Where Snow and Forest Meet: A Global Atlas, *Geophys. Res. Lett.*, 52, <https://doi.org/10.1029/2024GL113684>, 2025.
- Dharmadasa, V., Kinnard, C., and Baraër, M.: Topographic and vegetation controls of the spatial distribution of snow depth in agroforested environments by UAV lidar, *The Cryosphere*, 17, 1225–1246, <https://doi.org/10.5194/tc-17-1225-2023>, 2023.
- Dickerson-Lange, S. E., Vano, J. A., Gersonde, R., and Lundquist, J. D.: Ranking Forest Effects on Snow Storage: A Decision Tool for Forest Management, *Water Resour. Res.*, 57, e2020WR027926, <https://doi.org/10.1029/2020WR027926>, 2021.
- Dickerson-Lange, S. E., Howe, E. R., Patrick, K., Gersonde, R., and Lundquist, J. D.: Forest gap effects on snow storage in the transitional climate of the Eastern Cascade Range, Washington, United States, *Front. Water*, 5, <https://doi.org/10.3389/frwa.2023.1115264>, 2023.
- Dwivedi, R., Biederman, J. A., Broxton, P. D., Pearl, J. K., Lee, K., Svoma, B. M., van Leeuwen, W. J. D., and Robles, M. D.: How three-dimensional forest structure regulates the amount and timing of snowmelt across a climatic gradient of snow persistence, *Front. Water*, 6, <https://doi.org/10.3389/frwa.2024.1374961>, 2024.
- Ellis, C. R., Pomeroy, J. W., Essery, R. L. H., and Link, T. E.: Effects of needleleaf forest cover on radiation and snowmelt dynamics in the Canadian Rocky Mountains, *Can. J. For. Res.*, 41, 608–620, <https://doi.org/10.1139/X10-227>, 2011.
- Essery, R.: A factorial snowpack model (FSM 1.0), *Geosci. Model Dev.*, 8, 3867–3876, <https://doi.org/10.5194/gmd-8-3867-2015>, 2015.
- Essery, R., Pomeroy, J., Ellis, C., and Link, T.: Modelling longwave radiation to snow beneath forest canopies using hemispherical photography or linear regression, *Hydrol. Process.*, 22, 2788–2800, <https://doi.org/10.1002/hyp.6930>, 2008.
- BAFU: Auswirkungen des Klimawandels auf die Schweizer Gewässer. Hydrologie, Gewässerökologie und Wasserwirtschaft, Bundesamt für Umwelt BAFU, Bern, Umwelt-Wissen, Nr. 2101: 134 S, <https://www.bafu.admin.ch/de/gewaesser-im-klimawandel> (last access: 6 August 2025), 2021.
- Frazier, A. E. and Hemingway, B. L.: A Technical Review of Planet Smallsat Data: Practical Considerations for Processing and Using PlanetScope Imagery, *Remote Sens.*, 13, 3930, <https://doi.org/10.3390/rs13193930>, 2021.
- Geissler, J., Rathmann, L., and Weiler, M.: Combining Daily Sensor Observations and Spatial LiDAR Data for Mapping Snow Water Equivalent in a Sub-Alpine Forest, *Water Resour. Res.*, 59, e2023WR034460, <https://doi.org/10.1029/2023WR034460>, 2023.
- Gelfan, A. N., Pomeroy, J. W., and Kuchment, L. S.: Modeling Forest Cover Influences on Snow Accumulation, Sublimation, and Melt, *J. Hydrometeorol.*, 5, 785–803, [https://doi.org/10.1175/1525-7541\(2004\)005<0785:MFCIOS>2.0.CO;2](https://doi.org/10.1175/1525-7541(2004)005<0785:MFCIOS>2.0.CO;2), 2004.
- Haagmans, V., Mazzotti, G., Webster, C., and Jonas, T.: Wintertime forest surface temperature dynamics in boreal and sub-alpine forests, *Agr. Forest Meteorol.*, 372, 110703, <https://doi.org/10.1016/j.agrformet.2025.110703>, 2025.
- Harestad, A. S. and Bunnell, F. L.: Prediction of snow-water equivalents in coniferous forests, *Can. J. For. Res.*, 11, 854–857, <https://doi.org/10.1139/x81-126>, 1981.
- Harpold, A. A., Krogh, S. A., Kohler, M., Eckberg, D., Greenberg, J., Sterle, G., and Broxton, P. D.: Increasing the efficacy of forest thinning for snow using high-resolution modeling: A proof of concept in the Lake Tahoe Basin, California, USA, *Ecohydrology*, 13, e2203, <https://doi.org/10.1002/eco.2203>, 2020.
- Hedstrom, N. R. and Pomeroy, J. W.: Measurements and modelling of snow interception in the boreal forest, *Hydrol. Process.*, 12, 1611–1625, [https://doi.org/10.1002/\(SICI\)1099-1085\(199808/09\)12:10/11<1611::AID-HYP684>3.0.CO;2-4](https://doi.org/10.1002/(SICI)1099-1085(199808/09)12:10/11<1611::AID-HYP684>3.0.CO;2-4), 1998.
- Helbig, N., Schirmer, M., Magnusson, J., Mäder, F., van Herwijnen, A., Quéno, L., Bühler, Y., Deems, J. S., and Gascoin, S.: A seasonal algorithm of the snow-covered area fraction for mountainous terrain, *The Cryosphere*, 15, 4607–4624, <https://doi.org/10.5194/tc-15-4607-2021>, 2021.
- Hendrick, R. L., Filgate, B. D., and Adams, W. M.: Application of Environmental Analysis to Watershed Snowmelt, *J. Appl. Meteorol. Climatol.*, 10, 418–429, [https://doi.org/10.1175/1520-0450\(1971\)010<0418:AOEATW>2.0.CO;2](https://doi.org/10.1175/1520-0450(1971)010<0418:AOEATW>2.0.CO;2), 1971.
- Immerzeel, W. W., Lutz, A. F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B. J., Elmore, A. C., Emmer, A., Feng, M., Fernández, A., Haritashya, U., Kargel, J. S., Koppes, M., Kraaijenbrink, P. D. A., Kulkarni, A. V., Mayewski, P. A., Nepal, S., Pacheco, P., Painter, T. H., Pellicciotti, F., Rajaram, H., Rupper, S., Sinsalo, A., Shrestha, A. B., Viviroli, D., Wada, Y., Xiao, C., Yao, T., and Baillie, J. E. M.: Importance and vulnerability of the world's water towers, *Nature*, 577, 364–369, <https://doi.org/10.1038/s41586-019-1822-y>, 2020.
- Katsushima, T., Kato, A., Aiura, H., Nanko, K., Suzuki, S., Takeuchi, Y., and Murakami, S.: Modelling of snow interception on a Japanese cedar canopy based on weighing tree experiment in a warm winter region, *Hydrol. Process.*, 37, e14922, <https://doi.org/10.1002/hyp.14922>, 2023.
- Koshkin, A. L., Hatchett, B. J., and Nolin, A. W.: Wildfire impacts on western United States snowpacks, *Front. Water*, 4, <https://doi.org/10.3389/frwa.2022.971271>, 2022.
- Kostadinov, T. S., Schumer, R., Hausner, M., Bormann, K. J., Gaffney, R., McGwire, K., Painter, T. H., Tyler, S., and Harpold, A. A.: Watershed-scale mapping of fractional snow cover under conifer forest canopy using lidar, *Remote Sens. Environ.*, 222, 34–49, <https://doi.org/10.1016/j.rse.2018.11.037>, 2019.
- Koutantou, K., Mazzotti, G., Brunner, P., Webster, C., and Jonas, T.: Exploring snow distribution dynamics in steep forested slopes with UAV-borne LiDAR, *Cold Reg. Sci. Technol.*, 200, 103587, <https://doi.org/10.1016/j.coldregions.2022.103587>, 2022.
- Lewis, G., Harpold, A., Krogh, S. A., Broxton, P., and Manley, P. N.: The prediction of uneven snowpack response to forest thinning informs forest restoration in the central Sierra Nevada, *Ecohydrology*, 16, e2580, <https://doi.org/10.1002/eco.2580>, 2023.

- Link, T. E., Marks, D., and Hardy, J. P.: A deterministic method to characterize canopy radiative transfer properties, *Hydrol. Process.*, 18, 3583–3594, <https://doi.org/10.1002/hyp.5793>, 2004.
- López-Moreno, J. I. and Latron, J.: Spatial heterogeneity in snow water equivalent induced by forest canopy in a mixed beech–fir stand in the Pyrenees, *Ann. Glaciol.*, 49, 83–90, <https://doi.org/10.3189/172756408787814951>, 2008.
- López-Moreno, J. I. and Stähli, M.: Statistical analysis of the snow cover variability in a subalpine watershed: Assessing the role of topography and forest interactions, *J. Hydrol.*, 348, 379–394, <https://doi.org/10.1016/j.jhydrol.2007.10.018>, 2008.
- López-Moreno, J. I., Callow, N., McGowan, H., Webb, R., Schwartz, A., Bilish, S., Revuelto, J., Gascoin, S., Deschamps-Berger, C., and Alonso-González, E.: Marginal snowpacks: The basis for a global definition and existing research needs, *Earth-Sci. Rev.*, 252, 104751, <https://doi.org/10.1016/j.earscirev.2024.104751>, 2024.
- Lundquist, J. D., Dickerson-Lange, S. E., Lutz, J. A., and Cristea, N. C.: Lower forest density enhances snow retention in regions with warmer winters: A global framework developed from plot-scale observations and modeling, *Water Resour. Res.*, 49, 6356–6370, <https://doi.org/10.1002/wrcr.20504>, 2013.
- Lundquist, J. D., Dickerson-Lange, S., Gutmann, E., Jonas, T., Lumbrazo, C., and Reynolds, D.: Snow interception modelling: Isolated observations have led to many land surface models lacking appropriate temperature sensitivities, *Hydrol. Process.*, 35, e14274, <https://doi.org/10.1002/hyp.14274>, 2021.
- Magnusson, J., Gustafsson, D., Hüsler, F., and Jonas, T.: Assimilation of point SWE data into a distributed snow cover model comparing two contrasting methods, *Water Resour. Res.*, 50, 7816–7835, <https://doi.org/10.1002/2014WR015302>, 2014.
- Magnusson, J., Winstral, A., Stordal, A. S., Essery, R., and Jonas, T.: Improving physically based snow simulations by assimilating snow depths using the particle filter, *Water Resour. Res.*, 53, 1125–1143, <https://doi.org/10.1002/2016WR019092>, 2017.
- Marks, D., Kimball, J., Tingey, D., and Link, T.: The sensitivity of snowmelt processes to climate conditions and forest cover during rain-on-snow: a case study of the 1996 Pacific Northwest flood, *Hydrol. Process.*, 12, 1569–1587, [https://doi.org/10.1002/\(SICI\)1099-1085\(199808/09\)12:10<1569::AID-HYP682>3.0.CO;2L](https://doi.org/10.1002/(SICI)1099-1085(199808/09)12:10<1569::AID-HYP682>3.0.CO;2L), 1998.
- Marty, C., Tilg, A.-M., and Jonas, T.: Recent Evidence of Large-Scale Receding Snow Water Equivalents in the European Alps, *J. Hydrometeorol.*, 18, 1021–1031, <https://doi.org/10.1175/JHM-D-16-0188.1>, 2017.
- Matiu, M., Crespi, A., Bertoldi, G., Carmagnola, C. M., Marty, C., Morin, S., Schöner, W., Cat Berro, D., Chiogna, G., De Gregorio, L., Kotlarski, S., Majone, B., Resch, G., Terzago, S., Valt, M., Beozzo, W., Cianfarra, P., Gouttevin, I., Marcolini, G., Notarnicola, C., Petitta, M., Scherrer, S. C., Strasser, U., Winkler, M., Zebisch, M., Cicogna, A., Cremonini, R., Debernardi, A., Faletto, M., Gaddo, M., Giovannini, L., Mercalli, L., Soubeyroux, J.-M., Sušnik, A., Trenti, A., Urbani, S., and Weigluni, V.: Observed snow depth trends in the European Alps: 1971 to 2019, *The Cryosphere*, 15, 1343–1382, <https://doi.org/10.5194/tc-15-1343-2021>, 2021.
- Maxwell, J. and St. Clair, S. B.: Snowpack properties vary in response to burn severity gradients in montane forests, *Environ. Res. Lett.*, 14, 124094, <https://doi.org/10.1088/1748-9326/ab5de8>, 2019.
- Mazzotti, G., Currier, W. R., Deems, J. S., Pflug, J. M., Lundquist, J. D., and Jonas, T.: Revisiting Snow Cover Variability and Canopy Structure Within Forest Stands: Insights From Airborne Lidar Data, *Water Resour. Res.*, 55, 6198–6216, <https://doi.org/10.1029/2019WR024898>, 2019a.
- Mazzotti, G., Malle, J., Barr, S., and Jonas, T.: Spatially Continuous Characterization of Forest Canopy Structure and Subcanopy Irradiance Derived from Handheld Radiometer Surveys, *J. Hydrometeorol.*, 20, 1417–1433, <https://doi.org/10.1175/JHM-D-18-0158.1>, 2019b.
- Mazzotti, G., Essery, R., Webster, C., Malle, J., and Jonas, T.: Process-Level Evaluation of a Hyper-Resolution Forest Snow Model Using Distributed Multisensor Observations, *Water Resour. Res.*, 56, e2020WR027572, <https://doi.org/10.1029/2020WR027572>, 2020a.
- Mazzotti, G., Essery, R., Moeser, C. D., and Jonas, T.: Resolving Small-Scale Forest Snow Patterns Using an Energy Balance Snow Model With a One-Layer Canopy, *Water Resour. Res.*, 56, e2019WR026129, <https://doi.org/10.1029/2019WR026129>, 2020b.
- Mazzotti, G., Webster, C., Essery, R., and Jonas, T.: Increasing the Physical Representation of Forest-Snow Processes in Coarse-Resolution Models: Lessons Learned From Upscaling Hyper-Resolution Simulations, *Water Resour. Res.*, 57, e2020WR029064, <https://doi.org/10.1029/2020WR029064>, 2021.
- Mazzotti, G., Webster, C., Quéno, L., Cluzet, B., and Jonas, T.: Canopy structure, topography, and weather are equally important drivers of small-scale snow cover dynamics in subalpine forests, *Hydrol. Earth Syst. Sci.*, 27, 2099–2121, <https://doi.org/10.5194/hess-27-2099-2023>, 2023.
- MeteoSwiss: <https://www.meteoschweiz.admin.ch/klima/klima-der-schweiz.html>, last access: 19 July 2025a.
- MeteoSwiss: <https://www.meteoswiss.admin.ch/about-us/research-and-cooperation/projects/2012/cosmo-next.html>, last access: 5 August 2025b.
- Moeser, C. D., Broxton, P. D., Harpold, A., and Robertson, A.: Estimating the Effects of Forest Structure Changes From Wildfire on Snow Water Resources Under Varying Meteorological Conditions, *Water Resour. Res.*, 56, e2020WR027071, <https://doi.org/10.1029/2020WR027071>, 2020.
- Moeser, D., Stähli, M., and Jonas, T.: Improved snow interception modeling using canopy parameters derived from airborne LiDAR data, *Water Resour. Res.*, 51, 5041–5059, <https://doi.org/10.1002/2014WR016724>, 2015.
- Mott, R., Winstral, A., Cluzet, B., Helbig, N., Magnusson, J., Mazzotti, G., Quéno, L., Schirmer, M., Webster, C., and Jonas, T.: Operational snow-hydrological modeling for Switzerland, *Front. Earth Sci.*, 11, <https://doi.org/10.3389/feart.2023.1228158>, 2023.
- Mott, R., Winstral, A., Cluzet, B., Helbig, N., Magnusson, J., Mazzotti, G., Queno, L., Schirmer, M., Webster, C., and Jonas, T.: FSM2oshd (v1.1.2), Zenodo, <https://doi.org/10.5281/zenodo.18738194>, 2026.
- Painter, T. H., Berisford, D. F., Boardman, J. W., Bormann, K. J., Deems, J. S., Gehrke, F., Hedrick, A., Joyce, M., Laidlaw, R.,

- Marks, D., Mattmann, C., McGurk, B., Ramirez, P., Richardson, M., Skiles, S. M., Seidel, F. C., and Winstral, A.: The Airborne Snow Observatory: Fusion of scanning lidar, imaging spectrometer, and physically-based modeling for mapping snow water equivalent and snow albedo, *Remote Sens. Environ.*, 184, 139–152, <https://doi.org/10.1016/j.rse.2016.06.018>, 2016.
- Pflug, J. M., Yang, K., Cristea, N., Boudreau, E. T., Vuyovich, C. M., and Kumar, S. V.: Using Commercial Satellite Imagery to Reconstruct 3 m and Daily Spring Snow Water Equivalent, *Water Resour. Res.*, 60, e2024WR037983, <https://doi.org/10.1029/2024WR037983>, 2024.
- Planet Labs Inc.: PlanetScope Product Specifications, [https://assets.planet.com/docs/Planet\\_PSScene\\_Imagery\\_Product\\_Spec\\_letter\\_screen.pdf](https://assets.planet.com/docs/Planet_PSScene_Imagery_Product_Spec_letter_screen.pdf) (last access: 21 June 2025), 2023.
- Pomeroy, J., Fang, X., and Ellis, C.: Sensitivity of snowmelt hydrology in Marmot Creek, Alberta, to forest cover disturbance, *Hydrol. Process.*, 26, 1891–1904, <https://doi.org/10.1002/hyp.9248>, 2012.
- Pomeroy, J. W., Parviainen, J., Hedstrom, N., and Gray, D. M.: Coupled modelling of forest snow interception and sublimation, *Hydrol. Process.*, 12, 2317–2337, [https://doi.org/10.1002/\(SICI\)1099-1085\(199812\)12:15<2317::AID-HYP799>3.0.CO;2-X](https://doi.org/10.1002/(SICI)1099-1085(199812)12:15<2317::AID-HYP799>3.0.CO;2-X), 1998.
- Price, A. G.: Prediction of snowmelt rates in a deciduous forest, *J. Hydrol.*, 101, 145–157, [https://doi.org/10.1016/0022-1694\(88\)90032-7](https://doi.org/10.1016/0022-1694(88)90032-7), 1988.
- Quéno, L., Mott, R., Morin, P., Cluzet, B., Mazzotti, G., and Jonas, T.: Snow redistribution in an intermediate-complexity snow hydrology modelling framework, *The Cryosphere*, 18, 3533–3557, <https://doi.org/10.5194/tc-18-3533-2024>, 2024.
- Redding, T. and Devito, K.: Aspect and soil textural controls on snowmelt runoff on forested Boreal Plain hillslopes, *Hydrol. Res.*, 42, 250–267, <https://doi.org/10.2166/nh.2011.162>, 2011.
- Rinehart, A. J., Vivoni, E. R., and Brooks, P. D.: Effects of vegetation, albedo, and solar radiation sheltering on the distribution of snow in the Valles Caldera, New Mexico, *Ecology*, 1, 253–270, <https://doi.org/10.1002/eco.26>, 2008.
- Roth, T. R. and Nolin, A. W.: Forest impacts on snow accumulation and ablation across an elevation gradient in a temperate montane environment, *Hydrol. Earth Syst. Sci.*, 21, 5427–5442, <https://doi.org/10.5194/hess-21-5427-2017>, 2017.
- Roth, T. R. and Nolin, A. W.: Characterizing Maritime Snow Canopy Interception in Forested Mountains, *Water Resour. Res.*, 55, 4564–4581, <https://doi.org/10.1029/2018WR024089>, 2019.
- Safa, H., Krogh, S. A., Greenberg, J., Kostadinov, T. S., and Harpold, A. A.: Unraveling the Controls on Snow Disappearance in Montane Conifer Forests Using Multi-Site Lidar, *Water Resour. Res.*, 57, e2020WR027522, <https://doi.org/10.1029/2020WR027522>, 2021.
- Sanmiguel-Vallelado, A., Camarero, J. J., Morán-Tejeda, E., Gazol, A., Colangelo, M., Alonso-González, E., and López-Moreno, J. I.: Snow dynamics influence tree growth by controlling soil temperature in mountain pine forests, *Agr. Forest Meteorol.*, 296, 108205, <https://doi.org/10.1016/j.agrformet.2020.108205>, 2021.
- Seidl, R., Thom, D., Kautz, M., Martin-Benito, D., Peltoniemi, M., Vacchiano, G., Wild, J., Ascoli, D., Petr, M., Honkaniemi, J., Lexer, M. J., Trotsiuk, V., Mairota, P., Svoboda, M., Fabrika, M., Nagel, T. A., and Reyer, C. P. O.: Forest disturbances under climate change, *Nat. Clim. Change*, 7, 395–402, <https://doi.org/10.1038/nclimate3303>, 2017.
- Seyednasrollah, B. and Kumar, M.: How Surface Radiation on Forested Snowpack Changes across a Latitudinal Gradient, *Hydrology*, 6, 62, <https://doi.org/10.3390/hydrology6030062>, 2019.
- Sicart, J. E., Essery, R. L. H., Pomeroy, J. W., Hardy, J., Link, T., and Marks, D.: A Sensitivity Study of Daytime Net Radiation during Snowmelt to Forest Canopy and Atmospheric Conditions, *J. Hydrometeorol.*, 5, 774–784, [https://doi.org/10.1175/1525-7541\(2004\)005<0774:ASSODN>2.0.CO;2](https://doi.org/10.1175/1525-7541(2004)005<0774:ASSODN>2.0.CO;2), 2004.
- Stähli, M. and Gustafsson, D.: Long-term investigations of the snow cover in a subalpine semi-forested catchment, *Hydrol. Process.*, 20, 411–428, <https://doi.org/10.1002/hyp.6058>, 2006.
- Stewart, I. T., Cayan, D. R., and Dettinger, M. D.: Changes in Snowmelt Runoff Timing in Western North America under a “Business as Usual” Climate Change Scenario, *Clim. Change*, 62, 217–232, <https://doi.org/10.1023/B:CLIM.0000013702.22656.e8>, 2004.
- Strasser, U., Bernhardt, M., Weber, M., Liston, G. E., and Mauser, W.: Is snow sublimation important in the alpine water balance?, *The Cryosphere*, 2, 53–66, <https://doi.org/10.5194/tc-2-53-2008>, 2008.
- Strasser, U., Warscher, M., and Liston, G. E.: Modeling Snow-Canopy Processes on an Idealized Mountain, *J. Hydrometeorol.*, 12, 663–677, <https://doi.org/10.1175/2011JHM1344.1>, 2011.
- Sturm, M. and Liston, G. E.: Revisiting the Global Seasonal Snow Classification: An Updated Dataset for Earth System Applications, *J. Hydrometeorol.*, 22, 2917–2938, <https://doi.org/10.1175/JHM-D-21-0070.1>, 2021.
- Sturm, M., Holmgren, J., and Liston, G. E.: A Seasonal Snow Cover Classification System for Local to Global Applications, *J. Climate*, 8, 1261–1283, [https://doi.org/10.1175/1520-0442\(1995\)008<1261:ASSCCS>2.0.CO;2](https://doi.org/10.1175/1520-0442(1995)008<1261:ASSCCS>2.0.CO;2), 1995.
- Sun, N., Wigmosta, M., Zhou, T., Lundquist, J., Dickerson-Lange, S., and Cristea, N.: Evaluating the functionality and streamflow impacts of explicitly modelling forest-snow interactions and canopy gaps in a distributed hydrologic model, *Hydrol. Process.*, 32, 2128–2140, <https://doi.org/10.1002/hyp.13150>, 2018.
- Troendle, C. A.: The Potential for Water Yield Augmentation from Forest Management in the Rocky Mountain Region, *JAWRA J. Am. Water Resour. Assoc.*, 19, 359–373, <https://doi.org/10.1111/j.1752-1688.1983.tb04593.x>, 1983.
- Viviroli, D., Dürr, H. H., Messerli, B., Meybeck, M., and Weingartner, R.: Mountains of the world, water towers for humanity: Typology, mapping, and global significance, *Water Resour. Res.*, 43, <https://doi.org/10.1029/2006WR005653>, 2007.
- Webster, C., Rutter, N., and Jonas, T.: Improving representation of canopy temperatures for modeling sub-canopy incoming longwave radiation to the snow surface, *J. Geophys. Res. Atmospheres*, 122, 9154–9172, <https://doi.org/10.1002/2017JD026581>, 2017.
- Webster, C., Essery, R., Mazzotti, G., and Jonas, T.: Using just a canopy height model to obtain lidar-level accuracy in 3D forest canopy shortwave transmissivity estimates, *Agr. Forest Meteorol.*, 338, 109429, <https://doi.org/10.1016/j.agrformet.2023.109429>, 2023.