



Climate and topographic controls on pasture production

J. Lozano-Parra et al.

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

Climate and topographic controls on pasture production in a semiarid Mediterranean watershed with scattered tree cover

J. Lozano-Parra¹, M. P. Maneta², and S. Schnabel¹

¹Geoenvironmental Research Group, University of Extremadura, Avda, Universidad, 10071 Cáceres, Spain

²Geosciences Department, The University of Montana, 32 Campus Drive, Missoula, Montana, USA

Received: 22 November 2013 – Accepted: 22 November 2013
– Published: 11 December 2013

Correspondence to: J. Lozano-Parra (jlozano@unex.es)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

Natural grasses in semiarid rangelands constitute an effective protection against soil erosion and degradation, are a source of natural food for livestock and play a critical role in the hydrologic cycle by contributing to the uptake and transpiration of water. However, natural pastures are threatened by land abandonment and the consequent encroachment of shrubs and trees as well as by changing climatic conditions. In spite of their ecological and economic importance, the spatio-temporal variations of pasture production at the decadal to century scales over whole watersheds are poorly known. We used a physics-based, spatially-distributed ecohydrologic model applied to a 99.5 ha semiarid watershed in western Spain to investigate the sensitivity of pasture production to climate variability. The ecohydrologic model was run using a 300 yr long synthetic daily climate dataset generated using a stochastic weather generator. The data set reproduced the range of climatic variations observed under current climate. Results indicated that variation of pasture production largely depended on factors that also determined the availability of soil moisture such as the temporal distribution of precipitation, topography, and tree canopy cover. The latter is negatively related with production, reflecting the importance of rainfall and light interception, as well as water consumption by trees. Valley bottoms and flat areas in the lower parts of the catchment are characterized by higher pasture production. A quantitative assessment of the quality of the simulations showed that ecohydrologic models are a valuable tool to investigate long term (century scale) water and energy fluxes, as well as vegetation dynamics, in semiarid rangelands.

1 Introduction

Traditional Mediterranean agrosilvopastoral systems support high levels of biodiversity in a wide variety of coexisting natural and man-made habitats, such as grazing areas, agricultural lands, scrublands, forests or wildlife spaces (Joffre et al., 1988;

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



HESSD

10, 15167–15217, 2013

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Campos-Palacín, 2004). Natural grasses and pastures are an important element of cohesion between these habitats by supporting livestock and other fauna, by protecting the soil against erosion and degradation and by controlling the soil hydrologic and thermal regime (Schnabel, 1997; Paço et al., 2009). The economic importance of pasture incents the proper management and conservation of Mediterranean agrosilvopastoral systems, however owing to climate characteristics of semiarid Mediterranean environments, natural herbaceous production is highly variable with a pronounced seasonality, being highest in spring, low in autumn and winter, and nil during summer (Montero et al., 1998; Joffre and Rambal, 1993). Additionally, pasture yield is usually low and its spatiotemporal distribution is strongly conditioned by the balance of positive and negative effects of limiting factors such as water, light, or nutrients (Brooker et al., 2008).

Decreased pasture yields may upset the balance of habitats and threaten the sustainability of these Mediterranean systems due to changes in land use associated with the revision of economic priorities and management decisions. Indeed pastures in Mediterranean Europe have been experiencing land abandonment and consequent encroachment of shrubs and forest (Rivest et al., 2011; García-Ruiz and Lana-Renault, 2011; Lavado-Contador et al., 2004), which may lead to increased competition for resources, such as water and light, among different layers of vegetation (Cubera and Moreno, 2007a). The abandonment of traditional agrosilvopastoral systems may not only have important ecologic consequences but may also have a significant impact on regional economies and on food security by affecting forage quality and quantity and by affecting productivity and protection of the agricultural landscape against degradation.

Improved knowledge of the frequency of low and high pasture productivity periods and the expected variability of yields in different locations of a region permits making better informed management decisions that contribute to the sustainability of agrosilvopastoral systems, however, we still only have a partial understanding of the ecohydrological processes that control plant productivity across space and time (Asbjornsen et al., 2011).

**Climate and
topographic controls
on pasture
production**

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

From the mid 1990's there has been a growing interest in the complex interactions between ecological and hydrological processes at multiple scales (Viville and Littlewood, 1996; Rodríguez-Iturbe, 2000; Wang et al., 2012; Caylor et al., 2005, 2009; Porporato et al., 2002; Rodríguez-Iturbe et al., 1999). Because of the complex and non-linear interactions between vegetation and hydrology, few studies focus on the larger scales, such as landscapes or watersheds, where the processes are less understood (Asbjornsen et al., 2011). A limited number of models have been developed in the last decade to investigate ecohydrologic interactions at watershed and regional scales (e.g. Ivanov et al., 2008; Oleson et al., 2010; Tague and Band, 2004; Maneta and Silverman, 2013; Fatichi et al., 2012). Most of the studies using these models have focused on short-term studies because of the long run-times derived from their complexity and because the lack of existing extensive climate data sets (longer than a few decades) needed to force the models. These limitations have resulted in few studies producing detailed experiments that simulate the entire range of ecohydrological conditions that can be expected under current climate variability. These studies would be highly valuable to improve our understanding of the variability of pasture production and to inform grassland management.

Reproducing the entire range of ecohydrologic states at scales useful to gain insight into watershed scale processes require the ability to simulate extensive periods in the order of hundreds of years at small spatial (1–50 m) and temporal (daily) scales. Maneta and Silverman (2013) present a ecohydrologic model with a level of complexity that can make the simulation of extensive periods at detailed spatial and temporal scales tractable while maintaining a strong mechanistic description of the processes. The lack of extensive input datasets to the model can be overcome by producing synthetic datasets with stochastic weather generators (SWG). These tools have been successfully used since the early 1980's (Richardson, 1981) to generate long time-series of synthetic weather data that are statistically indistinguishable from observed shorter term climate records (Semenov and Barrow, 2002). SWGs have been used to simulate future scenarios of climate change (Fatichi et al., 2011; Semenov and Barrow, 1997),

crop yields (Semenov and Porter, 1995; Ivanov et al., 2007) or regional hydrologic response (Xia, 1996; Dubrovský et al., 2004).

In this paper we use a combination of mechanistic models and SWG to investigate the spatial-temporal variability of pasture production at watershed scales relevant for management. Questions that we seek to address include: How does pasture production respond to climate variability in combination with antecedent basin conditions? How sensitive is the production of pasture to the temporal distribution of precipitation during the year? How important are topographic controls vs climatic controls in determining the spatial and temporal dynamics of production in a watershed? Does the relative importance of these controls vary for different years and under different circumstances?

While abundant studies have applied numerical models to the study of grassland productivity (Montaldo et al., 2005; Istanbuluoglu et al., 2012) and some work has a focus on the spatio-temporal variability of pasture production over long periods (century scale) and large areas (Clark et al., 2003; Tubiello et al., 2007), to the authors' knowledge no studies have applied comprehensive mechanistic numerical models to address the questions posed above. Experimental or field studies have not addressed satisfactorily these questions either because pasture production over large areas is typically determined with a limited number of measurements commonly taken over a few years and at very specific locations (Plaixats et al., 2004; Santamaría et al., 2009). The limited number of samples could provide a skewed or erroneous estimate of the actual long-term pasture production of a region or farm because a short time interval may not properly capture weather variations, such as wet and dry periods, and the specific sampling location could not properly characterize the actual spatial variations. A modeling approach is therefore preferred in this study.

HESSD

10, 15167–15217, 2013

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2 Study area

General description

The study area is an experimental drainage basin located in the southwestern part of the Iberian Peninsula with an area of 99.5 ha (Fig. 1), characterized by an agrosilvopastoral land use system called *dehesa* in Spain. Geologically, the study area forms part of the Iberian Massif of Precambrian age, being the dominant rocks greywacke and schist, which were eroded giving rise to an erosion surface. Topography of the drainage basin is gently undulating with an average elevation of 394 m.a.s.l., being SSW the dominant aspect. Climate is Mediterranean with a high seasonal and inter-annual rainfall variability (Schnabel, 1998), which determines the available water content for plants, and a marked dry season during summer that can last four months or even more. Average annual precipitation for the period between 1999 and 2012 was 488 ± 149.5 mm (mean \pm standard deviation) and mean monthly temperatures ranged between 7.4 ± 1.7 °C in January to 26.4 ± 1.5 °C in July and August. Annual potential evapotranspiration is twice the annual rainfall amount. Vegetation is typically Mediterranean, characterized by a two-layered vegetation structure, with a layer of scattered trees (*Quercus ilex*) at low density (20 ± 18 individuals ha^{-1}), and a pasture layer. Natural pastures are composed of annual and perennial herbaceous plants, abounding especially annual grasses (such as *Vulpia bromoides*, *Bromus* sp. or *Aira caryophylllea*) and annual legumes (*Ornithopus compressus*, *Lathyrus angulatus* and several species of *Trifolium*), starting to grow with the first rainfall in autumn and reaching maximum production in spring. A layer of shrubs is also frequent (*Retama sphaerocarpa*), commonly eliminated by ranchers to facilitate pasture growth.

Soils in the catchment have a high bulk density, $\approx 1.5 \text{ g cm}^{-3}$, are poor in nutrients and have low organic matter content: $\approx 3\%$, except below tree cover where it is higher in the upper 5 cm (Schnabel et al., 2013b). Roots are concentrated in the upper soil layer (Moreno et al., 2005), favoring the higher porosity ($\approx 45\%$) of the topsoil. Two geomorphologic units can be distinguished in the catchment which determines the type

HESSD

10, 15167–15217, 2013

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Climate and topographic controls on pasture production

J. Lozano-Parra et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of soil and its hydrologic properties. The boundary between these units is marked by the 395 m contour (Fig. 1). The geomorphological unit above 395 m is the northern part of the catchment. It constitutes the slopes of a pediment with sandy loam soils classified as Luvisols (FAO, 1988), rich in rock fragments that provides it with a higher permeability and saturated hydraulic conductivity than the remaining soils (Van Schaik et al., 2008; Van Schaik, 2009). Soil depths in this unit are variable, often exceeding 1 m to bedrock and with an argillic B horizon. The other geomorphologic unit, flat to gently undulating, is located in the lower part of the basin. In this unit soils are very shallow (Cambisols and Leptosols), ranging between 20–50 cm, developed on impermeous bedrock of schist and greywacke, which frequently crops out. The lowest areas of this unit correspond with valley bottoms covered by alluvial sediments reaching a thickness of approximately 1 m in areas next to channels. The main channel is incised into these sediments, actively eroding at present and can be classified as a gully (Gómez-Gutiérrez et al., 2009). Owing to low permeability of these layers some sites are prone to ponding in wet periods (Cerdá et al., 1998; Van Schaik, 2009), which provide an extra water storage that may lengthen the phenological period of the herbaceous plants and that is totally dried in summer. A complete and detailed description of the study area can be found in Maneta (2006) and Van Schaik (2010).

3 Methods

3.1 Field data

3.1.1 Meteorological data

The study area is equipped with a meteorological station that collects information on precipitation, temperature, relative humidity, global radiation, net radiation, wind speed and direction at intervals of 5 min since year 2000. Rainfall is also measured in five

other locations (Fig. 1) with tipping bucket type rain gauges of 0.2 mm resolution. This information was aggregated in daily intervals for this study.

3.1.2 Soil moisture content and soil temperature

Volumetric soil water content was monitored by capacitive sensors (*Decagon Device, Inc.* model *EC-5*) at 5, 10, 15 and 30 cm depth every 30 min. Soil temperature was measured at 5 cm depth in line with the soil moisture probes (*Decagon Device, Inc.* model *RT-1*). The accuracy of the soil moisture sensors was improved by calibration following the method of Cobos and Chambers (2010). The sensors were grouped in soil moisture stations (SMS) at two sites: *Site 1* representative of hillslopes with Luvisols, and *Site 2* representative of the lower part of the catchment with shallow soils. A third SMS was installed in the eastern part of the catchment (Fig. 1). The selection of sites to install the SMSs were based on previous studies by Lavado-Contador et al. (2006), Maneta et al. (2007, 2008a, b) and Van Schaik et al. (2008, 2009). The SMSs in *Site 1* and *Site 2* began to register in March 2009, while SMS-3 started in May 2010. In each site there are sensors in open grass areas and under tree canopies. The overall soil moisture and soil temperature of each site was considered to be the depth-averaged soil moisture and soil temperature of the sensors under trees and in open areas, weighted by the relative canopy cover in its pixel.

3.1.3 Pasture production

We have measured natural pasture production at *Site 1* and *Site 2* for three hydrologic years (from Sept 2008 through August 2011). To prevent grazing, twelve 1 m × 1 m livestock exclusion cages were installed at midslope positions in open space. Only aerial (above-ground) production is considered in this study. Grasses and forbs were cut twice a year (at the end of winter and at the end of spring), dried during 48 h in an oven at 105 °C and weighted to determine aerial dry matter (DM) production (kg DM ha⁻¹).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Measurements of DM were augmented with measurements of pasture height. At each SMS, 16 measurements of plant height were taken biweekly during two hydrological years (from 1 March 2011 until 31 August 2012). The pasture production database was extended by estimating DM from pasture height measurements using their allometric relationship ($r^2 = 0.68$, $n = 12$).

3.2 Ecohydrologic model

To simulate water and energy exchanges and pasture production we used a spatially distributed ecohydrologic model as described in Maneta and Silverman (2013). This model couples a two layer (canopy and understory) vertical local closure energy balance scheme, a hydrologic model and a carbon uptake and vegetation growth component. The model was run using climate information from a stochastic weather generator as described below.

Vertical energy transfers are calculated using first-order closure profile equations for momentum, heat and mass under neutral stratification based on flux gradient similarity (Arya, 2001; Foken, 2008). The energy balance is solved for the canopy layer and then for the soil layer using canopy temperature and soil temperature as the closure variables, respectively. Canopy conductance is calculated with a Jarvis-type multiplicative model (Cox et al., 1998; Jarvis, 1976). Water infiltration into the soil is calculated using the Green and Ampt approximation to Richard's equation (Chow et al., 1988). Lateral water transfers in the soil are simulated using a 1-D kinematic wave model (Singh, 1997). Interception of water by canopies is simulated using a bucket model. The forest growth and carbon uptake components are based on 3-PG (Landsberg and Waring, 1997). See Maneta and Silverman (2013) for further details.

The ecohydrologic model by Maneta and Silverman (2013) was extended in this study with a new grass growth component. Net primary production of grass is related to the available radiation intercepted by the canopy and the water transpired:

$$\text{NPP} = C_{\text{NPP}} \cdot f(T_a) \cdot \sqrt{\alpha \cdot \text{PAR} \cdot \beta \cdot \text{Transp}} \quad (1)$$

$$\xi_{sd} = \min \left(1, \frac{T_c}{T_\xi} \right). \quad (3)$$

Parameter ϕ_a (Eq. 2a and b) controls the allocation of NPP to the aboveground (green leaves) and belowground (roots) pool of carbon based on the spare capacity of the land to carry aboveground biomass (Istanbulluoglu et al., 2012):

$$\phi_a = \left(1 - \frac{LAI_g}{LAI_{max} - LAI_d} \right) \quad (4)$$

where LAI_g , LAI_{max} , and LAI_d are green, maximum, and dead grass leaf area indices, respectively. The denominator of Eq. (4) indicates the space available to grow green leaves.

The transformation of the aboveground mass to leaf area index is done using the specific leaf area index for green and dead leaves:

$$LAI_g = \sigma_{LAI_g} M_g \quad (5a)$$

$$LAI_d = \sigma_{LAI_d} M_g \quad (5b)$$

$$LAI_t = LAI_g + LAI_d \quad (5c)$$

where σ_{LAI_g} and σ_{LAI_d} are the specific leaf area indices for green and dead leaves. Total leaf area index (LAI_t) is considered to be the sum of the green and dead leaf area indices.

3.3 Model set up

Hydrologic properties, land cover and vegetation parameters

The modeling domain was discretized with a 30 m × 30 m grid, as used in previous studies (Maneta et al., 2008). A digital elevation model (DEM) was used to delineate the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

daily rainfall as predictors ($r^2 = 0.75$). Wind velocity was obtained by repeating a series of 51 yr extracted from a station located at 24 km from the study site. Daily long wave radiation was estimated from air temperature using the method described by Swinbank (1964).

3.5 Model calibration, spin up and data analysis

The calibration runs were done running the period from 1 September 2008 to 31 August 2012 in a continuous loop using daily time steps. Model parameters listed in Table 2 were manually calibrated until soil moisture, soil temperature and pasture yield achieved steady state and satisfactorily matched the available measurements of soil moisture, soil temperature, and pasture yield based on height measurements. Calibration was based on trial and error systematically changing parameters one at a time. When available, the initial trial value was based on values cited in the literature or based on experience. Model performance was quantified using the coefficient of determination, root mean square error, bias and Nash–Sutcliffe efficiency coefficient between modeled and observed soil moisture, soil temperature and pasture yield. Once performance was satisfactory with parameter values within a realistic range the model was considered calibrated.

The calibrated model was used in a 300 yr long simulation at daily time steps resulting in 109 500 maps per state variable reported by the model. State variables analyzed included soil moisture, soil temperature, pasture production, pasture evaporation and transpiration, and tree evaporation and transpiration. Time averages and standard deviations for the entire simulation period were calculated for each variable, except for pasture production. For this latter variable, the average and standard deviations for 1 June were used in the analysis because this date corresponds to the end of the vegetative period of herbaceous plants and can be considered as the day of maximum accumulated production.

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



4 Results and discussion

4.1 Model performance

Annual mean precipitation for the simulated period was 508.8 mm with a standard deviation of 118.2 mm. Maximum and minimum annual rainfall were 934.1 and 188.2 mm, respectively. The longest dry spell spanned four years with annual rainfalls lower than 386.9 mm yr⁻¹, while the maximum wet period lasted three years with rainfall in excess of 693.4 mm yr⁻¹.

A comparison between simulated and observed atmospheric data indicated that the SWG was properly calibrated and that it successfully generated a synthetic times series that was statistically indistinguishable from the observations (Table 3) except for rainfall in July and August. This is because during these months precipitation volumes are insignificant and small fluctuations about the very low observed precipitation values have a relatively large influence in the K–S statistic. This is of minor importance because rainfall in these months is virtually zero. Further inspection of the results showed that the generated weather series present the seasonal and inter-annual variations typical of the Mediterranean climate.

An initial inspection of the graphs shown in Figs. 3 and 4 indicates that the model reproduced to a high degree the observed dynamic of soil moisture and temperature. The simulation captured the seasonal variations of soil moisture, including the wetting and recession rates, but also much of the observed high-frequency variation. Some mismatch can be observed in the reproduction of wetting peaks, such as those of *Site 1* (Fig. 3a). There is a general dampening of the amplitude of high frequency variations that may be due to the model representation of soil moisture as the average over the entire soil profile (Maneta and Silverman, 2013). However the standard goodness-of-fit statistics and descriptive statistic confirmed a satisfactory fit with high coefficients of determination ($r^2 \geq 0.80$), low RMSE ($\leq 0.047 \text{ m}^3 \text{ m}^{-3}$) and similar statistics for all measurement stations (Table 4). Further evaluation of the model performance show high Nash–Sutcliffe coefficients (≥ 0.75) and low prediction bias ($\leq 0.018 \text{ m}^3 \text{ m}^{-3}$).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The simulated soil temperature captured the high-frequency variation of observed soil temperature (Fig. 4). However, during the first year simulated temperatures were higher than observed in both study sites, which could be caused by uncommonly low pasture yields simulated that year and hence an overestimation of the amount of radiation reaching the bare soil, while actual ground covered by pasture was much higher at the SMS sites because they were protected against grazing. Efficiency statistics for soil temperature were satisfactory, with coefficients of determination $r^2 \geq 0.89$ and the Nash–Sutcliffe efficiency criterion 0.86, increasing our confidence on the capacity of the model to represent the energy fluxes in the study site (Table 4).

Simulated annual pasture production matched well the observed data at both field sites (Table 4). The average simulated value of production for both sites was $630.9 \text{ kg DM ha}^{-1}$, very similar to the observed $623.8 \text{ kg DM ha}^{-1}$. Other descriptive statistics (minimum, maximum, standard deviation) and goodness-of-fit statistics confirming the model in our research area are shown in Table 4. The model produced a satisfactory description of the spatio-temporal dynamics of production, which is supported by the high prediction efficiency of the model (Nash–Sutcliffe ≥ 0.75 ; $r^2 \geq 0.76$) and low residual errors (RMSE = $164.8 \text{ kg DM ha}^{-1}$).

The phenological cycle of the herbaceous plants in the study site (Fig. 5) is captured in the simulated data and includes low production in autumn although dependent on antecedent precipitation, scarce production in winter because of low air temperatures and available energy, high production in spring when water and energy are available and an absence of production in summer because of lack of water. It is important to note that once pasture is cut at the sites to measure its dry biomass, the exclusion cage is moved to a nearby location, which contributes to the difference between DM estimated from cuts (blue diamonds) and from vegetation height (green circles) since production is highly variable even at short distances (as indicated by the standard deviation of pasture cuts, Fig. 5). In contrast, plant height is always and consistently measured at the same location (SMS).

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A multiple regression analysis revealed that the most explanatory variables determining the spatial distribution of soil moisture are canopy cover, porosity, slope, and elevation. These variables explained 68 % of the observed variance and, with the exception of porosity, showed a negative correlation with soil moisture. Canopy cover showed a particularly strong negative relationship with soil moisture, indicating that the reduction of water reaching the ground due to rainfall interception and the additional water uptake by the trees was a more determinant control of soil moisture than the reduction of incident radiation and evaporation below tree canopies due to shading.

Low lying areas had greater average soil moisture (Fig. 7a). These areas correspond to the valley bottoms and flat footslopes, which show better conditions for water maintenance by the effect of topography (concentrating water) or thicker soils with a higher content of clay and silt particles and greater porosity (McGlynn et al., 2003; Jencso et al., 2009). In contrast, hillslopes and areas at greater altitude had lower soil moisture values, which could be attributed to smaller contributing areas, higher canopy cover and coarser soil textures. However, a small area in the north-eastern upper part of the catchment also showed high average soil moisture values, which could be explained by its low tree density and low canopy cover.

These results highlight the importance of trees in the spatial distribution of soil moisture. This has been observed in dehesa systems by Lavado-Contador et al. (2006), Martínez Fernández et al. (2007) or Moreno and Cubera (2008). Whether trees enhance or reduce soil moisture with respect to open areas seems to be dependent on the climatic conditions of the site (Lozano-Parra et al., 2011). Joffre and Rambal (1988) found higher water content beneath tree canopies in sub-humid ecosystems, which could explain enhanced pasture yields in these situations. Likewise, Gindel (1964) observed also higher water content beneath canopy than in open areas under subtropical and semi-desert conditions. In contrast, García-Estringana et al. (2013) measured lower soil moisture under forest cover in a Mediterranean mountain area, while Cubera and Moreno (2007b) and Gea-Izquierdo et al. (2009) found lower water contents

beneath canopy in semiarid conditions with scattered trees, which is in accordance with our results.

The variability of soil moisture is presented in Fig. 7b and shows a spatial distribution that correlates with the distribution of soil moisture averages. Higher temporal variability of soil moisture was observed in areas with high average soil moisture (e.g. valley bottoms). In contrast, areas with low mean soil water content such as hillslopes with high gradients showed less temporal moisture variability. An explanation for this behavior is that regions with intermediate and higher water contents and soils with good retention properties have more opportunities for soil moisture fluctuations than drier soils with poorer soil water retention capabilities that quickly drain and dry.

Simulated evapotranspiration was marked by the spatial distribution of vegetation cover and by topography (Fig. 7c). Maximum values were found in the valley bottoms where water content remained high during most of the year. High values were also observed in areas with high tree density, while they were lower in open areas where herbaceous vegetation dominates. Annual mean value of evapotranspiration for the whole catchment was 390 mm while annual mean precipitation was 508 mm. This implies that about 120 mm could become runoff or to be stored in the soil reservoirs (Fig. 1) or rock fractures of the impermeable bedrock of the catchment. In support of this, Schnabel et al. (2013a) measured in the same environment runoff values that oscillated between 10 and 190 mm depending on annual precipitation. The simulated annual evapotranspiration values in areas of relatively high tree density are similar to the 590 mm reported by Joffre and Rambal (1993) under tree cover in sub-humid Mediterranean rangelands. They found, however, higher annual values, 400 mm, in open spaces, which could be explained because their study was carried out in a wetter environment.

4.2.2 Pasture production: temporal dynamics

At *Site 1* annual average dry matter production was 338.0 kg ha^{-1} , with a standard deviation of 172.5 kg ha^{-1} , and maximum and minimum values of 977.6 and

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of the total annual precipitation. Years with low accumulated precipitation before June were less productive than years with higher accumulated precipitation (Table 6). For example, similar annual rainfall occurred in years 210 and 213, however in the year 213 the rainfall of the last four months prior to June was higher, which resulted in a greater yield. In the year 215 a large amount of rainfall occurred after May, but pasture production that year was low.

Antecedent rainfall of the last 120 days before June was the variable that explained best the annual pasture production ($r^2 = 0.73$ and $r^2 = 0.51$, for *Site 1* and *Site 2*, respectively). Shorter accumulation periods for antecedent precipitation had poorer correlations with yield, which can be explained because they are associated with less growing time and because as summer approaches there is an increase in evaporation losses.

4.2.3 Pasture production: spatial distribution

The spatial distribution of simulated pasture production varied greatly across the basin. Figure 10a presents the spatial distribution of average production in the catchment over the entire 300 simulated years. Areas of higher production tended to have higher variability in their production (Fig. 10b) as well as higher maximum and minimum productivities (Fig. 10c and d). Productivity areas were persistent in time, with distributions determined by physiographic characteristics of the basin and the distribution of trees. A multiple regression analysis of pasture production with different variables showed that soil moisture, slopes, tree density, canopy cover, and upslope catchment area were the best predictors of production ($r^2 = 0.81$).

The distribution, composition and structure of plant communities are directly conditioned by spatio-temporal patterns in water availability (Asbjornsen et al., 2011) which is strongly determined by topography. In the study catchment the spatial distribution of the natural pastures was clearly influenced by the distribution of soil moisture. Areas with higher water availability had greater yield (Fig. 11a). Low yields were obtained if average soil moisture was lower than $0.150 \text{ m}^3 \text{ m}^{-3}$. Slope also played a strong role

in the distribution of yield. Topographically, valley bottoms and flat areas of the catchment were characterized by higher pasture production. Production decreased rapidly as slope increased (Fig. 11b). This is because in semiarid regions higher slopes are associated with reduced infiltration, enhanced drainage and production of overland flow (Cerdá et al., 1998). The importance of physiographic controls on soil moisture distribution and hence of pasture production in the study region was clearly documented in Ceballos and Schnabel (1998) and Van Schaik (2009), who demonstrated the importance of soils in low lying areas as water storages and the fundamentally different hydrologic regimes of hilltops, hillslopes, low areas and valley bottoms.

Canopy cover exerted a strong control on pasture yield (Fig. 11c). An initial explanation is that pixels with high canopy coverage have higher interception of incident precipitation, more transpiration and therefore reduced soil moisture. This interpretation is however insufficient since the influence of trees on pasture production is a more complex issue that involves a number of processes not explicitly simulated in this study. For instance, trees may promote pasture production by enhancing soil fertility and structure or by providing a shaded and favorable microclimate. These factors were not explicitly simulated in this study. Still, it is known that in semiarid ecosystems, rainfall interception together with soil water uptake by trees in areas of high canopy cover would increase the competence for water resources between trees and pastures rather than enhance the production of pastures (Moreno, 2008). However, because the model used in this study does not incorporate many processes describing the overstory–pasture relationships such as the effect of vegetation on nutrients and on the soil microbial activity, we cannot conclude that tree canopy cover is strictly detrimental to the production of pastures. Indeed, several studies in the region show increased yield under trees as compared to open areas (Moreno, 2008). It has been observed that moderation of incident light could have a positive effect on crop production by altering the microclimate under trees, however this effect depends on antecedent conditions and the production of previous years (Gea-Izquierdo et al., 2009). Values of 13% of canopy cover with

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

24 trees ha⁻¹ were considered optimum for understory pasture production (Montero et al., 2008).

4.2.4 Climatic and physiographic factors

The degree to which the various controls discussed in the previous sections determine the distribution of pasture is not invariant. Precipitation is a main driver of total production (Fig. 12a) in almost a linear fashion, but the spatial distribution of pasture is to a large extent controlled by topography, since the spatial variability of precipitation in the study area is very small. In Fig. 12a we distinguish between low, medium, and high production years. These years are clearly related to total precipitation amounts during the February–June period (50 to 150 mm of precipitation are associated with years of low production, 150 to 250 mm correspond to years of medium production and more than 250 mm yields high production). Rainfall is related to pasture growth through an associated increase in soil moisture available for uptake. While precipitation is related to production in a somewhat linear relationship, soil moisture is related to pasture production in a nonlinear, approximately sigmoidal relationship (Fig. 12b) that starts to reveal the effects of the heterogeneity of the terrain. Figure 12b suggests that the precipitation amounts only have a scaling effect on the relationship between soil moisture and pasture production. The functional form of this relationship or the ability of soil moisture to explain pasture production remains relatively unchanged.

Unlike rainfall, the distribution of soil moisture is affected by the heterogeneity of the terrain, but the strength of this effect is proportional to the amount of soil moisture, which is partially controlled by the amount of precipitation. For instance low local slopes drive soil moisture by reducing flow velocity and by increasing the opportunity for infiltration, therefore high production tends to be found in flatter areas of the terrain (Fig. 12c). The effect of the slope, though, is stronger during wetter years when soil moisture is higher and there is more opportunity for overland and subsurface

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

redistribution of water. For drier years the ability of the local slope to explain the spatial variance of production decreases (Fig. 12c).

The relative position of a location in the drainage network, as defined by its upstream catchment area, is a non-local topographic control that also has a strong role in explaining the distribution of pasture production. More water is potentially drained at locations with a larger upstream catchment area, making them more prone to have a higher soil moisture content. Indeed, the production of a location increases with its upstream catchment area (Fig. 12d). Local drainage is defined by the small scale topographic features of the surface that form a convergent network. During years of low precipitation, concentration of moisture in converging areas of the drainage network produces a very contrasting spatial distribution of pasture production. The strength of this topographic control during dry years can be assessed by its relatively high explanatory power of the total spatial variance of pasture production. For increasingly wetter years, the strength of this topographic control wanes and with it its explanatory power (Fig. 12d). The contribution of upstream inflows to total local soil moisture decreases as incident precipitation increases. This reduces the influence of the non-local topographic controls.

Overall, during years of abundant production of pasture the importance of upstream water inflows tend to be overwhelmed by relatively large inputs of precipitation. In these conditions local topographic controls such as low slopes that reduce local water drainage rates have a relatively higher influence in the observed pasture productivity. As precipitation inputs are reduced the importance of the lateral redistribution of water becomes more relevant and non-local controls such as the upstream drainage area becomes increasingly more explanatory of the distribution of pasture.

HESSD

10, 15167–15217, 2013

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

5 Conclusions

Ecohydrological spatially-distributed models in conjunction with statistical weather generators are effective tools for simulating long-term pasture production dynamics and hydrologic conditions in semiarid rangelands, characterized by high spatial and temporal climatic and hydrologic variability. Results from this study contribute to insight into the hydrologic and climatic controls that determine the spatial and temporal distribution of grasses and the expected range of pasture production in different areas at the watershed scale.

This study aims at informing rangeland management and promoting the sustainability of grasslands. Spatially, the general physiographic characteristics of the terrain are good predictors of pasture yield, but the distribution of the canopy overstory is also important. Valley bottoms and flat areas adjacent to slopes, which tend to have relatively high soil moisture contents, had the highest production in the study area. Tree canopy cover was found to be negatively related with pasture production, reflecting the importance of rainfall and light interception, as well as water consumption by trees, in the development of a grassy understory in semiarid rangelands.

The simulated pasture production in the study catchment ranged from 21 to 1030.9 kg ha⁻¹ yr⁻¹, which ranks it as a medium to low productivity compared to other Mediterranean rangelands. With the calculated yields, the introduction of supplemental fodder is necessary to maintain livestock. Although the interannual distribution of precipitation is a strong control on the variability of pasture yield, its seasonal distribution during the year is as important. Specifically, years with low rainfall from February to May showed limited yield even for years with relatively high annual precipitation.

The importance of topographic structure of the landscape, as captured by the accumulated drainage area, becomes more relevant to explain the spatial distribution of pasture during years of low precipitation. This is because water inflows associated with lateral redistribution processes become a larger proportion of the total inflow into a location due to reduced precipitation inputs. The influence of lateral redistributions of

HESSD

10, 15167–15217, 2013

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Campos-Palacín, P.: Towards a sustainable global economics approach for Mediterranean agroforestry systems, in: Sustainability of Agrosilvopastoral Systems. Dehesas, Montados. Advances in Geocology, edited by: Schnabel, S. and Ferreira, A., Catena Verlag, Reiskirchen, Germany, 13–28, 2004.

5 Caylor, K. K., Manfreda, S., and Rodriguez-Iturbe, I.: On the coupled geomorphological and ecohydrological organization of river basins, Adv. Water Resour., 28, 69–86, doi:10.1016/j.advwatres.2004.08.013, 2005.

Caylor, K. K., Scanlon, T. M., and Rodriguez-Iturbe, I.: Ecohydrological optimization of pattern and processes in water-limited ecosystems: a trade-off-based hypothesis, Water Resour. Res., 45, W08407, doi:10.1029/2008wr007230, 2009.

10 Ceballos, A. and Schnabel, S.: Hydrological behaviour of a small catchment in the *dehesa* landuse system (Extremadura, SW Spain), J. Hydrol., 210, 146–160, doi:10.1016/S0022-1694(98)00180-2, 1998.

15 Cerdá, A., Schnabel, S., Ceballos, A., and Gómez-Amelia, D.: Soil hydrological response under simulated rainfall in the *dehesa* land system (Extremadura, SW Spain) under drought conditions, Earth Surf. Proc. Land., 23, 195–209, doi:10.1002/(SICI)1096-9837(199803)23:3<195::AID-ESP830>3.0.CO;2-I, 1998.

20 Clark, S. G., Austen, E. A., Prance, T., and Ball, P. D.: Climate variability effects on simulated pasture and animal production in the perennial pasture zone of south-eastern Australia, 1. Between year variability in pasture and animal production, Aust. J. Exp. Agr., 43, 1211–1219, doi:10.1071/EA02101, 2003.

Cobos, D. R. and Chambers, C.: Calibrating ECH2O Soil Moisture Sensors, Decagon Device, Application Note, available at: www.decagon.com (last access: November 2013), 2010.

25 Cox, P. M., Huntingford, C., and Harding, R. J.: A canopy conductance and photosynthesis model for use in a GCM land surface scheme, J. Hydrol., 212–213, 79–94, doi:10.1016/S0022-1694(98)00203-0, 1998.

Cox, P. M., Betts, R. A., Bunton, C. B., Essery, R. L. H., Rowntree, P. R., and Smith, J.: The impact of new land surface physics on the GCM simulation of climate and climate sensitivity, Clim. Dynam., 15, 183–203, doi:10.1007/s003820050276, 1999.

30 Cubera, E. and Moreno, G.: Effect of land use on soil water dynamic in *dehesas* of Central-Western Spain, Catena, 71, 298–308, doi:10.1016/j.catena.2007.01.005, 2007a.

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Cubera, E. and Moreno, G.: Effect of single *Quercus ilex* trees upon spatial and seasonal changes in soil water content in dehesas of central western Spain, *Ann. For. Sci.*, 64, 355–364, doi:10.1051/forest:2007012, 2007b.

5 Chow, V. T., Maidment, D. R., and Mays, L. W.: Applied Hydrology, McGraw-Hill, New York, xiii, 572 pp., 1988.

Dubrovský, M., Buchtele, J., and Žalud, Z.: High-frequency and low-frequency variability in stochastic daily weather generator and its effect on agricultural and hydrologic modelling, *Climate Change*, 63, 145–179, doi:10.1023/B:CLIM.0000018504.99914.60, 2004.

FAO: FAO-UNESCO soil map of the World, Technical Report 60, FAO, Rome, 1988.

10 Fatichi, S., Ivanov, V. Y., and Caporali, E.: Simulation of future climate scenarios with a weather generator, *Adv. Water Resour.*, 34, 448–467, doi:10.1016/j.advwatres.2010.12.013, 2011.

Fatichi, S., Ivanov, V. Y., and Caporali, E.: A mechanistic ecohydrological model to investigate complex interactions in cold and warm water-controlled environments: 1. Theoretical framework and plot-scale analysis, *J. Adv. Model. Earth Syst.*, 4, M05002, doi:10.1029/2011MS000086, 2012.

15 Foken, T.: Micrometeorology, Springer, Berlin, xix, 306 pp., 2008.

García-Estringana, P., Latron, J., Llorens, P., and Gallart, F.: Spatial and temporal dynamics of soil moisture in a Mediterranean mountain area (Vallcebre, NE Spain), *Ecohydrology*, 6, 741–753, doi:10.1002/eco.1295, 2013.

20 García-Ruiz, J. M. and Lana-Renault, N.: Hydrological and erosive consequences of farmland abandonment in Europe, with special reference to the Mediterranean region – a review, *Agr. Ecosyst. Environ.*, 140, 317–338, doi:10.1016/j.agee.2011.01.003, 2011.

Gea-Izquierdo, G., Montero, G., and Cañellas, I.: Changes in limiting resources determine spatio-temporal variability in tree–grass interactions, *Agroforest. Syst.*, 76, 375–387, doi:10.1007/s10457-009-9211-4, 2009.

25 Gindel, I.: Seasonal fluctuations in soil moisture under the canopy of xerophytes and in open areas, *Commonw. Forest. Rev.*, 43, 219–234, 1964.

Gómez-Gutiérrez, Á., Schnabel, S., and Lavado-Contador, J. F.: Gully erosion, land use and topographical thresholds during the last 60 years in a small rangeland catchment in SW Spain, *Land Degrad. Dev.*, 20, 535–550, doi:10.1002/ldr.931, 2009.

30 Gómez Gutiérrez, J. M. and Luis Calabuig, E.: Producción de praderas y pastizales, in: *El Libro de las Dehesas Salmantinas*, edited by: Gómez Gutiérrez, J. M., Junta de Castilla y León, Salamanca, 489–511, 1992.

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- González, F., Schnabel, S., Prieto, P. M., Pulido-Fernández, M., and Gragera-Facundo, J.: Pasture productivity in dehesas and its relationship with rainfall and soil, in: Nuevos retos de la ganadería extensiva: un agente de conservación en peligro de extinción, edited by: Canals Tresserras, R. M. and San-Emérito-Garciandía, L., Sociedad Española para el Estudio de los Pastos, Navarra, 37–43, 2012.
- Hoff, C. and Rambal, S.: An examination of the interaction between climate, soil and leaf area index in a *Quercus ilex* ecosystem, *Ann. For. Sci.*, 60, 153–161, doi:10.1051/forest:2003008, 2003.
- Houérou, H. N. and Hoste, C. H.: Rangeland production and annual rainfall relations in the Mediterranean Basin and in the African Sahelo-Sudanian Zone, *J. Range Manage.*, 30, 181–189, 1977.
- Infante, J. M., Domingo, F., Fernández-Alés, R., Joffre, R., and Rambal, S.: *Quercus ilex* transpiration as affected by a prolonged drought period, *Biol. Plantarum*, 46, 49–55, doi:10.1023/A:1022353915578, 2003.
- Istanbulluoglu, E., Wang, T., and Wedin, D. A.: Evaluation of ecohydrologic model parsimony at local and regional scales in a semiarid grassland ecosystem, *Ecohydrology*, 5, 121–142, doi:10.1002/eco.211, 2012.
- Ivanov, V. Y., Bras, R. L., and Curtis, D. C.: A weather generator for hydrological, ecological, and agricultural applications, *Water Resour. Res.*, 43, WR005364, doi:10.1029/2006WR005364, 2007.
- Ivanov, V. Y., Bras, R. L., and Vivoni, E. R.: Vegetation-hydrology dynamics in complex terrain of semiarid areas: 1. A mechanistic approach to modeling dynamic feedbacks, *Water Resour. Res.*, 44, W03429, doi:10.1029/2006WR005588, 2008.
- Jarvis, P. G.: The interpretation of the variations in leaf water potential and stomatal conductance found in canopies in the field, *Philos. T. Roy. Soc. B*, 273, 593–610, 1976.
- Jencso, K. G., McGlynn, B. L., Gooseff, M. N., Wondzell, S. M., Bencala, K. E., and Marshall, L. A.: Hydrologic connectivity between landscapes and streams: transferring reach and plot scale understanding to the catchment scale, *Water Resour. Res.*, 45, W04428, doi:10.1029/2008WR007225, 2009.
- Joffre, R. and Rambal, S.: Soil water improvement by trees in the rangelands of southern of Spain, *Oecologia*, 9, 405–422, 1988.
- Joffre, R. and Rambal, S.: How tree cover influences the water balance of mediterranean rangelands, *Ecology*, 74, 570–582, 1993.

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Joffre, R., Vacher, J., De los Llanos, C., and Long, G.: The dehesa: an agrosilvopastoral system of the mediterranean region with special reference to the Sierra Morena area of Spain, *Agroforest. Syst.*, 6, 71–96, doi:10.1007/BF02220110, 1988.
- Landsberg, J. J. and Waring, R. H.: A generalised model of forest productivity using simplified concepts of radiation-use efficiency, carbon balance and partitioning, *Forest Ecol. Manage.*, 95, 209–228, doi:10.1016/S0378-1127(97)00026-1, 1997.
- Lavado-Contador, J. F., Schnabel, S., and Trenado-Ordóñez, R.: Comparison of recent land use and land cover changes in two Dehesa agrosilvopastoral landuse systems, SW Spain, in: *Agrosilvopastoral Systems, Dehesas and Montados, Advances in Geocology*, 37, edited by: Schnabel, S. and Ferreira, A., Cáceres, Spain, 55–69, 2004.
- Lavado-Contador, J. F., Maneta, M., and Schnabel, S.: Prediction of near-surface soil moisture at large scale by Digital Terrain Modeling and Neural Networks, *Environ. Monit. Assess.*, 121, 211–230, doi:10.1007/s10661-005-9116-2, 2006.
- Lozano-Parra, F. J., Schnabel, S., and Ceballos-Barbancho, A.: Dinámica del agua del suelo en dehesa bajo diferentes cubiertas vegetales, resultados preliminares, in: *Estudios de la Zona No Saturada del Suelo*, edited by: Martínez-Fernández, J. and Sanchez Martín, N., Universidad de Salamanca, Salamanca, 47–52, 2011.
- Maneta, M.: Modelling of the hydrologic processes in a small semiarid catchment, Ph. D. Thesis, Geography and Land Planning Department, University of Extremadura, Cáceres, 278 pp., 2006.
- Maneta, M. and Silverman, N.: A spatially-distributed model to simulate water, energy and vegetation dynamics using information from regional climate models, *Earth Interact.*, 17, 1–44, doi:10.1175/2012EI000472.1, 2013.
- Maneta, M., Pasternack, G. B., Wallender, W. W., Jetten, V., and Schnabel, S.: Temporal instability of parameters in an event-based distributed hydrologic model applied to a small semiarid catchment, *J. Hydrol.*, 341, 207–221, doi:10.1016/j.jhydrol.2007.05.010, 2007.
- Maneta, M., Schnabel, S., and Jetten, V.: Continuous spatially distributed simulation of surface and subsurface hydrologic processes in a small semiarid catchment, *Hydrol. Process.*, 22, 2196–2214, doi:10.1002/hyp.6817, 2008a.
- Maneta, M., Schnabel, S., Wallender, W. W., Panday, S., and Jetten, V.: Calibration of an evapotranspiration model to simulate soil water dynamics in a semiarid rangeland, *Hydrol. Process.*, 22, 4655–4669, doi:10.1002/hyp.7087, 2008b.

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Martínez Fernández, J., Cano, A., Hernández-Santana, V., and Morán, C.: Evolución de la humedad del suelo bajo diferentes tipos de cubierta vegetal en la cuenca del Duero, Estudios en la Zona No Saturada VIII, University of Cordoba, Cordoba, 257–258, 2007.
- McGlynn, B. L., McDonnell, J., Stewart, M., and Seibert, J.: On the relationships between catchment scale and streamwater mean residence time, *Hydrol. Process.*, 17, 175–181, doi:10.1002/hyp.5085, 2003.
- Montaldo, N., Rondena, R., Albertson, J. D., and Mancini, M.: Parsimonious modeling of vegetation dynamics for ecohydrologic studies of water-limited ecosystems, *Water Resour. Res.*, 41, W10416, doi:10.1029/2005WR004094, 2005.
- Montero, G., San Miguel, A., and Cañellas, I.: System of Mediterranean silviculture “La Dehesa”, in: *Agricultura Sostenible*, edited by: Jiménez Díaz, R. M. and Lamo de Espinos, J., Mundi Prensa, Madrid, 1998.
- Montero, M. J., Moreno, G., and Bertomeu, M.: Light distribution in scattered-trees open woodlands in Western Spain, *Agroforest. Syst.*, 73, 233–244, doi:10.1007/s10457-008-9143-4, 2008.
- Moreno, G.: Response of understory forage to multiple tree effects in Iberian dehesas, *Agr. Ecosyst. Environ.*, 123, 239–244, doi:10.1016/j.agee.2007.04.006, 2008.
- Moreno, G. and Cubera, E.: Impact of stand density on water status and leaf gas exchange in *Quercus ilex*, *Forest Ecol. Manage.*, 254, 74–84, doi:10.1016/j.foreco.2007.07.029, 2008.
- Moreno, G., Obrador, J. J., Cubera, E., and Dupraz, C.: Fine root distribution in Dehesas of Central-Western Spain, *Plant Soil*, 277, 153–162, doi:10.1007/s11104-005-6805-0, 2005.
- Ogaya, R. and Peñuelas, J.: Phenological patterns of *Quercus ilex*, *Phillyrea latifolia*, and *Arbutus unedo* growing under a field experimental drought, *Ecoscience*, 11, 263–270, 2004.
- Oleson, K. W., Lawrence, D. M., Bonan, G. B., Flanner, M. G., and Kluzek, E.: Technical Description of version 4.0 of the Community Land Model (CLM), NCAR, Boulder, CO, doi:10.5065/D6FB50WZ, 2010.
- Paço, T. A., David, T. S., Henriques, M. O., Pereira, J. S., Valente, F., Banza, J., Pereira, F. L., Pinto, C., and David, J. S.: Evapotranspiration from a Mediterranean evergreen oak savannah: the role of trees and pasture, *J. Hydrol.*, 369, 98–106, doi:10.1016/j.jhydrol.2009.02.011, 2009.
- Panaïotis, C., Carcaillet, C., and M’Hamedi, M.: Determination of the natural mortality age of a holm oak (*Quercus ilex* L.) stand in Corsica (Mediterranean Island) *Oecologia*, 18, 519–530, 1997.

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Plaixats, J., Villareal, A., Bartolomé, J., and Espona, J.: Productivity characteristics of grassland in a dehesa system in Catalonia, Spain, in: Sustainability of Agrosilvopastoral Systems. Dehesas, Montados. Advances in Geocology, edited by: Schnabel, S. and Ferreira, A., Catena, Reiskirchen, Germany, 195–202, 2004.

5 Porporato, A., D'Odorico, P., Laio, F., Ridolfi, L., and Rodriguez-Iturbe, I.: Ecohydrology of water-controlled ecosystems, *Adv. Water Resour.*, 25, 1335–1348, doi:10.1016/s0309-1708(02)00058-1, 2002.

Richardson, C. W.: Stochastic simulation of daily precipitation, temperature, and solar radiation, *Water Resour. Res.*, 17, 182–190, doi:10.1029/WR017i001p00182, 1981.

10 Ricotta, C., Avena, G. C., and Teggi, S.: Relation between vegetation canopy surface temperature and the Sun-surface geometry in a mountainous region of central Italy, *Remote Sens.*, 18, 3091–3096, 1997.

Rivest, D., Rolo, V., López-Díaz, L., and Moreno, G.: Shrub encroachment in Mediterranean silvopastoral systems: *Retama sphaerocarpa* and *Cistus ladanifer* induce contrasting effects on pasture and *Quercus ilex* production, *Agr. Ecosyst. Environ.*, 141, 447–454, doi:10.1016/j.agee.2011.04.018, 2011.

Rodríguez-Iturbe, I.: Ecohydrology: a hydrologic perspective of climate–soil–vegetation dynamics, *Water Resour. Res.*, 36, 3–9, doi:10.1029/1999WR900210, 2000.

20 Rodríguez-Iturbe, I., D'Odorico, P., Porporato, A., and Ridolfi, L.: On the spatial and temporal links between vegetation, climate, and soil moisture, *Water Resour. Res.*, 35, 3709–3722, doi:10.1029/1999wr900255, 1999.

25 Sabaté, S., Gracia, C., and Sánchez, A.: Likely effects of climate change on growth of *Quercus ilex*, *Pinus halepensis*, *Pinus pinaster*, *Pinus sylvestris* and *Fagus sylvatica* forests in the Mediterranean region, *Forest Ecol. Manage.*, 5906, 1–15, doi:10.1016/S0378-1127(02)00048-8, 2002.

Santamaría, O., Poblaciones, M. J., Olea, L., Rodrigo, S., and García-White, T.: Efecto de fertilizantes alternativos al superfosfato de cal sobre la producción y calidad de pastos de dehesa en el suroeste de España, 5. Congreso Forestal Español, Ávila, 2009.

30 Schnabel, S.: Soil erosion and runoff production in small watershed under silvo-pastoral landuse (dehesas) in Extremadura, Spain, *Monografías Científicas, Geoforma, Logroño*, 163 pp., 1997.

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Schnabel, S.: La precipitación como factor en los procesos hidrológicos y erosivos, análisis de datos de Cáceres capital, in: *Hidrología y Erosión de Suelos*, edited by: Schnabel, S., Gómez-Amelia, D., and Ceballos-Barbancho, A., Norba, Revista de Geografía, Cáceres, 137–152, 1998.
- 5 Schnabel, S., Dahlgren, R. A., and Moreno, G.: Soil and water dynamics, in: *Mediterranean Oak Woodland Working Landscapes, Dehesas of Spain and Rangelands of California*, edited by: Campos, P., Huntsinger, L., Oviedo, J. L., Starrs, P. F., Díaz, M., Standiford, R., and Montero, G., Springer, New York, 2013a.
- Schnabel, S., Pulido Fernández, M., and Lavado-Contador, J. F.: Soil water repellency in range-
lands of Extremadura (Spain) and its relationship with land management, *Catena*, 103, 53–
61, doi:10.1016/j.catena.2011.11.006, 2013b.
- 10 Semenov, M. A. and Barrow, E. M.: Use of a stochastic weather generator in the development of climate change scenarios, *Climatic Change*, 35, 397–414, doi:10.1023/A:1005342632279, 1997.
- 15 Semenov, M. A. and Barrow, E. M.: LARS-WG: A Stochastic Weather Generator for Use in Climate Impact Studies, User Manual, available at: <http://www.rothamsted.ac.uk/mas-models/larswg.php> (last access: November 2013), 2002.
- Semenov, M. A. and Porter, J. R.: Climatic variability and the modelling of crop yields, *Agr. Forest Meteorol.*, 73, 265–283, doi:10.1016/0168-1923(94)05078-K, 1995.
- 20 Semenov, M. A., Brooks, R. J., Barrow, E. M., and Richardson, C. W.: Comparison of the WGEN and LARS-WG stochastic weather generators for diverse climates, *Clim. Res.*, 10, 95–107, 1998.
- Singh, V. P.: Kinematic wave modeling in water resources, in: *Environmental Hydrology*, Wiley Interscience, New York, 830 pp., 1997.
- 25 Swinbank, W. C.: Long-wave radiation from clear skies, *Q. J. Roy. Meteorol. Soc.*, 90, 488–493, doi:10.1002/qj.49708938105, 1964.
- Tague, C. L. and Band, L. E.: RHESSys: Regional Hydro-Ecologic Simulation System – an object-oriented approach to spatially distributed modeling of carbon, water, and nutrient cycling, *Earth Interact.*, 8, 1–42, doi:10.1175/1087-3562(2004)8<1:RRHSSO>2.0.CO;2, 2004.
- 30 Tubiello, F. N., Soussana, J. F., and Howden, S. M.: Crop and pasture response to climate change, *P. Natl. Acad. Sci. USA*, 104, 19686–19690, doi:10.1073/pnas.0701728104, 2007.
- Van Schaik, L.: Spatial variability of infiltration patterns related to site characteristics in a semi-arid watershed, *Catena*, 78, 36–47, doi:10.1016/j.catena.2009.02.017, 2009.

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Van Schaik, L.: The role of macropore flow from plot a catchment scale, *Faculteit Geowetenschappen, Universiteit Utrecht, Utrecht*, 174 pp., 2010.

Van Schaik, L., Schnabel, S., and Jetten, V.: The influence of preferential flow on hillslope hydrology in a semi-arid watershed (in the Spanish Dehesas), *Hydrol. Process.*, 22, 3844–3855, doi:10.1002/hyp.6998, 2008.

Vaz, M., Maroco, J., Ribeiro, N., Gazarini, L. C., Pereira, J. S., and Chaves, M. M.: Leaf-level responses to light in two co-occurring *Quercus* (*Quercus ilex* and *Quercus suber*): leaf structure, chemical composition and photosynthesis, *Agroforest. Syst.*, 82, 173–181, doi:10.1007/s10457-010-9343-6, 2011.

Viville, D. and Littlewood, I. G.: *Ecohydrologic Processes in Small Basins*, edited by: Viville, D. and Littlewood, I. G., UNESCO, Strasbourg, 199 pp., 1996.

Wang, L., Liu, J., Sun, G., Wei, X., Liu, S., and Dong, Q. (Eds.): *Water, climate, and vegetation: ecohydrology in a changing world*, Special Issue, *Hydrol. Earth Syst. Sci.*, 16, 4633–4636, doi:10.5194/hess-16-4633-2012, 2012.

White, M. A., Thornton, P. E., Running, S. W., and Nemani, R. R.: Parameterization and sensitivity analysis of the BIOME–BGC terrestrial ecosystem model: net primary production controls, *Earth Interact.*, 4, 1–85, doi:10.1175/1087-3562(2000)004<0003:PASAOT>2.0.CO;2, 2000.

Xia, J.: A stochastic weather generator applied to hydrological models in climate impact analysis, *Theor. Appl. Climatol.*, 55, 177–183, doi:10.1007/BF00864713, 1996.

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Set of model parameters included in the process of manual calibration.

Variable	Description	Unit	Final value		Source for initial values
			Tree	Pasture	
C_{NPP}	GPP to NPP conversion factor	–	0.25	0.35	Sabaté et al. (2002)
T_{opt}	Optimal Temperature for maximum plant growth	°C	15	18	Ogaya and Peñuelas (2004); and AEMET
T_{max}	Maximum Temperature for plant	°C	42.6	30	AEMET
T_{min}	Minimum Temperature for plant	°C	–5.6	2	AEMET
k_{sd}	Dry grass turnover rate	–	–	8.50E–07	adjusted
T_{ξ}	Temperature for enhanced grass decay	°C	–	18	adjusted
δ_t	Leaf Turnover Rate	s ^{–1}	1.40E–08	1.00E–07	Hoff and Rambal (2003)
σ_{LAI}	Specific Leaf Area	m ² (gC) ^{–1}	0.017	0.015	Vaz et al. (2011)
ξ_w	Vegetation water use efficiency	gCm ^{–1}	1150	6000	Hoff and Rambal (2003)
$X_{stor\ max}$	Maximum canopy water storage per unit LAI	m	0.00075	0.00015	White et al. (2000)
g_{cmax}	Maximum stomatal conductance	ms ^{–1}	0.0063	0.035	White et al. (2000)
θ_{wp}	Volumetric soil moisture content at wilting point	m ³ m ^{–3}	0.05	0.165	Van Schaik (2010)
K_{eff}^*	Effective hydraulic conductivity of the soil	ms ^{–1}	0.00479–0.00053		measured
η^*	Soil Porosity	0–1	0.50–0.26		measured
λ^*	Brooks and Corey exponent parameter	–	0.33–0.20		adjusted

* Values vary spatially.

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Table 3. Goodness-of-fit between observed and simulated weather data. K–S = Kolmogorov–Smirnov test; * = Example data: Obs. = Observed average values from the study catchment (2000–2012); Sim. = Simulated average values for 300 yr.

	Rainfall*		Rainfall Temperature		Maximum Temperature		Minimum Radiation		Short Wave	
	Obs.	Sim.	K–S	p value	K–S	p value	K–S	p value	K–S	p value
Jan	45.0	44.4	0.033	1.000	0.053	1.000	0.106	0.999	0.044	1.000
Feb	52.5	60.7	0.042	1.000	0.106	0.999	0.106	0.999	0.087	1.000
Mar	43.1	45.1	0.035	1.000	0.053	1.000	0.053	1.000	0.000	1.000
Apr	44.2	45.8	0.061	1.000	0.106	0.999	0.106	0.999	0.087	1.000
May	39.3	47.3	0.054	1.000	0.053	1.000	0.106	0.999	0.087	1.000
Jun	12.7	11.7	0.063	1.000	0.106	0.999	0.106	0.999	0.131	0.982
Jul	0.5	0.7	0.497	0.004	0.106	0.999	0.106	0.999	0.087	1.000
Aug	6.5	8.4	0.209	0.643	0.106	0.999	0.106	0.999	0.131	0.982
Sep	25.1	24.4	0.154	0.927	0.053	1.000	0.053	1.000	0.044	1.000
Oct	95.5	82.5	0.098	1.000	0.105	0.999	0.106	0.999	0.044	1.000
Nov	61.2	72.8	0.030	1.000	0.053	1.000	0.105	0.999	0.043	1.000
Dec	62.2	64.8	0.040	1.000	0.106	0.999	0.053	1.000	0.044	1.000

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Table 4. Descriptive statistics of observed (Obs.) and simulated (Sim.) series and quality parameters of the model. n = sample size; RMSE = Root Mean Square Error; * Values only showed for 2011 because it is the more monitored year.

	n	Average		Maximum		Minimum		Standard Deviation		r^2	RMSE	Bias	Nash–Sutcliffe
		Obs.	Sim.	Obs.	Sim.	Obs.	Sim.	Obs.	Sim.				
Soil Moisture ($\text{m}^3 \text{m}^{-3}$)													
Site 1	1268	0.219	0.202	0.417	0.430	0.060	0.075	0.108	0.091	0.85	0.047	0.018	0.81
Site 2	1267	0.222	0.212	0.451	0.440	0.074	0.083	0.114	0.094	0.90	0.040	0.010	0.88
SMS-3	848	0.165	0.151	0.312	0.349	0.066	0.068	0.069	0.061	0.80	0.034	0.014	0.75
Soil Temperature ($^{\circ}\text{C}$)													
Site 1	1274	18.0	19.8	37.0	47.1	−2.0	2.5	10.2	10.0	0.89	3.78	−1.8	0.86
Site 2	1267	18.1	19.0	33.4	42.7	3.2	1.9	8.2	9.5	0.91	3.08	−0.9	0.86
Pasture Production (kg DM ha^{-1})													
Site 1	20*	603.3	588.1	1319.3	1368.7	269.0	319.0	396.2	310.2	0.84	164.8	15.2	0.82
Site 2	20*	644.3	673.6	1392.7	1432.5	293.4	361.5	395.3	317.4	0.76	193.4	−29.3	0.75

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Climate and topographic controls on pasture production

J. Lozano-Parra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 5. Descriptive statistics for simulated rainfall (mm) and simulated average pasture production ($\text{kgDMha}^{-1}\text{yr}^{-1}$) for each site and 300 yr.

	<i>n</i>	Mean	Maximum	Minimum	Percentile			SD
					25	50	75	
Rainfall	300	508.7	934.1	188.9	426.7	503.7	571.9	118.2
Site 1	300	338.0	977.6	20.7	210.0	305.9	445.1	172.5
Site 2	300	456.0	1030.9	29.9	319.9	435.4	570.6	182.8

Climate and topographic controls on pasture production

J. Lozano-Parra et al.




Fig. 1. Location of the study catchment and the equipment.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2. Maps of catchment properties: **(A)** slope (mm^{-1}), **(B)** soil depth (m), **(C)** porosity (0–1), **(D)** flux accumulation (number of pixels that spill on another), **(E)** tree density (trees ha^{-1}), **(F)** tree canopy cover (0–1).

Climate and topographic controls on pasture production

J. Lozano-Parra et al.




Fig. 3. Observed and simulated soil moisture from March 2009 until September 2012. **(A)** Site 1; **(B)** Site 2; **(C)** SMS-3. Black line are measured values, and red line are simulated values.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Climate and topographic controls on pasture production

J. Lozano-Parra et al.




Fig. 4. Observed and simulated soil temperature from March 2009 until September 2012. **(A)** Site 1; **(B)** Site 2; Black line are measured values, and red line are simulated values.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Climate and topographic controls on pasture production

J. Lozano-Parra et al.




Fig. 5. Observed and simulated accumulated pasture production at **(A)** Site 1; and **(B)** Site 2. The red line represents simulated average pasture yield for whole pixels in every Site, with ± 1 standard deviation (green shade). Green circles represent average pasture production based on height measurements; blue rhombuses represent average pasture production based on plant cuts (moustaches correspond to \pm standard deviation).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Climate and topographic controls on pasture production

J. Lozano-Parra et al.




Fig. 6. Simulated transpiration during 4 hydrological years (2008–2012) for **(A)** *Quercus ilex* in *Site 1*, and **(B)** natural pastures in *Site 1* and *Site 2*.




Fig. 7. Spatial distribution of annual average soil moisture ($\text{m}^3 \text{m}^{-3}$) (**A**) and its standard deviation (**B**), and annual average evapotranspiration (**C**) (mm).

Climate and topographic controls on pasture production

J. Lozano-Parra et al.




Fig. 8. Simulated average pasture production and precipitation at *Site 1*, **(A)** at the annual timescale. **(B)** for ten years at the daily timescale (the green shade represents ± 1 standard deviation of pasture production, and the blue bars is the rainfall).

Climate and topographic controls on pasture production

J. Lozano-Parra et al.




Fig. 9. Simulated average pasture production and precipitation at *Site 2*, **(A)** at the annual timescale. **(B)** for ten years at the daily timescale (the green shade represents ± 1 standard deviation of pasture production, and the blue bars is the rainfall).

Climate and topographic controls on pasture production

J. Lozano-Parra et al.




Fig. 10. Spatial distribution of simulated pasture production (kg DM ha⁻¹): **(A)** average; **(B)** standard deviation; **(C)** maximum; **(D)** minimum.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Climate and topographic controls on pasture production

J. Lozano-Parra et al.




Fig. 12. Climate and physiographic factors that influence pasture production.

[Title Page](#)
[Abstract](#) | [Introduction](#)
[Conclusions](#) | [References](#)
[Tables](#) | [Figures](#)
[⏪](#) | [⏩](#)
[⏴](#) | [⏵](#)
[Back](#) | [Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)