

Impact of long-term drainage on summer groundwater flow patterns in the Mer Bleue peatland, Ontario, Canada

Benjamin J. Kopp¹, Jan H. Fleckenstein^{2,3}, Nigel T. Roulet⁴, Elyn Humphreys⁵, and Christian Blodau^{1,4,*}

[1] Limnological Research Station and Department of Hydrology, University of Bayreuth, Bayreuth, Germany

[2] Department of Hydrology, University of Bayreuth, Bayreuth, Germany

[3] present address: Department of Groundwater Research, Centre for Environmental research – UFZ, Halle-Leipzig, Germany

[4] Department of Geography and the Global Environmental and Climate Change Centre, McGill University, Montreal, Canada

[5] Department of Geography and Environmental Studies, Carleton University, Ottawa, Canada

* now at: Hydrology Group, Institute of Landscape Ecology, University of Münster, Germany


Correspondence to C. Blodau (christian.blodau@uni-muenster.de)

1 1 SUPPLEMENTARY INFORMATION

2 We installed additional piezometer nests at 45 m distance from the ditch and 30 m
3 distance from the transect to obtain information based on hydrologic triangles. In general,
4 identified groundwater flow patterns were supported by these piezometer nests. At 45 m
5 distance from the ditch at the bog side, hydraulic heads indicated a prevailing lateral flow
6 pattern combined with a vertical component alternating between upward and downward
7 gradients down to 2.0 m depth. Hydraulic potentials in 3.0 m depth were > 20 cm lower
8 than in 2.0 m depth. This indicated a downward orientated flow in deeper catotelm peat
9 (Fig. S 1) which is coherent with modelled hydraulic potentials using the MODFLOW
10 model as described in the paper. In the same distance from the ditch under forest the
11 hydraulic heads remained upward directed throughout the study period (Fig. S 2).


12 As a reference site, we installed an additional piezometer nest at a distance of 600 m
13 from the ditch to monitor hydraulic heads at a pristine part of the bog, which can thus be
14 assumed to be unaffected by drainage. Maximum water table fluctuations of 6.1 cm were
15 rather small and similar to manual measured fluctuations of 6.3 cm at the 200 m site
16 (data not shown). Hydraulic heads at the 600 m site were dominated by lateral flow with
17 a discharging vertical flow component until the end of August and slightly downward
18 orientated flow established in the beginning of September (Fig. S 3).

19



20


21 Fig. S 1 Manual measurements of hydraulic potentials at the piezometer nest in 45 m distance from
 22 the ditch in the bog.



23

24 Fig. S 2 Manual measurements of hydraulic potentials at the piezometer nest in 45 m distance from
 25 the ditch in the forest.

26



27

28 Fig. S 3 Manual measurements of hydraulic potentials at the piezometer nest in 600 m distance from
 29 the ditch in the open bog.

30

31

32