

1 Dear Dr. Schaepli (Editor), dear Prof. Dr. Leibundgut (Reviewer #1), dear Dr. Bakolowicz  
2 (Reviewer #2) and dear Reviewers #3 and #4.

3

4 *We<sup>(1)</sup> would like to thank you for your valuable comments and for the positive evaluation of*  
5 *our manuscript entitled “Identification of glacial melt water runoff in a karstic environment*  
6 *and its implication for present and future water availability. In the following we will comment*  
7 *on each evaluation separately, replying to every comment raised by the editor and the four*  
8 *reviewers. We are confident that after the requested minor revisions you will find our*  
9 *manuscript suitable for publication in HESS.*

10

## 11 **1 B. Schaepli (Editor)**

12 Received and published: 1 May 2013

13 All reviewers judge de significance, the scientific and presentation quality of this paper as  
14 being good to excellent. They raise an important number of detailed comments and critics, but  
15 overall, they recommend publication of this paper with minor revisions rather than a  
16 substantial re-writing or expansion of the manuscript.

17 Before preparing the revised manuscript, I would like to invite the authors to give a detailed  
18 response to each of the reviews. Hereafter, I give a short overview of the main comments:

19 - Reviewer 2 and 3 highlighted that the literature review seems incomplete.

20 - Reviewer 2 would also like to see a better presentation of the functioning of subglacial karst  
21 hydrology.

22 - Reviewer 3 suggest to improve the terminology and to better discuss the limitations and  
23 transferability of the results.

24 - Reviewer 4 made critical comments on the over-ambitious scope of the paper and  
25 recommends reducing the scope and setting a more modest aim of establishing the

26 hydrogeology of a glacierised karst depression. A better definition of the scope of the  
27 manuscript might also be required in view of the comments of M. Bakalowicz and the ensuing  
28 discussion.

29 - Besides many other detailed comments, reviewer 4 also asks for a better integration of all  
30 obtained results and avoiding conclusions that cannot be directly drawn from the analyses of  
31 this paper.

32 - Finally, several reviewers find the manuscript not yet very well structured and mention  
33 repetitions. And reviewer 1 would like to have more precise indications about the original  
34 contributions of this paper.

35

36 *We thank the editor for the positive evaluation and the concise summary of the four reviews.*

37 *The objective of our study is to identify current and future water availability in the Plaine*

38 *Morte region by combining meteo-hydrological observations, tracer experiments, isotopic*

39 *investigations, karst modeling and projections of glacier melt runoff. This interdisciplinary*

40 *approach represents a novel and heuristic approach of how numerous datasets from different*

41 *disciplines can be interpreted. While the results are surely only valid for the Plaine Morte*

42 *Region, the interdisciplinary approach can be applied to any similar mountainous study site*

43 *and certainly is a valuable asset for water resources research. During the revision process we*

44 *will do in particular the following:*

45 - *Incorporate suggested and relevant peer-reviewed literature*

46 - *Revise the description of the subglacial karst hydrology*

47 - *Revise the terminology and the discussion in order to clarify the limitations and*  
48 *transferability of our findings*

49 - *Revise the wording of the scope of the paper. It is our firm belief, that science lives*  
50 *from innovative approaches and that there should also be space for reasonable and*  
51 *well-founded hypothesis to assess future water availability. We will revise the*

52 *manuscript to demonstrate that our approach allows us to make assumptions about*  
53 *current and future flow paths and assess water availability in the Plaine Morte region.*  
54 *All climate change projections are just hypothesis based on founded assumptions.*  
55  
56

## 57 **2 C. Leibundgut**

58 Received and published: 8 April 2013

59 Overall assessment

60 Scientific Significance: excellent

61 The manuscript represents a substantial contribution to scientific progress within the scope of  
62 Hydrology and Earth System Sciences by its holistic approach dealing from hydrology up to  
63 the complex regional water resources systems.

64 Scientific Quality: excellent

65 The scientific approach and applied methods are valid. Also the results are discussed in an  
66 appropriate and balanced way.

67 Presentation Quality: excellent

68 The scientific results and conclusions are presented in a clear, concise, and wellstructured  
69 way. The number and quality of figures/tables is appropriate. I am not competent to perform  
70 an English language check.

71 Further aspects:

72 The paper address relevant scientific questions within the scope of HESS by presenting novel  
73 concepts (holistic), ideas, tools (combination of methods and techniques and a lot of new data.

74 There are substantial conclusions reached as well scientifically (i.e. karst model) and regionally  
75 (i.e. glacier runoff).

76 The scientific methods and assumptions are valid and clearly outlined. In particular the  
77 validation occurs with independent data (i.e. tracer data) which is a modern concept but not  
78 always used.

79 Without doubt the results are suitable and sufficient to support the given interpretations and  
80 conclusions.

81 The description of experiments and calculations is clear, supported by the tables and figures,  
82 is complete and precise to allow their reproduction by fellow scientists. The authors give  
83 proper credit to related work. The reference “Tracer experiments in temperate alpine glaciers”  
84 in Leibundgut Ch., P. Malozewski, Ch.Külls: Tracers in Hydrology. Wiley-Blackwell, 2009:  
85 310-321 would address directly the issue on page 2746.

86 It would be an improvement if they clearly indicate their own new/original contribution more  
87 explicitly.

88 The title reflects clearly the contents of the paper. The abstract provide a concise and  
89 complete summary.

90 The overall presentation is well structured and clear with a small restriction: there are some  
91 redundancies probably due to the single contributions provided by the single author groups.

92 All mathematical formulae, symbols, abbreviations, and units are correctly defined and used.

93 Should any parts of the paper (text, formulae, figures, tables) be clarified, reduced, combined,  
94 or eliminated? : no, except the mentioned reduncances.

95 The number and quality of references are appropriate. I made a proposal for a supplement  
96 above.

97 The amount and quality of supplementary material is appropriate.

98

99 *We would like to thank Prof. Dr. Leibundgut for his positive evaluation of our manuscript. It*  
100 *honors us that the significance, the quality and the topic presentation of our manuscript were*

101 *evaluated as being “excellent”. This evaluation is even more honoring as we are aware of the*  
102 *immense knowledge on karst hydrology of Prof. Dr. Leibundgut.*  
103 *Certainly, we will include the peer-reviewed reference Leibundgut (2009) and will*  
104 *incorporate its content into the manuscript. We will also outline and emphasis on the*  
105 *new/original contributions to the field of tracer hydrology by emphasizing the importance of*  
106 *interdisciplinary research.*

107

### 108 **3 M. Bakalowicz**

109 *We also would like to thank Dr. Bakalowicz for his valuable comments. As Dr. Bakalowicz*  
110 *has raised several concerns regarding our manuscript we will comment on each of his*  
111 *comments specifically below:*

112

113 General comments

114 This is a really interesting, but complicated issue, seldom considered, for two reasons:  
115 (i) combining glacial and karst hydrology looks to be a difficult challenge; and (ii) generally  
116 there is no practical interest what does not push scientists to work on. So the bibliography is  
117 relatively scarce. All the more reason to try to make a review as comprehensive as possible! I  
118 regret to tell that the authors seem to ignore a large part of the literature about karst and  
119 glacial hydrology. I think that they could give to their work a broader scope than in its present  
120 state. Glacial hydrology presents characteristics very close to karst hydrology to such an  
121 extent that the best synthesis on glacial hydrology was written by two karst specialists, A.  
122 Eraso and M. Pulina.  
123 - Eraso A. and Pulina M. 2011. Cuevas en hielo y rios bajo los glaciares. McGraw Hill, 2nd  
124 ed., 280 p. (and GLACKMA, 3rd ed., 300 p.). See <http://www.glackma.es/>

125 - Works of the International Committee “Glacier caves and karst in Polar Regions”  
126 (GLACKIPR) created in 1989 in Budapest, during the 10th International Congress of  
127 Speleology. They were published in symposium proceedings, the first one in 1990 in Madrid,  
128 Spain, and the 8th in 2007 in Katowice, Poland.

129 *We agree with Dr. Bakalowicz. During the revision of the ms we will perform an extensive*  
130 *literature research including the work by A. Eraso and M. Pulina. Nevertheless, in*  
131 *accordance with the comment of reviewer #3 and the HESS guidelines, we will primarily*  
132 *reference peer reviewed articles, and only refer to conference proceedings if specific aspects*  
133 *of our case study (area of Glacier de la Plaine Morte) make it necessary.*

134

135 C.C. Smart is probably the pioneer of subglacial karst hydrology. I agree that his PhD was  
136 unpublished, but he published some interesting papers which could have helped the authors in  
137 a better presentation of the relations between the glacier and the underneath karst and in a  
138 more generalized approach.

139 - Smart C.C. 1983. Hydrology of glacierised alpine karst. PhD, McMaster University, 343 p.

140 - Smart C.C. 1983. The hydrology of the Castelguard Karst, Columbia Icefields, Alberta,  
141 Canada. Arctic and Alpine Research, 15 (4): 471-486.

142 - Smart C.C. 1997. Hydrogeology of glacial and subglacial karst aquifers: Small River,  
143 British Columbia, Canada. Proc. 6th Conference on Limestone Hydrology and Fissured  
144 Media, La Chaux-de-Fonds, Switzerland, p. 315-318.

145 I also suggest to read the following references:

146 - Ford D.C. and Williams P.W. 2007. Karst hydrogeology and geomorphology. Wiley, 562 p.

147 See especially 10.3. The cold extreme: karst development in glaciated terrains, p. 410-421.

148 - Lauritzen S.E. 1984. Evidence of subglacial karstification in Glomdal, Svartisen. Norsk  
149 Geografisk Tidsskrift, 38 (3-4): 169-170.

150 - Lauritzen S.E. 1986. Kvithola at Fauske, northern Norway: an example of ice-contact  
151 speleogenesis. Norsk Geografisk Tidsskrift, 66: 153-161.

152 These references could help the authors in presenting in a better way the functioning of  
153 subglacial karst hydrology, and particularly the hydraulic connection through glacier  
154 “moulins” between the seasonal flows at the surface of the glacier and the karst sinkholes  
155 below it. This is clear from figure 11, but unclear at all in the 1st paragraph of 2.3 Hydrology,  
156 characteristic (i). This is a particularly important point which needs to be detailed, because  
157 some tracing tests combine glacial and karst flows and the contribution of glacial water looks  
158 important during summer melting.

159 *In accordance with the comment from reviewer #3 we will search the scientific databases for*  
160 *the work by Smart C.C and incorporate peer reviewed literature relevant to our study.*

161 *Nevertheless, in accordance to the comment posted by Prof. Dr. Leibundgut and in*  
162 *accordance with the scope of our study we cannot focus on subglacial karst hydrology only,*  
163 *as this is only one aspect of our scope. We recall that the scope of our study is to assess*  
164 *present and future water availability in the Plaine Morte region – not to investigate specific*  
165 *subglacial karst processes.*

166

167 Furthermore the manuscript is sometimes confusing, difficult to follow. I make some  
168 suggestions in order to help the authors in improving their manuscript. First, this Chapter 2.3  
169 Hydrology should give as an introduction the different flow types described in karst and  
170 glacial hydrology, from the literature.

171 Then the paragraph could be re-written for instance in that way:

172 “Discharge from snow and ice melt, as well as rainfall runoff is characterized by different  
173 mechanisms depending on field characteristics in the Plaine Morte area: (i) water storage in  
174 the glacier and the snow fields, (ii) rapid subsurface flow under gravel covered soil, (iii) low  
175 flow in fertile agricultural soil, (iv) retention in natural swamps, (v) storage in natural and

176 artificial lakes, (vi) seasonal streams running on the glacier, partly swallowed in “moulins”  
177 and crevices connected to sinkholes underneath the glacier, and (vii) a well-developed karst  
178 system which drains melt water to karstic springs at lower elevations.”

179 *We will edit the manuscript as follows:*

180 “Discharge from snow and ice melt, as well as rainfall runoff is characterized by different  
181 mechanisms depending on field characteristics in the Plaine Morte area: (i) water storage in  
182 the glacier and the snow fields, (ii) rapid subsurface flow under gravel-covered soil, (iii) low  
183 flow in fertile agricultural soil, (iv) retention in natural swamps, (v) storage in natural and  
184 artificial lakes, (vi) seasonal streams running on the glacier, primarily swallowed in “moulins”  
185 and crevasses partly connected to sinkholes underneath the glacier, and (vii) a well-developed  
186 karst system which drains melt water to karstic springs at lower elevations.”

187

188 In this aim, Chapter 2.2 Geologic setting should be re-named Geology and hydrogeology  
189 settings.

190 *We will edit the title as suggested.*

191

192 Chapter 3 “Methods and data” is quite surprising. What is named “Karst model” (Subchapter  
193 3.1), and defined as one of the tools used for studying the hydrology, is in fact a 3-D  
194 representation of the geology in which the flow paths inferred from tracing tests are indicated.  
195 This hydrogeological model (see fig. 3, which is not really easy to read) is an interpretation,  
196 synthesizing geological, hydrogeological and tracing data. It can’t be presented before tracing  
197 results. I think that it should be considered in Chapter 5 “Discussion” with sub-chapter 4.3  
198 “Karst model results” all included in sub-chapter 5.1 “Hydrogeological model”, and not in  
199 “Methods and data”, because it is an interpretation of all hydrogeological data.

200 *The two pathways presented in blue and coming from the injection points on the glacier are*  
201 *not a result of the tracer experiments, but rather a hypothesis for the possible pathways based*



202 on 3D geology and previous tracing tests. The karst model was used as a tool to identify the  
203 best locations on the glacier for tracer injections. In Figure 7 we present the modified  
204 pathways according to results of the tracer experiments on the glacier. Results of the tracer  
205 experiments reveal that the first guess of flow paths presented in Figure 3 are only partially  
206 correct. We will improve Figure 3 and 7, edit the ms and change the caption of Figure 3 as  
207 follows:

208 *“3-D view towards the East of the Urganian limestone basis (top of Valanginian Marls). Red*  
209 *and transparent planes are faults. Tracing experiments previously carried out are represented*  
210 *in green, including the expected underground flow paths. The Plaine Morte is visible in the*  
211 *middle (dark blue). Blue spheres are karstic springs. The blue surfaces with some*  
212 *transparency are karstic nappes. The two flow path (light blue) coming from potential*  
213 *injection points on the glacier are inferred from the 3D geology. This interpretation was used*  
214 *to design the strategy of the tracing experiment.”*

215

216 Sub-chapter 3.2 regards all climatic (not meteorological, which is related to weather forecast)  
217 and hydrological data.

218 *In our understanding, meteorological data are short term (seasonal) data of few months or a*  
219 *couple of seasons, while climate is described by long term observations of several decades.*

220 *Accordingly, we think meteorological data is more suitable.*

221

222 In Chapter 4, may tracing test data help in interpreting the observed differences in isotopic  
223 compositions of the springs, as a result of different contribution from ice melting?

224 *To some extent yes but only one tracer was found at the source of Loquesse (Duasyne), while*  
225 *no tracers were detected at the other sites used for isotopic monitoring. The isotopic*

226 *composition for the site at Loquesse at the time of the tracer breakthrough indicated that*

227 *karst water is dominated by glacier melt water during late summer. This is clearly also the*

228 *case for the other spring (Ertense), where glacier melt water dominates over the summer*  
229 *months. In contrast, the Tièche (small river) is likely to be dominated by surface runoff during*  
230 *the summer months. We will clarify this point by editing the ms as follows:*

231 *The temporal evolution of the isotopic composition in the two karst springs of Loquesse and*  
232 *Ertense as well as the river sampling site of Tièche are all dominated by snow melt water in*  
233 *spring and early summer. In contrast, once the snow has completely melted, the water in the*  
234 *two sources is dominated by glacial melt water and only the river gets dominated by surface*  
235 *run-off with isotopic compositions approaching those of the summer to autumn precipitation.*  
236 *For Loquesse the tracer breakthrough only about 17 to 24 hours after injection is also clearly*  
237 *transported by glacial melt water, as supported by the isotopic composition measured in*  
238 *August. As there was no rainfall directly before and during the tracer injection, both sources*  
239 *remain close to the isotopic compositions for the glacier and only the river sampling site of*  
240 *Tièche is dominated by the 18O-enriched August/September rain water.*

241 *In addition, the following sentence will be added to page 20 of the discussion:*

242 *Glacial melt water as the dominant transporting agent for the tracers to the karst springs is*  
243 *also clearly indicated by the isotopic compositions of the water sampled at Loquesse (and*  
244 *Ertense) in 2011.*

245

246 *Did you try to compare the results of the glacier melt modelling to the observed flows? I think*  
247 *that they could help in testing some of the assumptions of the models (hydrogeological*  
248 *functioning and glacier melting).*

249 *The modeled glacier runoff only refers to the surface of the actual glacier as catchment and*  
250 *not to the entire drainage basin. It would be beyond the scope of this paper to incorporate all*  
251 *the poorly understood processes (hydrogeology etc.) into a complete model at the basin scale*  
252 *which would allow direct comparison against runoff measurements. Our results on glacier*  
253 *melt runoff were, however, validated against direct measurements on the glacier such as*

254 *long-term ice volume change, and ablation measurements (Huss et al., submitted,*  
255 *Geographica Helvetica). This will be clarified in the revised ms. Furthermore, detailed*  
256 *hydrological models are being tested for the southern slopes of the Plaine Morte area within*  
257 *the PhD-thesis of M. Kauzlaric.*

258

259 In Chapter 6 “Conclusions” point 1 is interesting, because this is not really said in the text. In  
260 fact the interpretation of tracing test data (sub-chapter 4.2) should have explained that the  
261 tests show that the flows are typical of karst conduits, from the surface of the glacier to the  
262 springs.

263 *We will edit the manuscript and make sure that the discussion states clearly how we come to*  
264 *conclusion 1!*

265

266 Despite I am not a native English speaker, I think that the English writing should be revised,  
267 although the paper is understandable as it is.

268 *In accordance with reviewer #4 we will look for possibilities to let the manuscript be read by*  
269 *a native speaker.*

270 **Bruno, Rolf, is funding available for this?**

271

## 272 **4 Anonymous Referee #3**

273 Received and published: 19 April 2013

274 Summary and general recommendation: Finger et al. have studied a glacier-karst system in  
275 the Swiss Alps and have used different methods to estimate the possible impact of climate-  
276 change induced glacier retreat on karst groundwater resources. This is an interesting and  
277 relevant study. The paper is suitable for publication following moderate changes, mainly

278 concerning the use of terminology (sometimes incorrect) and missing information on some  
279 figures (that look good but are not really comprehensible).

280 Also some scientific aspects need to be improved, e.g. concerning the limitations and  
281 transferability of the results.

282

283 Specific and general comments:

284 line 6: The term “souterrain” is inappropriate. Do you mean underlying or underground?

285 Anyway, the word can be deleted without losing any information. In fact, the entire sentence  
286 is not logical. Yes, glacier retreat is an important issue for water resources. Yes, karst aquifers  
287 are relevant. However, it is not clear why glacier-karst systems require particular attention.

288 Either delete this statement or find a better justification why your test site / study are relevant.

289 In fact, most Swiss / alpine glaciers are not located on karst but on crystalline rock. Maybe  
290 you could indicate somewhere the proportion of glacier on karst and on other rock types?

291 *We deleted the term “souterrain” and replaced it by "underlying".*

292

293 24-25: I would also mention drinking water (small quantity, but very important) and irrigation  
294 (= irrigated agriculture). Not sure if snow production makes much sense here.

295 *We added drinking water and will edit the ms accordingly.*

296

297 2746: You cite 2 papers by Vivian Gremaud et al. who have studied the directly adjacent and  
298 very similar glacier-karst system of Tsanfleuron. You should refer again to these papers in  
299 your discussion or conclusions and compare your results with their results. Not all results are  
300 presented in the published 2 papers. There is more in Vivian’s PhD thesis, including an  
301 estimation of future water availability under conditions of glacier retreat. A third paper by  
302 Zeng, Gremaud et al. (2012) quantifies the efficiency of this glacier-karst system as a CO<sub>2</sub>  
303 sink under global warming.

304 Another reviewer has also observed that the literature review is incomplete and has agreed  
305 some relevant references. I agree, particularly concerning the pioneer work done by Chris  
306 Smart in the Canadian Rockies. However, I would not cite too much gray literature (old  
307 conference proceedings) but focus on papers in international journals and books – there is  
308 enough!

309 *As suggested we will integrate relevant peer-reviewed literature.*

310

311 2748, line 3: Completely snow-free: Very important observation! This means that there is NO  
312 accumulation, i.e. the glacier is not only retreating but disappearing. Say this! Similar  
313 situation at Tsanfleuron, reported by Gremaud et al.

314 *We will emphasize on our result that the glacier will disappear within the current century in*  
315 *the revised ms. This is also illustrated in Figure 10.*

316

317 Line 8: 1-1,5 %: Are you sure? I don't have better numbers, but it seems to me that many  
318 Swiss glaciers are much, much larger, so I would suppose a smaller number. Please check!

319 *The total ice volume in the Swiss Alps is currently (2013) between 50 and 65 km<sup>3</sup>. This*  
320 *number is confirmed by several recent studies (Farinotti et al., 2009; Linsbauer et al., 2012,*  
321 *Huss and Farinotti, 2012). With a volume of 0.8 km<sup>3</sup> Plaine Morte thus accounts for about*  
322 *1.5% of the total ice volume. A reference is given in the manuscript.*

323

324 2749, line 5: Urgonian = Schrattekalk (mention the name of this very famous limestone  
325 formation that hosts the two largest caves in the Alps)

326 *As suggested, we will include the names of the limestone formation.*

327

328 14: The term “land use” is inappropriate here, because much of the area is not used!

329 *We agree and will replace the term “land use” with “land types” or similar.*

330

331 2751, line 20-22: Correct in principle, but not as simple in this case, because folds and faults  
332 can create reservoir structures and phreatic conditions above the level of the springs.

333 *We will edit the manuscript pointing out that folds and faults can create reservoir structures*  
334 *and phreatic conditions above the level of the springs.*

335

336 2753, line 16: The injection quantities are crazily huge – Gremaud et al. used about 10-100  
337 times smaller quantities in the Tsanfleuron area.

338 *Indeed, this was one of our first findings: we expected most of the tracer to infiltrate into the*  
339 *karst and assumed that it would dilute a lot more. Accordingly, we drastically reduced tracer*  
340 *amounts during the 2<sup>nd</sup> and 3<sup>rd</sup> injection.*

341 *Please also note that the discharge rates of springs from the Tsanfleuron area are about 10*  
342 *times lower than those of the Plaine Morte area. The discharge rate of the Loquesse spring*  
343 *can reach up to 15 m<sup>3</sup>/s. The rate of the Siebenbrunnen spring can be at least 6.5 m<sup>3</sup>/s. The*  
344 *total amount of water flowing at the bottom of the glacier was unknown before the first*  
345 *tracing experiment. Quantities for the first test (three injections in 2011) were too high but*  
346 *this was adjusted in 2012 with smaller quantities. We will edit the ms to clarify this point.*

347

348 2757, line 23: The maximum concentration are enormous! Uranin concentrations are 20 times  
349 above the limit of visibility. At some place, you should mention that your injection quantities  
350 were extremely overdosed, otherwise you give a bad example for future tracer tests. I prefer  
351 µg/L over mg/m<sup>3</sup>, but that's a question of taste.

352 *We agree (see also reply above). Nevertheless the particular setting of the Plaine Morte*  
353 *requires tracer amounts that are also detectable at karst sources as well as in the surface*  
354 *runoff. We will edit the ms to clarify this issue.*

355

356 2758, line 7: “amount of tracer passing” => use the term (tracer) recovery (%)

357 *We will edit the ms as suggested.*

358

359 Discussion: Bette compare your findings with results from the literature, e.g. concerning flow  
360 velocities of subglacial, englacial and supraglacial melt waters. In fact, these important  
361 glaciological terms are not used in the entire paper. You should really read and cite more  
362 glacier (and karst) literature and use the relevant terms and concepts in your paper.

363 *We will incorporat relevant peer reviewed literature. If reviewer #3 has some specific*  
364 *suggestions we would be very pleased to consider them.*

365

366 Table 3: Amount of tracer => recovery (%)! See comment above.

367 *We will edit the ms as suggested.*

368

369 Table 4: Table heading makes no or little sense. Do you mean: Comparison of tracer  
370 recoveries and flow velocities obtained from the three tracer tests?

371 *We will edit the headings and change “Amount of tracer” to “recovery rate (%)”.*

372

373 Figure 1: A scale bar in figure b would be useful.

374 *We will add a scale bar in panel b.*

375

376 Figure 2: There should be a legend explaining the stratigraphy!

377 *We will add a description of the stratigraphy to Fig 2.*

378

379 Figure 3: The figure looks amazing, but in fact, it is incomprehensible: No vertical scale, no  
380 horizontal scale, no orientation, the relation between geology and topography is unclear: does

381 the figure shows surface geology (outcrops) or the internal geological structure or a bit of  
382 both? Inacceptable in the present form (although it looks good). Must be improved.

383 *We will revise Fig 3 and add a scale bar and orientation. We cannot show geology and*  
384 *topography because it's superimposed. However, we will rework it thoroughly making it*  
385 *easier to read.*

386

387 Figure 5 and 6: Such graphs are called (tracer) breakthrough curves! In general, you should  
388 use the correct terms from the glacier, karst and tracer literature.

389 *We will edit the ms as suggested.*

390

391 Figure 7: Similar problem as figure 3: What is the relation between the colorful parts of this  
392 figure and the non-colored part in the upper left corner?

393 *The objective Figure 3 is to demonstrate the possible flow pathways in the rocks. Standard*  
394 *hydrogeological maps would assume that flow paths follow the thalweg, ignoring the geologic*  
395 *underground. As stated above, we intend to illustrate in Figure 3 flow pathways based on the*  
396 *currently know geologic setting of the underground. In the revised ms we will revise the figure*  
397 *thoroughly.*

398

399 General comment: The limitations of your study should be addressed more clearly! Your  
400 results are (hopefully) true for your test site and for the neighboring test site studied by  
401 Gremaud et al.

402 *Tracer concentrations have been measured by our tracer experts Dr. Fischer and Dr. Wernli.*  
403 *The method proved to be very reliable in past studies. Of course they cannot be transmitted to*  
404 *a neighboring study site, but their interpretations and the system understanding can help to*  
405 *understand processes in other, similar settings.*

406



407 However, the transferability to other areas is very limited, even within the Alps: There are  
408 very few glacierised karst systems in the Alps, and very different general trends can be  
409 expected for non-glacierised areas and for areas including large glaciers (that will not  
410 disappear so quickly).

411 *Indeed, the results cannot be directly transferred to other study sites, as nature is very*  
412 *heterogeneous and each study site has its own characteristics. Nevertheless, the*  
413 *interdisciplinary approach and the process understanding can be transferred to any other*  
414 *case study. We will clarify this issue in the revised ms.*

415

## 416 **5 Anonymous Referee #4**

417

418 Received and published: 30 April 2013

419 “Identification of glacial melt water runoff in a karstic environment and its implication for  
420 present and future water availability” by D. Finger et al.

421 The problem of glacier-karst hydrogeology has largely been piecemeal studies, largely by the  
422 empirical investigation of field sites. Despite use of quantitative measurement and analytical  
423 tools, our understanding remains essentially qualitative. This paper therefore has an ambitious  
424 objective in attempting to further our understanding at a generalisable quantitative level, by  
425 incorporating field observations and predictive models for underground flow routing and  
426 future mass balance scenarios. Unfortunately, these ambitious aims are only met superficially.  
427 The models used are poorly developed and are not integrated into the overall study. The field  
428 work components also appear to have little overall integration. The study therefore is a useful,  
429 but essentially parochial (local) investigation of glacierkarst hydrogeology.

430 *We agree with the reviewer and are aware that our objective to assess current and future*  
431 *water availability in the Plaine Morte region is very ambitious. But shouldn't every scientific*

432 *project have the ambition to explore innovative techniques in order to establish new theories?*  
433 *We are convinced that our study presents valuable new insights into the understanding of*  
434 *processes influencing water availability in the region. While this is a valuable asset for local*  
435 *residents depending on the water resources, the presented holistic approach presents a new*  
436 *innovative approach in karst hydrology.*  
437 *As stated above, certainly our results are only valid for our study site. Actually, this applies to*  
438 *all case studies. Nevertheless, our innovative approach of combining different investigation*  
439 *techniques is applicable to any study site in the world.*

440

441 The Glacier de la Plaine Morte is perhaps the definitive example of a glacier-karst system as  
442 the glacier largely occupies a massive closed depression. However, it is far from typical and  
443 the ice is largely stagnant. (This probably means it hosts a more stable internal drainage  
444 system than more dynamic glaciers.) I suggest that extrapolating from this unusual site (and  
445 the limited data set) to global scale is over ambitious. The generalisation that loss of alpine  
446 glacier ice will reduce late summer flow is widely recognised, and poorly supported by the  
447 analyses presented here.

448 *We agree and are aware that our results are only valid for our study site. However, as stated*  
449 *above, our approach can be applied to any case study in the world. We will clarify this issue*  
450 *in the revised ms.*

451

452 The “karst model” utilised in designing the tracer experiments appears to be an excellent tool,  
453 particularly valuable in the complex tectonics of the Swiss Alps. It does appear to have some  
454 limitations, however. First, the relationship between lithology, structure and hydrogeology has  
455 to be explicit. (Here it is not apparent which units and discontinuities (faults) are considered  
456 susceptible to karst permeability.) Similarly, there is no obvious inclusion of glacier ice (or  
457 talus), so the predictions are tenuous (and indeed appear to have been misleading in missing

458 recharge from the major outlet stream). Finally, the tool's effectiveness relies on parallax-  
459 based three dimensional rendering and fails when printed on a page. It would have been much  
460 more effective to provide a clear map showing the inferred flow routes. Overall, the flow  
461 visualisation model seems a bit disappointing. The "predictions" (in text as figure 3 is  
462 unreadable) seemed to indicate various underground drainage routes. In contrast, the results  
463 seemed to show (again figure 4 is not readable) that the subglacial karst has surprisingly  
464 limited and conditional permeability (which we already know) and the tracer delivery more or  
465 less travels to the nearest spring. The dominant results of surface routing and subsequent  
466 capture on the Bernese side were not explicitly predicted by the model (as far as I can see.  
467 Although the likely capture of the surface stream to springs is fairly evident using Google  
468 Earth; to which I would add the possibility of a talus aquifer linking the Reitzliberg and  
469 Siebenbrunnen Springs). The routing model might be more effective if were used to make  
470 specific, well-illustrated and tested hypotheses.

471 *The karst model is continuously being developed by the Swiss Institute for Speleology and*  
472 *Karst Studies (ISSKA). The presented application represents a first application which will be*  
473 *improved in the future. As stated above, we will improve the two figures and the ms will be*  
474 *edited to clarify the raised concerns.*

475

476 The hydroclimatic data are quite substantial, but their discussion does not seem to relate to the  
477 main purpose of the paper. They are only used explicitly in attempting to characterise glacier  
478 melt, though presumably used in various ways in analysis and modelling.

479 *We performed tracer injections during three periods of distinct melting: i) just after the snow*  
480 *on the glacier disappeared, ii) during intense melting and iii) after the first snowfall in*  
481 *autumn. The objective of these three injections was to see tracer evolution during the three*  
482 *distinct meteorological conditions. Accordingly, the date relates directly to the results. We*  
483 *will emphasize on this topic in the revised ms.*

484

485 The artificial tracing work is interesting, must have taken considerable work and is worth  
486 publishing. However, I would recommend some revision to reach a reasonable professional  
487 standard. The description and analysis lack the substance I would expect in a work of this  
488 scale. The tracers and traces are not well described. I could not find data on the Duasyne, but  
489 as one of the “optical brightener” tracers, it will show quite different optical and tracer  
490 performance to the other two closely related compounds (eosine and uranine). Unfortunately,  
491 the use of a synchronous scanning strategy implied in the terse section on analysis is likely to  
492 fail quite badly on typical optical brightener tracers. “Blue” (and to some extent “green”)   
493 tracers are also expected to have a high natural background which is not apparent from the  
494 figures.

495 *The tracer analysis was carried out in our in house laboratory by our tracer experts Dr.*  
496 *Wernli and Dr. Fischer. We will revise the ms adding descriptions on all tracers used and*  
497 *revising the description of the laboratory procedures.*

498

499 The masses of tracer injected are exceptionally large and could result in prosecution in some  
500 jurisdictions.

501 *As stated in the response to reviewer #3 there are two reasons for the high amount of tracer*  
502 *injection: i) tracer has to be also detected in karstic sources, where dilution is about 100*  
503 *times higher than in the surface runoff and ii) during the first injection we did not expect the*  
504 *tracer to drain through the surface runoff. Tracer amounts were drastically reduced during*  
505 *2<sup>nd</sup> and 3<sup>rd</sup> injections. But we still wanted to be able to detect tracer concentration in karst*  
506 *sources, where concentration is expected to be more than 100 times lower.*

507

508 The injection descriptions imply random release on the ice surface. Tracer injection usually  
509 requires prior dissolution (typically to ppt level) and instantaneous injection without

510 perturbation of steady flow. This is virtually impossible with the masses used, even if a  
511 suitably large supraglacial stream were present. (I could see none larger than perhaps 1m<sup>3</sup>/s in  
512 Google Earth.) Did the stream then sink into a moulin, or run over the surface? I was also  
513 confused by the descriptions of the 2012 injection sites; were they close to or near the 2012 I2  
514 site?

515 *Tracer mass was diluted on site with melt water from the glacier. Runoff of the supraglacial*  
516 *meltwater streams was about 0.2-0.5 m<sup>3</sup>/s during the injection. The injection was performed*  
517 *in safe distance of 10-20m from moulins that drained the water quickly to a depth of >50m*  
518 *within the glacier ice.*

519 *Swiss national television reported live from the injection (see link below).*

520 *Link of TV report:*

521 <http://www.videoportal.sf.tv/video?id=8b441b32-ce13-4496-975b-1a204dc20bd7>

522 *We will edit the manuscript to clarify this issue.*

523

524 Why was the glacier outflow not monitored? It is surely the most critical monitoring point!

525 *The immediate outflow at the glacier terminus is not well accessible and does not allow the*  
526 *installation of sampling devices. However, Trübbach is the site which represents the glacier*  
527 *outflow well. Unless there is heavy precipitation almost all the water in Trübbach comes*  
528 *directly from the glacier. We will edit the ms to make this clearer.*

529

530 The tracer analysis remains fairly rudimentary (which is appropriate in a novel setting), but  
531 some clarification is required. To compensate for contrasts in fluorescence intensity, the tracer  
532 concentration can be readily normalised to concentration per mass (typically 100g for some  
533 reason).

534 *By giving the actual concentrations, we can discuss the efficiency of the karst system. This is*  
535 *an important asset in order to justify our conclusions about water availability. We do not see*

536 *an added value in normalizing the concentrations and how this could contribute to our*  
537 *objective.*

538

539 Tracer velocities require a characteristic travel time (first arrival, mean or peak?) and path  
540 length (straight line, sinuous or true path?). It is not clear what is being used here. Tracer  
541 recoveries are useful in general, but not meaningful when the breakthrough curve is poorly  
542 defined as it is for the main river traces in 2011.

543 *We agree with the reviewer and will clarify how travel time was computed. Nevertheless, this*  
544 *will not have any impact on our main conclusions on present and future water availability.*

545

546 The conditions of melt and routing through the glacier are fundamental to the design of the  
547 tracer tests, the analysis of results and the future modelling. It is therefore surprising that the  
548 glacier hydrology in the paper is fairly superficial. It is implied that peak melt develops with  
549 full ice cover. This is not normally the case, peak local runoff occurs in the late stage of  
550 snowpack loss. It is not clear what route the melt water was taking. Was it supraglacial,  
551 marginal or through a perennial or seasonal conduit system?

552 *We do not fully agree with the reviewer here: For catchments without glaciers the peak runoff*  
553 *occurs at the late stages of the snowpack loss. Due to the albedo contrast of snow and ice,*  
554 *melting is however much stronger after the disappearance of the snow cover. This effect is*  
555 *directly captured by the glacio-hydrological model applied which has been tested and*  
556 *validated in numerous studies on alpine glaciers.*

557 *Furthermore, the objective of our study is to investigate current and future water availability*  
558 *in the Plaine Morte area – not to describe specific processes of glacier hydrology.*

559 *Nevertheless, glacier hydrology is one important aspect of our investigation. Accordingly we*  
560 *treated it only to an extent that is necessary for our objective. We will edit the manuscript to*  
561 *give more emphasis on glacier hydrology.*

562

563 What was the likely state of this system at the time of tracing? How is the routing likely to  
564 change through the year and how will this influence recharge?

565 *In order to answer these questions we performed three injections during different hydro-*  
566 *meteorological conditions (this is why we need these observations, as stated in reply to the*  
567 *concerns raised above). We will emphasis on this issue in the revised manuscript.*

568

569 Is the absence of recharge really due to “silt” and “loess” at the bed, or is the melt water  
570 routed away from recharge windows? I have never seen basal silts beneath a glacier and loess  
571 (a wind-blown silt) seems improbable. Most deglaciated karst surfaces exhibit extensive  
572 solutional permeability and sediments are generally coarse (permable) and dominantly very  
573 late or post glacial in timing. So either some substantive evidence has to be advanced, or the  
574 absence of recharge maybe attributed to routing of melt through the ice. The concluding  
575 diagram (Figure 11) proposes a dual porosity model for the glacier. This might seem a  
576 reasonable proposition for testing, but seems poorly supported by the work.

577 *The results of our study clearly indicate that during the low flow season water penetrated*  
578 *primarily into the karst while during the melt season melt water is evacuated through the*  
579 *glacier to the north. This is a fact we could demonstrate with our tracer experiments and*  
580 *which is relevant for our objective to determine current and future water resources in the*  
581 *area. A detailed description of the recharge of the karst is certainly interesting, but is not the*  
582 *primary objective of our study. Nevertheless, our understanding of recharge to the karst is*  
583 *sufficient to draw conclusions regarding current and future water availability. We will clarify*  
584 *this issue in the revised ms.*

585

586 A potentiometric surface is often postulated in glacier hydrology despite lack of empirical  
587 evidence. Such a model is very difficult to sustain for impermeable glacier ice. Instead, a

588 seasonally and spatially variable sheet-cavity-conduit is more widely supported. It is most  
589 unlikely that a coherent water table exists in a glacier such as Plaine Morte. Some direct or  
590 indirect observations are needed to construct an appropriate glacier hydrology in a setting like  
591 this. In the light of the results provided, the only substantial evidence appear to be the high  
592 season tracer results (five traces in all) reported cessation of surface outflow (no data are  
593 given). No tracing has been undertaken under non-overflow conditions, so the Bernese  
594 subglacial routing remains hypothetical. The lesson learned from the substantial body of work  
595 on glacier tracing is that results vary dramatically with injection conditions, location, season  
596 and runoff (diurnal and storm driven). Glacier karst tracing is expected to be even more  
597 idiosyncratic because of the likely variation in recharge opportunity at the bed. So the only  
598 relevant tracer test of the key focus of this research is those injections (three in all) observed  
599 at Loquesse spring.

600 *The five tracer injections had two main purposes: i) the three simultaneous injections at three*  
601 *different locations on the glacier had the objective to determine flow paths from three*  
602 *representative parts of the glacier and ii) the three injection at location I2 during three*  
603 *different seasons (just after snow disappearance, during melt season and after the first snow*  
604 *in fall) had the objective to assess flow paths during different seasons.*

605

606 The stable isotopic data are too sparse to provide much insight. Two sites have not been  
607 proven connected to glacier. The key heavy signal at Tieche is presumably a transient rain  
608 event captured at that site. I would omit this section.

609 *Indeed, the stable isotope data is sparse but they nonetheless give a valuable insight to the*  
610 *connections between the glacier and the karst sources in the southern part of the Plaine*  
611 *Morte area. The August and September measurements for both karst sources are clearly*  
612 *compatible with melt water dominating in these sources at this time as there was no*  
613 *precipitation just prior to the injection of the tracer that was detected in Loquesse in August*



614 2011. In addition, all snow-melt was already drained. In contrast, the Tièche river is surface-  
615 water and hence precipitation dominated during the late summer and autumn, but this is NOT  
616 the case for the southern karst sources. As in this region no artificial tracer was observed  
617 after tracer injection in the Ertèense source, it must be assumed that snow and glacier melt  
618 from other regions of the glacier than the injection points must infiltrate into the karst and  
619 drain to the karst sources. The isotopic observations confirm this; accordingly, they present a  
620 valuable asset to the objective of assessing water availability in the region. Both in chapter  
621 4.4 and in the discussion, the text was rephrased in order to bring these points out more  
622 clearly (see above).

623

624 A glacier melt model is used to predict the future geometry of the glacier. It is not clear how a  
625 daily model is calibrated using two digital elevation models fifty years apart and validated  
626 using three years of limited accumulation and ablation data. Although it provides a crude  
627 linear trend, extrapolating this trend into the untested geometry of a closed basin seems very  
628 risky, especially when rather dire implications are drawn. It is not clear that the hydrology has  
629 been fully implemented. If the leaky closed depression model (figure 11) is used, then it  
630 seems that an increasingly large fraction of the melt would be retained in the depression and  
631 subsequently recharged. The data in this paper would seem to suggest that relatively little  
632 melt would travel north (Bern) and a greater fraction would be routed south (Valais).

633 *We are aware that model projections are highly uncertain. Besides the limited calibration*  
634 *data available this is also due to the uncertainties in future climate projections, and model*  
635 *assumptions as mentioned in the HESSD paper. In the revised ms we will clarify that the*  
636 *modeling approach used does not include the karst hydrology, but only provides the total*  
637 *quantities of glacier melt input into the system. Accordingly, the model results themselves do*  
638 *not allow an assessment of future water flow paths.*

639 *Calibration of a glacio-hydrological model in karstic environment is very difficult. Observed*  
640 *snow accumulation and ice melt data allow constraining the most important components of*  
641 *the water balance at time scales comparable to the projection period. We are aware of the*  
642 *uncertainties and clearly mention them in the paper. However, we are confident in the*  
643 *general results provided by this model that has proven to be well suited in modeling alpine*  
644 *runoff regime changes (see e.g. Farinotti et al., 2012).*

645

646 Overall, there are substantial weaknesses in all the components of the paper and in their  
647 integration. It is not clear that the conclusions are based on substantive analysis. This is  
648 unfortunate as the various components of the research are interesting and challenging, but  
649 lack adequate treatment in attempting to compose an integrated report. The results provided  
650 suggest that a test of underground flow predictions using dye tracing would be worth  
651 reporting. Similarly, the forward modelling looks promising. It is not clear that the results  
652 allow much advance in our understanding of glacier karst recharge. Therefore, the primary  
653 purpose of the paper is not adequately addressed.

654 *We can convincingly demonstrate current water flow paths: Our results show that i) glacier*  
655 *melt water from the eastern and northern injection points is evacuated rapidly through the*  
656 *glacier to the northern surface runoff (no tracer was found in the southern karst sources) ii)*  
657 *during melt season glacier water from the southern injection point is drained to approx. equal*  
658 *masses through the glacier to the northern runoff and to the Loquesse source. iii) at the*  
659 *beginning of the melt season glacier water from injection point I2 is drained also to the*  
660 *northern surface runoff as well as to the Loquesse source, however, travel velocities are*  
661 *almost halved to due lower melt production iv) at the start of the winter season (after the first*  
662 *snow) glacier water from injection point I2 is drained only to the Loquesse source, as no*  
663 *tracer was observed on the northern side. v) Although numerous small karst sources did not*  
664 *reveal any tracer concentrations, isotopic signatures indicate that water composition is*

665 *dominated by glacial water during melt season. The findings are based on the five tracer*  
666 *injections, hydro-meteorological data, isotopic investigations and satellite snow cover data.*  
667 *Based on these insights and climate change projections of the glacier melting, we make*  
668 *assumptions about future water availability. We are convinced that after revisions of the ms*  
669 *our conclusions will be fortified. We will edit the conclusions in order to clearly differentiate*  
670 *between facts and interpretation of facts.*

671

672 A final comment on the language: the English is good, but in places the technical usage is  
673 incorrect and possibly misleading. A editorial proof reading is advisable.

674 *We will revise the English and incorporate technical terms as suggested also by the other*  
675 *reviewers. An editorial proof reading would be welcomed.*

676

677 *(1) Side note to the author contribution:*

678 *Finger: writing of the ms, planning, coordination, supervision of project and first author*

679 *Hugentobler: wrote a MSc thesis about the tracer experiments*

680 *Huss: glacier modeling and contribution to text editing*

681 *Voinesco: wrote a MSc thesis about ice thickness and mass balance of Plaine Morte*

682 *Fischer: lab analysis of tracer concentrations*

683 *Weber: karst modeling and contribution to text editing*

684 *Jeannin: karst modeling and contribution to text editing*

685 *Kauzlaric: hydro-meteorological data and sampling after two injections in the south*

686 *Wirz: wrote a MSc thesis about the isotope investigations*

687 *Vennemann: isotope investigations and contribution to text editing*

688 *Hüsler: processing of MODIS satellite data*

689 *Schädler and Weingartner: project initiators, coordinators and PI of the MontanAqua*  
690 *project*