

Interactive comment on “Just two moments! A cautionary note against use of high-order moments in multifractal models in hydrology” by F. Lombardo et al.

D. Schertzer

daniel.schertzer@enpc.fr

Received and published: 10 July 2013

C3103

Multifractality: at least three moments !

D. Schertzer, I. Tchiguirinskaia, S. Lovejoy

10 July 2013

1 Summary of our comment

Multifractality of rainfall has been the central topic of theoretical hydro-meteorology for almost three decades and it is therefore important to improve the parameter estimation techniques of multifractal fields. Unfortunately, we believe that the present paper (Lombardo et al., 2013) brings more confusion than clarification on this issue. First, the conclusions of this paper, as well as part of its title, are misleading. The authors indeed claim that only the first and second statistical moments of a multifractal field are safely estimated. Unfortunately, these two moments are insufficient to determine the nonlinear scaling moment function $K(q)$ ¹ of a multifractal field R , e.g. the rain rate, observed at various resolutions² $\lambda = L/l$ (where L is the largest scale, l is the observation scale) and where $\langle . \rangle$ denotes a given ensemble average:

$$\langle R_\lambda \rangle \approx \lambda^{K(q)} \quad (1)$$

¹which has been often denoted $-\xi(q)$ for structure functions.

²these observations could be obtained by various means, e.g.: aggregation, fluctuation estimates.

Indeed, contrary to the linear case of a scaling moment function $K(q)$ of a uni/mono-fractal field, a third independent value is at least required to estimate the curvature of $K(q)$ for multifractal fields. Therefore, their conclusions would mean that the multifractal parameters could not be safely assessed and multifractals would be therefore of little interest to simulate rainfall.

Secondly, the present paper fully ignores the concept of second order multifractal phase transition, introduced years ago (Schertzer and Lovejoy, 1992; Schertzer et al., 1993; Schertzer and Lovejoy, 1994) (for recent reviews: Schertzer et al. (2010); Schertzer and Lovejoy (2011); Lovejoy and Schertzer (2013)). This phase transition explains not only in a straightforward manner the qualitative observations made by the authors that the estimates of the statistical moments of order $q \geq 3$ of their numerically simulated cascades seem to be spurious, but provides rigorous, analytical results. Indeed, this phase transition occurs at a critical moment order q_s that is analytically defined from $K(q)$ and such that the estimates over a sample of all the statistical moments of order $q > q_s$ are spurious. We show below that the discrete cascade model (Lombardo et al., 2012) used by the authors yields a theoretical critical order $q_s(k)$ that depends on the k^{th} level of the cascade. Its graph is displayed in Fig.?? for the parameter set chosen by the authors, but with a much larger number of steps to show its slow convergence to $q_s(\infty) \approx 2.582$. Because this asymptotic value is also an upper bound, the estimates of the moments of order $q > q_s(\infty)$ are all spurious. The simulations and qualitative observations of the present paper can be therefore seen as illustrations of this phase transition rather than new findings.

2 Multifractals and phase transitions

To go beyond the straightforward counter-argument that we can learn a lot on the curvature of $K(q)$ from statistical moments of non integer orders q 's, let us recall that a

C3105

convenient and physically meaningful choice of three independent multifractal parameters correspond to:

- the (Hurst) scaling exponent $H = -K(1)$ of the mean field. The value $H = 0$ corresponds to a 'conservative field", i.e. a field whose mean is strictly scale invariant, whereas $H \neq 0$ rather corresponds to a fractional integration/derivation of a conservative field;
- the mean intermittency measured by the codimension $C_1 = dK(q)/dq|_{q=1} + H$ of the support of the mean field; its lower bound $C_1 = 0$ corresponds to a homogeneous field;
- the index of multifractality $\alpha = d^2K(q)/dq^2|_{q=1}/C_1$ characterises both the curvature of $K(q)$ at $q = 1$ and how the intermittency relatively varies when departing from the mean field. Its lower bound $\alpha = 0$ corresponds to a uni/mono-fractal field, whereas its upper bound $\alpha = 2$ to the misnamed lognormal model.

These three characteristics are generally only local, i.e. characterise the statistics of a multifractal field only in the neighbourhood of its mean. They are indeed based on $K(q)$ and its two first derivatives at the point $q = 1$ and are therefore defined by moments of (non integer) orders $q \approx 1$. However, these characteristics became global, i.e. for all orders q 's, for the 'universal multifractals' (Schertzer and Lovejoy, 1987, 1997). This 3-parameter family of multifractals is not only large, but also stable and attractive; it corresponds to the limit set of a broad generalisation of the central limit theorem in a multiplicative framework. However, this global characterisation needs to take into account the generic phenomena of multifractal phase transitions, i.e. discontinuities of the effective scaling moment function $K^*(q)$ (estimated over a given number of samples), which is the analogue of a thermodynamic potential, at given critical order q^* , the analogue of the inverse of a critical temperature (θ denotes the Heaviside function):

C3106

$$K^*(q) = (1 - \theta(q - q^*))K(q) + \theta(q - q^*)(K(q^*) + \gamma^*(q - q^*)) K(q) = \frac{C_1}{\alpha - 1}(q^\alpha - q) \quad (2)$$

For a second order transition $\gamma^* = dK(q)/dq|_{q=q^*}$, whereas for a first order phase transition $\gamma^* > dK(q)/dq|_{q=q^*}$. In both cases, γ^* increases with the log of the number of samples, linearly for the latter case. For an infinite number of samples, the first order transition corresponds to a divergence of the theoretical moments. The authors mentioned this possibility, but ignored the second order phase transition that explains in a straightforward manner what they numerically observed. Indeed, recalling that in agreement with Eq.1 a multifractal field R has an infinite hierarchy of singularities γ 's:

$$R_\lambda \approx \lambda^\gamma \quad (3)$$

The physics of the second order multifractal transition corresponds to the fact that for any finite sample there is a supremum γ_s of the singularities γ 's present in this sample. This supremum being an isolated point, its support set has a zero dimension $D(\gamma_s)$, equivalently a codimension $c(\gamma_s) \equiv d - D(\gamma_s) = d$, where d is the dimension of the embedding space ($d = 1$ for a time series). Due to the fact that the codimension $c(\gamma)$ is the Legendre transform of the scaling moment function $K(q)$, and vice-versa, in analogy to the fact that the entropy is the Legendre transform of the thermodynamic potential, one obtains that the critical order q_s corresponding to γ_s is given by:

$$q_s = (d/C_1)^{1/\alpha} \quad (4)$$

To apply these ideas to the present model the only difficulty is to go through the cumbersome algebra of its scale dependent parameters, which implies a rather unusual scale dependence of the scaling moment function $K(q, k)$, see Fig.???. This complexity algebra is first due to the fact that the generator of this discrete cascade is obtained with the help of a discrete fractional gaussian noise that is finally transformed into a log

C3107

divergent generator with the resolution, as required to obtain a scaling field. To avoid confusion between the fractional integration involved in their cascade generator and a possible fractional integration over the cascade itself, as mentioned above, we denote their corresponding H parameter by h . The differences are further discussed in a companion comment ("Further (mono fractal) limitations of Climactograms", by Lovejoy et al.). In any case, it would have been simpler to directly define a log divergent generator, as usually done. A second complexity is introduced by an attempt to recover a translation invariance in a discrete cascade model. Let us briefly mention, without further discussion in this comment, that the interest and relevance of such an attempt can be brought into question by the fact that discrete cascades are intrinsically built within an ultrametric framework, not a metric one. Furthermore, there already exist well defined continuous (in scale) cascades that are not only translation invariant, but also respect causality, which remains beyond the discrete cascade framework (Marsan et al., 1996; Schertzer et al., 1997). In any case, the presented model has already the drawback that its scaling moment function is not scale invariant, but depends on the level k of the cascade. To simplify the discussion, let us focus on the lognormal case, i.e. $\alpha = 2$ in Eq. 2 with a codimension of the mean intermittency $C_1(k)$ depending on the k^{th} level of the cascade:

$$C_1(k) = Ln[2^{2k(1-h)}(\sigma_0/\mu_0)^2 + 1] \quad C_1(k) \approx 1 - h + 2Log[\sigma_0/\mu_0]/(2kLn2) \quad (5)$$

One may note that the mean intermittency (as measured by $C_1(k)$) increases with scale and therefore brings into question the physical relevance of such a model. Without discussing the (limited) significance of the involved parameters, we use the set of parameter values chosen by the authors ($\mu_0 = 1, \sigma_0 = 1.29, h = .85$) and thus obtain the figures Fig.?? and Fig.?? respectively for $q_s(k)$ and $C_1(k)$, as well as the asymptotic values $C_1(\infty) = .15$ and $q_s(\infty) \approx 2.582$, which are relevant for a "dressed" cascade, i.e. a cascade first downscaled to infinitely small scales, then upscaled back to a finite observation scale.

C3108

References

- Lombardo, F., Volpi, E., and Koutsoyiannis, D. (2012). Rainfall downscaling in time: theoretical and empirical comparison between multifractal and Hurst-Kolmogorov discrete random cascades. *Hydrolog. Sci. J.*, 57:1052–1066.
- Lombardo, F., Volpi, E., Koutsoyiannis, D., and Papalexiou, S. (2013). Just two moments! a cautionary note against use of high-order moments in multifractal models in hydrology. *Hydrol. Earth Syst. Sci. Discuss.*, 10:4627–4654.
- Lovejoy, S. and Schertzer, D. (2013). *The Weather and Climate: Emergent Laws and Multifractal Cascades*. Cambridge University Press.
- Marsan, D., Schertzer, D., and Lovejoy, S. (1996). Causal space-time multifractal processes: predictability and forecasting of rain fields. *J. Geophys. Res.*, 101(D21):26,333–26,346.
- Schertzer, D. and Lovejoy, S. (1987). Physical modeling and analysis of rain and clouds by anisotropic scaling multiplicative processes. *Journal of Geophysical Research D*, 8(8):9693–9714.
- Schertzer, D. and Lovejoy, S. (1992). Hard and soft multifractal processes. *Physica A*, 185:187–194.
- Schertzer, D. and Lovejoy, S. (1994). Multifractal Generation of Self-Organized Criticality *Fractals in the Natural and Applied Sciences* Novak, M. M. Ed., North-Holland, pages 325–340.
- Schertzer, D. and Lovejoy, S. (1997). Universal multifractals do exist! *J. Appl. Meteor.*, 36:1296–1303.
- Schertzer, D. and Lovejoy, S. (2011). Multifractals, generalized scale invariance and complexity in geophysics. *Int. J. Bifurcation Chaos*, 21(12):3417–3456.
- Schertzer, D., Lovejoy, S., and Lavallée, D. (1993). Generic multifractal phase transitions and self-organized criticality. *Cellular Automata: Prospects in Astronomy and Astrophysics*, Perdang, J. M. and et al. Eds, World Scientific, pages 216–227.
- Schertzer, D., Lovejoy, S., Schmitt, F., Tchiguirinskaia, I., and Marsan, D. (1997). Multifractal cascade dynamics and turbulent intermittency. *Fractals*, 5(3):427–471.
- Schertzer, D., Tchiguirinskaia, I., Lovejoy, S., and Hubert, P. (2010). No monsters, no miracles: in nonlinear sciences hydrology is not an outlier! *Hydrological Sciences J.*, 55(6):965–979.