

***Interactive comment on* “Post processing rainfall forecasts from numerical weather prediction models for short term streamflow forecasting” by D. E. Robertson et al.**

Anonymous Referee #3

Received and published: 24 July 2013

Post-processing methods are most important for removing bias and to improve the reliability of rainfall forecasts. In this paper the Bayesian joint probability (BJP) modelling approach is applied to generate forecast probability distributions and ensembles are produced using the Shaake shuffle. Different verification methods have been applied to the post-processed NWP rainfall predictions and the results are discussed. In general the paper is very well written and its worth to be published after some minor changes.

There are just a few comments:

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

1. The BJP modelling approach has been explained already in detail in several other papers by the authors (e.g. Wang, 2009), so this part could be short and removed in the Appendix, since there is no new information. What is really important is a thoroughly description how you keep the temporal and spatial correlations applying the Shaake shuffle, because these correlation structures are essential and are most often destroyed by the application of post-processing methods. Unfortunately it is not discussed in detail and from the results shown, it is difficult to see how the temporal correlation is reproduced. Furthermore it will be very interesting to see how the spatial correlation is reproduced, when the method is applied to more stations.

2. Regarding the forecast verification: If you have forecasts with lead-time up to 10 days available, why do you show only reliability diagrams for day 1 and day 2? At least you should mention the results for the other lead-times! Furthermore the sample size of the rainfall events exceeding the 5 mm threshold seems to be far too small to make some meaningful interpretations. In Fig. 10 you say on the one hand that for day 1 the forecast probability of a rainfall event of greater than 5 mm appears to be reliable. On the other hand for day 2 you say that the forecast is unreliable, because there are just a few forecasts falling into the two upper bins. When I look at the inserted histograms, I cannot see forecasts falling into the third bin at day 1 also.

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., 10, 6765, 2013.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper