

Development of a new classification for Cochin Estuary, West Coast of India

Shivaprasad A*, Vinita J., Revichandran C., Manoj N.T., Jayalakshmy K.V., Muraleedharan K.R.

National Institute of Oceanography Regional Centre, Dr. Salim Ali road, Kochi-18, Kerala, India

**Corresponding author: shivaprasadnio@gmail.com*

Abstract

Cochin estuary is a unique complex system along Indian coastline with a widespread area at the upstream. The fluctuations in salinity are of extreme kind ranging from entirely riverine to entirely saline. The high runoff months are characterized by monsoonal spells causing intense flushing. During the peak dry period, the runoff is less but steady providing a stable environment. The existing methods prove to be insufficient to represent the real salient features of this typical estuary. Arguments are also presented to illustrate the confusion in the names by which the estuary is commonly known. In this context, a new nomenclature is proposed as ‘Cochin Monsoonal Estuarine Bay’ embodying the physiographic, hydrographic and biological features of the estuary. This is achieved by collating past evidences and by examining the present characteristics of the estuary using recently acquired large comprehensive data sets. Several estuarine classification schemes based on relatively easily measurable parameters and hydrological factors like river runoff are also evaluated for the estuary to determine how well the classification schemes represent the reality. The constraints imposed by these classification schemes evidences the uniqueness of the region.

Key words: Estuary, classification, runoff, India

1. Introduction

Estuaries are always dynamic and often exhibit a gradient in conditions from absolute riverine to oceanic which makes estuarine classification a complex matter. For a specific estuary, the classifications dealing with one type may change from one type to another in consecutive tidal cycles, or from month to month and from season to season or even from one

35 location to another within the estuary. Additionally, the system may undergo changes under
36 the influence of natural hazards or even anthropogenic influences. Valle-Levinson, 2009 had
37 documented that the most widely accepted definition of an estuary was proposed by Cameron
38 and Pritchard (1963). Accordingly, an estuary is defined as a semi-enclosed coastal body of
39 water which has a free connection with the open sea and within which sea water is
40 measurably diluted with fresh water from land drainage. The above definition of an estuary
41 applies to temperate (classical) estuaries but is not applicable for arid, tropical and subtropical
42 basins. Arid basins and those forced intermittently by freshwater exhibit hydrodynamics that
43 are consistent with those of classical estuaries and yet have little or no freshwater influence.

44 Under this general definition, estuaries may be further separated into various
45 classifications based on their stratification or vertical structure of salinity (Pritchard (1955),
46 Cameron and Pritchard (1963), and later Dyer (1973, 1997)), water balance ((Valle-Levinson,
47 2009), geomorphology ((Fairbridge, 1980), tidal characteristics (Hayes, 1975, Dyer, 1995)
48 and combination of characteristics (Savenije, 2005). Indian estuaries are influenced by
49 monsoon rainfall and hence, are referred to as monsoonal estuaries (Vijith et al., 2009). A
50 realistic classification, representative of the true characteristics of an estuary can be done only
51 after understanding the dominant dynamic processes of an estuary. This demands rigorous
52 investigation in to the dynamics of each section of the estuary using comprehensive data sets.

53
54 Cochin estuary, situated along west coast of India, attained its present configuration as
55 a result of natural and man-made interventions. It has irregular topography and interspersed
56 by numerous islets and shoals of varied sizes and shapes. It was primarily a marine
57 environment bounded by an alluvial bar parallel to the coast line and interrupted by Arabian
58 Sea at intervals (Gopalan *et al.*, 1983). For the establishment of Cochin Port in 1936, the
59 “natural bar” was dredged out while deepening the channel to make the basin accessible for
60 ocean going vessels (Strikwerda., 2004). There were several ways in which Cochin estuary
61 was named in earlier studies. The estuary was sometimes called as a “lagoon” (Rao and
62 Balasubramaniam., 1996); or very often referred to as “backwaters” (Sankaranarayanan and
63 Qasim., 1969, Martin et al., 2008, Abhilash et al., 2012). Lagoons are shallow body of water
64 at least intermittently connected with sea or other larger body of water across a beach or
65 barrier (Snead 1982). Cochin estuary is permanently open to sea and is much larger and
66 deeper than a typical lagoon. The Webster dictionary defines ‘backwaters’ as part of river
67 water backed up in its course by an obstruction, an opposing current, or the tide. Being an
68 extraordinarily energetic and dynamic environment typified by strong currents (1.3m/s)

69 (Udaya Varma et al., 1981, Balachandran et al., 2008), the nomenclature 'backwaters'
70 remains subtle to this estuary.

71

72 The motive of this work is the different existing nomenclatures used for Cochin
73 estuary in previous literatures. Therefore, we in this paper attempt to establish a new
74 terminology for Cochin estuary that is representative of its behaviour as a whole. For the
75 present study, the runoff data for the year 2008-2009 is used to examine the runoff dynamics
76 of Cochin estuary. In the next section, the credibility of this runoff data is well established
77 using detailed statistical analyses with past data sets. In section 3, we illustrate the annual
78 cycle of salinity in the estuary followed by an evaluation of classification schemes based on
79 measurable parameters (Hansen and Rattray, 1966) and hydrological factors like river runoff
80 (Vijith *et al.*, 2009). In section 4, we determine the salinity steadiness of the estuary using
81 salt-balance equations during peak dry period. In section 5, we review the previous studies in
82 order to examine the physical-biological coupling in the estuary which ultimately leads to the
83 proposal for a new nomenclature described in section 6. Section 7 summarizes our
84 conclusions.

85

86

87 **Physiographic setting**

88

89 Cochin estuary is the largest estuarine system along the west coast of India. It is a part
90 of Vembanad-Kol wetland system, one among the Ramsar sites in Kerala (November 2002),
91 which extends from Munambam (10°10'N, 76°15' E) in the north to Alappuzha (09°30'N,
92 76°28'E) in the south at over 96.5km in length (**Figure 1a**). The estuary is characterized by its
93 major axis lying parallel to the coastline, with several small islands and interconnected
94 waterways, and it covers a surface area of about 300km². The width of the estuary varies
95 from 450m to 4km and the depths range from 15m at Cochin inlet to 3m near the head with
96 an average depth of 1.5m (depths are reduced to chart datum). The system is separated from
97 the Arabian Sea by barrier spits interrupted by tidal inlets at two places, namely (i)
98 Munambam in the north (inlet 1) and (ii) Cochin inlet in the middle (inlet 2). The Cochin
99 Port, situated on the Willingdon Island, is near the inlet 2, which provides the main entrance
100 channel to this system. Tides in the estuary are mixed, predominantly semi-diurnal type with
101 an average tidal range of 1m (Qasim and Gopinathan., 1969). Freshwater into estuary is
102 primarily contributed by six rivers. The branch of Periyar River feeds 30% of its discharges

103 into the northern parts of the estuary. The remaining 70% discharges directly into the Arabian
104 Sea through the inlet 1. Muvattupuzha River joins along the length of the channel whereas
105 Pampa, Achankovil, Manimala, and Meenachil join at the upstream end. During the dry
106 season, the runoff originating upstream is minimal which ensures strong saline intrusion to
107 the low-lying paddy fields located further upstream (Shivaprasad *et al.*, 2012) (**Figure 1a**).
108 Therefore, a salt water barrage called Thanneermukkam Barrage (TB) was constructed in
109 1976 which is thereafter kept closed during the dry season to facilitate paddy cultivation. The
110 flushing time of Cochin estuary ranged from 1 day to 2.5 days during wet season and 8.7
111 days during dry season (Vinita *et al.*, 2013).

112

113 For the present study, the region was divided into two parts (**Figure 1a**): the northern
114 arm extends from Cochin to Munambam and the southern arm extends from Cochin to
115 Thanneermukkam. Both the arms of the estuary receive significant amount of freshwater
116 throughout the year; larger in southern arm than the northern arm. When the TB was closed,
117 Muvattupuzha River contributed to the freshening of the southern arm. The two arms behave
118 differently in physiographical and hydrographical aspects and hence treated separately.

119

120

121 **2. Materials and methods**

122

123 **Data sets**

124

125 The study utilized runoff hydrological data obtained from Central Water Commission,
126 Government of India and physical parameters like salinity, velocity and water level measured
127 during several field campaigns.

128 The daily runoff data of six rivers for six gauging stations for three periods: 1978 –
129 2001; 1985-1989 and 2008-2009 were employed for statistical approach. To investigate the
130 runoff dynamics of Cochin estuary, 1 year runoff data (2008-2009) was used. The long term
131 runoff data sets were used for the validation, assessment of sufficiency and completeness of
132 the 1 year data. This is the most detailed hydrology of this estuary published to date.

133 Annual variation in salinity was monitored from the longitudinal transect
134 measurements covering ten stations from June 2008 to May 2009 (**Figure 1a**). CTD (SBE
135 Seabird 19 plus) casts of salinity (conductivity $\pm 0.001 \text{ Sm}^{-1}$) profiles were taken from a small
136 boat for every 8 km in the deepest part of the main channel during the spring tides of each

137 month. Stations, 1-4 were located in northern arm and the stations 5-10 were located in the
138 southern arm. A daily monitoring station near to the inlet 2 (**Figure 1**) was suitably chosen
139 and the vertical profiles of salinity were collected every day at 11.00 AM local time during
140 the same year (May 2008 to April 2009).

141 During the year 2009-2010, time series measurements of salinity and velocity were
142 conducted at five stations under three runoff conditions. Stations A and B were along
143 northern arm, stations D and E were along southern arm and station C represented inlet 2
144 (**Figure 1a**). Sampling was conducted on spring phases of October 2009, February 2010 and
145 August 2010. These months were representative of moderate, dry and high runoff periods
146 respectively. Each observation started at 9.00AM and finished at 9:00 AM of the next day.
147 For every 24 hours observation, CTD was lowered at 30 minutes interval. Current meters
148 (RCM-9) were moored and velocity was measured at 10 minutes interval from near surface
149 and bottom. Water level data for the five stations in February 2010 was obtained from
150 permanent mooring stations of the field program. The estuarine volume was estimated from
151 digitization of recently developed bathymetry charts using 3D Analysis tools in ArcGIS
152 software.

153

154 **Statistical analyses on river runoff data**

155

156 Statistical analyses were done to substantiate the credibility of the runoff data for the
157 year 2008-2009 which is used for the present study. For this purpose, the data of average
158 monthly runoff for 1978-2001 and 1985-1989 was obtained by calculating the arithmetic
159 means of daily runoff data. Among the 23 years (1978 to 2001) data set, data from four rivers
160 were missing. However, for the period 1985-1989 the data from all the six rivers were
161 available. Utilizing these past sets of data, monthly total runoff for the year 2008-2009 was
162 predicted using the best polynomial fitted for the average monthly runoff of past data sets
163 among a set of different polynomials (**Figure 2a**).

164 To determine the main contributing components to the river runoff, a multiplicative
165 time series model was fitted. Time series analyses were carried out for the complete data set
166 (1985-1989). The multiplicative Holt winter model (Pillai and Bagavathy, 2009) was chosen
167 in which the observed monthly runoff is equal to product of long term trend (T), seasonal
168 variation (S), cyclical component (C) and irregular variation (I) in the runoff

169

170

171 i.e., $O = T * S * C * I$(1)

172

173 Trend, 'T' was identified by centered moving average (MA) of period 2. Centered MA of
 174 period 2 implied that river runoff at a time point 't' was determined by runoff at t-1, t and
 175 runoff at t+1 with weights 1,2 and 1 respectively. This triplet was the best preferred one, since
 176 the plots of other periods (3 to 12) explained the observed runoff very poorly. River runoff
 177 was observed to follow the moving average trend of period 2 very precisely (Figure 2).
 178 Seasonal variation, 'S' in each month was explained by the seasonal index computed as the
 179 simple average of (O/T) over all the years for each month. Cyclical variation was computed
 180 as a percentage of moving average as

$$181 \quad c = \left[\left(\frac{O}{SI} \right) - MA(2) \right] * 100 / MA(2) \dots \dots \dots (2)$$

182 where SI is the average variation adjusted to 12 as

$$183 \quad SI = \left[\text{Average monthly} \left(\frac{O}{T} \right) * 12 \right] / \text{Total of all average monthly} \left(\frac{O}{T} \right) \dots \dots \dots (3)$$

184 and MA (2) is the moving average of period 2. Cycles in the variation was clearly explained
 185 by the cyclical variation with a period of 12 months for repeated cycle (**Figure. 2b**). Irregular
 186 variation gets removed while averaging at different stages. Then these three time series
 187 components were used as independent variables to determine the regression of runoff on
 188 these components.

189 The river runoff (Y) was regressed on moving average of period 2 (X_1), seasonal
 190 variation (X_2) and cyclical variation (X_3) and their first order interaction effects. Step up
 191 multiple regression method was applied to determine the $2^3 * 6$ models (Snedecor and
 192 Cochran, 1967, Jayalakshmy, 1998).

193 Multiple regression model fitted is of the form

$$194 \quad Y = a_0 + \sum_{i=1}^{i=k} a_i X_i + \sum_{i=1}^{i=k} \sum_{j=1, i < j}^{j=k} b_{ij} X_i X_j \dots \dots \dots (4)$$

195 where $a_i, b_{ij}, i, j = 1, 2, 3, \dots$ and $i < j$ are the regression coefficients of the individual effects
 196 and the corresponding interaction effects respectively. To determine the contribution levels of
 197 the components uniquely, first order and second order partial correlation coefficients were
 198 calculated (Snedecor and Cochran, 1967). First order partial correlation coefficient is

199

$$200 \quad r_{ij.k} = \frac{r_{ij} - r_{ik}r_{jk}}{\sqrt{(1-r^2_{ik})(1-r^2_{jk})}} \quad i, j, k = 1, 2, 3, 4 \dots \dots \dots (5)$$

201 where 1 = river runoff

- 202 2= MA (2)
 203 3= Seasonal variation ‘S’
 204 4= Cyclical variation ‘C’

205

206 Second order partial correlation coefficient is

207

$$208 \quad r_{ij.kl} = \frac{r_{ij.k} - r_{il.k}r_{rl.k}}{\sqrt{(1-r^2_{il.k})(1-r^2_{jl.k})}} \dots\dots\dots (6)$$

209 or

$$210 \quad r_{ij.kl} = \frac{r_{ij.l} - r_{ik.l}r_{jk.l}}{\sqrt{(1-r^2_{ik.l})(1-r^2_{jk.l})}} \dots\dots\dots (7)$$

211 These partial correlations have (n-3) and (n-4) degrees of freedom respectively for first order
 212 and second order.

213 The river runoff was also analyzed to determine the type of variations which influences the
 214 runoff of 2008-2009. Seasonal variation measured by seasonal index indicated up to what
 215 level, runoff was affected seasonally (**Table 1**). A seasonal index more (or less) than 100
 216 indicated that runoff was increased (or decreased) by an amount equal to that of seasonal
 217 index in excess (or deficit) of 100 implying a positive (or negative) effect of seasonal
 218 variation. A 100% seasonal index for any month implied that there was no effect of seasonal
 219 variation on the runoff. Cyclical variation provided the period of repetition of the peak of
 220 minimal runoff.

221

222 **3. Results and Discussion**

223

224 About 73 % of the total river runoff occurred during (wet season) characterized by
 225 monsoon. The mean inflows to the estuary varied from a maximum of 1000m³/s in July to a
 226 minimum of 49m³/s in March (**Figure 1b**). Based on river runoff, the annual seasonal cycle
 227 is distinguished as high runoff months characterised by Indian summer monsoon or ISM
 228 (June-September), moderate runoff months characterised by north-east monsoon or NEM
 229 (October-December) and low runoff months or dry period (January-May).

230

231 **3.1 Prediction from polynomial fitting**

232

233 A sixth degree polynomial was obtained as the best prediction equation for 1978-2001
 234 and 1985-1989 data sets (**Figure 2a**). The equations were

235 $Y = 0.485X^6 + 19.49X^5 - 300.3X^4 + 2205X^3 - 7802X^2 + 12214.0X - 6191.0$ (8) for 1978-
 236 2001 and

237 $Y = -0.321X^6 + 13.06X^5 - 204.3X^4 + 1523X^3 - 5456X^2 + 8624.0X - 4359.0$ (9) for 1985-
 238 1989

239 where Y is the total monthly runoff and X is the month number 1,2,3....12 from June to July.

240 Equation (8) could predict 2008-2009 runoffs with only 27.36% prediction efficiency
 241 whereas equation (9) could predict it with 83.69% prediction efficiency. The lower values for
 242 prediction efficiency for the 23 years data could be due to the missing data. Since total
 243 monthly runoff was predicted with high efficiency from the past data of 1985-1989, it
 244 followed that further analysis made in this study using the 2008-2009 runoff data could be
 245 generalised.

246 For the 1985-1989 monthly runoff data, time series components were calculated and
 247 the adjusted seasonal indices for June to July are 130.89, 108.28, 92.67, 115.88, 120.41,
 248 79.58, 76.86, 107.04, 111.85, 69.98, 69.33 and 117.23% respectively. From the $2^3 \times 6$ models
 249 (Jayalakshmy, 1998), ($2^{k \times r}$, where k is the number of independent parameters and r is the
 250 number of transformations for the dependent and independent variables) the one which
 251 explained the maximum variability and in which the independent variables were uncorrelated
 252 was chosen. The optimal model for this study was the simple model,

253 $\text{LOG}_{10}Y = -1.4453 \times 10^{-7} + 0.8839 \times \text{LOG}_{10}T + 0.2405 \times S + 0.002416 \times C$ (10)

254 It could explain about 99.86% of the variability in the river runoff distribution during 1985-
 255 1989. The other models were depicted in Table1. These regression models were fitted
 256 assuming that the three components were independent. From the regression models fitted,
 257 moving average of period 2 represented the observed runoff with 94.72% of precision (**Table**
 258 **1**).

259 In this study, seasonal variation could explain only 31.32% of the variability in the
 260 runoff (Table1). Based on 1985-1989 data sets, seasonal effect was positive on the river
 261 runoff of June, July, August, October, November, February and March. For the rest of the
 262 months, seasonal effect was negative on the average. The observed runoff was mostly
 263 controlled by the trend effects of the optimal period determined.

264 Cyclical variation could explain only <1% of the variations in the runoff. The period
 265 was unique with 12 months approximately (**Figure 2b**). Hence, it could be stated that the
 266 observed runoff was mostly controlled by the trend effect and to some extent by the seasonal

267 variations only. From the graph (**Figure 2b**), it could be understood that the cycles present
268 were removed along with the trend effect as the observed curve and the trend curves were
269 almost exact. The observed cycles presented for the MA were of period 12 months.

270 In order to study the contribution of 2 period centered moving average alone on the
271 river runoff, second order partial correlation coefficient using the non transformed data was
272 computed which was 0.96 ($P < 0.001$). Similarly, contribution of seasonal variation alone on
273 the river runoff was also high with second order partial correlation coefficient as 0.93
274 ($P < 0.001$). On the other hand, contribution of cyclical variation alone on the river runoff was
275 not significant, 0.30 ($P > 0.001$). Hence, river runoff was controlled by short term variations of
276 period 2 months, but not by long term variations with periods > 2 months.

277

278 **3.2 Salinity distribution**

279 *Annual variations*

280 **Figures 3 and 4** depict the longitudinal section of salinity distribution in estuary during one
281 year. With the onset of Indian summer monsoon on May 31, 2008, the mean runoff was
282 $356 \text{ m}^3/\text{s}$ in June 2008 (**Figure 3a**). As a result, oceanic salinities were confined to near-inlet
283 stations (1, 5, and 6) and the river-end stations (2, 3, 8, and 9) were brackish. When the runoff
284 peaked in July ($1000 \text{ m}^3/\text{s}$), the estuary transformed to a salt wedge type (**Figure 3b**). Higher
285 salinities (18-34) were found only in the bottom waters of stations 1, 5, and 6. The wedge
286 formation was more prominent at stations 5 and 6 than station 1 which could be attributed to
287 the greater depths at inlet 2. All the other stations remained well mixed until September 2008
288 with depth averaged salinity as low as 0.05 (**Figures 3b-3d**).

289 By October 2008, the salinity field expansion was established (**Figure 3e**). From October
290 to December, the runoff was moderate (on average $260 \text{ m}^3/\text{s}$) and an accumulation of fresh
291 water was observed only at the upstream regions (stations 8, 9, 10). However, during the dry
292 period, the river runoff decreased remarkably such that only $49 \text{ m}^3/\text{s}$ occurred in March.
293 Under limited river flows, the estuarine water column actively mixed and tended towards
294 extremely low horizontal and vertical salinity gradients (**Figures 4b-4f**). The salinity field
295 extended up to station 10 with maximum depth averaged salinity (15.12) attained in March
296 (**Figure 4d**). In May, there was a slight increase in runoff to 2.5% of the annual runoff. The
297 aftermath of an anomalous rainfall in the catchment of Periyar caused station 1 at the inlet 1
298 to be fresh water dominated (**Figure 4f**).

299 *Daily variations*

300

301 Figure 5 illustrates the daily salinity variations allowing to verify whether the daily
302 rainfall modifies the salinity pattern of the station significantly. The daily rainfall pattern
303 (**Figure 5a**) was characterised by spikes of high rainfall during the active spells of Indian
304 summer monsoon and North east monsoon. During the Indian summer monsoon, strong spate
305 occurred in July proceeding to the beginning of August too. Fresh water salinities occurred
306 for most of the time. Occasionally, high saline waters were also observed at the bottom due to
307 the intrusion of salt wedge. By the end of August, there was a lull in monsoon resulting in
308 intrusion of high saline waters. Consequently, a single vertical profile of salinity ranging
309 from 25 to 35 was noticed. Again by the second week of September, the monsoon regained its
310 strength causing freshening at the station. The same conditions were again observed only by
311 the end of October–November characterised by North east monsoon. In contrast, during the
312 rest of the year, high saline conditions (23-35) prevailed at the station. However Small peaks
313 in rainfall were sighted in April and May which could not however, bring any effect on the
314 salinity of that station.

315 **3.4 Estuarine classifications based on hydrodynamics and runoff**

316 *3.4.1 Hansen and Rattray characterization*

317 Hansen and Rattray (1966) developed a two-parameter system of estuarine classification
318 in which the classes are delineated by the magnitudes of the relative stratification and
319 circulation parameters associated with changes in the salt balance mechanism. The diagrams
320 represent $\partial S/S_0$, where ∂S is the difference in salinity between surface and bottom and S_0
321 is the depth mean salinity, both averaged over a tidal cycle, as the ordinate. The circulation
322 parameter U_s/U_f , where U_s is the surface velocity averaged over a tidal cycle and U_f is the
323 discharge velocity, that is the rate of river discharge divided by the cross-sectional area,
324 defines the abscissa. Here, the study exercised these parameters, calculated from the time
325 series observations. These were then plotted on the relevant portion of the stratification-
326 circulation diagram for three runoff conditions (**Figure 6**).

327

328 Figure 6 shows reasonable agreement with the longitudinal monthly salinity observations
329 discussed above. For high and moderate runoff months, the estuary exhibited similar
330 characteristics. High $\partial S/S_0$ values were found at station (C) near inlet 2 tending them to fall in

331 class “1b (stratified)” of the classification diagram. Station D occupied class “4” in the
 332 diagram suggesting a salt wedge type. This was because of the depth of station C so that the
 333 salt wedge thickness was higher reaching almost the surface. However, the wedge tapered
 334 towards station D allowing more freshwater to flow over it. Recorded U_s/U_f values were
 335 above 1 for all stations. Station B in the middle of the northern arm and upstream station E
 336 were fresh water dominated during both high and moderate runoff periods. In contrast, during
 337 the dry period, the system was well-mixed (classes “1a”). Whereas the values of $\partial S/S_0$ were
 338 below 0.1, U_s/U_f ratio was almost 1. This indicated an upstream transfer of salt by diffusion.

339

340 **3.4.2. Evaluation of runoff dynamics of Cochin estuary**

341

342 Vijith *et al.*, (2009) stated that estuaries that come under the influence of Indian
 343 Summer Monsoon (ISM) and for which the salinity is never in a steady state at any time of
 344 the year are generally shallow and convergent, i.e. the width decreases rapidly from mouth to
 345 head. In contrast, Cochin estuary is having a widespread area at the upstream and has no
 346 typical river mouth entrance (as discussed under section 1.1). Adding to the complexity it has
 347 dual inlets and the tidal range is 1 m which is lower than other Indian estuaries along west
 348 coast. These typical physical features lead to its uniqueness.

349 Vijith *et al.*, (2009) had documented that the monsoonal estuaries experience total
 350 annual runoff which is many times of the estuarine volume and that there is a high
 351 “peakiness” or seasonality in the runoff. They used the following equations to represent the
 352 above two features:

$$353 \quad \eta_R = \frac{R_a}{V_e} \dots\dots\dots (11)$$

354

355 where, R_a is the volume of total annual runoff (m^3) and V_e is the volume (m^3) with respect to
 356 mean sea level in the estuary. Higher the value of η_R , higher is the runoff. η_R was calculated
 357 as 42 for the Cochin estuary indicating the total volume of the estuary is exchanged 42
 358 times(s)/year. The equation for second parameter is

$$359 \quad \eta_T = \frac{\text{Maximum Monthly runoff}}{\text{Mean Monthly runoff}} \dots\dots\dots (12)$$

360

361 **Figure 7a** shows the mean monthly runoff to monsoonal estuaries in India (Vijith *et*
 362 *al.*, 2009). It can be plainly understood that while the runoff into other estuaries average to

363 zero for about eight month-long dry season, the average runoff into Cochin estuary is never
 364 zero. A steady runoff is maintained even during the peak dry period $\Gamma_T \sim 1$.

365

366 To zoom in the dynamics of the estuary, we reduce the above mentioned parameters
 367 into monthly scale. This will provide means to examine the seasonal variations in runoff.

368 We re-define the above classification parameters as written below:

369
$$Z_R = \frac{R_m}{V_e} \dots \dots \dots (13)$$

370

371
$$Z_T = \frac{\text{Total of the maximum among daily runoff of all rivers in a month}}{\text{Total of mean Daily runoff of all rivers in a month}} \dots \dots (14)$$

372

373 where R_m is the volume of total monthly runoff (m^3) and V_e is the volume (m^3) with respect
 374 to mean sea level in the system. R_m is computed from daily runoff values shown in **Figure**
 375 **1b**. Z_T represents the daily variations in runoff. The computed values are presented in **Figure**
 376 **7b**.

377

378 During June Z_R was 2.06 when Indian summer monsoon was in the progressing stage
 379 whereas for the rest of the months of wet season $Z_R > 5$. The observed maximum monthly
 380 runoff of wet season was $3.606 \times 10^9 m^3$ in July. For the moderate runoff months (October -
 381 December), the values were $1 < Z_R < 4$ and $1 < Z_T < 3$ (**Figure 7b**). From January-April, Z_R was
 382 about 0.3 and Z_T was 2. This indicated that although there were prominent daily runoff
 383 variations, for no single day of each month during the period, the runoff could flush the
 384 estuary. For it to occur, the runoff obtained should exceed to above 70% of the estuarine
 385 volume. During May, the runoff was higher which completed the annual cycle with Z_R and Z_T
 386 showing 0.8 and 2 respectively.

387

388 **Figure 7c**, shows the Z_R values of Cochin estuary with other estuaries in the world.
 389 The analysis showed that Z_R was an order of less than one for Tamar, Delaware, and Thames
 390 estuaries for all months. In the case of Columbia estuary, Z_R values were more or less
 391 comparable with Cochin estuary. However, the peak runoff in attained by Columbia in June
 392 was 6.5 which was less than that of Cochin estuary by 2. For estuary, the peak in July with a
 393 value of 8.5 was featured by Indian summer monsoon. The influence of North east monsoon
 394 on flushing of the estuary was negligible. The minimum Z_R of 0.3 occurred during peak dry.

395 Cochin estuary exhibited large range of Z_R values over the months compared to all other
 396 estuaries.

397

398 To explore the flushing nature more closely, Z_R ratio for the two arms of estuary were
 399 calculated separately (**Figure 7c**). It was found that, for July, with the Periyar River runoff in
 400 the northern arm Z_R ratio was 3.7. The runoff from all the other rivers was responsible for Z_R
 401 to go as high as 6.7 in the southern arm. The volume of southern arm was about 5 times larger
 402 than the northern arm. Notwithstanding this fact, the runoff into the south flushed the volume
 403 of the southern arm almost twice as that of northern arm. During August, the lull in monsoon
 404 (about $200 \text{ m}^3/\text{s}$ decrease from July) was characterised by an increase in runoff in the
 405 northern arm and a decrease in runoff into the southern arm. Consequently, an equal flushing
 406 of both arms ($Z_R \sim 5$ in both the arms) resulted in transforming the estuary into a river. This
 407 implied that the uniform flushing of all the sections of the estuary could not be directly
 408 related to the ‘peakiness’ of monsoonal spell and the subsequent runoff.

409

410 **4. Cochin estuary in a quasi-steady state**

411

412 Implicit in several estuarine classification schemes commonly used for understanding
 413 estuarine dynamics is a steady state assumption. By the term "steady state" is meant that the
 414 average of the salinity concentration over a tidal cycle does not change from tide to tide if the
 415 river flow remains constant (Stommel 1953). In such cases, during each tidal cycle the
 416 salinity at any location varies with the stage of the tide, but on successively similar tidal
 417 stages the salinity returns to substantially the same value (Ketchum and Rawn, 1951). In an
 418 estuary like Cochin estuary, such a steady state can be expected during the peak dry period
 419 (January-April). In order to establish this fact, we use the salt balance equations to determine
 420 the salinity steadiness in the Cochin estuary.

421 The general unsteady salt balance is given by:

422

$$423 \frac{\partial}{\partial t} \int_x^{x_r} S_{(x)} A_{(x)} dx + RS = K_{unst} A \frac{\partial S}{\partial x} \dots\dots\dots(15)$$

424

425 where $S_{(x)}$ is the salinity integrated over the volume of the estuary, and A is the cross sectional
 426 area, R is the river runoff, S is the average salinity. K_{unst} is the unsteady horizontal diffusion

427 coefficient computed in the axial direction from x until the upstream location x_r .
 428 With the steady state assumption, the time dependent term of equation (15) vanishes. The
 429 equation can then be re-written as:

430

$$431 \quad RS = K_{st} A \frac{\partial S}{\partial x} \dots\dots\dots(16)$$

432

433 K_{st} is the horizontal diffusion coefficient under equilibrium (steady state) conditions.

434 If the estuary is in a steady state, the total salt content of the estuary does not change, so
 435 the same volume R will have to leave the estuary at its mouth during one tidal cycle. Thus, by
 436 comparing K_{unst} with K_{st} , the steadiness of the salt balance can be diagnosed roughly.
 437 Dividing equation (14) by (15), the ratio of K_{unst} to K_{st} can be obtained as:

438

439

$$440 \quad \frac{K_{unst}}{K_{st}} = \frac{\frac{\partial}{\partial t} \int_x^{x_r} S_{(x)} A_{(x)} dx}{RS} + 1 \dots\dots\dots(17)$$

441

$$= \Phi + 1 \dots\dots\dots(18)$$

442

443 The steadiness of the salt balance was diagnosed for the months, January-April, when Φ was
 444 continuously > 0 . The integral term in (17) was estimated using longitudinal salinity
 445 measurements (**Figures 4-5**) from x to the upstream location x_r for two consecutive months.
 446 The averages of salinity S and runoff R for these two months were used. The ratios were
 447 computed for all sections from x (station 1) to x_r (station 10).

448 The analyses proved that the ratios approached 1 most of the time throughout the
449 estuary. Occasionally, a maximum value of 1.5 was also obtained (**Figure 8**). This suggested
450 a steady state or rarely a quasi-steady state. The total salt content remained constant for the
451 peak dry period. The period from March to April was in an acute steady state even at the
452 upstream. Specifically, along sections from stations 5 to 7, the balance was better achieved
453 than the other locations. This is possible as Muvattupuzha joins between the regions which
454 supplied a constant runoff. It is the only river that caused freshening in the southern arm
455 during the period. The upstream salt flux was balanced by this runoff induced oceanward
456 advective flux asserting a steadiness in salt balance.

457

458 **Figure 9** illustrates the water level and salinity variations over a tidal cycle (depth
459 averaged mean salinity from the CTD profiles over a tidal cycle) at five stations during
460 February 2010. In each case the salinity at successive high tides returned to the value
461 previously observed approximately. Therefore, Hansen Rattray classification holds well for
462 this particular steady state of the estuary. Whatever be the runoff occurred during the period,
463 it is not sufficient to bring the salinity at the upstream to zero. This typical feature is due to
464 the diverging geometry of the estuarine channel unlike other Indian estuaries such as
465 Mandovi and Zuari channels which are strongly convergent at the upstream regions (Manoj *et al.*,
466 *2009*). For the Mandovi and Zuari, although the tidal flushing times are in the order of
467 days during the dry season, so much of freshwater remains available at the upstream and
468 these systems always lag behind steady state (Vijith *et al.*, 2009).

469

470 The steadiness in salinity during dry period is even reflected in the abundance of
471 zooplankton species which showed little variations during tidal cycles (Mathupratap *et al.*,
472 1977). They had opined that these species appear to develop behavioural mechanisms in
473 response to tidal changes which keep it in the water of same salinity throughout the tidal
474 cycle by having some kind of biological clock or signal. So, we conclude that estuary is in a
475 steady state for some time during a year and deserves to be placed under a ‘special’ category
476 among the monsoonal estuaries.

477

478 **5. The Physical-biological coupling**

479

480 Cochin estuary is one of the largest productive ecosystems along west coast of India with
481 an estimated annual gross production of nearly 300gC/m^2 (Qasim *et al* 1969). Its bioceonosis
482 can be recognized as a physically controlled community. It may be called as a "tropical
483 monsoonal estuary" due to the pronounced influence of monsoon on the ecology of the
484 system bringing about a total change in the environment and fauna (Madhupratap *et al.*,
485 1977). In such estuaries, the seasonality in salinity is a key feature as the ecosystems have to
486 adjust accordingly. CMEB is more productive at all levels during dry season. The salinity
487 gradient during the period favoured large species richness, species diversity and species
488 evenness in zooplankton (Jyothibabu *et al.*, 2006). Whereas in monsoon, the abundance of
489 phytoplankton grazers (zooplankton) is reduced and this altered the trophic food web of the
490 estuary resulting in substantial amount of unconsumed carbon at primary level (Madhu *et al.*,
491 2010). A qualitative shift in phytoplankton composition (Qasim, 2003) and an increase in its
492 biomass owing to high residence times (Shivaprasad *et al.*, 2012, 2013) were also reported
493 during peak dry conditions. In essence, the dry season provides a biotope supporting the
494 survival of various high species as competitors, expanding their overlapping niches in space
495 with time because of the facility provided by salinity intrusion. The impact of monsoonal
496 effluxes and high flushing evokes its elimination and an 'essential' cleanup of the estuary.

497

498 **6. A new nomenclature: Cochin Monsoonal Estuarine Bay**

499

500 The present analyses manifested that the assumptions implicit in the classification
501 schemes discussed above limits their applicability to Cochin estuary. There arises a need for a
502 comprehensive classification system representing all the dominant conditions of the estuary.
503 Such an approach was suggested by Whitefield (1992) for African estuaries using a
504 combination of physiographic, hydrographic and salinity features. According to him,
505 estuarine bays are estuaries that may be either natural or partly artificial due to dredging
506 activities in the mouth and harbour region. They have a large tidal prism exceeding 10×10^6
507 m^3 and tides are the dominant force driving mixing of water column. The salinity ranges
508 from 20-35 and near marine conditions may extend even to the upper reaches.

509 Cochin estuarine system is partly artificial due to the anthropogenic activities like land
510 reclamations (Gopalan *et al.*, 1983) and dredging at inlet 2 (Balchand and Rasheed 2000),
511 frequently modifying its geomorphology. Also, the tidal prism of Cochin inlet is estimated at
512 $107.8 \times 10^6 \text{m}^3$ during Indian summer monsoon, $18.6 \times 10^6 \text{m}^3$ during moderate runoff
513 months (October to December) and $31.5 \times 10^6 \text{m}^3$ during the dry season (Rama Raju *et al.*,

514 1979). The salinity conditions of a bay are found in the lower reaches only during dry period.
515 Meanwhile, the maximum salinity observed at the upstream is never greater than 15. Hence, a
516 salinity gradient from mouth to head persists throughout the dry period. Peak monsoonal
517 spells and runoff may entirely change the estuary from an estuarine bay to a riverine system.
518 This transformation plays a fundamental role in the ecology of the system. Thus, 'Monsoonal
519 Estuarine Bay' seems to be an appropriate term for this estuary.

520

521 ***7. Synthesis and Conclusion***

522

523 The runoff into estuary is never zero at any time of the year. It is a unique divergent
524 estuary with a widespread area at the upstream. During the wet season and moderate runoff
525 months, the salinity field is extremely sensitive to the drastic variations in river runoff even
526 on daily time scales. Saline water creeps in slowly during moderate runoff months, but then
527 persists unabatedly in the following peak dry season. During peak dry period, the salinity
528 values are high throughout the system with a gradient from mouth to head and the variations
529 in runoff is slow. The lower reaches behave like an extension of the coastal waters and
530 salinity ranging from 10-12 is observed at the upstream and the water column is well mixed.
531 The runoff that enters is only 30% of the estuarine volume so that zero salinity is never
532 attained at the upstream. The 'little but constant' runoff is mainly contributed by
533 Muvattupuzha River flowing into southern arm which is not sufficient to flush the large
534 upstream volume.

535 Fluctuations in the estuary are of extreme nature with regard to salinity. The new
536 terminology 'Monsoonal Estuarine Bay' encapsulates the salinity gradient of the Cochin
537 estuary ranging from completely riverine to completely saline. The term 'Monsoonal'
538 succinctly describes the unsteadiness of salinity of wet season. The possibility of the estuary
539 turning to a river cannot be ruled out. 'Bay' conditions are accomplished during peak dry
540 season when the estuary is in a steady state with little constant runoff. During the rest of the
541 year, the system behaves only as a true estuary. The gist of the previous studies is that the
542 ecosystem and ecology respond well to this varying salinity and environment. The
543 terminology may be used for future works due to its significance. It provides basic
544 information about the physiographic, hydrographic, salinity and ecological features of the
545 system.

546

547 **8. Acknowledgement**

548 The authors are thankful to the Director, National Institute of Oceanography, Goa, and
 549 also to the Scientist-in-Charge of the Regional Center, for encouragement. Shivaprasad
 550 Amaravayal is also grateful to CSIR, New Delhi, for financial support in the form of Senior
 551 Research fellowship. This work is a part of A.Shivaprasad's doctoral research. Manoj N. T. is
 552 thankful to DST, Govt. of India for financial support under SERC-fast track scheme. This
 553 study would not have been possible without the active support of the project team; we
 554 express our sincere thanks to them. This work forms part of project "Ecosystem modeling of
 555 Cochin estuary" funded by ICMAM. This is NIO contribution no. 5349.

556

557 **References**

- 558 Abhilash, K.R., Raveendran, T.V., Limna Mol, V.P. Deepak, M.P., 2012. Sediment Oxygen
 559 Demand in Cochin backwaters, a tropical estuarine system in the south-west coast of
 560 India. *Marine Environmental Research* 79,160-166
- 561 Balachandran, K.K., Reddy, G.S., Revichandran, C., Srinivas, K., Vijayan, P.R., Thottam, T.J.,
 562 2008. Modelling of tidal hydrodynamics for a tropical ecosystem with implications for
 563 pollutant dispersion (Cochin Estuary, Southwest India). *Ocean Dynamics* 58, 259–273
- 564 Balchand, A.N.; Rasheed, K., 2000. Assessment of short term environmental impacts on
 565 dredging in a tropical estuary. *Terra et Aqua* 79,16-26
- 566 Cameron, W. M., and Pritchard, D. W., 1963. Estuaries. In: M. N. Hill (editor), *The Sea*. John
 567 Wiley and Sons, New York.2, 306 - 324.
- 568 Dyer, K.R., 1973. *Estuaries, a physical introduction*. John Wiley, London, 140 pp.
- 569 Dyer, K.R., 1995. Sediment transport processes in estuaries. In: G.M.E. Perillo (ed.).
- 570 Dyer, K.R., 1997. *Estuaries, a physical introduction, second edition*. John Wiley, London, 195
 571 pp.
- 572 Fairbridge, R.W., 1980. The estuary: its definition and geodynamic cycle. In: E. Olausson, I.
 573 Cato (Eds.), *Chemistry and biogeochemistry of estuaries*. John Wiley, Chichester, pp. 1–35.
- 574 Gopalan, U.K., Doyil, T.V., Udaya Varma, P., Krishnankutty, M., 1983. The shrinking
 575 backwaters of Kerala. *Journal of Marine Biological Association India* 25,131-141.

- 576 Hansen, D.V., Rattray, M., 1966. New dimensions in estuary classification. *Limnology and*
577 *Oceanography* 11 (3), 319–326
- 578 Hayes, M.O., 1975. Morphology of sand accumulation in estuaries. In:
579 L.E.Cronin(ed.), *Estuarine Research*, Vol. 2. Academic Press, New York, pp. 3–22.
- 580 Jayalakshmy, K.V., 1998. Biometric studies on trophic level relations in the Indian Ocean,
581 PhD Thesis, Cochin University of science and technology.
- 582 Jyothibabu, R., Madhu, N.V. Jayalakshmi, K.V. Balachandran, K.K. . Shiyas, C.A Martin,
583 G.D., Nair, K.K.C., 2006. Impact of freshwater influx on microzooplankton mediated food
584 web in a tropical estuary (Cochin backwaters e India), *Estuarine, Coastal and Shelf*
585 *Science* 69, 505-518
- 586 Ketchum, B.H., Rawn, A. M., 1951. The Flushing of Tidal Estuaries [with Discussion]
587 *.Sewage and Industrial Wastes* 23 (2),198-209
- 588 Madhu, N. V., Balachandran, K. K., Martin, G. D., Jyothibabu, R., Shoji, D. T., Nair, M,
589 Joseph, T., Kusum, K. K., 2010. Short-term variability of water quality and its
590 implications on phytoplankton production in a tropical estuary (Cochin backwaters –
591 India. *Environmental Monitoring and Assessment* 170 (1-4), 287-300
- 592 Madhupratap , M., Rao, T.S.S., Haridas, P., 1977. Secondary production in the Cochin
593 backwaters, a tropical monsoonal estuary. *Proceedings of the Symposium on Warm Water*
594 *Zooplankton*, Special Publication, UNESCO/NIO: 515-519
- 595 Manoj, N.T,Unnikrishnan, A.S., Sundar, D., 2009. Tidal Asymmetry in the Mandovi and Zuari
596 Estuaries, the west coast of India . *Journal of Coastal Research* 25(6), 1187-1197
- 597 Martin, G.D., Vijay, J.G., Laluraj, C.M1., Madhu, N.V., Joseph, T., Nair, M., Gupta, G.V.M2.,
598 Balachandran, K.K., 2008. Fresh water influence on nutrient stoichiometry in a tropical
599 estuary,southwest coast of India. *Applied Ecology And Environmental Research* 6(1), 57-
600 64

- 601 Pritchard, D.W., 1955. Estuarine circulation patterns. Proceedings Amer. Soc. Civil
602 Eng.(ASCE) 81, Paper no. 717, 11 pp.
- 603 Pillai, R.S.N. and Bagavathy, V., 2009. Statistics theory and practice.published by S.chands
604 company Ltd, 868pp
- 605 Qasim, S.Z., 2003. Indian Estuaries. Allied Publication Pvt. Ltd., Heriedia Marg, Ballard
606 estate, Mumbai, 259
- 607 Qasim, S.Z.; Gopinathan, C.K. et al., 1969. Tidal cycles and the environmental features of
608 Cochin backwater (A tropical estuary), Proceedings of the Indian academy of sciences,
609 Bangalore.
- 610 Rama Raju, V.S., Udaya Varma, P., Pylee, A., 1979. Hydrographic characteristics and tidal
611 prism at the Cochin harbour mouth. Indian Journal of Marine Sciences 8, 78-84
- 612 Rao, K.K., Balasubramanian., 1996, Distribution of foraminifera in the Cochin
613 estuary.Journal of marine biology association of India 38 (1&2), 50-57.
- 614 Sankaranarayanan, V.N., Qasim, S.Z., 1969. Nutrients of the cochin backwaters in relation
615 to environmental characteristics. Marine biology 2,236-247
- 616 Savenije, H.H.G., 2005. Salinity and tides in alluvial estuaries. Elsevier, Amsterdam, 197 pp.
- 617 Shivaprasad, A., Vinita, J., Revichandran, C., Manoj, N.T., Srinivas, K., Reny, P.D., Ashwini,
618 R., Muraleedharan, K.R., 2012.Influence of saltwater barrage on Tides, Salinity and
619 Chlorophyll a in Cochin Estuary, India. Journal of Coastal Research DOI:
620 10.2112/JCOASTRES-D-12-00067.1
- 621 Shivaprasad, A., Vinita, J., Revichandran, C.,Reny, P. D., Deepak, M. P., 2013.
622 Muraleedharan, K. R., and Naveen Kumar, K. R.:Seasonal stratification and property
623 distributions in a tropical estuary (Cochin estuary, west coast, India),_Hydrol. Earth Syst.
624 Sci._17, 187-199
- 625 Snead, E.R., 1982. Coastal landforms and surface features.Hutchinson Ross publishing
626 company,Stroudsburg,Pennsylvania:ISBN:0-87933-052-X

- 627 Snedcor, G.W., Cochran, W.G., 1975. Statistical methods, 6th edn. Oxford IBH Publication
628 Co. New Delhi
- 629 Stommel, H., 1953. Computation of Pollution in a Vertically Mixed Estuary. Sewage and
630 Industrial Wastes 25 (9), 1065-1071
- 631 Strikwerda, M., 2004. Cochin Estuary morphological modeling and coastal zone
632 management. Ph.D Thesis, University of Delft, Germany
- 633 Udaya Varma, P., Pylee, A., Rama Raju, V.S., 1981. Tidal influence on the seasonal variation
634 in current and salinity around Willington Island. Mahasagar 4(4), 225-237
- 635 Valle-Levinson, A., 2010. Contemporary Issues in Estuarine Physics, Cambridge University
636 Press, Cambridge, UK, 2010, 326 pp.
- 637 Vijith, V., Sundar, D., Shetye, S.R., 2009. Time-dependence of salinity in monsoonal
638 estuaries. Estuarine, Coastal and Shelf Science 85, 601-608.
- 639 Vinita J., Shivaprasad A., Revichandran C., Manoj N.T., Muraleedharan K.R., Binzy J.
640 2013 (in press) Journal of Coastal research. DOI: 10.2112/JCOASTRES-D-13-00038.1
- 641 Whitefield, A.K., 1992. Characterization of South African Estuarine systems. African Journal
642 of aquatic sciences 18(1/2) 89-103

643

644 **List of figures**

645

646 **Fig. 1(a)** The Cochin estuary (West coast, India), showing rivers and extent of the system,
647 having two inlets to Arabian Sea, Munambam Inlet (north) and Cochin inlet (middle of the
648 extent of the system). Daily station is located 5 km away from Cochin inlet. Monthly
649 longitudinal and time series stations are discerningly marked. **1 (b)** Runoff from 6 rivers for
650 the period of 1 year (June 2008 to May 2009).

651 **Fig. 2 (a)** Polynomials of different degrees for the monthly total runoff **(b)** Spline smoothing
652 of Time series components of the river runoff data.

653 **Fig. 3** Longitudinal distribution of salinity measured monthly once during June - November
654 2008. The Cochin inlet is at the coordinate origin. The northern / southern arm stations are at

655 positive /negative distances, respectively. Times of each station appear along the lower x-
656 axis. Grey shaded area shows the bathymetry of the estuary.

657 **Fig. 4** Longitudinal distribution of salinity measured monthly once during December 2008, to
658 May 2009. The Cochin inlet is at the coordinate origin. The northern / southern arm stations
659 are at positive /negative distances, respectively. Times of each station appear along the lower
660 x-axis. Grey shaded area shows the bathymetry of the estuary.

661 **Fig. 5 (a)** The daily rainfall pattern (May 2008-June 2009) **(b)** The daily salinity pattern of
662 the station situated 5km away from Cochin Inlet.

663 **Fig. 6** Hansen –Rattrey classification diagram for Cochin Estuary.

664 **Fig. 7 (a)**Mean monthly runoff to monsoonal estuaries(Vijith et al.,2009).(b) Positions of
665 each month of Cochin estuary on the (Z_R , Z_T) plane.(c)comparison of Z_R of major estuaries in
666 the world with Cochin estuary

667

668 **Fig. 8** The ratios of $K_{unsteady}$ to K_{steady} calculated as shown in equation (7).

669 **Fig9** Average salinity variations during a tidal cycle for monthly time series stations during
670 the dry period.

671 **List of tables**

672

673 **Table 1** Multiple regression model results based on log transformed runoff, log transformed
674 trend and non transformed seasonal and cyclical variations

675