

1 **Impacts of climate variability on**
2 **wetland salinization in the North**
3 **American Prairies**

4
5
6 U. Nachshon^{1*}

7 A. Ireson¹

8 G. van der Kamp²

9 S. R. Davies³

10 H. S. Wheeler¹

11
12 ¹*School of Environment and Sustainability and the Global Institute for Water Security, University of*
13 *Saskatchewan, 11 Innovation Boulevard, Saskatoon, Saskatchewan S7N 3H5, Canada.*

14 ²*Environment Canada, 11 Innovation Boulevard, Saskatoon, Saskatchewan S7N 3H5, Canada.*

15 ³*Imperial College London, Department of Civil and Environmental Engineering.*

16
17
18
19 * *Corresponding author (unachshon@gmail.com)*

20

21 **Abstract**

22 The glaciated plains of the North American continent, also known as the ‘prairies’, are a
23 complex hydrological system characterized by hummocky terrain, where wetlands, containing
24 seasonal or semi-permanent ponds, occupy the numerous topographic depressions. The prairie
25 subsoil and many of its water bodies contain high salt concentrations, in particular sulfate salts,
26 which are continuously cycled within the closed drainage basins. The period between 2000 and
27 2012 was characterized by an unusual degree of climatic variability, including severe floods and
28 droughts, and this had a marked effect on the spatial distribution, water levels and chemical
29 composition of wetland ponds. Understanding the geochemical and hydrological processes under
30 changing environmental conditions is needed in order to better understand the risk and mitigate
31 the impacts of future soil and water salinization.

32 Here we explore salt dynamics in the prairies using field observations from St. Denis,
33 Saskatchewan, taken over the last 40 years. Measurements include meteorological data, soil
34 moisture, soil salinity, groundwater levels and pond water volume, salinity, and chemical
35 composition. The record includes periods of exceptional snow (1997, 2007) and periods of
36 exception rainfall (2010, 2012), both of which resulted in unusually high pond water levels.
37 However, severe salinization only occurred in response to extreme summer rainfall. We
38 hypothesize that since rainfall and snowmelt activate different hydrological pathways, they have
39 markedly different impacts on salinization. We propose that a wet condition associated with high
40 snowmelt conditions does not pose a strong threat to salinization, which has important
41 implications for agricultural planning. Whilst this hypothesis is consistent with our conceptual
42 understanding of the system, it needs to be tested further at a range of field sites in the prairies.

43

44

45 **1. Introduction**

46 Surface water and shallow ground water salinization is a problem affecting agriculture, water
47 resources and ecosystem health in many areas of the world, including Australia (Dehaan and
48 Taylor, 2002; Rengasamy, 2006), the Aral Sea region (Micklin, 2007), playas and sabkhat
49 environments (Tyler et al., 2006) and many other areas (Rengasamy, 2006). Salt dynamics are
50 driven by hydrological processes, which cycle seasonally but also change over long time scales
51 as a result of climate variability and change, and changes in land use and land management
52 practices. The physical mechanisms that lead to salinization operate in different ways in different
53 areas, largely as a function of local climatological, hydrological and geological conditions. This
54 paper takes the salt rich glaciated plains of North America, known as the prairies, as a case study
55 to explore how recent climate variability has dramatically affected the salinity of ponds. The
56 prairies spread from Alberta, Saskatchewan and Manitoba in Canada, through Montana and the
57 Dakotas in U.S.A. The landscape is characterized by hummocky terrain, where wetlands and
58 ponds occupy the ubiquitous depressions (Winter, 1989; van der Kamp and Hayashi, 2009). The
59 region supports a diverse community of wildlife species and major agricultural industry that are
60 highly sensitive to the hydrological conditions (Wienhold et al., 1989). Hydrological processes in
61 the prairies are complex and unusual, characterized by closed basins isolated from any regional
62 drainage network, with drainage via a spill and fill sequence into terminal ponds (Shook and
63 Pomeroy, 2011); Snowmelt runoff from surrounding uplands and precipitation on the wetlands
64 are the dominant hydrological inputs into the wetland systems together with occasional runoff
65 events due to high-intensity rainfall, and snow distribution over the landscape is an important
66 control on the hydrology (Shaw et al., 2012; Spence and Woo, 2003); semi-arid conditions result
67 in minimal shallow groundwater recharge from uplands, while ponds drain to shallow
68 groundwater beneath the depressions (van der Kamp and Hayashi, 2009). Nachshon et al. (2013),
69 based on previous work (e.g., Keller et al., 1991; Hayashi et al., 1998b; Berthold et al., 2004;
70 Heagle et al., 2013) presented a conceptual model to describe the major salt dynamics occurring
71 within the glacial till portion of the prairies under various land use and climatic conditions. In
72 humid environments, streamflow is an integrated variable that aggregates the effect of the
73 climatic boundary conditions and the land use/management mediated hydrological processes

74 occurring within the watershed (Wheater et al., 1993). In the prairies, where there is often no
75 regional streamflow, these factors are expressed in the pond storage, and therefore studying the
76 ponds is a good way to understand the system sensitivity to a particular change. This work
77 explores pond salinization using a 20 year record of observations from a field site in the
78 Canadian prairies, which in the past decade has been subject to extremely variable climatic
79 conditions – including both extreme droughts and floods. In particular this study explores the
80 impacts of extreme precipitation on salinity, considering separately summer rainfall and winter
81 snowfall.

82 **2. Methods**

83 **2.1 Field site**

84 The field site is located at the St Denis National Wildlife Area (106°05'0.20" W, 52°12'31.32"N),
85 approximately 40 km east of Saskatoon, Saskatchewan, Canada (**Figure 1A**). The St. Denis area
86 has a hummocky topography and consists of a clayey glacial till, with typical hydrogeology for
87 the glacial till portion of the prairies (Hayashi et al., 1998a). The main part of this study focuses
88 on ponds 107, 108A, and 109 (**Figure 1**). Pond 109 is a semi-permanent pond, i.e., the pond
89 remains wet all year for most years. Ponds 107 and 108A are ephemeral ponds that dry out by
90 the end of most summers. Most of the area is cultivated, with the exception of the area to the
91 north of pond 109 which is a natural grassland, and the numerous wetlands (that is, the ponds
92 and their riparian zones). Pond 109 has a substantial riparian zone with a ‘willow ring’ with a
93 width of ~10 m around the pond, covered by trembling aspen trees, balsam poplar and willows
94 rising up to ~8 m. The pond 107 and 108A willow rings are minor compared to pond 109, with a
95 width of 1-2 m, and covered mainly by cat-tails (*Typha*) rising up to ~1.5 m. A surface runoff
96 flow path exists from ponds 107 through pond 108A to pond 109, and pond 109 can be
97 considered as the terminal pond of this local watershed, at least with respect to surface runoff.

98 **2.2 Precipitation data**

100 Precipitation data used in this study are from the nearest climate station with continuous records
101 from 1993 to 2012, the Saskatchewan Research Council Saskatoon climate reference station
102 (Beaulieu and Wittrock, 2013), about 35 km west of the St Denis site. The data are presented as

103 cumulative amounts of summer rain (April – September) and winter precipitation (October –
104 March), mostly snow. The winter data are not corrected for the wind under-catch effect, for
105 example as described by Mekis and Vincent (2011). Comparison with the corrected precipitation
106 data, available to 2007, indicates that the actual winter precipitation is under-reported by ~40 to
107 50%. However, for the purpose of this paper these readings are sufficient to indicate inter-annual
108 variability; the focus is on the differences between the reported winter precipitation in every year
109 (from the autumn of previous year (October) to the spring of the reported year (March)) and the
110 long-term average.

111 **2.3 Groundwater and surface water data**

112 Ground water levels at a number of piezometers scattered throughout the site have been
113 measured. For this work data from piezometers 94W7 (hereafter the “upland piezometer”) and
114 802P1 (hereafter the “pond piezometer”) were used (**Figure 1C**). The upland piezometer, located
115 north of pond 109, is screened from 1.8 – 5.3 m bgl (below ground level) and the pond
116 piezometer, located within Pond 109, is screened from 5.8 – 7.6 m bgl. Water table depths were
117 measured continually from 1997 to 2012, on a daily basis. The depth at the deepest point of pond
118 109 was measured manually on a monthly basis during the summers from 1968 to the present
119 (Conly et al., 2004). Vertical hydraulic gradients between the pond piezometer and the pond
120 water were small. Here, a continuous pond water level time series was constructed from 2007
121 onwards, by using the daily logged piezometer data to temporally infill between the monthly
122 manual surface water level measurements. The pond’s water volume V (m³) was computed based
123 on the work of Hayashi and van der Kamp (2000):

$$124 \quad V = 1420h^{2.24} \quad (1)$$

125 where h is the depth of water at the centre of the pond. These authors limited and validated
126 **Equation 1** for maximal h of 1.2 m, since that was the deepest measurement of the water pond
127 depth at that time. Since over recent years deeper pond water levels were recorded, the original
128 bathymetric survey data of Hayashi and van der Kamp (2000), which is archived at Environment
129 Canada, was used to compute V for $1.2 < h < 1.8$ m and to revalidate **Equation 1** for these depths. It
130 was found (data not shown) that **Equation 1** is valid for these depths with errors smaller than
131 5%.

Comment [MJ1]: Were water table wells installed in addition to piezometers? Piezometers measure hydraulic head (not water table) whereas observation wells measure height of unconfined water table where pressure head is atmospheric.

132 **2.4 Pond water salinity**

133 Electrical Conductivity, EC, was measured at 14 ponds at St Denis (**Figure 1B**) several times a
134 year during the summer months from 2009 to 2012. In addition, in pond 109 EC measurements
135 were taken every few weeks in the summer months from 1993 to 2012.

136 Major ion analysis was conducted annually for pond 109 water from 1994 to 2009, and in 2012.
137 Data from 2007-2009 and 2012 are used in this study.

138 Pond 109 EC and volume measurements permit an estimate of the mass of dissolved salts (M_{salt})
139 (kg) in the water, based on an approximate relationship between EC and dissolved salt mass
140 (Rhoades, 1996):

141
$$M_{salt} = 0.00064 \cdot EC \cdot V \quad (2)$$

142 where M_{salt} is in kg, EC is in $\mu\text{S}/\text{cm}$ and V is in m^3 .

143 **2.5 Transect measurements**

144 Spatially-detailed manual measurements of EC and water levels along a transect from Pond 107
145 to pond 108A were obtained over a rainy 25 day period in July 2012. A series of 16 mini
146 observation wells were installed to measure shallow water table changes beneath the upland
147 between pond 107 and 108A (**Figure 1C**). The transect was located along the shortest path
148 between the ponds, in the lowest part of the landscape. The spill elevation of pond 107 is
149 approximately 1 m higher than that of pond 108A and the lateral distance between the ponds is
150 35 m. Each mini observation well consists of a PVC tube, 75 cm long, inner diameter of 1.27 cm,
151 perforated along its entire length. Each tube's lower end was sealed and they were inserted into
152 the ground by direct push, to a depth of ~60 cm. Water levels in the mini observation wells were
153 measured manually with a ruler and EC measurements were taken *in situ* by a portable EC probe
154 (Cole-Parmer, 1481-60, Canada). These measurements were taken on July 5, 6, 11, 19 and 24.
155 For this period, precipitation (rain) measurements were taken hourly at the climate station
156 located within the St. Denis field site (operated by Environment Canada). In addition, on July 24
157 an EM38 probe (Geonics Ltd, Canada) was used to measure soil EC around pond 108A and pond
158 109. Measurements were taken at 1 m intervals along the pond's edge. Since the EM38 readings
159 are sensitive both to water content and salinity, calibration had to be done to obtain an estimate

160 of pore water EC. Since the soil around the pond is fully saturated, it was assumed that the
161 changes in the EM38 readings were due to salinity differences. Manual EC measurements of the
162 pore water at specific points around the pond were taken using the EC probe, and these data were
163 used to calibrate the EM38 readings.

164 3. Results and discussion

165 3.1 Field scale changes in pond water levels

166 In the prairies, changes in pond storage are a response to a large number of often confounding
167 processes, including rainfall amount, timing and intensity, snow spatial distribution and amount,
168 timing of snowmelt, the spatiotemporal pattern of surface runoff versus infiltration (strongly
169 affected by soil freeze-thaw processes, as well as land use), and the spatiotemporal pattern of
170 evapotranspiration (again strongly affected by land use). A particularly important factor is pre-
171 freeze up soil moisture content: if the soils are very wet when they freeze they have a very low
172 infiltration capacity, and hence runoff over frozen soils during the subsequent melt period is
173 more intense. This mechanism explains the widespread flooding in Saskatchewan and Manitoba
174 in 2011, which was attributed to high rainfall in the summer of 2010 (see **Figure 2**), leading to
175 high antecedent soil moisture. Another factor is the timing of snowfall and accumulation. Earlier
176 snowfall means that the relatively warm soils are more effectively insulated, and hence the extent
177 of soil freezing is less, and a larger proportion of snowmelt is expected to go to infiltration versus
178 runoff. This discussion serves to demonstrate that the hydrological processes in the prairies are
179 highly complex, and rainfall-runoff type of responses that apply elsewhere, do not tend to work
180 in this environment. **Figure 2** contrasts summer and winter precipitation (**Figure 2A**) with an
181 integrated measure of annual maximum pond level across the site (**Figure 2B**). The pond level
182 measurement was calculated by taking the mean of the normalized depths of 12 ponds (ponds 1,
183 2, 20, 25, 26, 35, 36, 37, 50, 60, 109, 120). Each pond depth time series (limited to the open-
184 water season) was normalized by dividing the depth by the maximum depth observed in the
185 period 1993 – 2012. The fact that the averaged level is never 100% indicates that different ponds
186 reached their maximum level in different years. For the purposes of this discussion, water levels
187 of 70% and above are assumed to represent unusually wet conditions, and these were measured
188 in 2006, 2007, 2010, 2011 and 2012. It is clear from **Figure 2** that there is a considerable delay

Deleted: then

190 before precipitation extremes are translated into responses in the pond levels, and also that there
191 is a different sensitivity to snow versus rainfall. For all of the wet years, this is summarized,
192 qualitatively, in **Table 1**. It is also useful to this discussion to consider the years 1994, when
193 current and antecedent conditions were at or slightly below the average for the period of record,
194 and 1997, where the snowpack was deepest, and the pond levels were high, though not up to the
195 70% threshold.

196 **3.2 Field scale changes in pond salinity**

197 Salinity of ponds in the prairies is highly variable (Euliss et al., 2004). Stewart and Kantrud
198 (1972) and Millar (1976) distinguished between fresh water ponds (EC, < 500 $\mu\text{S}/\text{cm}$),
199 moderately brackish ponds (EC between 500 – 5000 $\mu\text{S}/\text{cm}$) and brackish-saline ponds (EC >
200 5000 $\mu\text{S}/\text{cm}$). The reasons for these differences in salinity are understood to be a function of how
201 the ponds interact with the groundwater and surface runoff. Fresh “recharge” ponds lose water to
202 groundwater, brackish-saline “discharge” ponds gain water from groundwater and surface water
203 (Nachshon et al., 2013), and the moderately brackish ponds are a more complex combination of
204 input and output of surface and subsurface water of various degrees of salinity. The St Denis
205 field site is only 1.6 x 2.4 km, yet contains ponds with salinities that cover this entire spectrum.
206 To examine the impact of wet conditions on pond salinity, EC measurements from 14 ponds at
207 St. Denis were taken from 2009 – 2012, shown in **Figure 3**. A longer record from only one pond
208 is explored in more detail below. There is an almost completely consistent pattern in the
209 response, with fresh water ponds becoming salinized over the wet period from 2010 onwards,
210 brackish-saline ponds becoming diluted, and moderately-brackish ponds having relatively stable
211 EC values. The water flushed into the ponds may have a varying salinity over the landscape in
212 the different ponds, but these data would suggest that the salinity of this water is of a similar
213 order to the moderately-brackish ponds. Therefore note that while the brackish ponds are diluted,
214 there may still be a considerable mass input of salt into the ponds. Heagle et al. (2013) showed
215 this for pond 50 at St. Denis (a brackish-saline pond), where from 2009 to 2011 the mass of SO_4
216 in the pond water increased by more than 50% whilst the EC reduced by ~20%. Overall, the data
217 in **Figure 3** suggest that all ponds in the landscape were enriched in salt mass under wet
218 conditions, assuming that the pond volumes increased over the wet conditions period, as
219 indicated by **Figure 2B**.

220 Ponds 60 and 117 are fresh-water ponds that maintained a relatively constant EC over 2009 -
221 2012, in contrast to all of the other fresh water ponds where an increase in EC was observed.
222 Ponds 60 and 117 are the highest and most isolated ponds in the site, and unlike all the other
223 fresh water ponds, they do not have elevated ponds adjacent to them, from which they may
224 receive salts, by surface or subsurface pathways.

225 **3.3 Changes in water level in a terminal recharge pond**

226 Pond 109 at St Denis is a terminal recharge pond (i.e. it collects surface water but does not spill,
227 and loses water and salt to groundwater), which is well-studied, with a good archived dataset.
228 **Figure 4** presents pond 109 water depths from 1969 to 2012. A depth of 1.27 m (70% of the
229 deepest recorded pond depth, shown in **Figure 4**) was used to indicate unusually wet conditions.
230 Consequently, and in good agreement with the precipitation data and the regional pond depths
231 (**Figure 2**), wet conditions in pond 109 persist in 1997, 2007 and 2010-2012. A distinct
232 difference between the wet conditions associated with snowy winters (1997, 2007) and rainy
233 summers (2010, 2012) is observed. In wet years associated with snowy winters, as in non-wet
234 years, the pond maximal depths occurred in spring time, following the snowmelt. For wet years
235 associated with rainy summers, the pond water levels continually rose up from spring to late
236 summer. In 2010, pond levels rose from June to September. The year 2012 also had high water
237 levels throughout the entire summer from March to October, with positive increases in the pond
238 level from March to July. For 2011, as indicated in **Table 1**, the high water levels were due to the
239 high antecedent water storage from 2010, with the (unexceptional) snowmelt event superimposed
240 on this to produce what were then unprecedented water levels. This unprecedented pond level
241 was matched and slightly exceeded in 2012, this time due to high summer rainfall.

242 **3.4 Changes in salinity in a terminal recharge pond**

243 EC measurements for pond 109 have been taken since 1993, excluding the drought years (1999-
244 2004) when the pond dried out completely, and 1995 and 2006 due to logistical problems.
245 **Figure 5** presents pond 109 estimated pond water volume, based on the depth-volume
246 relationship in **Equation 1 (A)**; measured pond water EC and estimated salt mass based on
247 **Equation 2 (B)**; and changes in pond volume, EC, and salt mass over selected summer periods
248 (C).

249 It can be seen that until 2010 the pond EC was of the order of 100-500 $\mu\text{S}/\text{cm}$, meaning the pond
250 would be classified as fresh. The salt mass in the pond during this period was of the order of 500
251 kg. Average conditions are exemplified by 1994 in **Figure 5**. Here, the pond water volume
252 increases in spring time following the snowmelt runoff from the adjacent uplands which causes a
253 strong dilution. Over the summer months, the water levels drop due to a combination of
254 infiltration and evaporation, and the salinity increases, due to evaporation. The mass of salt in the
255 ponds steadily decreases, due to infiltration, and salts accumulate in the soils, especially in the
256 saline ring around the pond (Hayashi et al., 1998b; Heagle et al, 2007; Nachshon et al., 2013).
257 This is the steady salt cycle that operates continuously in the closed drainage basins of the
258 prairies. However, to complete the cycle, salts must re-enter the pond at some point. The re-entry
259 mechanism is harder to observe, but is thought to be associated with the flushing of near surface
260 salts from the soil by the snowmelt via surface or shallow subsurface pathways (Hayashi et al.,
261 1998b). It is also possible that diffusion might play a role in returning salts to the ponds from the
262 soils.

263 During the two wet years associated with exceptional snowmelt (1997 and 2007) the pond
264 salinity was unexceptional, and pond salt mass was only marginally elevated. This is consistent
265 with the snowmelt re-entry mechanism, but shows that this effect has a negligible and short-lived
266 impact on the salt cycle.

267 The first wet year associated with extreme summer rainfall (which fell in that year) was 2010,
268 and here a dramatic increase in pond salinity (EC rose from 261 to 801 $\mu\text{S}/\text{cm}$) and salt mass
269 (M_{salt} rose from 260 to 1350 kg) was initiated, making the pond moderately brackish. In 2011,
270 water levels were even higher, but this was largely due to the antecedent water levels from the
271 previous year. After the melt event and through the summer, the water level dropped, whilst the
272 salt mass increased, perhaps suggesting that the pond was functioning as a flow-through pond
273 during this period (Nachshon et al., 2013), with a highly saline inflow at some point, and a more
274 dilute outflow somewhere else. This will be explored further below. The EC during this period
275 steadily rose from 527 to 846 $\mu\text{S}/\text{cm}$ due to evaporation. Finally in 2012, which had both high
276 antecedent water levels and salt mass, and extreme high summer rainfall, the salt mass and
277 salinity continued to rise rapidly to unprecedented levels (EC up to 1061 $\mu\text{S}/\text{cm}$, salt mass
278 peaking at 3800 kg).

Deleted: 2010 was

Deleted: t

Comment [MJ2]: Whilst or and?

281 3.5 Changes in water chemistry in a terminal recharge pond

282 The major ion analysis of pond 109 water over the years of 2007-2009 and 2012 reveals a
283 significant change in the cations composition of the pond water. In 2007-2009 the major cations
284 in the pond water were Ca, Mg, and Na with molar fractions of approximately 60%, 35%, and
285 5%, respectively (**Figure 6**). In 2012, at the end of the measured wet period, the pond water
286 cation composition was of 35%, 50%, and 15% for Ca, Mg, and Na, respectively, indicating
287 enrichment of the pond water with Mg and Na. This enrichment is likely due to dissolution and
288 migration of mirabilite ($\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$), bloedite ($\text{Na}_2\text{Mg}[\text{SO}_4]_2 \cdot 4\text{H}_2\text{O}$), and epsomite
289 ($\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$) which are the more soluble salts that persists in the prairie tills. Under normal
290 conditions, these salts accumulate at the more distant parts of the uplands, away from the ponds,
291 according to the Hardie-Eugster model (Miller et al., 1989; Miller and Brierley, 2011; Skarie et
292 al., 1987; Timpson et al., 1986; Nachshon et al., 2013). The fact that under high summer rainfall
293 these cations migrate into the pond indicates subsurface water flows from high and distant parts
294 of the uplands that flush these salts from the subsurface into the pond.

Comment [MJ3]: Is Fig. 6 a Piper diagram?

295 3.6 Groundwater-surface water interactions

296 The measurements obtained by the piezometers (**Figure 1**) indicate the direction of subsurface
297 exchanges between Pond 109 and the upland to the north of the pond. **Figure 7** contrasts
298 differences in the magnitude and direction of this gradient with changes in the mass of salt (M_{salt})
299 in the pond water, for 2008 – 2012. It can be seen that most of the time the hydraulic head
300 gradient is from the pond to the uplands, indicating flow out of the pond. However, on several
301 occasions, most prominently in the summer of 2010, the hydraulic head gradients were reversed,
302 indicating fluxes from the uplands back into the pond, with the potential of transporting
303 dissolved salts from the uplands into the pond. For more than 85% of the data points, these data
304 behave consistently – that is an inflow to the pond is associated with an increasing salt mass and
305 an outflow from the pond is associated with a reducing salt mass. However, for ~15% of the data
306 points, a positive buildup of salt mass coincided with apparent flow out of the pond. These
307 anomalous measurements (red symbols in **Figure 7**) were observed mainly in the summer of
308 2012. These points can be explained either as slow mixing within the pond, or, again, as
309 evidence of non-uniform inflows and outflows to/from the pond.

310 3.7 Salinity of the riparian zone

311 Mini observation wells combined with EM38 readings provide estimates of EC around pond
312 108A (the end of the transect discussed below) and pond 109 (the terminal recharge pond
313 discussed above) (**Figure 8**). Around pond 109 average EC was 3342 $\mu\text{S}/\text{cm}$ with a maximum
314 EC of 5500 $\mu\text{S}/\text{cm}$ in the eastern side of the pond. The soil was 3 to 5 times more saline than the
315 pond water, which was 1020 $\mu\text{S}/\text{cm}$ at the time of measurement. For pond 108A average EC at
316 pore water around the pond was equal to 3200 $\mu\text{S}/\text{cm}$ with maximum EC of 4000 $\mu\text{S}/\text{cm}$ in the
317 southern side of the pond. These values are ~ 1.5 times more saline than the pond water (equal to
318 2300 $\mu\text{S}/\text{cm}$). It follows that exchanges of water between the pond and soils/groundwater are far
319 more efficient at advecting salts into the pond than out. For example, for pond 109, in order to
320 remove the salts added to the pond for every unit of inflow (i.e. exfiltration of groundwater), 3 –
321 5 units of outflow (i.e. infiltration of pond water) would be required. It is also evident from
322 **Figure 8** that the spatial distribution of salt in the saline ring is non-uniform with distinctive
323 regions of high and low salt concentrations. We speculate that this may be related to the slope
324 steepness and the spatial distribution of the elevated - adjacent ponds. However, this hypothesis
325 and the associated processes should be explored in future studies.

Comment [MJ4]: Or "...at transporting salts by advection into the pond than out."

Comment [MJ5]: The logic is hard to follow in this sentence?

326 3.8 Response to summer rainfall along a transect

327 **Figure 9** shows the water levels and EC data observed along a transect between ponds 107
328 (shown on the left in **Figure 9 B & C**, where $x = 0$ m) and 108A (shown on the right in **Figure 9**
329 **B & C**, where $x = 35$ m) for a 20 day period in July 2012, which included a large rainfall event
330 on July 15. For the first two observation times (5 and 6 July, row 1 and 2 in **Figure 9**),
331 conditions were relatively dry. The water table pattern reflects the general conceptual model for
332 the prairies (Fig 6 in van der Kamp and Hayashi, 2009), whereby both ponds are close to
333 equilibrium with the adjacent groundwater, which drops lower beneath the uplands. With respect
334 to salinity; under these conditions the highest EC values were measured in the pore water close
335 to pond 108A, which is the local saline ring of this pond. On July 11 (row 3 in **Figure 9**), a
336 minor rainfall event resulted in a moderate increase in the water table, which also allowed for
337 more points to be sampled for EC. The only significant difference in EC was measured at $x=31$
338 m (~ 5 m from pond 108A), where EC readings rose from ~ 2000 $\mu\text{S}/\text{cm}$ to well over 6000 $\mu\text{S}/\text{cm}$.
339 This may indicate leaching of salts from upper parts of the unsaturated profile by the infiltrating

340 rain water, but the affect appears highly localized above the saline ring of Pond 108A. The next
341 set of measurements on July 19 (row 4 in **Figure 9**), followed a large rain event on July 15 and
342 16. The water table came to ground surface along much of the transect, and in these parts of the
343 transect seepage of the water above ground surface was visibly observed. We can be certain that
344 this was a saturation excess mode of runoff, and moreover no fill-and-spill of pond 107 into pond
345 108A was observed. Since the piezometers were screened along their entire length, it is not
346 possible to determine whether this was local perching, or groundwater recharge bringing the
347 water table up to the surface, but due to the high infiltration capacity of the fractured till, the
348 latter is believed to be more plausible. During this period the EC was reduced across the transect,
349 due to dilution, except on the right hand edge, closest to Pond 108A. This edge would have
350 received much of the salts being flushed laterally from the transect, as the head gradients show,
351 and hence here the EC rose up from ~2620 $\mu\text{S}/\text{cm}$ to 3310 $\mu\text{S}/\text{cm}$. For the final set of
352 measurements on July 24 (row 5 in **Figure 9**), taken after five days with no rain, the water table
353 depth declined along the entire transect, but not quite to depths as low as the antecedent levels
354 (row 1 and 2 in **Figure 9**). Hydraulic gradients from the earlier observations showed two ponds
355 that were disconnected from one another, whereas the later observations show a more-or-less
356 continuous hydraulic gradient from Pond 107 to Pond 108A, suggesting the ponds were then
357 connected, albeit perhaps temporarily. As the pulse of rain water left the profile, the EC profile
358 returned to levels very similar to the antecedent conditions. Adjacent to Pond 108A the salinity
359 dropped again, probably in response to mixing with other fresher water in the pond.

Comment [MJ6]: Do you mean observation well?

Comment [MJ7]: If pipe is screened along entire length, shouldn't this be an observation well? Piezometers are generally screened along shorter intervals.

Deleted: reduced

360 **4. Conclusions**

361 Field measurements collected over the past 20 years from St. Denis in central Saskatchewan shed
362 new light on salt dynamics of prairie wetlands. It was shown that under wet conditions associated
363 with rainy summers, large fluxes of salts from the subsurface are flushed into the ponds. The
364 corresponding change in pond salinity (i.e. salt concentration) depends on the antecedent salinity,
365 the volume of water flushed into the ponds and the salinity of the water flushed into the pond.
366 However, the general tendency is quite uniform, with fresh-water ponds becoming more saline,
367 moderately brackish ponds remaining largely unchanged, and brackish ponds becoming diluted.
368 Most of the time, however, summer rainfall is small compared with potential evaporation, and
369 thus the dominant (or only) source of infiltration is depression-focused recharge of snowmelt

371 beneath the ponds. Significantly, it was seen that wet conditions associated with this kind of
372 infiltration, i.e. unusually snowy winters, had a negligible impact on the salt dynamics and pond
373 salinization. Moreover, the rain-associated wet conditions lasted for much longer durations
374 (months) compare to the snow-associated wet conditions (weeks), therefore, increasing the
375 potential impact of the unusual rain conditions on the subsurface salt transport processes.

376 The relatively high permeability of the top few meters of the prairies till results in a quick
377 response in the upland subsurface storage that leads to relatively quick transport of salts from the
378 upland subsurface to the ponds, on time scales of the order of few days only. Even quicker
379 responses in water flows and pond levels may occur by surface fill and spill events, but usually
380 the surface waters are less saline than the pore water, thus salt transport is more limited under
381 these conditions.

382 It is hard to predict what would be the climatic conditions in the near and far future, but recent
383 years' evidences, as well as climatic models indicate that the variability of the climate is
384 expected to increase, with a high chance of extreme rain periods, as well as extreme droughts
385 (Frelich and Reich, 2010; Semenov and Stratonovitch, 2010). If a series of extremely wet years,
386 where salt is being accumulated in the ponds, is followed by a severe drought, the high
387 evaporation rates of the drought period will result in further increase in the pond salinity and
388 eventually, all of the salts within the pond water will be precipitated, following the complete
389 drying of the pond. This scenario may result in salt concentrations in the upper soil horizons of
390 the wetlands at concentrations that were never measured before. Since both the wildlife of the
391 prairies, as well as agricultural activities, depend on the wetlands' physical and chemical
392 conditions it is critical to improve our understanding of the geo-chemical processes and to be
393 prepared to cope with salinization processes in the prairies.

394 **References:**

395 Beaulieu, C. and V. Wittrock: Climate reference station Saskatoon annual summary 2012.
396 Saskatchewan Research Council Publication No. 10440 - 1E13.

397 Berthold, S., Bentley, L.R. and Hayashi, M.: Integrated hydrogeological and geophysical study
398 of depression-focused groundwater recharge in the Canadian prairies, *Water Resour. Res.*, 40,
399 W06505, doi:10.1029/2003WR002982, 2004.

400 Conly, F.M., Su, M., van der Kamp, G. and Millar, J. B.: A practical approach to monitoring
401 water levels in prairie wetlands. *Wetlands* 24, 1, 219-226, 2004.

402 Dehaan, R. and Taylor, G.R.: Field-derived spectra of salinized soils and vegetation as indicators
403 of irrigation-induced soil salinization, *Rem. Sens. Environ.* 80, 406–417, 2002.

404 Euliss, N. H., LaBaugh, J.W., Fredrickson, L.H., Mushet, D.M., Laubhan, M.K., Swanson,
405 G.A., Winter, T.C., Rosenberry, D.O. and Nelson, R.D.: The wetland continuum: a conceptual
406 framework for interpreting biological studies, *Wetlands* 24, 448–458, 2004.

407 Frelich, L. E. and Reich, P. B.: Will environmental changes reinforce the impact of global
408 warming on the prairie-forest border of central North America?, *Front. Ecol. Environ.*, 8, 7, 371-
409 378, 2009.

410 Hayashi, M. van der Kamp, G. and Rudolph, D.L.: Mass transfer processes between a prairie
411 pothole and adjacent uplands, 1: water balance, *J. Hydrol.*, 207, 42–55, 1998a.

412 Hayashi, M., van der Kamp, G. and Rudolph, D.L.: Mass transfer processes between a prairie
413 pothole and adjacent uplands, 2: chloride cycle, *J. Hydrol.*, 207, 56–67, 1998b.

414 Hayashi, M. and van der Kamp, G.: Simple equations to represent the volume–area–depth
415 relations of shallow wetlands in small topographic depressions, *J. Hydrol.*, 237, 1-2, 74–85,
416 doi:10.1016/S0022-1694(00)00300-0, 2000.

417 Heagle, D.J., Hayashi, M. and van der Kamp, G.: Use of solute mass balance to quantify
418 geochemical processes in a prairie recharge wetland, *Wetlands*, 27, 806–818, 2007

419 Heagle, D.J., Hayashi, M. and van der Kamp, G.: Surface-subsurface salinity distribution and
420 exchange in a closed-basin prairie wetland, *J. Hydrol.*, 478, 1-14, 2013.

421 Keller, C.K., van der Kamp, G. and Cherry, J.A.: Hydrogeochemistry of a Clayey Till: 1. Spatial
422 Variability, *Water Resour. Res.*, 27, 10, 2543–2554, 1991.

423 Mekis, É. and Vincent, L. A.: An overview of the second generation adjusted daily precipitation
424 dataset for trend analysis in Canada. *Atmosphere-Ocean*, 49, 2, 163-177, 2011.

425 Micklin, P.: The Aral sea disaster, *Annu. Rev. Earth Planet. Sci.*, 35, 47-72, 2007.

426 Millar, J. B.: Wetland classification in Western Canada, *Can. Wildl. Rep. Ser.* 37, 1, 1976.

427 Miller, J.J., Pawluk, S. and Beke, G.J.: Evaporite mineralogy, and soil solution and groundwater
428 chemistry of a saline seep from southern Alberta, *Can. J. Soil Sci.* 69, 273-286, 1989.

429 Miller, J.J. and Brierley, J.A.: Solonchic soils of Canada: Genesis, distribution, and
430 classification. *Can. J. Soil Sci.* 91, 889-902, 2011.

431 Nachshon, U., Ireson, A., van der Kamp, G. and Wheeler, H.: Sulfate salt dynamics in the
432 glaciated plains of North America, *J. Hydrol.*, 499, 188–199, 2013.

433 Rengasamy, P.: World salinization with emphasis on Australia, *J. Exp. Bot.*, 57.5, 1017-1023,
434 2006.

435 Rhoades, J. D., Sparks, D. L., Page, A. L., Helmke, P. A., Loeppert, R. H., Soltanpour, P. N. and
436 Sumner, M. E.: Salinity: electrical conductivity and total dissolved solids. *Methods of soil*
437 *analysis. Part 3-chemical methods.*, 417-435, 1996.

438 Semenov, M. A. and Stratonovitch, P.: Use of multi-model ensembles from global climate
439 models for assessment of climate change impacts, *Climate research*, 41, 1, 1-14, 2010.

440 Shaw, D.A., van der Kamp, G., Conly, M., Pietroniro, A. and Lawrence, M.: The fill-spill
441 hydrology of prairie wetland complexes during drought and deluge, *Hydrol. Process.*, 26, 20,
442 3147–3156, 2012.

443 Shook, K.R. and Pomeroy, J.W.: Memory effects of depressional storage in Northern Prairie
444 hydrology. *Hydrol. Process.* 25, 3890–3898, 2011.

445 Skarie, R.L., Richardson, J.L., McCarthy, G.J. and Maianu, A.: Evaporite mineralogy and
446 groundwater chemistry associated with saline soils in eastern North Dakota, *Soil Sci. Soc. Am. J.*
447 51, 137-1377, 1987.

448 Spence, C. and Woo, M.K.: Hydrology of subarctic Canadian Shield: Soil-filled valleys, *J.*
449 *Hydrol.*, 279, 151–166, 2003.

450 Stewart, R. E. and Kantrud, H. A.: Vegetation of prairie potholes, North Dakota, in relation to
451 quality of water and other environmental factors. US Government Printing Office, 1972.

452 Timpson, M. E., Richardson, J. L., Keller, L. P. and McCarthy, G.J.: Evaporite mineralogy
453 associated with saline seeps in southwestern North Dakota, *Soil Sci. Soc. Am. J.*, 50, 490-493,
454 1986.

455 Tyler, S. W., Munoz, J. F., and Wood, W. W: The response of playa and sabkha hydraulics and
456 mineralogy to climate forcing, *Ground water*, 44, 3, 329-338, 2006.

457 van der Kamp, G. and Hayashi, M.: Groundwater-wetland ecosystem interaction in the semiarid
458 glaciated plains of North America, *Hydrogeol. J.*, 17, 203–214, 2009.

459 Wheater, H. S., Jakeman, A. J. and Beven, K. J.: Progress and directions in rainfall-runoff
460 modeling, in: *Modeling change in environmental systems*, John Wiley & sons, 101-132, 1993.

461 Wienhold, C. E. and van der Valk, A. G.: The impact of duration of drainage on the seed banks
462 of northern prairie wetlands, *Can. J. Bot.*, 67, 6, 1878-1884, 1989.

463 Winter, T.C.: Hydrologic studies of potholes in the northern prairies, in: *Northern prairie*
464 *wetlands*, Iowa State University Press, 17–54, 1989.

465

466 **Table 1: Factors influencing the pond levels during wet conditions.**

Year	High pond level in previous year	High summer rainfall in previous year	High winter snowpack	High summer rainfall in current year
2006		X	X	X
2007	X	X	X	
2010				X
2011	X	X		
2012	X			X

467

468

469 **Figure captions:**

470 **Figure 1: (A) Regional location of St. Denis; (B) St. Denis site; (C) Enlargement of pond 109**
471 **area. In (C) the dots between pond 107 and 108A indicate the location where salinity and**
472 **ground water levels were measured along the transect. Stars represent piezometers; thick**
473 **solid line around pond 109 indicates the water level of the pond in July 2012; and the**
474 **dashed line at pond 109 indicates the average location of pond 109 water level. White**
475 **numbers indicate selected ponds numbers.**

476 **Figure 2: (A) total summer rainfall plus winter snow per hydrologic year.(B) Averaged and**
477 **normalized depths of 12 ponds at St. Denis.**

478 **Figure 3: Averaged EC of selected ponds in St. Denis from 2009 to 2012. Solid, dot, and**
479 **dashed lines indicate on brackish-saline, moderately brackish, and fresh water wetlands,**
480 **respectively.**

481 **Figure 4: Pond 109 water depths from 1969 to 2012.**

482 **Figure 5: (A) Pond 109 volume; (B) measured EC and calculated M_{salt} ; and (C) summer-**
483 **time changes in volume, EC and M_{salt} of the pond. Separated (lower) figures in (A) and (B)**
484 **are zoom in for selected years with various climatic conditions.**

485 **Figure 6: Major cations molar fraction in pond 109 for the years of 2007-2009 and 2012.**
486 **Marker size is proportional to the pond EC.**


487 **Figure 7: (A) Water level differences between upland and pond 109 and M_{salt} . Red symbols**
488 **indicate on the anomalous measuring points, where a disparity was observed between the**
489 **direction of the subsurface water flows and the changes in M_{salt} .**

490 **Figure 8: Salinity around ponds 109 and 108A.**


491 **Figure 9: Measured precipitation (A); groundwater table depth (B); and EC (C) along the**
492 **107-108A transect. The dashed lines in (A) indicate on the day at which measurements**
493 **were taken for (A), (B), and (C) in the same row. In (B) the black line is ground surface and**
494 **the blue curve is the location of the water table. Arrows indicate on flows directions. In (C)**
495 **blue lines indicate on the measured EC along the transect. Red curves indicate on**
496 **measured EC at previous measurement, to emphasize the change of EC with time.**

497


498 **Figures:**




499 **Figure 1: (A) Regional location of St. Denis; (B) St. Denis site; (C) Enlargement of pond 109**
500 **area. In (C) the dots between pond 107 and 108A indicate the location where salinity and**
501 **ground water levels were measured along the transect. Stars represent piezometers; thick**
502 **solid line around pond 109 indicates the water level of the pond in July 2012; and the**
503 **dashed line at pond 109 indicates the average location of pond 109 water level. White**
504 **numbers indicate selected ponds numbers.**
505




506
 507 **Figure 2: (A) total summer rainfall plus winter snow per hydrologic year.(B) Averaged and**
 508 **normalized depths of 12 ponds at St. Denis.**
 509



510
 511 **Figure 3: Averaged EC of selected ponds in St. Denis from 2009 to 2012. Solid, dot, and**
 512 **513 dashed lines indicate on brackish-saline, moderately brackish, and fresh water wetlands,**
 514 **respectively.**




515
516 **Figure 4: Pond 109 water depths from 1969 to 2012.**
517




518
 519 **Figure 5: (A) Pond 109 volume; (B) measured EC and calculated M_{salt} ; and (C) summer-**
 520 **time changes in volume, EC and M_{salt} of the pond. Separated (lower) figures in (A) and (B)**
 521 **are zoom in for selected years with various climatic conditions.**


Comment [MJ8]: Need to define NC and WC?




522
 523 **Figure 6: Major cations molar fraction in pond 109 for the years of 2007-2009 and 2012.**
 524 **Marker size is proportional to the pond EC.**
 525



526
 527 **Figure 7: (A) Water level differences between upland and pond 109 and M_{salt} . Red symbols**
 528 **indicate on the anomalous measuring points, where a disparity was observed between the**
 529 **direction of the subsurface water flows and the changes in M_{salt} .**



530
531 **Figure 8: Salinity around ponds 109 and 108A.**



532
 533 **Figure 9: Measured precipitation (A); groundwater table depth (B); and EC (C) along the**
 534 **107-108A transect. The dashed lines in (A) indicate on the day at which measurements**
 535 **were taken for (A), (B), and (C) in the same row. In (B) the black line is ground surface and**
 536 **the blue curve is the location of the water table. Arrows indicate groundwater flow**
 537 **directions. In (C) blue lines indicate the measured EC along the transect. Red curves are**
 538 **measured EC at previous measurement, to emphasize the change of EC with time.**
 539

Deleted: on
 Deleted: s
 Deleted: on
 Deleted: indicate
 Deleted: on