~~A Scaling Approach,~~ Predicting the ~~Continuous Form of~~ Soil Moisture ~~Characteristics~~Retention Curve, from Soil Particle Size Distribution and Bulk Density Data using a Packing Density Scaling Factor

F. Meskini-Vishkaee[1], M. H. Mohammadi[1,2] and M. Vanclooster[2]

[1] {Department of Soil Science, Faculty of Agriculture, University of Zanjan post code 313, Zanjan, (Iran)}

[2] {Earth and Life Institute, Environmental Sciences, Univ. Catholique de Louvain, Croix du Sud 2, Bte 2, 1348 Louvain-la-Neuve, Belgium}

Correspondence to:F. Meskini-Vishkaee (fatemeh.meskini@yahoo.com)

ABSTRACT

A substantial number of models, predicting the Soil Moisture Characteristic Curve (SMC) from Particle Size Distribution (PSD) data, underestimate the dry range of the SMC especially in soils with high clay and organic matter contents. In this study, we applied a continuous form of the PSD model to predict the SMC and subsequently, we developed a physically based scaling approach to reduce the model's bias at the dry range of the SMC. The soil particle~~s~~ packing ~~parameter~~density~~, obtained from the porosity~~ was considered as a metric of soil structure and used to ~~asacharacteristic length~~define a soil particle packing scaling factor. This factor was subsequently integrated in the conceptual SMC prediction model. The model was tested ~~by using~~ on eighty-two soils~~ samples~~, selected from the UNSODA database. The results~~showed~~ show that the scaling approach properly estimates the SMC for all soil samples. In comparison to ~~i) the formerly used physically based~~ the original conceptual SMC model without scaling, the scaling ~~proposed~~ approach ~~improved~~ improves the model estimations ~~by an~~on average ~~of~~ by30%~~ for all soil samples~~. ~~However, the advantage of this new approach was larger~~Improvements were particularly significant for the fine and medium textured soils~~than that for the coarse textured soil~~. ~~In view that in~~ Since ~~this~~ the scalingapproach is parsimonious and ~~there is~~do not rely on ~~no further need for~~ additional

empirical parameters, we conclude that this approach may be used for ~~could become applicable for~~ estimating SMC at the larger field scale from basic soil data.

1 Introduction

Increasing contamination of the groundwater resources, have profoundly accentuated the need for accurate predictions of subsurface flow and chemical transport.Water flow and subsequent chemical transport are largely determined by the soil hydraulic properties, such as the Soil Moisture Characteristics curve (SMC) (Wang et al, 2002;Mohammadi et al., 2009). Measuring the soil hydraulic properties is still difficult, labor intensive, and expensive. Therefore, many researchers havemade an attempt to develop an indirect method as an alternative to the direct measurement of hydraulic properties. For the SMC, indirect methods are classified into conceptual methods(Nimmo et al., 2007; Mohammadi and Vanclooster, 2011),semi-physical methods(e.g. Arya and Paris, 1981; Haverkamp and Parlange, 1982; Wu et al. 1990; Smetten and Gregory, 1996) and empirical methods (e.g. Saxton et al. 1986; Schaap et al. 1998).

The semi-physical methods are mainly based on shape similarity between the SMC and the Particle Size Distribution (PSD) curve (Zhung et al., 2001; Schaap, 2005; Haverkamp et al., 2005; Hwang and Choi, 2006), implying that the pore size distribution (PoSD) is closely related to the PSD (Arya et al., 2008). Arya and Paris (1981) did a pioneering work (AP model) for the development of ~~ing~~ semi-physical models. They showed that~~,~~ the pore size, which is associated with a pore volume, is determined by scaling the pore length, using a scaling factor, α. They demonstrated that~~,~~ an average value of 1.38 for α, scales the pore lengths based on spherical particles to natural pore lengths properly. However, later investigations by Arya et al. (1982), Tyler and Wheatcraft (1989), Basile and D'Urso (1997) and Vaz et al. (2005) revealed that α value varies between 1.02 - 2.97 for fine and coarse textured soils, respectively. A slight error in the estimation of α may result in considerable error in predicting the SMC (Schuh et al. 1988). Schuh et al. (1988) found that the value of α varies with soil texture and suction head, especially in the wet range of sandy soils. Using three formulations of α, Arya et al. (1999) modeledthe parameter α as a function of particle sizes and showed that α was not constant. ~~and therefore i~~It decreased with increasing particle size, especially for the coarse fractions. Tyler and

Wheatcraft(1989) showed that the parameter α is equivalent to the fractal dimension of a tortuous fractal pore.

Although, the empirical methods have been developed extensively (e.g. Puhlmann and von Wilpert, 2012), the performance of an empirical method will depend on the databases, being used for the model calibration and testing (Tietje and Tapkenhinrichs, 1993; Kern, 1995; Schaap and Leij, 1998; Schaap et al., 2004; Haverkamp et al., 2005; Hwang and Choi, 2006, Weynants et al., 2009). Moreover, direct measurementsof SMC are often integrated as predictor variables of the continuous SMC function. Many attempts have ~~so far~~ been made to reduce the sensitivity of the indirect methods to empirical and database-dependent parameters. For instance, Mohammadi and Vanclooster (2011), proposed a conceptual robust model (MV) that~~,~~ does not include~~d~~ an empirical parameter and is independent of the databases that are being used. The disadvantages of ~~the most~~ semi-physical or conceptual models such as the AP and MV models are~~,~~ the use of "bundle of cylindrical capillaries" (BCC) concept to represent the pore space geometry and the lack of consideration of surface forces (Or and Tuller, 1999; Tuller et al., 1999; Mohammadi and Meskini-Vishkaee, 2012). These conceptual problems often ~~which~~ lead~~s~~ to the underestimation of the dry range of the SMC (Arya et al., 1999; Hwang and Choi, 2006; Mohammadi and Vanclooster, 2011). Such underestimations would result in large ~~error in~~ modeling errors of hydraulic dependent soil functions such ~~of~~ as mechanical ~~properties of unsaturated soil~~ resistance functions (Gras et al., 2010), ~~and biological processes including~~ plant water uptake functions (Ryel et al., 2002) ~~and~~ microbial activity functions (Jamieson et al., 2002; Santamarı´a and Toranzos, 2003), in particular in ~~especially in~~ arid environments.

To predict continuous SMC, Naveed et al. (2012) parameterized the van Genuchten model based on the SMC data points predicted ~~by their proposed model using~~ from organic matter, clay, silt, ~~and~~ fine sand and coarse ~~sands~~sand content. Mohammadi and Meskini-Vishkaee (2013) integrated the MV model with the van Genuchten (VG) model (van Genuchten, 1980) to predict the continuous SMC curve (MV-VG model) from PSD data. They ~~and~~ found that ignoring the residual moisture content (θ_r) is the main source of systematic error ~~resulted fromthe~~in the MV model. They further tested and compared four approaches to predict the θ_r, and showed that the incorporation of ~~the safely estimated value of the~~ predicted θ_r, will improve the MV-VG prediction results considerably. However, the estimation of θ_r has some limitations, due to the

lack of a conceptual underpinning and the poor predictability of θ_r (Leij et al., 2002). Tuller and Or (2005) suggested that, the introduction of θ_r as a fitting parameter in most SMC models often makes the physical representation of key processes in the dry soils vague. Moreover, they pointed out that, the dry range of the SMC shows remarkable scaling behavior. ~~In recent years, the scaling theory has been widely used as an effective tool to describe the variation of the soil hydraulic properties (Sharma and Luxmoore, 1979; Ahuja et al., 1984; Eching et al., 1994; Kosugi and Hopmans, 1998; Oliveira et al., 2006; Nasta et al., 2009). The concept of geometric similitude and similar media was used to develop scaling theory in soil physics (Miller and Miller, 1956). Scaling methods provide a means to relate hydraulic properties of different soils to those of a reference soil, using scaling factors (Nasta et al., 2009). On the other hand, some attempts have been made to modify the original scaling method (Warrick et al., 1977; Vogel et al., 1991; Deurer and Duijnisveld, 2000; Das et al., 2005).~~ Arya et al. (2008) developed a procedure to scale natural pore lengths, directly from straight pore lengths. They ~~exhibited~~ showed that the~~,~~ scaling approach is less sensitive to uncertainties in model parameters and provides better predictions of the SMC, compared with the original AP model.

Kosugi (1996) showed that the SMC can be expressed by a lognormal pore-size distribution function, while Kosugi and Hopmans (1998) found that the set of scaling factors is lognormally distributed when PoSD curve is lognormal. Hayashi et al. (2007) used the Kosugi model (Kosugi, 1996) to evaluate the effectiveness of three kinds of scaling factors obtained by the microscopic characteristic length, standard deviation of pore-size distribution and the porosity. They indicated that in the natural forested hillslope soils, the variability in the SMC is characterized by variability in the effective soil pore volume. Nasta et al. (2009) concluded that the scaling of the PSD curves provides for adequate characterization of the mean and variance of SMCs, which allows for characterization of the soil spatial variability.

Many researchers developed empirical models for expressing the SMC~~;~~ since the parameters of these models do not address the physical significance of the medium. Hence the spatial variability in the pore structure of soils is not fully understood (Hayashi et al., 2007). Likewise, the conventional scaling approaches are based on empirical curve fitting, without considering the physical meaning of the scaling factor (Perfect, 2005; Milla´n and Gonzalez-Posada, 2005). To apply these models, one needs to determine the scaling factor, where the complexity of

measurements of the pore–size and pore–volume distributions easily nullifies the estimation of the scaling factors. Nevertheless, some efforts have been made to relate the scaling factor to the soil texture (Tuli et al., 2001; Milla´n et al., 2003).

From this brief review~~,~~ we conclude that~~ although~~ the scaling approaches improve~~s~~the modeling and prediction of the SMC. Yet, most scaling approaches imply empirical parameters and a robust fully conceptual approach for the estimation of the SMC from easily measurable properties is still lacking~~s~~.

The MV model~~,~~ underestimates~~d~~ the moisture content in the dry range of the SMC, because of the simplified conceptualization of the pore geometr~~ic concepts~~y. ~~, i~~In particular the packing parameter ~~which~~ does not effectively reflect the pore geometry. The general aim of ~~our current work on th~~this study ~~is subject is to augment and~~is to improve the accuracy of the model proposed by Mohammadi and Meskini-Vishkaee (2013) using a scaling approach.

Therefore the objectives of this study ~~were~~ are (i) to formulate a robust and physically-based model to scale the SMC from the PSD and porosity, and (ii) to compare the model performance with the results from the existing MV-VG model, using soils documented in the UNSODA database (Nemes et al., 2000). We also evaluate the overall model performance with the results from the full empirically SMC prediction software, ROSETTA (Schaap et al., 2001).

2 Theory

Because of the close similarity between the shapes of the PSD and SMC curves, many researchers expressed aSMC model in terms of a PSD model (Haverkamp and Parlange, 1982; Fredlund et al., 2000; Zhuang et al., 2001). The SMC model developed by van Genuchten (1980) is very flexible, widely used and given by:

$$S_e = \left[\frac{1}{1+(\alpha h)^n} \right]^m \qquad (1)$$

$$S_e = \frac{\theta - \theta_r}{\theta_s - \theta_r} \qquad (2)$$

where θ ($L^3 L^{-3}$) is the soil moisture content, S_e (-) is effective saturation degree and θ_s ($L^3 L^{-3}$) and θ_r ($L^3 L^{-3}$) are saturated and residual soil moisture contents, respectively. The parameters n, m and α (L^{-1}) are fitting coefficients, and h (L) is the suction head.

The suction head, h_i(L), corresponding to the particle radius of the i-th fraction R_i (L) is given by (Mohammadi andVanclooster, 2011):

$$h_i = \frac{0.543\times10^{-4}}{R_i}\zeta \quad (3)$$

where ζ (-) is a coefficient depending on the state of soilparticles packing and is defined as:

$$\zeta = \frac{1.9099}{1+e} \quad (4)$$

where e (-) is the void ratio given by:

$$e = \frac{\rho_s - \rho_b}{\rho_s} \quad (5)$$

where the ρ_s(ML^{-3})and ρ_b (ML^{-3}) are soil particle and bulk densities respectively.

Arya and Paris (1981) suggested that the moisture content, θ_i (L^3L^{-3}), can be obtained from PSD and θ_s (L^3L^{-3}), as:

$$\theta_i = \theta_s \sum_{j=1}^{j=i} w_j; \qquad i = 1,2,3,...,k \qquad (6)$$

where $w_j$is the mass fraction of particles (-) in the j-th particle-size fraction. Considering that the

$$P_i = \sum_{j=1}^{j=i} w_j \quad (7)$$

would result in:

$$\theta_i / \theta_s = S \quad (8)$$

where S(-) is the saturation degree and P_i (-) is the cumulative mass fraction of soil particles. It is obvious, that if $\theta_r=0$ then, $S_e=S$ and subsequently $S=P_i$.Arya and Paris (1981) however, ignored the residual moisture content, while it may be a considerable value for many types of soil and clayey soils in particular. Combining Eq. (1) and (3) with Eq. (7) yields:

$$P_i = \left[\frac{1}{1+(\alpha\frac{0.543\times10^{-4}}{R_i}\zeta)^n}\right]^m \qquad (9)$$

In Eq. (9), the cumulative mass fraction, P_i, is substituted with the Se in Eq. (1). Hence, fitting of ~~the~~ Eq. (9) to the PSD data, enables one to directly predict the SMC parameters (n, m and α). Moreover, these coefficients allow expression of the continuous form of predicted SMC. Since,

assuming that θ_r=0 would result in model underestimation in dry range of the SMC (Mohammadi and Meskini-Vishkaee, 2013), we developed a conceptual scaling approach to reduce the model bias.

2.1 Scaling approach

~~Following the Miller and Miller (1956) scaling theory, we assume that, the geometrically identical soils are characterized bythe similarity of PoSD and PSD, but differ in their microscopic length scale, which is defined as follows (Nasta et al., 2009):~~

$$\cancel{\beta = \frac{\bar{\gamma}}{\gamma} \qquad (10)}$$

~~where β is a scaling factor, $\bar{\gamma}$ and γ represent the microscopic characteristic lengths of the reference soil and actual soil respectively. For instance, Kosugi and Hopmans (1998) proposed the median suction head, as the macroscopic characteristic length to scale SMC.~~

Following Hayashi et al. (2007), we suggest that the porosity is an appropriate property for inferring a characteristic ~~length~~scaling factor. Since, the soil porosity is linked to the packing parameter, ~~ζ~~ξ, in the MV model (Eq. (4)), we hypothesize that ~~ζ~~ξ is the characteristic ~~length~~ scale of the soil.

We assume that the reference soil is the one, which consists of uniform-size spherical particles that are arranged in random close packing state, leading to minimum porosity (known as the Kepler conjecture in literature of crystallography). Literature suggests that the porosity of this packing state is 0.259 (Hopkins and Stillinger, 2009). Subsequently, the maximum value of packing parameter, ζ_{max}, would equal to 1.41432 for reference soil. Hence the scaling factor, λ, for each soil sample can be ~~defined~~ suggested by:

$$\lambda = \frac{\zeta}{\zeta_{max}} \qquad (10\cancel{11})$$

In general, the values of the pore size distribution index, n, (Eqs. (1) and (9)) and ζ are large for the coarse-textured soils and small for the fine-textured soils. We suggest that the λ can scale the parameter 'n', obtained from fitting Eq. (9) to the PSD data, to the ~~index manifested by~~ 'n' parameter in the SMC model (Eq. (1)) (hereafter n^*) as follows:

$n^* = \lambda \times n$ (11~~12~~)

where n^* is scaled to the PoSD index in VG model. Hence, the modified model is

$$\frac{\theta}{\theta_s} = \left[\frac{1}{1+(\alpha h)^{n^*}}\right]^m$$

(12~~13~~)

In summary, given a known $\theta_s$we can calculate ζξ and subsequently λ using Eq. (10~~11~~). The soil parameter 'α' and 'm' are obtained from fitting Eq. (9) to the PSD data and, n^*, is estimated by Eq. (11~~12~~), and consequently the SMC is predicted directly by Eq. (12~~13~~).

3 Material and Methods

Eighty-two soil samples, with a wide range of physical properties that contained at least five PSD data, were selected from the UNSODA hydraulic properties database (Nemes et al., 2000).UNSODA is a database of basic soil and hydraulic properties from 790 samples, gathered from all over the world, and compiled by the U.S. Department of Agriculture. All soils are summarized in Table 1.

In this procedure, volumetric moisture contents corresponding to the i-th fraction were computed, using Eq. (6) and suction heads were predicted, using Eq. (3), in which the parameter ζ was calculated with Eq. (4). In this study, we assumed that, the porosity is equivalent to θ_s. For soils, that neither provide a porosity nor a θ_s, the first point of the SMC data that corresponds to the lowest suction head were used as θ_s (Chan and Govindaraju, 2004).

We fitted Eq. (9) to the PSD data. We used nonlinear regression analysis to fit Eq. (9) to the PSD, using Matlab7.1 software (Matlab 7.1, The Mathworks Inc., Natick, MA) and the Marquardt–Levenberg algorithm (Marquardt, 1963). We calculated for each soil the scaling factor, using either the bulk density or the available saturated soil moisture content, and predicted the SMC.

((Table 1))

For each prediction, the agreement between the predicted moisture content $\theta_{i(p)}$ and measured moisture content $\theta_{i(m)}$ was expressed in terms of the root mean square errors (RMSEs), given by:

$$RMSE = \sqrt{\frac{1}{N}\sum_{i=1}^{n}\left(\theta_{i(p)} - \theta_{i(m)}\right)} \quad (13\cancel{14})$$

in which N~~n~~ is the number of observed data points. The relative improvement (RI) resulted from the scaling approach rather than MV-VG model was calculated as follows (Minasny and McBartney, 2002):

$$RI\,(\%) = \frac{RMSE_M - RMSE_S}{RMSE_M} \times 100 \quad (14\cancel{15})$$

where, RI is the Relative Improvement, $RMSE_M$ and $RMSE_s$ are RMSE values of the MV-VG model and the current scaledmodel respectively. Obviously, ~~the~~ a negative RI value~~s~~ indicates that the scaling approach would diminish the accuracy level of the prediction of the SMC in comparison with the MV-VG model.

Wealso fitteda cubic polynomial function to the overall predicted data and,calculated the area between the fitted polynomial and the 1:1 line fromthe difference of the numerical integrals of these two functions (do Carmo, 1976).

Moreover, to consider and compare the reliability of the scale MV-VG model ~~scaling approach~~ with a fully-empirical SMC prediction ~~method~~model, we compared the estimations of scaled MV-VG model ~~recent approach~~ with the estimations of the ROSETTA software. In this software, we used the ~~(M~~SSCBD model option, i.e. we used ~~: SSCBD,~~textural percentages and bulk density as model predictors~~)~~ (Schaap et al., 2001).

4 Results and Discussion

~~To analyze the performance of the proposed scaling approach, the RMSE was used as an index, to compare models for each soil and textural class.~~ Table 2~~depicts~~ gives ~~this~~ the comparison between the MV-VG model, the ROSETTA model and the scaled MV-VG model in terms of RMSE, R^2 and RI. Table 2 ~~and~~ demonstrates the significant improved accuracy of the ~~presented~~ scaled MV-VG approach as compared to the original MV-VG model and ROSETTA. The RMSEs of the predicted and measured moisture contents, ranged from 0.0223 to 0.1502 for

the original MV-VG model (average 0.086), ~~and~~ from 0.0169 to 0.1122 (average 0.0601) for the scaled approach and from 0.0188 to 0.2453 (average 0.745) for the ROSETTA software. In terms of RMSEs, the scaled approach performed better than Schaap et al., 1998 and the Schaa~~p~~p and Leij (1998) models with similar predictor variables.The results showed that there is a significant difference between performance of scaled MV-VG approach and ROSETTA (p=0.05). Despite the pure statistical and empirical nature of the ROSETTA approach, it provides worse results than the approach based on the current scaling technique. The improvement of the scaled approach is also reflected by RI in Table 2. Except for soils no. 3033 (Clay loam) and 3090 (Silt loam), the scaling approach resulted in more accurate predictions for all soils. Table 2 also indicates that the scaling approach can improve the model estimations of the original MV-VG model by 30 %.

For the fine and medium textured soils, the values of RI are larger than for the coarse textured soil. This result was expected, because the MV and MV-VG models underestimate the dry range moisture content for the fine texture soils (Mohammadi and Vanclooster, 2011; Mohammadi and Meskini, 2013) and subsequently the scaling approach was more effective for these soils.

We examined the possible relations between the RI and soil physical properties. Among all parameters, the saturated moisture content and scaling factor show strong relations with the RI. Fig. 1a shows that the RI values increase significantly with the saturated moisture content of the soils, i.e. the scaling approach would more effectively improve the model accuracy for the fine texture soils with higher θ_s. This result can be confirmed with Fig. 1b, which ~~exhibits~~ shows that the scaling factor is inversely correlated with the IR factor (Fig. 1b). Indeed, the soils with high porosity commonly have abundant amount of clay materials and organic matter, characterized with high surface energy. These attributes are the main sources of errors of the MV and MV-VG models.

Typical examples of measured vs. predicted SMCs with the MV-VG model, ~~and~~ the scaling approach and ROSETTA for clay, sandy loam, loam, and silt loam textures are presented in Fig. 2 (a-f). For the clay (codes: 2340 and 4681), sandy loam (code: 3180 and 3200), loam (code: 3191) and silt loam (code: 3090) soils, the scaling approach fits the data well and ~~out~~ performs better than the MV-VG model at entire range of the SMC. For the silt loam soil (code:3090), the scaling approach slightly overestimates the moisture content through the entire range of suction heads and the MV-VG model underestimates the moisture content at low suction heads.~~As~~ Fig 2

(a-f) shows that the ROSETTA software performs worse in the wet part of SMC. Overall, the scaling approach performs better than MV-VG model and ROSETTA software for all soil samples (Table 2~~ and Fig.3~~) but, the performance of scaling approach did not suitably respond for two soil samples (codes: 3033, 3090). The residual model error may be related to the simplified representation of the total porosity which is considered equal to the saturated volumetric moisture content. The swelling properties and high organic carbon content of these soils (>4%, 3.85% respectively) may partially be a source of these errors. We further suspect that the complexity of the relationship between PSD, PoSD and pore connectivity can be effective in the model performance (Zhang et al., 2001). The assumption of the similarity between PSD and PoSD does not perform equally well to all soils.

We tentatively conclude that the scaling of the PSD curves using the parameter ζ generally performs better in predicting the SMC as compared to the original MV-VG model. The un-scaled MV-VG model underestimates the moisture content at high suction heads.

The most semi-physical based methods for predicting SMC rely on the use of empirical parameters to improve the SMC estimates from PSD (Lilly and Lin, 2004).Hydraulic properties are indeed affected by both the soil texture and the soil structure (Haverkamp et al., 2002). The MV-VG model uses the packing parameter, ζ, derived from soil bulk density as a metric of soil structure. Moreover, the scaling parameter that is inferred from the packing state is integrated in the scaled MV-VG model. Hence the soil structural features are integrated in the MV-VG model at two levels. First at the MV-VG model to convert moisture into pressure head, and second, to correct the SMC model prediction. The good performance of scaling approach in the wet range and dry range of SMC suggests a convenient of soil structural features in the SMC prediction.

Fig. 3 compares all estimated moisture contents, using MV-VG model and scaling approach, respectively, with the measured soil moisture content for all the 82 soil samples. The overall predictability of the two methods is evaluated by comparing the experimental data and the predicted soil moisture content on a 1:1 plot. Linear regression of the measured and estimated moisture contents, using the two methods for all the soil samples showed that the slope values were 0.7675 and 0.8484 and the coefficients of determination (R^2) between the estimated results and measured data for all soils were 0.765 and 0.8565 for MV-VG model and scaling approach, respectively. Hence, the MV-VG model and the scaling approach, underestimate the moisture content by about 23% and 15% respectively, while the bias of the scaling approach is smaller ~~less~~

than the MV-VG model. Regarding the R^2, the scaling method still remains the most preferable method. Comparing the overall predictability of the two methods, the correlation coefficient of linear regression~~, as a statistical summary, cannot exclusively~~ should not replace the visual examination of the data. We therefore, use the cubic polynomial function to adequately express the data variations. The fitted polynomial functions are drawn as shadowed red curves in Fig. 3(a, b).We further suggest that the area between the fitted curve and the 1:1 line (AE), is an expression of the systematic error. The values of AE were 0.0369 and 0.0250 for the MV-VG model and the scaling approach respectively. It confirms that the level of systematic errors of the scaling approach is about 33% less than that of MV-VG model. This result can be confirmed by the comparison of the R^2 values obtained when predicting the SMC for each soil with two methods (Table 2 columns 5 and 6). We conclude that the scaled PSD curve will result in a more accurate prediction of the SMC as compared to the un-scaled PSD data. ~~Moreover, the physically based scaling approach allows the upscaling of the physically based parameters which is essential forthe parameterization of the soil hydraulic properties for large study areas from individual soil samples (Kosugi and Hopmans, 1998).~~

To scale the SMC, Tuller and Or (2005) used the soil specific surface area (SA) and the thickness of water film to express the moisture content in dry range of the SMC. Despite, their reasonable model performance, the application of their procedure was limited, due to difficulty in the measurement or the estimation of SA.

Hayashi et al. (2007) found that in natural forested hillslope soils, the variability in the SMC is scaled and characterized by the variability in effective porosity. Nevertheless, the determination of the effective porosity is also difficult.

The scaling factor proposed in current study is defined by using the index of packing state which can be determined easily from the bulk density of soil and particle density. As compared to prediction models that rely on the measured attributes as suggested above, or prediction models that rely on measured SMC data, our approach is based on a robust metric of the soil structure: the packing density. It does not rely on any other additional empirical parameter. The scaled MV-VG model is therefore very parsimonious and robust. We therefore conclude that the scaled MV-VG model may be appropriate for predicting SMC from basic soil data.

5 Conclusion

Using a new scaling approach, the current study showed that the continuous form of SMC curve can be predicted from knowledge of PSD, as modeled by the van Genuchten (1980) model and particle packing state. In this approach it was assumed that the scaling factor can be defined as the ratio of packing state of a soil sample and the packing state of a reference soil. Results showed that the proposed approach can adequately predict the SMC of 82 soil samples selected from the UNSODA database. It was further found ~~further~~ that the scaling approach provides better predictions of the SMC than MV-VG model and ROSSETA software, especially in the dry range of the SMC. For soils for which the error was important, we attributed the proposed scaling approach error to high organic carbon content and swelling properties of the soil. Indeed, in these soils the soil pore structure and porosity is changing in time, leading to uncertainty in the scaling factor based on the soil porosity.

In summary, we concluded that the main advantages of the proposed scaling approach as compared to many SMC prediction models are:i) the applied scaling factor is determined easily from soil bulk and particle densities; ii) the scaling factor has physical meaning, which doesn't depend on soil database and empirical parameters; iii) the proposed approach predicts a continuous form of the SMC; and, iv) this approach estimates the SMC more appropriately in comparison with many other models. Considering that, there is no further need for empirical parameters, we conclude that this approach may be useful in estimating the SMC ~~at larger~~for regional scale soil hydrological studies ~~field scales~~.

References

~~Ahuja, L. R., Nancy, J. W., and Nielsen, D. R.: Scaling soil water properties and infiltration modeling, Soil Sci. Soc. Am. J., 48, 970-973, 1984.~~

Arya, L. M., Bowman, D. C., Thapa, B. B., and Cassel, D. K.: Scaling soil water characteristics of golf course and athletic field sands from particle-size distribution, Soil Sci. Soc. Am. J., 72, 25-32, 2008.

Arya, L. M., Leij, F. J., van Genuchten, M. T., and Shouse, P. J.: Scaling parameter to predict the soil water characteristic from particle-size distribution, Soil Sci. Soc. Am. J., 63, 510-519, 1999.

Arya, L. M., and Paris, J. F.: A physicoempirical model to predict the soil moisture characteristic from particle-size distribution and bulk density data, Soil Sci. Soc. Am. J., 45, 1023-1030, 1981.

Arya, L. M., Ritchter, J. C., and Davidson, S. A.: A comparison of soil moisture characteristic predicted by the Arya–Paris model with laboratory-measured data, AgRISTARS Tech. Rep. SM-L1-04247, JSC-17820, NASA-Johnson Space Center: Houston, TX, 1982.

Basile, A., and D'Urso, G.: Experimental corrections of simplified methods for predicting water retention curves in clay-loamy soils from particle-size determination, Soil Technol., 10, 261-272, 1997.

Chan, T. P., and Govindaraju, R. S.: Soil water retention curves from particle-size distribution data based on polydisperse sphere systems, Vadose Zone J., 3, 1443-1454, 2004.

~~Das, B.S., Haws, N.W., andRao, P.S.C.:Defining geometric similarity in soils, Vadose Zone J.,4, 264-270, 2005.~~

~~Deurer, M., andDuijnisveld, W.H.M.:Spatial analysis of water characteristic functions in a sandy podzol under pine forest, Water Resour. Res.,36, 2925-2935, 2000.~~

do Carmo, M.: Differential geometry of curves and surfaces, Prentice-Hall, pp.98, 1976.

~~Eching, S., Hopmans, J., and Wallender, W.: Estimation of in situ unsaturated soil hydraulic functions from scaled cumulative drainage data,Water Resour. Res., 30, 2387-2394, 1994.~~

Fredlund, M. D., Fredlund, D. G., and Wilson, G. W.: An equation to represent grain-size distribution, Canadian Geotechnology J., 37, 817-827, 2000.

Gras, J-P., Delenne, J-Y., Soulie, F., and ElYoussoufi, M. S.: DEM and experimental analysis of the water retention curve in polydisperse granular media, Power Tech., 12, 231-238, 2010.

Havayashi, Y., Kosugi, K., and Mizuyama, T.: Soil water retention curves characterization of a natural forested hillslope using a scaling technique based on a lognormal pore-size distribution, Soil Sci. Soc. Am. J., 73, 55-64, 2007.

Haverkamp, R., and Parlange, J. Y.: Comments on "A physicoemperical model to predict the soil moisture characteristic from particle-size distribution and bulk density data", Soil Sci. Soc. Am. J., 46, 1348-1349, 1982.

Haverkamp, R., Reggiani, P., Nimmo, J. R.: Property-Transfer Models. In:Methods of soil analysis, Part 4,Dane, J. H.andTopp, G. C. (Eds.) SSSA Book Series No. 5, SSSA, Madison, WI, 759–782, 2002.

Haverkamp, R., Leij, F. J., Fuentes, C., Sciortino, A., and Ross, P. J.: Soil water retention: I. Introduction of a shape index, Soil Sci. Soc. Am. J., 69, 1881-1890, 2005.

Hopkins, A. B., and Stillinger, F. H.: Dense sphere packings from optimized correlation functions, Phys. Rev.E., 79, 031123, doi:10.1103/PhysRevE.79.031123, 2009.

Hwang, S. I., and Choi, S. I.: Use of a lognormal distribution model for estimating soil water retention curves from particle-size distribution data, J. Hydrol., 323, 325-334, 2006.

Jamieson, R. C., Gordon, R. J., Sharples, K. E., Stratton, G. W., and Madani, A.: Movement and persistence of fecal bacteria in agricultural soils and subsurface drainage water: A review, Can. Biosyst. Eng., 44, 1-9, 2002.

Kern, J. S.: Estimation of soil water retention models based on soil physical properties, Soil Sci. Soc. Am. J., 59, 1134-1141, 1995.

Kosugi, K.: Lognormal distribution model for unsaturated soil hydraulic properties,WaterResour. Res., 32, 2697-2703, 1996.

Kosugi, K., and Hopmans, J. W.: Scaling water retention curves for soils with lognormal pore-size distribution, Soil Sci. Soc. Am. J., 62, 1496-1505, 1998.

Leij, F. J., Schaap, M. G., and Arya, L. M.: Indirect methods, in: Methods of Soil Analysis, Part 4, Dane, J. H. and Topp, C. G. (Eds.), SSSA Book Series 5, SSSA, Madison, 1009-1045. 2002.

Lilly, A., and Lin, H.: Using soil morphological attributes and soil structure in pedotransfer functions, in: Development of pedotransfer functions in soil hydrology, Pachepsky, Ya., and Rawls, W. (Eds.), Dev. Soil Sci. 30. Elsevier, Amsterdam, 115–141. 2004.

Marquardt, D. W.: An algorithm for least-squares estimation of nonlinear parameters. SIAM Journal on Applied Mathematics, 431-441, 1963.

Milla´n, H., and Gonza´lez-Posada, M.: Modelling soil water retention scaling. Comparison of a classical fractal model with a piecewise approach, Geoderma, 125, 25-38, 2005.

Milla´n, H., Gonza´lez-Posada, M., Aguilar, M., Domınguez, J., and Cespedes, L.: On the fractal scaling of soil data. Particle-size distributions, Geoderma, 117, 117-128, 2003.

~~Miller, E. E., and Miller, R. D.: Physical theory of capillary flow phenomena,J. SOC. Induster. Apple. Math., 27, 324-332, 1956.~~

Minasny, B., and McBartney, A. B.: The neuro-m method for fitting neural network parametric pedotransfer functions, Soil Sci. Soc. Am. J., 66, 352-361, 2002.

Mohammadi, M. H., and Meskini-Vishkaee, F.: Predicting the film and lens water volume between soil particles using particle size distribution data, J. Hydrol., 475, 403-414, 2012.

Mohammadi, M.H., and Meskini-Vishkaee, F.: Evaluation of soil moisture characteristics curve from continuous particle size distribution data, Pedosphere, 23(1), 70-80, 2013.

Mohammadi, M. H., and Vanclooster, M.: Predicting the soil moisture characteristic curve from particle size distribution with a simple conceptual model, Vadose Zone J., 10, 594-602, 2011.

Mohammadi, M. H., Neishabouri, M. R., and Rafahi, H.: Predicting the solute breakthrough curve from soil hydraulic properties, Soil Sci., 174, 165-173, 2009.

Nasta, P., Kamai, T., Chirico, G. B., Hopmans, J. W., and Romano, N. :Scaling soil water retention functions using particle-size distribution, J. Hydrol., 374, 223-234, 2009.Naveed, M., Moldrup, P., Tuller, M., Ferre, T. P. A., Kawamato, K., Komatso, T., and de Jong, L. W.: Prediction of the soil water characteristic from soil particle volume fractions,Soil Sci. Soc. Am. J., 76, 1946–1959, doi:10.2136/sssaj2012.0124, 2012.

Nemes, A., Schaap, M. G., and Leij, F. J.: The UNSODA unsaturated soil hydraulic property database, version 2.0, available at: http://www.ars.usda.gov/Services/docs.htm?docid=8967 (last access: November 2013), 2000.

Nimmo, W., Herkelrath, N., and Luna, A. M. L.: Physically based estimation of soil water retention from textural data: general framework, new models, and streamlined existing models, Vadose Zone J., 6, 766-773, 2007.

~~Oliveira, L. I., Demond, A. H., Abriola, L. M., and Goovaerts, P.: Simulation of solute transport in a heterogeneous vadose zone describing the hydraulic properties using a multistep stochastic approach,Water Resour. Res.,42, W05420, doi:10.1029/2005WR004580, 2006.~~

Or, D., and Tuller, M.: Liquid retention and interfacial area in variably saturated porous media: Upscaling from single-pore to sample-scale model, Water Resour. Res., 35(12), 3591-3605, 1999.

Perfect, E.: Modeling the primary drainage curve of prefractal porous media, Vadose Zone J., 4, 959-966, 2005.

Puhlmann, H., and von Wilpert, K.: Pedotransfer functions for water retention and unsaturated hydraulic conductivity of forest soils, J. Plant Nutr. Soil Sci., 175(2), 221-235, 2012.

Ryel, R. J., Caldwell, M. M., Yoder, C. K., Or, D., and Leffler, A. J.: Hydraulic redistribution in a stand of Artemisia tridentata: Evaluation of benefits to transpiration assessed with a simulation model, Oecologia, 130, 173-184, 2002.

Santamarı´a, J., and Toranzos, G. A.: Enteric pathogens and soil: A short review, Int. Microbiology, 6, 5-9, doi:10.1007/s10123-003-0096-1, 2003.

Saxton, K. E., Rawls, W. J., Romberger, J. S., and Papendick, R. I.: Estimating generalized soil-water characteristics from texture, Soil Sci. Soc. Am. J., 50, 1031-1036, 1986.

Schaap, M. G.: Models for indirect estimation of soil hydraulic properties, in: Encyclopedia of Hydrological Sciences, Anderson, M. G. (Ed.), Wiley, pp.1145-1150, 2005.

Schaap, M. G., and Leij, F. J.: Database related accuracy and uncertainty of pedotransferfunctions, Soil Sci., 163, 765-779, 1998.

Schaap, M. G., Leij, F. J., and van Genuchten, M. Th.: Neural network analysis for hierarchical prediction of soil water retention and saturated hydraulic conductivity, Soil Sci. Soc. Am. J., 62, 847-855, 1998.

Schaap, M.G., Leij, F. J., and van Genuchten, M. Th.: ROSETTA: A computer program for estimating soil hydraulic parameters with hierarchical pedotransfer functions, J. Hydrol., 251, 163–176, 2001.

Schaap, M. G., Nemes, A., and van Genuchten, M. Th.: Comparison of models for indirect estimation of water retention and available water in surface soils, Vadose Zone J., 3, 1455-1463, 2004.

Schuh, W. M., Cline, R. L., and Sweeney, M. D.: Comparison of a laboratory procedure and a textural model for predicting in situ soil water retention, Soil Sci. Soc. Am. J., 52, 1218-1227, 1988.

~~Sharma, M. L., and Luxmoore, R. J.: Soil spatial variability and its consequences on simulated water balance, Water Resour. Res., 15, 1567-1573, 1979.~~

Smettem, K. R. J., and Gregory, P. J.: The relation between soil water retention and particle-size distribution parameters for some predominantly sandy Western Australian soils, Aust. J. Soil Res., 34, 695-708, 1996.

Teitje, O., and Tapkenhinrichs, M.: Evaluation of pedo-transfer functions, Soil Sci. Soc. Am. J., 57, 1088-1095, 1993.

Tuli, A., Kosugi, K., and Hopmans, J. W.: Simultaneous scaling of soil water retention and unsaturated hydraulic conductivity functions assuming lognormal pore-size distribution, Adv. Water Resour., 24, 677-688, 2001.

Tuller, M., and Or, D.: Water films and scaling of soil characteristic curves at low water contents, Water Resour. Res., 41, W09403, doi:10.1029/2005WR004142, 2005.

Tuller, M., Or, D., and Dudley, L. M.: Adsorption and capillary condensation in porous media: liquid retention and interfacial configurations in angular pores, Water Resour. Res., 35, 1949-1964, 1999.

Tyler, S., and Wheatcraft, S.: Application of fractal mathematics to soil water retention estimation, Soil Sci. Soc. Am. J., 53, 987-996, 1989.

van Genuchten, M. Th.: A closed-form equation for predicting the hydraulic conductivity of unsaturated flow, Soil Sci. Soc. Am. J., 44, 892-898, 1980.

Vaz, C., lossi, M. D. F., Naimo, J. d. M., Macero, A., Reichert, J. M., Reinert, D. J., and Cooper, M.: Validation of the Arya and Paris water retention model for Brazilian soils, Soil Sci. Soc. Am. J., 69, 577-583, 2005.

~~Vogel, T., Cislerova, M., and Hopmans, J.: Porous media with linearly variable hydraulic properties,Water Resour. Res., 27, 2735-2741, 1991.~~

Wang, Q. J., Horton, R., and Lee, J. A.: Simple model relating soil water characteristic curve and soil solute breakthrough curve, Soil Sci., 167, 436-443, 2002.

~~Warrick, A. W., Mullen, G. J., and Nielsen, D. R.: Scaling field-measurement soil hydraulic properties using a similar media concept,Water Resour. Res., 13, 355-362, 1977.~~

Weynants, M., Vereecken, H., and Javaux, M.: Revisiting Vereecken Pedotransfer Functions: Introducing a Closed-Form Hydraulic Model, Vadose Zone J., 8(1), 86-95, 2009.

Wu, L., Vomocil, J. A., and Childs, S. W.: Pore size, particle size, aggregate size, and water retention, Soil Sci. Soc. Am. J., 54, 952-956, 1990.

Zhuang, J., Jin, Y., and Miyazaki, T.: Estimating water retention characteristic from soil particle-size distribution using a non-similar media concept, Soil Sci., 166, 308-321, 2001.

Symbols and Abbreviations

SMC: Soil Moisture Characteristics curve
PSD: Particle Size Distribution.
PoSD: Pore Size Distribution.
MV: Mohammadi and Vanclooster (2011) model
BCC: Bundle of Cylindrical Capillaries
VG model: van Genuchten model
MV-VG model: integrated the MV model with the van Genuchten model
AP: Arya and Paris (1981)
PTF: Pedotransfer Function.
RMSE: Root Mean Square Error.
θ: the soil moisture content
S_e: effective saturation degree
θ_s: saturated moisture contents
θ_r: residual moisture contents
n: fitting coefficients
m: fitting coefficients
α: fitting coefficients
h: suction head
ξ: a coefficient depending on the state of soilparticles packing
e: the void ratio
ρ_s: soil particle density
ρ_b: soil bulk density
w_j: the mass fraction of particles in the *j*th particle-size fraction
S: saturation degree
P_i: cumulative mass fraction of soil particles
R_i: particle radius of the *i*th fraction
β: scaling factor
$\bar{\gamma}$: microscopic characteristic lengths of the reference
γ: microscopic characteristic lengths of the subjected soil
ξ_{max}: maximum value of packing parameter
n: pore size distribution index
λ: scaling factor
n^*: scaled the PoSD index in VG model
$\theta_{i(p)}$: predicted moisture content
$\theta_{i(m)}$: measured moisture content
RI: relative improvement
$RMSE_M$:RMSE values of MV-VG model
$RMSE_s$: RMSE values of scaling approach

AE: area between the fitted curve and 1:1 line

Table1. Textural classes and UNSODA codes for soils used for testing and evaluating the approach.

Textural class	Clay	Clay loam	Loam	Silt loam	Silty clay	Silty clay loam	Loamy sand	Sand	Sandy clay loam	Sandy loam
UNSODA codes	1400 2340 2361 2362 4120 4680 4681 2360	3033	3221 1211 1260 1261 2530 3190 3191 3222	2000,3090 3213,3261 4042,4070 4180,4181 2464,1341 1342,1350 1351,1352 2001,2002 2010,2011 2012,	3030 1360	3100 3101 1371	1160 2102 2103 3130 3150 3152 3160 3161 3170 3171 4251	1050,1240,1460 1464,1466,2100 3133,3134,3140 3141,3144,3155 3162,3163,3164 3165,3172,3340 4051,4152,4263 4272,4282,4441 4520,4650,4000	3202	1130 3180 3200 3290

Table2. Average Root Mean Square Error (RMSE), coefficients of determination (R^2), Relative Improvement (RI) compared to the MV-VG model and hydraulic parameters for each soil textural group, with standard deviations in parentheses.

Soil texture	*Number of soil*	*RMSE*					*R^2*	*RI value (%)*	*Hydraulic parameters*					
		MV-VG model	Scaling approach	ROSETTA	MV-VG model	Scaling approach	ROSETTA		θ_s ($L^3 L^{-3}$)	α (L^{-1})	m (-)	n (-)	n^* (-)	λ (-)
Clay	8	0.088 (0.014)	0.041 (0.020)	0.1150 (0.0403)	0.973 (0.017)	0.977 (0.020)	0.9153 (0.0710)	53.87 (16.74)	0.51 (0.04)	0.043 (0.085)	0.128 (0.104)	2.457 (2.195)	1.467 (1.127)	0.6228 (0.0706)
Clay loam	1	0.027	0.017	0.1468	0.725	0.872	0.9790	38.15	0.58	0.002	0.248	1.634	1.040	0.6365
Loam	8	0.078 (0.019)	0.045 (0.015)	0.0546 (0.0333)	0.896 (0.100)	0.913 (0.088)	0.8986 (0.0667)	41.04 (17.74)	0.45 (0.06)	0.042 (0.030)	0.157 (0.079)	3.115 (1.452)	2.187 (1.100)	0.6970 (0.0389)
Silt loam	19	0.082 (0.026)	0.059 (0.020)	0.0512 (0.0222)	0.922 (0.043)	0.950 (0.033)	0.9598 (0.0230)	25.63 (20.28)	0.44 (0.04)	0.019 (0.011)	0.233 (0.365)	4.937 (2.593)	3.637 (1.982)	0.7189 (0.0414)
Silty clay	2	0.076 (0.056)	0.061 (0.022)	0.0868 (0.0427)	0.932 (0.040)	0.941 (0.010)	0.9166 (0.0348)	3.01 (43.07)	0.51 (0.09)	0.040 (0.028)	0.195 (0.157)	1.573 (1.402)	1.091 (1.039)	0.6628 (0.0699)
Silty clay loam	1	0.129	0.093	0.1080	0.887	0.924	0.9083	28.15	0.43	0.020	0.116	2.548	1.870	0.7339
Loamy sand	11	0.093 (0.037)	0.060 (0.022)	0.0862 (0.0359)	0.893 0.062	0.926 (0.038)	0.9067 (0.0537)	32.63 (11.64)	0.40 (0.07)	0.067 (0.038)	0.179 (0.058)	5.488 (1.357)	4.048 (1.088)	0.7339 (0.0445)
Sand	27	0.093 (0.030)	0.073 (0.024)	0.0254 (0.0626)	0.854 (0.102)	0.893 0.081	0.8853 0.0760	20.74 (7.51)	0.37 (0.04)	0.052 (0.033)	0.458 (0.444)	5.592 (2.018)	4.380 (1.531)	0.8228
Sandy clay loam	1	0.084	0.065	0.0653	0.957	0.967	0.9702	23.07	0.36	0.043	0.054	8.000	6.582	0.7031 (0.0630)
Sandy loam	4	0.073 (0.014)	0.035 (0.015)	0.0776 (0.0547)	0.950 (0.028)	0.971 (0.008)	0.9364 (0.0278)	51.63 (18.12)	0.46 (0.05)	0.066 (0.020)	0.093 (0.032)	5.676 (1.526)	3.980 (1.137)	0.7641 (0.0444)
Average	**82**	**0.086 a (0.028)**	**0.060 b (0.024)**	**0.0745 a (0.0417)**	**0.898 (0.084)**	**0.927 (0.065)**	**0.9276 (0.064)**	**30.14 (18.88)**	**0.42 (0.07)**	**0.044 (0.040)**	**0.272 (0.338)**	**4.726 (2.329)**	**3.519 (1.816)**	**0.7177 (0.0532)**

[a] Different lowercase letters indicate significant differences at $P < 0.05$

Figure captions

Figure 1.The efficiency of scaling approach, % RI, defined with RMSE (Eq. (~~15~~14)) as function of (a) the saturated moisture content and (b) scaling factor for all soil samples. **: significant at P= 0.01.

Figure 2. Examples of measured vs. predicted soil moisture characteristics curve (SMC) for each texture using the integrated MV-VG model (Eq. (9)), scaling approach (Eq. (12~~3~~)) and ROSETTA: for (a) clay soil, (b) sandy loam soil, (c) loam soil, (d) sandy loam soil, (e) clay soil and (f) silt loam soil.

Figure 3. Comparisons of the measured and estimated moisture contents for 82 selected soils using the MV-VG model (Eq. (9)) and scaling approach (Eq. (~~13~~12)). Dashed lines: the 1:1 line. Solid lines: linear-regression line, shadowed solid red line: nonlinear-regression-line.