

1
2
3
4
5
6 MILLENNIAL SCALE VARIABILITY IN HIGH MAGNITUDE FLOODING ACROSS
7 BRITAIN
8
9
10

11 NEIL MACDONALD
12
13
14
15 Department of Geography and Planning, University of Liverpool, Liverpool, L69 7ZT
16
17
18
19
20
21 e: Neil.Macdonald@liverpool.ac.uk
22
23 t: 0151 794 2510
24
25

26 **Abstract**

27 The last decade has witnessed severe flooding across much of the globe, but have these floods
28 really been exceptional? Globally, relatively few instrumental river flow series extend beyond 50
29 years, with short records presenting significant challenges in determining flood risk from high-
30 magnitude floods. A perceived increase in extreme floods in recent years has decreased public
31 confidence in conventional flood risk estimates; the results affect society (insurance costs),
32 individuals (personal vulnerability) and companies (e.g. water resource managers). Here we show
33 how historical records from Britain have improved understanding of high magnitude floods, by
34 examining past spatial and temporal variability. The findings identify that whilst recent floods are
35 notable, several comparable periods of increased flooding are identifiable historically, with periods
36 of greater frequency (flood-rich periods). Statistically significant relationships between the British
37 flood index, the Atlantic Meridional Oscillation and the North Atlantic Oscillation Index are
38 identified. The use of historical records identifies that the largest floods often transcend single
39 catchments affecting regions and that the current flood rich period is not unprecedented.

40

41 **Keywords:** flood, historic, flood-rich, spatial and temporal variability, Atlantic Meridional
42 Oscillation, North Atlantic Oscillation, Britain

43

44 **1 INTRODUCTION**

45 One of the greatest challenges presently facing river basin managers is the dearth of reliable long-
46 term data on the frequency and severity of extreme floods, with an average gauged record length of
47 ~40 years in the UK (Marsh and Lees, 2003). Historical accounts represent a precious resource
48 when considering the frequency and risks associated with high-magnitude low-frequency floods
49 (Williams and Archer 2002). Historical flood records are found in a variety of forms, directly or
50 indirectly chronicling historic floods (Brázdil et al., 2005); sources include, documentary accounts
51 e.g. journals, newspapers, diaries (McEwen, 1987; Brázdil et al., 2012); flood stones (markers
52 indicating the greatest spatial flood extent) and epigraphic markings (inscribed water levels on
53 structures; see Macdonald, 2007) for sites around the globe (Popper, 1951; Camuffo and Enzi,
54 1996; Brázdil 1998; Demarée, 2006; Bürger et al., 2007). Historical accounts often contain
55 important details including incidence, magnitude, frequency (comparable to other historic events)
56 and seasonality. With records becoming more frequent since AD 1500 and coeval descriptions
57 permitting account corroboration (Brázdil et al., 2006). Historic centres often retain the most
58 complete series of historical records, as the presence of literate individuals associated with
59 important monastic, trade and/or governmental functions provide detailed flood accounts
60 (Macdonald et al., 2006), an important aspect in the preservation of early materials. This paper
61 presents the first coherent large scale national analysis undertaken of historical flood chronologies,
62 providing an unparalleled network of sites (Fig. 1), permitting analysis of the spatial and temporal
63 distribution of high-magnitude flood patterns and the potential mechanisms driving periods of
64 increased flooding at a national scale (Britain) over the last 800 years.

65

66 **2 SERIES CONSTRUCTION**

67 Site inclusion within this study is dependent on the availability of detailed historical accounts and
68 the presence of relatively long instrumental river flow/level series (>40 years in length). Historical
69 accounts were collated and augmented onto existing instrumental series, with historical flood levels
70 estimated based on documented descriptions (see Wetter et al., 2011), physical evidence or
71 epigraphic markings, providing estimates of flow (Herget & Meurs, 2010), with greater significance
72 placed on ranking event severity than on precise discharge estimation (Payrastre et al., 2011) (Fig.
73 2). Only those floods (historical and instrumental) exceeding the 90th percentile based on the
74 instrumental period are included, thus ensuring only the largest events are considered, providing a
75 threshold of events comparable to those likely to have been recorded within the historical period.
76 The largest flood events are unlikely to be significantly impacted by moderate anthropogenic driven
77 changes within catchments (Mudelsee et al., 2003; Macdonald and Black, 2010; Hall et al., 2015);

78 where significant catchment/channel and floodplain (see Lewin, 2010) changes have occurred (e.g.
79 channel cross-section, land use, urbanisation, etc.), the impact, where possible, has been accounted
80 for using available information (Elleder et al., 2013), with greater confidence in comparable
81 catchment form for the later period (c.1750-), compared to earlier periods (Macdonald et al., 2014).
82 The data used within this paper focusses on single locations, as merging of historical data over
83 whole catchments is fraught with difficulties (Böhm, et al., 2015), as such ‘stable’ sections of
84 channel are selected, where possible, at sites with long detailed historical flood records.
85

86 **3 CATCHMENT CHARACTERISTICS**

87 A brief summary of the catchment conditions and anthropogenic influence on each of the systems is
88 provided below, detailed in depth discussions of local histories and land-use practices are provided
89 in the cited papers.
90

91 **3.1 River Findhorn, Forres**

92 The River Findhorn drains the Monadhliath Mountains in Central Scotland, with a predominantly
93 metamorphic bedrock, including granitic intrusions, extensive blanket peat coverage with
94 agricultural activities along the coastal strip and much heath and mountainous land, with limited
95 anthropogenic development within the catchment. Instrumental series are available at Shenachie
96 (07001; 1960-) and Forres (07002; 1958-), with an upstream catchment area of 782km²; the
97 Findhorn has received considerable attention within a British context (NERC 1975; Newson 1975;
98 Acreman 1989) as it includes one of the best documented ‘extreme’ floods of the nineteenth
99 century. Sir Thomas Dick Lauder’s *‘An account of the great floods of August 1829 in the province
100 of Moray and adjoining districts* (1830) provides a detailed eyewitness account of the floods and
101 the destruction across the region, with detailed information permitting the reconstruction of the
102 flood. Throughout Lauder’s account he frequently comments on human modification of the
103 landscape, partly attributing the severity of the 1829 flood to agricultural improvement and drainage
104 undertaken in the decades of the late eighteenth and early nineteenth centuries within the catchment.
105 Within the instrumental period the flood of 1970 is the largest, estimated at 2402 m³s⁻¹, but has
106 subsequently been reduced to 1113 m³s⁻¹ following considerable reanalysis; with the 1829 flood
107 estimated to be between 1500-1800 m³s⁻¹ (McEwen and Werritty, 2007). The present river channel
108 consists of a number of bedrock sections, particularly within the upper catchment, with alluvial
109 highly mobile sections within the lower catchment susceptible to lateral avulsion, though McEwen
110 and Werritty (2007) note limited migration in most channel sections since the present channel was
111 excavated during the 1829 flood. There are no severe flood events recorded on the Findhorn

112 between the flood of 1829 and the start of the instrumental series in the 1950s from which reliable
113 estimates can be derived, though it is notable that several floods are described in the period between
114 1914 and 1924, though accurate estimates of their discharge are not achievable from the available
115 records.

116

117 3.2 River Tay, Perth

118 The River Tay has the largest mean discharge of any British river ($165 \text{ m}^3\text{s}^{-1}$; Marsh and Lees,
119 2003) with a mean annual flood of $990 \text{ m}^3\text{s}^{-1}$. Although relatively small by European standards, the
120 Tay catchment is one of Britain's largest, draining 4690 km^2 of the Scottish Highlands, with several
121 mountain peaks $>1000 \text{ mAOD}$ (meters Above Ordnance Datum). Annual precipitation in excess of
122 3000 mm a^{-1} in the western highlands is not uncommon as a result of high elevation and westerly
123 situation (Roy, 1997); by contrast lowland sections of the catchment (around Perth) have an average
124 annual rainfall of $\sim 700 \text{ mm a}^{-1}$ (Jones *et al.*, 1997) and annual evapotranspiration losses of
125 $\sim 450 \text{ mm a}^{-1}$ (Harrison, 1997). The River Tay has six major tributaries: the Almond, Earn, Garry,
126 Isla, Lyon and Tummel, with a tidal limit near the Tay-Almond confluence approximately 4 km
127 upstream of Perth. The longest gauged flow record is at Caputh (15003; since 1947), despite being
128 shorter, the Ballathie (15006) record (1952-) includes flows from the River Isla tributary, $\sim 6 \text{ km}$
129 upstream of the city of Perth and as such provides a better comparison to epigraphic flood levels in
130 the city, most notably those on Smeaton's Bridge (Macdonald *et al.*, 2006). Generally, the
131 catchment is characterised by thin soils and impermeable bedrock with high runoff rates; while
132 Lochs Tummel, Tay and Lyon significantly reduce flooding by attenuating flood peaks. The
133 development of major hydro-schemes in the Tay catchment completed in 1957 (Payne, 1988),
134 incorporates 42.2% of the upper Tay catchment area (Marsh and Lees, 2003). The development
135 consists of two power schemes, i) the Tummel-Garry scheme (1649 km^2) to Pitlochry Dam
136 (including inflows from a further 130 km^2 of the headwaters of the River Spey; Marsh and Lees,
137 2003); ii) to the south, the Breadalbane scheme which controls a further 511 km^2 draining to
138 Comrie Bridge, at the Tay-Lyon confluence. Four sets of flood information are used at Perth for
139 constructing the flood series: i) a gauged record since 1952 at Ballathie; ii) epigraphic makings on
140 Smeaton's Bridge in central Perth (intermittent since 1814); iii) a series of river level readings from
141 the old waterworks in Perth (intermittent since 1877); and, episodic documentary accounts which
142 extend back to AD 1210. A rating curve constructed from peak flows at the Ballathie gauging
143 station and sites in Perth enables estimated discharges to be assigned to historic flood flows
144 (Macdonald *et al.*, 2006). The rich documentary sources reporting floods in Perth permits extension
145 of the record back to 1210, these accounts are compared to contemporary events within the

146 augmented series with reference to relative extent in relation to buildings or road junctions. For the
147 purpose of this analysis it is assumed that the relationship between stage and discharge at the site of
148 Smeaton's Bridge and Ballathie has not changed over the intervening period; see Macdonald et al.
149 (2006) and Werritty et al., (2006) for a more detailed discussion of the flood history and flood series
150 reconstruction, hydrological changes and landscape change within the catchment.

151

152 3.3 River Tweed

153 The River Tweed rises at Tweed's Well in the Lowther Hills flowing east through the Scottish
154 Boarders before entering Northumberland in Northeast England and flowing into the North Sea at
155 Berwick-on-Tweed. The River Tweed consists of two principal rivers, the Tweed and Teviot
156 draining from the west and southwest respectively and the Whiteadder draining from the northwest
157 entering the Tweed c.3km upstream of Berwick. The reservoirs in the headwaters have limited
158 impact on the river discharges downstream, with ~30% lowland agriculture and ~70% upland given
159 to moorland and upland hill pasture (Marsh and Hannaford, 2008), with few urban centres, the
160 exceptions being the towns of Berwick-upon-Tweed, Coldstream and Kelso. The geology is of
161 mixed bedrock, predominantly impervious Palaeozoic formations with thick superficial deposits.
162 Annual precipitation in c. 790mm^a⁻¹ (Marchmount House) (McEwen, 1989).

163

164 There is a long well documented flood history for the town of Kelso, a historic strategically
165 important town on the English – Scottish boarder, which was held by both countries several times
166 during the various conflicts. The town was important commercially as a market town, with a
167 number of historic monastic centres nearby (e.g. Lindisfarne and Kelso Monastery) and was on one
168 of the main routes between London and Edinburgh for much of the period. The town of Berwick on
169 the coast contained the oldest bridging points on the Tweed, with the earliest recorded flood dating
170 from 1199 resulting in the loss of the bridge and subsequent rebuilding costs, the second bridge was
171 destroyed by the English in 1216, with the third lost in 1294 again to a flooding. A forth wooden
172 bridge was built in 1376 and subsequently replaced by the present stone Berwick Old Bridge in
173 1634 (ICE, undated). The first bridge at Kelso was built in 1754 and replaced a ferry, in October
174 1756 part of the bridge collapsed during a flood killing six, the bridge was repaired but a storm in
175 October 1797 lead to its collapse, with a replacement bridge constructed in 1803 by the engineer
176 John Rennie.

177

178 Gauged river flow series for Sprouston (21021) and Norham (21009) exists from 1969- present and
179 1959-present respectively; with Sprouston located ~1km downstream of Kelso and Norham located

180 ~9km upstream of the Whiteadder confluence draining from the north (~12km upstream of the
181 coastal town of Berwick-upon-Tweed). Historical flood series have been produced by McEwen
182 (1990) for the rivers Tweed, Teviot, Whiteadder and Leader (Tweed tributary) from 1750, with a
183 longer series for the Tweed starting in AD218, but early records are of questionable reliability as
184 the original sources are unknown. The long chronology produced by McEwen (1990) notes that the
185 largest floods prior to 1750 occurred in August 1294, March 1296, 1523, 1631, 1646, and possibly
186 three other events in 834/6, 1327/8 and 1333, but again with lower confidence; with major floods
187 since 1750 at Kelso in ranked order (1) 1948, (2) 1831, (3) 1846 and 1881, (4) 1891, 1926 and 1982
188 ($1452 \text{ m}^3\text{s}^{-1}$), (5) 1956, 1962 ($1174 \text{ m}^3\text{s}^{-1}$, Norham) and 1977 ($1269 \text{ m}^3\text{s}^{-1}$); discharge at Sprouston
189 unless stated. Flood events since the publication of McEwen's 1990 study of a comparable
190 magnitude ($>1250 \text{ m}^3\text{s}^{-1}$) have occurred in 2002 ($1444 \text{ m}^3\text{s}^{-1}$) and 2005 ($1436 \text{ m}^3\text{s}^{-1}$), which appear
191 of a comparable discharge to that of 1982 and as such are placed in the rank four category.
192 Additional floods not identified by McEwen occurred in 22nd October 1756 and 26 October 1797,
193 both of which led to the loss of the bridge, the former resulting in several lost lives (Star,
194 26/10/1797). Based on the descriptive accounts and details provided in McEwen (1990) and other
195 sources, discharges for the historic events are presented as $1850 \text{ m}^3\text{s}^{-1}$ for 1948 (rank 1); $1750 \text{ m}^3\text{s}^{-1}$
196 for 1831 (rank 2); $1650 \text{ m}^3\text{s}^{-1}$ for 1846 and 1881 (rank 3); $1450 \text{ m}^3\text{s}^{-1}$ for 1891 and 1926 (rank 4)
197 and $1250 \text{ m}^3\text{s}^{-1}$ for 1956 (rank 5). For the floods of 1294; 1296; 1523; 1631; 1646, 1756 and 1797
198 an estimated discharge of $1450 \text{ m}^3\text{s}^{-1}$ is used within the analysis, each event was worthy of
199 description, with several attributed to the loss of bridges, life or other notable structures and
200 therefore are likely to be of equivalent or greater than rank 4.

201

202 3.4 River Tyne

203 The River Tyne in Northeast England consists of two principal rivers, the South and North Tyne
204 rivers which join near the town of Hexham to form the River Tyne. The geology of the upper
205 catchment is characterised by Carboniferous Limestone and Millstone Grit, with a thick layer of
206 alluvial drift material covering the lower catchment, with land-use predominantly upland farming,
207 grassland and woodland with relatively little urban development. The North Tyne rises near the
208 Scottish boarder before flowing south east, the principal water body on the tributary is Kielder
209 Water. The construction of Kielder Water, the largest UK reservoir with the potential to hold 200B
210 litres has considerably altered flood discharges within the North Tyne since its completion in 1982,
211 attenuating $\sim 240 \text{ km}^2$ (11%) of the catchment. Archer et al. (2007) identifies that the impact on
212 flood discharges ranges from $114 \text{ m}^3\text{s}^{-1}$ at mean annual flood to $225 \text{ m}^3\text{s}^{-1}$ at the 20 year return
213 period, and is liable to increase with less frequent events. The South Tyne's source on Alston Moor

214 in Cumbria, before flowing north-northeast through the Tyne Gap, there are no notable
215 impoundment structures on the river, before reaching its confluence with the North Tyne near
216 Hexham.

217

218 The River Tyne (~321km length and 3296km² catchment) has undergone extensive river channel
219 modification over recent centuries as a result of gravel extraction (Rumsby and Macklin 1994), with
220 Archer (1993) estimating approximately 4.5M tons having been extracted during the period 1890-
221 1970, from 15 sites along the rivers course. The level of gravel extraction has had a considerable
222 impact on the channel and bedform of the river within the lower reaches, altering the stage-
223 discharge relationships, as such the creation of a reliable long flood series is challenging. Extensive
224 analysis of available historical information was undertaken by Archer (1992) for his book *Land of*
225 *Singing Waters*, and subsequent book *Tyne and Tide* (Archer 2003). The discharge series for the
226 gauging station at Bywell is used (1956-), but earlier flows modified after Archer (2007) to account
227 for gravel extraction (1955-61) and the construction of Kielder Water. The gauge at Bywell was
228 installed following severe flooding in January 1955, with an estimated discharge of 1520m³s⁻¹
229 (Archer Pers. Comm. 2005). Notable historical flood discharges on the Tyne have previously been
230 estimated, particularly the 1815 (1700m³s⁻¹) and 1771 (3900m³s⁻¹) floods, with an uncertainty of
231 c.20% (Archer, 1993), the latter being the most devastating flood event recorded, not just on the
232 Tyne but regionally, with many rivers losing bridges during this event (e.g. see Archer, 1987). The
233 1771 flood appears to be the largest within the flood record since AD1200, with 1815 ranked third,
234 with the flood of 1339 ranked second. The information available for the flood of 1339, is limited,
235 though the Chronicle of Lanercost, 1272-1346 (translated by Maxwell, 1913) describes the event as:
236 “...on the third day before the feast of the Assumption of the Glorious Virgin [14th August]
237 a marvellous flood came down by night upon Newcastle-on-Tyne, which broke down the
238 town-wall at Walkenow for a distance of six perches, where 160 men, with seven priests and
239 others, were drowned”.

240 Jervoise (1931) notes that a stone bridge built at Newcastle by the Newcastle Corporation and
241 the Bishop of Durham in AD1248 survived a flood when 90 years old (c. 1339), but suffered
242 severe flood damage with the loss of 120 lives and was eventually destroyed during the 1771
243 flood. The severity of the floods of 1771 and 1815 led to the production of a book ‘*An account of*
244 *the great floods in the rivers Tyne, Tees, Wear, Eden, &c. in 1771 and 1815*’ in 1818 by William
245 Garret, documenting the impacts of the floods across Northern England. A number of additional
246 accounts document floods between 1722 and the start of the gauged series in 1956, these include
247 1722, 1763, 1782, 1831, 1856, 1881, 1903, within this study these are estimated to have

248 discharges between (1225-1375m³s⁻¹), making them broadly comparable to the 2005 flood (1370
249 m³s⁻¹) on the River Tyne. As Archer (2007) notes when commenting on the 1955 and 2005
250 floods, it is conceivable that the floods of 1722, 1763, 1782, 1831 and 1856 may have been
251 greater, as the estimation of historical discharges on the River Tyne are particularly challenging
252 as a result of the uncertainties in estimation. The recent December 2015 flood on the Tyne is
253 likely to be greatest since 1771, with a level exceeding the 1815 event by 0.4m, but below that of
254 1771, with a provisional discharge of approximately 1700m³s⁻¹ (Parry et al., 2016).

255

256 **3.5 River Eden**

257 The River Eden in Northwest England, has a catchment area of c.2300km², it flows north-northwest
258 direction for much of its course from its source at Black Fell Moss, Mallerstang, in the Yorskhire
259 Dales through to the Solway Firth. It has four principal tributaries, the Eamont, Irthing, Petteril and
260 Caldew. The Eamont drains the upland area of the eastern Lake District with a confluence with the
261 Eden near Penrith, followed by the Irthing tributary joining from the east-northeast c.10km
262 upstream of Carlisle, with the River Petteril confluence c.1km upstream and the Caldew confluence
263 adjacent to the city of Carlisle. The catchments geology consists of Carboniferous Limestones to the
264 east and impervious Lower Palaeozoics of the Lake District massif to the west, with extensive
265 Permo-Triassic sandstone within the Vale of Eden (Marsh and Hannaford, 2008). The land-use is
266 predominantly rural, with moorland and upland grazing at elevation and grasslands at lower
267 elevations and limited urban coverage except for the towns of Appleby, Penrith and the city of
268 Carlisle. Precipitation can exceed 2000 mma⁻¹ at elevation in the Lake District, with an average
269 precipitation at Carlisle of 787mma⁻¹ (Todd et al., 2015).

270

271 Severe flood events have affected Carlisle in recent years (2015, 2005), with three people killed and
272 ~2700 properties affected in 2005. A rich detailed history of flooding exists for Carlisle, with a
273 combination of existing reconstructions (Smith and Tobin, 1979; Macdonald 2006; Patterson and
274 Lane, 2012), a series of flood marks on Eden Bridge since 1822 and descriptive accounts from
275 multiple sources augment the instrumental series from Sheepmount gauging station (1967-present),
276 a gauged series is also available from Warwick bridge from 1959, but this is upstream of the
277 confluence with the Irthing. The 2005 (1516m³s⁻¹) flood event is recorded by the EA as one meter
278 higher than the previous highest mark 1822, with the flood of 2015 (1680m³s⁻¹) 0.6m higher than
279 2005 (Environment Agency 2015), the recent flood of December 2015 is provisionally estimated at
280 approximately 1700m³s⁻¹ (Parry et al., 2016). Following the severe floods of 1968, Smith and Tobin
281 (1979) mapped the flood extent of all known flood events between 1800 and 1968, producing a

282 ranked series of 49 major floods at Carlisle, of which 1822, 1856, 1925 and 1968 are the largest,
283 these are all also marked on Eden Bridge. The flood of 1771, whilst notable does not appear as
284 extreme as witnessed in catchments on the eastern side of northern England, accounts of bridges
285 being lost over several of the principal tributaries are documented in Garret (1818), with livestock
286 lost at Hole Farm near Carlisle. Notable floods prior to 1771 include 1360 and 1767, with floods
287 also recorded in 1684, 1685, 1710 and 1763 (Chronology of British Hydrological Events, Black and
288 Law, 2004); the snowmelt flood of 1767 is documented in the weather accounts of the Bishop of
289 Carlisle as discussed by Todd et al., (2015). There are limited materials documenting the 1360
290 flood, though Jervoise (1931) notes that the old bridge over the Eden was destroyed by a flood in
291 1360, as Bishop Welton of Carlisle gave indulgences to those helping towards the repair of the
292 bridge (*Testamenta Karleolensia* – translated by Ferguson 1893), with several accounts
293 commenting on the remains of the bridge still being visible into the early nineteenth century.

294

295 **3.6 River Ouse, York**

296 The Yorkshire Ouse located in northeast England has a catchment area of 3315 km² upstream of
297 Skelton, the site of the present gauging station (27009), on the northern outskirts of the city of
298 York. Upstream of the city the main tributaries of the River Ouse are the Rivers Swale, Ure and
299 Nidd, together draining much of the Northern Pennines. Precipitation totals vary throughout the
300 catchment, ranging from in excess of 1800 mm a⁻¹ in upland areas to less than 600 mm a⁻¹ in the
301 Vale of York and adjacent lowland regions (Meteorological Office, 2002). The geology of the
302 upper catchment is characterised by Carboniferous Limestone and Millstone Grit, with a thick layer
303 of alluvial drift material covering the lower catchment in the Vale of York. Land use varies
304 throughout the catchment, with predominantly arable and pastoral farming in lowland areas (Dennis
305 et al., 2003), with increasing levels of grassland, rough grazing, heathland and moorland at higher
306 altitudes. The influence of drainage and particularly gripping in the Upper Pennines is unlikely to
307 significantly influence flooding in the lower catchment, as relatively small changes within the
308 headwaters are aggregated out by the time flood waters reach the lower catchment (Longfield and
309 Macklin, 1999). The principal flood generating mechanisms within the catchment during the
310 instrumental period (1960s-present) are persistent rainfall over a saturated catchment associated
311 with westerly and cyclonic systems and combined rainfall - snowmelt events (Macdonald, 2012).
312 The tidal limit of the Yorkshire Ouse is downstream of present day York.

313

314 The historical flood record for the City of York is one of the most detailed in Britain (Macdonald
315 and Black, 2010). The instrumental series is unique in that it provides the longest continuous

316 Annual Maximum (AM) flow series in Britain, derived from river level data obtained from adjacent
317 stageboards (all within ~200 m) at Ouse Bridge (1877-1892), Guildhall (1893-1963) and the Viking
318 Hotel (from 1963), producing an augmented stage series. These stage records were coupled with
319 data from the gauging station at Skelton (1969-present) to produce a rating curve, allowing a
320 continuous series of AM flows to be produced from 1877- (Macdonald and Black, 2010). Based on
321 the analysis of historical documents, the channel cross section has remained stable throughout the
322 city reach during the last two hundred and fifty years, as the area is confined within a walled section
323 with occasional landings (see Rocque's map of 1750). The City of York has three main bridges,
324 the most recently constructed, Skeldergate Bridge (1882) and Lendel Bridge (1863) are both new
325 bridge sites; the Ouse Bridge which was reconstructed in 1821, is the fifth bridge following Roman,
326 Viking, medieval (destroyed during the flood of 1564) and 16th century bridges. The influence of
327 the historical bridges at high flow is difficult to estimate as little information remains (other than an
328 engraving of the 1565-1810 bridge); whilst the impact of the contemporary bridges appears
329 minimal, as during the floods of 2000 some localised backing-up of flow at Ouse Bridge was
330 observed with little impact on the overall water-levels upstream and downstream. Analysis of
331 epigraphic flood markings (inscribed markings, Macdonald, 2007) inside the basement of the old
332 Merchant Venturers' Hall in central York illustrates how the city has built up over the original
333 floodplain during the centuries. Although the ground level in York has been raised, analysis of
334 historical maps and documentary accounts show little evidence of change in base river level during
335 the historical period, though bathymetric surveys post large floods suggests that bed excavation of
336 up to 2m may occur at York, as seen post 1892 and 2000 floods (Macdonald, 2004). Documentary
337 accounts provide a detailed record of flooding at York from the early eighteenth century (e.g.
338 Drake, 1736); prior to this they are more sporadic in nature, often documenting only notable large
339 floods (Macdonald and Black, 2010). A detailed discussion of the flood history and flood series
340 reconstruction is provided by Macdonald and Black (2010) and historical flood seasonality by
341 Macdonald (2012).

342

343 **3.7 River Dee**

344 The River Dee's source is in Snowdonia on the eastern slope of Dduallt (the Black Hill), the river
345 then flows down to Llyn Tegid (Lake Bala), a natural lake with an area of 1.6km², the largest
346 natural water body in the Dee catchment, before flowing eastwards through a broad valley and the
347 Vale of Llangollen, meandering northwards (Gurnell et al, 1994) through the Cheshire plain to its
348 tidal limit at Chester Weir (NRA, 1993). Llyn Tegid has a long management history, with the level
349 raised in the 1790s to support the Ellesmere Canal (constructed Thomas Telford) and subsequently

350 for water resources, in the 1960s the original Telford sluices were bypassed and the lake level
351 lowered, with new sluices constructed downstream at the confluence of the Afon (river) Tryweryn,
352 this enabled 18Mm³ storage within Llyn Tegid, permitting up to 0.235m³ for abstraction daily and
353 additional flood storage (NRA, 1993). In 1967 the construction of Llyn Celyn (6.5km²; 81Mm³)
354 was completed in the headwaters of the Afon Tryweryn, which can supply an additional flood
355 attenuation and hydropower and is operated in conjunction with the Bala Lake Scheme. In the
356 1900s and 1920s the Alwen reservoir was constructed 8 km downstream of Llyn Alwen to supply
357 water to Birkenhead, near Liverpool, with subsequent inclusion into the Bala Lake Scheme in the
358 1960s; in 1979 Llyn Brenig (3.7km²) was constructed and became part of the Dee regulation
359 scheme with a capacity of 60Mm³; both Llyn Alwen and Brenig are located on the Afron Alewn, a
360 tributary of the Dee (NRA, 1993). The geology of the upper catchment is lower Palaeozoic rocks
361 with the lower catchment (below Llangollen) Carboniferous Limestones and sandstone outcrops.
362 The land-use of the upper Dee catchment is predominantly upland grazing and moorland, while the
363 lowlands are grassland and mixed agriculture, with limited urban development, with the exception
364 of Bala, Llangollen and Chester (Marsh and Hannaford, 2008).

365

366 The city of Chester has its origins in a settlement that developed around the Roman fort of *Devia*
367 *Victrix*, quickly becoming an important port town. By the late-seventh century Chester had become
368 an important regional town, which thrived during the medieval period the town thrived, though the
369 port by the fifteenth century had become silted, with deepening of the channel in 1755 to allow
370 navigation. The Old Dee Bridge was built about 1387 (widened in 1826), following the loss of
371 several wooden bridges from flooding (1227, 1280, 1297 and 1353) and withstood the flood of 16
372 the January 1551 (Stewart-Brown, 1933), with a Letters Patent granted to the citizens on the 25 July
373 1387 by Richard II for the purpose of the construction of a bridge, following the destruction of a
374 previous bridge. The earliest account of a bridge over the Dee comes from the Domesday Book
375 (1085), which notes the bridge at Chester, though this likely follows earlier bridges and a Roman
376 fording point. A sandstone weir was built in 1093 just upstream of site of the Old Dee Bridge for
377 the Benedictine Abbey of St Werburgh's (now Chester Cathedral), to power a set of mills, which
378 were demolished in 1910, with the weir converted to producing hydroelectric from 1913-1939
379 (Historic England, 2015), today the weir maintains its role as a tidal point preventing tidal
380 transgression upstream. The rural and low population density for much of the catchment limits the
381 likely recording of events, particularly in the earlier period, were the Welsh language and an oral
382 tradition are prominent in weather recording in the uplands (Macdonald et al., 2010), as such many
383 of the records consulted focus on the lowland areas.

384

385 A long stage series is available for Chester Weir (67020) since 1894, though the weir drowns at
386 c.280m³s⁻¹, a discharge series is available for Chester Suspension Bridge since 1994, with longer
387 gauged series available from Manley Hall (1937-present) and Erbistock Rectory (1923-1970) with
388 the pre-1970 series at Manley Hall calculated from the Erbistock series, but both sites are located
389 c.50km upstream of Chester, with notable flood attenuation in the lower Dee floodplain (Marsh and
390 Hannaford, 2008), which accounts for an apparent reduction in discharge between Manley Hall and
391 Chester Weir. The estimation of discharges at Chester is challenging as there has been considerable
392 catchment management and change over the last millennium, with extensive regulation in the
393 headwaters over the last c.200 years (Lambert, 1988). As such early estimations of discharge at
394 Chester (pre c.1750) have considerable uncertainty and should be used as indicative of a large
395 flood. A series compiled for Chester Weir is presented, checked against the series for Manley Hall,
396 with notable floods being those exceeding c.325m³s⁻¹, during the instrumental series events
397 exceeding this threshold are 1899, 1946, 1964, 2000, 2004 and 2011. It is worth noting that the
398 series at Chester Weir begins just after a severe flood in 1890, as British Rainfall reported (Symons,
399 1890, 5).

400

401 3.8 River Trent, Nottingham

402 The River Trent has five major tributaries: the Tame, Soar, Ryton, Derwent and Dove, draining a
403 large section (7486 km²) of central England, with a mean annual discharge of 84.3 m³s⁻¹ at Colwick
404 (28009), approximately 5 km downstream of the city of Nottingham (Marsh & Lees, 2003).
405 Nottingham presents one of the longest and most detailed flood histories within the Britain; with
406 epigraphic markings indicating the level of the largest floods from 1852 inscribed into the abutment
407 of Trent Bridge, an AM series at Trent Bridge from 1877 until 1969, descriptive accounts since the
408 thirteenth century and a gauging record from Colwick since 1958. The wealth of records reflects
409 the prominent role the city had as a trade and commercial centre, a site of strategic military
410 importance historically and as an important bridging point. The catchment lies predominantly
411 beneath the 250 m contour (Hains & Horton, 1969), with exceptions in the Peak District near the
412 source of the Rivers Derwent and Dove at over 450 mAOD (Edwards & Trotter, 1954). Bedrock
413 varies throughout the catchment with the Peak District and higher altitudes predominantly Millstone
414 Grit and Carboniferous Limestone with lowland areas covered by superficial alluvial deposits,
415 beneath which are red sandstones and historically significant Coal Measures. Land use is varied
416 with rural hilly areas, forestry, pasture and rough grazing to the north; while arable farming
417 dominates lowland areas. There are considerable population centres, namely Birmingham located

418 on the River Tame in the upper catchment, Nottingham on the River Trent, Derby on the River
419 Derwent and Leicester on the River Soar; providing a total urbanised coverage of around 11 %
420 (Marsh & Hannaford, 2008). Precipitation is largely determined largely by elevation, with northern
421 sections of the catchment (Peak District) receiving >1000 mm a^{-1} , reducing to ~ 550 mm a^{-1} in
422 eastern areas, with an average of ~ 750 mm a^{-1} (Kings and Giles, 1997). The upper River Derwent
423 flow is modified by three important impoundment structures, the Derwent (holding $c.9.5$ Mm 3),
424 Howden ($c.9$ Mm 3) and Ladybower ($c.28.5$ Mm 3) reservoirs (Potter, 1958). Their role in reducing
425 the magnitude of flood peaks in the lower catchment at Nottingham is minor, as the proportion of
426 the catchment controlled by these reservoirs at Colwick is small $\sim 1.7\%$ (IH, 1999). The present tidal
427 limit of the Trent is Cromwell lock, ~ 25 km downstream of Nottingham.

428

429 The borough and city of Nottingham have a unique series of scrolls documenting the period AD
430 1303-1455; with the first map of Nottingham drawn by the notable cartographer John Speed in 1610
431 with subsequent maps in 1675 (Richard Hall), 1744 (Badder and Peat), 1835 (Sanderson) and 1844
432 (Drearden) detailing city development and changes to the areas adjacent to the River Trent,
433 including channel improvements (e.g. construction of the Nottingham Canal running from the River
434 Trent to the town centre in 1793). The canal construction and navigable depth of the Trent produced
435 an intensely industrialised area. The planform of the River Trent in the map of 1844 indicates
436 stability within the channel, post $c.1800$, with industrial development along the northern bank, in
437 the area historically known as 'the meadows' (Beckett, 1997). The River Trent has some of the
438 oldest channel management in Britain (pre-roman), with banking of several breaches in a series of
439 sand dunes (Spalford Bank) between Girton in Nottinghamshire through to Marton Cliff, in
440 Lincolnshire; these represent an important geomorphic structure as when breached the floodwaters
441 can travel into the Witham Valley, the city of Lincoln and subsequently into the Fens, causing
442 substantial damage (e.g. the flood of 1795, St James Chronicle, 1795). Floods breaking through the
443 defences of the Spalford Bank can be used as indicative of flood magnitude, as breaching of
444 Spalford Bank occurs at discharges of ~ 1000 m 3 s $^{-1}$ (Brown *et al.*, 2001). A detailed discussion of
445 the flood history and flood series reconstruction is provided by Macdonald (2013).

446

447 **3.9 River Severn**

448 The River Severn is the longest river in the British Isles (220 km), its source is on Plynlimon in the
449 Cambrian Mountains of mid-Wales. The major tributaries are the Vyrnwy, Clywedog, Teme, Avon
450 (Warwickshire) and Stour, with the River Wye draining into the Severn estuary. The upland areas
451 in mid-Wales are predominantly given to upland grazing and moorland, with little urban

452 development except for the towns of Newtown and Welshpool. The development of impoundment
453 structures can have a notable impact on discharges, particularly at low flow, though these are more
454 limited during high flow (Marsh and Hannaford, 2008); the most significant being Lake Vyrnwy
455 built in 1880 to supply water to the city of Liverpool (~60000 Ml) and Clywedog built in 1967
456 which supplies water to the city of Birmingham and can hold 50Ml. The lower catchment is
457 predominantly given to arable and cattle grazing, with large urban centres of Shrewsbury,
458 Worcester, Gloucester. Whilst there has been an extensive history of land-use and river
459 modification the implications on the largest flows appear limited as the impact is aggregated out, a
460 view supported by Archer (2007) when looking at the upper Severn catchment (Wales-England
461 border).

462

463 The towns of Shrewsbury, Worcester, Tewksbury and Gloucester all have a long history of
464 flooding, with each representing historically important ports on the River Severn, in addition the
465 UNESCO world heritage site at Ironbridge Gorge (an early Industrial Revolution site) is located
466 c.19km downstream of Shrewsbury. These towns were important commercial, military and religious
467 centres during the early period (Macdonald, 2006) and maintained important commercial roles
468 through to the present, with each of the docks maintaining long water level data series, the earliest
469 from 1827-present, which are currently being transcribed for further analysis. A number of bridges
470 crossed the River Severn by the fourteenth century, including at Gloucester, Worcester and
471 Bridgnorth (between Bewdley and Shrewsbury), with the earliest accounts indicating that a bridge
472 was present at Worcester in the eleventh century. Unlike most major British river systems there
473 appears to have been few losses of bridges, with most damage to the early bridges arising from
474 conflicts between the English and Welsh armies. For the purpose of this study, the site of
475 Gloucester will not be discussed in further detail, as the city and port are located on the Avon just
476 upstream of the Severn confluence, with a historical flood chronology constructed for the city by
477 Bayliss and Reed (1999). Historic flood levels have been recorded at both Worcester and
478 Shrewsbury since the late Seventeenth century, with flood levels recorded on the Watergate at
479 Worcester Cathedral since 1672, with 20 floods since marked on the wall, the most recent being the
480 flood of December 2014. During the medieval period the River Severn remained tidal beyond
481 Worcester, but the tidal limit was subsequently moved below the city with the instillation of the
482 weir at Diglis in 1844 (Herbert, 1988). To reduce the uncertainties presented by the tidal signal the
483 flood reconstruction is undertaken for Bewdley, situated between the cities of Worcester and
484 Shrewsbury and the site of the long gauged series (1921-present), an additional long series is
485 available for Welsh Bridge at Shrewsbury (1911-present).

487 A rating curve constructed from flood marks at Worcester and the gauged flows at Bewdley is used
 488 to estimate the discharges for flows before 1921 back to 1672 (seven marks), with the cross section
 489 at Worcester considered to be relatively stable through this period based on analysis of historic
 490 maps, including John Speeds' of 1610. The flood of 1795 is notable for its absence on the
 491 Watergate, Green (1796) notes the flood waters "rose to precisely those of 1672" and that a plate
 492 marking the level was added to the wall of North Parade; while the 'New Bridge' built in 1781,
 493 became jammed with ice and caused extensive local flooding. The flood is documented at
 494 Gloucester as reaching within 6 inches (15cm) of the level achieved in 1770 (Star, 1795). A number
 495 of historic floods are also documented on the Severn prior to 1672, namely the 'Duke of
 496 Buckingham's water' flood of October 1484, with the flood waters preventing his rebellion against
 497 Richard III. The earliest well documented flood is that noted in the Annals of Tewksbury, which
 498 notes the 'covering of the country beyond the banks of the River Severn from Shrewsbury to
 499 Bristol'; the years 1377-81, are noted by Abbot Boyfield as having frequent inundations (translated
 500 by Luard, 1864). The floods of 1258 and 1484 are given notional discharges of $450 \text{ m}^3\text{s}^{-1}$, as they
 501 likely exceeded the threshold and are clearly documented in reliable sources as notable events, but
 502 exact estimation of the events is difficult based on the available materials. A number of additional
 503 severe floods (1236, 1338, 1348, 1576, 1587/8, 1594, 1607, 1611 and 1620) are purported to have
 504 occurred in local chronicles produced in the nineteenth century, but original sources have not been
 505 located (e.g. Fosbroke 1819), as such estimates for these events are not included.

507 **3.10 River Thames**

508 The River Thames presents one of the most heavily managed and modified river systems within the
 509 Europe. An extensive historical chronology of flooding is available for London, but this is a
 510 particularly challenging site to reconstruct a flood series for as tidal influences are particularly
 511 strong and over the last millennium development of both banks, and loss of surface tributary
 512 systems have changed the hydraulics of the system, as such an attempt to reconstruct a complete
 513 flood history of the Thames at London would be a colossal task (see Galloway (2009) for the period
 514 1250-1450), though the historical archive is unparalleled within a British context, with over 2000
 515 accounts known. The current tidal extent of the Thames is Teddington weir/lock which dates from
 516 1811, with a gauged series from 1883-present (39001), a catchment area of 9948 km^2 and average
 517 annual rainfall of 710 mm a^{-1} . Historically, the tidal extent was a weir constructed between the Old
 518 London Bridge (1209-1831) arches, on replacing the bridge seawater could subsequently reach
 519 Teddington Lock. The Thames catchment land-use consists of extensive arable farming in the

520 headwaters and a number of urban centres upstream of London, including Reading, Swindon and
521 Oxford. The geology consist of Jurassic limestone and chalk outcrops, with thick alluvium and
522 clays inn the vales (Marsh and Hannaford, 2008).

523

524 Teddington Lock contains one of the most studied gauged series in the British Isles, with the largest
525 gauged flow that of 1894, originally estimated by Symons and Chatterton (1894) as 20135.7M
526 gal/day (equivalent to $1064 \text{ m}^3 \text{s}^{-1}$), within which a spatial analysis of the contributing tributaries and
527 the relative ranking of 1894 on these systems and throughout the catchment was undertaken. This
528 discharge was subsequently reassessed by Marsh et al. (2004) based on an extensive review of the
529 information available for the flood and the channel geometry, with a revised discharge estimate of
530 $806 \text{ m}^3 \text{s}^{-1}$. Whilst 1894 is the largest gauged flow, a number of historic floods can be attributed
531 heights relative to this event, with 1593 (substantially exceeded 1894), 1774 (about 12 inches
532 higher), 1809 (12 inches higher) and 1821 (10 inches higher) all noted as being greater than that of
533 1894 by Marsh and Harvey (2012).

534

535 The first recorded flood of the Thames is that of 9AD, though likely a tidally influenced, with the
536 first fluvial flood that of AD48, where the Thames overflowed with “waters extending through four
537 countries, 10,000 persons were drowned and much property destroyed” (Griffiths, 1969), this high
538 number of deaths appears somewhat unrealistic and reflects a wider exaggerated recording practice
539 for events in this period. The first well documented flood is that of 1091/2, which witnessed the loss
540 of the ‘Old London bridge’ (it is worth noting that several bridges across the Thames acquired this
541 name over two millennia). A stone bridge was built in 1209, replacing several earlier timber
542 structures, the channel during this period was much wider and shallower, as Jones (2008) describes
543 and as illustrated with *The Frozen Thames* by Abraham Hondius (1677) and in Claude de Jongh
544 (1632) *View of London Bridge* in which the weir beneath the arches is evident. By the late
545 seventeenth century the city of London was starting to develop its quays and docks along the banks
546 and as such confine the river, as evident in Morgan’s map of the Whole of London (1682); by John
547 Rocques map of 1746, it is evident that the channel is increasingly confined, particularly around
548 London Bridge. The map of Bacon (1868), clearly illustrates the development of the Embankment
549 reach with further constriction of the river and extensive development and expansion of the city of
550 London both up and downstream of the bridge area, which influenced the channel hydraulics, with
551 the constriction of the channel resulted in channel deepening.

552

553 The flood of 1894 is the most severe in the instrumental record, but as noted by Marsh et al., (2004)
554 a number of events prior to this have reached a greater height, including 1593, 1774, 1809 and
555 1821, a view supported by Beran and Field (1988) for the later three events; other notable floods
556 also occurred in 1555, 1742, 1765, 1768, 1770, 1795, 1852, 1875 and 1877, as identified by
557 Symons and Chatterton (1895; see Table 1) and flood levels given for some relative to 1894. An
558 analysis of the descriptive accounts indicates that the largest flood is likely to have been that of
559 1593, a view supported by Marsh et al. (2005), a notional discharge of $900\text{m}^3\text{s}^{-1}$ is used in this study
560 solely for the purpose of identifying it as the largest event, though there is considerable uncertainty
561 present in this estimation. An earlier event in 1555 appears to have resulted from heavy rains, but
562 also possible tides, though unclear. The flood of 1809 is ranked second based on the descriptive
563 account, a notional discharge of $875\text{ m}^3\text{s}^{-1}$ is used to illustrate its relative position. A number of
564 epigraphic flood marks have been identified around London; the 1774 flood mark appears to be the
565 earliest, on the wall at Radnor Gardens, Twickenham, with G.B Laffan giving the 1774 level as
566 being 0.85m higher than that of 1894, though recognising a tidal influence was present, though at
567 Windsor 1894 was considered higher (Symons and Chatterton, 1895), as such a notional discharge
568 of $850\text{ m}^3\text{s}^{-1}$, is used. The floods of 1795 and 1821 appear relatively similar in description, with
569 both appearing to be fractionally greater than 1894 in the lower catchment, as such a notional
570 discharge of $825\text{ m}^3\text{s}^{-1}$ is used for both events. It is worth noting that Beran and Field (1988),
571 considered the 1821 event to be the largest of the three events to have exceeded that of 1894. This is
572 followed by the 1894 flood at $806\text{ m}^3\text{s}^{-1}$, with the event of 1742 estimated at $750\text{ m}^3\text{s}^{-1}$, as records
573 upstream at Windsor (Griffiths, 1969) suggest it was only slightly lower than 1894, though few
574 records exist further downstream. The remaining historical floods 1768, 1770, 1852, 1875 appear
575 to be similar in magnitude, some slightly higher/lower in some reaches, but similar once past
576 Windsor, as such for the purposes of this paper are all given a notional discharge of $650\text{ m}^3\text{s}^{-1}$ based
577 on the descriptive accounts, lower than that recorded in 1947 ($714\text{ m}^3\text{s}^{-1}$) but greater than 1968 ($600\text{ m}^3\text{s}^{-1}$). The flood of 1673, whilst severe at Reading and Oxford receives little mention in the lower
579 Thames, as such no estimate is provided for the event. Channel changes, river modification and
580 uncertainties involved in estimating discharges makes the ranking of events challenging, as such
581 these are notional magnitudes based on apparent ranking of events for the area around Kingston
582 upon Thames and should only be used as indicative.

583

584 **3.11 River Ouse (Sussex)**

585 The Sussex Ouse flows south through Downs into the English Channel at New Haven, past the
586 principal settlements of Uckfield and Lewes. The catchment is predominantly rural, consisting

587 almost entirely of ground beneath 150 mAOD, with established forestry in the upper catchment.
588 Few notable impoundment structures are present within the Sussex Ouse catchment, the exceptions
589 being Ardingly Reservoir constructed in 1978 (impounding ~20km²) in the headwaters and the
590 Ashdown and Barcombe reservoirs located between the forest of St Leonards and the lowland
591 floodplain (~5 km upstream of Lewes). The tidal limit is at Barcombe Mills (~6.5 km upstream of
592 Lewes) above the confluence of the Sussex Ouse and River Uck, with mean high water 3.5 km
593 downstream of Lewes. The lower Sussex Ouse valley consists of thick alluvium overlying chalk,
594 with an underlying mixed geology in the upper catchment. Precipitation is largely determined by
595 elevation, with northern sections along the South Downs receiving ~1000 mm a⁻¹ and the coastal
596 region receiving ~730 mm a⁻¹. A long history of river management downstream of Lewes exists,
597 reflecting the active shingle spit which episodically impedes drainage of the lower Ouse through to
598 the Channel, with phases of extensive flooding and drainage documented (Brandon and Short,
599 1990; Woodcock, 2003). The numerous activities culminated in the 1790 Ouse Navigation Act,
600 which would straighten (canalise) the Sussex Ouse at various points, in addition to providing
601 drainage structures which would prevent sediment supply to the shingle spit. The eventual result of
602 the canalisation was 35km of canalisation channel, 19 locks and a 1.3km branch, with navigation up
603 to Balcombe. The consequence on the hydraulic capacity of the channel during high flow events is
604 poorly detailed, though historical accounts continue to document overbank flooding during events
605 comparable to that described by Pearce (2002) of extensive flood plain storage upstream of Lewes
606 during flooding in 2000. The town of Lewes also floods from the Winterbourne Stream, which
607 emerges from the chalk aquifer during periods of high groundwater, as such, it can flood in
608 combination with, or independently of, the Sussex Ouse.

609

610 Three bridges cross the Sussex Ouse in central Lewes: i) Cliffe Bridge, which is the oldest bridge
611 and is the site of several historical bridges in Lewes (commonly known as Ouse Bridge) which
612 probably reflects the location of a ford, ferry and roman bridge (Dunvan, 1795; Salzman, 1940); ii)
613 Willey's Footbridge (opened in 1965); and, iii) the Phoenix Causeway (a larger road bridge built in
614 the early 1970s). The modern A27 trunk road crosses the Sussex Ouse to the south of Lewes,
615 together with a railway bridge, but these have limited impact on the hydrology at Lewes. Accounts
616 detailing the repair of a bridge in Lewes exist from AD 1159, with the bridge rebuilt in 1561 and
617 repaired in 1652, both coincide with accounts of extensive flooding (Dunvan, 1795). Historical
618 accounts detail the bridges destruction in 1726 (Sawyer, 1890); with the current single stone arch
619 structure dating from 1727, with widening work undertaken in 1932 (Salzman, 1940). The adjacent
620 wharf was constructed in 1770-71 and subsequently repaired in 1802 (Salzman, 1940), suggesting

621 little change in the channel cross-section at Lewes during the intervening period; the first Ordnance
622 Survey map (1875) of Lewes shows little change in channel location and adjacent structures to the
623 present day. Inevitably the potential for channel cross section modification during the historical
624 period represents a challenge when estimating historical flows; consequently this study considers
625 only the largest floods for the period since 1772. Although intermittent records are available prior to
626 this date, less confidence can be placed in the cross sectional area of the channel at Lewes; with
627 greater confidence also placed in record completeness post c.1750, a timeframe comparable to that
628 selected in previous studies (e.g. Parent and Bernier, 2003; Macdonald, 2013). The historical
629 accounts of flooding from documentary sources provide detailed descriptive accounts of past flood
630 extents which can be converted into levels, augmenting the discharge readings from 1960 for the
631 Isfield (41006; Uck) and Gold Bridge (41005; Ouse) gauging stations (m^3s^{-1}). A detailed discussion
632 of the flood history and flood series reconstruction is provided by Macdonald (2014).

633

634 **3.12 River Exe**

635 The River Exe drains the upland regions of Dartmoor, Exmoor and the Blackdown Hills in
636 Southwest England (Fig.1), with most of the catchments underlying geology consisting of relatively
637 impermeable Carboniferous shales and slates (British Geological Survey, 1995). Exeter is the
638 principal settlement on the River Exe, with a history predating Roman times (Hoskings, 1960). The
639 city of Exeter is situated at the tidal extent, with an extensive history of human activity on the
640 floodplain (Brown 2010), including historic fording and medieval bridges, the oldest dating from
641 the end of the twelfth century; a detailed discussion of bridging at Exeter is provided by Brierley
642 (1979), which includes a discussion of historic bridge damage and maintenance closely tied to flood
643 events. Catchment land-use is predominantly agricultural and rough grazing, with limited urban
644 development. The River Exe at Exeter consists of three principal sub-catchments, the Exe flowing
645 from the north ($\sim 600\text{km}^2$), the Culm which enters the Exe just upstream of Exeter from the west
646 with a catchment area of $\sim 250\text{km}^2$ and the Creedy which flows from the east and also enters the Exe
647 just upstream of Exeter, with a catchment area of $\sim 260\text{km}^2$. The only significant impoundment
648 structure in the headwaters of the Exe is Wimbleball lake in the River Haddeo sub-catchment,
649 which was constructed in 1979 and has a volume of $\sim 21,000$ Ml and a catchment area of 29 km^2
650 (Webb and Walling, 1996). Precipitation is greatest ($>1400\text{mm a}^{-1}$) over the uplands, dropping to
651 $\sim 850\text{mm a}^{-1}$ at Exeter Airport at the coast (> 500 mAOD). The geology and relatively steep gradient
652 have resulted in a fluvial system with a flashy flood regime, a detailed discussion of channel form is
653 provided by Bennet et al. (2014), including copies of the city maps from John Hooker's map of
654 1587 through to those of the early nineteenth century, detailing the instability within the lower

655 channel with high rates of channel movement across the floodplain, with greater stability since the
656 nineteenth century.

657

658 A set of gauged records exists for the River Exe at Thorverton (45001) since 1956, ~11km upstream
659 of Exeter (Marsh and Hannaford, 2008); the Culm at Wood Mill (45003) since 1962, ~15km
660 upstream of Exeter and at Cowley (45012) on the Creedy since 1964, ~3km² upstream of Exeter.
661 These gauged series are combined to generate a single series for the site, instantaneous peak flow
662 (ipf) data is used where available, where gaps are present mean daily flow (mdf) is included, whilst
663 under-representing peak flow this provides a conservative discharge estimate, with only two years
664 recording no ipf at any station, where ipf are within 1-day of each other at the sites these are used as
665 they provide a better depiction of the highest flows. It should be noted that the two tributaries
666 (Creedy and Culm) have flashy regimes, which can produce high ipf, but may still have relatively
667 low mdf, whereas the main River Exe has a less flashy discharge regime. The highest combined
668 flow during the instrumental period is 722m³s⁻¹ (2000), which using the descriptive accounts as a
669 guide was initially estimated at 700m³s⁻¹ at Exeter. A number of well documented flood events
670 during the gauged series, particularly 1960 with subsequent events in 1974, 1985, 2000 and 2002
671 providing useful guidance on past event magnitudes at Exeter, with a number of historical events
672 being documented to a high level e.g. the flood of January 1866, for which the local newspaper *The*
673 *Exeter and Plymough Gazette* (19 January, 1866) produced a separate supplement detailing the
674 extent and impact of flood events around the country in both urban (Exeter) and rural areas (Figure
675 3). Izacke (1676) provides the first discussion of flooding at Exeter with a number of floods
676 detailed, with the first reported (unsupported) in 12AD. As at previous sites, greater confidence can
677 be placed in the discharge estimates post 1800 as channel form is more stable, with high magnitude
678 events before this date worthy of note included, but with greater uncertainty attached to their
679 estimates. As at previously described, sites the estimated discharges of the pre-instrumental series
680 are derived from the relative extent, level and damage caused by historic floods relative to the
681 associated damage and extent of floods within the gauged period.

682

683 4 SERIES COMPOSITION

684 The absence of flood record(s) for any given year does not necessarily indicate flooding did not
685 occur, simply that no record of flooding remains, or the account(s) included insufficient detail to
686 provide an estimation of the flow. However, it is likely that the largest events have been included
687 since c.AD 1750, as record density increases and becomes more systematic nearing the present,
688 with greater confidence given to high-magnitude flood event inclusion after AD 1750. The period

689 AD 1500-1749 includes a number of high-magnitude flood events, but when compared to the period
690 after 1750 it is clear that the frequency of events is considerably lower. Whilst this may be a
691 function of climatic variability, the significant growth in flood recording during the mid-eighteenth
692 century (Fig. 4) corresponding to newspaper distribution growth (Williams, 2009), suggests that an
693 increase in recording is actually detected, as such this increase in recording needs to be accounted
694 for.

695

696 Documentary flood records frequently include basic information concerning date, height or
697 magnitude of events, and often the causative mechanism i.e. rain, thaw or a combination of the two
698 (McEwen, 1987). The presence of long flood records result from several influences, namely the
699 presence of literate individuals linked to monastic, political and economic activities within the
700 cities; a detailed discussion of sources are provided in Archer (1999) and Macdonald (2004, 2007)
701 among others.

702

703 **4.1 Flood thresholds**

704 Whilst much research has focussed on the impact of land use on relatively small flood events (e.g.
705 Climent-Soler, 2009), little research, either modelled or field instrumented, has attempted to
706 undertake this analysis with rarer high-magnitude events. Wheater and Evans (2009) postulate that
707 the impact of urbanisation is potentially reduced during large flood events, whilst O'Connell et al.,
708 (2005) identify that there is very limited evidence that local changes in runoff result propagate
709 downstream. Knowledge of the conditions (climate, channel form, anthropogenic influence,
710 upstream catchment activity, etc) from which events were recorded is important in considering the
711 value of contemporary or historical flood information. When dealing with extreme flooding at
712 York, Macdonald and Black (2010) identified that there have been a number of phases of increased
713 flooding (flood rich) and periods of reduced flooding (flood poor) throughout the historical record.
714 As such, the argument has been made that once long periods are considered ($> \sim 250$ years)
715 variability becomes inescapable, and that inclusion of flood rich and flood poor periods leads to
716 more robust flood frequency estimates. The changing nature of climate and catchment land use
717 throughout the historical period may have caused many changes within the river regime, potentially
718 manifesting as 'flood rich' and/or 'flood poor' periods (Starkel, 2002; Benito and Thorndycraft,
719 2005). However at York, Macdonald and Black (2010) identified a phase of increased flooding
720 around AD 1625, but no significant change in flood frequency over the period AD 1800-2000. A
721 view supported at a European scale by Mundeljee et al., (2003), but contrasting to the findings by

722 Macklin and Rumsby (2007) when examining British upland catchments, as they identified a
723 decrease in flood frequency based on geomorphologically-inferred flood events over the last 50
724 years.

725

726 5 FLOOD INDICES (FI)

727 The issue of increased recording of floods nearing the present represents a challenge when
728 attempting to analyse long time-periods, as the availability and recording frequency increase. A new
729 method was developed to adjust the data based on its frequency and distribution over time. This
730 allows for growth in record number but does not assume linear growth. First, two distinct
731 timeframes are identified within the records AD 1200-1750 and AD 1750-2012 which are treated
732 separately. The AD 1750-2012 timeframe has considerable growth in flood recording with a 10-
733 year count rising from 0 records (AD 1752) to 22 records in AD 1968 and 1969. The FI [Equation
734 1] is calculated to determine periods of increased flood incidence in the most extreme events (flood
735 rich periods), with those years exceeding the 0.9 percentile of FI considered to represent flood rich
736 years:

737

$$738 FI_t = \bar{z}_t^{(10)} \left(1 - \frac{t}{n} \left(\frac{\max(z) - \min(z)}{n} \right) \right), \quad t = 1, 2, \dots, n \quad [Equation 1]$$

739

740 Where:

741 z the number of flood recordings in each year

742 \bar{z}^{10} the mean number of flood records within the preceding 10-year period

743 n the total number of years within the study period t

744 t the number of years after the start of the period (e.g. 1760 is 10)

745 e total number of extreme events over study period

746

747 6 SPATIAL AND TEMPORAL FLOOD VARIABILITY

748 The flood series are compiled from archival materials and previously published series for the rivers
749 Findhorn (McEwen and Werritty 2007), Tay (Werritty et al., 2006; Macdonald et al., 2006), Tweed
750 (McEwen, 1990), Tyne (Archer et al., 2007), Eden (Macdonald, 2006; Patterson and Lane, 2012),
751 Dee, Yorkshire Ouse (Macdonald and Black, 2010), Trent (Macdonald, 2013), Severn, Thames,
752 Sussex Ouse (Macdonald et al., 2014) and Exe (Fig. 5). An additional chronology for the River
753 Kent in the southern Lake District has been constructed, but is relatively short compared to those
754 presented here and is therefore not included. In each case the estimated discharges are derived from

755 historical accounts and records, where previous studies have been conducted the original archive
756 materials were considered, a detailed review of the different materials and chronologies for each
757 site is beyond the scope of this paper (please refer to the site specific studies for further information
758 where available). These series represent the sites for which the most detailed and complete
759 historical series exist; the Thames reconstruction is based at Teddington above the tidal limit, as
760 determining the influence of tidal input to the historical floods in London is challenging, though the
761 potential of the historical flood record at London is considerable (~1500 accounts to date have been
762 collected), see specific detailed comments above relating to the Thames system.

763

764 The individual flood series are compiled into grouped series at a range of spatial scales: national (all
765 sites); east (Tay, Tweed, Tyne, Ouse-Yorkshire, Trent, Thames) and west (Findhorn, Eden, Dee,
766 Trent, Severn, Exe) draining catchments; and, Wales (Dee and Severn), Scotland (Findhorn, Tay,
767 Tweed), northern (Eden, Tyne, Ouse-Yorkshire, Trent) and southern England (Thames, Exe, Ouse-
768 Sussex), permitting further detailed regional analysis (Fig. 5). The focus on relatively large
769 catchments, within a British context, inevitably constrains the generating mechanisms that are likely
770 to result in high-magnitude floods; which are likely to be either snowmelt, or persistent/heavy
771 rainfall on saturated/frozen ground, or a combination of the two (Black and Werritty, 1997); intense
772 rainfall events generally have greater impact on small catchments with high relief, although sub-
773 catchments of those studied may contain high relief, these are unlikely to result in significant flood
774 events at the sites examined. The potential role of snowmelt as a flood generating mechanism since
775 AD 1800 with the Yorkshire Ouse was examined (Macdonald 2012), the ratio of floods deriving a
776 snowmelt component were found to be consistent, though potential changes in accumulation within
777 the upper catchment may vary (no records exist of snow depth). The role of ice jamming in Britain
778 as a cause for significant flood events is limited, with only the 1814 flood on the River Tay clearly
779 exacerbated by ice floes (jamming under Smeaton's Bridge, see Macdonald et al., 2006); though
780 historical accounts identify a number of ice fairs over the period of study, that were held on several
781 of the rivers, including the Thames, Trent and Severn.

782

783 Of the sites considered within this paper, no site incorporates a large groundwater component
784 during extreme events, with the Thames and Sussex Ouse potentially including a greater
785 groundwater contribution than other sites as detailed above. The Thames catchment may experience
786 localised groundwater flooding, but this is small relative to the flows within the main channel and
787 localised within the catchment; similarly the Sussex Ouse receives limited groundwater flooding,

788 with groundwater flooding from the Winterbourne stream tributary affecting a specific area of
789 Lewes downstream of the point considered within this study.

790

791 **6.1 Flood rich and poor phases**

792 Discernible flood-rich periods are identified at a national scale, across multiple catchments and
793 within specific catchments during the last 814 years (1200-2014; Fig. 2). The Flood Index (FI – Fig.
794 5) generated for Britain correspond well to events/periods recorded elsewhere within the literature
795 as containing significant flood events e.g. c.1200s and c.1600, while long periods with a low FI, e.g.
796 drier phases c.1300-1550s, correspond well with proxy series e.g. the peat wetness record
797 (Charman, 2010).

798

799 The regional FIs (Fig. 5) show both coherent flood-rich phases (e.g. 1770s) across most catchments,
800 but also regionally specific flood rich periods (e.g. Wales, c.1883). The division of Britain east –
801 west shows similar patterns in the FI, with some subtle differences, e.g. stronger flooding signal
802 c.1770 in eastern Britain, though overall it illustrates that there are not considerable differences in
803 flooding on an east-west basis. Division into four regions provides more variability and permits an
804 assessment of spatial variability, with clear differences in FI for Scotland and Wales, with the flood
805 peak around 1883 in Wales not evidenced in Scotland and a lower FI score for the 1853 event in
806 Wales than Scotland. The northern and southern Britain divisions also show considerable
807 differences, particularly for the period since 1950, with considerably more events in the north
808 during this period. Consideration of the regional flood rich periods, as indicated by the black boxes
809 on the right vertical axis (Fig. 5) illustrates the temporal and spatial variability of flood rich periods
810 across Britain. Flood-rich periods can be determined within individual catchments, e.g. River Tay
811 AD 1567-1621 (seven floods) and River Ouse (Yorkshire) with five notable events (AD 1564,
812 1614, 1623, 1625 and 1636) (Fig. 2) and these can be identified at a national level.

813

814 National flood-rich periods are identified during the periods 1570-1640, 1850s and 2000-present,,
815 with several short flood-rich phases: 1690-1700, 1765-80, 1850s, late-1940s, and mid-1960s. High-
816 magnitude floods in the mid-to-late nineteenth century are widely documented across Britain (e.g.
817 Brookes and Glasspoole, 1928), with the period AD 1875-1885 identified as including a number of
818 years with severe floods (Marsh et al., 2005), though this period is not identified when applying a
819 0.9 percentile threshold, if the threshold is lowered to 0.8, this period appears as flood rich (Fig. 5).
820 The current flood-rich period (2000-) is of particular interest with several extreme events
821 documented in recent years, though it should be noted from a historical perspective that these are

822 not unprecedented, with several periods with comparable FI scores during the last 260 years, it
823 remains unclear at present whether the current period represents a short or long flood-rich phase. It
824 is notable that the current flood-rich phase is more evident in northern rivers than those of the south,
825 though several of the southern rivers examined recorded high flows in winter 2014. The severe
826 floods of December 2015 are not included within the series, as data are unavailable for all sites, but
827 resulted in record breaking discharges in several of the catchments, it is worth noting that gauged
828 discharges on the Eden and Tyne are the highest recorded (est. $\sim 1700 \text{m}^3 \text{s}^{-1}$) and third highest on the
829 Tweed (est. $\sim 1361 \text{m}^3 \text{s}^{-1}$; CEH, 2016), all of which are northern England catchments. The spatial
830 coherence of the FI varies, illustrating the importance of good spatial coverage, and suggests that an
831 understanding of flood rich periods needs to be undertaken first at a catchment scale, with
832 subsequent studies examining larger areas/regions. The spatial variability in the series suggests that
833 regions are behaving differently, with periods of synchronous (e.g. national 1770s) and non-
834 synchronous (e.g. regional 1920s) activity.

835

836 In the context of the long historical flood series available for mainland Europe, flooding appears to
837 be synchronous and asynchronous during different phases in comparison to the British series.
838 Benito et al. (2004) identified flood rich periods for the Tagus river in southern Spain during the
839 periods 1590–1610, 1730–1760, 1780–1810, 1870–1900, 1930–1950 and 1960–1980 (underlined
840 coinciding with British flood-rich periods at 0.9 threshold). Sheffer et al. (2008) study of the
841 Gardon river in southern France identifies several flood rich phases: 1740–1750, 1765–1786, 1820–
842 1846, 1860–1880 and 1890–1900; with Llasat et al. (2005) identifying flood-rich phases for
843 Catalonia in 1580–1620, 1760–1800 and 1830–1870. Comparison of the British FI to the historical
844 flood series presented by Glaser et al. (2010) for central Europe shows a more complex story, with a
845 number central European systems appearing to be asynchronous in relation to the British (e.g.
846 Vistula), whilst others provide similar flood-rich and -poor phases (e.g. Rhine). The flood-rich
847 phase c. 1600 identified in Britain though is identified at a central European scale from 1540–1610,
848 and the mid-late eighteenth century flood-rich phase in Britain coincides with a longer flood-rich
849 phase in central Europe from 1730–1790 (Glaser et al., 2010), the other phases identified (1640–
850 1700, the last 10 years being the exception, and 1790–1840) coincide with periods of little flooding
851 in Britain. Brazdil et al. (2005) identified a series of flood phases on the Vltava at Prague, with
852 peaks 1560–1600, c.1750, c.1825, 1840–1860, 1890, 1940–1950 and 1975–1990, again these show
853 some overlap with flood-rich periods witnessed in Britain, but also periods of little flood activity
854 e.g. 1975–1990. Wetter et al. (2011) identify a number of large floods for the Rhine: c.1350, 1560–
855 1600, c. 1740–1791, 1850–1880, 1994–2007, of the published flood series this shows good

856 comparison to the British FI. The flood peak identified c.1350, does not relate to a British FI, but
857 closer inspection of the Rhine series shows two events, AD 1342 and 1374, with examination of the
858 British series also identifying two events, AD 1352 (Dee) and 1360 (Eden), though analysis of the
859 early records is restricted because of the limited detailed data, it may suggest that this period may be
860 one for further examination as a number of descriptive accounts for which estimates were not
861 derived detail the loss of bridges during the 1340s, 1350s and 1370s for catchments around Britain.
862 Few studies have examined the flood history of Irish rivers, an account of the history of Dublin
863 (Dixon 1953) identifies a number of floods in the mid-eighteenth century (1726, 1728, 1739, 1745
864 and 1749) often associated with bridge damage/destruction, with subsequent events in 1794, 1802,
865 1807, 1851 and 1931, though it is difficult to ascertain any further information from these, accounts
866 other than event occurrence. Tyrell and Hickey (1991) identify the three most severe floods in Cork,
867 southern Ireland as 1789, 1853 and 1916, with increases in flood frequency in the 1920s, 1930s and
868 1960s. Whilst both the Tyrell and Hickey (1991) and Dixon (1953) studies provide some
869 information for Ireland it is challenging to determine whether these are small- or wide-scale flood-
870 rich periods, with the flood-rich phase in Dublin of the mid-eighteenth century occurring before that
871 in Britain, the increased frequency in Cork in both 1920s (apparent at 0.8 threshold) and the 1960s
872 and large flood of 1853 both coincide with those identified in the British FI.

873

874 7 FLOOD DRIVERS

875 During much of the Holocene, three forms of natural forcing of climate are evident: orbital (Esper et
876 al., 2012), solar (Lean, 2000; Vaquero, 2004) and volcanic (Brázdil et al., 2010), these have
877 influenced the global climate, and as such potential flood generating mechanisms. Orbital forcing
878 over the last millennium has changed little.

879

880 Solar forcing can manifest itself in a variety of different ways on flood patterns through
881 modification of the climate (Benito et al., 2004); it is notable that no flood events are recorded in
882 the British FI during the Sporer minimum (1450-1550), with relatively few accounts documenting
883 floods during this period, the exceptions being 1484 on the Severn and Trent (also a year of
884 drought) and 1545 on the Severn. The increase in high magnitude floods in central and southern
885 Europe c. AD 1700, linked to the cold and dry climate of the Late Maunder Minimum (AD 1675-
886 1720) (Mudelsee et al., 2004) have not previously been identified within British flood chronologies,
887 with the British FI identifying the period 1689-1698 as including a number of extreme events (Fig.
888 5), with two clearly identified as snowmelt events, though generally the period 1650-1750 is
889 characterised as flood-poor, with few floods recorded in Scotland, Northern or Eastern England

890 (Fig. 5). Several series (Fig. 5) indicate increased flood frequency during the late eighteenth century
891 corresponding to the Dalton minima (AD 1790-1830), with notable flooding across catchments in
892 the eight-year period AD 1769-1779, a climatic period considered to include the sharpest phases of
893 temperature variability during the ‘Little Ice Age’ (Lamb, 1995; Wanner et al., 2008). The spatial
894 and temporal variability in relation to these events may suggest that snowmelt becomes a more
895 important driver for flooding relative to heavy precipitation, suggesting that flood response to solar
896 forcing may be regionally and temporally heterogeneous (Benito et al., 2004) (Fig. 5). The flood-
897 rich phase in different catchments around Britain (except Wales) during the late sixteenth and early
898 seventeenth century corresponds to a phase of increased storminess in the North Atlantic (Lamb and
899 Frydendahl, 1991) and increased solar activity (Muscheler et al., 2007), and is evidenced in flood
900 accounts from catchments across southern and central Europe (e.g. Brazdil et al., 1999) suggesting a
901 wider flood-rich period, which relates to a particularly strong phase of positive solar forcing (Fig.
902 5). A positive significant relationship exists ($p>0.95$) between solar irradiance (Lean, 2000) and FI
903 national and North, East, Scotland, Wales regions (AD 1200-2014; Fig. 5; $p= <0.03$). A significant
904 positive correlation between Atlantic Meridional Oscillation (AMO; 1850-present; Enfield et al.,
905 2001 updated by NOAA) and national FI is identified ($p= <0.01$), with significant positive regional
906 correlations also identified for the North, South, Scotland and West FI at both annual and
907 winter/summer half years ($p= <0.01$). Analysis of dendro-chronological reconstruction of AMO
908 (Gray et al., 2004) over the last millennium identifies significant positive correlations with regional
909 FI West and FI Wales, but not for other regions, or nationally
910

911 A significant negative correlation between North, South, Scotland and national FI and winter North
912 Atlantic Oscillation Index (NAOI) over the last millennium are identified, with the West exhibiting
913 a positive correlation ($p= <0.02$; Trouet et al., 2009). These findings correspond to previous studies
914 which have attributed flood-rich phases to both positive (Dixon et al., 2006; Hannaford and Marsh,
915 2008) and negative (Macklin and Rumsby, 2007; Folland et al., 2009; Foulds et al., 2014) phases of
916 the NAOI, though these studies have used different river flow series, with those evidencing positive
917 NAOI relationships often using short instrumental series (c.1960-), conversely those evidencing
918 negative relationships have applied palaeo-historic-geomorphic flood series for several centuries.
919 This suggests that the relationship between NAOI and flooding is complex, with potentially
920 different flood generating mechanisms responding to different NAOI states, with different levels of
921 threshold of inclusion being used in the different datasets considered. The relationship identified
922 within this paper suggests that historical high magnitude floods occur during phases of negative
923 NAOI (Fig. 5), with the exception of the western catchments; specific flood-rich periods identified

924 in the British FI correspond to negative (e.g. c.1600) and positive (e.g. c.1770) phases of NAOI.
925 The significant correlations identified above indicate that warming of the Atlantic through solar
926 forcing has potentially resulted in changes to flood phases, with the presence of flood-rich phases
927 across multiple catchments suggesting abrupt changes in flood frequency/magnitude, reflecting
928 wider climatic variability, permitting an assessment of regional palaeoclimatic change (e.g.
929 Schillereff et al., 2014). This represents an interesting an important finding, with potential future
930 implications for flood type, with a warmer Atlantic potentially leading to greater potential energy
931 which may result in an increase in intense precipitation leading to high-magnitude floods affecting
932 Britain, with areas particularly vulnerable being coastal uplands in the southwest, southern Wales
933 and the Lake District, with recent notable floods (2005, 2009 and 2015) in the latter.

934

935 Aerosol optical depth was used as a proxy for volcanic forcing (Crowley and Unterman 2012), with
936 no relationship evident to the British FI, with only the Wales FI presenting a significant correlation
937 ($p=0.03$). The British FI fails to identify a relationship between large volcanic events and flooding
938 in Britain (e.g. Laki Fissure, 1784; Krakatoa, 1883 and Tarawera 1886; Fig. 5), when the threshold
939 is lowered to 0.8 percentile, these years do appear within a flood rich phase, but so does the
940 preceding year 1882, suggesting that this may have been a wetter period prior to these eruptions .
941 The clear peak in AOD following the Tambora (Indonesia) eruption of 1815 results in elevated
942 AOD for several years (Fig. 5), whilst there have been clearly documented impacts felt across
943 Europe in relation to temperature, with the ‘year without a summer’ (Oppenheimer, 2003), no
944 evidence is presented from the British flood chronologies of any associated change in flood
945 magnitude or frequency. The widespread flooding documented across much of Central Europe
946 during the winter of AD 1783-84 following the Laki fissure (Iceland) eruption is not widely
947 evidenced within British catchments (Brázil et al., 2010). Overall, there appears to be little
948 evidence in British systems of volcanic forcing influencing flood events directly during the period
949 of study.

950

951 8 SUMMARY

952 The apparent increase in flooding witnessed over the last decade appears in consideration of the
953 long term flood record not to be unprecedented, whilst the period since 2000 is considered as flood-
954 rich, the period 1970-2000 is relatively ‘flood poor’, which may partly explain why recent floods
955 are often perceived as extreme events. The much publicised (popular media) apparent change in
956 flood frequency since 2000 may reflect natural variability, as there appears to be no shift in long
957 term flood frequency (Fig. 5). In reviewing the flood series for European systems for which long

958 flood series have been reconstructed, a complex picture is identified, whilst flood rich phases appear
959 synchronous across many systems (c.1600 and 1765-1780) others show less synchronicity (1920s),
960 whilst a number of prominent flood-rich phases at a European scale appear subdued or are not
961 evident in the British FI (e.g. c.1740-1750).

962

963 The principal finding of this work is that of the strong correlation between flood-rich phases and
964 solar magnetic activity, indicating a clear driver for flooding patterns across Britain, what is still
965 unclear is the relationship between the spatial/temporal distribution of flood clusters and solar
966 activity. This work suggests that high magnitude flood-rich periods relate to negative NAOI across
967 much of the country, in western catchments with a stronger westerly airflow signal significantly
968 correlating to positive NAOI, with reasonable correspondence with previously diagnosed periods of
969 climatic variability identified from individual series from across Europe. It also identifies the
970 importance of the Atlantic Multi-decadal Oscillation as a clear correlation is shown between higher
971 North Atlantic sea temperatures and increased severe flood events across much of Britain. It is
972 worth noting that when the threshold is reduced to the 0.8 percentile of events, significant
973 correlations remain between the British FI and summer, winter, annual AMO (1850-) and NAOI
974 (Trouet et al., 2009). The inclusion of historical flood information provides a better understanding
975 of long-term flood patterns. The detection of flood-rich periods and attribution to periods of
976 climatic change are tentative. The historical records still hold a wealth of untapped information
977 within the records for which specific discharges cannot be estimated, but from which indices could
978 be extracted in the future (Barriendos & Coeur, 2004). The wealth of information presented by the
979 historical records presents valuable new information for flood risk assessment and management
980 (Kjeldsen et al., 2014); as new flood chronologies become available, more detailed and complete
981 indices based chronologies will improve the resolution and enhance understanding of flood-rich and
982 -poor periods, presenting a more complete depiction of the role of climate and extreme floods.

983

984 **Data availability**

985 Discussions are currently ongoing concerning the deposition of final datasets, this is in-part
986 constrained by the requirements of data ownership of the gauged hydrological data.

987

988 **Acknowledgements**

989 This research was in part funded by RGS-IBG/EPSRC Geographical Research Grant EPSRC 3/05
990 and supported in part by a NERC-CASE studentship to the author. The assistance of the National
991 River Flow Archive at CEH Wallingford and Scottish Environment Protect Agency in provision of

992 contemporary gauged river flow data is greatly appreciated. Thanks to Thomas Kjeldsen for reading
993 a draft of the paper. Thanks are also given to the anonymous reviewers who provided thoughtful
994 comments.

995

996 **Declaration of interests**

997 The author declares that they have no conflict of interest.

998 **REFERENCES**

999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 Acreman, M.C. Extreme Historical UK floods and maximum flood estimation, *Water Environ. J.*, 3: 404-412, 1989

Archer, D.R. Improvement in flood estimates using historical flood information on the River Wear at Durham, *British Hydrolo. Soc. Symp. First National symposium*, Hull. 1987.

Archer, D.R. *Land of Singing Waters: Rivers and Great Floods of Northumbria*. Spredden Press, Stocksfield, Northumbria, UK. 1992

Archer, N., Discharge estimate for Britain's greatest flood: River Tyne, 17th November 1771, *BHS 4th National Hydrolo. Symp.*, Cardiff. 1993.

Archer, D.R. Practical application of historical flood information in flood estimation. In: *Hydrological Extremes: Understanding, Predicting, Mitigating* (ed. by L. Gottschalk, J.-C. Olivry, D. Reed & D. Rosbjerg, 191–199. IAHS Publ .255. IAHS Press, Wallingford, UK. 1999.

Archer, D.R. *Tyne and Tide: A celebration of the River Tyne*, Daryan Press, Ovingham. 2003

Archer, D.R. Leesch F. and Harwood K., Assessment of severity of the extreme River Tyne flood in January 2005 using gauged and historical information, *Hydrol. Sci. J.*, 52: 992-1003, 2007.

Bacon, G.W. *Map of London*. G.W. Baco, London. 1868.

Barriendos, M. and Coeur, D., Flood data reconstruction in historical times from non-instrumental sources in Spain and France. In: *Systematic, Palaeoflood and Historical Data for the Improvement of Flood Risk Estimation. Methodological Guidelines* (ed. by G. Benito & V. R. Thorndycraft), 29–42. Centro de Ciencias Medioambientales, Madrid, Spain. 2004.

Bayliss, A.C. & Reed, D.W. The use of historical data in flood frequency estimation, Centre for Ecology and Hydrology (NERC), Wallingford. 2001

Beckett, J. *Nottingham: An illustrated History*, Manchester University Press, Manchester, UK, 1997

Benito, G., Diez-Herrero A. and de Villalta M.F., Response to Solar Signal in the Number of Floods Recorded for the Tagus River Basin over the Last Millennium, *Climate Change*, 66:27-28, 2004.

Benito, G., and Thorndycraft, V.R. Palaeoflood hydrology and its role in applied hydrological sciences. *Journal of Hydrology*, 313(1), 3-15, 2005

Bennett, J.A. Brown A.G. Reed S.J. Holocene fluvial geomorphology of the River Exe (UK) from archaeological and historical data and implications for urban form, *Proc. Geol. Ass.*, 125:639-648, 2014

Beran, M and Field, E.K., Maidenhead, Windsor and Eton Flood Study, *Inst. of Hydrol.*, Wallingford. p.48. 1988.

Black, A.R. and Werritty A., Seasonality of flooding: a case study of North Britain, *J Hydrol* 195:1–25, 1997.

1048 Black, A.R., Law F.M. 2004. Development and utilisation of a national web-based chronology of
1049 hydrological events. *Hydro. Sci. J.*, 49(2): 237–246.

1050

1051 Böhm, O., Jacobbeit J., Glaser R., and Wetzel K.-F., Flood History of the Bavarian Alpine Foreland
1052 since the late Middle Ages in the context of internal and external climate forcing factors, *Hydrol*
1053 and *Earth Syst. Sci.* 19:4721-4734, 2015.

1054

1055 Brázdil, R., The History of Floods on the Rivers Elbe and Vltava in Bohemia, *Erfurter Geogr.*
1056 *Studien*, 7:93–108, 1998

1057

1058 Brazdil, R. Glaser, R. Pfister, C. Dobrovolny, P. Antoine, J.M. Barriendos, M. Camuffo, D. Deutsch
1059 M. Enzi S. Guidoboni E. Kotyza O. and Rodrigo F.S., Flood events of selected European rivers in
1060 the sixteenth century, *Climate Change*, 43:239-285, 1999.

1061

1062 Brázdil, R., Pfister, C., Wanner, H., Storch, H. Von and Luterbacher, J. Historical climatology in
1063 Europe – the state of the art. *Climate Change*, 70: 363-430, 2005.

1064

1065 Brázdil, R., Kundzewicz Z.W. Benito G., Historical hydrology for studying flood risk in Europe.
1066 *Hydrol. Sci. J.*, 51:739-764, 2006.

1067

1068 Brázdil, R., Demarée G.R., Deutsch M., Garnier E., Kiss A., Luterbacher J., Macdonald N., Rohr C.,
1069 Dobrovolný P., Kolář P. and Chromá K., European floods during the winter 1783/1784: scenarios
1070 of an extreme event during the 'Little Ice Age', *Theo. and Ap. Clim.*,100 (1-2):163-189, 2010.

1071

1072 Brázdil, R., Kundzewicz Z.W., Benito G., Demarée G., Macdonald N. and Roald L.A., Historical
1073 floods in Europe in the past millennium, In: *Changes of Flood Risk in Europe*, Kundzewicz Z.W.
1074 (Ed.), Chapter 7, 121-168, IAHS Press, Wallingford, UK, 2012.

1075

1076 Brierley, J., The mediaeval Exe bridge, *Proc. Inst. Civ. Engineers*, 66: 127-139, 1979

1077

1078 British Geological Survey, Exeter, England and Wales Sheet 325. Solid and Drift Geology 1:50
1079 000. Keyworth, Nottingham, 1995.

1080

1081 Brandon, P. and Short, B., *The South-East from AD 1000*. Longman, London, 1990

1082

1083 Brookes, C.E.P. and Glasspoole J., *British floods and droughts*, Ernest Benn Ltd, London, 1928.

1084

1085 Brown, A.G., Cooper, L., Salisbury, C.R. and Smith, D.N. Late Holocene channel changes of the
1086 middle Trent: channel response to a thousand year flood record, *Geomorphology*, 39: 69-82, 2001

1087

1088 Brown, A.G., Basell, L.S., Toms, P.S., Bennett, J.A., Hosfield, R.T. and Scrivner, R.C. Later
1089 Pleistocene evolution of the Exe valley: A chronstratigraphic model of terrace formation and its
1090 implications for Palaeolithic archaeology. *Quaternary Science Reviews*, 29, (7-8), 897-912, 2010

1091

1092 Bürger, K., Seidel J., Dostal P., Glaser R., Sudhaus D. and Mayer H., Extreme floods on the 19th
1093 century in southwest Germany, *La Houille Blanche* 1-2007: 67-73, 2007.

1094

1095 Camuffo, D and Enzi S., The analysis of two bi-millennial series: Tiber and Po river floods,
1096 *Climatic Variations and Forcing Mechanisms of the Last 2000 Years*, NATO ASI Series, 41: 433-
1097 450, 1996

1098
1099 CEH (Centre for Ecology and Hydrology), Briefing note: Severity of the December 2015 floods - -
1100 preliminary analysis, Wallingford, UK, p.5, 2016
1101
1102 Charman, D.J., Centennial climate variability in the British Isles during the mid-late Holocene,
1103 *Quat. Sci. Rev.*, 29:1539-1554, 2010.
1104
1105 Charman, D.J., Blundell A., Chiverrell R.C., Hendon D., and Langdon P.G., Compilation of non-
1106 annually resolved Holocene proxy climate records: Stacked Holocene peatland palaeo-water table
1107 reconstructions from northern Britain. *Quat. Sci. Rev.*, 25, 336-350, 2006.
1108
1109 Climent-Soler, D., Holman, I. P., & Archer, D. R. Application of flow variability analysis to
1110 identify impacts of agricultural land-use change on the River Axe, southwest England. *Hydrology*
1111 *Research*, 40(4), 380-393, 2009
1112
1113 Crowley, T.J. and Unterman M. B., Technical details concerning development of a 1200-yr proxy
1114 index for global volcanism. *Earth Syst. Sci. Data*, 5:187-197, 2012.
1115
1116 Demarée, G.R., The catastrophic floods of February 1784 in and around Belgium – a Little Ice Age
1117 event of frost, snow, river ice... and floods, *Hydrol. Sci. J.*, 51:878-898, 2006.
1118
1119 Dennis, I.A. Macklin M.G., Coulthard T.J. and Brewer P.A., The impact of the October – November
1120 2000 floods on the contaminant metal dispersal in the River Swale catchment, North Yorkshire,
1121 UK, *Hydrol. Processes*, 17:1641-1657, 2003
1122
1123 Dixon, F.E., Weather in Old Dublin, *Dublin Historical Record*, 13(3-4), 94–107, 1953.
1124
1125 Dixon, H., Lawler D.M. and Shamseldin A.Y., Streamflow trends in western Britain, *Geophys. Res.*
1126 *Lett.*, 33:L1940, 2006.
1127
1128 Drake, F. *Eboracum (The history and antiques of the City of York)*, William Bowyer and Sons,
1129 London, 1736
1130
1131 Dunvan, P. *Ancient and Modern History of Lewes and Brighthelmston: In which are compressed*
1132 *the most interesting events of the county at large, under the Regnian, Roman, Saxon and Norman*
1133 *Settlements*, W. Lee, London, 1795.
1134
1135 Edwards, W. & Trotter F.M. (3rd ed.), *British Regional Geology: The Pennines and adjacent areas*,
1136 Institute of Geological Sciences (NERC), London, 1954
1137
1138 Elleder, L., Herget J., Roggenkamp T., Nießen A., Historic floods in the city of Prague – a
1139 reconstruction of peak discharges for 1481–1825 based on documentary sources. *Hydrol. Res.*, 44,
1140 202–214, 2013.
1141
1142 Enfield, D.B Mestas-Nunez, A.M. and Trimble, P.J., The Atlantic Multidecadal Oscillation and its
1143 relationship to rainfall and river flows in the continental U.S., *Geophys. Res. Lett.*, 28: 2077-2080.
1144 2001
1145
1146 Environment Agency, *Carlisle Flood Investigation Report*, Carlisle, UK, p.52, 2016
1147

1148 Esper J., Frank D.C., Timonen M., Zorita E., Wilson R.J.S., Luterbacher J., Holzkämper S., Fischer
1149 N., Wagner S., Nievergelt D., Verstege A. and Büntgen U., Orbital forcing of tree-ring data, *Nature*
1150 Climate Change

1151 Ferguson, R.C. *Testamenta karleolensia. The series of wills from the præ-reformation registers of*
1152 *the bishops of Carlisle. 1353-1386*, T. Wilson Kendal. 1893

1153

1154 Folland, C. K., Knight J., Linderholm H.W., Fereday D., Ineson S. and Hurrell J.W., *The Summer*
1155 *North Atlantic Oscillation: Past, Present, and Future.*, *J. Climate*, 22, 1082–1103, 2009.

1156

1157 Foulds, S.A., Griffiths H.M., Macklin M.G. and Brewer P.A., *Geomorphological records of*
1158 *extreme floods and their relationship to decadal-scale climate change*, *Geomorphology*, 216:193–
1159 207, 2014.

1160

1161 Galloway, J.A. *Storm flooding, coastal defence and the land use around the Thames estuary and*
1162 *tidal river c. 1250-1450*, *J. of Medieval History*, 35:171–188. 2009.

1163

1164 Garret, W., *An Account of the Great Floods in the Rivers Tyne, Tees, Wear, Eden, &c. in 1771 and*
1165 *1815*. E. Charnley, Newcastle, 1818

1166

1167 Glaser, R., Riemann D., Schönbein J., Barriendos M., Brázil R., Bertolin C., Camuffo D., Deutsch
1168 M., Dobrovolný P., van Engelen A., Enzi S., Halíčková M., Koenig S.J., Kotyza O., Limanówka D.,
1169 Macková J., Sghedoni M., Martin B. and Himmelsbach I., *The variability of European floods since*
1170 *AD 1500*, *Climatic Change*, 101:235–256, 2010.

1171

1172 Gray, S.T. Graumlich, L.J., Betancourt, J. L., and Pederson G. T. *A tree-ring based reconstruction*
1173 *of the Atlantic Multidecadal Oscillation since 1567 A.D.* *Geophys. Res. Lett.*, 31, L12205,
1174 doi:10.1029/2004GL019932. 2004

1175

1176 Green, V. *City and suburbs of Worcester*, Hulmer and Co. London. 1796

1177

1178 Griffiths, P.P. *The Two Extremes (Floods and Droughts)* Thames Conservancy. 1969.

1179

1180 Gurnell, A., Downward S. and Jones R *Channel planform change on the River Dee meanders, 1876-*
1181 *1992, Regulated Rivers: Research and Management*. 9:187–204. 1994

1182

1183 Hall, J., Arheimer B., Borga R., Brazdil R., Claps P., Kiss A., Kjeldsen T., Kriaučiūnienė Z.,
1184 Kundzewicz Z., Lang M., Llasat M.C., Macdonald N., McIntyre N., Mediero L., Merz B., Merz R.,
1185 Molnar P., Montanari A., Neuhold C., Parajka J., Perdigao R., Plavcová L., Rogger M., Salinas J.,
1186 Sauquet E., Scähr J., Szolgay J., Viglione A and Blöschl G., *Understanding Flood Regime Changes*
1187 *in Europe: A state of the art assessment*, *Hydrol. Earth Sys. Sci.*, 18: 2735–2772, 2014

1188

1189 Hains, B.A. and Horton, A. *British Regional Geology- Central England*, British Regional Geology
1190 Series, Nottingham, 1969

1191

1192 Hannaford, J. and Marsh, T.J., *High-flow and flood trends in a network of undisturbed catchment in*
1193 *the UK*, *Int. J. Hydrol.*, 28(10):1325–1338, 2008.

1194

1195 Harrison, J. *Central and Southern Scotland*, In Wheeler D. and Mayes J. (eds.), 1997, *Regional*
1196 *climates of the British Isles* Routledge, London, 1997

1197
1198 Herbert N. M. (ed) Gloucester: Quay and docks', In A History of the County of Gloucester: Volume
1199 4, the City of Gloucester, London pp. 251-258, 1988

1200
1201 Herget, J. & Meurs H., Reconstructing peak discharges for historic flood levels in the city of
1202 Cologne, Germany, Global and Planetary Change, 70, 108–116, 2010

1203
1204 Historic England. Chester Weir and Salmon Leap, LEN: 1375691.
1205 (<https://www.historicengland.org.uk/listing/the-list/list-entry/1375691>) 2015

1206
1207 Hondius, A. The Frozen Thames, Museum of London. 1677

1208
1209 Hoskins, W.G., Two thousand years in Exeter, James Townsend and Sons Ltd, Exeter, 1960

1210
1211 ICE, Historic Boarder Bridges, Institution of Civil Engineers North-East, ICE, London, p.4. undated

1212
1213 Jervoise, E., The ancient bridges of the North of England, Architectural Press, London, p.146, 1931

1214
1215 Jones, P. Historical climatology – a state of the art review, Weather, 2008.

1216
1217 Jones P. Conway D. and Briffa K. Precipitation variability and drought Hulme M. and Barrow E.
1218 (Eds.) 1997 Climates of the British Isles: Past, present and future Routledge, London, 1997

1219
1220 Jongh, C de. View of London Bridge. V&A Museum, London, 1632

1221
1222 Izacke, R., Antiquities of the City of Exeter, Richard Marriott, London, 1676

1223
1224 Kings, J., and Giles, B. The Midlands, In, Wheeler D. and Mayes J., 1997, Regional Climates of the
1225 British Isles, Routledge, London, 1997

1226
1227 Kjeldsen, T.R., Macdonald N., Lang M., Mediero L., Albuquerque T., Bogdanowicz E., Brazdil R.,
1228 Castellarin A., David V., Fleig A., Gül G.O., Kriauciuniene J., Kohnova S., Merz B., Nicholson
1229 O., Roald L.A., Salinas J.L., Sarauskiene D., Sraj M., Strupczewski W., Szolgay J., Toumazis A.,
1230 Vanneuville W., Veijalainen N. and Wilson D., Documentary evidence of past floods in Europe and
1231 their utility in flood frequency estimation, J. of Hydrol., 517: 963-973, 2014.

1232
1233 Lauder, Sir T.D. An account of the great floods of August 1829, in the province of Moray and the
1234 adjoining districts, (3rd ed. reprint 1873), J.M. Gillivray & Sons, Elgin, Scotland, 1830

1235
1236 Lamb, H.H., (2nd ed) Climate history and the modern world, Routledge, London, 1995.

1237
1238 Lamb, H.H. and Frydendahl K., Historic storms of the North Sea, British Isles and Northwest
1239 Europe, Cambridge University Press, Cambridge, UK, 1991.

1240
1241 Lambet, A. Regulation of the River Dee, Regul River, 2:293-308. 1988

1242
1243 Luard, H.R. 'Annales Monasterii de Theokesburia' Annals of Tewkesbury 1066-1263, Annales
1244 Monastici 1:36, Rolls Series 1864-1869, London. 1864.

1245

1246 Llasat, M.C., Barriendos M., Barrera A. and Rigo T, Floods in Catalonia (NE Spain) since the 14th
1247 century. Climatological and meteorological aspects from historical documentary sources and old
1248 instrumental records, *J. of Hydro.* 313:32-47, 2005.

1249

1250 Lean, J., Evolution of the Sun's Spectral Irradiance Since the Maunder Minimum. *Geophys. Res.*
1251 Lett., 27(16):2425-2428, 2000.

1252

1253 Lewin, J., Medieval Environmental Impacts and Feedbacks: The Lowland Floodplains of England
1254 and Wales, *Geoarchaeology Review Paper*, 25:267-311, 2010

1255

1256 Longfield, S.A. and Macklin, M.G., (1999) The influence of recent environmental change on
1257 flooding and sediment fluxes in the Yorkshire Ouse basin, *Hydrol. Process.*, 13:1051-1066

1258

1259 Luterbacher, J., Xoplaki E., Dietrich D., Jones P.D., Davies T.D., Portis D., Gonzalez-Rouco J.F.,
1260 von Storch H., Gyalistras D., Casty C., and Wanner H., Extending North Atlantic Oscillation
1261 Reconstructions back to 1500. *Atmos. Sci. Lett.*, 2:114-124, 2002.

1262

1263 Macdonald, N. The application of historical flood information in reassessing flood frequency in
1264 Britain, Unpublished Ph.D. thesis, University of Dundee, Dundee, UK. 2004

1265

1266 Macdonald, N., An underutilized resource: historical flood chronologies a valuable resource in
1267 determining periods of hydro-geomorphic change, In: Rowan J.S., Duck R.W. & Werritty A. (Eds.),
1268 IAHS/ICCE International Symposium on Sediment Dynamics and the Hydromorphology of Fluvial
1269 Systems [IAHS Red Book], IAHS Publ. 306:120-127, 2006

1270

1271 Macdonald, N. Epigraphic records: a valuable resource in re-assessing flood risk and long-term
1272 climate variability, *Environ. Hist.*, 12:136-140, 2007.

1273

1274 Macdonald, N., Jones C.A., Davies S. & Charnell-White C. Developing a historical climatology of
1275 Wales from English and Welsh language sources, *Weather*, 65(3):72-81. 2010

1276

1277 Macdonald, N., Trends in flood seasonality of the River Ouse (northern England), from archive and
1278 instrumental sources since AD 1600, *Climatic Change*, 110(3): 901-923, 2012

1279

1280 Macdonald, N., Reassessing flood frequency for the River Trent, Central England, since AD 1320,
1281 *Hydrol. Res.*, 44(2): 215-233, 2013.

1282

1283 Macdonald, N. Werritty A. Black A.R. and McEwen L.J., Historical and pooled flood frequency
1284 analysis for the River Tay at Perth, Scotland, *Area*, 38:34-46, 2006.

1285

1286 Macdonald, N. and Black A.R., Reassessment of flood frequency using historical information for
1287 the River Ouse at York, UK, *Hydrol. Sci. J.*, 55: 1152-1162, 2010

1288

1289 Macdonald, N., Kjeldsen T.R. Prosdocimi I. and Sangster H., Reassessing flood frequency for the
1290 Sussex Ouse through the inclusion of historical flood information since AD 1650, *Natural Hazards*
1291 and *Earth Sys. Sci.*, 14: 2817-2828 (doi:10.5194/nhessd-1-7615-2013), 2014

1292

1293 Macklin, M.G. and Rumsby, B.T., Changing climate and extreme floods in the British uplands,
1294 *Trans. Inst. of British Geog.*, 32:168-186, 2007.

1295

1296 Marsh, T.J. and Lees, M.L., Hydrological data UK: Hydrometric register and statistics 1996-2000,
1297 Centre for Ecology and Hydrology, Wallingford, Oxon, 2003

1298

1299 Marsh, T.J. and Hannaford, J., UK Hydrometric register, Centre for Ecology and Hydrology,
1300 Wallingford, Oxon, 2008

1301

1302 Marsh, T. and Harvey, C. L. The Thames flood series: a lack of trend in flood magnitude and a
1303 decline in maximum levels. *Hydrol Res*, 43: 203-214. 2012

1304

1305 Marsh, T.J., Greenfield B.J. and Hannaford J., The 1894 Thames flood – a reappraisal, *Proceedings*
1306 of the Institution of Civil Engineers: Water Management, 158:103-110, 2005.

1307

1308 Maxwell, H Sir. The Chronicle of Lanercost, 1272-1346 (translated with notes), J. Maclehose,
1309 Glasgow. p.416. 1913.

1310

1311 McEwen, L.J., Sources for establishing a historic flood chronology (Pre 1940) within Scottish River
1312 Catchment, *Scottish Geographical Mag.*, 103:132-140, 1987.

1313

1314 McEwen, L.J., Extreme rainfall and its implications for flood frequency: A case study of the Middle
1315 River Tweed Basin, Scotland. *Trans. Inst. British Geog.* 147:287-298. 1989

1316

1317 McEwen, L.J., The establishment of a historical flood chronology for the River Tweed catchment,
1318 Berwickshire, Scotland, *Scottish Geographical Mag.*, 106: 37-47, 1990.

1319

1320 McEwen, L.J. and Werritty, A., 'The Muckle Spate of 1829': the physical and societal impact of a
1321 catastrophic flood on the River Findhorn, Scottish Highlands, *Trans. Inst. British Geog.*, 32:66-89,
1322 2007.

1323

1324 Meteorological Office (UK Annual Rainfall web-site), Accessed 2002,
1325 <http://www.met-office.gov.uk/climate/uk/averages/images/RainAnnual6190.gif>, 2002

1326

1327 Morgan, W. Morgan's Map of the Whole of London in 1682, British History
1328 Online <http://www.british-history.ac.uk/no-series/london-map-morgan/1682/map> [accessed 6 June
1329 2016]. 1682

1330

1331 Mudelsee, M. Börngern, M., Tetzlaff, G. and Grünwald, U., No upward trends in the occurrence of
1332 extreme floods in central Europe, *Nature*, 425:166-170, 2003.

1333

1334 Mudelsee, M. Börngern M., Tetzlaff G. & Grünwald U., Extreme floods in central Europe over the
1335 last 500 years: Role of cyclone pathway "Zugstrasse Vb", *J. Geophys. Res.*, 109: D23101, 2004.

1336

1337 Muscheler, R., Joos, F. Beer J., Müller S.A., Vonmoos M., and Snowball I., Solar activity during
1338 the last 1000 yr inferred from radionuclide records. *Quat. Sci. Rev.*, 26:82-97, 2007.

1339

1340 NERC, Flood Studies Report, Institute of Hydrology, Wallingford, 1975

1341

1342 Newson, M.D. Flooding and flood hazard in the United Kingdom, p59, 1975

1343

1344 NRA, The Dee Regulation Scheme. National River Authority. Cardiff. p4. 1993

1345

1346 O'Connell, P.E., Beven, K. J., Carney, J. N., Clements, R. O., Ewen, J., Fowler, H.,
1347 Harris, G. L., Hollis, J., Morris, J., O'Donnell, G. M. Packman J. C., Parkin A., Quinn P. F.,
1348 Rose S. C. Shepherd M. and Tellier S. Review of impacts of rural land use and management on
1349 flood generation Impact study report R&D Technical Report FD2114/TR 2004, Joint Defra/EA
1350 Flood and Coastal Erosion Risk Management R&D Programme, London, p.142, 2004

1351

1352 Oppenheimer, C., Climatic, environmental and human consequences of the largest known historic
1353 eruption; Tambora volcano (Indonesia) 1815, *Prog. Phys. Geog.*, 27:230-259, 2003.

1354

1355 Parent, E. and Bernier, J. Bayesian POT modelling for historical data, *J. Hydrol.*, 274:95-108, 2003

1356

1357 Parry, S., Barker, L., Prosdocimi, I., Lewis, M., Hannaford, J., Clemas, S. Hydrological summary
1358 for the United Kingdom: December 2015. Wallingford, UK, NERC/Centre for Ecology &
1359 Hydrology, 12pp. 2016

1360

1361 Patterson, I. and Lane, S., The relationship between Lamb weather types and long-term changes in
1362 flood frequency, *River Eden*, UK. *Int. J. Clim.* 32 :1971-1989. 2012

1363

1364 Payne, P.L. The Hydro Aberdeen University Press, Aberdeen, 1988

1365

1366 Payrastre, O., Gaume, E., Andrieu, H., Usefulness of historical information for flood frequency
1367 analyses: developments based on a case study. *Water Res. Res.*, 47, W08511,
1368 doi:10.1029/2010WR009812, 2011.

1369

1370 Pearce, J. Hydrologist experiences in times of flood: Sussex 2000, BHS 8th National Hydrology
1371 2000, Symposium, Birmingham, 2002

1372

1373 Popper, W. 1951. *The Cairo Nilometer*, The University of California Press, Berkley, 269 pp

1374

1375 Rocque, J., *Map of London*, Pine & Tinney, London 1746

1376

1377 Rocque, J., A plan of the City of York; In Murray H. (1997) Scarborough, York and Leeds: The
1378 town plans of John Cossins 1697-1743, Yorkshire Architectural and York Archaeological Society
1379 Ebor Press, York p.53 1750

1380

1381 Roy, M.G. *Highland and Islands Scotland* In Wheeler D. and Mayes J. (eds.) 1997 *Regional*
1382 *climates of the British Isles* Routledge, London, 1997

1383

1384 Rumsby, B. T. and Macklin, M. G. Channel and floodplain response to recent climate change: the
1385 Tyne basin, northern England. *Earth Surf. Processes Landf.* **19**, 499–515. 1994

1386

1387 Salzman, L.F., The borough of Lewes: Introduction and history, *A History of the County of Sussex:*
1388 Volume 7: The rape of Lewes, pp. 7-19, 1940.

1389

1390 Schillereff, D.N. Chiverrell, R.C., Macdonald N., Hooke J.M., Flood stratigraphies in lake
1391 sediments: a review, *Earth Sci. Reviews*, 135: 17-37, 2014.

1392

1393 Sheffer, N. A., Rico, M., Enzel, Y., Benito, G., and Grodek, T. The palaeoflood record of the
1394 Gardon river, France: a comparison with the extreme 2002 flood event. *Geomorphology*, 98(1), 71-
1395 83, 2008

1396
1397 Smith, K., and Tobin, G., *Human Adjustment of the flood hazard*, Longman, London, 1979
1398
1399 St James Chronicle, 1795, 14-17 February, Issue 5769, London, England
1400
1401 Star, 1795, 23 February, Issue 2032, London, England
1402
1403 Star, 1797, 26 October, Issue 2868, London, England
1404
1405 Starkel, L. Change in the frequency of extreme events as the indicator of climatic change in the
1406 Holocene (in fluvial systems). *Quatern. Int.*, 91(1), 25-32, 2002
1407
1408 Stewart Brown, R., *The Old Dee Bridge at Chester*. *J. Chester Arch. Soc.* NS30 64-65. 1933
1409
1410 Symons, G.R. *British Rainfall 1890*, G. Shield, London, p.5 1891
1411
1412 Symons, G. J. and Chatterton M. A. The November floods of 1894 in the Thames valley. *Quarterly*
1413 *Journal of the Royal Meteorological Society*, 12, 189–208. 1895
1414
1415 The Exeter and Plymouth Gazette, 19 January 1866. Page 9. Supplement title: Great Flood, London
1416 © The British Library Board. 1866
1417
1418 Todd, B. Macdonald, N. and Chiverrell R.C. Revision and extension of the composite Carlisle
1419 rainfall record, NW England: 1757-2012, *Int. J. Clim.*, 35: 3593-3607. 2015.
1420
1421 Trouet, V., Esper, J., Graham, N.N.E., Baker, A., Scourse, J.D. and Frank, D.C., Persistent Positive
1422 North Atlantic Oscillation Mode Dominated the Medieval Climate Anomaly, *Science*,
1423 324(5923):78-80, 2009.
1424
1425 Tyrrell, J.G. and Hickey, K.J., A flood chronology for Cork city and its climatological background,
1426 *Irish Geography*, 24(2), 81–90, 1991.
1427
1428 Vaquero, J.M., Solar Signal in the Number of Floods Recorded for the Tagus River Basin over the
1429 Last Millennium, *Climate Change*, 66:23-26, 2004.
1430
1431 Wanner, H., Beer J., Bütkofer J., Crowley T.J., Cubasch U., Flückiger J., Goosse H., Grosjean M.,
1432 Joos F., Kaplan J.O., Küttel M., Müller S.A., Prentice I.C., Solomina O., Stocker T.F., Tarasov P.,
1433 Wagner M., Widmann M., Mid- to Late Holocene climate change: an overview, *Quat. Sci. Rev.* 27:
1434 1791–1828, 2008.
1435
1436 Webb, B.W. Walling D.E., Longtern variability in the thermal impact of river impoundment and
1437 regulation, *Appl Geogr*, 16(3):211-223, 1996
1438
1439 Werritty, A., Paine J., Macdonald N., Rowan J.S. & McEwen L.J., Use of proxy flood records to
1440 improve estimates of flood risk: lower River Tay, Scotland, *Catena*, 66:107-119, 2006.
1441
1442 Wetter, O., Pfister, C., Weingartner, R., Luterbacher, J., Reist, T., and Trösch, J., The largest floods
1443 in the High Rhine basin since 1268 assessed from documentary and instrumental evidence, *Hydrol.*
1444 *Sci. J.*, 56:5, 733-758, 2011
1445

1446 Wheater, H., & Evans, E. Land use, water management and future flood risk. *Land Use Policy*, 26,
1447 S251-S264. 2009

1448

1449 Williams, A. and Archer, D., The use of historical flood information in the English Midlands to
1450 improve risk assessment, *Hydrol. Sci. J.*, 47:67-76, 2002.

1451

1452 Williams, K., *Read All About It!: A History of the British Newspaper*, Routledge, Abingdon, UK
1453 p.422, 2009.

1454

1455 Woodcock, A., The archaeological implications of coastal change in Sussex. p.1-16 in Rudling, D.,
1456 (ed.), *The Archaeology of Sussex to AD2000*. Heritage, University of Sussex, 2003.

1457

1458

1459




Figure 1: Catchments for which historical flood reconstruction has been undertaken, where a county is included in brackets multiple catchments exhibit the same name.




Figure 2: Historical flood chronologies for sites across Britain, showing events that exceed the 0.9 percentile (based on the instrumental record; river discharges are given as $\text{m}^3 \text{s}^{-1}$). River chronologies (l-r) Findhorn; Tay; Tweed; Tyne; Eden; Ouse (Yorkshire); Dee (Wales); Trent; Severn; Thames; Ouse (Sussex); Exe; and Flood Indices (Britain) 1200-2014.




Figure 3: Example of a newspaper supplement detailing the extent of localised flooding of the River Exe, January 1866. © The British Library Board, The Exeter and Plymouth Gazette, 19 January 1866. Page 9. Supplement title: Great Flood




Figure 4: Number of floods with a recorded/estimated discharge exceeding the 0.9 threshold.




Figure 5: Historical flood chronologies (grey) by region and associated flood-rich periods (black): Britain (1200-2012); West Britain FI; East Britain FI; Scotland FI; Wales FI; Northern England FI; Southern England FI; NAO reconstruction (with 10-year running mean; Luterbacher et al., 2002), extended with CRU data; winter NAO (Trouet et al., 2009); solar magnetic (MeV, Muscheler et al., 2007); AMO grey (Gray, 2004) and black (Enfield 2001); annual stacked peat water level (10-year running mean; Charman et al., 2006); volcanic signal derived from aerosol optical depth (AOD; Crowley and Unterman, 2012) and national flood phases, using a 0.9 threshold (black) and 0.8 (grey).