

**VWC of rice and  
spatial  
characteristics  
analysis in China**

L. J. Zhang et al.

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

# Determination of virtual water content of rice and spatial characteristics analysis in China

**L. J. Zhang, X. A. Yin, Y. Zhi, and Z. F. Yang**

State Key Laboratory of Water Environmental Simulation, School of Environment,  
Beijing Normal University, Beijing, China

Received: 23 December 2013 – Accepted: 27 December 2013 – Published: 22 January 2014

Correspondence to: X. A. Yin (yinxinan@bnu.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

## Abstract

China is a water-stressed country, and agriculture consumes the bulk of its water resources. Assessing the virtual water content (VWC) of crops is one important way to develop efficient water management measures to alleviate water resources conflicts among different sectors. In this research, the VWC of rice, as a major crop in China, was assessed and the spatial characteristics were analyzed. In addition to the calculation of green, blue and grey water – the direct water in VWC – the indirect water use of rice was also calculated, using the Input–Output model. The percentages of direct green, blue, grey and indirect water in the total VWC of rice in China were 43.8, 28.2, 27.6, and 0.4 %. The total VWC of rice generally showed a three-tiered distribution, and decreased from southeast to northwest. The higher values of direct green water of rice were mainly concentrated in Southeast and Southwest China, while these values were relatively low in Northwest China and Inner Mongolia. The higher direct blue water values were mainly concentrated in the eastern and southern coastal regions and Northwest China, and low values were mainly concentrated in Southwest China. Grey water values were relatively high in Shanxi and Guangxi provinces and low in Northeast and Northwest China. The regions with high values for indirect water were randomly distributed but the regions with low values were mainly concentrated in Northwest and Southwest China. For the regions with relatively high total VWC the high values of blue water made the largest contribution, although for the country as a whole the direct green water is the most important contributor.

## 1 Introduction

The term *virtual water* was first proposed by Allan (1994) and defined as the water embodied in the traded products. In water-stressed regions, limited water resources should be used efficiently by not allocating the majority of resources to the production of water-intensive products, and should be made available for other economic purposes

HESSD

11, 1047–1072, 2014

### VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that can contribute more to the regional value-added by consuming less water (Allan, 2002; Chapagain and Hoekstra, 2008). Assessing the virtual water content (VWC) of products is the basis for developing such water resource management practices.

Hoekstra and Hung (2002) estimated the VWC of crops for many countries of the world. In their research, the crop VWC was determined through estimating the accumulated crop evapotranspiration over the growing period, and the VWC was not divided into subtypes. To better understand the VWC of crops, many scientists divide the VWC into subtypes and calculate them separately. The calculation of green (effective precipitation) and blue water (irrigation water withdrawn from ground or surface water) was first proposed in the studies of crop VWC. Research has been performed at global, national, provincial, and river basin scales. For example, Rost et al. (2008) made a global estimate of agricultural green and blue water consumption. Siebert and Döll (2010) computed the green and blue VWC of crops at a global scale, and found the global average VWC of cereal crops was  $1109 \text{ m}^3 \text{ t}^{-1}$  of green water and  $291 \text{ m}^3 \text{ t}^{-1}$  of blue water. Scientists have added the grey water to the VWC, defined as freshwater that is required to assimilate the pollutant load based on natural background concentrations and existing ambient water quality standards. Chapagain et al. (2006) first calculated the grey water in VWC of crops, finding that the global VWC of rice was an average of  $1325 \text{ m}^3 \text{ t}^{-1}$  and, further, that grey water occupied about 8% of the total VWC (Chapagain and Hoekstra, 2011). Mekonnen and Hoekstra (2011a) quantified grey VWC of global crop productions for the period 1996–2005, and found that green, blue, and grey water accounted for 78, 12, and 10% in the total VWC of crops.

China is one of the world's 13 most water-poor countries (Yu et al., 2006). Per capita use of water resources of China is only  $2300 \text{ m}^3$  (by population in 1997), less than the 1/4 of the world per capita consumption. Agriculture is the largest water user in China, accounting for nearly 70% of total water withdrawals (Ministry of Water Resources, 2012). Studies on the VWC of crops in China are relatively limited. Liu et al. (2007) through the perspective of crop water productivity estimated the virtual water use of winter wheat. Sun et al. (2013a) calculated the average VWC of wheat and maize for

## HESSD

11, 1047–1072, 2014

### VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





where  $VWC_{total}$  is the total VWC of crop ( $m^3 kg^{-1}$ );  $VWC_{indirect}$  is the indirect water use of the crop ( $m^3 kg^{-1}$ );  $VWC_{direct}$  is the direct water use of the crop ( $m^3 kg^{-1}$ );  $VWC_{direct, green}$  is the direct green water use ( $m^3 kg^{-1}$ );  $VWC_{direct, blue}$  is the direct blue water use ( $m^3 kg^{-1}$ ); and  $VWC_{direct, grey}$  is the direct grey water use ( $m^3 kg^{-1}$ ).

### 2.1.1 Indirect water of crops

The Input–Output model represents the monetary trade of products and services among different sectors of an economic system (Leontief, 1941), and is adopted to calculate the indirect virtual water of crops supplied by each economic sector. The calculations are as follows (Chen, 2000; Kanada, 2001; Zhao et al., 2009; Zhang et al., 2011):

#### (1) Direct consumption coefficient matrix

The input-output table is used because it reflects the contact of the material and technical. This contact is reflected through the direct consumption coefficient.

$$\mathbf{A} = |a_{ij}| = |x_{ij}/x_i| \quad (2)$$

where  $\mathbf{A}$  is the direct consumption matrix;  $a_{ij}$  is the direct consumption coefficient;  $x_{ij}$  is the monetary volume of products from sector  $j$  consumed by sector  $i$  in its production process (RMB); and  $x_i$  is the output of sector  $i$  (RMB).

#### (2) Complete consumption coefficient matrix

Compared with the direct consumption coefficient, the complete consumption coefficient can more accurately measure the direct and indirect costs (the sum of the direct and indirect costs is completely consumed) of products or services from other sectors.

$$\mathbf{B} = |b_{ij}| = (\mathbf{I} - \mathbf{A})^{-1} - \mathbf{I} \quad (3)$$

where  $\mathbf{B}$  is the complete consumption coefficient matrix;  $b_{ij}$  is the complete consumption coefficient; and  $\mathbf{I}$  is a unit diagonal matrices.

#### (3) Water use coefficient

To account for indirect water, it is necessary to compute the water use coefficient of different sectors, which is the water needed to produce one monetary unit (RMB).

$$DWC_i = w_j / x_i \quad (4)$$

5 where  $DWC_i$  is the direct water coefficient of sector  $i$  ( $m^3 RMB^{-1}$ ) and  $w_j$  is the direct water consumption of sector  $i$  ( $m^3$ ).  $DWC_i$  is the amount of direct water intake to produce one monetary unit of production.

#### (4) Indirect water of agriculture

10 Indirect water consumption of agriculture is the amount of total water input from other sectors.

$$VW_{\text{indirect}}^a = \sum_{i=1}^n (DWC_i \cdot b_{ia}) \cdot C_a \quad (5)$$

15 where  $VW_{\text{indirect}}^a$  is the indirect water consumption of agriculture ( $m^3$ );  $b_{ia}$  is the complete consumption coefficient of sector  $i$  for agriculture; and  $C_a$  is the total consumption of agriculture (RMB).

#### (5) Indirect water consumption of a crop

$VWC_{\text{indirect}}^i$  ( $m^3 kg^{-1}$ ) is calculated according to the proportion of indirect water use of crop  $i$  in the total indirect water consumption of agriculture.

$$20 \quad VWC_{\text{indirect}}^i = \frac{VW_{\text{indirect}}^a \alpha_i}{SA_j \cdot Y} \quad (6)$$

where  $VWC_{\text{indirect}}^i$  is the indirect water consumption of crop  $i$  ( $m^3 kg^{-1}$ );  $\alpha_i$  is the proportion of indirect water use of crop  $i$  in the total indirect water consumption. Because of lack of data, we assume that the planting cost is proportional to the indirect water use.  $SA_j$  is the sown area of crop  $i$  (ha); and  $Y$  is the crop yield per unit area ( $kg ha^{-1}$ ).

## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Thus  $\alpha_j$  can be calculated as follows:

$$\alpha_j = \frac{PC_j \cdot SA_j}{\sum_{i=1}^n (PC_i \cdot SA_i)} \quad (7)$$

where  $PC_j$  is the planting cost of crop  $j$  per unit area ( $\text{RMBha}^{-1}$ ).

### 2.1.2 Direct green water of crops

Direct green water use is the lesser of potential crop evapotranspiration and effective precipitation. Effective precipitation is defined as the amount of precipitation that enters the soil and will be available in the soil for crop growth (Sun et al., 2013b).

$$\text{VWC}_{\text{direct, green}} = \frac{10 \min(\text{ET}_c, P_e)}{\gamma} \quad (8)$$

where  $\text{ET}_c$  is the crop evapotranspiration during the growing period (mm) and  $P_e$  is the effective precipitation over the crop growing period (mm), calculated by the CROPWAT model using monthly climatic data (mm) (Clarke, 1998; FAO, 2003).

Crop evaporation during the growing period is calculated as follows (Allen et al., 1998):

$$\text{ET}_c = \text{ET}_o \cdot k_c \quad (9)$$

where  $k_c$  is the crop coefficient, reflecting the differences in physical and physiological factors between the actual and reference crops and  $\text{ET}_o$  is the soil evaporation of the reference underlying surface ( $\text{mmd}^{-1}$ ), calculated by the FAO Penman–Monteith formula (Allen et al., 1998).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

### 2.1.3 Direct blue water of crops

The direct blue water of a crop is calculated using the actual irrigation water consumption,  $I_c$ . The direct blue water is calculated according to the proportion of irrigation water consumption of crop  $i$  in the total irrigation water consumption of the irrigation district (Sun, 2013b).

$$WVC_{\text{direct, blue}}^i = I_c^i / Y = \frac{W_A \beta_i}{SA_i \cdot Y} \quad (10)$$

where  $I_c^i$  is the irrigation water consumption of crop  $i$  per unit area ( $\text{m}^3 \text{ha}^{-1}$ );  $W_A$  is the actual total irrigation water consumption of the irrigation district ( $\text{m}^3$ ) and  $\beta_i$  is the proportion of irrigation water use of crop  $i$  in the total irrigation water consumption of the irrigation district.  $\beta_i$  can be calculated as follows:

$$\beta_i = \frac{(ET_c^i - P_e^i) \cdot SA_i}{\sum_{i=1}^n [(ET_c^i - P_e^i) \cdot SA_i]} \quad (11)$$

where  $ET_c^i$  is the crop evapotranspiration of crop  $i$  during the growing period (mm);  $P_e^i$  is the effective precipitation over the crop  $i$  growing period (mm); and  $SA_i$  is the sown area of crop  $i$  (ha).

### 2.1.4 Direct grey water of crops

In this study we quantify direct grey water related to nitrogen use only. The direct grey water is calculated by multiplying the fraction of nitrogen that leaches or runs off by the nitrogen application rate and dividing this by the difference between the maximum acceptable concentrations of nitrogen and the natural concentration of nitrogen in the receiving water body and by the actual crop yield (Mekonnen and Hoekstra, 2011a).

HESSD

11, 1047–1072, 2014

**VWC of rice and  
spatial  
characteristics  
analysis in China**

L. J. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Because of lack of data, the natural nitrogen concentrations were assumed to be 0. On average, 10% of the applied nitrogen fertilizer is lost through leaching (Chapagain et al., 2006). The maximum value of nitrate in surface and ground water recommended by the United States Environmental Protection Agency is  $10 \text{ mgL}^{-1}$  (Chapagain et al., 2006).

$$\text{VWC}_{\text{direct, grey}} = \frac{N_c \cdot 10\%}{10 - N_n} / Y \quad (12)$$

where  $N_c$  is the amount of nitrogen fertilizer consumption per hectare ( $\text{g ha}^{-1}$ ) and  $N_n$  is the natural concentration of nitrogen in the receiving water body ( $\text{mgL}^{-1}$ ).

## 2.2 Data

The 2007 climate data for 29 regions, including monthly average maximum temperature, monthly average minimum temperature, relative humidity, wind speed, precipitation, and sunshine hours, are taken from the National Climatic Center (NCC) of the China Meteorological Administration (CMA, 2011). The agricultural data, including crop yield and sown area, are taken from the China Agricultural Yearbook (Ministry of Agriculture of the People's Republic of China, 2008). The average amount of fertilization of crops per unit area is taken from Li et al. (2010). The irrigation water consumption and irrigation water supply for the 29 regions are taken from the Water Resources Bulletins (2007) of 29 regions. Water data for various sectors in the regions are from the Statistical Yearbooks (2008) of 29 regions. The IO data for the 29 regions come from the official IO tables (2007) of 29 regions.

# HESSD

11, 1047–1072, 2014

## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 3 Results

### 3.1 Indirect water of rice

The  $VWC_{\text{indirect}}$  of rice varied between 0.001 and  $0.010 \text{ m}^3 \text{ kg}^{-1}$ . The average  $VWC_{\text{indirect}}$  was  $0.004 \text{ m}^3 \text{ kg}^{-1}$ . The input to agriculture in relatively underdeveloped regions was relatively small; relatively developed regions invest more money in agriculture. Northwest and Southwest China are relatively underdeveloped with relatively low  $VWC_{\text{indirect}}$  of rice. Beijing, Guangdong, and Shanghai are the three most developed regions and had the highest  $VWC_{\text{indirect}}$  of rice in 2007 (Fig. 1). We found that  $VWC_{\text{indirect}}$  of rice is directly related to the degree of regional economic development.

Of all sectors, the sector of forestry, animal husbandry, and fishery contributed the most  $VWC_{\text{indirect}}$  of rice, accounting for 52.2%. The electricity and heat sector contributed 21.6% and the chemical industry sector contributed 9.6%. The rest of the sectors contributed less.

### 3.2 Direct green water of rice

The regional differences in  $VWC_{\text{direct, green}}$  for rice were significant, owing to differences in climatic conditions and crop yields. The  $VWC_{\text{direct, green}}$  of rice for 29 regions in 2007 ranged from 0.10 to  $0.90 \text{ m}^3 \text{ kg}^{-1}$ . The average  $VWC_{\text{direct, green}}$  of rice was  $0.59 \text{ m}^3 \text{ kg}^{-1}$ . The regions with higher  $VWC_{\text{direct, green}}$  values were concentrated in the Southeast China and Southwest China (Fig. 2). The high  $VWC_{\text{direct, green}}$  in these regions is a result of high ratio between effective precipitation and rice yield. The regions in the Southeast China and Southwest China had relatively low rice yields, much lower than the national average, and relatively high effective precipitation, more than 400 mm. For example, the  $VWC_{\text{direct, green}}$  of rice was more than  $0.80 \text{ m}^3 \text{ kg}^{-1}$  in Hainan (Southeast China), Guangxi (Southwest China), and Yunnan (Southwest China). The  $VWC_{\text{direct, green}}$  of rice in Guizhou (Southwest China) was also relatively high, more

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

than  $0.8 \text{ m}^3 \text{ kg}^{-1}$ . This is mainly because the effective precipitation over the growing period of rice in this region was more than 590 mm, although rice yield in this region was higher than the national average.

The  $\text{VWC}_{\text{direct, green}}$  of rice in Northwest China and Inner Mongolia were relatively low (Fig. 2). Because of high yields and low effective precipitation. Xinjiang (Northwest China) had the lowest  $\text{VWC}_{\text{direct, green}}$  ( $0.10 \text{ m}^3 \text{ kg}^{-1}$ ) because the rice yield in Xinjiang was 1.32 times higher than the national average and the effective precipitation over the growing period was small, less than 90 mm.

### 3.3 Direct blue water of rice

The differences in blue water requirements, actual irrigation water consumption, and rice yields resulted in significantly different  $\text{VWC}_{\text{direct, blue}}$  values between regions. The  $\text{VWC}_{\text{direct, blue}}$  values ranged from 0.07 to  $1.65 \text{ m}^3 \text{ kg}^{-1}$ , and the average was  $0.42 \text{ m}^3 \text{ kg}^{-1}$ . The regions with higher  $\text{VWC}_{\text{direct, blue}}$  of rice were mainly concentrated in the eastern and southern coastal regions and in Northwest China (Fig. 3). The  $\text{VWC}_{\text{direct, blue}}$  of rice was high in the municipalities of Beijing, Tianjin, and Shanghai; more than  $0.83 \text{ m}^3 \text{ kg}^{-1}$ . These three municipalities have developed economies with relatively more developed agricultural irrigation system, so irrigation water consumption was relatively larger than that of other regions. Regions in Northwest China had higher  $\text{VWC}_{\text{direct, blue}}$  values, perhaps because the effective precipitation in these regions was limited, making it necessary to increase the irrigation water supply for crops.

The  $\text{VWC}_{\text{direct, blue}}$  values for rice in Southwest China were relatively low (Fig. 3). The  $\text{VWC}_{\text{direct, blue}}$  values in Chongqing, Guizhou, Sichuan, and Yunnan were less than  $0.16 \text{ m}^3 \text{ kg}^{-1}$  because in these four regions the effective precipitation can almost meet the water requirements of rice, so the actual irrigation water consumption was limited.

### 3.4 Direct grey water of rice

The  $VWC_{\text{direct, grey}}$  values of rice ranged from 0.21 to  $0.64 \text{ m}^3 \text{ kg}^{-1}$  with an average  $0.37 \text{ m}^3 \text{ kg}^{-1}$ . Regional differences in  $VWC_{\text{direct, grey}}$  for rice were insignificant (Fig. 4). Because nitrogen use is similar between regions,  $VWC_{\text{direct, grey}}$  mainly depends on rice yield. The rice yield of regions in Northeast China and Northwest China were relatively high, making the  $VWC_{\text{direct, grey}}$  of rice in Northeast China and Northwest China relatively low. The highest  $VWC_{\text{direct, grey}}$  values for rice were in Shanxi and Guangxi, because the rice yields of Shanxi and Guangxi in 2007 were the two lowest of all the regions studied.

### 3.5 Total VWC of rice

The  $VWC_{\text{total}}$  values of rice for 29 regions are calculated and shown in Fig. 5. There were large differences in  $VWC_{\text{total}}$  between regions, with values ranging from 0.80 to  $2.59 \text{ m}^3 \text{ kg}^{-1}$ . The average  $VWC_{\text{total}}$  of rice was  $1.39 \text{ m}^3 \text{ kg}^{-1}$ . The  $VWC_{\text{total}}$  values show a three-tiered distribution, decreasing gradually from southeast to northwest of China. The  $VWC_{\text{total}}$  values in eastern coastal China, Southeast China, Beijing, and Tianjin were relatively high. The regions with lower  $VWC_{\text{total}}$  values were mainly concentrated in the Northeast China and Northwest China.

## 4 Discussion

### 4.1 Comparison of calculations of total VWC of rice by different frameworks

Here, we compare four frameworks for total VWC determinations (Table 1). (1) In the crop water requirement (CWR) calculation framework, the total virtual water content of crops is divided into green and blue water. Green water use is the lesser of the potential crop evapotranspiration and the effective precipitation. Blue water use is the irrigation water requirement, which is the potential crop evapotranspiration minus green water

HESSD

11, 1047–1072, 2014

### VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## WVC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

use (Siebert and Döll, 2010; Sun et al., 2013a). (2) In the green, blue and grey water (GBG) calculation framework, the total virtual water content of crops is divided into green, blue, and grey water. The calculation methods for green and blue water are the same as in the CWR framework. Grey water is calculated by multiplying the fraction of nitrogen that leaches or runs off by the nitrogen application rate and dividing this by the difference between the maximum acceptable concentration of nitrogen and the natural concentration of nitrogen in the receiving water body (Chapagain et al., 2006; Bulsink et al., 2010; Mekonnen and Hoekstra, 2011b). (3) In Sun's framework, the total virtual water content of crops is also divided into green and blue water. Blue water is calculated according to actual irrigation water consumption (Sun et al., 2013b). (4) In our framework, the total virtual water content of crops is divided into direct green, direct blue, direct grey and indirect water. The calculations for green and grey water are the same as in the GBG framework and the calculation method for blue water is the same as in Sun's framework. As described previously, we add indirect water to the total VWC of crops.

The VWC of rice of Heilongjiang province in 2007 is used as an example to compare results among the four different frameworks. As shown in Table 2,  $VWC_{total}$  calculated by our framework is  $0.97 \text{ m}^3 \text{ kg}^{-1}$ , which accounts for 66.9, 57.7, and 133.8% of the  $VWC_{total}$  of rice under the CWR framework, GBG framework, and Sun's framework, respectively. The  $VWC_{total}$  calculated by our framework is lower than the  $VWC_{total}$  calculated by CWR framework and GBG framework. Because the  $VWC_{direct, blue}$  is calculated by the crop irrigation water requirement under CWR framework and GBG framework and calculated by the actual irrigation consumption under our framework. According to the calculation in our method, the actual irrigation consumption can not meet the irrigation water requirement of rice in Heilongjiang. The  $VWC_{direct, blue}$  value calculated from the actual irrigation consumption under our framework is much lower than the  $VWC_{direct, blue}$  values calculated under the CWR and GBG frameworks. Adding the  $VWC_{direct, grey}$  value resulted in the  $VWC_{total}$  calculated by our framework being larger than the  $VWC_{total}$  calculated by Sun's framework. The contribution of  $VWC_{direct, grey}$  is

very important, accounting for 22.9% of the  $\text{VWC}_{\text{total}}$  of rice under our framework, and cannot be ignored. The contribution of  $\text{VWC}_{\text{indirect}}$  as calculated under our framework is limited and has little effect on the  $\text{VWC}_{\text{total}}$  of rice in Heilongjiang.

## 4.2 Analysis of VWC structures of rice in China

5 Regional differences in  $\text{VWC}_{\text{total}}$  values for rice were significant, as were regional differences in the structure of the  $\text{VWC}_{\text{total}}$  of rice. The average  $\text{VWC}_{\text{total}}$  of rice in China was  $1.39 \text{ m}^3 \text{ kg}^{-1}$ . The percentages of  $\text{VWC}_{\text{direct, green}}$ ,  $\text{VWC}_{\text{direct, blue}}$ ,  $\text{VWC}_{\text{direct, grey}}$ , and  $\text{VWC}_{\text{indirect}}$  in the  $\text{VWC}_{\text{total}}$  for rice in China were 43.8, 28.2, 27.6, and 0.4%, respectively.

10 Generally speaking, the regions with lower  $\text{VWC}_{\text{direct, green}}$  values usually had higher  $\text{VWC}_{\text{direct, blue}}$  values, and vice versa. For example, in Beijing the percentage of  $\text{VWC}_{\text{direct, green}}$  was 19.8% and the percentage of  $\text{VWC}_{\text{direct, blue}}$  was 63.2%, in Tianjin the percentage of  $\text{VWC}_{\text{direct, green}}$  was 14.7% and the percentage of  $\text{VWC}_{\text{direct, blue}}$  was 69.1%, in Chongqing the percentage of  $\text{VWC}_{\text{direct, green}}$  was 67.6% and the percentage of  $\text{VWC}_{\text{direct, blue}}$  was 6.6%, and in Sichuan the percentage of  $\text{VWC}_{\text{direct, green}}$  was 53.7% and the percentage of  $\text{VWC}_{\text{direct, blue}}$  was 12.4%. Three regions were exceptions, where the percentages of  $\text{VWC}_{\text{direct, green}}$  and  $\text{VWC}_{\text{direct, blue}}$  of rice were consistent: in Inner Mongolia the percentage of  $\text{VWC}_{\text{direct, green}}$  was 29.7% and the percentage of  $\text{VWC}_{\text{direct, blue}}$  was 27.2%, in Gansu the percentage of  $\text{VWC}_{\text{direct, green}}$  was 33.5% and the percentage of  $\text{VWC}_{\text{direct, blue}}$  was 37.9%, and in Shanghai the percentage of  $\text{VWC}_{\text{direct, green}}$  was 41.4% and the percentage of  $\text{VWC}_{\text{direct, blue}}$  was 40.8%. Beijing, Tianjin, Shanghai, and Fujian had the highest  $\text{VWC}_{\text{total}}$  of rice and also had the highest  $\text{VWC}_{\text{direct, blue}}$  of rice. This shows that rice grown in the regions with relatively high  $\text{VWC}_{\text{total}}$  values had greater dependence on direct blue water. But for the country as a whole, the percentage of  $\text{VWC}_{\text{direct, green}}$  of rice (43.8%) was far above the  $\text{VWC}_{\text{direct, blue}}$  value (28.2%), indicating that rice growth is mainly dependent on direct green water in China.

### VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Regional differences in  $VWC_{\text{direct, grey}}$  values were insignificant. This shows that the grey water in all regions plays an important role in  $VWC_{\text{total}}$ . Agricultural pollution is an important issue in every region that cannot be ignored. Because the direct grey water estimated only considers chemical fertilizer pollution, and not the effect of pesticides and herbicides on water quality, the result of this estimation is a conservative estimate.

The indirect water values in the VWC of rice for the 29 regions were comparatively less, ranging from 0.001 to  $0.010 \text{ m}^3 \text{ kg}^{-1}$ . The average indirect water contribution to the VWC of rice was  $0.004 \text{ m}^3 \text{ kg}^{-1}$ . The region with the highest  $VWC_{\text{indirect}}$  value was Beijing and the region with the lowest value was Ningxia. Because the contribution of  $VWC_{\text{indirect}}$  was quite small, in future research on the VWC of rice, indirect water may not be considered. For some agricultural products such as potatoes, cotton, and fruits, the  $VWC_{\text{indirect}}$  is expected to be higher and should be included in the  $VWC_{\text{total}}$ .

## 5 Conclusions

Faced with increasingly severe pressure on water resources, virtual water theory provides a feasible solution to improve global water use efficiency. Research on the VWC of crops can provide the basis for agricultural water resources management and help to improve the efficiency of agricultural water use. Rice is the most important food crop in China and also one of the largest water consumers, so it is important to study the total VWC of rice in China. Previous research constructed calculation frameworks using direct green water, direct blue water, and direct grey water of crops. Building on that previous research, we also considered the indirect water of VWC. In this paper, we calculated the virtual water content of rice for 29 regions of China in 2007. The following conclusions were reached.

1. The total VWC of rice ranged from 0.804 to  $2.586 \text{ m}^3 \text{ kg}^{-1}$ . The average total VWC of rice of China in 2007 was  $1.39 \text{ m}^3 \text{ kg}^{-1}$ . The percentages of direct green, blue, grey and indirect water were 43.8, 28.2, 27.6, and 0.4% in the total VWC of rice in China, respectively. Analysis showed that the total VWC of rice in China decreased gradually from southeast to northwest.

## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

2. The indirect VWC of rice for 29 regions in 2007 was comparatively lower, ranging from  $0.001$  to  $0.010 \text{ m}^3 \text{ kg}^{-1}$ , with an average of  $0.004 \text{ m}^3 \text{ kg}^{-1}$ . The regions with high indirect water are randomly distributed. The indirect VWC of rice in Northwest China and Southwest China was relatively low.
3. The direct green water of rice for 29 regions in 2007 ranged from  $0.10$  to  $0.90 \text{ m}^3 \text{ kg}^{-1}$ , with an average of  $0.59 \text{ m}^3 \text{ kg}^{-1}$ . The regions with higher direct green water were mainly concentrated in the Southeast China and Southwest China. The direct green water of rice for Northwest China and Inner Mongolia was relatively low.
4. The direct blue water of rice for 29 regions in 2007 ranged from  $0.07$  to  $1.65 \text{ m}^3 \text{ kg}^{-1}$ , with an average of  $0.42 \text{ m}^3 \text{ kg}^{-1}$ . The regions with higher direct blue water of rice were mainly concentrated in the eastern and southern coastal regions of China and in Northwest China. The direct blue water of rice of Southwest China was relatively low.
5. Rice grown in regions with relatively high total VWC depended greatly on direct blue water. In the country as a whole, the percentage of direct green water of rice was far above that of direct blue water. Therefore, rice growth is mainly dependent on direct green water in China.
6. The direct grey water of rice ranged from  $0.21$  to  $0.64 \text{ m}^3 \text{ kg}^{-1}$ , with an average of  $0.37 \text{ m}^3 \text{ kg}^{-1}$ . The direct grey water of rice in Northeast China and Northwest China was relatively low. But in all regions, grey water occupies a very important position in total VWC.

*Acknowledgements.* We thank the National Science Foundation for Innovative Research Group (No. 51121003), the International Science & Technology Cooperation Program of China (No. 2011DFA72420), the National Basic Research Program of China (No. 51309012) for their financial support.

## References

- Allan, J. A.: Overall perspectives on countries and regions, in: Water in Arab World: Perspectives and Progress, edited by: Rogers, P. and Lydon, P., Harvard University Press, Cambridge, UK, 65–100, 1994.
- 5 Allan, J. A.: The Middle East Water Question: Hydropolitics and the Global Economy, Tauris Publishers, London, UK, 2002.
- Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration – guidelines for computing crop water requirements, FAO Irrigation and Drainage Paper 56, FAO, Rome, 1998.
- 10 Bultink, F., Hoekstra, A. Y., and Booij, M. J.: The water footprint of Indonesian provinces related to the consumption of crop products, Hydrol. Earth Syst. Sci., 14, 119–128, doi:10.5194/hess-14-119-2010, 2010.
- Chapagain, A. K. and Hoekstra, A. Y.: The global component of fresh water demand and supply: an assessment of virtual water flows between nations as a result of trade in agricultural and industrial products, Water Int., 33, 19–32, 2008.
- 15 Chapagain, A. K. and Hoekstra, A. Y.: The blue, green and grey water footprint of rice from production and consumption perspectives, Ecol. Econ., 70, 749–758, 2011.
- Chapagain, A. K., Hoekstra, A. Y., Savenije, H. H. G., and Gautam, R.: The water footprint of cotton consumption: an assessment of the impact of worldwide consumption of cotton products on the water resources in the cotton producing countries, Ecol. Econ., 60, 186–203, 2006.
- 20 Chen, X. K.: Shanxi water resource input-occupancy-output table and its application in Shanxi Province of China, in: The 13th International Conference on Input-output Techniques, Macerata, Italy, 21–25 August 2000, 2–11, 2000.
- 25 China Meteorological Administration: China Meteorological Data Sharing Service System, Beijing, China, available at: <http://cdc.cma.gov.cn> (last access: 6 September 2011), 2011.
- Clarke, D.: CROPWAT for Windows: User Guide, University of Southampton, UK, 1998.
- FAO (Food and Agriculture Organization: Land and Water Development Division, CROPWAT model, Food and Agriculture Organization, Rome, Italy, available at: [http://www.fao.org/nr/water/infos\\_databases\\_cropwat.html](http://www.fao.org/nr/water/infos_databases_cropwat.html), last access: 9 June 2011, 2003.
- 30

## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Hoekstra, A. Y. and Hung, P. Q.: Virtual water trade: a quantification of virtual water flows between nations in relation to international crop trade, Value of Water Research Report Series No. 11, UNESCO-IHE, Delft, the Netherlands, 2002.
- Hoekstra, A. Y., Chapagain, A. K., Aldaya, M. M., and Mekonnen, M. M.: The Water Footprint Assessment Manual: Setting the Global Standard, Earthscan, London, UK, 2011.
- Kanada, N.: Land Resources and International Trade, Taga Shuppan, Tokyo, 2001.
- Leontief, W.: The Structure of the American Economy, Oxford University Press, Oxford, UK, 1941.
- Li, H. L., Zhang, W. F., Zhang, F. S., Du, F., and Li, L. K.: Chemical fertilizer use and efficiency change of main grain crops in China, Journal of Plant Nutrition and Fertilizer, 16, 1136–1143, 2010 (in Chinese).
- Liu, J. G., Wiberg, D., Zehnder, A. J. B., and Yang, H.: Modelling the role of irrigation in winter wheat yield, crop water productivity, and production in China, Irrigation Science, 26, 21–33, 2007.
- Mekonnen, M. M. and Hoekstra, A. Y.: The green, blue and grey water footprint of crops and derived crop products, Hydrol. Earth Syst. Sci., 15, 1577–1600, doi:10.5194/hess-15-1577-2011, 2011a.
- Mekonnen, M. M. and Hoekstra, A. Y.: National water footprint accounts: the green, blue and grey water footprint of production and consumption, Value of Water Research Report Series No. 50, UNESCO-IHE, Delft, the Netherlands, 2011b.
- Ministry of Agriculture of the People's Republic of China: Chinese agricultural statistical data, Ministry of Agriculture of the People's Republic of China, Beijing, 2008.
- Ministry of Water Resources: China Water Resources Bulletin 2011, China Water & Power Press, Beijing, 2012.
- Rost, S., Gerten, D., Bondeau, A., Lucht, W., Rohwer, J., and Schaphoff, S.: Agricultural green and blue water consumption and its influence on the global water system, Water Resour. Res., 44, W09405, doi:10.1029/2007WR006331, 2008.
- Siebert, S. and Döll, P.: Quantifying blue and green virtual water contents in global crop production as well as potential production losses without irrigation, J. Hydrol., 384, 198–207, 2010.
- Sun, S. K., Wu, P. T., Wang, Y. B., and Zhao, X. N.: The virtual water content of major grain crops and virtual water flows between regions in China, J. Sci. Food Agr., 93, 1427–1437, 2013a.

# HESSD

11, 1047–1072, 2014

## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Sun, S. K., Wu, P. T., Wang, Y. B., Zhao, X. N., Liu, J., and Zhang, X. H.: The impacts of interannual climate variability and agricultural inputs on water footprint of crop production in an irrigation district of China, *Sci. Total Environ.*, 444, 498–507, 2013b.

Yu, P. P., Zhang, J. Z., and Lin, C. G.: Agricultural development strategy of China in the early 21st century, *Resour. Environ. Dev.*, 7, 21–27, 2006 (in Chinese).

Zhao, X., Chen, B., and Yang, Z. F.: National water footprint in an IO framework: a case study of China 2002, *Ecol. Model.*, 220, 245–253, 2009.

Zhang, Z. Y., Yang, H., and Shi, M. J.: Analyses of water footprint of Beijing in an interregional input-output framework, *Ecol. Econ.*, 70, 2494–2502, 2011.

## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.

**Table 1.** Comparison of VWC calculating frameworks.

Method	VWC			
	VWC <sub>direct, green</sub>	VWC <sub>direct, blue</sub>	VWC <sub>direct, grey</sub>	VWC <sub>indirect</sub>
Our method	$10 \min(ET_C, P_e)/Y$	$VWC_{\text{direct, blue}} = I_c/Y = \frac{W_A \alpha_i}{A_i}/Y$	$VWC_{\text{direct, grey}} = \frac{N_c \cdot 10\%}{10 - N_n}/Y$	$VWC_{\text{indirect}} = \frac{VW_{\text{blue, indirect}}^{\beta_i}}{A_i}/Y$
Sun's method	$10 \min(ET_C, P_e)/Y$	$VWC_{\text{direct, blue}} = I_c/Y = \frac{W_A \alpha_i}{A_i}/Y$	–	–
GBG method	$10 \min(ET_C, P_e)/Y$	$10 \max(0, ET_c - P_e)/Y$	$VWC_{\text{direct, grey}} = \frac{N_c \cdot 10\%}{10 - N_n}/Y$	–
CWR method	$10 \min(ET_C, P_e)/Y$	$10 \max(0, ET_c - P_e)/Y$	–	–

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



## WVC of rice and spatial characteristics analysis in China

L. J. Zhang et al.




Fig. 1. Indirect water of rice ( $\text{m}^3 \text{ kg}^{-1}$ ).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.



**Fig. 2.** Direct green water of rice ( $\text{m}^3 \text{kg}^{-1}$ ).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.



**Fig. 3.** Direct blue water of rice ( $\text{m}^3 \text{kg}^{-1}$ ).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## VWC of rice and spatial characteristics analysis in China

L. J. Zhang et al.



**Fig. 4.** Direct grey water of rice ( $\text{m}^3 \text{kg}^{-1}$ ).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## WVC of rice and spatial characteristics analysis in China

L. J. Zhang et al.




Fig. 5. Total VWC of rice (m<sup>3</sup> kg<sup>-1</sup>).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion