

1

2 **Sensitivity of potential evaporation estimates to 100 years**  
3 **of climate variability**

4 **R.P. Bartholomeus<sup>1</sup>, J.H. Stagge<sup>2</sup>, L.M. Tallaksen<sup>2</sup>, J.P.M. Witte<sup>1,3</sup>**

5 1 KWR Watercycle Research Institute, P.O. Box 1072, 3430 BB Nieuwegein, The  
6 Netherlands

7 2 Department of Geosciences, University of Oslo, P.O. box 1047, Blindern, 0316 Oslo,  
8 Norway

9 3 VU University, Institute of Ecological Science, Department of Systems Ecology, de  
10 Boelelaan 1085, 1081 HV Amsterdam, The Netherlands

11 Correspondence to: R.P. Bartholomeus (ruud.bartholomeus@kwrwater.nl)

12

13

## Abstract

Hydrological modeling frameworks require an accurate representation of evaporation fluxes for appropriate quantification of e.g. the water balance, soil moisture budget, recharge and groundwater processes. Many frameworks have used the concept of potential evaporation, often estimated for different vegetation classes by multiplying the evaporation from a reference surface ('reference evaporation') with crop specific scaling factors ('crop factors'). Though this two-step potential evaporation approach undoubtedly has practical advantages, the empirical nature of both reference evaporation methods and crop factors limits its usability in extrapolations under non-stationary climatic conditions. In this paper, rather than simply warning about the dangers of extrapolation, we quantify the sensitivity of potential evaporation estimates for different vegetation classes using the two-step approach when calibrated using a non-stationary climate. We used the past century's time series of observed climate, containing non-stationary signals of multi-decadal atmospheric oscillations, global warming, and global dimming/brightening, to evaluate the sensitivity of potential evaporation estimates to the choice and length of the calibration period. We show that using empirical coefficients outside their calibration range may lead to systematic differences between process-based and empirical reference evaporation methods, and systematic errors in estimated potential evaporation components. Quantification of errors provides a possibility to correct potential evaporation calculations and to rate them for their suitability to model climate conditions that differ significantly from the historical record, so-called no-analogue climate conditions.

## 1    **1    Introduction**

2    Evaporation from the vegetated surface is the largest loss term in many, if not the most, water  
3    balance studies on earth. As a consequence, an accurate representation of evaporation fluxes  
4    is required for appropriate quantification of surface runoff, the soil moisture budget,  
5    transpiration, recharge and groundwater processes (Savenije, 2004). However, despite being a  
6    key component of the water balance, evaporation figures are usually associated with large  
7    uncertainties, as this term is difficult to measure (Allen et al., 2011) or estimate by modeling  
8    (Wallace, 1995).

9    Research attempting to model the evaporation process has a long history (Shuttleworth,  
10    2007). This research took two parallel tracks, with the meteorological community developing  
11    process-based models of surface energy exchange and the hydrological community  
12    considering evaporation as a loss term in the catchment water balance (Shuttleworth, 2007).  
13    To quantify the evaporation loss term, many hydrological modeling frameworks have used  
14    the concept of potential evaporation (Federer et al., 1996; Kay et al., 2013; Zhou et al., 2006),  
15    defined as the maximum rate of evaporation from a natural surface where water is not a  
16    limiting factor (Shuttleworth, 2007). With the progression from catchment-scale lumped  
17    models (such as HBV (Bergström and Forsman, 1973)) to distributed models with increasing  
18    spatial resolution and spatially resolved data (such as SHE (Abbott et al., 1986)), the explicit  
19    representation of land surface water budgets also increased (Ehret et al., 2014; Federer et al.,  
20    1996). To this end, estimation of evaporation from a variety of land surfaces within the  
21    simulated domain is needed (Federer et al., 1996). More models were developed that included  
22    vegetation explicitly, commonly by describing the stomatal conductance of the vegetation as a  
23    function of environmental drivers (see Shuttleworth (2007) and references therein). However,  
24    until now these models are rarely used in practice and merely have a scientific meaning.

25    Parallel to this development, the irrigation engineering community refined the traditional  
26    potential evaporation approach (Shuttleworth, 2007). They developed the ‘two-step approach’  
27    (Doorenbos and Pruitt, 1977; Penman, 1948; Zhou et al., 2006; Feddes and Lenselink, 1994;  
28    Vázquez and Feyen, 2003; Hupet and Vanclooster, 2001), in which the potential evaporation  
29    of a specific crop or vegetation class is estimated by multiplying the evaporation from a  
30    reference surface with empirical crop specific scaling factors: ‘crop factors’. This  
31    development was mainly driven by the need for a relatively simple approach using commonly  
32    available data from climate stations. The two-step approach has even expanded outside the

field of irrigation engineering into hydrological modeling frameworks. Crop factors are now being applied in 1D hydrological models (e.g. Tiktak and Bouten (1994)), spatially lumped models (e.g. Driessen et al. (2010); Calder (2003)), and spatially distributed hydrological models (e.g. Ward et al. (2008); Shabalova et al. (2003); Trambauer et al. (2014); Van Roosmalen et al. (2009); Lenderink et al. (2007); Bradford et al. (1999); Guerschman et al. (2009); Sperna Weiland et al. (2012); Van Walsum and Supit (2012); Vázquez and Feyen (2003)).

With the development of the two-step potential evaporation approach, different equations to simulate reference evaporation have been suggested (Federer et al., 1996; Bormann, 2011; Shuttleworth, 2007) for use in both regional and global hydrological models (e.g. Sperna Weiland et al. (2012); Haddeland et al. (2011)). However, due to their empirical nature, these equations are limited in their transferability in both time and space (Feddes and Lenselink, 1994; Wallace, 1995). Since the increasing need for predictions under global change (land use and climate) (Ehret et al., 2014; Coron et al., 2014; Montanari et al., 2013), the empirical nature of most commonly used potential evaporation approaches is a serious drawback (Hurkmans et al., 2009; Wallace, 1995; Shuttleworth, 2007; Witte et al., 2012). Thus, although the two-step approach may be warranted for practical reasons, both the reference evaporation and estimated crop factors include a series of empirical parameters that may affect the validity and general applicability of the estimated potential evaporation for a specific vegetation class.

Since the term “potential evaporation” has been used by the hydrologic community to refer to several different combinations of evaporation components in the past, it is important to re-introduce these definitions and to be very specific about nomenclature in future evaporation research. Total evaporation ( $E_{\text{tot}}$ ) from a vegetated surface is the sum of three fluxes: transpiration ( $E_t$ ), soil evaporation ( $E_s$ ) and evaporation of intercepted water ( $E_i$ ).  $E_t$  and  $E_s$  occur at a potential rate when the availability of water (soil moisture or interception) is not limiting. As we will only focus on potential rates in this paper, all values should be interpreted as potential, unless stated otherwise. Reference evaporation ( $E_{\text{ref}}$ ) is defined as the rate of evaporation from an extensive surface of green grass, with a uniform height of 0.12 m, a surface resistance of  $70 \text{ s m}^{-1}$ , an albedo of 0.23, actively growing, completely shading the ground and with adequate water (Allen et al., 1998). By definition,  $E_i$  is not part of reference evaporation, as it is defined for a plant surface which is externally dry (Federer et al., 1996;

Allen et al., 1998). Often, the term reference evapotranspiration is used instead, which is the sum of transpiration ( $E_t$ ) and soil evaporation ( $E_s$ ). By definition (Allen et al., 1998) the reference crop completely shades the ground and hence  $E_s$  will be zero and  $E_{ref}$  equals  $E_t$  of the reference crop (at least for daily estimates, when the soil heat flux can be assumed zero). This is in agreement with the definition of Penman (1956) who also stated that the often-used expansion of the term “reference evaporation” to “evapotranspiration” was unnecessary.

$E_{ref}$  is used in the two-step method to estimate the potential evaporation,  $E_p$ , of a crop or vegetation stand.  $E_p$  will reduce to the actual evaporation,  $E_a$ , in case of water shortage or waterlogging. Here, we focus on the estimation of  $E_p$  from  $E_{ref}$ , by multiplying  $E_{ref}$  with a crop factor  $K$  (Allen et al., 2005; Feddes, 1987; Allen et al., 1998; Penman, 1956). Different applications of crop factors exist:

- $K_t$  corrects for potential transpiration of a crop with a dry canopy only, i.e.  $E_t = K_t \times E_{ref}$ . This corresponds to the basal crop factors defined by Allen (2000), which are equivalent to the approach of Penman (1956).
- $K_{ts}$  corrects for both potential transpiration and potential soil evaporation for a crop with a dry canopy, i.e.  $E_t + E_s = K_{ts} \times E_{ref}$ . This corresponds to the single crop factors defined by Allen et al. (1998).
- $K_{tot}$  corrects for potential total evaporation, i.e. transpiration, soil evaporation, and interception. Using  $K_{tot}$  with  $E_{ref}$  directly gives  $E_{tot}$ , i.e.  $E_{tot} = K_{tot} \times E_{ref}$ .

$K_{tot}$  holds for crop factors that have been derived by soil water balance experiments, and especially from sprinkling experiments in the field, where water is applied in such quantities that soil water is not limiting for plant growth (Feddes, 1987). Sprinkling, however, leads to interception. So, crop factors like those of Feddes (1987) implicitly involve  $E_i$ . Therefore Feddes (1987) emphasizes that the presented crop factors “are averages taken over a population of ‘average’, ‘dry’, and ‘wet’ years, that will certainly not be homogeneously distributed”. The crop factor approach by Feddes (1987) is different from the single crop factor approach of Allen et al. (1998), as crop factors from the latter are by definition applied to correct for  $E_t + E_s$ , or for  $E_t$  only (Allen, 2000). However, Allen et al. (1998) indicate that their crop factors should be multiplied with a factor 1.1-1.3 if interception, due to sprinkling irrigation for example, is involved (i.e. if interception is not simulated explicitly). This indicates that  $E_i$  could significantly affect potential evaporation from a vegetated surface. As  $E_i$  is largely driven by precipitation, a term that is generally not incorporated in  $E_{ref}$  methods,

1 it has already been stated that the crop factor approach only makes sense in times of drought,  
2 when interception does not contribute to the total evaporation (De Bruin and Lablans, 1998).  
3 This condition is especially relevant for tall forests, which intercept a higher percentage of  
4 rain water, under climatological conditions with significant rainfall (De Bruin and Lablans,  
5 1998). Nevertheless, this crop factor approach is used in practice (Van Roosmalen et al.,  
6 2009).

7 The objective of this paper is to assess the sensitivity of potential evaporation estimates for  
8 different vegetation classes using the commonly used two-step approach when calibrated  
9 based on a non-stationary climate. To this end, we use century long meteorological  
10 observations representing the historic variability in climatic conditions at the De Bilt, The  
11 Netherlands climate monitoring station. The past century's global warming, dimming and  
12 brightening periods (Suo et al., 2013; Stanhill, 2007; Wild, 2009; Wild et al., 2005), and their  
13 effects on evaporation provide an opportunity to evaluate the robustness of the two-step  
14 estimation of potential evaporation for non-stationary conditions. Given the 20<sup>th</sup> century  
15 climate induced variability in  $E_{\text{ref}}$  and the projected increase for the near future, which has no  
16 historical analogue, (Fig. 1), it is of great importance to recognize the limitations of applying  
17 empirical coefficients outside their calibration range (i.e. extrapolation). This applies not only  
18 to transferring coefficients in space, as between climatic regions (Allen et al., 1998), but also  
19 in time.

20 The 20<sup>th</sup> century global surface temperature can be characterized by two major warming  
21 periods; the first one from about 1925-1945, followed by a period of cooling, and a second  
22 starting in about 1975 and continuing to the present (Jones and Moberg, 2003; Yamanouchi,  
23 2011). While the variations in temperature until the 1970s can be related to changes in global  
24 radiation, i.e. global dimming and brightening, this relationship no longer holds for the rapid  
25 warming since 1975 (Wang and Dickinson, 2013). Empirical equations for reference  
26 evaporation that use either radiation or temperature implicitly assume a relationship between  
27 the two variables. Given the nonlinearity of evaporation components, it is not only  
28 questionable whether empirical equations for reference evaporation will be applicable under  
29 future climatic conditions (Shaw and Riha, 2011), but also whether they are applicable for the  
30 recent past.

31 Although the limitations of using empirical coefficients to calculate evaporation are generally  
32 well known, the potential errors that could be made by using such coefficients in evaporation

calculations have, as far as we know, never been quantified. Thus, there is a need to raise the awareness of the uncertainty that may result applying such an empirical estimation method outside its valid area (site and time specific). In this study we systematically unravel the use of the two-step approach to simulate potential evaporation and identify and actually quantify systematic errors that may be introduced when empirical coefficients are applied outside their calibration period. Such extrapolations of time-variant model parameters are not only relevant for the calculation of potential evaporation, but also for hydrological modeling in general, thus limiting the temporal robustness of hydrological models (Ehret et al., 2014; Karlsson et al., 2014; Coron et al., 2014; Seibert, 2003). Quantification of errors, as demonstrated in this study, provides the possibility to i) derive uncertainty ranges for the parameters, ii) quantify the errors that are introduced by a specific method and set of parameters, and iii) correct for the errors when they are predictable.

## 2 Methods

### 2.1 General approach

We use 108 years of meteorological observations to quantify the sensitivity of potential evaporation when calibrated using a non-stationary climate for various natural vegetation classes using the two-step approach. We investigate how empirical  $E_{\text{ref}}$ -methods and empirical  $K$ -values affect the validity of the estimated potential evaporation for different vegetation classes, by applying empirical coefficients outside their calibration period. We vary the calibration period in both length (2-30 years) and reference period (in 1906-2013).

First (section 2.3), we simulate reference evaporation according to the process-based Penman-Monteith equation ( $E_{\text{ref\_PM}}$ ), which is considered the international standard method for estimating reference evaporation (Allen et al., 1998). In addition, we apply four empirical equations that contain constants derived for a calibration period (Fig. 2: §2.3). From these simulations, we identify deviations between each empirical  $E_{\text{ref}}$  method and the  $E_{\text{ref\_PM}}$  (Fig. 2: §2.3).

Secondly (section 2.4), we generate time series of the main components of potential evaporation, i.e. synthetic series of  $E_t$ ,  $E_s$  and  $E_i$ , for five different vegetation classes, using the Soil-Vegetation-Atmosphere Transfer (SVAT) scheme SWAP (Kroes et al., 2009; Van Dam et al., 2008) (Fig. 2: §2.4). SWAP allows users to simulate potential evaporation for

different vegetation classes directly (i.e. one-step approach), by parameterizing the Penman-Monteith equation for each vegetation class implicitly rather than using crop factors. These synthetic series are considered ‘observations’ throughout the paper for all comparisons with estimates from the two-step approach.

Finally (section 2.5), we derive monthly crop factors for each vegetation type (5x) and for each  $E_{\text{ref}}$  method (5x) based on the synthetic data of  $E_t$ ,  $E_s$  and  $E_i$  for a calibration period (e.g. 1906-1935) to simulate crop factor estimation using field measurements (Fig. 2: §2.5). We use different (3x) definitions of crop factors: for transpiration ( $K_t$ ), for transpiration plus soil evaporation ( $K_{ts}$ ) and for total evaporation ( $K_{\text{tot}}$ ). Next, we apply the two-step approach, using  $E_{\text{ref}}$  and crop factors from the calibration period to calculate daily ‘predicted’ evaporation components (3x) for each vegetation class (5x) and each  $E_{\text{ref}}$  method (5x) for the entire period (1906-2013) (Fig. 2: §2.6). Doing so, the empirical  $E_{\text{ref}}$  methods and crop factors are applied outside their calibration range. From these simulations we quantify the deviations introduced by the use of  $E_{\text{ref}}$  and  $K$ , by comparing the evaporation components obtained with the two-step approach to the synthetic ‘observations’ (Fig. 2: §2.6). Each of these steps, which are executed for all calibration periods during the period 1906-2013 (2697x), are described in greater detail in subsequent sections.

Although SWAP may be expected to provide adequate evaporation values, its absolute accuracy is not discussed in this paper, because we focus on the sensitivity of the two-step approach using synthetic (hypothetical) data only. Therefore, the actual accuracy of SWAP is irrelevant for this paper. To ensure that our analysis is not biased by a specific choice of SWAP parameter settings, we considered different vegetation classes ranging from grasses to shrubs and forests. Doing so, we include different parameter sets that affect the variability and relative proportions of the calculated  $E$  components. For a detailed discussion of the SWAP model and its accuracy, please refer to Kroes et al. (2009) and Van Dam et al. (2008). By comparing potential evaporation components obtained from the two-step approach with the synthetic ‘observations’ as simulated using the physical SWAP model, we are able to quantify the deviations introduced by using different  $E_{\text{ref}}$  methods in combination with crop factors, as no other source of uncertainty is involved.



## 2.2 Meteorological data

We use meteorological data from De Bilt, The Netherlands, covering the period 1906-2013, which was provided by the Royal Netherlands Meteorological Institute (KNMI). De Bilt (longitude = 5.177° east, latitude = 52.101° north, altitude = 2 m) is the main meteorological site of the KNMI, located in the center of the Netherlands. Daily records are available for minimum and maximum temperature, sunshine hours, wind speed, and precipitation from 1906 onwards, and for global radiation from 1957. The observations are continuous, except for April 1945, where values from April 1944 are used instead. All required input variables are calculated for the period 1906-2013 following Allen et al. (1998). Observed global radiation was used to derive the Angstrom coefficients needed to calculate daily global radiation (Allen et al., 1998) from 1906 onwards. For consistency we only use these simulated values for further analysis, which agree very well with observations (1957-2013,  $R^2_{adj} = 0.96$ ). Wind speed, measured at different heights, was scaled to the reference height of 2 meter following Allen et al. (1998) and corrected for systematic differences between measurement periods. Fig. 3 shows the annual values and the 30 year moving averages of the variables used to calculate evaporation from De Bilt.

Although the results are only valid for the site and period they were developed, the times series of radiation for De Bilt station resembles the global trends of global dimming/brightening. Values of global radiation ( $R_s$ ) from De Bilt show a similar trend to the observations for Stockholm, as presented in Wild (2009). The data (Fig. 3) show an increase in temperature consistent with previous studies (Solomon et al., 2007) and a pattern of sunshine duration consistent with dimming and brightening for northwestern Europe identified by Sanchez-Lorenzo et al. (2008).

Long time series of meteorological observations will, to some extent, not be homogeneous, for example due to changes in measurement devices over time. However, this does not affect the calculations herein, as the aim is to investigate the sensitivity of the two-step potential evaporation methodology to non-stationary climate, rather than to produce an exact reconstruction of the last century's climate conditions. In this way, changes in measurement accuracy with time simply represent another non-stationary trend in this dataset.

## 2.3 Reference evaporation

Several methods are available for calculating reference evaporation, differing in complexity and empiricism (Sperna Weiland et al., 2012; Bormann, 2011; Federer et al., 1996). Here we analyze five of these methods, given in Table 1: the physically-based Penman-Monteith equation (PM), the radiation based methods of Makkink (Mak) and Priestley-Taylor (PT), and the temperature based methods of Hargreaves (Har) and Blaney-Criddle (BC).

The FAO-56 method (Allen et al., 1998), using PM parameterized for reference grass, is recommended as the international standard for calculation of  $E_{\text{ref}}$ . Given the physical basis of PM, it can be used globally, without the need to estimate or calibrate its parameters (Droogers and Allen, 2002). In contrast, the methods of Mak, PT, Har, and BC contain empirical coefficients, derived for specific meteorological conditions and sites. Following Farmer et al. (2011) we consider  $E_{\text{ref\_PM}}$  as the best approximation of  $E_{\text{ref}}$ . In order to reduce any systematic differences between  $E_{\text{ref}}$  values, we estimate the empirical factors  $C_1$ ,  $C_0$ ,  $\alpha'$ ,  $\beta$ ,  $a$ ,  $b$ ,  $c$ ,  $d$  of the other four  $E_{\text{ref}}$  methods (Table 1) by least squares regression against the simulated daily  $E_{\text{ref\_PM}}$ , for a specific calibration period. Subsequently, daily values of  $E_{\text{ref}}$  are calculated for each method during the full period, i.e. 1906-2013, and deviations between the empirical  $E_{\text{ref}}$  methods and  $E_{\text{ref\_PM}}$  are calculated. The sensitivity of  $E_{\text{ref}}$  to the choice of calibration period is evaluated for each of the methods using  $E_{\text{ref\_PM}}$  as a basis.

## 2.4 Synthetic evaporation series

Synthetic time series of the three evaporation components are derived to systematically unravel the use of empirical crop factors. The synthetic time series are based on the physical model SWAP (Van Dam et al., 2008; Kroes et al., 2009) from which  $E_t$ ,  $E_s$  and  $E_i$  can be simulated separately. From these simulations we derive monthly  $K$ -values for each  $E_{\text{ref}}$  method (5x) and vegetation class (5x) (Fig. 2: §2.5), which are subsequently used to derive the corresponding potential evaporation components (5x5x3) using the two-step approach (Fig. 2: §2.6).

Standard values for the vegetation classes and their schematization are taken from the National Hydrologic Instrument (NHI, [http://www.nhi.nu/nhi\\_uk.html](http://www.nhi.nu/nhi_uk.html); De Lange et al. (2014)) of The Netherlands. The vegetation schematization is constant throughout the period 1906-2013, i.e. dynamic vegetation is not simulated. We consider five natural vegetation classes: grassland (height = 0.5 m and no full soil cover, i.e. not to be confused with the

reference grass), heather, deciduous forest, pine forest and spruce forest. Parameters are chosen following NHI (2008) and are provided in the supplementary material. It should be noted that we do not discuss the exact validity of the parameter values used, as we are only concerned with evaporation sensitivity to non-stationary climate within the range of typical vegetation.

SWAP simulates the potential evaporation components of a crop or vegetation class based on the aerodynamic resistance, height, Leaf Area Index (LAI), and albedo. SWAP uses the Penman-Monteith equation, parameterized for each vegetation class to simulate  $E_t$  (potential transpiration) and  $E_s$  (potential soil evaporation). In case of intercepted precipitation, the values of  $E_t$  and  $E_s$  are reduced (Van Dam et al., 2008). Interception, which partly evaporates ( $E_i$ ) and partly drips to the ground, is estimated following Von Hoyningen-Hüne (1983) and Braden (1985) for short vegetation and Gash et al. (1995) for forests. For an extended description of SWAP and the procedures for calculating  $E_t$ ,  $E_s$  and  $E_i$ , we refer to Kroes et al. (2009) and Van Dam et al. (2008). Given the international recognition of the SWAP model and successful testing, we assume that the model is able to produce representative synthetic estimates of each evaporation component.

As  $K_t$  and  $K_{ts}$  are defined for a vegetated surface with a dry canopy (i.e. without interception) and  $K_{tot}$  includes interception (see introduction), two different SWAP runs are performed for each vegetation class, without and with interception. Throughout the paper,  $E_t$  and  $E_s$  are valid for conditions with a dry canopy, whereas  $E_{tot}$  includes interception and its limiting effect on transpiration and soil evaporation.

## 2.5 Derivation of $K_t$ , $K_{ts}$ and $K_{tot}$

We derive  $K_t$ ,  $K_{ts}$  and  $K_{tot}$  for each vegetation class (5x) and  $E_{ref}$  method (5x) based on the synthetic  $E_t$ ,  $E_s$  and  $E_{tot}$  time series, and the equations given in Table 2. Similar to the calibration of  $E_{ref}$  methods,  $K$ -values are derived for a specific calibration period, (e.g. 1906-1935).  $K$ -values for each vegetation class and  $E_{ref}$  method are derived as monthly averages over the calibration period.

## 2.6 Calculation of potential evaporation components using the two-step approach

Potential evaporation components,  $E_t$ ,  $E_t$  plus  $E_s$  (hereafter  $E_t \& E_s$ ) and  $E_{tot}$ , for each vegetation class and method are calculated from the daily  $E_{ref}$  values by multiplying it with the corresponding  $K$ -values, respectively  $K_t$ ,  $K_{ts}$  and  $K_{tot}$ , for each vegetation class. Using these three definitions of crop factors separately allows quantifying the error that is made by correcting for each evaporation component.

$E_{ref}$  estimates that are calibrated for a specific period, combined with  $K$ -values determined for the same period, are used to calculate daily values of  $E_t$ ,  $E_t \& E_s$  and  $E_{tot}$  for the full period, i.e. 1906-2013. This procedure corresponds to what is commonly done using the two-step approach, where the empirical parameters of an  $E_{ref}$  method are fixed for the region in question, along with the corresponding  $K$ -values. Here, we determine the deviation that is potentially introduced when this approach is applied outside its calibration range (period and region/site) in a changing environment, by comparing  $E_t(E_{ref}, K_t)$ ,  $E_t \& E_s(E_{ref}, K_{ts})$  and  $E_{tot}(E_{ref}, K_{tot})$  obtained by the two-step approach with the synthetic ‘observed’  $E_t$ ,  $E_t \& E_s$  and  $E_{tot}$  series.

## 3 Results

### 3.1 Calibration period and reference evaporation

Fig. 4 shows the 30-year backwards-looking moving average  $E_{ref}$  according to PM, Mak, PT, Har and BC, with the four latter models calibrated to fit the simulated  $E_{ref\_PM}$  for the first 30-year period, i.e. the calibration period 1906-1935. The minor differences seen between all 30-year mean  $E_{ref}$  values during the calibration period (Fig. 4A, year 1935) indicate that each method was calibrated successfully. Using the calibrated equations,  $E_{ref}$ ’s are calculated for the period 1906-2013, i.e. also outside the calibration period. All empirical models are evaluated with respect to the physically based  $E_{ref\_PM}$ , which was also used when calibrating the empirical coefficients. The radiation based methods, Mak and PT, deviate only slightly from PM on average with no consistent bias (Fig. 4D), which can be explained by the relatively strong correlations between the trend in 30-year average  $E_{ref\_PM}$  (Figure 4A) and  $R_s$  and  $R_n$  (Figure 3F and G); Pearson’s  $r = 0.85$  and  $0.70$  for  $E_{ref\_PM}$  vs.  $R_s$  and  $E_{ref\_PM}$  vs.  $R_n$  respectively. The temperature based methods, Har and BC, deviate systematically from PM,

which can be explained by the relatively weak correlation between the trend in 30 year average  $E_{\text{ref\_PM}}$  (Figure 4A) and  $\bar{T}$  (Figure 3B); Pearson's  $r = 0.17$  for  $E_{\text{ref\_PM}}$  vs.  $\bar{T}$ . This shows that, as the energy used for evaporation mainly comes from direct solar radiation and to a lesser extent from air temperature, temperature based models fail if radiation and temperature trends are weakly correlated (see Figure 3; Pearson's  $r$  for 30-year average  $R_s$  vs.  $\bar{T} = 0.22$ ). Additionally, Har and BC, deviate in different directions (Fig. 4B and D): Har consistently underestimates  $E_{\text{ref}}$ , whereas BC consistently overestimates  $E_{\text{ref}}$ . This opposite trend is related to the decreasing trend in  $T_{\text{max}}-T_{\text{min}}$  (used in Har), while  $\bar{T}$  increases (Figure 3). All four empirical models are unable to reproduce the extreme high evaporation values predicted by PM, especially Har and BC (Fig. 4C). The deviations from  $E_{\text{ref\_PM}}$  are considerably larger for individual years (Fig. 4D) than for the 30-year moving average (Fig. 4B).

In practice, 30 year observed time series of evaporation are rarely available for calibration. Therefore, Fig. 5 shows the effect of calibration period length on estimates of  $E_{\text{ref}}$  for the current climate (1984-2013). This effect is expressed as the maximum absolute deviation of the 30-year average with respect to  $E_{\text{ref\_PM}}$ . Fig. 5 was compiled by first calibrating the empirical  $E_{\text{ref}}$  coefficients for all possible calibration periods (in 1906-2013) with a given length (2-30 years) and then simulating  $E_{\text{ref}}$  for the period 1984-2013 using the calibrated coefficients. The largest deviations occur for shorter calibration periods, as expected. Specific years may cause large deviations when the obtained empirical coefficients are applied outside the calibration period. Deviation decreases notably with increasing calibration periods, suggesting that using more calibration data should result in more stable and accurate  $E_{\text{ref}}$  estimates. As the calibration period length decreases, deviations in the 30-year average  $E_{\text{ref}}$  for 1984-2013 increase exponentially.

It should be noted that only deviations in 30-year averages are shown for varying calibration period lengths; deviations in the underlying yearly values are larger, as indicated by Fig. 4D. Additionally, the amplitude of the deviations shown in Fig. 4B and D would increase when calibrated using periods shorter than 30 years (Fig. 5).

### 3.2 Crop factors and potential evaporation components

Fig. 6 gives monthly average synthetic evaporation components  $E_t$ ,  $E_s$  and  $E_{\text{tot}}$  which were used to derive monthly crop factors (three methods: Table 2) for five vegetation classes and

five  $E_{\text{ref}}$  methods (Table 1), i.e. 3x5x5 crop factors for each calibration period. In contrast to the reference grass surface, the grassland of Fig. 6 does not fully cover the soil, which results in higher  $E_s$  and lower  $E_t$ . Fig. 7 shows simulated  $E_t$  (the two-step approach) for the period 1906-2013, using empirical coefficients for each  $E_{\text{ref}}$  method and matching  $K_t$ -values, all calibrated on the period 1906-1935. The general patterns in  $E_t$  correspond to those of  $E_{\text{ref}}$  (Fig. 4B), meaning that the deviations introduced by the two-step approach are mainly determined by the empirical coefficients in the  $E_{\text{ref}}$  methods.

The deviation introduced for  $E_t$  derived from  $E_{\text{ref\_PM}}$  and  $K_{t\_PM}$  is relatively minor compared to what is found for the empirical  $E_{\text{ref}}$  methods, especially for short vegetation. Apparently,  $E_{\text{ref\_PM}}$  follows the trend in  $E_t$  (also obtained using the Penman-Monteith equation, but parameterized for each vegetation class, section 2.4) and the ratio of  $E_t$  and  $E_{\text{ref\_PM}}$ , used to estimate  $K_{t\_PM}$ , changes little with time for short vegetation. More significant effects of  $K_{t\_PM}$  are seen for taller vegetation, as climate induced temporal changes in  $E_{\text{ref\_PM}}$  show a height dependent nonlinear relation to changes in  $E_t$  (Allen et al., 1998). Therefore, the deviation introduced when using  $E_{\text{ref\_PM}}$  is larger for forests than for the short vegetation classes (Fig. 7). Similar to what is seen in Fig. 4, the deviations for individual years can be considerably larger than the climatic averages.

Fig. 8 shows the sensitivity of crop factors,  $K$ , with respect to the calibration period length for heather and spruce forest. The variation in  $K$  decreases with increasing calibration length for all methods, but, except for  $E_{\text{ref\_Mak}}$ , the variability of  $K$  values for the empirical  $E_{\text{ref}}$  methods is larger than for  $E_{\text{ref\_PM}}$ . These differences are especially notable for forests and illustrate that a poor relationship between the  $E_{\text{ref}}$  method and the synthetic potential evaporation component (section 2.4) is compensated by  $K$  values that thus show a larger variation over time. Remarkable is the low variability in  $K_{\text{tot}}$  values for heather (Fig. 8C), which indicates that the variability seen for  $K_t$  (Fig. 8A) is reduced by interception. However, for spruce forest, for which interception is much more dominant, interception increases the variability in  $K_{\text{tot}}$ .

From Fig. 8A and D, it can be concluded that the deviations shown in Fig. 7 will increase when shorter calibration periods are used, irrespective of the applied  $E_{\text{ref}}$  method. Fig. 9 shows the effect of period (years and length) on the maximum absolute deviation made by the two-step approach for each  $E_{\text{ref}}$  method and for  $K_t$ ,  $K_{ts}$  and  $K_{\text{tot}}$ . Fig. 9 confirms that deviations in climatic average evaporation components obtained by applying the two-step approach will

generally increase when shorter calibration periods are used. Additionally, Fig. 9 illustrates that deviations are i) larger for tall vegetation than for short vegetation and ii) larger for  $K_{\text{tot}}$  than for  $K_t$  and  $K_{\text{ts}}$  for vegetation classes with high interception, as is the case for spruce forest. The large deviations for  $E_{\text{tot}}$  for spruce forest confirm the remark by De Bruin and Lablans (1998), that for wet forest evaporation, the crop factor approach will not be sufficient. Nevertheless, when derived for a sufficiently long time series, the deviations level out and there is no detectable bias.

### 3.3 Propagation of dimming/brightening periods

In contrast to Fig. 9, which only shows the maximum absolute deviations for the 30-year average potential evaporation components for the years 1984-2013 as a function of the calibration period length, Fig. 10 includes the results of all underlying deviations for heather and spruce using  $E_{\text{ref\_PT}}$ . Fig. 10 demonstrates that climate variability induces systematic overestimation or underestimation of the calculated potential evaporation components, depending on the calibration period used. The sign of the error strongly varies with the calibration period, and the inclusion of a single anomalous year can change the sign of the error.

Fig. 10 further shows that anomalous years or multi-annual climate patterns tend to propagate considerable errors outside the calibration period to the current climate (1984-2013). The patterns of deviations from the synthetic ‘observations’ show similarities to the global dimming and brightening periods (see Introduction): the first warming period (about 1925-1945) causes a systematic overestimation up to calibration lengths of 30 years, although specific calibration years may result in an underestimation for shorter calibration lengths. The succeeding period of cooling leads to a systematic overestimation, while the second warming period (starting around 1975) results in a more variable pattern. The latter may be linked to the finding of Wang and Dickinson (2013) that, in contrast to the years until the 1970’s, there is no significant relationship between variations in temperature and global radiation in following years.

The patterns are comparable for  $E_t$ ,  $E_t + E_s$  and  $E_{\text{tot}}$ , based on  $K_t$ ,  $K_{\text{ts}}$  and  $K_{\text{tot}}$  respectively, for short vegetation classes. However, for tall vegetation classes with high interception capacity, e.g. spruce (Fig. 10F), using  $K_{\text{tot}}$  results in a noisier pattern due to specific years of high precipitation. Additionally, including interception may shift the sign of the error.

## 4 Discussion

### 4.1 Temporal robustness in hydrological modeling

In this paper we systematically unraveled and quantified how empirical coefficients in the two-step approach affect estimates of potential evaporation. We used the past century's time series of observed climate containing non-stationary signals of multi-decadal atmospheric oscillations, global warming, and global dimming/brightening (Suo et al., 2013; Stanhill, 2007; Wild, 2009; Wild et al., 2005) to evaluate the sensitivity of the two-step approach to both the length of the reference calibration period and the reference years. To this end we calibrated the empirical coefficients of the two-step approach based on different periods and showed that using the thus obtained empirical coefficients outside their calibration range may lead to systematic differences between  $E_{\text{ref}}$ -methods, and to systematic errors in estimated potential  $E$  components. The signs of the errors for calculated climatic average evaporation components differ, depending on the  $E_{\text{ref}}$  method used, and on the specific period (length and years) of calibration. Hooghart and Lablans (1988) stated that, for the two-step approach, the correctness of empirical coefficients for the estimation of  $E_{\text{ref}}$  are of minor importance, as these are compensated by  $K$ . However, here we have shown that while this may be true within the calibration period, this statement does not hold when extrapolating. As potential evaporation is a key input in hydrological models, input errors will propagate to estimates of related processes, such as the soil moisture budget, droughts, recharge and groundwater processes.

These results are important because the two-step approach, including extrapolating empirical coefficients, is frequently applied in hydrological modeling studies, as mentioned in the introduction. Ehret et al. (2014) state that “in hydrological modeling, it is often conveniently assumed that the variables presenting climate vary in time while the general model structure and model parameters representing catchment characteristics remain time-invariant”. There is a clear parallel of this statement with the approach presented herein where meteorological conditions vary in time, while climate-dependent (empirical) parameters are often fixed values.

In practice, long time series of observed evaporation are rare and not evenly distributed spatially. As such, for many applications, hydrologists must rely on incomplete calibration



data, use analogous stations with similar characteristics, or simply default to published values for crop factors and  $E_{\text{ref}}$  model parameters. Such published values for empirical factors of the different  $E_{\text{ref}}$  methods (Table 1) are  $C_1=0.65$ ,  $C_0=0$  (De Bruin, 1987),  $\alpha'=1.3$ ,  $\beta=0$  (De Bruin and Lablans, 1998),  $a=0.0023$ ,  $b=17.8$  (Droogers and Allen, 2002),  $c=0$ ,  $d=1$  (Sperna Weiland et al., 2012). Besides absolute values of the calibrated empirical factors, our analysis provides insight into the sensitivity of the results, i.e. the parameter values, to the calibration period. Calibrated model parameters for 30 year calibration periods are (standard deviations between brackets):  $C_1=0.64(0.01)$ ,  $C_0=0.37(0.03)$ ,  $\alpha'=1.06(0.01)$ ,  $\beta=0.57(0.02)$ ,  $a=0.0022(0.002)$ ,  $b=20.7(3.0)$ , and  $c=-2.31(0.06)$ ,  $d=1.73(0.03)$ . For a more realistic calibration period of e.g. three years, the standard deviations increase by a factor 2-10, depending on the  $E_{\text{ref}}$  method used.

For the Netherlands, published crop factors ( $K_{\text{tot}}$  for  $E_{\text{ref\_Mak}}$  with coefficients  $C_1 = 0.65$  and  $C_0 = 0$ ) are 1.0, 0.8, 1.1, 1.2 and 1.3 for grass, heather, deciduous forest, pine forest and spruce forest, respectively (Spieksma et al., 1996). Values from our study for e.g. the mean 30 year  $K_{\text{tot}}$  values, were 0.8, 1.03, 1.02, 1.07 and 1.25, respectively. Problems arise on the applicability of the published and frequently re-used crop factors, as the climatic conditions used for fitting are rarely documented. The analysis herein provides insight in the uncertainty ranges that could be expected using published empirical coefficients (Figure 8) and their potential impact on simulated potential evaporation components.

This study has shown that potential evaporation estimates are most accurate and stable with a long calibration period. However, even when using a long observed record, estimates may include errors due to the assumption of constant empirical coefficients in a non-stationary climate, i.e. the calibration period not being representative of current conditions. Evaporation estimates outside the calibration period are even more susceptible to non-stationarity when the calibration period is relatively short, as with areas where observed evaporation data are sparse. Finally, estimating evaporation based on published typical values without calibration is most susceptible to errors, as these parameters are typically global averages but also contain the non-stationary reference period issues identified in this paper. To remove bias by systematic input errors, as in e.g. evaporation, it is common practice to tune models by calibration (Ehret et al. (2014) and references therein). Although model calibration may compensate for biased input data, resulting in more accurate results and comparable model efficiencies, such calibration limits the general applicability of models when the bias is not

constant over time (Andréassian et al., 2004). Figs. 4 and 7 show that such non-constant bias occurs for both  $E_{\text{ref}}$  and potential  $E$  estimates, thus limits their application outside the calibration range.

Although extrapolations to future periods will always include uncertainty, it is important to quantify and limit this uncertainty. This analysis provides such a quantification, identifying the sensitivity of evaporation estimates to extrapolation and representing information on ways to reduce potential errors; e.g. Figure 7 quantifies the error that is made in extrapolations. A similar modeling approach as presented here could be used to identify climate induced changes in potential evaporation components. Moreover, such information can be used to reduce potential errors, if the errors can be explained from e.g. differences in climatic conditions between the periods of calibration and application. We did so for the errors identified for potential evaporation in Figure 7, and found that for all  $E_{\text{ref}}$  methods except for Har, and for all vegetation classes, the error correlates well with differences in relative humidity ( $R^2 > 0.78$ ) (Figure 3). This makes the errors predictable, and provides opportunities to correct for them.

Although we advocate using process-based evaporation simulations where possible, it should be emphasized that the two-step approach still can be a valuable concept, especially in regions with limited data availability. However, some considerations may strengthen the robustness of the two-step approach. First, our results show that applying radiation based methods are preferred over temperature based methods. Second, ideally, independent of the type of empirical method used, the coefficients should be recalibrated against measurements. Third, as such recalibration will practically often not be feasible, we advocate to identify changes in climatic conditions for the period of application and the calibration period, and to quantify using a sensitivity analysis, how they may impact potential evaporation estimates. This provides uncertainty ranges that advance the interpretation of modeling exercises.

## **4.2 Implications for climate change impact studies**

Poor transferability of parameter estimates made during calibration can have potentially large impacts for studies in non-stationary conditions (Coron et al., 2014), e.g. for climate change impact studies (Bormann, 2011; Karlsson et al., 2014). To improve the temporal robustness of hydrological modeling, Coron et al. (2014) propose, while adding it to the framework of the new IAHS Scientific Decade “Panta Rhei” (Montanari et al., 2013), to especially advance our

abilities to estimate temporal variations in evaporation fluxes. This study contributes to this larger objective.

For climate change impact studies, applications of empirical models are particularly problematic, as empirical methods closely approximate observations of natural processes, but do not capture the underlying physics. When extrapolating to new climate regimes, these assumptions are not guaranteed to remain valid (Kay and Davies, 2008; Bormann, 2011; Arnell, 1999). Similar to our findings, simulating historic non-stationary climatic conditions, Kay and Davies (2008) demonstrate that  $E_{\text{ref\_PM}}$  and temperature based  $E_{\text{ref}}$  methods give different projected evaporation estimates when applied to future climate model data. Additionally, Haddeland et al. (2011) show, using the WATCH climate forcing data (Weedon et al., 2011), that global hydrological models that differ in their choice of evaporation schemes, show significantly different evaporation estimates. These large discrepancies in an important part of the water cycle may have a large effect on the modeled hydrological impacts of climate change and increases the uncertainty of impact estimates (Bormann, 2011; Kay and Davies, 2008; Haddeland et al., 2011).

To show the implications of using different empirical  $E_{\text{ref}}$  methods in hydrological applications under recent climate change, without the need for numerous extensive model runs, we calculated the Standardized Precipitation and Evaporation Index (SPEI) for the period 1906-2013, with the empirical coefficients calibrated for the 30-year period 1906-1935.

The SPEI (Beguería et al., 2013; Vicente-Serrano et al., 2010) is a commonly used meteorological drought index, which is a variant of the WMO-recommended Standardized Precipitation Index SPI (Guttman, 1999; Hayes et al., 2011; McKee et al., 1993). Unlike the SPI, which calculates precipitation accumulated over a period and then normalizes the accumulated value based on typical seasonal conditions, the SPEI instead normalizes the accumulated difference of the climatic water balance, defined as the difference between precipitation and  $E_{\text{ref}}$ . This produces a time series of normalized values, such that an SPEI of 0 refers to typical conditions, an SPEI of negative one refers to a condition where  $\Sigma(P - E_{\text{ref}})$  is one standard deviation drier than typical, and vice versa for positive one. For this example, the SPEI6 was calculated, normalizing the climatic water balance summed over the preceding six months, following the fitting procedures outlined in Stagge et al. (2014a) and Gudmundsson and Stagge (2014).

Fig. 11 shows the results of this analysis, with the assumed accurate SPEI6, based on  $E_{\text{ref\_PM}}$ , shown at the top and the difference between this and SPEI6 for all other empirical reference evaporation models shown below. As with the results of  $E_{\text{ref}}$  simulations, the Mak and PT models are closest to the observed signal (differences in the range of -0.2 to 0.2), while the Har and BC models produce greater variability ( $\Delta\text{SPEI6} = -0.5$  to 0.5). Differences of this magnitude can make a large difference when interpreting drought risk. For example, the year 1947 produced a severe drought at the De Bilt site ( $\text{SPEI6} = -2.2$ ); however all other methods underestimate  $E_{\text{ref}}$ , producing SPEI6 values between -1.5 and -1.9. This in turn, changes the interpretation of this drought from an event expected to occur once every 72 years to an event expected to occur once every 15-35 years. This is a significant difference in risk level which can be attributed to differences among the evaporation methods and a potentially non-representative calibration period. SPEI sensitivity to  $E_{\text{ref}}$  method is analyzed in greater detail in Stagge et al. (2014b).

## 5 Conclusion

In this study we thoroughly analyzed the robustness of the two-step approach to simulate potential evaporation. We quantified the magnitude of the systematic errors that may be introduced when empirical coefficients are applied outside their calibration period, depending on differences in climate, the period, and the length of the calibration period. Our hydrological models are to varying extent regression models, which limits their general applicability, and the estimation of potential evaporation is closely linked to climate variability. With our analysis, we want to raise awareness and to provide a quantification of possible systematic errors that may be introduced in estimates of potential evaporation and in hydrological modeling studies due to straightforward application of i) the common two-step approach for potential evaporation specifically, and ii) fixed instead of time-variant model parameters in general.

## 6 Acknowledgements

This study was carried out under the auspices of the joint research program of the Dutch Water Utility sector. We would like to thank the three anonymous reviewers for their valuable suggestions to improve the manuscript.

## 2 **References**

- 3 Abbott, M. B., Bathurst, J. C., Cunge, J. A., O'Connell, P. E., and Rasmussen, J.: An  
4 introduction to the European Hydrological System -- Systeme Hydrologique Europeen,  
5 "SHE", 2: Structure of a physically-based, distributed modelling system, J Hydrol, 87, 61-77,  
6 1986.
- 7 Allen, R. G., and Pruitt, W. O.: Rational use of the FAO Blaney-Criddle formula, Journal of  
8 Irrigation and Drainage Engineering, 112, 139-155, 1986.
- 9 Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration - Guidelines for  
10 computing crop water requirements FAO Irrigation and drainage paper, FAO - Food and  
11 Agriculture Organization of the United Nations, Rome, 1998.
- 12 Allen, R. G.: Using the FAO-56 dual crop coefficient method over an irrigated region as part  
13 of an evapotranspiration intercomparison study, J Hydrol, 229, 27-41, 2000.
- 14 Allen, R. G., Pereira, L. S., Smith, M., Raes, D., and Wright, J. L.: FAO-56 Dual crop  
15 coefficient method for estimating evaporation from soil and application extensions, Journal of  
16 Irrigation and Drainage Engineering, 1, 2-13, 2005.
- 17 Allen, R. G., Pereira, L. S., Howell, T. A., and Jensen, M. E.: Evapotranspiration information  
18 reporting: I. Factors governing measurement accuracy, Agric Water Manag, 98, 899-920,  
19 <http://dx.doi.org/10.1016/j.agwat.2010.12.015>, 2011.
- 20 Andréassian, V., Perrin, C., and Michel, C.: Impact of imperfect potential evapotranspiration  
21 knowledge on the efficiency and parameters of watershed models, J Hydrol, 286, 19-35, 2004.
- 22 Arnell, N. W.: The effect of climate change on hydrological regimes in Europe: a continental  
23 perspective, Global Environmental Change, 9, 5-23, [http://dx.doi.org/10.1016/S0959-](http://dx.doi.org/10.1016/S0959-3780(98)00015-6)  
24 [3780\(98\)00015-6](http://dx.doi.org/10.1016/S0959-3780(98)00015-6), 1999.
- 25 Beguería, S., Vicente-Serrano, S. M., Reig, F., and Latorre, B.: Standardized precipitation  
26 evapotranspiration index (SPEI) revisited: parameter fitting, evapotranspiration models, tools,  
27 datasets and drought monitoring, International Journal of Climatology, doi: 10.1002/joc.3887,  
28 2013.
- 29 Bergström, S., and Forsman, A.: Development of a conceptual deterministic rainfall-runoff  
30 model, Nord Hydrol, 4, 147-170, 1973.
- 31 Blaney, H. F., and Criddle, W. P.: Determining water requirements in irrigated areas from  
32 climatological and irrigation data, USDA (SCS), 48, 1950.
- 33 Bormann, H.: Sensitivity analysis of 18 different potential evapotranspiration models to  
34 observed climatic change at German climate stations, Clim Change, 104, 729-753,  
35 [10.1007/s10584-010-9869-7](http://dx.doi.org/10.1007/s10584-010-9869-7), 2011.
- 36 Braden, H.: Ein Energiehaushalts- und Verdunstungsmodell for Wasser und  
37 Stoffhaushaltsuntersuchungen landwirtschaftlich genutzter Einzugsgebiete, Mittlungen  
38 Deutsche Bodenkundliche Gesellschaft, 42, 294-299, 1985.
- 39 Bradford, R. B., Ragab, R., Crooks, S. M., Bouraoui, F., and Peters, E.: Simplicity versus  
40 complexity in modelling groundwater recharge in Chalk catchments, Hydrology and Earth  
41 System Sciences, 6, 927-937, [10.5194/hess-6-927-2002](http://dx.doi.org/10.5194/hess-6-927-2002), 1999.

1 Calder, I. R.: Assessing the water use of short vegetation and forests: Development of the  
2 Hydrological Land Use Change (HYLUC) model, *Wat Resour Res*, 39, 1318,  
3 10.1029/2003wr002040, 2003.

4 Coron, L., Andréassian, V., Perrin, C., Bourqui, M., and Hendrickx, F.: On the lack of  
5 robustness of hydrologic models regarding water balance simulation: a diagnostic approach  
6 applied to three models of increasing complexity on 20 mountainous catchments, *Hydrology  
7 and Earth System Sciences*, 18, 727-746, 10.5194/hess-18-727-2014, 2014.

8 De Bruin, H. A. R., and Holtslag, A. A. M.: A simple parameterization of the surface fluxes  
9 of sensible and latent heat during daytime compared with the Penman-Monteith concept,  
10 *Journal of Applied Meteorology*, 21, 1610-1621, 10.1175/1520-  
11 0450(1982)021<1610:aspots>2.0.co;2, 1982.

12 De Bruin, H. A. R.: From Penman to Makkink, in: *Evaporation and weather*, edited by:  
13 Hooghart, C., Proceedings and Information. Comm. Hydrological Research TNO, The Hague,  
14 5-32, 1987.

15 De Bruin, H. A. R., and Lablans, W. N.: Reference crop evapotranspiration determined with a  
16 modified Makkink equation, *Hydrological Processes*, 12, 1053-1062, 1998.

17 De Lange, W. J., Prinsen, G. F., Hoogewoud, J. C., Veldhuizen, A. A., Verkaik, J., Oude  
18 Essink, G. H. P., van Walsum, P. E. V., Delsman, J. R., Hunink, J. C., Massop, H. T. L., and  
19 Kroon, T.: An operational, multi-scale, multi-model system for consensus-based, integrated  
20 water management and policy analysis: The Netherlands Hydrological Instrument,  
21 *Environmental Modelling & Software*, 59, 98-108,  
22 <http://dx.doi.org/10.1016/j.envsoft.2014.05.009>, 2014.

23 Doorenbos, J., and Pruitt, W. O.: *Crop water requirements*; Irrigation and drainage paper No.  
24 24, FAO, Rome, 1977.

25 Driessen, T. L. A., Hurkmans, R. T. W. L., Terink, W., Hazenberg, P., Torfs, P. J. J. F., and  
26 Uijlenhoet, R.: The hydrological response of the Ourthe catchment to climate change as  
27 modelled by the HBV model, *Hydrology and Earth System Sciences*, 14, 651-665,  
28 10.5194/hess-14-651-2010, 2010.

29 Droogers, P., and Allen, R. G.: Estimating reference evapotranspiration under inaccurate data  
30 conditions, *Irrigation and Drainage Systems*, 16, 33-45, 10.1023/a:1015508322413, 2002.

31 Ehret, U., Gupta, H. V., Sivapalan, M., Weijs, S. V., Schymanski, S. J., Blöschl, G., Gelfan,  
32 A. N., Harman, C., Kleidon, A., Bogaard, T. A., Wang, D., Wagener, T., Scherer, U., Zehe,  
33 E., Bierkens, M. F. P., Di Baldassarre, G., Parajka, J., van Beek, L. P. H., van Griensven, A.,  
34 Westhoff, M. C., and Winsemius, H. C.: Advancing catchment hydrology to deal with  
35 predictions under change, *Hydrology and Earth System Sciences*, 18, 649-671, 10.5194/hess-  
36 18-649-2014, 2014.

37 Farmer, W., Strzepek, K., Schlosser, C. A., Droogers, P., and Gao, X.: A method for  
38 calculating reference evapotranspiration on daily time scales, MIT Joint Program on the  
39 Science and Policy of Global Change, 2011.

40 Feddes, R. A.: Crop factors in relation to Makkink reference-crop evapotranspiration, in:  
41 *Evaporation and weather*, edited by: Hooghart, C., Proceedings and Information. Comm.  
42 Hydrological Research TNO, The Hague, 33-47, 1987.

1 Feddes, R. A., and Lenselink, K. J.: Evapotranspiration, in: Drainage principles and  
2 applications, edited by: Ritzema, H. P., International Institute for Land Reclamation and  
3 Improvement ILR, Wageningen, 1994.

4 Federer, C. A., Vörösmarty, C., and Fekete, B.: Intercomparison of methods for calculating  
5 potential evaporation in regional and global water balance models, *Water Resour. Res.*, 32,  
6 2315-2321, 10.1029/96wr00801, 1996.

7 Gash, J. H. C., Lloyd, C. R., and Lachaud, G.: Estimating sparse forest rainfall interception  
8 with an analytical model, *J Hydrol*, 170, 79-86, 1995.

9 Gudmundsson, L., and Stagge, J. H.: SCI: Standardized Climate Indices such as SPI, SRI or  
10 SPEI. R package version 1.0-1, 2014.

11 Guerschman, J. P., Van Dijk, A. I. J. M., Mattersdorf, G., Beringer, J., Hutley, L. B., Leuning,  
12 R., Pipunic, R. C., and Sherman, B. S.: Scaling of potential evapotranspiration with MODIS  
13 data reproduces flux observations and catchment water balance observations across Australia,  
14 *J Hydrol*, 369, 107-119, <http://dx.doi.org/10.1016/j.jhydrol.2009.02.013>, 2009.

15 Guttman, N. B.: Accepting the Standardized Precipitation Index: A calculation algorithm, in,  
16 Wiley Online Library, 1999.

17 Haddeland, I., Clark, D. B., Franssen, W., Ludwig, F., Voß, F., Arnell, N. W., Bertrand, N.,  
18 Best, M., Folwell, S., Gerten, D., Gomes, S., Gosling, S. N., Hagemann, S., Hanasaki, N.,  
19 Harding, R., Heinke, J., Kabat, P., Koirala, S., Oki, T., Polcher, J., Stacke, T., Viterbo, P.,  
20 Weedon, G. P., and Yeh, P.: Multimodel estimate of the global terrestrial water balance:  
21 Setup and first results, *Journal of Hydrometeorology*, 12, 869-884, 10.1175/2011jhm1324.1,  
22 2011.

23 Hayes, M., Svoboda, M., Wall, N., and Widhalm, M.: The Lincoln declaration on drought  
24 indices: universal meteorological drought index recommended, *Bull Am met Soc*, 92, 485-  
25 488, doi: 10.1175/2010BAMS3103.1, 2011.

26 Hooghart, J. C., and Lablans, W. N.: van Penman naar Makkink: een nieuwe  
27 berekeningswijze voor de klimatologische verdampingsgetallen, KNMI, De Bilt, 1988.

28 Hupet, F., and Vanclooster, M.: Effect of the sampling frequency of meteorological variables  
29 on the estimation of the reference evapotranspiration, *J Hydrol*, 243, 192-204,  
30 [http://dx.doi.org/10.1016/S0022-1694\(00\)00413-3](http://dx.doi.org/10.1016/S0022-1694(00)00413-3), 2001.

31 Hurkmans, R. T. W. L., Terink, W., Uijlenhoet, R., Moors, E. J., Troch, P. A., and Verburg,  
32 P. H.: Effects of land use changes on streamflow generation in the Rhine basin, *Wat Resour*  
33 *Res*, 45, W06405, 10.1029/2008wr007574, 2009.

34 Jones, P. D., and Moberg, A.: Hemispheric and large-scale surface air temperature variations:  
35 An extensive revision and an update to 2001, *Journal of Climate*, 16, 2003.

36 Karlsson, I. B., Sonnenborg, T. O., Jensen, K. H., and Refsgaard, J. C.: Historical trends in  
37 precipitation and stream discharge at the Skjern River catchment, Denmark, *Hydrology and*  
38 *Earth System Sciences*, 18, 595-610, 10.5194/hess-18-595-2014, 2014.

39 Kay, A. L., and Davies, H. N.: Calculating potential evaporation from climate model data: A  
40 source of uncertainty for hydrological climate change impacts, *J Hydrol*, 358, 221-239,  
41 <http://dx.doi.org/10.1016/j.jhydrol.2008.06.005>, 2008.

1 Kay, A. L., Bell, V. A., Blyth, E. M., Crooks, S. M., Davies, H. N., and Reynard, N. S.: A  
2 hydrological perspective on evaporation: historical trends and future projections in Britain,  
3 Journal of Water and Climate Change, 4, 193-208, doi:10.2166/wcc.2013.014, 2013.

4 Klein Tank, A. M. G., and Lenderink, G.: Klimaatverandering in Nederland; Aanvullingen op  
5 de KNMI'06 scenario's, KNMI, De Bilt, 2009.

6 Kroes, J. G., Van Dam, J. C., Groenendijk, P., Hendriks, R. F. A., and Jacobs, C. M. J.:  
7 SWAP version 3.2, Theory description and user manual, Wageningen University and  
8 Research Centre, WageningenAlterra report 1649 (update 02), 2009.

9 Lenderink, G., Buishand, A., and van Deursen, W.: Estimates of future discharges of the river  
10 Rhine using two scenario methodologies: direct versus delta approach, Hydrology and Earth  
11 System Sciences, 11, 1145-1159, 10.5194/hess-11-1145-2007, 2007.

12 Makkink, G. F.: Testing the Penman formula by means of lysimeters, Journal of the  
13 Institution of Water Engineers, 11, 277-288, 1957.

14 McKee, T. B., Doesken, N. J., and Kleist, J.: The relationship of drought frequency and  
15 duration to time scales, Proceedings of the 8th Conference on Applied Climatology, 1993,  
16 179-183,

17 Montanari, A., Young, G., Savenije, H., Hughes, D., Wagener, T., Ren, L., Koutsoyiannis, D.,  
18 Cudennec, C., Toth, E., and Grimaldi, S.: "Panta Rhei—Everything Flows": Change in  
19 hydrology and society—The IAHS Scientific Decade 2013–2022, Hydrological Sciences  
20 Journal, 58, 1256-1275, 2013.

21 Monteith, J.: Evaporation and environment, Symp. Soc. Exp. Biol, 1965, 4,

22 NHI, N. H. I.: Modelrapportage; Deelrapport Gewassenmerken. Versie:  
23 NHI\FASE\_1+\2008\DR12\v2 2008.

24 Penman, H. L.: Natural evaporation from open water, bare soil and grass, Proceedings of the  
25 Royal Society of London. Series A. Mathematical and Physical Sciences, 193, 120-145,  
26 10.1098/rspa.1948.0037, 1948.

27 Penman, H. L.: Evaporation: An introductory survey, Neth J Agric Sci, 4, 9-29, 1956.

28 Sanchez-Lorenzo, A., Calbó, J., and Martin-Vide, J.: Spatial and temporal trends in sunshine  
29 duration over Western Europe (1938–2004), Journal of Climate, 21, 6089-6098,  
30 10.1175/2008jcli2442.1, 2008.

31 Savenije, H. H. G.: The importance of interception and why we should delete the term  
32 evapotranspiration from our vocabulary, Hydrological Processes, 18, 1507-1511,  
33 10.1002/hyp.5563, 2004.

34 Seibert, J.: Reliability of model predictions outside calibration conditions, Nord Hydrol, 34,  
35 477-492, 10.2166/nh.2003.028, 2003.

36 Shabalova, M. V., Deursen, W. P. A. v., and Buishand, T. A.: Assessing future discharge of  
37 the river Rhine using regional climate model integrations and a hydrological model, Clim  
38 Res, 23, 233-246, 10.3354/cr023233, 2003.

39 Shaw, S. B., and Riha, S. J.: Assessing temperature-based PET equations under a changing  
40 climate in temperate, deciduous forests, Hydrological Processes, 25, 1466-1478,  
41 10.1002/hyp.7913, 2011.



1 Shuttleworth, W. J.: Putting the 'vap' into evaporation, *Hydrology and Earth System Sciences*,  
2 11, 210-244, 2007.

3 Solomon, S., Qin, D., Manning, M., Alley, R. B., Berntsen, T., Bindoff, N. L., Chen, Z.,  
4 Chidthaisong, A., Gregory, J. M., Hegerl, G. C., Heimann, M., Hewitson, B., Hoskins, B. J.,  
5 Joos, F., Jouzel, J., Kattsov, V., Lohmann, U., Matsuno, T., Molina, M., Nicholls, N.,  
6 Overpeck, J., Raga, G., Ramaswamy, V., Ren, J., Rusticucci, M., Somerville, R., Stocker, T.  
7 F., Whetton, P., Wood, R. A., and Wratt, D.: Technical Summary, in: *Climate Change 2007:*  
8 *The physical science basis. Contribution of working group I to the fourth assessment report of*  
9 *the intergovernmental panel on climate change*, edited by: Solomon, S., Qin, D., Manning,  
10 M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge  
11 University Press, Cambridge, United Kingdom and New York, NY, USA, 2007.

12 Sperna Weiland, F. C., Tisseuil, C., Dürr, H. H., Vrac, M., and van Beek, L. P. H.: Selecting  
13 the optimal method to calculate daily global reference potential evaporation from CFSR  
14 reanalysis data for application in a hydrological model study, *Hydrology and Earth System*  
15 *Sciences*, 16, 983-1000, 10.5194/hess-16-983-2012, 2012.

16 Spijksma, J. F. M., Dolman, A. J., and Schouwenars, J. M.: De parameterisatie van de  
17 verdamping van natuurterreinen in hydrologische modellen, 1996.

18 Stagge, J. H., Tallaksen, L. M., L., G., Van Loon, A. F., and Stahl, K.: Pan-European  
19 comparison of candidate distributions for climatological drought indices (SPI and SPEI),  
20 *International Journal of Climatology*, In Review, 2014a.

21 Stagge, J. H., Tallaksen, L. M., Xu, C.-Y., and Van Lanen, H. A. J.: Standardized  
22 precipitation-evapotranspiration index (SPEI): Sensitivity to potential evapotranspiration  
23 model and parameters. *Proceedings of FRIEND-Water 2014 (IAHS Publ. 363, 2014).*  
24 Montpellier, France, October 2014., 2014b,

25 Stanhill, G.: A perspective on global warming, dimming, and brightening, *Eos, Transactions*  
26 *American Geophysical Union*, 88, 58-58, 10.1029/2007eo050007, 2007.

27 Suo, L., Otterå, O. H., Bentsen, M., Gao, Y., and Johannessen, O. M.: External forcing of the  
28 early 20th century Arctic warming, *Tellus A*, 65, 2013.

29 Tiktak, A., and Bouten, W.: Soil water dynamics and long-term water balances of a Douglas  
30 fir stand in the Netherlands, *J Hydrol*, 156, 265-283, [http://dx.doi.org/10.1016/0022-](http://dx.doi.org/10.1016/0022-1694(94)90081-7)  
31 [1694\(94\)90081-7](http://dx.doi.org/10.1016/0022-1694(94)90081-7), 1994.

32 Trambauer, P., Dutra, E., Maskey, S., Werner, M., Pappenberger, F., van Beek, L. P. H., and  
33 Uhlenbrook, S.: Comparison of different evaporation estimates over the African continent,  
34 *Hydrology and Earth System Sciences*, 18, 193-212, 10.5194/hess-18-193-2014, 2014.

35 Van Dam, J. C., Groenendijk, P., Hendriks, R. F. A., and Kroes, J. G.: Advances of modeling  
36 water flow in variably saturated soils with SWAP, *Vadose Zone Journal*, 7, 640-653,  
37 DOI:10.2136/vzj2007.0060, 2008.

38 Van den Hurk, B., Klein Tankink, A., Lenderink, G., Van Ulden, A., Van Oldenborgh, G. J.,  
39 Katsman, C., Van den Brink, H., Keller, F., Bessembinder, J., Burgers, G., Komen, G.,  
40 Hazeleger, W., and Drijfhout, S.: KNMI Climate change scenarios 2006 for the Netherlands,  
41 KNMI, De Bilt, 82 pp., 2006.

42 Van Roosmalen, L., Sonnenborg, T. O., and Jensen, K. H.: Impact of climate and land use  
43 change on the hydrology of a large-scale agricultural catchment, *Water Resour. Res.*, 45,  
44 doi:10.1029/2007WR006760, 2009.

- 1 Van Walsum, P. E. V., and Supit, I.: Influence of ecohydrologic feedbacks from simulated  
2 crop growth on integrated regional hydrologic simulations under climate scenarios,  
3 *Hydrology and Earth System Sciences*, 16, 1577-1593, 10.5194/hess-16-1577-2012, 2012.
- 4 Vázquez, R. F., and Feyen, J.: Effect of potential evapotranspiration estimates on effective  
5 parameters and performance of the MIKE SHE-code applied to a medium-size catchment, *J*  
6 *Hydrol*, 270, 309-327, 2003.
- 7 Vicente-Serrano, S. M., Beguería, S., and López-Moreno, J. I.: A multiscalar drought index  
8 sensitive to global warming: the standardized precipitation evapotranspiration index, *Journal*  
9 *of Climate*, 23, 1696-1718, 2010.
- 10 Von Hoyningen-Hüne, J.: *Die Interception des Niederschlags in landwirtschaftlichen*  
11 *Beständen*, 1983.
- 12 Wallace, J. S.: Calculating evaporation: resistance to factors, *Agric For Meteorol*, 73, 353-  
13 366, DOI:10.1016/0168-1923(94)05084-J, 1995.
- 14 Wang, K., and Dickinson, R. E.: Contribution of solar radiation to decadal temperature  
15 variability over land, *Proceedings of the National Academy of Sciences*, 110, 14877-14882,  
16 10.1073/pnas.1311433110, 2013.
- 17 Ward, P. J., Renssen, H., Aerts, J. C. J. H., van Balen, R. T., and Vandenberghe, J.: Strong  
18 increases in flood frequency and discharge of the River Meuse over the late Holocene:  
19 impacts of long-term anthropogenic land use change and climate variability, *Hydrology and*  
20 *Earth System Sciences*, 12, 159-175, 10.5194/hess-12-159-2008, 2008.
- 21 Weedon, G. P., Gomes, S., Viterbo, P., Shuttleworth, W. J., Blyth, E., Österle, H., Adam, J.  
22 C., Bellouin, N., Boucher, O., and Best, M.: Creation of the WATCH forcing data and its use  
23 to assess global and regional reference crop evaporation over land during the Twentieth  
24 Century, *Journal of Hydrometeorology*, 12, 823-848, 10.1175/2011jhm1369.1, 2011.
- 25 Wild, M., Gilgen, H., Roesch, A., Ohmura, A., Long, C. N., Dutton, E. G., Forgan, B., Kallis,  
26 A., Russak, V., and Tsvetkov, A.: From dimming to brightening: Decadal changes in solar  
27 radiation at earth's surface, *Science*, 308, 847-850, 10.1126/science.1103215, 2005.
- 28 Wild, M.: Global dimming and brightening: A review, *Journal of Geophysical Research:*  
29 *Atmospheres*, 114, D00D16, 10.1029/2008jd011470, 2009.
- 30 Witte, J. P. M., Runhaar, J., van Ek, R., van der Hoek, D. C. J., Bartholomeus, R. P.,  
31 Batelaan, O., van Bodegom, P. M., Wassen, M. J., and van der Zee, S. E. A. T. M.: An  
32 ecohydrological sketch of climate change impacts on water and natural ecosystems for the  
33 Netherlands: bridging the gap between science and society, *Hydrology and Earth System*  
34 *Sciences*, 16, 3945-3957, 10.5194/hess-16-3945-2012, 2012.
- 35 Yamanouchi, T.: Early 20th century warming in the Arctic: A review, *Polar Science*, 5, 53-71,  
36 <http://dx.doi.org/10.1016/j.polar.2010.10.002>, 2011.
- 37 Zhou, M. C., Ishidaira, H., Hapuarachchi, H. P., Magome, J., Kiem, A. S., and Takeuchi, K.:  
38 Estimating potential evapotranspiration using Shuttleworth-Wallace model and NOAA-  
39 AVHRR NDVI data to feed a distributed hydrological model over the Mekong River basin, *J*  
40 *Hydrol*, 327, 151-173, 2006.

## 1 Tables

2 Table 1: Equations used to calculate daily values of reference evaporation  $E_{\text{ref}}$  [mm/d] for the  
3 period 1906-2013 at De Bilt meteorological station.

Abbreviation	Method	Equation
$E_{\text{ref\_PM}}$	Penman-Monteith (Monteith, 1965)	$E = \frac{1}{\lambda} \left( \frac{\Delta(R_n - G) + \rho_a c_p \frac{(e_s - e_a)}{r_a}}{\Delta + \gamma \left( 1 + \frac{r_s}{r_a} \right)} \right)$
$E_{\text{ref\_Mak}}$	Makkink (Makkink, 1957)	$E = \frac{1}{\lambda} \left( C_1 \frac{\Delta}{\Delta + \gamma} R_s + C_0 \right)$
$E_{\text{ref\_PT}}$	Modified Priestley-Taylor (De Bruin and Holtslag, 1982)	$E = \frac{1}{\lambda} \left( \alpha' \frac{\Delta(R_n - G)}{\Delta + \gamma} + \beta \right)$
$E_{\text{ref\_Har}}$	Hargreaves (Droogers and Allen, 2002; Farmer et al., 2011)	$E = \frac{1}{\lambda} \left( a R_a (\bar{T} + b) (T_{\text{max}} - T_{\text{min}})^{0.5} \right)$
$E_{\text{ref\_BC}}$	Blaney and Criddle (1950) as in Allen and Pruitt (1986)	$E = \frac{1}{\lambda} \left( c + d \left( p (0.46 \bar{T} + 8.13) \right) \right)$

Where  $E$  = evaporation [ $\text{kg d}^{-1} \text{m}^{-2}$ ],  $\lambda$  = latent heat of vaporization [ $\text{MJ kg}^{-1}$ ],  $\Delta$  = slope of the vapor pressure curve [ $\text{kPa } ^\circ\text{C}^{-1}$ ],  $R_a$  = extraterrestrial radiation [ $\text{MJ m}^{-2} \text{day}^{-1}$ ],  $R_s$  = solar radiation [ $\text{MJ m}^{-2} \text{day}^{-1}$ ],  $R_n$  = net radiation [ $\text{MJ m}^{-2} \text{day}^{-1}$ ],  $G$  = soil heat flux [ $\text{MJ m}^{-2} \text{day}^{-1}$ ],  $\rho_a$  = mean air density [ $\text{kg m}^{-3}$ ],  $c_p$  = specific heat of the air [ $\text{MJ kg}^{-1} ^\circ\text{C}^{-1}$ ],  $\gamma$  = psychrometric constant [ $\text{kPa } ^\circ\text{C}^{-1}$ ],  $r_s$  = surface resistance [ $\text{s m}^{-1}$ ],  $r_a$  = aerodynamic resistance [ $\text{s m}^{-1}$ ],  $(e_s - e_a)$  = saturation vapour pressure deficit [ $\text{kPa}$ ],  $\bar{T}$ ,  $T_{\text{max}}$  and  $T_{\text{min}}$  = mean, maximum and minimum temperature [ $^\circ\text{C}$ ],  $p$  = mean daily percentage of annual daytime hours [%].  $C_1$ ,  $C_0$ ,  $\alpha'$ ,  $\beta$ ,  $a$ ,  $b$ ,  $c$ ,  $d$  are the coefficients adjusted in the calibration.

4

5

1 Table 2: Equations used to calculate monthly average crop factors for each vegetation class  
 2 and  $E_{\text{ref}}$  method.

Crop factor	Description	Equation
$K_t(E_{\text{ref}})$	crop factor for potential transpiration	$K_t = E_t / E_{\text{ref}}$
$K_{\text{ts}}(E_{\text{ref}})$	crop factor for potential transpiration + soil evaporation	$K_{\text{ts}} = (E_t + E_s) / E_{\text{ref}}$
$K_{\text{tot}}(E_{\text{ref}})$	crop factor for total evaporation	$K_{\text{tot}} = E_{\text{tot}} / E_{\text{ref}}$

3

4

# 1 Figures






Figure 1: Yearly and 30-year moving average  $E_{\text{ref}}$  according to Penman-Monteith for De Bilt, the Netherlands and projected  $E_{\text{ref}}$  values for the period 2036-2065. Projections are based on national climate scenarios (Van den Hurk et al., 2006) developed by the Royal Netherlands Meteorological Institute (KNMI). Two of the scenarios have been found to be most likely (Klein Tank and Lenderink, 2009) and are presented here: scenario W (blue line) and W+ (red line). Both comprise a +2K global temperature increase, but with respectively unchanged and changed (+) air circulation patterns in summer and winter. The scenarios were used to transfer the climatic conditions of 1976-2005 to the period 2036-2065 (Van den Hurk et al., 2006).




2

3 Figure 2: Flow chart of the methodology followed.  $E_{ref\_x} = E_{ref}$  of the empirical methods Mak,  
 4 Har, PT and BC; R = number of reference evaporation methods, V = number of vegetation  
 5 classes, C = number of evaporation components. For the explanation of the other  
 6 abbreviations we refer to the introduction.




1

2 Figure 3: Annual and 30-year moving average variables for De Bilt meteorological station. A:  
3 precipitation, B: mean temperature, C: mean relative humidity, D: sunshine duration, E: mean  
4 wind speed, F: global radiation, G: net radiation, H: vapour pressure deficit. A-E are  
5 observations, F-H are calculated following Allen et al. (1998).




1  
2 Figure 4:  $E_{ref}$  values for five methods for the period 1906-2013. Each empirical method  
3 calibrated on daily  $E_{ref\_PM}$  for the period 1906-1935. A: 30 year moving average  $E_{ref}$ , B:  
4 deviation of 30 year moving average  $E_{ref}$  from  $E_{ref\_PM}$ . C: yearly variability in  $E_{ref}$  for each  
5 method. D: yearly deviation of each  $E_{ref}$  with  $E_{ref\_PM}$ . The boxplots show the minimum, first  
6 quartile, median, third quartile, maximum and outliers of the annual data.






1

2 Figure 5: Maximum absolute deviation in 30-year average  $E_{ref}$  from 30-year average  $E_{ref\_PM}$   
 3 for the period 1984-2013, as a function of the length of the calibration period.



1  
2 Figure 6: Illustration of synthetic ‘observed’ potential evaporation components simulated with  
3 SWAP. The lines give monthly means over the period 1906-2013.  $E_t$  and  $E_s$  hold for a  
4 vegetation stand with a dry canopy only;  $E_{tot}$  includes interception.



1  
 2 Figure 7:  $E_t$  calculated for each vegetation class using each  $E_{ref}$  method and matching  $K_t$   
 3 calibrated on the 30-year period 1906-1935. Presented are 30-year moving averages in mm  
 4 (top), and deviations with the synthetic  $E_t$  for both the 30 year moving averages (center) and  
 5 annual values (bottom).




Figure 8: Standard deviation (sd) for  $K_t$  (A,D),  $K_{ts}$  (B,E) and  $K_{tot}$  (C,F) normalized to their mean values, for heather (A-C) and spruce (D-F), as function of the length of the calibration period.  $K$ -values are derived for each  $E_{ref}$  method;  $K$  and  $E_{ref}$  are calibrated on the same periods.





Figure 9: Maximum absolute deviation with synthetic ‘observations’ in mean  $E_t$  (A,D),  $E_t \& E_s$  (B,E) and  $E_{tot}$  (C,F) for the period 1984-2013, for heather (A-C) and spruce (D-F), obtained by the two-step approach, as function of the length of the calibration period. Presented as in Fig. 5, though using  $E_{ref}$  and crop factors ( $K_t$ ,  $K_{ts}$  and  $K_{tot}$ ) to derive  $E_t$ ,  $E_t \& E_s$  and  $E_{tot}$ .

1



2

3 Figure 10: Deviations with synthetic ‘observations’ in  $E_t$  (left),  $E_t \& E_s$  (centre) and  $E_{tot}$  (right)  
 4 for the last 30 year period (i.e. 1984-2013), due to different reference years and lengths of  
 5 calibration periods for both  $E_{ref}$  and  $K_t$ ,  $K_{ts}$  and  $K_{tot}$ . Results for PT and heather (top) and  
 6 spruce (bottom).




Figure 11: SPEI6 time series with  $E_{ref}$  based on PM (A). The subsequent figures show differences in SPEI with  $E_{ref}$  based on Mak (B), PT (C), Har (D) and BC (E), calibrated on the period 1906-1935.