

# Time-Series Analysis of the Long-Term Hydrologic Impacts of Afforestation in the Águeda Watershed of North-Central Portugal

D. Hawtree<sup>1</sup>, J. P. Nunes<sup>2</sup>, J. J. Keizer<sup>2</sup>, R. Jacinto<sup>2</sup>, J. Santos<sup>2</sup>, M. E. Rial-Rivas<sup>2</sup>, A. -K. Boulet<sup>2</sup>, F. Tavares-Wahren<sup>1</sup>, and K. -H. Feger<sup>1</sup>

[1]{Technische Universität Dresden, Institute of Soil Science & Site Ecology, Piener Str. 19 D-01737 Tharandt, Germany}

[2]{University of Aveiro, CESAM & Department of Environment and Planning, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal}

Correspondence to: D. Hawtree (daniel.hawtree@mailbox.tu-dresden.de)

## Abstract

The north-central region of Portugal has undergone significant land-cover change since the early 1900s, with large-scale replacement of natural vegetation types with plantation forests. This transition consisted of an initial conversion primarily to *Pinus pinaster*, followed by a secondary transition to *Eucalyptus globulus*. This land-cover change is likely to have altered the hydrologic functioning of this region; however these potential impacts are not fully understood. To contribute to a better understanding of the potential hydrologic impacts of this land cover change, this study examines the temporal trends in 75 years of data from the Águeda watershed (part of the Vouga Basin) over the period of 1936 to 2010. A number of hydrometeorological variables were analyzed using a combined Thiel-Sen / Mann-Kendall trend testing approach, to assess the magnitude and significance of patterns in the observed data. These trend tests indicated that there have been no significant reductions in streamflow over either the entire test period, or during sub-record periods, despite the large-scale afforestation which has occurred. This lack of change in streamflow is attributed to the specific characteristics of the watershed and land cover change. By contrast, a number of significant trends were found for baseflow index, with positive trends in the early data record (primarily during *Pinus pinaster* afforestation), followed by negative trends later in the data

1 record (primarily during *Eucalyptus globulus* afforestation). These trends are attributed to  
2 land-use and vegetation impacts on streamflow generating processes, both due to species  
3 differences and to alterations in soil properties (i.e. infiltration capacity, soil water  
4 repellency). These results highlight the importance of considering both vegetation  
5 types/dynamics and watershed characteristic when assessing hydrologic impacts, in particular  
6 with respect to soil properties.

7

## 8 **1 Introduction**

9 Water resource management is inherently tied to watershed-scale land dynamics, and proper  
10 management requires understanding how changes in land cover/use will impact hydrological  
11 processes (Calder, 2005). A key land cover type are forests, as changes in forest cover have  
12 the potential to significantly affect watershed-scale hydrologic processes, particularly by  
13 altering interception, evaporation, and streamflow. Changes in water availability due to  
14 afforestation/deforestation are driven by several factors controlling the water consumption of  
15 different vegetation species, in particular canopy interception and evapotranspiration rates,  
16 which are typically higher in tree species than in shrub and herbaceous species (Calder, 1998).

17 Meta-analyses of paired catchments studies have found that afforestation typically results in  
18 decreased streamflow while deforestation typically leads to increased streamflow (e.g. Bosch  
19 and Hewlett, 1982; Brown et al., 2005). However, the hydrologic response to deforestation is  
20 in general more consistent than the response to afforestation. This difference may be due to  
21 higher variability in land cover following afforestation compared to deforestation, and the  
22 effects of different transitional species and/or changes in forest physiology (Andréassian,  
23 2004). In a global synthesis of afforestation studies, Farley et al., (2005) found that  
24 afforestation of grasslands or shrublands will lead, on average, to reductions of one-third to  
25 two-thirds of streamflow, with these reductions occurring rapidly after planting (i.e. within  
26 the first 5 years) and reaching their maximum reduction 15 to 20 years following planting.

27 Changes in forest cover can also impact hydrologic processes by altering physical soil  
28 conditions, for example by reducing soil bulk density, increasing macro-porosity, or changing  
29 soil water repellency. Forested areas tend to have higher infiltration and groundwater recharge  
30 rates than alternate land cover types (e.g. Bruijnzeel, 2004). Higher infiltration rates will  
31 increase soil moisture levels, and therefore increase water availability as well as streamflow  
32 during dry periods (e.g. Scott and Lesch, 1997). The increased infiltration capacity of forested

1 areas may also help mitigate storm-driven peak flows, and therefore reduce potential flood  
2 damage; however, this effect may be subordinate to other watershed characteristics,  
3 particularly during severe flooding events (Calder, 2005; Wahren et al., 2012).

4 While the hydrologic impacts of forests at the watershed scale are generally well understood,  
5 predicting the effects of forest land-cover change for a specific watershed requires  
6 consideration of both the physical site conditions and vegetation types involved. In this  
7 respect, Andréassian (2004) identifies several prerequisite conditions that need to be met in  
8 order to observe hydrologic impacts at the watershed scale. These include climatic (i.e.  
9 periods of hydrologic surplus / deficit), pedological (i.e. soil depth) and eco-physiological (i.e.  
10 forest age-dependence) conditions.

11 The European Mediterranean region has undergone significant land cover changes over its  
12 long history of human habitation, which has left only an estimated 4.7 % of primary  
13 vegetation unaltered (Geri et al., 2010). These land cover changes are likely to have altered  
14 hydrologic processes at multiple scales, and the impacts of these changes are often not well  
15 understood. Gaining a better understanding of these past changes is critical for predicting the  
16 impact of future land-cover changes, particularly given widespread concerns over potential  
17 water shortages in this region due to changing temperature and rainfall regimes (Giorgi and  
18 Lionello, 2008). Some of the most significant land cover/use changes observed in the  
19 European Mediterranean region in recent decades have been: increased rural abandonment, a  
20 decrease in traditional agricultural/pastoral activities, and widespread planting of fast-growing  
21 tree species (Geri et al., 2010; Serra et al., 2008).

22 These regional trends are representative of the changes which have taken place in north-  
23 central Portugal, where traditional rural agrosilvopastoral activities have been widely replaced  
24 by plantations of the tree species *Pinus pinaster* and *Eucalyptus globulus* (Jones et al., 2012;  
25 Moreira et al., 2001). Both of these tree species have relatively high consumptive water  
26 demand and the potential to substantially reduce local water availability. Bosch and Hewlett  
27 (1982) estimated that pine and eucalypt forests cause an average decrease of over 40 mm/yr in  
28 water yield per 10 % change in land cover, while Farley et al. (2005) reported that  
29 afforestation with pines and eucalypts lead to reductions in streamflow of 40 % ( $\pm$  3 %) and  
30 75 % ( $\pm$  10 %), respectively. Rodríguez-Suárez et al. (2011) found that afforestation with  
31 *Eucalyptus globulus* caused a drop in water table depth as well as a decrease in streamflow

1 during the summer period, which they attributed to the higher transpiration capacity of the  
2 eucalypt plantations compared to the original crop lands.

3 In addition to consumptive water use through transpiration, evaporation from canopy  
4 interception is an important component of water use by Mediterranean forests. Interception  
5 rates have been found to vary widely in this region, depending on the tree species, canopy  
6 density, and climatic conditions. With respect to *Pinus pinaster*, Ferreira, (1996) reported  
7 interception rates of 15-18 % in the Águeda watershed of north-central Portugal (mean  
8 precipitation  $\approx$  1700 mm/yr), while (Valente et al., 1997) found similar rates of 17 % in a  
9 drier region of central Portugal (mean precipitation  $\approx$  600 mm/yr). For *Eucalyptus globulus*,  
10 both Ferreira (1996) and Valente et al. (1997) observed lower rates, amounting to 10-14 %  
11 and 11 %, respectively. By contrast, much higher interception rates have been found for other  
12 tree species in different parts of the Mediterranean, with values near and even exceeding 50  
13 %. For example, Scarascia-Mugnozza et al. (1988) found canopy interception rates of 68 %  
14 for a mature *Quercus cerris* forest in central Italy (mean precipitation 1006 mm/yr), Iovino et  
15 al. (1998) found rates of 58 % for a mature *Pinus negra* forest in southern Italy (mean  
16 precipitation 1179 mm/yr), and Tarazona et al. (1996) observed rates of 48 % for a mature  
17 *Pinus sylvestris* forest in northern Spain (long-term mean precipitation of 895 mm/y, 1253  
18 mm/yr during the study period).

19 A further hydrologic factor relevant to afforestation in north-central Portugal is the potential  
20 for impacts on soil water repellency (SWR). Both pine and eucalyptus tree species can  
21 promote SWR in the topsoil due to the considerable amount of resins, waxes, and aromatic  
22 oils contained in their organic matter (Benito and Santiago, 2003; Doerr and Thomas, 2000;  
23 Ferreira et al., 2000; Keizer et al., 2005a, 2005b). SWR is a key factor in triggering land  
24 degradation processes due to reductions in infiltration capacity and increased overland flow  
25 (Benito and Santiago, 2003; Doerr and Thomas, 2000; Keizer et al., 2005b; Shakesby et al.,  
26 2000). While in many regions SWR is associated primarily with post-fire soil conditions,  
27 Doerr et al. (1996) demonstrated that SWR is a widespread characteristic of both burned and  
28 unburned soils in the Águeda watershed during dry periods, in particular for stands of  
29 *Eucalyptus globulus*. Santos et al. (2013) examined temporal patterns in topsoil  
30 hydrophobicity in the Águeda watershed between July 2011 and June 2012 in unburnt pine  
31 and eucalypt plantations. Their findings suggested that the breakdown of SWR following dry  
32 summer conditions occurs through different mechanisms in the pine and eucalypt stands. In

1 the pine stands, SWR breakdown occurred from the top-down (i.e. vertically downwards),  
2 while in the eucalypt stands, breakdown occurred from the bottom-up (i.e. vertically  
3 upwards). Unpublished results indicated that this contrast reflected varying infiltration  
4 patterns, with infiltration occurring relatively slowly (i.e. matrix flow) in pine stands, as  
5 opposed to much faster (i.e. macropore flow) in eucalypt stands. This contrast in infiltration  
6 patterns appeared to be a product of SWR induced alterations in flow pathways.

7 Despite the well-documented potential for hydrologic impacts from afforestation in the  
8 Mediterranean region, there has been little investigation into the long-term effects in north-  
9 central Portugal. This is in part due to a lack of long-term streamflow records that allow for  
10 historical analyses. A notable exception to this lack of data is the Águeda watershed in the  
11 Caramulo Mountains, where streamflow data records are available from 1936 until the  
12 present.

13 Afforestation/deforestation studies typically focus on small paired watersheds, of which one  
14 has undergone fairly abrupt and well-recorded changes in land cover (e.g. Bosch and Hewlett,  
15 1982). By contrast, this study is conducted on a meso-scale watershed (404 km<sup>2</sup>), where  
16 afforestation has occurred progressively over an extended period of time. Furthermore, the  
17 present study case lacks a nearby watershed to serve as a paired site, which has a similarly  
18 long data record, similar physical-environmental characteristics, or a land use history without  
19 similar land cover changes (to serve as a control site).

20 To assess the hydrologic impacts of afforestation in the Águeda watershed, this study  
21 therefore adopts a data-driven and exploratory approach, which conducts multiple trend  
22 analyses on the 75-years of hydrometeorological data available from 1936 to 2010. This  
23 assessment is conducted over the entire data record as well as over multiple (overlapping)  
24 sub-periods for both annual and seasonal trends. The significant trends detected through this  
25 analysis are then considered with respect to the regional afforestation trends, and discussed in  
26 the context of previous field-studies conducted in this watershed. Therefore, the objective of  
27 this study is to apply a trend-testing methodology to a long-term data set in a watershed which  
28 has undergone progressive afforestation over a 75-year period, to assess what significant  
29 trends can be detected, and to relate these changes to the afforestation which has occurred  
30 there.

## 1 2 Methods

### 2 2.1 Watershed Description

3 The Águeda watershed is located in the Caramulo Mountains of north-central Portugal, east of  
4 the coastal city of Aveiro (Fig. 1). From the streamflow gauging point of Ponte Águeda, the  
5 watershed area is approximately 404 km<sup>2</sup>. The Águeda River is a left bank tributary to the  
6 Vouga River, which terminates at the coastal wetland of the Ria de Aveiro lagoon. This  
7 region of Portugal is categorized as a wet Mediterranean climate zone, with pronounced  
8 seasonal differences in temperature and precipitation between dry summer and wet winter  
9 seasons (Fig. 2). The Serra do Caramulo Mountains, which forms the source area of the  
10 Águeda river network, receives a substantial amount of annual rainfall, which can range from  
11 1 000 to 2 500 mm/yr. Topographically the landscape is dominated by steep hill-slopes with  
12 stony and shallow soils (< 0.5 m), which have a long history of anthropogenic impacts. These  
13 shallow soil were characterized by Ferreira et al. (2000) as stony, sandy loam, weakly  
14 structured Umbric Leptosols.

15 North-central Portugal has undergone substantial land cover/use changes over the past  
16 centuries, which have fundamentally altered the vegetative landscape of this region. From the  
17 1800s until the 1980s, the region had a general trend towards both increased agricultural and  
18 forest land cover, with reductions in natural vegetation types (e.g. Matos shrublands and  
19 mixed forests). This trend was primarily driven by the adoption of fertilizers and  
20 mechanization, as well as the abolition of feudal land systems (Estêvão, 1983; Jones et al.,  
21 2011; Silva et al., 2004). The period between 1930 and 1980 saw particularly rapid  
22 afforestation, due to incentives from the establishment of related government regulations and  
23 subsidies.

24 A key driver was the enactment of legislation in 1938 which encouraged afforestation of areas  
25 classified as “uncultivated/wasteland”, which often consisted of areas of matos (shrublands),  
26 mountain ranges, and sand dunes (Coelho et al., 1995; Estêvão, 1983; Ferreira et al., 2010;  
27 Jones et al., 2011; Silva et al., 2004). The primary species planted during this earlier period  
28 was *Pinus pinaster*, and beginning in the 1970s *Eucalyptus globulus* became the preferred  
29 species due to its faster growth and higher profitability for use in the paper pulp industry.  
30 During this period, eucalypt plantations began to replace pine forests as these were harvested,

1 as well as being widely introduced into remaining areas of shrublands and in recently burned  
2 areas (Jones et al., 2011).

3 Wildfire is another important factor in land cover/use change in Portugal, which has some of  
4 the highest rates of wildfire in Europe. Figure 3 shows the burned area of the Águeda  
5 watershed from 1975 to 2010, during which a total of 30 790 hectares burned, with some  
6 single years having wildfire over more than 10 % of the watershed (i.e. 1986 and 1995;  
7 Instituto da Conservação da Natureza e das Florestas, 2014). Wildfire can have significant  
8 short-term impacts on hydrologic functions in the study region, such as decreased infiltration  
9 and increased surface runoff / erosion (Malvar et al., 2011; Prats et al., 2012; Shakesby et al.,  
10 1993). In addition to these short-term impacts, wildfire can have potential long-term impacts  
11 by promoting changes in vegetation type. Wildfire has been a major driver of land-cover  
12 change in north-central Portugal in this respect, by allowing land-owners to convert from pine  
13 to eucalyptus plantations in the post-fire period.

14 This region-wide trend of the afforestation of shrubland with *Pinus pinaster*, followed by a  
15 secondary transition from *Pinus pinaster* to *Eucalyptus globulus* plantations, is representative  
16 of the land cover changes in the Vouga basin as a whole, and for the Águeda watershed in  
17 particular. From this regional pattern, and from afforestation maps of the Serra do Caramulo  
18 Mountains (Rego, 2001), a general timeline of land-cover change in the Águeda watershed  
19 during the period of investigation can be approximated, which is summarized in Table 1.

20 The current land cover in the Águeda watershed reflects this large-scale transition towards  
21 eucalyptus forests. According to the Corine Land Cover classification of 2006, approximately  
22 46 % of the watershed was covered by broad-leaf forest - which is predominantly eucalyptus  
23 (Corine Land Cover, 2010). Other land cover types with significant areal coverage in 2006  
24 include: 22 % mixed forest (mostly mixed stands of eucalypt and pine), 14 % agriculture, 10  
25 % pine forest, 6 % mato shrubland, 2 % urban, and 1% grasslands (Fig. 1).

## 26 **2.2 Hydrometeorological Data**

27 Hydrometeorological records for the Águeda watershed were compiled from hydrological  
28 year 1935/36 (i.e. Oct 1st 1935 to Sep 30th 1936) until hydrological year 2009/10, for the  
29 variables: precipitation, temperature, potential evapotranspiration, streamflow quantity,  
30 streamflow yield, baseflow quantity, and baseflow index. Table 2 provides an overview of the  
31 hydrometeorological variables used in this study.

1 Precipitation data were obtained from the rain-gauge “Campia”, of the ‘Sistema Nacional de  
2 Informação de Recursos Hídricos’ (SNIRH, 2013), which consists of 24 hour rainfall totals  
3 collected at 9:00 each day. The SNIRH provides a reliability ranking for the data in the range  
4 of 5 – 15, for which Campia is ranked as 14 (highly reliable). Data gaps occurred with the  
5 greatest frequency between 1997 and mid-2003, which were filled by linear regression with  
6 the nearby rain-gauges “Varzielas” ( $r^2 = 0.82$ ) and “Barragem de Castelo Burgães” ( $r^2 = 0.79$ ).

7 Temperature data was compiled using data from the gauge “Campia” of the Instituto  
8 Portugues do Mar e Atmosfera (IPMA, 2014). When data for “Campia” was not available, the  
9 time-series gaps were filled using linear regression with the temperature gauge “Coimbra” ( $r^2$   
10 = 0.93) which is part of the Global Historical Climate Network available at the National  
11 Climatic Data Center (NCDC).

12 Potential evapotranspiration (PET) was estimated using the Thornthwaite equation (1948),  
13 using the temperature data from the gauge “Campia”. The Thornthwaite equation was selected  
14 for PET, as opposed to more sophisticated equations (e.g. Hargreaves, Penman–Monteith), as  
15 there was insufficient data available over the entire time-series to support calculations from  
16 these equations. Another option considered for long-term PET values is the gridded Penman–  
17 Monteith based dataset available from the Climatic Research Unit (CRU) at the University of  
18 East Anglia (Harris et al., 2014). To assess the suitability of this dataset to the study  
19 watershed, the monthly PET values of the CRU data and the locally derived Thornwaite  
20 values were compared against locally derived Penman-Monteith values, over the period of  
21 January 2002 to September 2010. This assessment indicated that the locally derived  
22 Thornwaite values are better correlated than the gridded CRU dataset with local Penman-  
23 Monteith values. This may be due to the mountainous terrain of the study watershed, and the  
24 relatively large grid size of the CRU dataset (0.5 degrees) being unable to capture smaller  
25 scale impacts on PET. Based on this assessment, the locally derived Thornwaite PET values  
26 were identified as the most reliable and representative data source available for assessing  
27 long-term trends.

28 Streamflow data consists of daily average discharge measurements from the gauging station  
29 “Ponte Águeda” of the ‘Sistema Nacional de Informação de Recursos Hídricos’ (SNIRH,  
30 2013). This station was operational from June 1935 until the end of September 1990, and was  
31 then reactivated in October 1999. Streamflow for the interim period (1990/91 until 1998/99)  
32 was estimated by linear regression with the upstream gauges “Ribeiro” ( $r^2 = 0.76$ ) and “Ponte



1 Redonda” ( $r^2 = 0.75$ ). However, the streamflow estimates from the hydrologic years of  
2 1999/2000 through 2002/03 were eliminated from the dataset due to low data quality, owing  
3 to the absence of an adequate stage-discharge curve during this period.

4 In addition, a number of smaller streamflow gaps occurred throughout the daily streamflow  
5 dataset. When they occurred during periods with little or no precipitation, the gaps were filled  
6 by fitting a logarithmic decay curve (traditional linear reservoir with a semi-log fitting) to the  
7 streamflow recession. If gaps occurred during a precipitation event, then this approach was  
8 not applied and the gaps were left unfilled. If the number of gaps was greater than 5 % of the  
9 total record, then the entire period was removed from analysis, which was the case for the  
10 hydrologic years 1954/55 and 1975/76. Finally, data for the driest months of the year (i.e.  
11 June to September) during the period from before 1963 and after 2004 had very high  
12 uncertainty, due to unreported and variably occurring impoundments of streamflow during  
13 these months. Therefore, this four month period had to be removed from the streamflow  
14 analysis for the entire data record, to keep the inter-annual comparisons consistent. After the  
15 streamflow gaps were filled, the ratio of precipitation which becomes streamflow was  
16 calculated, to allow potential changes in the streamflow-precipitation relationship to be  
17 assessed. This ratio is defined as the “streamflow yield”, which is the total streamflow divided  
18 by total precipitation, with the period of summation determined by the period being  
19 considered (*i.e.* the annual or the seasonal ratio).

20 The final data set utilized in this study is a baseflow time-series calculated with the Eckhardt  
21 digital filter (Eckhardt, 2005) using the daily streamflow dataset. Baseflow corresponds to the  
22 portion of streamflow which does not come directly from a precipitation event, and can be  
23 used as a proxy of the sustained streamflow contribution from slow-flow. The relative  
24 proportion of baseflow from each day of streamflow was estimated, which was then  
25 aggregated to the time periods used for analysis. To assess the baseflow time-series calculated  
26 using the Eckhardt digital filter, a supplementary data set from 2001 to 2009 was also utilized,  
27 which calculates baseflow contribution using conductivity data from the SNIRH streamflow  
28 data using the ‘Conductivity Mass-Balance Method’ (Stewart et al., 2007).

### 29 **2.3 Thiel-Sen / Mann-Kendall Trend Testing Approach**

30 To examine the magnitude and significance of potential trends in the hydrometeorological  
31 time-series, a multi-step trend-testing approach was applied, following the general approach

1 presented in (Yue et al., 2002). This approach first determines the magnitude (i.e. slope) of  
2 any potential trend in the data using the non-parametric Thiel-Sen slope estimator (Sen,  
3 1968). This value is determined by calculating the median slope among the set generated  
4 between all sample points. This method also estimates the 95 % confidence intervals of the  
5 true slope, based on the set of slopes from the sample points, which provides a measure of  
6 uncertainty of the median Thiel-Sen value. If a potential trend is detected by the Thiel-Sen  
7 test (i.e. a non-zero slope), then the data is processed using the ‘Trend Free Pre-whitening’  
8 procedure of (Yue et al., 2002). This step reduces the over-estimation of significance which  
9 can occur in time-series data that exhibits positive serial correlation, as is typically the case  
10 for streamflow time-series data.

11 After the “Trend Free Pre-whitening procedure”, a Mann-Kendall test was applied to assess  
12 the statistical significance of any non-zero slope identified by the Thiel-Sen test. The Mann-  
13 Kendall test is a widely used, rank-based significance test, where the null hypothesis is that  
14 there is no trend in the observed data (Helsel, 1993). For this study, statistical significance  
15 was determined using an  $\alpha$  value of 0.05.

16 For each hydrometeorological variable, this trend testing procedure was applied over 12  
17 different time periods with varying start/end dates and lengths (Fig. 4). The longest period  
18 tested contains the entire 75-year data record (hydrologic year 1936-2010), followed by two  
19 periods of 50 years, three periods of 35 years, and six periods of 25 years. These overlapping  
20 periods of different lengths aim to thoroughly sample the potential range of years, while still  
21 allowing enough years of data to produce a robust significance test within each test period  
22 (i.e. a minimum of 25 years). Figure 4 provides an overview of the testing periods, and their  
23 temporal correspondence with the afforestation periods listed in Table 1.

24 When conducting multiple simultaneous hypothesis tests, it is necessary to correct for the  
25 false discovery rate (FDR). FDR corresponds to the expected proportion of incorrectly  
26 rejected null hypotheses, and therefore a method is needed to reduce the chance of receiving  
27 false-positive results (i.e. type I errors). A number of different methods can be applied to  
28 control for FDR, however given the overlapping time periods examined in this study, a  
29 method is needed which can deal with FDR under the assumption of positive dependence.  
30 Therefore, the Benjamini–Hochberg–Yekutieli procedure was applied to the trend-testing  
31 output from each individual ‘analysis set’ (Benjamini and Yekutieli, 2001). An analysis set  
32 corresponds to a group of tests which are expected to exhibit mutual positive dependence,

1 which in this case are the 12 overlapping periods over which each hydrometeorological  
2 variable was tested for the different annual and seasonal periods (i.e. Fig. 4 for a given  
3 variable and period).

4 Over the time periods shown in Fig. 4, the trend testing was conducted over both annual and  
5 seasonal time periods. The seasonal breakdown corresponds to the prevailing precipitation  
6 patterns of the study site, which consists of: a “Wet Season” from October to January when  
7 the largest amount of precipitation occurs, a “Transitional Season” from February to May  
8 when precipitation rates are reduced, and a “Dry Season” from June to September when  
9 precipitation is lowest. Due to gaps in the streamflow record (discussed in section 2.2), the  
10 hydrologic years 1999/2000 through 2002/03 were unavailable for the trend testing for both  
11 the annual and seasonal time periods, and the hydrologic years 1954/55 and 1975/76 were  
12 unavailable for the annual and transitional season.. In addition, the trend tests were not  
13 conducted during the “Dry Period” for streamflow (and therefore also baseflow), due to the  
14 uncertain data quality during these months.

15

## 16 **3 Results**

### 17 **3.1 Summary of the Seasonal Breakdown**

18 To characterize the hydrometeorological conditions of the three seasons’ used in this study;  
19 the median values of the hydrometeorological variables during the study period are presented  
20 in Table 3. This summary shows the strong climatic pattern in the watershed, with distinctly  
21 contrasting precipitation, temperature, and potential evapotranspiration values between  
22 seasons. With respect to streamflow, the values are similar during the wet and transitional  
23 seasons, however both streamflow yield and baseflow index are higher during the transitional  
24 season, which reflects the sustained streamflow carried over from the wet season  
25 precipitation, and the lower proportion of streamflow coming directly from precipitation  
26 events.

### 27 **3.2 Analysis of the Elimination of the Dry Season Streamflow**

28 As discussed in the data section, the months of June to September had to be removed from all  
29 streamflow analyses, due to uncertainty related to unrecorded seasonal impoundments during  
30 this part of the year. To quantify the percentage of streamflow that this excluded from the

1 analysis, an assessment was made over the years when streamflow impoundments did not  
2 occur (45 % of years). During these years, approximately 6.5 % of streamflow occurred  
3 between the months of June to September (Fig. 5, monthly mean values presented).

### 4 **3.3 Assessment of the Baseflow Calculations**

5 The baseflow index used in this study is estimated by the Eckhardt digital filter, which  
6 contains two parameters:  $BFI_{max}$  (maximum value of the baseflow index) and  $a$  (a filter  
7 parameter). These parameters were set at  $BFI_{max} = 0.80$ , and  $a = 0.98$ , based on an  
8 examination of different recommended values provided by Eckhardt (2005). To provide a  
9 check on the baseflow values estimated using this method, the results were then compared  
10 against baseflow values calculated using conductivity data from 2001 to 2009 with the  
11 ‘Conductivity Mass-Balance Method’ (Stewart et al., 2007).

12 At the monthly time-scale, the two compared baseflow data-sets have a Pearson’s correlation  
13 coefficient of 0.96 for all months (Fig. 6a), and 0.83 for months with less than 100 mm of  
14 baseflow (Fig. 6b), which indicates that the Eckhardt method agreed well with the more  
15 empirical Conductivity Mass-Balance Method. This in itself does not confirm the accuracy of  
16 the baseflow values utilized, but it does indicate their consistency over the study period, and  
17 thus their suitability for the time-series trend analysis.

### 18 **3.4 Thiel-Sen / Mann-Kendall Trend Testing Results**

19 The results for the Thiel-Sen/Mann-Kendall trend tests for the variables with the most  
20 noteworthy results (i.e. precipitation, temperature, potential evapotranspiration, streamflow  
21 yield, and baseflow index) are presented by Fig. 7. The full test results for all  
22 hydrometeorological variables and test periods are provided in the supplementary material.

23 For the precipitation data, three significant trends were identified during the transitional  
24 season. All trends corresponded to decreases in precipitation:  $-7.9$  mm/yr trend over the 50  
25 years from 1961 to 2010,  $-11.3$  mm/yr trend over the 35 years from 1976 to 2010, and  $-14.3$   
26 mm/yr trend over the 25-year period from 1976 to 2000. These trends indicate that there was  
27 a pattern of decreasing precipitation totals during the transitional season (February to May)  
28 starting during the P2 land-cover period, and this pattern continued through the E1 and E2  
29 land-cover periods (cf. Table 1).

1 Three significant trends were also found for potential evapotranspiration (PET) during the  
2 transitional season: a  $-0.8$  mm/yr trend over the 50 years from 1936 to 1985, a  $-1.3$  mm/yr.  
3 trend over the 25 years from 1956 to 1980, and a  $1.7$  mm/yr trend over the 25-year period  
4 from 1976 to 2000. Therefore the PET data shows a pattern of negative trends throughout the  
5 P1, P2, and into the E1 land-cover periods, which reverses and becomes positive during the  
6 E1 period and into the E2 land-cover period (cf. Table 1).

7 For the streamflow data record, no significant trends were found for either streamflow  
8 quantity or streamflow yield. No significant trends were found for baseflow quantity either,  
9 however a number of significant trends were found for baseflow index (BFI). For the annual  
10 test period, four significant trends were found in total: including significant positive trends of  
11  $0.16\%/yr$  for the 35 year period from 1936 to 1970 and of  $0.31\%/yr$  for the 25 year period  
12 from 1946 to 1970; and negative trends of  $-0.22\%/yr$  for the 35 year period from 1956 to  
13 1990 and a  $-0.46\%/yr$  trend for the 25 year period from 1966 to 1990. Two significant trends  
14 were found for BFI during the wet season: a  $0.28\%/yr$  trend for the 35 year period from 1936  
15 to 1970 and a  $-0.33\%/yr$  trend for the 25 year period from 1966 to 1990. Therefore, the BFI  
16 data showed an overall pattern of positive trends during the P1 and P2 land-cover periods,  
17 which reverse to negative trends during the P2 period and throughout the E1 land-cover  
18 period (cf. Table 1).

19

## 20 **4 Discussion**

### 21 **4.1 Streamflow Trends**

22 The streamflow trend tests revealed that there were no significant trends for either quantity or  
23 yield over any of the periods tested (Fig. 7). These results therefore contrast with the overall  
24 pattern found in meta-analysis studies dealing with the hydrologic impacts of  
25 afforestation/deforestation, which indicate that afforestation tends to reduce streamflow (e.g.  
26 Bosch and Hewlett, 1982; Brown et al., 2005; Farley et al., 2005). However, there are a  
27 number of individual cases within these meta-analyses studies which show contrasting trends  
28 to the overall pattern. These cases are difficult to directly compare to the current study  
29 however, as most were conducted at the plot to micro-catchment scale, which underwent  
30 relatively rapid land-cover change. By contrast, this study was conducted on a  $404\text{ km}^2$   
31 watershed, which underwent relatively gradual land-cover change over a 75 year period. In

1 this case, any potential changes in hydrologic processes are likely to be far more diffuse and  
2 difficult to detect, when compared to the paired catchment studies.

3 Despite this limitation, some comparisons can be made to sites with similar site conditions, in  
4 terms of having winter-dominant precipitation and shallow soils. Across a number of  
5 catchments with winter-dominant rainfall, Brown et al. (2005) found that afforestation led to  
6 much larger proportional reductions in summer flows compared to winter flows, which they  
7 attributed to the afforestation-induced changes in interception and evapotranspiration. Among  
8 these catchments, those of Gallart et al. (2001) and Lewis et al. (2000) demonstrated the  
9 importance of soil depth in controlling the hydrological response of Mediterranean mountain  
10 catchments in the Pyrenees and California, respectively. Other studies with somewhat similar  
11 site conditions (i.e. Bari et al., 1996; Van Lill et al., 1980) were conducted at very different  
12 temporal and spatial scales than the present study, making comparisons to their findings  
13 difficult. In spite of the lack of comparable studies for direct comparison, the absence of a  
14 marked reduction in streamflow was an unexpected finding, given the scale of afforestation in  
15 the Águeda watershed.

16 A potential explanation for this lack of observed impact could be the presence of offsetting  
17 climatic trends over the same period. Either an increase in water availability due to higher  
18 precipitation (P) and / or a reduction in atmospheric demand due to lower potential  
19 evapotranspiration (PET) could compensate for any land-cover induced changes. While no  
20 significant trends were found for either P or PET at the annual time scale, or during the wet or  
21 dry seasons, significant trends were found during the transitional season, which may have  
22 impacted water availability.

23 With respect to increasing water availability during the transitional season, negative trends in  
24 PET were found from 1936 to 1985 and from 1956 to 1980 (Fig 7). These trends occur  
25 primarily during the periods of pine afforestation (P1, P2) and partially during the transition  
26 to eucalyptus (E1; Cf. Table 1). The trends in PET would lead to a reduction in atmospheric  
27 demand during this period, and therefore could be responsible for offsetting an increase in  
28 consumptive demand that occurred from afforestation.

29 With respect to reductions in water availability during the transitional season, negative trends  
30 in P were found from 1961 to 2010, 1976 to 2010, and 1976 to 2000; and a positive trend in  
31 PET was found from 1976 to 2000 (Fig. 7). These trends indicate movement toward a  
32 relatively more arid environment, which could therefore lead to a reduction in water

1 availability. However, no corresponding trends in streamflow were found during this period.  
2 This lack of change is particularly noteworthy given that these trends occurred during the  
3 eucalyptus afforestation periods (E1, E2; Cf. Table 1), which would also be expected to  
4 increase consumptive demand, and would therefore amplify, rather than offset, an increase in  
5 atmospheric demand.

6 Given the lack of significant climate trends at the annual time scale, and the contrasting  
7 findings during the transitional season, offsetting climatic trends do not appear to be an  
8 adequate explanation for the overall lack of observed streamflow changes in the Águeda  
9 watershed. However, given that the observed climate trends occurred during the transitional  
10 season, there may have been streamflow impacts during the (following) dry season. This can  
11 only be speculated on however, since no assessment can be made on streamflow during the  
12 dry season, due to the limitations in the streamflow data (i.e. the summer streamflow  
13 impoundments). Therefore, no comparison could be made with the findings of Rodríguez-  
14 Suárez et al. (2011), who found dry season reductions in the water table and streamflow  
15 discharge following afforestation with eucalyptus; or to Brown et al. (2005) which found that  
16 afforestation led to much larger proportional reductions in summer flows compared to winter  
17 flows.

18 An alternate explanation for the lack of streamflow change could relate to the specific  
19 characteristics of the watershed, which may make it less responsive to changes in forest land-  
20 cover than is typical. With respect to watershed characteristics, (Andréassian, 2004) identifies  
21 several prerequisites conditions necessary to observe hydrologic impacts, including soil,  
22 climatic, and eco-physiological factors.

23 With respect to soil conditions, the characteristics of the soils of the Águeda watershed may  
24 be a key factor in the lack of a reduction in streamflow. Under conditions of well-developed  
25 soils, the deeper rooting depths of trees will give greater access to soil moisture, allowing for  
26 more transpiration, resulting in higher water consumption. However, the soils of the Águeda  
27 watershed tend to be fairly shallow, being typically less than 1 meter deep and often as  
28 shallow as 20-30 cm (Santos et al., 2013). These depths are less than the maximum rooting  
29 depth of pine and eucalypt trees, and therefore are likely to be a constraint to deep rooting for  
30 both species (Canadell et al., 1996). In addition, the schist and granite bedrock in this  
31 watershed is relatively impermeable and not easily penetrated by tree roots, which restricts the  
32 access of tree species to groundwater reserves as well. Therefore, the capability of the fast-

1 growing pine and eucalypt trees to access deeper sources of soil moisture than the original  
2 shrub and slow-growing tree species is likely much less relevant in this watershed than it  
3 would be in a location with deeper soils. In the case of the Águeda watershed, the most  
4 important soil related factor in water consumption appears to be the low moisture storage  
5 capacity of the soils, severely off-setting the potential impact of widespread planting of trees  
6 with higher water consumptive capacity.

7 A second factor which could contribute to the lack of reductions in streamflow is the  
8 Mediterranean climate regime of the study area. In all Mediterranean-type climates, the period  
9 of peak sunlight and temperature, and therefore potential evapotranspiration, is out of phase  
10 with the maximum precipitation period (Brown et al., 2005). Given the low amount of  
11 summer precipitation, and the shallowness of soils in this watershed, there will typically be  
12 little soil water available for summer evapotranspiration (David et al., 1997; Doerr and  
13 Thomas, 2000). In this regard, the climatic conditions of the Águeda catchment may have an  
14 amplifying effect on the impacts of the shallow soils, by further reducing the higher  
15 evapotranspiration potential of fast-growing trees species.

16 With respect to eco-physiological conditions, the specific land-cover changes in the Águeda  
17 watershed may also be a factor in the lack of an observed reduction in streamflow. One of the  
18 primary drivers of increased consumptive water use by tree species is their typically high  
19 canopy interception capacity (Domingo et al., 1994; Scarascia-Mugnozza et al., 1988;  
20 Tarazona et al., 1996). In the Águeda watershed, however, the interception rates appear to be  
21 comparatively low for pine and eucalypt species (Coelho et al., 2008; Ferreira, 1996; Valente  
22 et al., 1997), while the interception capacity of Mediterranean shrublands can be relatively  
23 high. Garcia-Estringana et al. (2010) found that Mediterranean shrub species can have  
24 interception capacities similar to those of forests. In addition, interception rates are  
25 particularly high in shrublands growing in dense stands (Llorens and Domingo, 2007). These  
26 characteristics apply to the ‘matos’ shrubland that was the most common vegetation type in  
27 the Águeda watershed prior to pine afforestation, as it has a relatively high leaf-area index and  
28 the tendency to grow in very dense stands (Asner et al., 2003). By contrast, given the poor  
29 soil conditions of the study site, the densities of the tree plantations are not as high as they  
30 could be on well-developed soils. Average tree density from unpublished plot assessments put  
31 the density of unevenly spaced eucalyptus stands (< 15 yr old) at 1 600 trees/ha, of evenly  
32 spaces eucalyptus stands on terraces (< 5 years old) at 1,500 trees/ha, of eucalyptus on flat



1 terrain (< 5 yr old) at 2,600 trees/ha, and of unevenly spaced pines (< 30 yr old) at 500  
2 trees/ha. Therefore, the land cover/use change from shrubland to pine/eucalypt forest might  
3 not have resulted in large changes in either transpiration rates or canopy interception rates.

4 Therefore, the Águeda watershed does not meet the prerequisites conditions identified by  
5 Andréassian (2004) for observing afforestation-driven streamflow changes at the watershed  
6 scale. Given this lack of prerequisites conditions, and the absence of offsetting climate trends  
7 as an alternative explanation, the streamflow findings of this study appear to be primarily a  
8 function of watershed characteristics, with soil properties as the most important factor.

## 9 **4.2 Baseflow Trends**

10 No significant trends were found for baseflow quantity (BF) over any of the periods or  
11 seasons tested. However a number of trends were found for baseflow index (BFI), for both the  
12 annual data and the wet season data, which includes both positive and negative trends over  
13 different parts of the data record.

14 Positive trends in BFI were found from 1936 to 1970 for the annual data and the wet season,  
15 and from 1946 to 1970 for the annual data (Fig. 7). These trends correspond with the pine  
16 afforestation land-cover periods P1 and P2 (Cf. Table 1). These trends could be an indication  
17 that the pine afforestation promoted slower flow pathways, by increasing the amount of water  
18 entering the soil matrix via infiltration, and reducing surface flow and fast subsurface flow  
19 (i.e. via macropores). However, given that previous studies in Águeda watershed have found  
20 soil water repellent (SWR) conditions at pine stands (during dry periods), pine afforestation  
21 would not normally be expected to increase matrix infiltration in this location (Keizer et al.,  
22 2005a, 2005b; Santos et al., 2013). However, the land-cover state during the initial conversion  
23 to pine forests were significantly different from the state during these studies, which may  
24 have led to a more positive impact on infiltration rates. This is due to the ground preparation  
25 and planting operations used, which would have the effect of breaking-up the repellent topsoil  
26 layer and creating sinks for overland flow, both of which would promote infiltration. This  
27 effect would be reduced over time, and eventually SWR would recover in established stands.

28 Negative BFI trends were found from 1956 to 1990 for the annual data, and from 1966 to  
29 1990 for the wet season (Fig. 7). This corresponds with the early part of the P2 land-cover  
30 period, and the entirety of the first eucalyptus afforestation period (E1, Cf. Table 1).  
31 Therefore, the negative BFI trends occur during the period when *Pinus pinaster* plantations

1 had reached greater maturity and (after logging) were being rapidly replaced with *Eucalyptus*  
2 *globulus*. The reductions in baseflow during this period may therefore be related to high rates  
3 of soil water repellency (SWR) in the established pine stands and the newly established  
4 eucalypt stands. An increase in SWR could lead to an increase in quick flow, particularly via  
5 fast sub-surface flow from macropore infiltration, and lead to more rapid conversion of  
6 precipitation into streamflow.

7 The temporal correspondence between the significant trends in BFI and land cover changes  
8 which could affect hydrologic flow pathways indicate there may be a relationship between  
9 afforestation and changes in baseflow index in the Águeda watershed. These findings are  
10 further supported by field studies conducted in the watershed, which show the strong impact  
11 of SWR in pine and (particularly) eucalyptus stands on hydrologic flow pathways (Santos et  
12 al., 2013). However, given that there is no field data available to verify the site conditions  
13 during the time of the observed trends, the attribution of the changes in BFI to land-cover  
14 change is necessarily speculative. To test this hypothesis, further field studies would be  
15 needed to examine baseflow dynamics under land-cover conditions which replicate the  
16 historic conditions.

17

## 18 **5 Conclusions**

19 This study did not detect statistically significant – negative or positive – trends in streamflow  
20 quantity or yield in the Águeda watershed of north-central Portugal over the 75 year period  
21 examined, despite the large scale afforestation with *Pinus pinaster* and later *Eucalyptus*  
22 *globulus* which has taken place there. While these findings differ from the general conclusion  
23 of afforestation/deforestation meta-analysis studies, such as Bosch and Hewlett (1982),  
24 Brown et al. (2005), and Farley et al. (2005), they do support the assertion of Andréassian  
25 (2004) that there are requisite climatic, pedological, and eco-physiological watershed  
26 conditions that are necessary to observe hydrologic impacts at the watershed scale. These  
27 conditions are not present in the Águeda watershed, and the lack of soil moisture holding  
28 capacity is likely the primary controlling factor.

29 With respect to baseflow trends, the initial conversion from more natural land-cover types  
30 (i.e. matos shrublands, mixed forests) to pine plantations appears to have had a significant –  
31 initial – positive impact on baseflow index, while the substitution of pine plantations by  
32 eucalypt plantations had a negative impact on baseflow index. The positive trends are

1 attributed to the impact of the site preparation methods applied during the initial pine planting  
2 on soil infiltration capacity, while the negative baseflow trends are attributed to the onset of  
3 soil water repellency (SWR) under the mature pine and eucalypt stands. Therefore, from the  
4 standpoint of promoting well-regulated streamflow (i.e. higher baseflow) the impacts of the  
5 afforestation with pine appear generally positive, while those of re-/afforestation with  
6 eucalypts were generally negative.

7 However, it is important to stress that the pine and eucalypt planting in the study catchment  
8 took place on dissimilar types of land cover. Pines were primarily replacing naturally  
9 occurring shrublands, which was followed by the replacement of the planted pines by  
10 eucalypts. Therefore, a direct comparison between the impacts of widespread planting with  
11 pine or with eucalypt cannot be drawn from this study. In addition, these baseflow findings  
12 are based on a statistical / historical analysis, with no field data available for validation. To  
13 further test this hypothesis, field studies would be needed to examine baseflow dynamics  
14 under different land-cover conditions replicating the historic conditions.

15

## 16 **Acknowledgements**

17 The authors would like to express their gratitude to Professor Celeste Coelho at the University  
18 of Aveiro, Centre for Environmental and Marine Studies (CESAM) for providing her  
19 expertise knowledge on the background and land use history of the study region. Thanks are  
20 also given to Dr. Ameer Manceur and Dr. Robert Schlicht, for providing assistance on the  
21 statistical methods. Funding for this research was provided by the ‘Erasmus Mundus - Forest  
22 and Nature for Society (FONASO) Joint Doctoral Program’ ; the European Regional  
23 Development Fund (through COMPETE), the European Social Fund and the Portuguese  
24 Republic (through FCT), both by individual fellowships (references:  
25 SFRH/BPD/87571/2012; SFRH/BPD/64425/2009; SFRH/BD/61451/2009) and research  
26 projects (references: FCOMP-01-0124-FEDER-009308; FCOMP-01-0124-FEDER-008534;  
27 PTDC/CTE-ATM/111508/2009); and the German Academic Exchange Service (DAAD) for  
28 sponsoring a bilateral collaboration action between Universidade de Aveiro and TU Dresden.

29

## 1 **References**

- 2 Andréassian, V.: Waters and forests: from historical controversy to scientific debate, *J.*  
3 *Hydrol.*, 291(1-2), 1–27, doi:10.1016/j.jhydrol.2003.12.015, 2004.
- 4 Asner, G. P., Scurlock, J. M. O. and A. Hicke, J.: Global synthesis of leaf area index  
5 observations: implications for ecological and remote sensing studies, *Glob. Ecol. Biogeogr.*,  
6 12(3), 191–205, doi:10.1046/j.1466-822X.2003.00026.x, 2003.
- 7 Bari, M. A., Smith, N., Ruprecht, J. K. and Boyd, B. W.: Changes in streamflow components  
8 following logging and regeneration in the southern forest of Western Australia, *Hydrol.*  
9 *Process.*, 10(3), 447–461, doi:10.1002/(SICI)1099-1085(199603)10:3<447::AID-  
10 *HYP431*>3.0.CO;2-1, 1996.
- 11 Benito, E. and Santiago, J. L.: Deforestation of water repellent soils in Galicia (NW Spain):  
12 effects on surface runoff and erosion under simulated rainfall, *Earth Surf. Process. Landf.*,  
13 28(2), 145 – 155, doi:10.1002/esp.431, 2003.
- 14 Benjamini, Y. and Yekutieli, D.: The control of the false discovery rate in multiple testing  
15 under dependency, *Ann. Stat.*, 29, 1165–1188, 2001.
- 16 Bosch, J. M. and Hewlett, J. D.: A review of catchment experiments to determine the effect of  
17 vegetation changes on water yield and evapotranspiration, *J. Hydrol.*, 55(1-4), 3–23,  
18 doi:10.1016/0022-1694(82)90117-2, 1982.
- 19 Brown, A. E., Zhang, L., McMahon, T. A., Western, A. W. and Vertessy, R. A.: A review of  
20 paired catchment studies for determining changes in water yield resulting from alterations in  
21 vegetation, *J. Hydrol.*, 310(1-4), 28–61, doi:10.1016/j.jhydrol.2004.12.010, 2005.
- 22 Bruijnzeel, L. A.: Hydrological functions of tropical forests: not seeing the soil for the trees?,  
23 *Agric. Ecosyst. Environ.*, 104(1), 185–228, doi:10.1016/j.agee.2004.01.015, 2004.
- 24 Calder, I. R.: Water use by forests, limits and controls, *Tree Physiol.*, 18(8-9), 625 –631,  
25 doi:10.1093/treephys/18.8-9.625, 1998.
- 26 Calder, I. R.: Blue revolution: integrated land and water resource management, Earthscan,  
27 London., 2005.
- 28 Canadell, J., Jackson, R. B., Ehleringer, J. B., Mooney, H. A., Sala, O. E. and Schulze, E.-D.:  
29 Maximum rooting depth of vegetation types at the global scale, *Oecologia*, 108(4), 583–595,  
30 doi:10.1007/BF00329030, 1996.
- 31 Coelho, C. O. A., Shakesby, R. A. and Walsh, R. P. D.: Effects of Forest Fires and Post-fire  
32 Land Management Practice on Soil Erosion and Stream Dynamics, Águeda Basin, Portugal,  
33 Soil and groundwater research report V, European Commision, Luxembourg., 1995.
- 34 Coelho, C. O. A., Ferreira, A. J. D., Prats, S. A., Tomé, M., Soares, P., Cortiçada, A., Tomé,  
35 J. A., Salas, G. R., Páscoa, F. and Amaral, A.: Assessment of climatic change impact on water  
36 resources and CO<sub>2</sub> fixation in fast growing stand in Portugal, Final Report Silvaqua Project  
37 POCTI/MGS/49210/2002., 2008.

- 1 Corine Land Cover: Corine Land Cover 2006 raster data, European Environment Agency  
2 (EEA). [online] Available from: [http://www.eea.europa.eu/data-and-](http://www.eea.europa.eu/data-and-maps/data/ds_resolveuid/a645109f7a11d43f5d7e275d81f35c61)  
3 [maps/data/ds\\_resolveuid/a645109f7a11d43f5d7e275d81f35c61](http://www.eea.europa.eu/data-and-maps/data/ds_resolveuid/a645109f7a11d43f5d7e275d81f35c61), 2010.
- 4 David, T. S., Ferreira, M. I., David, J. S. and Pereira, J. S.: Transpiration from a mature  
5 *Eucalyptus globulus* plantation in Portugal during a spring-summer period of progressively  
6 higher water deficit, *Oecologia*, 110(2), 153–159, doi:10.1007/PL00008812, 1997.
- 7 Doerr, S. H. and Thomas, A. D.: The role of soil moisture in controlling water repellency:  
8 new evidence from forest soils in Portugal, *J. Hydrol.*, 231–232, 134–147,  
9 doi:10.1016/S0022-1694(00)00190-6, 2000.
- 10 Doerr, S. H., Shakesby, R. A. and Walsh, R. P. D.: Soil hydrophobicity variations with depth  
11 and particle size fraction in burned and unburned *Eucalyptus globulus* and *Pinus pinaster*  
12 forest terrain in the Águeda Basin, Portugal, *CATENA*, 27(1), 25–47, doi:10.1016/0341-  
13 8162(96)00007-0, 1996.
- 14 Domingo, F., Puigdefabregas, J., Moro, M. J. and Bellot, J.: Role of vegetation cover in the  
15 biogeochemical balances of a small afforested catchment in southeastern Spain, *J. Hydrol.*,  
16 159(1–4), 275–289, doi:10.1016/0022-1694(94)90261-5, 1994.
- 17 Eckhardt, K.: How to construct recursive digital filters for baseflow separation, *Hydrol.*  
18 *Process.*, 19(2), 507–515, doi:10.1002/hyp.5675, 2005.
- 19 Estêvão, J. A.: A florestação dos baldios, *Análise Soc.*, XIX: 77-79, 1983.
- 20 Farley, K. A., Jobbágy, E. G. and Jackson, R. B.: Effects of afforestation on water yield: a  
21 global synthesis with implications for policy, *Glob. Change Biol.*, 11(10), 1565–1576,  
22 doi:10.1111/j.1365-2486.2005.01011.x, 2005.
- 23 Ferreira, A. J. D., Coelho, C. O. A., Walsh, R. P. D., Shakesby, R. A., Ceballos, A. and Doerr,  
24 S. H.: Hydrological implications of soil water-repellency in *Eucalyptus globulus* forests,  
25 north-central Portugal, *J. Hydrol.*, 231–232, 165–177, doi:10.1016/S0022-1694(00)00192-X,  
26 2000.
- 27 Ferreira, C. G.: Erosão hídrica em solos florestais: estudo em povoamentos de *Pinus Pinaster*  
28 e *Eucalyptus Globulus* em macieira de Alcôba-Águeda, *Rev. Fac. Let. – Geogr. Sér.*, vol  
29 XII/XIII, pp. 145–244, 1996.
- 30 Ferreira, R. V., Cerqueira, M. A., de Melo, M. T. C., de Figueiredo, D. R. and Keizer, J. J.:  
31 Spatial patterns of surface water quality in the Cértima River basin, central Portugal, *J.*  
32 *Environ. Monit. JEM*, 12(1), 189–199, doi:10.1039/b914409a, 2010.
- 33 Gallart, F., Llorens, P., Latron, J. and Regüés, D.: Hydrological processes and their seasonal  
34 controls in a small Mediterranean mountain catchment in the Pyrenees, *Hydrol Earth Syst Sci*,  
35 6(3), 527–537, doi:10.5194/hess-6-527-2002, 2001.
- 36 Garcia-Estringana, P., Alonso-Blázquez, N. and Alegre, J.: Water storage capacity, stemflow  
37 and water funneling in Mediterranean shrubs, *J. Hydrol.*, 389(3–4), 363–372,  
38 doi:10.1016/j.jhydrol.2010.06.017, 2010.

- 1 Geri, F., Amici, V. and Rocchini, D.: Human activity impact on the heterogeneity of a  
2 Mediterranean landscape, *Appl. Geogr.*, 30(3), 370–379, doi:10.1016/j.apgeog.2009.10.006,  
3 2010.
- 4 Giorgi, F. and Lionello, P.: Climate change projections for the Mediterranean region, *Glob.*  
5 *Planet. Change*, 63(2–3), 90–104, doi:10.1016/j.gloplacha.2007.09.005, 2008.
- 6 Harris, I., Jones, P. d., Osborn, T. j. and Lister, D. h.: Updated high-resolution grids of  
7 monthly climatic observations – the CRU TS3.10 Dataset, *Int. J. Climatol.*, 34(3), 623–642,  
8 doi:10.1002/joc.3711, 2014.
- 9 Helsel, R. M. H. D. R.: *Statistical Methods in Water Resources*, *Technometrics*, 36(3),  
10 doi:10.1080/00401706.1994.10485818, 1993.
- 11 Instituto da Conservação da Natureza e das Florestas: Mapas de áreas ardidadas entre 1990-  
12 1996, 1997-2004, 2010, 2011, 2012 e 2013, [online] Available from:  
13 <http://www.icnf.pt/portal/florestas/dfci/inc/mapas> (Accessed 1 July 2014), 2014.
- 14 Iovino, F., Cinnirella, S., Veltri, A. and Callegari, G.: Processus hydriques dans des  
15 écosystèmes forestiers, *Ecologie*, 29, 369–375, 1998.
- 16 IPMA: Instituto Portugues do Mar e Atmosfera, IPMA. [online] Available from:  
17 <http://snirh.pt>, 2014.
- 18 Jones, J. A., Creed, I. F., Hatcher, K. L., Warren, R. J., Adams, M. B., Benson, M. H., Boose,  
19 E., Brown, W. A., Campbell, J. L., Covich, A., Clow, D. W., Dahm, C. N., Elder, K., Ford, C.  
20 R., Grimm, N. B., Henshaw, D. L., Larson, K. L., Miles, E. S., Miles, K. M., Sebestyen, S. D.,  
21 Spargo, A. T., Stone, A. B., Vose, J. M. and Williams, M. W.: Ecosystem Processes and  
22 Human Influences Regulate Streamflow Response to Climate Change at Long-Term  
23 Ecological Research Sites, , doi:10.1525/bio.2012.62.4.10, 2012.
- 24 Jones, N., de Graaff, J., Rodrigo, I. and Duarte, F.: Historical review of land use changes in  
25 Portugal (before and after EU integration in 1986) and their implications for land degradation  
26 and conservation, with a focus on Centro and Alentejo regions, *Appl. Geogr.*, 31(3), 1036–  
27 1048, doi:10.1016/j.apgeog.2011.01.024, 2011.
- 28 Keizer, J. J., Coelho, C. O. A., Matias, M. J. S., Domingues, C. S. P. and Ferreira, A. J. D.:  
29 Soil water repellency under dry and wet antecedent weather conditions for selected land-cover  
30 types in the coastal zone of central Portugal, *Soil Res.*, 43(3), 297–308, 2005a.
- 31 Keizer, J. J., Coelho, C. O. A., Shakesby, R. A., Domingues, C. S. P., Malvar, M. C., Perez, I.  
32 M. B., Matias, M. J. S. and Ferreira, A. J. D.: The role of soil water repellency in overland  
33 flow generation in pine and eucalypt forest stands in coastal Portugal, *Soil Res.*, 43(3), 337–  
34 349, 2005b.
- 35 Lewis, D., Singer, M. J., Dahlgren, R. A. and Tate, K. W.: Hydrology in a California oak  
36 woodland watershed: a 17-year study, *J. Hydrol.*, 240(1–2), 106–117, doi:10.1016/S0022-  
37 1694(00)00337-1, 2000.
- 38 Van Lill, W. S., Kruger, F. J. and Van Wyk, D. B.: The effect of afforestation with  
39 *Eucalyptus grandis* Hill ex Maiden and *Pinus patula* Schlecht. et Cham. on streamflow from

- 1 experimental catchments at Mokobulaan, Transvaal, *J. Hydrol.*, 48(1–2), 107–118,  
2 doi:10.1016/0022-1694(80)90069-4, 1980.
- 3 Llorens, P. and Domingo, F.: Rainfall partitioning by vegetation under Mediterranean  
4 conditions. A review of studies in Europe, *J. Hydrol.*, 335(1–2), 37–54,  
5 doi:10.1016/j.jhydrol.2006.10.032, 2007.
- 6 Malvar, M. C., Prats, S. A., Nunes, J. P. and Keizer, J. J.: Post-fire overland flow generation  
7 and inter-rill erosion under simulated rainfall in two eucalypt stands in north-central Portugal,  
8 *Environ. Res.*, 111(2), 222–236, doi:10.1016/j.envres.2010.09.003, 2011.
- 9 Moreira, F., Rego, F. C. and Ferreira, P. G.: Temporal (1958-1995) pattern of change in a  
10 cultural landscape of northwestern Portugal: implications for fire occurrence, *Landsc. Ecol.*,  
11 16(6), 557–567, 2001.
- 12 Prats, S. A., MacDonald, L. H., Monteiro, M., Ferreira, A. J. D., Coelho, C. O. A. and Keizer,  
13 J. J.: Effectiveness of forest residue mulching in reducing post-fire runoff and erosion in a  
14 pine and a eucalypt plantation in north-central Portugal, *Geoderma*, 191, 115–124,  
15 doi:10.1016/j.geoderma.2012.02.009, 2012.
- 16 Rego, F. C.: *Florestas Públicas*, Ministério da Agricultura, Desenvolvimento Rural e Pescas,  
17 Lisbon, Portugal., 2001.
- 18 Rodríguez-Suárez, J. A., Soto, B., Perez, R. and Diaz-Fierros, F.: Influence of Eucalyptus  
19 globulus plantation growth on water table levels and low flows in a small catchment, *J.*  
20 *Hydrol.*, 396(3–4), 321–326, doi:10.1016/j.jhydrol.2010.11.027, 2011.
- 21 Santos, J. M., Verheijen, F. G. A., Tavares Wahren, F., Wahren, A., Feger, K.-H., Bernard-  
22 Jannin, L., Rial-Rivas, M. E., Keizer, J. J. and Nunes, J. P.: Soil Water Repellency Dynamics  
23 in Pine and Eucalypt Plantations in Portugal – a High-Resolution Time Series, *Land Degrad.*  
24 *Dev.*, n/a–n/a, doi:10.1002/ldr.2251, 2013.
- 25 Scarascia-Mugnozza, G., Valentini, R. and Spinelli, R.E.G.: Osservazioni sul ciclo dell’acqua  
26 in un bosco ceduo di *Quercus cerris* L., *Ann. Dell’Accademia Ital. Sci. For.*, XXXVII, 3–21,  
27 1988.
- 28 Scott, D. F. and Lesch, W.: Streamflow responses to afforestation with *Eucalyptus grandis*  
29 and *Pinus patula* and to felling in the Mokobulaan experimental catchments, South Africa, *J.*  
30 *Hydrol.*, 199(3–4), 360–377, doi:10.1016/S0022-1694(96)03336-7, 1997.
- 31 Sen, P. K.: Estimates of the Regression Coefficient Based on Kendall’s Tau, *J. Am. Stat.*  
32 *Assoc.*, 63(324), 1379–1389, doi:10.1080/01621459.1968.10480934, 1968.
- 33 Serra, P., Pons, X. and Saurí, D.: Land-cover and land-use change in a Mediterranean  
34 landscape: A spatial analysis of driving forces integrating biophysical and human factors,  
35 *Appl. Geogr.*, 28(3), 189–209, doi:10.1016/j.apgeog.2008.02.001, 2008.
- 36 Shakesby, R., Coelho, C., Ferreira, A., Terry, J. and Walsh, R.: Wildfire Impacts on Soil-  
37 Erosion and Hydrology in Wet Mediterranean Forest, Portugal, *Int. J. Wildland Fire*, 3(2),  
38 95–110, 1993.

- 1 Shakesby, R. A., Doerr, S. H. and Walsh, R. P. D.: The erosional impact of soil  
 2 hydrophobicity: current problems and future research directions, *J. Hydrol.*, 231–232, 178–  
 3 191, doi:10.1016/S0022-1694(00)00193-1, 2000.
- 4 Silva, G., Antunes, A., Escada, A. and Marques, J.: *Relações Agricultura/Floresta Ambiente*,  
 5 Gabinete de Planeamento e Política Agro-Alimentar (GPPAA), Ministério da Agricultura, do  
 6 Desenvolvimento Rural e das Pescas, Lisbon, Portugal., 2004.
- 7 SNIRH: Sistema Nacional de Informação de Recursos Hídricos, Agência Portuguesa do  
 8 Ambiente, Lisbon. [online] Available from: <http://snirh.pt>, 2013.
- 9 Stewart, M., Cimino, J. and Ross, M.: Calibration of base flow separation methods with  
 10 streamflow conductivity, *Ground Water*, 45(1), 17–27, doi:10.1111/j.1745-  
 11 6584.2006.00263.x, 2007.
- 12 Tarazona, T., Santa Regina, I. and Calvo, R.: Interception, throughfall and stemflow in two  
 13 forest of the “Sierra de la Demanda” in the province of Burgos [Spain], *Pirin. Espana* [online]  
 14 Available from: [http://agris.fao.org/agris-](http://agris.fao.org/agris-search/search.do?f=1998%2FES%2FES98017.xml%3BES1998000049)  
 15 [search/search.do?f=1998%2FES%2FES98017.xml%3BES1998000049](http://agris.fao.org/agris-search/search.do?f=1998%2FES%2FES98017.xml%3BES1998000049) (Accessed 4 May  
 16 2014), 1996.
- 17 Thornthwaite, C. W.: An Approach toward a Rational Classification of Climate, *Geogr. Rev.*,  
 18 38(1), 55–94, doi:10.2307/210739, 1948.
- 19 Valente, F., David, J. S. and Gash, J. H. C.: Modelling interception loss for two sparse  
 20 eucalypt and pine forests in central Portugal using reformulated Rutter and Gash analytical  
 21 models, *J. Hydrol.*, 190(1–2), 141–162, doi:10.1016/S0022-1694(96)03066-1, 1997.
- 22 Wahren, A., Schwärzel, K. and Feger, K.-H.: Potentials and limitations of natural flood  
 23 retention by forested land in headwater catchments: evidence from experimental and model  
 24 studies, *J. Flood Risk Manag.*, 5(4), 321–335, doi:10.1111/j.1753-318X.2012.01152.x, 2012.
- 25 Yue, S., Pilon, P., Phinney, B. and Cavadias, G.: The influence of autocorrelation on the  
 26 ability to detect trend in hydrological series, *Hydrol. Process.*, 16(9), 1807–1829,  
 27 doi:10.1002/hyp.1095, 2002.

28

## 29 **Tables**

30 Table 1. Land-cover periods and dominant afforestation trends in Águeda watershed from  
 31 1935 to 2010.

<i>Land-Cover Period</i>	<i>Time Period</i>	<i>Dominant Afforestation Trend</i>
<b>P1</b>	<b>1935 - 1950</b>	Large scale replacement of shrubland with <i>Pinus pinaster</i> .
<b>P2</b>	<b>1950 - 1970</b>	Continuing afforestation with <i>Pinus pinaster</i> , but at a slower rate.



<b>E1</b>	<b>1970 - 1990</b>	Rapid reforestation with <i>Eucalyptus globulus</i> (particularly post '86 wildfire), replacement of <i>Pinus pinaster</i> .
<b>E2</b>	<b>1990 - 2010</b>	Relatively stable forested area, with continued replacement of <i>Pinus pinaster</i> with <i>Eucalyptus globulus</i> .

1

2 Table 2. Summary of hydrometeorological variables.

<b>Hydrometeorological Variables</b>			
<b>Variable</b>	<b>Description</b>	<b>Data Source</b>	<b>Unit</b>
P	Precipitation	SNIRH Gauge Data	mm
T	Temperature	IPMA Gauge Data	°C
PET	Potential Evapotranspiration	Thornthwaite Equation	mm
Q	Streamflow Quantity	SNIRH Gauge Data	mm
Q <sub>yld</sub>	Streamflow Yield	$\sum Q_{mm} / \sum P$	%
BF	Baseflow Quantity	Recursive Digital Filter	mm
BFI	Baseflow Index	$\sum BF_{mm} / \sum Q_{mm}$	%

3

4


1 Table 3. Seasonal and annual median values of the hydrometeorological variables in Águeda  
 2 watershed from 1936 - 2010.

Median Values: 1936 - 2010								
Season	Months	P (mm)	T (°C)	PET (mm)	Q (mm)	Q <sub>yl</sub> (%)	BF (mm)	BFI (%)
Wet	Oct - Jan	965	11.7	145	301	30 %	149	55 %
Transitional	Feb - May	626	12.6	198	281	43 %	184	63 %
Dry	Jun - Sep	193	19.3	390	NA	NA	NA	NA
Annual	All*	1 787	14.7	732	565	36 %	320	59 %

3 \* The months of June to September are not included for Q (mm), Q<sub>yl</sub> (%), BF (mm), and BFI (%).


4  
5  
6  
7

### Figures



8  
9 Figure 1. Location and Land-Cover of the Águeda watershed.


10



1

2 Figure 2. Average monthly precipitation and temperature in the Águeda watershed from 1971  
 3 to 2000.

4




5

6 Figure 3. Burned area in the Águeda watershed from 1975 to 2010; total watershed area is  
 7 404 km<sup>2</sup>.

8


Timeline	1935	1940	1945	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000	2005
Afforestation Period	P1			P2			E1			E2					
75 yr Trend Test	1936 to 2010														
50 yr Trend Tests	1936 to 1985														
						1961 to 2010									
35 yr Trend Tests	1936 to 1970														
						1956 to 1990									
						1976 to 2010									
25 yr Trend Tests	1936 to 1960														
				1946 to 1970											
					1956 to 1980										
							1966 to 1990								
									1976 to 2000						
													1986 to 2010		

1  
2 Figure 4. Timeline of the trend-testing periods and their correspondence with the different  
3 afforestation periods.



5  
6 Figure 5. Monthly mean streamflow during the years without seasonal impoundments; the  
7 boxed off period (June - September) indicates the period removed from the streamflow and  
8 baseflow analysis.

9



1

2 Figure 6. Monthly plots of baseflow from the Conductivity Mass-Balance (CMB) and  
 3 Eckhardt digital filter calculations; 5a includes all months ( $r^2 = 0.96$ ) and 5b includes months  
 4 with less than 100 mm of baseflow ( $r^2 = 0.83$ ).

5

6

1

2 Figure 7. Summary of the trend testing results, with the afforestation periods (P1, P2, E1, E2: cf. Table 1) overlain for comparison. Significant  
3 trends are indicated with dashes lines.

