

A Baseline Probabilistic Drought Forecasting Framework Using Standardized Soil Moisture Index: Application to the 2012 United States Drought

A. AghaKouchak¹

[1] University of California, Irvine

Correspondence to: A. AghaKouchak (amir.a@uci.edu)

Abstract

The 2012 drought was one of the most extensive drought events in half a century, resulting in over \$12 billion in economic loss in the United States, and substantial indirect impacts on global food security and commodity prices. An important feature of the 2012 drought was rapid development and intensification in late spring/early summer, a critical time for crop development and investment planning. Drought prediction remains a major challenge because dynamical precipitation forecasts are highly uncertain, and their prediction skill is low. Using a probabilistic framework for drought forecasting based on the persistence property of accumulated soil moisture, this paper shows that the U.S. drought of summer 2012 was predictable several months in advance. The presented drought forecasting framework provides the probability occurrence of drought based on climatology and near-past observations of soil moisture. Our results indicate that soil moisture exhibits higher persistence than precipitation, and hence improves drought predictability.

1 Introduction

According to United States Department of Agriculture (USDA) estimates, about 80 percent of U.S. agricultural land experienced drought in 2012 which made the event more extensive than any since 1950 (USDA, 2012). A striking aspect of the 2012 drought was rapid increase in

1 severity in early July during a critical time of crop development (USDA, 2012). The quick
2 onset of the drought in the central plains during late spring led to a so-called “flash drought”
3 (Hoerling et al., 2013). A drought early warning system with seasonal predictions of drought
4 onset, severity, persistence, and spatial extent in a timely manner would provide invaluable
5 information to decision-makers and stakeholders. There are a number of research and
6 operational drought (or hydrologic) prediction systems (Pozzi et al., 2013; Mishra and Singh,
7 2010), including the Climate Prediction Center Seasonal Drought Outlook (Steinemann,
8 2006), the University of Washington’s Surface Water Monitor (Wood and Lettenmaier, 2006;
9 Wood, 2008), Princeton University’s drought forecast system (Luo and Wood, 2007; Li et al.,
10 2008; Sheffield et al., 2008), U.S. - Mexico Drought Prediction Tool (Lyon et al., 2012), and
11 the Global Integrated Drought Monitoring and Prediction System (GIDMaPS; Hao et al.,
12 2014). Despite all these efforts, a community White Paper by the World Climate Research
13 Program identified sub-seasonal to seasonal drought prediction as one of the major research
14 gaps in hydroclimatology (WCRP, 2010).

15 Drought forecasting is generally based on drought indicators computed using dynamic or
16 statistical model simulations of drought-related variables (e.g., Mishra et al., 2009; Madadgar,
17 and Moradkhani, 2013). Droughts are classified as agricultural (soil moisture deficit),
18 meteorological (precipitation deficit), and hydrological (streamflow/groundwater deficit), and
19 various drought indicators based on soil moisture, precipitation and runoff have been
20 developed to describe different aspects of droughts (Heim, 2002; Wood et al., 2002; Wood
21 and Lettenmaier, 2006; Mo, 2008; Shukla and Lettenmaier, 2011; Hao and AghaKouchak,
22 2013). Most drought prediction studies are based on the Standardized Precipitation Index
23 (SPI; McKee et al., 1993) with the input precipitation derived from dynamical
24 weather/climate models (Yoon et al., 2012; Mwangi et al., 2013; Dutra et al., 2013, 2014a,
25 2014b). While dynamic models provide valuable information, precipitation forecasts are
26 subject to high uncertainty and models exhibit very low skill in predicting precipitation with a
27 few months lead time (Goddard et al., 2003; National Research Council, 2006; Livezey and
28 Timofeyeva, 2008; Lavers et al., 2009). A baseline probability method is proposed for
29 meteorological drought forecasting based on persistence of the SPI (Lyon et al., 2012),
30 indicating that a statistical persistence-based model could lead to a good seasonal drought
31 forecasting skill (Quan et al., 2012). Hao et al., 2014 developed a multivariate method for
32 statistical drought prediction using a persistence-based approach.

1 Soil moisture is often used as an indicator of agricultural drought monitoring, and has been
2 used in different forms (Samaniego et al., 2013) including the soil moisture percentile (Luo
3 and Wood, 2007; Wood, 2008; Shukla et al., 2011), normalized soil moisture (Dutra et al.,
4 2008), and soil moisture anomaly (Sheffield and Wood, 2007; Sheffield and Wood, 2008).
5 Typically, precipitation and temperature forecasts, either from dynamic models or
6 climatology resampling (i.e., Ensemble Streamflow Prediction, ESP method; Mo et al., 2012),
7 are used to force land-surface/hydrologic models for predicting soil moisture conditions and
8 drought (e.g., Luo and Wood, 2007; Luo and Wood, 2008; Trambauer et al., 2013). The
9 uncertainty of dynamic soil moisture forecasts is even higher than the climate forcings
10 (precipitation and temperature) because in addition to input uncertainty, model errors and
11 uncertainty also propagate into soil moisture simulations. For this reason, different statistical
12 methods such as conditional ESP resampling have been explored for soil moisture prediction
13 (Wood, 2008).

14 Persistence is a distinctive characteristic of the soil moisture as it exhibits less variability
15 relative to precipitation (Hao and AghaKouchak, 2014). Mo et al., (2012) emphasized the
16 importance of the persistence of soil moisture in improving drought forecasting skill. Great
17 strides have been made to explore soil moisture persistence properties, and results reveal that
18 the persistence of soil moisture memory spans weeks to a couple of months (Vinnikov and
19 Yeserkepova, 1991; Entin et al., 2000; Seneviratne et al., 2006; Koster et al., 2010). Though
20 the persistence property of soil moisture has been well documented, the properties of
21 accumulated soil moisture and its potential use for drought forecasting has less been
22 investigated. In this study, a probabilistic drought prediction framework is proposed using the
23 Standardized Soil moisture Index (SSI) as the drought indicator, which allows for the
24 description of soil moisture across different time scales (e.g., 3-, 6-, 12 month). In other
25 words, soil moisture is treated in a similar fashion to precipitation accumulation across
26 different time scales relative to the corresponding long-term climatology (McKee et al.,
27 1993). Given the temporal integration of data, SSI leads to even higher persistence compared
28 with the commonly used soil moisture percentiles or soil moisture anomaly.

29

30 **2 Data**

31 The data sets used in this study include the monthly precipitation and soil moisture from the
32 NASA Modern-Era Retrospective analysis for Research and Applications (MERRA-Land),

1 available on a $2/3^\circ \times 1/2^\circ$ grid from January 1, 1980 onwards (Reichle et al., 2011; Rienecker
2 et al., 2011). MERRA data sets have been used in numerous studies in different climatic
3 regions (Bosilovich et al., 2011; Golian et al., 2014; Wong et al., 2011). Uncertainties in
4 MERRA data sets have been evaluated against different observations (e.g., Yi et al., 2011;
5 Kennedy et al., 2011). The results show that MERRA provides valuable information
6 consistent with observations especially in middle-latitudes, while uncertainties in high
7 latitudes are often large (Yi et al., 2011; Reichle et al., 2011).

8

9 **3 Methodology**

10 The Standardized Soil Moisture Index (SSI; Hao and AghaKouchak, 2014) can be defined in
11 a similar way to the commonly used Standardized Precipitation Index (SPI; Mckee et al.,
12 1993) that has been used in a wide variety of studies (Dutra et al., 2013; Damberg and
13 AghaKouchak, 2013). Here, the SSI is estimated using a nonparametric approach in which the
14 empirical probability (p) of the historical soil moisture data is derived using the empirical
15 Gringorten plotting position (Gringorten, 1963). In other words, instead of fitting a
16 distribution function to soil moisture data, the probabilities (p) are obtained empirically using
17 the empirical Gringorten approach: $(i-0.44)/(n+0.12)$ where n denotes the sample size and i
18 refers to the rank of soil moisture data from from the smallest.

19 The empirical probabilities, derived from the Gringorten plotting position, are then
20 transformed into the standard normal distribution function: $SSI = \Phi^{-1}(p)$, where Φ is the
21 standard normal distribution function. In this approach, one can avoid making a decision
22 about the parametric distribution function of accumulated soil moisture at different time
23 scales. Assume that soil moisture for the month i is S_i . Then the 6-month accumulation of the
24 soil moisture A_i for the month i can be expressed as:

$$25 \quad A_i = S_{i-5} + S_{i-4} + S_{i-3} + S_{i-2} + S_{i-1} + S_i, \quad (1)$$

26 In this study, the Ensemble Streamflow Prediction (ESP) method (Twedt et al., 1977; Day,
27 1985) is used for resampling from historical records of soil moisture to obtain monthly
28 moisture at the target season with the 6-month SSI as the drought indicator. Assume the l -
29 month lead forecasting is needed based on the monthly soil moisture observations with
30 forecast initialization at month i . Then the l month ($1 \leq l \leq 5$) ahead forecasting of the
31 accumulated soil moisture A_{i+l} can be expressed:

$$A_{i+l} = S_{i+l-5} + S_{i+l-4} + S_{i+l-3} + S_{i+l-2} + S_{i+l-1} + S_{i+l} \quad (2)$$

Assume that one month lead forecasting (i.e., $l=1$) based on the 6-month SSI is needed. The unknown S_{i+1} is predicted by resampling the soil moisture from the historical record of the target month (i.e., $i+1$). As a result, an ensemble of m (i.e., the length of observation in the historical record) sequence of the monthly soil moisture in the target season can be obtained from the observed monthly soil moisture. In this manner, m sequences of accumulated 6-month soil moisture for the l month lead time can be generated by blending the observed and predicted monthly soil moisture. For example, for $l=1$, the blended sequences of accumulated 6-month soil moisture can be expressed as:

10

$$A_{i+1}^{(1)} = S_{i-4} + S_{i-3} + S_{i-2} + S_{i-1} + S_i + S_{i+1}^{(1)}$$

$$A_{i+1}^{(2)} = S_{i-4} + S_{i-3} + S_{i-2} + S_{i-1} + S_i + S_{i+1}^{(2)} \quad (3)$$

13

$$A_{i+1}^{(m)} = S_{i-4} + S_{i-3} + S_{i-2} + S_{i-1} + S_i + S_{i+1}^{(m)}$$

where S_{i-4}, \dots, S_i are the observed soil moisture prior to the target month in the 6-month window, while $S_{i+1}^{(1)}, \dots, S_{i+1}^{(m)}$ are the sequences of sampled monthly soil moisture from the observations in the historical record for the target month (here, S_{i+1}). Note for the first 5 months of the year, some of the observed soil moisture prior to the target month (S_{i-4}, \dots, S_i) in the past 6-month will be sampled from one year prior to the target year.

Each sequence of the blended 6-month soil moisture $A^{(j)}$, $j=1, 2, \dots, m$, in equation (3) can be combined with the observed 6-month accumulated soil moisture in the past years to derive the corresponding SSI^(j). Here, the probability of drought is defined as the probability that a future drought condition (SSI) is lower than an alarm threshold (e.g., SSI < -0.8 corresponding to ~20th percentile). The empirical probability is estimated by dividing the number of the forecasted values below the threshold (e.g., -0.8) by the number of the ensemble members.

26

1 4 Results

2 First it is shown that the accumulated soil moisture typically exhibits much higher persistence
3 compared to precipitation, and hence can be used for drought forecasting with up to several
4 months lead time. Then, the 2012 summer drought conditions are predicted using the SSI with
5 different lead times. The SSI is obtained using predicted soil moisture information using the
6 ESP concept based on long-term climatology and near-past observations (see Section
7 Methodology). The study focuses on the drought prediction for May to August which is an
8 important period for agricultural decision-making.

9 Understanding the persistence property of soil moisture is fundamental to drought forecasting.
10 It is hypothesized that using accumulated soil moisture would improve persistence-based
11 drought forecasting relative to using accumulated precipitation. First, the persistence property
12 of accumulated soil moisture is evaluated against the accumulated precipitation that has been
13 used for meteorological drought prediction (Lyon et al., 2012; Quan et al., 2012; Hao et al.,
14 2014; Yoon et al., 2012). The monthly precipitation and soil moisture data from MERRA-
15 Land (Reichle et al., 2011; Rienecker et al., 2011) in California and Texas are used to
16 examine the persistence of accumulated soil moisture relative to precipitation. Both states are
17 among the most important producers of agricultural products, and have experienced
18 severe/extreme drought conditions in the past decade. The autocorrelations of accumulated 6-
19 month precipitation and soil moisture for 1- to 6-month time lags and four different initial
20 conditions (March, April, May and June) for summer drought prediction are provided in
21 Figure 1. In the figure, the term *initial* is defined similar to initial condition in Section
22 Methodology. For example, March corresponds to precipitation and soil moisture from Oct.
23 2011 through March 2012. The boxplots present the median, 25th, 75th percentiles, and
24 whiskers of the autocorrelations. Lyon et al., 2012 showed that variance of the accumulated
25 precipitation can enhance or diminish the persistence of the SPI at different start times,
26 mainly due to seasonality of precipitation. As shown, the autocorrelation of the accumulated
27 soil moisture (or SSI) is generally higher than that of accumulated precipitation (or SPI) for
28 the four different initial conditions. The figure shows that the autocorrelations of the
29 accumulated 6-month soil moisture decay at a slower rate than the accumulated 6-month
30 precipitation in both California (Figure 1a) and Texas (Figure 1b). For example, in California
31 and for the initial condition in April, the medians of the autocorrelation coefficients are higher
32 than 0.6 even at a 5-month lag. However, the medians of the autocorrelations of the 6-month

1 SPI drop below 0.6 after a 4-month lag. The higher persistence of the SSI relative to SPI
2 implies that a persistence-based model based on SSI would lead to better predications as
3 compared to a similar model based on SPI (see also Changnon 1987).

4 The 6-month SSI is used as the drought indicator to monitor and predict the 2012 (May-
5 August) U.S. drought. Figure 2a shows observed drought conditions from May to August
6 2012. As shown, the drought develops and intensifies quickly, affecting most of the
7 continental U.S. including the Great Plains, the Midwest, and west and southeast. By August,
8 a large portion of the country experienced severe, extreme, or exceptional drought conditions.
9 In operational drought early warning, the severe drought condition is of critical concern. In
10 this paper, the proposed methodology is tested for predicting the moderate and severe drought
11 conditions in summer 2012. Following the U.S. Drought Monitor (USDM), D-scale, the
12 moderate drought is defined as SSI below -0.8 (corresponding to nonexceedance probability
13 of ~ 0.2), whereas the severe drought is defined as SSI below -1.3 (or nonexceedance
14 probability of ~ 0.1) (Svoboda et al., 2002). The observed drought conditions below the
15 severe level (D2) for May-August are shown in Figure 2b.

16 The 1-month and 2-month lead drought ($SSI < -0.8$) forecasts for May-August 2012 are
17 presented in Figures 3a and 3b, respectively. The 1-month lead forecasted SSI maps for
18 different initializations resemble the observed SSI well in terms of the spatial extent (compare
19 Figure 3a with Figure 2a). As shown, the regions with high probability of drought (e.g., above
20 approx. 90%) are in very good agreement with the observations. For example, the outlined
21 methodology predicts high probability of drought over the western U.S. and high plains in
22 August, which is consistent with observations. Furthermore, as the 2012 drought intensifies,
23 the area with high probability of drought (Figure 3a) increases in a similar manner to the
24 observations (Figure 2a). A visual comparison of the two month lead drought forecasts
25 (Figure 3b) and observations (Figure 2a) reveal that the predicted drought conditions are in
26 very good agreement with probabilities higher than 0.8 in most regions. The 1-month and 2-
27 month lead severe drought ($SSI < -1.3$) forecasts for May-August 2012 are presented in Figure
28 4a and Figure 4b. The 1-month lead forecasts of May-August severe drought conditions are in
29 very good agreement with observations. As shown, the severe drought condition from May-
30 August in northern Texas, and the western U.S. are captured in the predictions. Figure 4b
31 highlights that even at a 2-month lead, the proposed model predicts the 2012 summer drought
32 reasonably well.

1 The predicted drought probability maps for July and August 2012 for 3-month and 4-month
2 lead time are presented in Figures 5a (SSI<-0.8) and 5b (SSI<-1.3). One can see that the 3-
3 month and 4-month lead forecasts capture the observed drought conditions with probabilities
4 ranging from 0.1 to 0.8. The prediction skill of the model is higher in the western U.S. where
5 drought conditions are predicted at higher probabilities relative to the Midwest. A review of
6 Figures 3 and 4 indicate that the predicted probabilities in longer leads (i.e., 3- and 4-month)
7 are typically lower than those of shorter (1- and 2-month) lead forecasts. Basically, in
8 persistence-based models, as the lead month increases, one expects the forecast probabilities
9 to decrease as well. This can be partly explained from the autocorrelations of accumulated soil
10 moisture presented in Figure 1. As shown, in the western U.S., the 4-month lead forecasted
11 drought probabilities for July and August 2012 are relatively high and in fairly good
12 agreement with observations. In the Midwest and eastern U.S., the proposed model indicates
13 relatively low probabilities of drought for 3- and 4-month lead forecasts. While the forecasted
14 drought probabilities are lower at a 4 month lead, still they provide valuable information by
15 showing the drought signal. While the 3- and 4- month lead forecasted probabilities of severe
16 droughts are substantially less compared to the 2-month lead forecasts, the drought signal in
17 the western U.S. is still strong (see Figure 5b).

18 It should be noted that the seasonal climate predictions based on weather/climate models
19 initialized in April and May 2012 revealed limited drought information for May-July and
20 June-August 2012 (Hoerling et al., 2013). This highlights that improvements in just two 2-
21 month lead forecast could be very important for risk assessment and decision making. The
22 presented persistence-based model with the SSI as the drought indicator provides potential
23 capability to predict droughts that would of great value to agricultural planning.

24 The quality and the latency of predictions rely on the quality and availability of input data
25 sets. Currently, limited observations of soil moisture are available across the globe, and soil
26 moisture estimation relies on model simulations. The Soil Moisture Ocean Salinity (SMOS)
27 and the upcoming Soil Moisture Active and Passive (SMAP) mission may provide the
28 opportunity to integrate near real-time satellite data with long-term climate data records such
29 as MERRA to improve drought monitoring and prediction.

30

1 **5 Conclusions**

2 Using the Standardized Soil moisture Index (SSI) as the drought indicator, a persistence-
3 based drought prediction method is presented and used for predicting the 2012 United States
4 drought. It is shown that because of high persistence property of soil moisture, the SSI can be
5 used for seasonal drought forecasting. The presented statistical approach predicted the May –
6 August drought conditions relatively well, especially for 1- and 2-month lead forecasts. The
7 3- and 4-month lead forecasts of the western U.S. were in good agreement with observations.
8 However, the drought prediction signal in the eastern U.S. was not as strong at 3- and 4-
9 month lead time. Given the persistence-based nature of the methodology, uncertainties of
10 predictions increase with lead time. Similar behaviour has been observed in persistence-based
11 drought recovery assessment (Pan et al., 2013). However, even 1- and 2-month lead
12 information is valuable to some end-users including farmers and commodity investors.

13 It is acknowledged that, similar to other methods, both the presented modelling framework
14 and input data sets are subject to uncertainties. The presented model is based on near past soil
15 moisture conditions and log-term climatology. Soil moisture responds to precipitation with
16 some delay, and for this reason, the methodology may not capture rapid developments.
17 Furthermore, this methodology relies on historical observations and because of limited
18 samples of extreme conditions in historical records, it should not be used for predicting
19 extreme droughts.

20 It is stressed that the proposed approach is not meant to replace the currently available
21 dynamic drought forecasting models. Rather, the persistence-based predictions should be used
22 as additional information that can potentially improve drought predictability. Finally, it
23 should be pointed out that SSI is not suggested as an alternative to use of SPI (or other
24 indicators) for seasonal drought prediction. The best choice of index or the best set of
25 indicators depends on the problem in hand and the climate of the study area. It is our view
26 that drought monitoring and prediction should be based on multiple sources of information
27 (data and indicator) as well as models (e.g., dynamic, statistical).

28

29 **Acknowledgements**

30 The financial support for this study is provided by the National Science Foundation Award
31 No. EAR-1316536 and the Hellman Foundation.

32

1 **References**

- 2 AghaKouchak A., and Nakhjiri N., 2012, A Near Real-Time Satellite-Based Global Drought
3 Climate Data Record, *Environmental Research Letters*, 7(4), 044037, doi:10.1088/1748-
4 9326/7/4/044037.
- 5 Changnon, S. A. (1987). Detecting drought conditions in Illinois (No. 163-170). Illinois State
6 Water Survey.
- 7 Day, G. N. (1985). Extended streamflow forecasting using NWSRFS. *Journal of Water*
8 *Resources Planning and Management*. 111(2): 157-170.
- 9 Dutra, E., P. Viterbo and P. M. Miranda (2008). ERA-40 reanalysis hydrological applications
10 in the characterization of regional drought. *Geophysical Research Letters*. 35(19):
11 L19402, doi: 10.1029/2008GL035381.
- 12 Dutra, E., Wetterhall, F., Di Giuseppe, F., Naumann, G., Barbosa, P., Vogt, J., ... &
13 Pappenberger, F. (2014a). Global meteorological drought–Part 1: Probabilistic
14 monitoring. *Hydrology and Earth System Sciences Discussions*, 11(1), 889-917.
- 15 Dutra, E., Pozzi, W., Wetterhall, F., Di Giuseppe, F., Magnusson, L., Naumann, G., ... &
16 Pappenberger, F. (2014b). Global meteorological drought–Part 2: Seasonal forecasts.
17 *Hydrology and Earth System Sciences Discussions*, 11(1), 919-944.
- 18 Dutra, E., Di Giuseppe, F., Wetterhall, F., and Pappenberger, F. (2013): Seasonal forecasts of
19 droughts in African basins using the Standardized Precipitation Index, *Hydrol. Earth*
20 *Syst. Sci.*, 17, 2359-2373, doi:10.5194/hess-17-2359-2013.
- 21 Damberg L., AghaKouchak A., 2013, Global Trends and Patterns of Droughts from Space,
22 *Theoretical and Applied Climatology*, doi: 10.1007/s00704-013-1019-5.
- 23 Entin, J. K., Robock, A., Vinnikov, K. Y., Hollinger, S.E., Liu S., Namkhai, A. (2000).
24 Temporal and spatial scales of observed soil moisture variations in the extratropics.
25 *Journal of Geophysical Research*. 105(D9): 11865-11811,11877.
- 26 Goddard, L., A. Barnston and S. Mason (2003). Evaluation of the IRI's “Net assessment”
27 seasonal climate forecasts: 1997-2001. *Bulletin of the American Meteorological Society*.
28 84(12): 1761-1781.

- 1 Gringorten, I. I. (1963). A Plotting Rule for Extreme Probability Paper. *Journal of*
2 *Geophysical Research*. 68(3): 813-814.
- 3 Hao Z., AghaKouchak A., Nakhjiri N., Farahmand A., (2014), Global Integrated Drought
4 Monitoring and Prediction System, *Scientific Data*, 1:140001, 1-10, doi:
5 10.1038/sdata.2014.1.
- 6 Hao, Zengchao, Amir AghaKouchak, 2014: A Nonparametric Multivariate Multi-Index
7 Drought Monitoring Framework. *J. Hydrometeor*, 15, 89–101. doi:
8 <http://dx.doi.org/10.1175/JHM-D-12-0160.1>.
- 9 Hao, Z. and A. AghaKouchak (2013). Multivariate Standardized Drought Index: A multi-
10 Index parametric approach for drought analysis. *Advances in Water Resources*, Volume
11 57: Pages 12–18.
- 12 Heim, R. R. (2002). A review of twentieth-century drought indices used in the United States.
13 *Bulletin of the American Meteorological Society*. 83(8): 1149-1166.
- 14 Hoerling, M., S. Schubert, K. Mo, et al. (2013). An Interpretation of the Origins of the 2012
15 Central Great Plains Drought, Assessment Report, NOAA Drought Task Force.
- 16 Kennedy, A. D., X. Dong, B. Xi, et al. (2011). A comparison of MERRA and NARR
17 reanalyses with the DOE ARM SGP data. *Journal of climate*. 24(17): 4541-4557.
- 18 Koster, R. D., Mahanama, S. P. P., Yamada, T. J., Balsamo, G., Berg, A. A., Boisserie, M.,
19 Dirmeyer, P. A., Doblas-Reyes, F. J., Drewitt, G., Gordon, C. T., Guo, Z., Jeong, J.-H.
20 Lawrence, D. M., Lee, W.-S., Li, Z., Luo, L., Malyshev, S., Merryfield, W. J.,
21 Seneviratne, S. I., Stanelle, T., van den Hurk, B. J. J. M., Vitart, F., E. F. Wood (2010).
22 Contribution of land surface initialization to subseasonal forecast skill: First results from
23 a multi-model experiment. *Geophysical Research Letters*. 37(2): L02402.
- 24 Lavers, D., L. Luo and E. F. Wood (2009). A multiple model assessment of seasonal climate
25 forecast skill for applications. *Geophysical Research Letters*. 36(23),
26 10.1029/2009GL041365.
- 27 Li, H., L. Luo and E. F. Wood (2008). Seasonal hydrologic predictions of low-flow
28 conditions over eastern USA during the 2007 drought. *Atmospheric Science Letters*. 9(2):
29 61-66.

- 1 Livezey, R. E. and M. M. Timofeyeva (2008). The first decade of long-lead US seasonal
2 forecasts: Insights from a skill analysis. *Bulletin of the American Meteorological Society*.
3 89(6): 843-854.
- 4 Luo, L. and E. F. Wood (2007). Monitoring and predicting the 2007 US drought. *Geophysical*
5 *Research Letters*. 34(22): L22702, doi: 10.1029/2007GL031673.
- 6 Luo, L. and E. F. Wood (2008). Use of Bayesian merging techniques in a multimodel
7 seasonal hydrologic ensemble prediction system for the eastern United States. *Journal of*
8 *Hydrometeorology*. 9(5): 866-884.
- 9 Lyon, B., Bell, M.A., Tippett M.K., Kumar, A., Hoerling, M.P., Quan, X.-W, Wang, H.
10 (2012). Baseline probabilities for the seasonal prediction of meteorological drought.
11 *Journal of Applied Meteorology and Climatology*. 51(7): 1222-1237.
- 12 Madadgar, Shahrbanou, Hamid Moradkhani, 2013: A Bayesian Framework for Probabilistic
13 Seasonal Drought Forecasting. *J. Hydrometeor*, 14, 1685–1705. doi:
14 <http://dx.doi.org/10.1175/JHM-D-13-010.1>.
- 15 McKee, T. B., N. J. Doesken and J. Kleist (1993). The relationship of drought frequency and
16 duration to time scales. Eighth Conference on Applied Climatology, Am. Meteorol. Soc.,
17 Anaheim, CA.
- 18 Mishra, A. K., & Singh, V. P. (2010). A review of drought concepts. *Journal of Hydrology*,
19 391(1), 202-216.
- 20 Mishra, A. K., Singh, V. P., & Desai, V. R. (2009). Drought characterization: a probabilistic
21 approach. *Stochastic Environmental Research and Risk Assessment*, 23(1), 41-55.
- 22 Mwangi, E., Wetterhall, F., Dutra, E., Giuseppe, F. D., & Pappenberger, F. (2013).
23 Forecasting droughts in East Africa. *Hydrology and Earth System Sciences Discussions*,
24 10(8), 10209-10230.
- 25 Mo, K. C. (2008). Model-Based Drought Indices over the United States. *Journal of*
26 *Hydrometeorology*. 9(6): 1212-1230.
- 27 Mo, K.C., Shukla, S., Lettenmaier, D.P., Chen, L.-C. (2012). Do Climate Forecast System
28 (CFSv2) forecasts improve seasonal soil moisture prediction? *Geophysical Research*
29 *Letters*. 39(23), doi: 10.1029/2012GL053598.

- 1 National Research Council (2006). Completing the Forecast: Characterizing and
2 Communicating Uncertainty for Better Decisions Using Weather and Climate Forecasts,
3 The National Academies Press.
- 4 Pan, M., Yuan, X., & Wood, E. F. (2013). A probabilistic framework for assessing drought
5 recovery. *Geophysical Research Letters*, 40(14), 3637-3642.
- 6 Pozzi, W.; Sheffield, J.; Stefanski, R.; Cripe, D.; Pulwarty, R.; Vogt, J. V.; Heim Jr., R. R.;
7 Brewer, M. J.; Svoboda, M.; Westerhoff, R.; Van Dijk, A. I. J. M.; Lloyd-Hughes, B.;
8 Pappenberger, F.; Werner, M.; Dutra, E.; Wetterhall, F.; Wagner, W.; Schubert, S.; Mo,
9 K.; Nicholson, M. (2013). Toward global drought early warning capability. *Bulletin of*
10 *the American Meteorological Society*, 94(6).
- 11 Quan, X.-W., Hoerling, M. P., Lyon, B., Kumar, A., Bell, M. A., Tippett, M. K., Wang, H.
12 (2012). Prospects for dynamical prediction of meteorological drought. *Journal of Applied*
13 *Meteorology and Climatology*. 51(7): 1238-1252.
- 14 Reichle, R.H., Koster, R.D., De Lannoy, G. J. M., Forman, B.A., Liu, Q., Mahanama, S. P.
15 P., Touré, A. (2011). Assessment and enhancement of MERRA land surface hydrology
16 estimates. *Journal of climate*. 24(24): 6322-6338.
- 17 Rienecker, Michele M.; Suarez, Max J.; Gelaro, Ronald; Todling, Ricardo; Bacmeister, Julio;
18 Liu, Emily; Bosilovich, Michael G.; Schubert, Siegfried D.; Takacs, Lawrence; Kim, Gi-
19 Kong; Bloom, Stephen; Chen, Junye; Collins, Douglas; Conaty, Austin; da Silva,
20 Arlindo; Gu, Wei; Joiner, Joanna; Koster, Randal D.; Lucchesi, Robert; Molod, Andrea;
21 Owens, Tommy; Pawson, Steven; Pegion, Philip; Redder, Christopher R.; Reichle, Rolf;
22 Robertson, Franklin R.; Ruddick, Albert G.; Sienkiewicz, Meta; Woollen, Jack (2011).
23 MERRA: NASA's modern-era retrospective analysis for research and applications.
24 *Journal of climate*. 24(14): 3624-3648.
- 25 Samaniego, L., Kumar, R., & Zink, M. (2013). Implications of parameter uncertainty on soil
26 moisture drought analysis in Germany. *Journal of Hydrometeorology*, 14(1), 47-68.
- 27 Seneviratne SI, Koster RD, Guo Z, Dirmeyer PA, Kowalczyk E, Lawrence D, Liu P, Lu C-H,
28 Mocko D, Oleson KW, Verseghy D, (2006). Soil moisture memory in AGCM
29 simulations: analysis of global land-atmosphere coupling experiment (GLACE) data.
30 *Journal of Hydrometeorology*. 7(5): 1090-1112.

1 Sheffield, J. and E. F. Wood (2007). Characteristics of global and regional drought, 1950–
2 2000: Analysis of soil moisture data from off-line simulation of the terrestrial hydrologic
3 cycle. *Journal of Geophysical Research*. 112(D17): D17115, doi: doi:
4 10.1029/2006JD008288.

5 Sheffield, J. and E. F. Wood (2008). Global trends and variability in soil moisture and drought
6 characteristics, 1950-2000, from observation-driven simulations of the terrestrial
7 hydrologic cycle. *Journal of climate*. 21(3): 432-458.

8 Sheffield, J., E. Wood, D. Lettenmaier, A. Lipponen, 2008, Experimental drought monitoring
9 for Africa, *GEWEX News* 18 (3), 4-6.

10 Shukla, S. and D. Lettenmaier (2011). Seasonal hydrologic prediction in the United States:
11 understanding the role of initial hydrologic conditions and seasonal climate forecast skill.
12 *Hydrology and Earth System Sciences*. 15(11): 3529.

13 Shukla, S., A. C. Steinemann and D. P. Lettenmaier (2011). Drought Monitoring for
14 Washington State: Indicators and Applications. *Journal of Hydrometeorology*. 12(1): 66-
15 83.

16 Steinemann, A. C. (2006). Using climate forecasts for drought management. *Journal of*
17 *Applied Meteorology and Climatology*. 45(10): 1353-1361.

18 Svoboda, Mark; Lecomte, Doug; Hayes, Mike; Heim, Richard; Gleason, Karin; Angel, Jim;
19 Rippey, Brad; Tinker, Rich; Palecki, Mike; Stooksbury, David; Miskus, David; Stephens,
20 Scott (2002). The drought monitor. *Bulletin of the American Meteorological Society*. 83:
21 1181-1190.

22 Trambauer, P., Maskey, S., Winsemius, H., Werner, M., & Uhlenbrook, S. (2013). A review
23 of continental scale hydrological models and their suitability for drought forecasting in
24 (sub-Saharan) Africa. *Physics and Chemistry of the Earth, Parts A/B/C*, 66, 16-26.

25 Twedt, T. M., Schaake Jr., J. C., and Peck, E. L., 1977: National Weather Service extended
26 streamflow prediction, in: *Proc. 45th Western Snow Conference*, Albuquerque, NM,
27 Colorado State University, 52–57.

28 USDA (2012). U.S. Drought 2012: Farm and Food Impacts.
29 [http://www.ers.usda.gov/topics/in-the-news/us-drought-2012-farm-and-food-](http://www.ers.usda.gov/topics/in-the-news/us-drought-2012-farm-and-food-impacts.aspx)
30 [impacts.aspx](http://www.ers.usda.gov/topics/in-the-news/us-drought-2012-farm-and-food-impacts.aspx).

1 Vinnikov, K. Y. and I. Yeserkepova (1991). Soil moisture: Empirical data and model results.
2 Journal of climate. 4(1): 66-79.

3 WCRP (2010). WCRP White Paper on ‘Drought Predictability and Prediction in a Changing
4 Climate: Assessing Current Capabilities, User Requirements, and Research
5 Priorities,Tech. rep., World Climate Research Programme. , Barcelona, Spain.

6 Wood, A. W. (2008). The University of Washington Surface Water Monitor: An experimental
7 platform for national hydrologic assessment and prediction. American Meteorology
8 Society annual meeting, 22nd conference on hydrology, New Orleans.

9 Wood, A. W. and D. P. Lettenmaier (2006). A test bed for new seasonal hydrologic
10 forecasting approaches in the western United States. Bulletin of the American
11 Meteorological Society. 87(12): 1699-1712.

12 Wood, Andrew W.; Maurer, Edwin P.; Kumar, Arun; Lettenmaier, Dennis P. (2002). Long-
13 range experimental hydrologic forecasting for the eastern United States. J. Geophys. Res.
14 107(D20): 4429, doi: 10.1029/2001JD000659.


15 Yi, Y., J. S. Kimball, L. A. Jones, et al. (2011). Evaluation of MERRA land surface estimates
16 in preparation for the Soil Moisture Active Passive Mission. Journal of climate. 24(15):
17 3797-3816.

18 Yoon, J. H., K. Mo and E. F. Wood (2012). Dynamic-Model-Based Seasonal Prediction of
19 Meteorological Drought over the Contiguous United States. Journal of
20 Hydrometeorology. 13(2): 463-482.

21 Wong, S., Fetzer, E. J., Kahn, B. H., Tian, B., Lambriksen, B. H., & Ye, H. (2011). Closing
22 the Global Water Vapor Budget with AIRS Water Vapor, MERRA Reanalysis, TRMM
23 and GPCP Precipitation, and GSSTF Surface Evaporation. Journal of Climate, 24(24).

24
25

1




2

3 Figure 1 Boxplots of autocorrelation coefficients (up to 6 month) of accumulated 6-month
4 precipitation (blue) and soil moisture (red) from MERRA-Land for different initial month for
5 (a) California and (b) Texas. The boxplots show the median (center), 25th (lower) and 75th
6 (upper) percentiles edges.

7

8

1




2

3 Figure 2 (a) Observed 6-month SSI for May-August 2012; (b) Observed 6-month SSI with
4 severe drought condition ($SSI < -1.3$) for May-August 2012.

5

1



2


3 Figure 3 (a) One month and (b) two months lead drought probability predictions for May-

4 August 2012 for $SSI6 < -0.8$.

5

1


2



3 Figure 4 (a) One month and (b) two months lead drought probability predictions for May-
4 August 2012 for $SSI6 < -1.3$.

5

1



2 Figure 5 Three and four month lead time predictions of drought probability for July-August

3 2012; (a) SSI6 < -0.8; (b) SSI6 < -1.3

4

5