

1           **Quantitative contribution of climate change and human**  
2           **activities to runoff changes in the Wei River basin, China**

3  
4           C.S. Zhan<sup>2</sup>, S.S. Jiang<sup>3\*</sup>, F.B. Sun<sup>2</sup>, Y.W. Jia<sup>1</sup>, C.W. Niu<sup>1</sup>, W.F. Yue<sup>3</sup>

5  
6           1. State Key Laboratory of Simulation and Regulation of Water Cycle in River Basin,  
7           China Institute of Water Resources and Hydropower Research, Beijing 100038, China

8           2. Key Laboratory of Water Cycle Relate Land Surface Processes, Institute of  
9           Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences,  
10           Beijing, 100101, China;

11           3. College of Water Sciences, Beijing Normal University, 100875, Beijing, China;

12  
13           \* Corresponding author: Shanshan Jiang, 19 Xijiekou Wai, Haidian, Beijing ,  
14           100875, China

15           Tel: +86(10)6488-8975

16           Fax: +86(10)6488-8975

17           Email: [jiangshanshan@mail.bnu.edu.cn](mailto:jiangshanshan@mail.bnu.edu.cn)

21 **Abstract**

22 Surface runoff from the Wei River basin, the largest tributary of the Yellow River in  
23 China, has dramatically decreased over last 51 years from 1958 to 2008. Climate  
24 change and human activities have been identified as the two main reasons for the  
25 decrease in runoff. The study period is split into two sub-periods (1958-1989 and  
26 1990-2008) using the Mann-Kendall jump test. This study develops an improved  
27 climate elasticity method based on the original climate elasticity method, and  
28 conducts a quantitative assessment of the impact of climate change and human  
29 activities on the runoff decrease in the Wei River basin. The results from the original  
30 climate elasticity method show that climatic impacts contribute 37%~40% to the  
31 decrease in runoff, while human impacts contribute 60%~63%. In contrast, the results  
32 from the improved climate elasticity method yield a climatic contribution to runoff  
33 decrease of 22%~29% and a human contribution of 71%~78%. A discussion of the  
34 simulation reliability and uncertainty concludes that the improved climate elasticity  
35 method has better mechanism and can provide more reasonable results.

36

37

38

39

40

## 41 **1. Introduction**

42 Climate change is expected to extensively alter global hydrological cycles (Legesse et  
43 al., 2003; Milly, 2005; Piao et al., 2007) by primarily changing the pattern of  
44 precipitation (IPCC, 2013; Sun et al., 2012). After precipitation falling into a basin,  
45 human activities including land use change, dam construction, river diversion, and  
46 other engineering and management practices will modify hydrological cycles locally  
47 and therefore temporal and spatial distribution of water resources (Govinda, 1995;  
48 Milly et al., 2005). Quantitatively assessing the influence of climate change and  
49 human activities on surface runoff is vital for sustainable water resources  
50 management.

51 Quantitative evaluation of the effects of climate change and human activities on  
52 runoff has yielded significant results, but with complex regional patterns. For instance,  
53 Zhang et al. (2008) used the sensitivity of runoff to precipitation and potential  
54 evaporation to study the response of runoff to changes in climate and land use/cover  
55 in the Loess Plateau of China and pointed out that LUCC( Land Use and Land Cover  
56 Change ) accounted for over 50% of the reduction in mean annual runoff in 8 out of  
57 11 catchments. Bao et al. (2012) discussed the reasons for runoff changes in Haihe  
58 River Basin, and analyzed the influence of human activities through the VIC model  
59 and then proved human activities were the main driving force for the reduction of  
60 water resources. Wang et al. (2009) established a distributed monthly water balance  
61 model (DTVGM) to analyze the Chaobai River Basin upstream of Miyun Reservoir in  
62 north China, and concluded that human activities were therefore the main cause of

63 runoff changes. Ma et al. (2008) estimated that the effects of climate change  
64 accounted for over 64% of the mean annual runoff reduction in the Shiyang River in  
65 the arid region of Northwest China. For the same basin, human activities and climate  
66 change may have different influences on runoff for different periods. For example,  
67 Qiu et al. (2012) analyzed the influence of climate change and human activities on  
68 water resources in North China and found that in the 1970s-80s the effects of climate  
69 change were dominant, but in the late 1980s and early 1990s the effects of the two  
70 factors were similar, and since the 90s the influence of human activities has been  
71 slightly higher at around 55%. Guo et al. (2008) employed the SWAT model to  
72 analyze the annual and seasonal runoff variability caused by climate change and  
73 human activities and found that the main influencing factor on annual runoff was  
74 climate change, but that changing land use was the main influence on seasonal runoff  
75 changes.

76       So far, there are many different methods used to evaluate and separate the effects  
77 of the two factors. One such method is based on physical processes or physical  
78 mechanisms. This method controls the evolution of the various elements and analyzes  
79 the changes in driving factors and contributions in a physical process simulation. The  
80 approach may be physically sound but requires major efforts on model calibration and  
81 can lead to remarkably different results because of uncertainty in model structure and  
82 parameter estimation (Nash and Gleick, 1991; Revelle and Waggoner, 1983; Schaake,  
83 1990; Vogel et al., 1999a). Other methods can be classified as statistical data analysis  
84 methods, such as the climate elasticity method used in this paper. The statistical data

85 analysis methods are based on mean annual change trend in long time series, and  
86 provide generalized relationships which do not consider species differences. This  
87 approach incorporates measured or observed data via a variety of data validation  
88 techniques, and analyzes the contributions of different factors on different processes.  
89 The method is relatively simple, but requires a large volume of high-quality data  
90 (Risbey and Entekhabi, 1996).

91 Many studies have proved that the climate elasticity method is reasonable and  
92 credible. Chiew (2006) evaluated rainfall elasticity of streamflow in 219 catchments  
93 across Australia using the nonparametric climate elasticity estimator and compared  
94 the estimates with results obtained from the conceptual rainfall-runoff model  
95 SIMHYD, showing a consistent relationship between climate elasticity values  
96 estimated using the rainfall-runoff model and the nonparametric estimator. Ma et al.  
97 (2010) used a distributed hydrological model (GBHM) and a climate elasticity model  
98 to conduct a quantitative assessment of the impacts of climate change and human  
99 activities on inflow into a reservoir. The GBHM simulation and climate elasticity  
100 model showed that climate change accounts for about 55% and 51% of the reservoir  
101 inflow reduction, respectively. Hu et al. (2012) analyzed the impacts of climate  
102 change and human activities on the Baiyangdian upstream runoff, using two  
103 assessment methodologies (climate elasticity and hydrological modeling). The climate  
104 elasticity method was implemented at the annual scale and was computationally  
105 relatively simple; it needed fewer data and parameters to calculate the impacts of  
106 climate change on annual runoff. The hydrological model was implemented at the

107 daily scale, and therefore needed more data and parameters but yielded more detailed,  
108 high temporal resolution results. These two independent methods based on different  
109 time scales could obtain consistent results. Thus, the climate elasticity method is  
110 considered to be an important indicator for quantifying the sensitivity of runoff to  
111 climate change and for separating the effects of natural and anthropogenic factors at a  
112 catchment scale (Dooge et al., 1999; Fu et al., 2007; Milly and Dunne, 2002;  
113 Sankarasubramanian et al., 2001; Schaake, 1990; Zheng et al., 2009). However, the  
114 main poor point for the climate elasticity method previous research indicates is that  
115 the method is used to separate the sensitivity of runoff to climate change without  
116 considering the human activities directly, and furthermore the accuracy of the original  
117 climate elasticity method should be improved.

118 In this paper we improve the climate elasticity approach by adding the influence  
119 of human activities to evaluate the hydrological consequences of climate change and  
120 human activities. To compare the original and improved climate elasticity approaches,  
121 we choose the Wei River basin as a case study. The application results have a great  
122 strategic meaning in the regional economic development and the development of West  
123 China (Song et al., 2007). The rest of the paper is structured as follows: Section 2  
124 describes the study area and data; Section 3 presents the methodology; Results and  
125 discussion are in Section 4. The conclusions are presented in section 5.

## 126 **2. Study Area and Data**

### 127 **2.1 Study Area**

128 The Wei River is the largest tributary in the Yellow River. It originates from Niaoshu  
129 Mountain, and runs into the Yellow River at Tongguan. The basin is located between  
130 104 °00'E to 110 °20'E and 33 °50'N to 37 °18'N with a length of 818 km and a drainage  
131 area of  $1.35 \times 10^5 \text{ km}^2$ .

132 The Wei River basin is located in temperate continental monsoon climate region.  
133 The climate is cold, dry and rainless in winter controlled by the Mongolia high, while  
134 hot and rainy in summer affected by the West Pacific subtropical high. The mean air  
135 temperature is 7.8 to 13.5 °C, the mean annual precipitation is 400 to 800 mm , , and  
136 the mean annual potential evapotranspiration is 800 to1000 mm. The mean annual  
137 runoff in depth is 450 to 550 mm with the coefficient of variation falling within 0.1 to  
138 0.2 (He et al., 2009).

### 139 **2.2 Data Description**

140 This study uses the continuous daily series data from 1958 to 2008 at 7 national  
141 meteorological observatory stations in and around the Wei River basin. Observed  
142 daily mean air temperature, precipitation and solar shortwave radiation could largely  
143 reflect climatic change in the region (Fig. 1). In addition, observational runoff data of  
144 1958-2008 comes from Hua Country hydrological station which located in the outlet  
145 of river downstream.

146 **Figure 1** Location of the meteorological and hydrological stations used in this study

147 The potential evapotranspiration within the watershed can be calculated using  
148 Hargreaves method, Hargreaves formula is an empirical formula which was derived  
149 using the permeameter by Hargreaves and Samani (1982) to estimate potential  
150 evapotranspiration, and is most appropriate in all radiation and temperature estimation  
151 methods in North China (Luo and Rong, 2007). The expression is as follows:

$$152 \quad E_0 = 0.0135(T + 17.8) \frac{R_s}{\lambda} \quad (1)$$

153 Where  $E_0$  represents potential evapotranspiration, mm/day;  $T$  represents mean air  
154 temperature, °C;  $R_s$  represents solar shortwave radiation, MJ/(m<sup>2</sup> day);  $\lambda$  represents  
155 latency for vapouring water,  $\lambda = 2.45$  MJ/kg.

### 156 **3. Methodology**

#### 157 **3.1 Detection of hydrologic changes**

##### 158 **3.1.1 Trend analysis**

159 In this study, the Mann-Kendall trend test is used to test the long-term trends of  
160 precipitation, evaporation and runoff in the Wei River basin. The Mann-Kendall trend  
161 test is a non-parametric statistical test method. It does not need the sample to follow  
162 any particular distribution, and is not subject to interference from a small number of  
163 outliers. Moreover, the method is relatively simple (Mann H B, 1945; Kendall M G,  
164 1975).

165 For a time series  $X$  which has  $n$  samples, construct variable  $S$ :

166 
$$S = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \text{sgn}(x_i - x_j) \quad (2)$$

167 Where  $x_i$  and  $x_j$  represent the values in years  $i$  and  $j$ ,  $i > j$ ,  $n$  is the record length of  
 168 the series, and  $\text{Sgn}(x_i - x_j)$  is a characterization of the function.

169 The statistical test value  $Z_c$  is calculated by the following formula:

170 
$$Z_c = \begin{cases} \frac{S - 1}{\sqrt{\text{Var}(S)}}, & S > 0 \\ 0, & S = 0 \\ \frac{S + 1}{\sqrt{\text{Var}(S)}}, & S < 0 \end{cases} \quad (3)$$

171 When  $|Z_c| \leq Z_{1-\alpha/2}$ , we accept the null hypothesis, which indicates the sequence  
 172 does not have a trend. If  $|Z_c| > Z_{1-\alpha/2}$ , we reject the null hypothesis and conclude that  
 173 the sequence does have a significant trend.  $Z_{1-\alpha/2}$  is obtained from standard normal  
 174 distribution function, and  $\alpha$  is the significance level of the test.

175 To test the trend of the sequence in the Mann-Kendall test, it is usually necessary  
 176 to estimate the slope of the monotonic trend, estimated as follows:

177 
$$\beta = \text{M e d i a} \left( \frac{x_i - x_j}{i - j} \right), (\forall j < i, 1 \leq j < i \leq n) \quad (4)$$

178 A positive value of  $\beta$  indicates a rising trend (positive rate of change with time),  
 179 and vice versa for negative  $\beta$ .

### 180 **3.1.2 Change-point analysis**

181 Identifying the change-point for runoff substantially influenced by human activities is  
 182 significant. Depending upon the change-point, the hydrology series could be divided  
 183 into two period, before change-point, it was regarded as no human activities effects  
 184 and after change-point it was believed that human activities sharply affected runoff.

185 Under the hypothesis that the time series is independent and stochastic, we can  
 186 compute the following statistics:

$$187 \quad UF_k = \frac{S_k - E(S_k)}{\sqrt{\text{Var}(S_k)}} \quad (k = 1, 2, \dots, n) \quad (5)$$

188 Where  $E(S_k)$  and  $\text{Var}(S_k)$  represent the mean and variance of  $S_k$ , respectively.

189 Next, the time series order is reversed (i.e.,  $x_n, x_{n-1}, \dots, x_1$ ), and the above process  
 190 is repeated to yield the statistical variables  $UB_k (k=n, n-1, \dots, 1)$ , such that

$$191 \quad UB_k = -UF_k \quad (6)$$

192 Next, the curves of  $UB_k$  and  $UF_k$  are drawn, and if the two curves have an  
 193 intersection point and if the value of  $U$  at this point satisfies  $|U| < 1.96$ , then that point  
 194 is regarded as a change point, with a confidence level  $\alpha$  of 0.05.

195 Because the time series length in this study is 51 years, we use a significance test  
 196 of t-test.

### 197 **3.2 Original climate elasticity method**

198 Runoff ( $R$ ) can be expressed as a function of climate variables ( $C$ ) and other  
 199 characteristics ( $H$ ) (Hu et al., 2012):

$$200 \quad R = f(C, H) \quad (7)$$

201 While the parameter  $H$  represents the combined results of terrain, soil, land use /  
 202 land cover and human activities (such as artificial water transfer). If the topography  
 203 and soil in the study area remain constant during the study period, then  $H$  can  
 204 represent human activities. So the runoff change can be expressed as:

$$205 \quad \Delta R = \Delta R_C + \Delta R_H \quad (8)$$

206 While the parameter  $\Delta R$  represents the total runoff change; and  $\Delta R_C$ ,  $\Delta R_H$   
 207 represent the runoff changes caused by climate change and human activities,  
 208 respectively.

209 The total runoff change can be obtained from the formula  $\Delta R = R_{obs1} - R_{obs2}$ , where  
 210  $R_{obs1}$  and  $R_{obs2}$  represent the measured runoff before and after the change point,  
 211 respectively.

212 Schaake(1990) first introduced the climate elasticity method to analyze the  
 213 sensitivity of runoff to climate change. Climate elasticity of runoff ( $\varepsilon_X$ ) can be  
 214 defined as the proportional change in runoff ( $R$ ) relative to the change in climatic  
 215 variables ( $X$ ) (such as changes in precipitation or potential evapotranspiration) (Fu et  
 216 al., 2007):

$$217 \quad \varepsilon_X = \frac{\partial R / R}{\partial X / X} \quad (9)$$

218 According to the long-term water balance equation ( $R = P - E$ ), we assume that  
 219 the runoff response to climate factors is mainly caused by the precipitation and  
 220 potential evapotranspiration. According to the theory of total differential equations,  
 221 the differential form is as follows:

$$222 \quad \Delta R_C = \varepsilon_P \frac{R}{P} \Delta P + \varepsilon_{E_0} \frac{R}{E_0} \Delta E_0 \quad \text{and} \quad \varepsilon_P + \varepsilon_{E_0} = 1 \quad (10)$$

223 Where  $\Delta R_c$  represents runoff change caused by climate change,  $\Delta P$  and  $\Delta E_0$  are  
 224 the change of precipitation and potential evapotranspiration, and  $\varepsilon_P$  and  $\varepsilon_{E_0}$  are the  
 225 precipitation and potential evapotranspiration elasticities of runoff, respectively.

226 According to the Budyko hypothesis, actual evapotranspiration ( $E$ ) is a function  
 227 of the dryness indices ( $\phi = E_0 / P$ ), specifically  $E = P * F(\phi)$ , and the precipitation and  
 228 potential evapotranspiration elasticities of stream flow can be expressed as:

$$229 \quad \varepsilon_p = 1 + \phi F'(\phi) / (1 - F(\phi)) \quad \varepsilon_{E_0} = -\phi F'(\phi) / (1 - F(\phi)) \quad (11)$$

230 The following formulae (one with a parameter and the others without) for the  
 231 Budyko hypothesis are often used to estimate  $F(\phi)$ , as shown in Tab. 1.

232 **Table 1** Different Formulae for the Budyko Hypothesis

233 According to Tab.1, the precipitation elasticity ( $\varepsilon_p$ ) and potential  
 234 evapotranspiration elasticity ( $\varepsilon_{E_0}$ ) can be determined, allowing the runoff change  
 235 caused by climate change ( $\Delta R_c$ ) to be calculated, and thus the contribution of climate  
 236 change can be assessed. Calibration of Zhang's (2001) formula using land cover and  
 237 land use conditions yielded a parameter value of 1.5.

### 238 **3.3 Improved climate elasticity method**

239 At the catchment scale, and over a long time period, the water balance equation can  
 240 be simplified as  $P = E + R$ . Here the mean annual runoff  $R$  can be divided into  
 241 observed runoff and changing runoff caused by human activities, i.e.  $R = R_{obs} + R_H$ ,  
 242 where  $R_{obs}$  is observed runoff, and  $R_H$  refers to mainly water consumption or water  
 243 intake by human activities which mainly include measures of water and soil  
 244 conservation, river dam construction, water intake from rivers, water transfer and so  
 245 on. So the water balance equation can be expressed as  $P = R_{obs} + R_H + E$ , which in  
 246 differential form is:

$$247 \quad dP = dR_{obs} + dR_H + dE \quad (12)$$

248 Meanwhile, according to the Budyko hypothesis  $E = PF(\phi)$ ,  $\phi = E_0 / P$ , the total  
 249 differential form can be expressed as:

$$250 \quad dE = [F(\phi) - \phi F'(\phi)] dP + F'(\phi) dE_0 \quad (13)$$

251 When substituted into Equation (12), this leads to:

$$252 \quad dR_{obs} = [1 - F(\phi) + \phi F'(\phi)] dP - F'(\phi) dE_0 - dR_H \quad (14)$$

253 After dividing equation (14) by  $R_{obs}$ , we obtain the following equation:

$$254 \quad \frac{dR_{obs}}{R_{obs}} = [1 - F(\phi) + \phi F'(\phi)] \frac{P}{R_{obs}} \frac{dP}{P} - F'(\phi) \frac{E_0}{R_{obs}} \frac{dE_0}{E_0} - \frac{R_H}{R_{obs}} \frac{dR_H}{R_H} \quad (15)$$

255 The climate and anthropic elasticities are calculated as follows:

$$256 \quad \varepsilon_P = [1 - F(\phi) + \phi F'(\phi)] \frac{P}{R_{obs}}$$

$$257 \quad \varepsilon_{E_0} = -F'(\phi) \frac{E_0}{R_{obs}}$$

$$258 \quad \varepsilon_H = -\frac{R_H}{R_{obs}} \quad (16)$$

259 Which satisfy  $\varepsilon_P + \varepsilon_{E_0} + \varepsilon_H = 1$ .

260 From Tab. 1 and Eq. (16), the elastic coefficients  $\varepsilon_P, \varepsilon_{E_0}, \varepsilon_H$  which represent  
 261 the precipitation, evapotranspiration, and human activities elasticities, can be  
 262 calculated. Next, the runoff change  $R_H$  caused by human activities is computed. The  
 263 contribution of human activities to runoff can be expressed by  $P_H$ ,  
 264 where  $P_H = R_H / \Delta R$ , thus the contribution of climate change can also be calculated.

## 265 4. Results and Discussion

### 266 4.1 Detection of hydrologic changes

267 In this study, the Mann–Kendall trend test is adopted to determine the significance of  
268 the trends in runoff, precipitation and potential evapotranspiration, and to analyze the  
269 trends in meteorological factors and corresponding runoff changes over nearly 50  
270 years. Figure 2 shows time series of precipitation, potential evapotranspiration and  
271 runoff from 1958-2008, and the runoff series are observed at the hydrological gauging  
272 station located the river basin outlet. Qualitative inspection shows that the trends in  
273 precipitation and potential evapotranspiration are not obvious, while runoff notably  
274 decreases. The fluctuation range of potential evapotranspiration is not obvious in  
275 different years, but the fluctuation range of precipitation is significant, and its overall  
276 trends are stable. Thus, the decrease of runoff implies that precipitation and potential  
277 evapotranspiration are not the only influencing factors on runoff, and instead human  
278 activities may have had the main influence on the decreasing runoff.

279 **Figure 2** Time series of annual precipitation, annual potential evapotranspiration and annual  
280 runoff in the Wei river basin from 1958-2008

281 The Mann-Kendall test is also used to analyze the change point of the runoff in  
282 the Wei River basin, at a confidence level set to  $\alpha=0.05$ . The normal distribution  
283 shows that the critical value was  $U_{\alpha/2}=1.96$ . The result of change point test is  
284 presented in Fig. 3. It can be seen in Figure 3 that the two curves intersect in 1990,  
285 and the intersection is within the critical value range  $U_{\alpha/2}=\pm 1.96$ . The result illustrates  
286 that an abrupt change of runoff occurred in 1990.

287 **Figure 3** The result of the abrupt change point test in the Wei river basin

288 According to the results of trend analysis and change-point analysis, the monthly  
289 runoff data in the periods of 1958-1990 and 1990-2008 is used to plot the Flow  
290 Duration Curve (FDC) that indicates the runoff change of the basin in different  
291 periods, and the monthly runoff series are observed at the hydrological gauging  
292 station located the river basin outlet. Vogel and Fennessey (1994) provides the details  
293 of FDC method, which represents the relationship between the magnitude and  
294 frequency of runoff, providing an estimate of percentage of time a given runoff is  
295 equal or exceeded over a historical period. The relationship between the magnitude  
296 and frequency of monthly average runoff are shown in Fig.4, which indicates the  
297 percentage of time runoff is exceeded in the period of 1958-1990 is larger than that in  
298 1990-2008, and the runoff relative change for the two periods only have a large  
299 fluctuations at the percentage that is less than 10% and more than 90%. The Fig.4 also  
300 implies the decrease of runoff in 1990-2008 has the correlations with human activities,  
301 and the influence on the decreasing runoff is easier to happen in the high-flow and  
302 low-flow periods at which the percentage time runoff is exceeded is less than 10%  
303 and more than 90%.

304 **Figure 4** Flow duration curves under different periods in the Wei river basin

## 305 **4.2 Results of the original and improved climate elasticity methods**

### 306 **4.2.1 Original climate elasticity method**

307 In order to evaluate the influence on runoff caused by climate change, firstly,  
308 Equation (11) is used to calculate the elasticities of precipitation and potential

309 evapotranspiration for 1958-2008, which can be expressed as  $\varepsilon_p$  and  $\varepsilon_{E_0}$ ,  
310 respectively, as shown in Tab. 2. These results reveal that if precipitation decreases by  
311 10%, runoff will decrease by 25.8%~27.7%, and if potential evapotranspiration  
312 decreases by 10%, runoff will increase by 15.8%~17.7%. Then, according to the  
313 calculated  $\varepsilon_p$ ,  $\varepsilon_{E_0}$  and Eq. (10), the runoff decrease caused by climate change can be  
314 computed. The total contribution of precipitation and potential evapotranspiration to  
315 the runoff decrease is 32.1 mm. Therefore, the contribution of climate change to the  
316 runoff decrease is 37%~40%, and the contribution of human activities is 60%~63%.

317 **Table 2** Results of the original climate elasticity method

#### 318 **4.2.2 Improved climate elasticity method**

319 In order to evaluate the influence on runoff caused by human activities, firstly,  
320 Equation (16) is used to calculate the elasticities of precipitation, potential  
321 evapotranspiration and human activities for 1958-2008, which can be expressed as  $\varepsilon_p$ ,  
322  $\varepsilon_{E_0}$  and  $\varepsilon_H$ , respectively, as shown in Tab.3. We realize that if annual runoff decreases  
323 by 32.1 mm, the decrease caused by human activities is 22.9~24.9 mm. So the  
324 contribution of human activities to runoff decrease is 71%~78%, and the contribution  
325 of climate change is 22%~29%.

326 **Table 3** Results of the improved climate elasticity method

#### 327 **4.2.3 Comparison of the simulation results**

328 In this paper, two methods are used to analyze the causes of the runoff decrease in the  
329 Wei River basin. One is the original climate elasticity method, and the other is the  
330 improved climate elasticity method. Each method adopts two formulas based on the

331 Budyko hypothesis (noting that one formula includes parameters and the other does  
332 not). Results are compared with each other and with those calculated by other  
333 methods, and the precision of the two methods is analyzed. Both methods are  
334 implemented at the annual time scale, and require relatively simple computation. The  
335 contributions of climate change and human activities to runoff variability can thus be  
336 computed by fewer data and parameters compared with other methods. The results of  
337 the original climate elasticity method show that the contributions of climate changes  
338 and human activities to runoff variability are 37%~40% and 60%~63%, respectively.  
339 Meanwhile, corresponding contributions calculated using the improved climate  
340 elasticity method are 22%~29% and 71%~78%, respectively. Early studies showed  
341 that during 1970-1995 the contribution of human activities to runoff decrease was  
342 58.3% in the Wei River basin (Zhang and Wang, 2007). In recent years, human  
343 activities have intensified, so by 2008 the contribution of human activities should  
344 have increased, and may now exceed 60%. Gao et al. (2013) found that the  
345 contribution of human activities to reduced stream flow in the Wei River basin was  
346 even as high as 82.80%. Zhan et al.(2014) used the SIMHYD model to partition the  
347 effect of climate change and human activities on surface runoff in the Wei River basin  
348 and found that the contribution rate of human activities to stream flow change was  
349 more than 65%.The results of the improved climate elasticity method are closer to the  
350 existing results than those of the original, suggesting that the improved climate  
351 elasticity method, which is more adaptable and easier to implement, is much more  
352 reliable and practical.

353 Moreover, it is important to note that the improved climate elasticity method is  
354 the first to introduce human activities elasticity  $\varepsilon_H$ . Without the trend analysis and  
355 change-point test, the strength of the influence of human activities on runoff changes  
356 can be calculated. When calculating the contribution of human activities to runoff  
357 changes, the change-point test is needed so that the total runoff decrease can be  
358 calculated.

359 The improved climate elasticity method broadens the concept of climate  
360 elasticity, and provides more intuitive and practical formula for calculating the  
361 contribution of human activities to runoff changes. Compared with the hydrological  
362 simulation method, the climate elasticity method not only needs fewer data and  
363 parameters, and is more reliable and easier to implement, but can also be easily  
364 extended. However, its temporal resolution is low and it lacks a physical basis. There  
365 is a trend towards coupling hydrological simulations with the more reliable  
366 hydrological and meteorological statistical methods, to quantitatively study  
367 hydrological responses to climate change and human activities.

### 368 **4.3 Discussion**

369 In the paper it is assumed that over a long period of time, change in catchment storage  
370 can be neglected so that the water balance equation can be expressed as  $P=E+R$ . And  
371 it is also assumed that the Budyko curve can comparably precisely estimate mean  
372 annual evaporation. In fact the two assumptions are fundamental assumptions and  
373 commonly used in Budyko-type elasticity studies for long term average (Gentine et al.,  
374 2012). We would not think there is any approach that can estimate the mean annual

375 evaporation “precisely” but if any, the Budyko curve would be comparable at least for  
376 long term mean.

377 The maximum daily precipitation and also averaged the top five maximum daily  
378 precipitation for each year are estimated as shown in Fig.5. The results show that  
379 there is no steady decreasing trend in these two measures of extreme rainfall and  
380 distribution, while the steady decrease in runoff investigated in this study would  
381 require a steady decrease in rainfall intensity if the change in distribution is the cause.  
382 Those results are consistent with our experience about this catchment that  
383 precipitation and potential evapotranspiration are not the only influencing factors on  
384 runoff, and instead human activities have had the main influence on the decreasing  
385 runoff.

386 **Figure 5** The top five maximum daily precipitation and the maximum daily precipitation curves

387 The impacts of human activities on runoff are reflected in land use and land  
388 cover changes. Land use and land cover change is a gradual process, and the impacts  
389 on runoff also accumulate gradually. We can decrease the uncertainty of quantitative  
390 predictions by analyzing the LUCC changes in the Wei River basin in 1980, 1990,  
391 2000, 2005 and 2007, and then checking whether the results are reasonable. The main  
392 type of land cover in the Wei River basin is cultivated land, which covers more than  
393 50% of the total area, followed by woodland and grassland.

394 By analyzing the changing areas of cultivated land and woodland and grassland  
395 in the Wei river basin in 1980, 1990, 2000, 2005 and 2007, it can be concluded that  
396 the year 1990 is the turning point in cultivated land area, since the area decreases

397 during 1980-1990, and then begin to increase again after 1990. The year 1990 is also  
398 the turning point in the area of woodland and grassland area, but the corresponding  
399 trends are opposite to those of the cultivated area. Because the area of cultivated land  
400 and woodland and grassland reaches 90%, and considering the turning points of the  
401 three types of land, the year 1990 can be regarded as a more general turning point in  
402 surface characteristics (Song et al., 2012). This decreases the uncertainty regarding  
403 1990 as the runoff change point.

404 Major sources of uncertainty in the simulation associated with the climate  
405 elasticity may arise from the input data, classification of the stages, and the parameter  
406 in the Zhang (2001) formula. Precipitation data used in the models are from 7 rain  
407 gauges and meteorological stations in and around the study catchment. The flow data  
408 are measured during 1958-2008 from the Hua County hydrological station, which is  
409 located at the downstream end of the basin, but may not sufficiently represent the  
410 whole basin. Even though the breakpoint test is found to be reasonable, there remain  
411 some uncertainties which may be caused by the test method and artifacts when  
412 reading the results in the chart. The parameter which is very sensitive in the Zhang  
413 (2001) formula is calibrated according to the land cover and land use conditions.

414 Furthermore, it is essential to point out that climate change and human activities  
415 are supposed to be two mutually independent variables when separating their impacts  
416 on runoff; however, we note that land use and land cover can be influenced by both  
417 climate change and by human activities.

418 In this study  $R$  is divided into  $R_{obs}$  and  $R_H$  because the water intake directly from

419 rivers is significant amount in almost all rivers in China. However in the original

420 Budyko-type elasticity this part is not considered directly and the main formulae are

421  $\Delta R = \varepsilon_P \frac{d_P}{P} + \varepsilon_{E_0} \frac{d_{E_0}}{E_0}$  and  $\Delta R = \Delta R_C + \Delta R_H$ . Then that framework is extended by

422 including the direct influence from water intake to adapt to catchments with intense

423 water consumption and intake, and the main formula is

424  $\Delta R = \varepsilon_P \frac{d_P}{P} + \varepsilon_{E_0} \frac{d_{E_0}}{E_0} + \varepsilon_H \frac{d_{R_H}}{R_H}$ . We believe this is a new contribution over the climate

425 elasticity method reported in literatures.

## 426 **5. Conclusions**

427 Decreasing runoff in many rivers in China has been reported in recent years, and the

428 Wei River basin is one of the most serious cases. This paper is aiming at developing a

429 new approach to quantifying the impact of climate variations and human activities on

430 this decreasing runoff in the Wei River basin. The man-made changes here include

431 land use, vegetation, and other land surface conditions, while climate change and

432 climate variability are reflected in precipitation and potential evapotranspiration. This

433 study uses the Mann-Kendall test to assess the temporal trends in precipitation,

434 potential evapotranspiration and runoff, and also analyzes the point of abrupt change.

435 On this basis, the original climate elasticity method and improved climate elasticity

436 method are used to analyze the quantitative hydrological effects of climate change and

437 human activities; these findings are then compared to existing results from the

438 hydrological simulation method. The study shows as following:

439 In the last 50 years, the runoff from the Wei River basin has obviously decreased,  
440 but the precipitation and potential evapotranspiration have shown no clear trend.  
441 Therefore it can be seen that climate factors have not obviously contributed to the  
442 runoff decrease.

443 From 1958 to 2008, the runoff in the Wei River basin shows an abrupt change in  
444 1990, effectively dividing the total runoff into natural runoff process and runoff  
445 process affected by human activities. At the same time, this change shows that human  
446 activities have significant effects on runoff.

447 Human activities resulted in a shift in land use and land cover in 1990, and the  
448 type of land use and land cover has a great influence on runoff. This illustrates that  
449 human activities, especially those causing land use and land cover change, are the  
450 main reason for the runoff decrease.

451 The original climate elasticity method shows that the contributions of climate  
452 change and human activities to runoff decrease are 37%~40% and 60%~63%,  
453 respectively, but the improved climate elasticity method indicated that the  
454 contributions of climate change and human activities to runoff decrease are 22%~29%  
455 and 71%~78%, respectively. The result of the improved climate elasticity method is  
456 closer to those existing results concluded based on comprehensive hydrological  
457 models, thereby demonstrating that human activities are the main reason for the  
458 runoff decrease in the Wei River basin.

459

460 **Acknowledgements**

461 This work was partially supported by the National Natural Science Foundation of  
462 China (41271003, 41371043, 50939006), Key Project for the Strategic Science Plan  
463 in IGSNRR, Chinese Academy of Sciences (2012ZD003), and the Open Research  
464 Fund of State Key Laboratory of Simulation and Regulation of Water Cycle in River  
465 Basin (China Institute of Water Resources and Hydropower Research)  
466 (IWHR-SKL-201311).

467

468 **References**

- 469 Bao, Z.X., Zhang, J.Y., Wang, G.Q. Fu, G.B., He, R.M., Yan, X.L., Jin, J.L., Liu, Y.L., and Zhang,  
470 A.J.:Attribution for decreasing runoff of the Haihe River basins, northern China: Climate  
471 variability or human activities?, *Journal of Hydrology* ,460-461, 117-129, 2012.
- 472 Chiew, F. H. S.: Estimation of rainfall elasticity of streamflow in Australia, *Hydrol. Sci. J.*, 51,  
473 613–625, 2006.
- 474 Dooge, J.C.I., Bruen, M., and Parmentier, B.: A simple model for estimating the sensitivity of  
475 runoff to long-term changes in precipitation without a change in vegetation, *Advance in Water*  
476 *Resources*, 23,153–163, 1999.
- 477 Fu, G., Charles S. P., and Chiew, F. H. S.: A two-parameter climate elasticity of runoff index to  
478 assess climate change effects on annual runoff, *Water Resources Research*,43,W11419,  
479 doi:10.1029/2007WR005890, 2007.
- 480 Gao, P., Geissen, V., Ritsema, C.J., Mu, X.M. and Wang, F.: Impact of climate change and  
481 anthropogenic activities on streamflow and sediment discharge in the Wei River basin,

482 China, *Hydrology and Earth System Sciences*, 17,961-972, 2013.

483 Gentine, P., P. D'Odorico, B. R. Lintner, G. Sivandran, and G. Salvucci, Interdependence of  
484 climate, soil, and vegetation as constrained by the Budyko curve, *Geophysical Research Letters*,  
485 39, L19404, doi:10.1029/2012GL053492, 2012.

486 Govinda R.P. :Effect of climate change on stream flows in the Mahanadi river basin, India, *Water*  
487 *Int*, 20,205–212, 1995.

488 Guo H., Hu Q., and Jiang T.: Annual and seasonal runoff responses to climate and land-cover  
489 changes in the Poyang Lake basin, China, *Journal of Hydrology*, 355, 106-122, 2008.

490 Hargreaves, G. H.,and Samani Z. A.: Estimating of potential evapotranspiration, *Journal of*  
491 *irrigation and Drainage Divdide*, *Proceedings of the American Society of Civil*  
492 *Engineers*,108,223-230, 1982.

493 He, H.M., Zhang, Q.F., Zhou, J., Fei J., and Xie X.P. : Coupling climate change with hydrological  
494 dynamic in Qinling Mountains, China, *Climatic Change*, 94, 409–427, 2009.

495 Hu, S.S., Zheng, H.X., Liu, C.M., Yu, J.J., and Wang Z.G.: Assessing the Impacts of Climate  
496 Variability and Human Activities on Streamflow in the Water Source Area of Baiyangdian Lake,  
497 *Acta Geographic Sinica*, 67,62-70, 2012.

498 IPCC, I. 2013. *Climate Change 2013: The Physical Science Basis*. Retrieved from  
499 <http://www.buildingclimatesolutions.org/view/article/524b2c2f0cf264abcd86106a>.

500 Kendall, M.G. : *Rank Correlation Measures*, Charles Griffin,London, 202, 1975.

501 Legesse, D., Vallet-Coulomb, C., and Gasse, F.:Hydrological response of a catchment to climate  
502 and land use changes in Tropical Africa: case study South Central Ethiopia, *Journal of*  
503 *Hydrology*, 275, 67–85, 2003.

504 Luo, J., and Rong, Y.S.: Evaluation on several Empirical methods for estimating potential  
505 evapotranspiration in North China[OL], 2007. (in Chinese).

506 Ma, H., Yang, D.W., Tan, S.K., Gao, B., and Hu, Q.F. :Impact of climate variability and human  
507 activity on streamflow decrease in Miyun Reservoir catchment, Journal of  
508 Hydrology ,389,317–324, 2010.

509 Ma, Z.M., Kang, S.Z., Zhang, L., Tong, L.,and Su, X.L. : Analysis of impacts of climate change  
510 and human activity on runoff for a river basin in arid region of northwest China, Journal of  
511 Hydrology, 352,239–249, 2008.

512 Mann, H. B.:Nonparametric tests against trend. *Econometrica*, Journal of the Econometric Society  
513 13(3), 245-259, 1945.

514 Milly, P. C. D., Dunne, K. A., and Vecchia, A. V. : Global pattern of trends in runoff and water  
515 availability in a changing climate, *Nature* ,438,347–350, 2005.

516 Milly, P.C.D.,and Dunne KA. : Macro-scale water fluxes 2. Water and energy supply control of  
517 their inter-annual variability. *Water Resources Research*, 38,1206, doi:10.1029/2001WR000760,  
518 2002.

519 Nash, L. L., and Gleick, P. H.: Sensitivity of streamflow in the Colorado basin to climatic changes,  
520 *Journal of Hydrology*, 125, 221–241, doi:10.1016/0022-1694(91)90030-L, 1991.

521 Piao, S. L., Friedlingstein, P., Ciais, P., de Noblet-Ducoudre, N., Labat, D., and Zaehle, S.:  
522 Changes in climate and land use have a larger direct impact than rising CO<sub>2</sub> on global river  
523 runoff trends, *Proc. Natl. Acad. Sci*, 104(39),15242-15247, 2007.

524 Pike, J. G. The estimation of annual run-off from meteorological data in a tropical climate, *Journal*  
525 *of Hydrology*, 2,116–123, 1964.

526 Qiu, G. Y., Yin, J., and Shu, G.: Impact of climate and land-use changes on water security for  
527 agriculture in Northern China, *Journal of Integrative Agriculture* ,11, 144-150, 2012.

528 Revelle, R. R., and Waggoner , P. E.: Effects of a carbon dioxide-induced climatic change on  
529 water supplies in the western United States, in *Changing Climate*, edited by Carbon Dioxide  
530 Assess. Comm. et al., pp. 419–432, Natl. Acad., Washington, D. C., 1983.

531 Risbey, J. S., and D. Entekhabi. :Observed Sacramento basin stream-flow response to precipitation  
532 and temperature changes and its relevance to climate impact studies, *Journal of Hydrology*  
533 184,209–223, doi:10.1016/0022-1694(95)02984-2, 1996.

534 Sankarasubramanian, A., Vogel, R. M., and J. F. Limbrunner. :Climate elasticity of runoff in the  
535 United States,*Water Resources Research*, 37,1771–1781, doi:10.1029/2000WR900330, 2001.

536 Schaake, J. C.:From climate to flow, in *Climate Change and U.S. Water Resources*.edited by  
537 Waggoner, pp. 177–206, John Wiley, New York, 1990.

538 Song, J.X., Xu, Z.X., Liu, C.M.,and Li, H.E.:Ecological and environmental instream flow  
539 requirements for the Wei River—the largest tributary of the Yellow River, *Hydrological*  
540 *Processes*, 21, 1066–1073, 2007.

541 Song,W.N., Zhan C.S., Li J.Y., Xu Z.X., Wang H.X.:Remote Sensing Analysis on the  
542 Spatial-temporal Characteristics of LUCC in the Guanzhong Plain Region of the Wei River  
543 Basin in the Past 30 Years, *China Land Sciences*, 2,56-62, 2012(in Chinese).

544 Sun, F., Roderick ,M. L., and Farquhar, G. D.: Changes in the variability of global land  
545 precipitation, *Geophys. Res. Lett.*,39, L19402, doi:10.1029/2012GL053369, 2012.

546 Turc, L. Le bilan d'eau des sols. Relations entre les precipitations,l'evaporation et l'ecoulement,  
547 *Ann. Agronomy*, 5,491–596, 1954.

548 Vogel R.M. and Fennessey N.M.: Flow-duration curves. I: New interpretation and confidences  
549 intervals, *Journal of Water Resources Planning and Management*, 120,485-504, 1994.

550 Vogel, R. M., I. Wilson, and C. Daly . Regional regression models of annual streamflow for the  
551 United States, *J. Irrig. Drain. Eng.*, 125,148–157,  
552 doi:10.1061/(ASCE)0733-9437(1999)125:3(148), 1999a.

553 Wang, G.S., Xia, J., and Chen, J.: Quantification of effects of climate variations and human  
554 activities on runoff by a monthly water balance model: a case study of the Chaobai River basin  
555 in northern China, *Water Resources Research* ,45, W00A11, 2009.

556 Zhan, C.S., Zeng S.D., Jiang S.S., Wang H.X., Ye W.: An integrated approach for partitioning the  
557 effect of climate change and human activities on surface runoff. *Water Resources Management*,  
558 WARM-D-14-00199R1, doi: 10.1007/s11269-014-0713-0, 2014.

559 Zhang, J.Y., and Wang, G.Q. Research on climate change impact on hydrology and water  
560 resources[M], 2007 (in Chinese).

561 Zhang L., Dawes W.R. and Walker G.R.: Response of mean annual evapotranspiration to  
562 vegetation changes at catchment scale, *Water Resources Research* 37, 701-708, 2001.

563 Zhang, X.P., Zhang, L., Zhao, J., Rustomji, P., and Hairsine, P.: Response of streamflow to  
564 changes in climate and land use/cover in the Loess Plateau, China, *Water Resources*  
565 *Research* ,44,W00A07, 2008.

566 Zheng, H., Zhang L., Zhu R., Liu C., Sato Y., and Fukushima Y.: Responses of runoff to climate  
567 and land surface change in the headwaters of the Yellow River Basin, *Water Resources*  
568 *Research* ,45,W00A19, doi:10.1029/2007WR006665, 2009.

569 **Table 1.** Different Formulae for the Budyko Hypothesis

Expression	$F(\phi)$	$F'(\phi)$
Turc (1954); Pike (1964)	$(1 + \phi^{-2})^{-0.5}$	$1 / \left[ \phi^3 \left( 1 + (1/\phi)^2 \right)^{1.5} \right]$
Zhang (2001)	$(1 + w\phi) / (1 + w\phi + 1/\phi)$	$(w + 2w/\phi - 1 + 1/\phi^2) / (1 + w\phi + 1/\phi)^2$

570

571

**Table 2.** Results of the original climate elasticity method

Period	Formula	$P$ /mm	$E_0$ /mm	$\varepsilon_p$	$\varepsilon_{E_0}$	$\Delta R_c$ /mm	$\Delta R_c$ /%	$\Delta R_H$ /%
1958-2008	Turc (1954); Pike (1964)	569.3	923.1	2.77	-1.77	11.9	37	63
1958-2008	Zhang (2001)	569.3	923.1	2.58	-1.58	12.9	40	60

572

573

574

**Table 3.** Results of the improved climate elasticity method

Period	Formula	$P$ /mm	$E_0$ /mm	$\varepsilon_p$	$\varepsilon_{E_0}$	$\varepsilon_H$	$\Delta R_H$ / mm	$\Delta R_C$ / %	$\Delta R_H$ / %
1958-2008	Turc (1954); Pike (1964)	569.3	923.1	3.53	-2.16	-0.37	22.9	29	71
1958-2008	Zhang (2001)	569.3	923.1	3.89	-2.49	-0.40	24.9	22	78

575

576

577

578


579

580

581

582


583



584

585

**Figure 1** Location of the meteorological and hydrological stations used in this study



586

587


**Figure 2** Time series of annual precipitation, annual potential evapotranspiration and annual

588

runoff in the Wei river basin from 1958-2008


589

590




591  
592  
593

**Figure 3** The result of the abrupt change point test in the Wei river basin



594  
595  
596  
597  
598

**Figure 4** Flow duration curves under different periods in the Wei river basin



599

600 **Figure 5** The top five maximum daily precipitation and the maximum daily precipitation curves