

Impacts of
ENSO/PDO on
precipitation and
streamflow in China

R. Ouyang et al.

Linkages between ENSO/PDO signals and precipitation, streamflow in China during the last 100 years

R. Ouyang^{1,2}, W. Liu³, G. Fu⁴, C. Liu¹, L. Hu⁵, and H. Wang⁶

¹Key Laboratory of Water Cycle & Related Land Surface Processes, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

²Bureau of Comprehensive Development Ministry of Water Resources, Beijing 100053, China

³Key Laboratory of Tibetan Environmental Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China

⁴CSIRO Land and Water, Private Bag 5, Wembley, WA 6913, Australia

⁵Zhejiang Provincial Hydrology Bureau, Hangzhou, 310009, China

⁶Graduate School of INSA, Rennes, 35000, France

Received: 4 March 2014 – Accepted: 31 March 2014 – Published: 15 April 2014

Correspondence to: W. Liu (liuw@itpcas.ac.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

This paper investigates the single and combined impacts of the El Niño-Southern Oscillation (ENSO) and the Pacific Decadal Oscillation (PDO) on precipitation and streamflow in China over the last century. Results indicate that the precipitation and streamflow overall decrease during the El Niño periods/PDO warm phase while mainly increase during the La Niña periods/PDO cool phase in the majority of China, although there are still regional and seasonal differences. Specifically, the precipitation/streamflow in the Yellow River basin, Yangtze River basin and Pearl River basin are more significantly influenced by El Niño and La Niña events compare to those in the Songhua River basin among different months, especially in the October and November. Moreover, the significant influences of ENSO on streamflow in the Yangtze River mainly occur in summer and autumn while that in the Pearl River primary occur in the winter and spring. The precipitation/streamflow are relatively more in the warm PDO phase in the Songhua River basin and several parts of Yellow River basin while are relatively less in the Pear River basin and the most parts of the northwest China compare to those in the cool PDO phase, though there are rarely significances clarified using the Wilcoxon signed ranks test. When considering the combined influences of ENSO and PDO, the responses of precipitation/streamflow are shown to be opposite from northern China to southern China, with the ENSO-related precipitation/streamflow enhance in the northern China during the warm PDO phases and that increase in the southern China in the cool PDO phases. This study conducted would beneficial for understanding how the precipitation/streamflow responses to the changing climate and would correspondingly provide valuable references for the water resources prediction and management over China.

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



1 Introduction

It is well known that El Niño-Southern Oscillation (ENSO) is an important factor influencing the interannual climate variability over East Asia (Zhou and Wu, 2010). The warm ENSO, which is also called El Niño, is usually accompanied by a weaker than normal East Asian winter monsoon (Zhang et al., 1996; Wang et al., 2008) and consequently induces a warmer and wetter climate over East Asia during El Niño winters (Li, 1990; Wen et al., 2000). As an example, the ENSO influences can persist to the following summer, with significantly abundant precipitation and annual maximum streamflow over the Yangtze River valley during the decaying stage of El Niño event (Huang and Wu, 1989; Zhang et al., 2007). However, the aforementioned anomalies are generally reverse during the cool ENSO phase, namely La Niña events (Wang et al., 2008).

Some previous studies (Latif and Barnett, 1996; Mantua et al., 1997; Cayan et al., 1998; Nigam et al., 1999; Higgins and Shi, 2000; Minobe, 2000; Neal et al., 2002; Krishnan and Sugi, 2003; Wang et al., 2008) have indicated that the interannual relationship between ENSO and global climate is not stationary and the Pacific Decadal Oscillation (PDO), which is a largely interdecadal oscillation, could modulate the interannual ENSO-related teleconnections. For instance, the already enhanced precipitation and streamflow in eastern Australia are demonstrated to be even further magnified during La Niña events that occurred in the PDO/IPO (Interdecadal Pacific Oscillation) cool phase (Verdon et al., 2004). Additionally, the precipitation patterns showed different responses in the El Niño periods for Southeastern South America and Myanmar during the PDO warm/cool phase (Silva et al., 2011; Sen Roy and Sen Roy, 2011). These studies mentioned indicated that the in phase/out-of-phase of ENSO and PDO usually have distinct effects on precipitation and streamflow in different regions, and thus, the discussions considering the influences of ENSO in association with PDO are quite necessary in the related studies. There are various studies extensively documented the linkages between ENSO/PDO and annual/seasonal precipitation over

HESSD

11, 4235–4265, 2014

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

China during the past several decades (Liu and Ding, 1995; Gong and Wang, 1999; Zhang et al., 1999; Wu et al., 2003; Zhu and Yang, 2003; Xu et al., 2004; Li et al., 2005; Chan and Zhou, 2005; Ma and Shao, 2006; Hao et al., 2008; Zhou and Wu, 2010). For example, Zhou and Wu (2010) revealed that the warm ENSO mainly led to lower-level southwesterly winds deflect from the southeast coast of China and consequently influenced the winter precipitation in southern China. In addition, Chan and Zhou (2005) found that there was less precipitation over South China Monsoon Region during the period of high PDO index and vice versa. However, majority of aforementioned studies did not consider the combined influences of both ENSO and PDO on regional precipitation. On the other hand, streamflow, as a comprehensive integrator of rainfall over basin areas, also related to the variations of ENSO and PDO signals. If a strong relationship between river discharge and ENSO/PDO can be quantified, the streamflow forecasting, which is vital for effective water resource management, would be highly improved. Although many studies have been conducted nowadays on the relations between river streamflow and ENSO/PDO nowadays in China (Chen and Xu, 2005; Fu et al., 2007; Xu et al., 2007; Zhang et al., 2007; Lü et al., 2011), as far as the authors are aware, there has not been a related study documenting the combined influences of both ENSO and PDO signals on streamflow in the major large rivers over China. Considering all of the above, in this paper, the possible influences of ENSO and PDO, coupled and separately, on the annual/monthly precipitation and streamflow are conducted over China. Additionally, the precipitation and annual streamflow datasets adopted in this study were extended to the last 100 years (1901–2009) and full seasonal cycles were considered for presenting more reliable climate variability. The paper is organized as follows. Section 2 introduces the datasets and methodologies used. Section 3 examines the relationships among PDO, ENSO, precipitation and streamflow, and finally, the conclusions and proposed future research are presented in Sect. 4.

2 Data and method

2.1 Data

The precipitation data (1901–2009) were extracted from the newest Climatic Research Unit (CRU) Time Series (TS) 3.10 high resolution gridded datasets (http://badc.nerc.ac.uk/view/badc.nerc.ac.uk__ATOM__dataent_1256223773328276) at the University of East Anglia (Mitchell and Jones, 2005). The monthly CRU TS3.10 datasets, which were calculated on high-resolution ($0.5^\circ \times 0.5^\circ$) grids based on more than 4000 weather stations distributed around the world (with more than 160 meteorological station from China), were validated well matched with the observations over China except for the western Tibetan Plateau (Ma and Shao, 2006). In addition, one hundred years of continuous annual streamflow data and fifty to a hundred years of monthly streamflow data were collected in this study from the gauging stations located at the four major river basins, namely Songhua River basin, Yellow River basin, Yangtze River basin and Pearl River basin over China (Fig. 1 and Table 1). The basins selected covers approximately from the north to south of China, and are expected to be able to present the streamflow variability over China under climate change.

The ENSO index is represented by the Niño 3.4 SST defined as the January to March SST anomaly averages over the region ($5^\circ \text{S} - 5^\circ \text{N}$, $90 - 150^\circ \text{W}$), which is downloaded from the National Oceanic and Atmospheric Administration (NOAA, http://www.cgd.ucar.edu/cas/catalog/climind/Nino_3_3.4_indices.html) (Trenberth, 1997). The PDO index “is the leading empirical orthogonal function (EOF) of SST anomalies (January–March) in the North Pacific Ocean, poleward of 20°N ” (Mantua et al., 1997; Chan and Zhou, 2005) and is available at the Joint Institute for the Study of the Atmosphere and Ocean (JISAO) website: <http://jisao.washington.edu/pdo/> (Barnett et al., 1999).

HESSD

11, 4235–4265, 2014

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.2 Method

2.2.1 Precipitation and streamflow stratification according to El Niño and La Niña

ENSO is a quasi-periodic climate pattern that occurs across the tropical Pacific Ocean every several years (three to seven years' recurrence) which always couples two variations: the warm oceanic phase (El Niño) accompanies high air surface pressure in the western Pacific and the cold phase (La Niña) accompanies low air surface pressure in the western Pacific (Trenberth et al., 2007). Generally, it has been very difficult to define an El Niño/La Niña event and there is no universal single definition (Trenberth and Hoar, 1997; Anthony and Stewart, 2001; Fu et al., 2007). In this study, the definition of Trenberth (1997) is adopted that is "... an El Niño can be said to occur if 5 month running means of sea temperature (SST) anomalies in the Niño 3.4 region (5° N–5° S, 120–170° W) exceed 0.4 °C for 6 months or more.". Similarly, La Niña, the opposite event of El Niño, can simply be said to occur if 5 month running mean of SST anomalies below the threshold –0.4 °C (see Fig. 2a).

In this paper, the periods of El Niño events and La Niña events were used to stratify the precipitation and streamflow time series for analyzing the influences of El Niño and La Niña on hydro-climatic variables in China. The precipitation/streamflow time series were firstly extracted for each calendar month conditioned by El Niño/La Niña events, for instance, the multiyear mean value of January precipitation occurs during El Niño periods was treated as "January precipitation in El Niño". Finally, the sum of monthly precipitation from January to December in El Niño/La Niña months was treated as "annual" precipitation in El Niño/La Niña year (Fu et al., 2007).

HESSD

11, 4235–4265, 2014

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.2.2 Precipitation/streamflow stratification according to the PDO cool/warm phase

The PDO is a pattern of Pacific climate variability that shifts phases usually on at least 20–30 years' inter-decadal time scale (Mantua et al., 1997). It is detected as warm/cool surface water in the Pacific Ocean (north of 20° N), during a “warm” or “positive” phase, the west Pacific becomes cool and part of the eastern ocean warms while during a “cool” or “negative” phase, the opposite pattern occurs. The cool and warm PDO phases (Fig. 2) are identified from the PDO index series in accordance with the approach used in Mantua and Hare (2000) and Sen Roy (2011). Over the past century, the PDO was in a cool phase approximately during the periods 1901–1924, 1947–1976, and 1998–2009, and warm phase PDO regimes existed during the periods 1925–1946 and 1977–1997 (see Fig. 2b). It should be noted that these multi-decade epochs sometimes contain intervals of up to a few years in length in which the polarity of the PDO is reversed (e.g. the cool phase in 1998–2009 showed a warm phase in 2002–2005).

The precipitation/streamflow spanning the period 1901–2009 are stratified into two segments conditioned on the PDO warm/cool phase. Further, the series in warm PDO–El Niño, warm PDO–La Niña, Cool PDO–El Niño, and Cool PDO–La Niña are stratified used the method similar to the Sect. 2.2.1 from the precipitation/streamflow series extracted for PDO warm/cool phase, separately. Additionally, Wilcoxon signed ranks test were adopted to determine if average precipitation/streamflow received during PDO warm phases/La Niña periods was statistically different from that received during PDO cool phase/El Niño periods. It is a nonparametric test equivalent to the dependent t test, which does not assume normality in the data and could be used for the case that there are only small number of samples available for analysis (Kolivras and Comrie, 2007).

HESSD

11, 4235–4265, 2014

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3 Results and discussion

3.1 Perspective impacts of ENSO on precipitation and streamflow over China

3.1.1 Precipitation impacts of El Niño and La Niña events

Compare to the long-term average (1901–2009), the “annual” precipitation changes in El Niño and La Niña periods are spatially opposite (Fig. 3). For example, the overall “annual” precipitation increase in the North China Plain, southwest China as well as the Tibetan Plateau while decline in the northeast China, southeast China and northwest China during the La Niña periods (Fig. 3). However, the trends in the El Niño periods over these regions are obviously reversed. The Yangtze River can be spatially treated as a dividing line of ENSO influences on precipitation for eastern China, with the “annual” average precipitation is obviously less in the southern regions of Yangtze River in La Niña periods rather than that in El Niño years ($< -5\%$) while more ($> 5\%$) in the northern regions (including the Yellow River, Hai River and Huai River). It should also be noted that the results obtained in the Yellow River basin (similar to North China Plain) are consistent with many previous studies (Gong and Wang, 1999; Fu et al., 2007; Hao et al., 2008).

The influences of El Niño and La Niña on precipitation are found have obviously seasonal-cycle and monthly characteristics (Fig. 4). For instance, the ENSO impacts on precipitation in summer and autumn are more significant rather than winter and spring, especially for September, October and November. Moreover, the precipitation in southeast China (including parts of Pearl River and Yangtze River) are relatively larger during El Niño winter and spring while lower during El Niño summer and autumn compare to that during correspondingly La Niña periods. The possible reason is that southern coast of the South China are always influenced by different anomalous circulation systems between wet season and dry seasons (Wu et al., 2003). In addition, the percentage changes for the wet season precipitation (June–September) between

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

El Niño and La Niña periods are similar to that for “annual” precipitation, because that more than 40 % of the total annual precipitation falls in summer (Zhang et al., 2009).

The influences of El Niño and La Niña events on precipitation are also spatial unevenly distributed and are different from month to month over the entire China (Fig. 4). Although monthly precipitation changes between two ENSO phases over majority of regions do not statistically significant at the 0.05 level, some consistent and interesting results are still drawn. The overall influences of El Niño and La Niña on precipitation are more significant in the eastern and southern China (including the Pearl River, Yangtze River and Yellow River) rather than in the western and northern China (including the Songhua River). Correspondingly, the ENSO influences become increasingly weaker from Pearl River, Yangtze River and Yellow River to Songhua River. The reason maybe that the eastern and southern portions of China near the ocean with more total precipitation which is significantly influenced by the East Asian Monsoon and South Asian Monsoon. More specifically, the precipitation from November to March received from La Niña events are less than that received from El Niño events over almost the entire China and the tendencies reverse in the remaining seven months, especially in the wet seasons (June, July, August, September). While in October, the trends found above are reversed in most parts of Yellow river and Yangtze River.

In addition, precipitation responses patterns to El Niño/La Niña events are also discrepant among different parts of basin. As an example, the ENSO influences in the lower basin of Songhua River are opposite to the head and middle basin, which to some extent would lead to uniformly streamflow responses among different sub-basins. The difference responses for four river basins (or even for different parts of basin) to ENSO properly attribute to the spatially diverse influences of the different monsoon circulations and mid-latitudinal circulations. For example, the Pearl River basin is impacted by the retreating monsoon, East Asian Winter monsoon as well as the Taifoon-season, consequently, the precipitation-streamflow regime in sub-basin is considerable complex when response to the ENSO influences (Jiang et al., 2007; Zhang et al., 2011). Moreover, Li et al. (2010) indicated that the East Asia

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Summer Monsoon exhibited a southward shift in its major components due to the meridional asymmetric warming, which would maybe weaken the influences of East Asian Summer Monsoon on Songhua River basin and result in difference response to ENSO for Songhua River basin and other three basins.

3.1.2 Streamflow impacts of El Niño and La Niña events

The “annual” streamflow changes overall more in the La Niña periods relatively to that in the El Niño periods for all four basins, especially for the Yellow River basin (Fig. 5). Moreover, the ENSO influences on streamflow are spatial-temporally consistent with that on precipitation for the major river basins over China with obviously differences among months and basins. On the whole, the streamflow in the Yellow River basin, Yangtze River basin and Pearl River basin are more significantly influenced by El Niño and La Niña events compare to those in the Songhua River basin among different months, especially in October and November. The reason maybe those basins locate at southern or eastern China is proximity to the ocean, where generally larger precipitation amount are received compare to the inland regions due to the significantly influences of the East Asian Monsoon, South Asian Monsoon, and ENSO (Zhang et al., 1996). Although the streamflows in Songhua River basin for all twelve months in La Niña periods consistently increase while those in majority months (eight in twelve) in El Niño periods decrease compare to the multiyear average monthly streamflow during the past 100 years, only the La Niña impacts on August are found statistically significant. While in the Yellow River basin, the monthly streamflow trends influenced by El Niño/La Niña events are basically coincident with streamflow relatively lower than normal in El Niño periods and higher in La Niña periods almost for all months, though the statistical significance tests do not exhibit obviously seasonal characteristics (Fig. 5). The overall percentage difference between El Niño-related and La Niña-related streamflow is 32.1 %, and varies monthly from 10.1 % (March) to 59.7 % (November) (Fig. 5). Specifically, the streamflow in January, February, April, July, October and November change significantly between El Niño and La Niña events. Moreover, the

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



percentage changes of monthly streamflow are relatively smaller in spring (March–May) during the La Niña periods while larger in other seasons (especially in autumn), which are consistent with Fu et al. (2007) and Lü et al. (2011).

The significant influences of ENSO on streamflow in the Yangtze River mainly occur in summer and autumn while that in the Pearl River primary occur in the winter and spring. The spatial variability of streamflow is responsible for both the influences of El Niño mature phase on precipitation in summer when the intensified western Pacific subtropical high covers the southeastern periphery of China and the weakening of the Indian monsoon provides less moisture inflow to the northern part of China (Zhang et al., 1999). The streamflow responses to ENSO for Yangtze River basin (Hankou station) exhibit obvious seasonal variations (Fig. 5). For example, the streamflow are relatively higher in El Niño periods relatively to that in La Niña periods in winter (December–February) and spring while reverse in summer and autumn (September–November). Especially, compare to correspondingly average monthly streamflow, the differences of La Niña-related streamflow and El Niño-related streamflow change significantly in June, July, August and September. In the Pearl River basin (Wuzhou Station), the ENSO impacts seem to be more complicated. The absolute percentage difference of streamflow between La Niña and El Niño periods are all more than 10% from October to March, as well as July. In September, the streamflow in La Niña month exceeds that in El Niño month and their percentage difference exceed 63.0%. Different to Yangtze River basin, the ENSO influences in the Pearl River are only statistically significant (0.05 level) on autumn and winter streamflow, which possibly because that the regions Pearl River locates at is in tandem with the strengthening and weakening of sea surface temperature (SST) in western Pacific (Juneng and Tangang, 2005).

HESSD

11, 4235–4265, 2014

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

3.2 Perspective impacts of PDO on precipitation and streamflow

3.2.1 Variability of precipitation due to PDO impacts

The percentage changes of “annual” precipitation also show spatially opposite responses to the PDO warm phase and cool phase, although only changes over a few regions are statistically significant at 0.05 levels (Fig. 6). Specifically, the “annual” precipitation in most parts of northeast China and northwest China tend to be higher during the PDO warm phase relatively to that in the cool phase, especially in the Songhua River basin and in the inland watersheds of Yellow River (blue regions in Fig. 6b). The results obtained are consistent with Zhu and Yang (2003), which indicates that the summer precipitation (account for more than 50 % of total annual precipitation) in the northeast and northwest China increase during a warm PDO phase due to the weakening of East Asian Summer monsoon and the southward shift of Western Pacific Subtropical High. In contrast, the “annual” streamflow responses are found opposite over the North China Plain, southwest China and Central China with the precipitation to be less during the warm PDO phase and to be more during the cool phase (Yang et al., 2005; Fu et al., 2009). The results in the northern China areas maybe because that they always dominated by high pressure and experiencing precipitation decrease when the Pacific is in warm phase, with the sea temperature over tropical mid-eastern Pacific rises and that over the central part of northern Pacific is lower than normal (Yang et al., 2005). Additionally, the precipitation over the Yellow River basin (Fu et al., 2004), Yangtze River basin and Pearl River basin decrease from the mid and late 1970s to 1990s when the PDO is in a persistent warming phase, while increase after 2000 when PDO entered into an unstable cool phase.

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



3.2.2 Variability of streamflow due to PDO impacts

The “annual” streamflow changes shown in Fig. 7 are basically consistent with those for precipitation during warm and cool PDO phases against the long-term average, although there are no significant trends tested. The PDO influences in Songhua River basin are opposite to that in other three basins with the streamflow are obviously higher than the long-term average (6.1 %) in the PDO warm phase while are lower in the PDO cool phase (−4.0 %). The streamflow results related to the PDO warm/cool phase correspond to the division of streamflow dry/wet stages by Song et al.(2010), which indicated both the streamflow (Harbin station) for Songhua River basin significantly changed in the following stages: 1900–1907, 1915–1928, 1975–1980, and 1999–2005 as four dry stages, 1970–1974 as the medium water stage, and 1908–1914, 1929–1969, and 1981–1998 as the three wet stages. Instead, in the Yellow River, Yangtze River and Pearl River, the streamflow are relatively lower in the PDO warm phase and higher in the PDO cool phase and their percentage differences become increasing small from north to south. The results are consistent with Gordon and Giulivi (2004), which indicated that the high (low) runoff in the Yangtze River and Yellow River correspond to the PDO negative (positive) phase. Additionally, similar results are also found when replace the 100 years streamflow observations by 50 years for analyzing the connections between streamflow and PDO in Songhua River and Yangtze River (not shown). It should be indicated that the streamflow tendency in the downstream of Yellow River continued to decrease even in the PDO cool phase after 2000 maybe due to the influences of the human activities (Ren et al., 2002), for example, water withdrawal attributed to more than 60 % of the streamflow decrease in the downstream of Yellow River after 2000 (Zhang et al., 2011).

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



3.3 Combined influences of ENSO and PDO on both streamflow and precipitation

Many evidences (Chan and Zhou, 2005; Andreoli and Kayano, 2005) indicated that the PDO could modulate the ENSO teleconnections, which always effects the precipitation coupled with ENSO acting constructively (strong and well-defined anomalies) when they are in phase and destructively (weak and noisy anomalies) when they are out-of-phases. In this study, the precipitation/streamflow in El Niño periods are compared to that in La Niña periods during the PDO warm/cool phase, respectively (Fig. 8 and Fig. 9). Results show that the “annual” precipitation changes in El Niño/La Niña period compare to multi-year average in cool PDO phase are quite similar to Fig. 3, which indicates that the cool PDO phase do not significantly modulate the ENSO influences on precipitation. However, in the warm PDO phase, the percentage changes are obviously for the precipitation related to the El Niño/La Niña. For instance, in the northeast China and northwest China, the precipitation received from La Niña periods is obviously higher than that received from El Niño periods during the PDO warm phase while reverse during the PDO cool phase. Instead, these precipitation responses to the two PDO phases are almost opposite in the south China and central China, including the Yangtza River basin and the upper stream of the Pearl River basin.

The El Niño/La Niña-related streamflow in the four basins show different responses when simultaneously considering the PDO influences (Fig. 9). During the PDO cool phase, the streamflow in all basins tend to be higher in La Niña periods and be lower in El Niño periods. The results obtained are quite similar to the single impacts of El Niño/La Niña shown in Fig. 5, which indicate that the cool PDO also do not obviously change the El Niño/La Niña influences on streamflow anomalies. However, the cool PDO phase still acts both more negative anomalies in El Niño-related streamflow and more positive anomalies in La Niña-related streamflow in south China (including the Yangtze River basin and the Pearl River basin) and induces both less negative anomalies in El Niño-related streamflow and less positive anomalies in

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

La Niña-related streamflow in north China (including the Songhua River basin and the Yellow River basin). Moreover, it should also be noted that the streamflow and precipitation responses to El Niño/La Niña are opposite during the PDO cool phase in the Songhua River basin, which maybe because that the Harbin station, locates at the middle stream cannot fully represent the entire basin.

During the PDO warm phase, the streamflow received from La Niña periods is higher than that received from El Niño periods in Songhua River basin and Yellow River basin, with the change percentages 9.7 % and 44.1 %, respectively. Obviously, the warm PDO enhances the anomalies in both two basins during the La Niña periods and the El Niño periods. However, the situations are different in the southern China. For example, fewer differences between the El Niño-related and La Niña-related streamflow in Yangtze River basin are found with the overall percentage changes only 0.7 %, which indicates that compare to the cool PDO phase, the warm PDO phase weakens the ENSO influences in the Yangtze River basin. In the Pearl River basin, the La Niña-related streamflow tends to be lower than El Niño-related streamflow with the percentage difference of -21.8% . In addition, compares to the percentage changes in the PDO cool phase (10 %) and in the long-term average without considering the impacts of PDO (0.6 %), the warm PDO is proved increase the El Niño-related steamflow and decrease La Niña-related streamflow. In other words, similar to Andreoli and Kayano (2005), the warm PDO acts constructively influences in the north China and destructively influences in the south China. Overall, the El Niño/La Niña-related precipitation/streamflow experience similar variability during the warm/cool PDO phase except for the Songhua River basin in the cool PDO phase. Moreover, the streamflow, which is also influenced by many other factors such as global SST, longwave radiation, snow and human activities (Xu et al., 2007), seems to be more sensitive than the precipitation during the El Niño/La Niña periods in both warm and cool PDO phases (Fig. 9). However, the general influence patterns of the combined effects are basically consistent. Compare to the ENSO impacts, although the PDO indicator do not show significantly prediction capacity for annual streamflow probably

because of its multi-decadal cycles, the modulation effects on ENSO still deserve to be included in the researches that considering the long-term influences of ENSO on annual/seasonal/monthly water resources.

4 Summary and conclusions

This study investigated the single and combined impacts of ENSO and PDO on the precipitation/streamflow over China during the last century, which would enrich our knowledge for understanding their complex spatial-temporal teleconnections and provide a scientific basis for water resources prediction using ENSO/PDO as a potential predictor. The following conclusions can be drawn:

Overall, the El Niño events mainly decrease while La Niña events increase the precipitation/streamflow over China. However, there are considerable differences exist among months and basins, for example, the precipitation/streamflow changes in the Yellow River basin, Yangtze River basin and Pearl River basin are more significantly influenced by El Niño and La Niña events compare to those in the Songhua River basin among different months, which properly because that the precipitation/streamflow in the regions/basins close to the ocean seem more significantly influenced due to the mixed impacts of ENSO and other factors such as the East Asian Monsoon, South Asian Monsoon, and the Typhoon systems. Additionally, due to the influences of different circulation systems (Wu et al., 2003), the significant influences of ENSO on streamflow in the Yangtze River mainly occur in summer and autumn while that in the Pearl River primary occur in the winter and spring.

Although rarely significantly changes are detected, the influences of the PDO warm/cool phases on precipitation/streamflow are basically similar as but less than that of El Niño/La Niña. Spatially, the precipitation/streamflow in the Songhua River basin and parts of Yellow River basin are larger during the warm PDO phase than those during the cool PDO phase, while in the Pear River basin and the most parts of the northwest China these responses are reversed. When considering both the

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

influences of the PDO and ENSO, the responses for precipitation/streamflow are shown to be opposite between northern China and southern China. The El Niño-related precipitation/streamflow decrease while the La Niña-related precipitation/streamflow increases during the PDO warm phase in the northern China (including Songhua River basin and Yellow River basin), and the cool PDO phase do not change the El Niño/La Niña influences obviously on positive-negative streamflow anomalies.

The results obtained indicate that the monthly/seasonal ENSO could be a potential predictor for streamflow prediction in the Yangtze River, Pearl River or even for the Yellow River; however, further researches on the physical mechanism driving these relations are still necessary. Firstly, the influences should be further quantitatively conducted to enhance the forecast abilities of the ENSO/PDO indicator for streamflow and water resource modeling and forecasting. Additionally, there are also other factors influencing the streamflow changes which should be comprehensively considered in our future studies such as the East Asia Summer Monsoon (Wu and Wang, 2002) global SST, outgoing longwave radiation, sea level pressure, snow as well as human activities (Xu et al., 2007). Finally, ENSO/PDO events can be predicted one to two years in advance using the physical based coupled ocean–atmosphere models (Lü et al., 2011), their potential future states and influences of ENSO/PDO could be further conducted considering as much as possible aforementioned factors by coupling atmospheric/oceanic/land surface models with a proper distributed physical-based hydrological model.

Acknowledgements. This study is financially supported by the National Natural Science Foundation for Young Scholar of China (41101030), Postdoctoral Science Foundation of China (20100480443 and 201104141), the National Basic Research Program of China (2010CB428406), Strategic Priority Research Program of the Chinese Academy of Sciences (XDA05090309) and the Australia 2012 Endeavour Award.

References

- Andreoli, R. V. and Kayano, M. T.: ENSO-related rainfall anomalies in South America and associated circulation features during warm and cold Pacific decadal oscillation regimes, *Int. J. Climatol.*, 25, 2017–2030, 2005.
- 5 Anthony, S. and Stewart, W. F.: On the identification of ENSO-induced rainfall and runoff variability: a comparison of methods and indices, *Hydrolog. Sci. J.*, 46, 715–727, 2001.
- Barnett, T. P., Pierce, D. W., Latif, M., Dommenges, D., and Saravanan, R.: Interdecadal interactions between the tropics and midlatitudes in the Pacific basin, *Geophys. Res. Lett.*, 26, 615–618, 1999.
- 10 Cayan, D. R., Dettinger, M. D., Diaz, H. F., and Graham, N. E.: Decadal variability of precipitation over western North America, *J. Climate*, 11, 3148–3166, 1998.
- Chan, J. C. L. and Zhou, W.: PDO, ENSO and early summer monsoon rainfall over south China, *Geophys. Res. Lett.*, 32, L08810, doi:10.1029/2004GL022015, 2005.
- Chen, Y. N. and Xu, Z. X.: Plausible impact of global climate change on water resources in the
- 15 Tarim River Basin, *Sci. China Ser. D*, 48, 65–73, 2005.
- Fu, G. B., Chen, S. L., Liu, C. M., and Shepard, D.: Hydro-climatic trends of the Yellow River basin for the last 50 years, *Climatic Change*, 65, 149–178, 2004.
- Fu, G. B., Charles, S. P., Viney, N. R., Chen, S. L., and Wu, J. Q.: Impacts of climate variability on stream-flow in the Yellow River, *Hydrol. Process.*, 21, 3431–3439, 2007.
- 20 Fu, G. B., Charles, S. P., Yu, J. J., and Liu, C. M.: Decadal climatic variability, trends, and future scenarios for the North China Plain, *J. Climate*, 22, 2111–2123, 2009.
- Gong, D. Y. and Wang, S. W.: Impacts of ENSO on rainfall of global land and China, *Chinese Sci. Bull.*, 44, 852–857, 1999.
- Gordon, A. L. and Giulivi, C. F.: Pacific decadal oscillation and sea level in the Japan/East sea, *Deep-Sea Res. Pt. I*, 51, 653–663, 2004.
- 25 Hao, Z. X., Zheng, J. Y., and Ge, Q. S.: Precipitation cycles in the middle and lower reaches of the Yellow River (1736–2000), *J. Geogr. Sci.*, 18, 17–25, 2008.
- Higgins, R. W. and Shi, W.: Dominant factors responsible for interannual variability of the summer monsoon in the south-western United States, *J. Climate*, 13, 759–775, 2000.
- 30 Huang, R. H. and Wu, Y. F.: The influence of ENSO on the summer climate change in China and its mechanism, *Adv. Atmos. Sci.*, 6, 21–32, 1989.

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Jiang, T., Su, B. D., and Hartmann, H.: Temporal and spatial trends of precipitation and river flow in the Yangtze River Basin, 1961–2000, *Geomorphology*, 85, 143–154, 2007.
- Juneng, L. and Tangang, F. T.: Evolution of ENSO-related rainfall anomalies in Southeast Asia region and its relationship with atmosphere–ocean variations in Indo-Pacific sector, *Clim. Dynam.*, 25, 337–350, 2005.
- Kolivras, K. N. and Comrie, A. C.: Regionalization and variability of precipitation in Hawaii, *Phys. Geogr.*, 28, 76–96, 2007.
- Krishnan, R. and Sugi, M.: Pacific decadal oscillation and variability of the Indian summer monsoon rainfall, *Clim. Dynam.*, 21, 233–242, 2003.
- Latif, M. and Barnett, T. P.: Decadal climate variability over the North Pacific and North America: dynamics and predictability, *J. Climate*, 9, 2407–2423, 1996.
- Li, C. Y.: Interaction between anomalous winter monsoon in East Asia and El Niño events, *Adv. Atmos. Sci.*, 7, 36–46, 1990.
- Li, J. P., Wu, Z. W., Jiang, Z. H., and He, J. H.: Can global warming strengthen the East Asian Summer Monsoon?, *J. Climate*, 23, 6696–6705, 2010.
- Li, Q., Yang, S., Kousky, V. E., Higgins, R. W., Lau, K.-M., and Xie, P.: Features of Cross-Pacific climate shown in the variability of China and US precipitation, *Int. J. Climatol.*, 25, 1675–1696, 2005.
- Liu, Y. Q. and Ding, Y. H.: Reappraisal of the influence of ENSO events on seasonal precipitation and temperature in China, 19, 200–208, 1995 (in Chinese).
- Lü, A., Jia, S., Zhu, W., Yan, H., Duan, S., and Yao, Z.: El Niño–Southern Oscillation and water resources in the headwaters region of the Yellow River: links and potential for forecasting, *Hydrol. Earth Syst. Sci.*, 15, 1273–1281, doi:10.5194/hess-15-1273-2011, 2011.
- Ma, Z. G. and Shao, L. J.: Relationship between dry/wet variation and the Pacific Decade Oscillation (PDO) in northern China during the last 100 years, *Chin. J. Atmos. Sci.*, 30, 464–474, 2006 (in Chinese).
- Mantua, N. J. and Hare, S. R.: The Pacific decadal oscillation, *J. Oceanogr.*, 58, 35–44, 2002.
- Mantua, N. J., Hare, S. R., Zhang, Y., Wallace, J. M., and Francis, R. C.: A Pacific interdecadal climate oscillation with impacts on salmon production, *B. Am. Meteorol. Soc.*, 78, 1069–1080, 1997.
- Minobe, S.: Spatio-temporal structure of the pentadecadal variability over the North Pacific, *Prog. Oceanogr.*, 47, 381–408, 2000.

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Mitchell, T. D. and Jones, P. D.: An improved method of constructing a database of monthly climate observations and associated high-resolution grids, *Int. J. Climatol.*, 25, 693–712, 2005.
- 5 Neal, E., Todd Walter, M., and Coffeen, C.: Linking the pacific decadal oscillation to seasonal stream discharge patterns in Southeast Alaska, *J. Hydrol.*, 263, 188–197, 2002.
- Nigam, S., Barlow, M., and Berbery, E. H.: Analysis links Pacific decadal variability to drought and streamflow in the United States, *EOS T. Am. Geophys. Un.*, 80, 621–625, 1999.
- Ren, L. L., Wang, M. R., Li, C. H., and Zhang, W.: Impacts of human activity on river runoff in the northern area of China, *J. Hydrol.*, 261, 204–217, 2002.
- 10 Sen Roy, S.: Identification of periodicity in the relationship between PDO, El Niño and peak monsoon rainfall in India using S-transform analysis, *Int. J. Climatol.*, 31, 1507–1517, 2011.
- Sen Roy, S. and Sen Roy, N.: Influence of Pacific decadal oscillation and El Niño Southern oscillation on the summer monsoon precipitation in Myanmar, *Int. J. Climatol.*, 31, 14–21, 2011.
- 15 Silva, G. A. M., Drumond, A., and Ambrizzi, T.: The impact of El Niño on South American summer climate during different phases of the Pacific Decadal Oscillation, *Theor. Appl. Climatol.*, 106, 307–319, 2011.
- Song, X., Mu, X., Gao, P., Wang, F., and Wang, S.: Analysis on historical evolution and driving force of rainfall and runoff of Harbin Station in Songhua River, *Science of Soil and Water Conservation*, 8, 46–51, 2010 (in Chinese).
- 20 Trenberth, K. E.: The Definition of El Niño, *B. Am. Meteorol. Soc.*, 78, 2771–2777, 1997.
- Trenberth, K. E. and Hoar, T. J.: El Niño and climate change, *Geophys. Res. Lett.*, 24, 3057–3060, 1997.
- Trenberth, K. E., Jones, P. D., Ambenje, P., Bojariu, R., Easterling, D., Tank Klein, A., Parker, D., Rahimzadeh, F., Renwick, J. A., Rusticucci, M., Soden, B., and Zhai, P. M.: Observations: surface and atmospheric climate change, in: *Climate Change 2007: the Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chedvn, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK, 235–336, 2007.
- 30 Verdon, D. C., Wyatt, A. M., Kiem, A. S., and Franks, S. W.: Multidecadal variability of rainfall and streamflow: Eastern Australia, *Water Resour. Res.*, 40, W10201, doi:10.1029/2004WR003234, 2004.

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Wang, L., Chen, W., and Huang, R. H.: Interdecadal modulation of PDO on the impact of ENSO on the east Asian winter monsoon, *Geophys. Res. Lett.*, 35, L20702, doi:10.1029/2008GL035287, 2008.
- Wen, C., Graf, H. F., and Huang, R. H.: The interannual variability of East Asian winter monsoon and its relation to the summer monsoon, *Adv. Atmos. Sci.*, 17, 48–60, 2000.
- Wu, R., Hu, Z. Z., and Kirtman, B. P.: Evolution of ENSO-related rainfall anomalies in East Asia, *J. Climate*, 16, 3742–3758, 2003.
- Xu, K. Q., Brown, C., Kwon, H., Lall, U., Zhang, J. Q., Hayashi, S., and Chen, Z. Y.: Climate teleconnections to Yangtze river seasonal streamflow at the Three Gorges Dam, China, *Int. J. Climatol.*, 27, 771–780, 2007.
- Xu, Z. X., Takeuchi, K., and Ishidaira, H.: Correlation between El Niño Southern Oscillation (ENSO) and precipitation in Southeast Asia and the Pacific region, *Hydrol. Process.*, 18, 107–123, 2004.
- Yang, X. Q., Xie, Q., Zhu, Y. M., Sun, X. Q., and Guo, Y. J.: Decadal-to-interdecadal variability of precipitation in North China and associated atmospheric and oceanic anomaly patterns, *Chinese J. Geophys.*, 48, 789–797, 2005.
- Zhang, Q., Xu, C. Y., Jiang, T., and Wu, Y. J.: Possible influence of ENSO on annual maximum streamflow of the Yangtze River, China, *J. Hydrol.*, 333, 265–274, 2007.
- Zhang, Q., Xu, C. Y., Zhang, Z., Chen, Y. D., and Liu, C. L.: Spatial and temporal variability of precipitation over China, 1951–2005, *Theor. Appl. Climatol.*, 95, 53–68, 2009.
- Zhang, Q., Singh, V. P., Sun, P., Chen, X., Zhang, Z. X., and Li, J. F.: Precipitation and streamflow changes in China: changing patterns, causes and implications, *J. Hydrol.*, 410, 204–216, 2011.
- Zhang, R., Sumi, A., and Kimoto, M.: Impact of El Niño on the East Asian monsoon: a diagnostic study of the '86/87 and '91/92 events, *J. Meteorol. Soc. Jpn.*, 74, 49–62, 1996.
- Zhang, R., Sumi, A., and Kimoto, M.: A diagnostic study of the impact of El Niño on the precipitation in China, *Adv. Atmos. Sci.*, 16, 229–241, 1999.
- Zhou, L. T. and Wu, R. G.: Respective impacts of the East Asian winter monsoon and ENSO on winter rainfall in China. *J. Geophys. Res.*, 115, D02107, doi:10.1029/2009JD012502, 2010
- Zhu, Y. M. and Yang, X. Q.: Relationships between Pacific Decadal Oscillation (PDO) and climate variabilities in China, *Acta Meteorol. Sin.*, 61, 641–654, 2003 (in Chinese).

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.

Table 1. List of streamflow gauging stations used in this study.

River basin	Station	Drainage area (km ²)	Annual streamflow record period	Monthly streamflow record period
Songhua River (I)	Harbin	390 526	1901–2009	1901–1948, 1953–2004
Yellow River (II)	Sanmenxia	688 421	1901–2009	1953–2004
Yellow River (II)	Huayuankou	730 036	1901–2009	1950–2004
Yangtze River (III)	Hankou	1 488 036	1901–2009	1901–2004
Pearl River (IV)	Wuzhou	329 705	1901–2009	1950–2004

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.




Fig. 1. Map of China showing four major river basins (I: Songhua River basin; II: Yellow River basin; III: Yangtze River basin and IV: Pearl River basin) and streamflow gauging stations used in this study.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.




Fig. 2. The definition of ENSO events (El Niño and La Niña) from 5 month running mean series of Niño 3.4 SST index **(a)** and the partition of warm/cool phase PDO from monthly PDO index **(b)**.

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.




Fig. 3. Percentage changes of annual precipitation in El Niño months and La Niña months over the long-term average (1901–2009): **(a)** annual precipitation occurred in La Niña events; **(b)** annual precipitation occurred in El Niño events; and **(c)** annual precipitation occurred in La Niña events minus that occurred in El Niño events.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.




Fig. 4. Monthly precipitation changes in La Niña months over China compared to El Niño months. Regions change significantly at the 0.05 level based on the Wilcoxon signed ranks test are shown with shadow.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.




Fig. 5. Monthly and annual streamflow changes over four major river basins in China in El Niño months and La Niña month over the monthly/annual long-term average (1901–2004). The asterisks indicate the statistical significance based on Wilcoxon signed ranks test (lower than 0.05 is **, lower than 0.10 is * and otherwise is nothing). It should also be noted that the monthly streamflow changes in Yellow River basin and Pearl River basin were only calculated during the period 1950–2004 due to their limited availability of monthly streamflow data.

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.




Fig. 6. Percentage changes of annual precipitation in PDO warm phase and cool phase over the long-term average (1901–2009): **(a)** annual precipitation in POD cool phase; **(b)** annual precipitation in POD warm phase; **(c)** annual precipitation in POD cool phase minus that in POD warm phase. Regions change significantly at the 0.05 level based on the Wilcoxon signed ranks test are shown with shadow.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.




Fig. 7. Percentage changes of annual streamflow over four major river basins in China in PDO warm phase and cool phase over the long-term average (1901–2009). *P* values based on Wilcoxon signed ranks test are showed on the top/bottom of each bar.

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.




Fig. 8. Percentage changes of precipitation between El Niño periods and La Niña periods during the PDO warm phase (left panels) and the PDO cool phase (right panels).

Impacts of ENSO/PDO on precipitation and streamflow in China

R. Ouyang et al.




Fig. 9. Percentage changes of streamflow between El Niño period and La Niña periods during the PDO warm phase **(a)** and the PDO cool phase **(b)**.