

1 **Flood sensitivity of the Bavarian Alpine Foreland since the**
2 **late Middle Ages in the context of internal and external cli-**
3 **mate forcing factors**
4
5

6 O. Böhm¹, J. Jacobeit¹, R. Glaser² & K.-F. Wetzel¹
7
8

9 *1 Institute of Geography, University of Augsburg, Universitätsstraße 10, D-86135, Germany*
10 *oliver.boehm@geo.uni-augsburg.de, jucundus.jacobeit@geo.uni-augsburg.de,*
11 *karl-friedrich.wetzel@geo.uni-augsburg.de*
12

13
14
15 *2 Institute of Geography, Albert-Ludwigs-Universität Freiburg, Werthmannstraße 4, D-79085,*
16 *Germany*
17 *ruediger.glaser@geographie.uni-freiburg.de*
18
19

20
21 **Abstract**
22

23 This paper describes the flood sensitivity of the Bavarian part of the Alpine Foreland of Germany
24 and addresses different questions concerning climate variability and flood frequencies, from the 13th
25 century until today. Focal point of the paper is the flood frequencies of the superordinate spatial unit
26 of the Bavarian Foreland but not the ones of its single time series. Will recent climatic change modi-
27 fy the flood frequencies within the Bavarian Alpine Foreland or have the flood frequencies ~~been~~
28 varying due to altering climatic conditions since historical times? In the context of recent discus-
29 sions whether man-made climate change will modify the present state of flood frequencies, a look
30 back into the past is essential to understand the occurrence of floods in general and of recent floods
31 in particular. In order to understand climatic variability and changes in a comprehensive way, it is
32 necessary to review long time series. A perceived increase of summer floods in eastern Germany
33 and Bavaria since 1997 requires examination of long time series to estimate changes in flood fre-
34 quencies in a proper way. In view of the annual distribution of flood events within the Alpine Fore-
35 land of Germany, summer floods prove to be ^{the} most important. Based on written historical sources,
36 the flood history of the Alpine Foreland of Germany can be reconstructed back to the 14th century.
37 One major result is the occurrence of 'flood-rich' and 'flood-poor' episodes in almost cyclical se-
38 quences. Flood-rich periods, before the 16th century based on ~~weak amount of~~ ^{limited} available data, were
39 recorded in the periods 1300 – 1335, 1370 – 1450, 1470 – 1525, 1555 – 1590, 1615 – 1665, 1730 –
40 1780, 1820 – 1870, 1910 – 1955 as well as in a ninth period beginning in 1980. The flood-rich peri-
41 ods are characterized by longer flood duration. Most of the flood-rich and flood-poor periods (in
42 particular the beginning and the end of them) can be connected to changes in natural climate varia-
43 bility. These include changing sunspot numbers (as a measure of solar activity), so-called Little Ice
44 Age Type Events (LIATEs) as well as changes in the North Atlantic Oscillation (NAO). Climate
45 signals from external forcing factors, which could be used to explain the changing flood frequencies
46 in the Bavarian Alpine Foreland, end in 1930. Relationships within the climate system such as the
47 correlation of flood frequencies with the NAO have changed during the transition from the post Lit-
48 tle Ice Age period to the Modern Climate Optimum around 1930. Natural climate variability might
49 have been outperformed by anthropogenic climate change.
50

51 **Key words** Bavarian Alpine Foreland, flood history, flood frequencies, climate signals, forcing fac-
52 tors
53
54

This is not clear, what do you mean?

The

?

55

56

57 **1 Introduction**

58

59 Historical climatology, especially ^{with} the branch addressing historical floods, has gained
 60 increasing interest during the ~~last~~ ^{recent} decades. Different parts of Central Europe have
 61 been investigated, Schmoecker-Fackel & Naef (2010), Pfister (1984, 1996, 1999) and
 62 Wetter et al. (2010) ~~in previous instance~~ analyzed ^{the} the flood history of Switzerland,
 63 Brázdil et al. (2005) ~~have~~ examined the flood history of the Czech Republic, Böhm &
 64 Wetzel (2006), Mudelsee et al. (2004), Deutsch & Pörtge (2001, 2002), Deutsch et al.
 65 (2004) and Glaser (2008) have investigated different parts or catchments ~~areas~~ ^{basins} of Ger-
 66 many, while Rohr (2008, 2013) has examined extreme natural events in Austria;
 67 ~~Sturm et al. (2001) and Glaser & Stangl (2003a, b) have~~ analyzed
 68 ~~European~~ ^{Central} European flood histories ~~with a focus on Central Europe~~ ^{and} Kiss &
 69 Laszlovszky (2013) ~~have~~ examined parts ~~of the~~ ^{of the} Danube for the western and
 70 central Carpathian Basin.

71 Nevertheless, the flood history of the entire Bavarian Alpine Foreland has not been
 72 systematically analyzed until now (cf. Böhm 2011). Additionally, the Bavarian Alpine
 73 Foreland represents a region with a high susceptibility to climatic changes (cf. Auer et
 74 al. 2007). The Bavarian part of the Alpine Foreland of Germany (hereafter termed
 75 Bavarian Foreland) has experienced flood events on a regular basis, but the return pe-
 76 riods, e. g. for ~~strong~~ ^{strong} floods as well as for flood-rich or flood-poor periods, cannot be
 77 derived from ^a standard reference periods ~~of 30 years~~ ^{of 30 years}. All ~~of the~~ ^{of the} recent major summer
 78 floods have been triggered by cyclones following a special pathway (cf. van Bebber
 79 1891). This so-called Vb cyclone track seems to be the main precondition causing
 80 catastrophic flood events in the Bavarian Foreland, currently and also in the past (yet
 81 not every Vb cyclone affects the ^{whole} investigation area ~~in total~~). Reconstructions of historic
 82 weather patterns show the ~~emersion~~ ^{emersion} of this phenomenon in the past (cf. Böhm
 83 2011). The recent period, starting ⁱⁿ 1997, has experienced numerous floods triggered by
 84 Vb-conditions during the summer months. The so-called "(River) Oder Flood 1997"
 85 and the "Pfungsthochwasser (Whitsun Flood) 1999", both of which took place at the
 86 end of May, can be compared to the following summer floods of 2002 (Elbe/Danube
 87 Flood) and 2005 (Alps Flood). All these floods (despite their special naming) have
 88 affected the Bavarian Foreland.

89

90 To generate ~~a~~ long time series, it was necessary to integrate three different periods of
 91 flood documentation. The oldest pieces of information originate from the so-called
 92 period of documentary evidences and have been obtained from historical recordings
 93 such as chronicles and compilations from the late Middle Ages. These written records
 94 can be statistically analyzed, as depicted in Glaser (2008). From the year 1826 on-
 95 ward, data are available from the so-called early instrumental period (EIP) (cf. Jaco-
 96 beit et al. 1998). ~~From~~ ^{From} at least one representative gauging station in the lower sections
 97 of every alpine river recorded historic water levels, ^{can} be evaluated. From the begin-
 98 ning of the 20th century ~~and~~ ^{and} modern instrumental data (water level and dis-
 99 charge measurements) are available. The separately evaluated flood histories of the
 100 rivers Iller, Lech (with tributary Wertach) Isar, Inn and its tributary Salzach have been
 101 merged ^{for} one overall time series. The merging of the single time series should re-
 102 veal the flood-susceptibility of a superordinate spatial unit based on recent administra-

to form

Central

What is a strong flood? extreme?


emersion - new word.

103 tive borders, ^{with} ~~under~~ consideration ^{to} ~~of~~ climatic parameters. In the methods' section the
104 merging of the ~~single~~ time series is more elaborately described. Single flood events as
105 well as quantification of flood events do not stand in the limelight of the current pa-
106 per. The time line of the flood history of the Bavarian Foreland includes 584 individu-
107 al flood events (see methods' section).

108 **2 Investigation area**

109

110 In this paper the Bavarian Foreland is defined by the lower sections of the catchment
111 areas of the aforementioned rivers, from west to east the research area is bordered by
112 the river Iller in the west and the rivers Inn and Salzach in the east, as depicted in Fig.
113 1. The headwaters, however, are located in the Alps so that all of the rivers are north-
114 ern-alpine mountain rivers (apart from the river Wertach which should be regarded as
115 a foreland river). The main rivers Iller, Lech, Isar and Inn are Alpine tributaries of the
116 Danube river.
117



118

119 **Figure 1.** Investigation area "Bavarian Foreland" is bordered by the rivers Iller, Danube,
120 Inn/Salzach and the Alpine border (dashed line). Red spots are locating outstanding historical
121 locations and gauges.
122

123 All the rivers coming from the Northern Limestone Alps traverse the Flysch Zone,
124 enter the area of the faulted Molasse sediments and cross the belt of Pleistocene mo-
125 raines and gravel fields. The substratum of the lower sections of the rivers is formed
126 of ~~by~~ sandy sediments of the Molasse trough. All traversed geological formations differ
127 in their east-to-west extension. The geological formations of the outer-alpine stream
128 segments are of particular interest because of their texture. Due to the texture of the
129 subsoil and anthropogenic encroachments from 1850 on, the riverbeds have become
130 deeper. About 1850, Bavarian Administration systematically started riverbed correc-
131 tions in order to prevent floods, to protect infrastructure like railways and roads and to
132 support agriculture to the fertile plains and meadows supplying the growing popula-
133 tion. In Fig. 2a one can see the beginning of ~~the~~ anthropogenic encroachments ~~due to~~ *impacting on*
134 the ~~results of~~ gauge measurements, starting around 1860. The gauges' neutral points *river flow*
135 have not been changed, staff gauges have been prolonged into the negative measure- *through*
136 ment range. These circumstances affect hydrological interpretations concerning the

137 EIP. To assure homogeneity in ~~the~~ comparison of flood events of individual time se-
 138 ries, a high-pass-filter has been applied (see methods). The regional distribution of
 139 annual precipitation ~~differs~~ ^{varies} from around 600mm/a in the region of the Danube river to
 140 2500mm/a and more within the high~~est~~ elevations ~~of the investigation area~~. The share
 141 of alpine catchment is important for the runoff of the headwaters into the Bavarian
 142 Foreland due to its function as ^{temporary} water storage reservoir and orographic bar-
 143 rier. In Table 1 the reference data of the relevant rivers are listed.

144
 145 **Table 1.** Reference data of the Bavarian Foreland rivers, modified after different authors. The
 146 information "Medium Discharge Summer" refers to the lowest official downstream
 147 gauge

River	Headwaters	Rivers Length	Overground Catchment Area	Medium Discharge Summer
Iller	Allgäu High-alps (Germany)	~ 160 km	2215 km ²	79,3 m ³ /s (Gauge Wiblingen)
Wertach	Allgäu Alps (Germany)	~ 135 km	1290 km ²	16,7 m ³ /s (Gauge Türkheim)
Lech	Rhaetic Alps (Austria)	~ 250 km	4162 km ²	136 m ³ /s (Gauge Augsburg)
Isar	Karwendel Mountains (Austria & Germany)	~ 260 km	8960 km ²	191 m ³ /s (Gauge Plattling)
Inn	Maloja Saddle (Switzerland)	~ 520 km	26100 km ²	972 m ³ /s (Gauge Passau)
Salzach	Kitzbühl Alps (Austria)	~ 220 km	6700 km ²	332 m ³ /s (Gauge Burghausen)

148
 149 In the Bavarian Foreland we can differentiate spatial and chronological aspects of
 150 flood genesis as a function of hypsometric distribution and thereto linked the snow
 151 retention into the individual catchments. We can observe a west-east gradient con-
 152 trasting the chronological annual mean discharge maximum starting in the western
 153 part with the investigation area with a spring peak (rivers Iller and Wertach). The
 154 catchment areas of the rivers Lech and Isar, located in the central part of the investiga-
 155 tion area, are denoted by a distinctly marked summer peak, followed by the eastern
 156 part with prolonged summer peaks (rivers Inn and Salzach). This spatial distribution is
 157 reflected by the seasonal distribution of the flood events indicated in Table 2.

158
 159 **Table 2.** Seasonal distribution of flood events (percentages refer to the individual outer-alpine
 160 river sections) in the Bavarian Foreland for the period of 14th century – 2008

	Iller	Wertach	Lech	Isar	Inn	Salzach
Summer	42%	39%	57%	58%	79%	73%
Spring	16%	20%	14%	14%	10%	6%
Winter	28%	23%	12%	12%	2%	7%
Autumn	14%	18%	17%	16%	9%	14%

161

Be consistent in formatting of tables.

162 Due to the spatial distribution of the catchment areas within the Northern Limestone
163 Alps and the Bavarian Foreland, the highest proportion of summer floods occur in
164 the eastern river sections of Inn (79%) and Salzach (73%), followed by Isar (58%) and
165 Lech (57%). Due to the lower extent of the alpine catchment area, the rivers Iller and
166 Wertach only have about 40% of ~~the~~ floods during summer. In total, a dominance of
167 summer floods can be stated for the annual time series. Hereafter the use of annual
168 information is owed to minimized data quality and information before the 16th centu-
169 ry. The flood-rich periods of the whole year (depicted in Fig. 6 by blue graph) and of
170 the summer months (depicted in Fig. 8 by blue graph) correspond substantially to
171 each other.

172
173 The investigation period of 1300 – 2008 covers the Little Ice Age (1300 – 1850) and
174 the transition period to the Modern Climate Optimum as it is today. According to
175 Wanner et al. (2000), it is advisable to differentiate the Little Ice Age into so-called
176 Little Ice Age Type Events (LIATEs) addressing three major periods of extended
177 glacier tongues. The periods, starting with LIATE3, are depicted for the years 1300 –
178 1380, 1570 – 1640 and 1810 – 1850.

179 180 **3 Database**

181
182 The body of historical source material corresponds to the wide range of settlements
183 along the northern alpine mountain rivers. All of the above-mentioned rivers host at
184 least one notable historical site (among a multitude of other sites, e.g. Kempten (Iller),
185 Augsburg (Lech/Wertach), Munich (Isar), Wasserburg (Inn), Burghausen (Salzach).
186 Within the framework of a DFG- funded (German Research Foundation) research pro-
187 ject a database called IBT (Inundationes Bavariae Thesaurus) (cf. Böhm 2011) was
188 developed in cooperation with HISKLID (Historische Klimadatenbank Deutschland -
189 historical climatic database of Germany - cf. Glaser 2008). The former HISKLID has
190 meanwhile migrated to tambora.org. Tambora is the acronym for the climate and en-
191 vironmental history collaborative research environment.

192 The IBT itself contains more than 32.000 flood events within (Central-) Europe, all of
193 them with a temporal relationship to the 584 independent flood events identified for
194 the Bavarian Foreland (see below). The first investigation period was the period of
195 documentary evidences from the 14th century to the year 1880. The data set of the pe-
196 riod of written evidences includes more than 15.000 flood events. The evaluated writ-
197 ten evidences originated from manuscripts and chronicles (e.g. the “Historische
198 Kommission der Bayerischen Akademie der Wissenschaften/Historical Commission
199 of the Bavarian Academy of Sciences” has published 37 volumes of city chronicles
200 between 1862 and 1968), annals, historical dailies, of the investigated area (cf. e.g.
201 Augsburger Postzeitung 1833-1935, Innsbrucker Nachrichten 1854-1945), compila-
202 tions (e.g. Sonklar 1883, Weikinn 1958 – 1963, Alexandré 1987, Fliri 1998, Stahleder
203 1995 – 2005, Börngen & Tetzlaff 2000 – 2002, Brázdil 2005), un-edited historical
204 leaflet database (Schorn † 1937, Ferdinandum Innsbruck Administration of Inher-
205 itance) and already existing databases (Mitzner 1998, Glaser 2008) which were re-
206 examined with a focus on the Bavarian Foreland. Due to the approach explained be-
207 low ~~every~~ written evidence of the middle reaches and tail waters has been considered.

208

diary's

209 To highlight some selected sources the 'Chroniken deutscher Städte' (Chronicles of
210 German Cities 1862 – 1968) focused to the city of Augsburg upon ^{the} river Lech and the
211 publications of Stahleder (1995 – 2005) for the city of Munich upon river Isar will be
212 introduced briefly. ~~Within 'Chroniken deutscher Städte' the chronicles of city of~~
213 ~~Augsburg must be highlighted especially.~~ Inside the ^{superior} 'Chroniken deutscher
214 Städte' the history of Augsburg is organized into 'Die Chroniken der schwäbischen
215 Städte' (The chronicles of Swabian Cities). In total nine volumes are existent for the
216 second oldest city of Germany, including substantial information about ^{the} river Lech
217 floods. Within these nine volumes the following chronicles have been edited: Volume
218 1 (1865) contains the 'Augsburger Anonyme Chronik' from 1368 – 1406 with pro-
219 ceedings until 1447, the chronicle by Erhard Wahraus from 1126 – 1445 with supple-
220 ments until 1462 and the chronicle from foundation of the city of Augsburg until
221 1469. Volume 2 (1866) contains the chronicle by Burkard Zink from 1368 – 1468.
222 Volume 3 (1892) contains the chronicle by Hector Müllich 1348 – 1487 and the anon-
223 ymous chronicle from 991 – 1483. Volume 4 (1894) includes the chronicle from ^{the} old-
224 est time of the city until 1536 plus proceedings of the chronicle by Hector Müllich.
225 Volume 5 (1896) contains 'Cronica newer geschichten' by Wilhelm Rem 1512 – 1527,
226 Johannes Franks 'Augsburger Annalen' from 1430 until 1462 and supplements con-
227 cerning the chronicle by Clemens Sender. Volume 6 (1906) contains the chronicle of
228 Georg Preu from 1512 until 1537. Volume 7 (1917) contains two chronicles by appar-
229 itor Paul Hektor Mair. Volume 8 contains 'The Diary of Paul Hektor Mair' from 1560
230 – 1563 and the second chronicle by Paul Hektor Mair 1547 – 1565. ~~And~~ Volume 9
231 contains the weaver chronicle by Clemens Jäger from 955 – 1545.

232 Helmuth Stahleder, ex-alternate director of the 'Stadtarchiv München' (city archive
233 Munich) evaluated data within the city archive of Munich to compensate the missing
234 of history of Munich within 'Chroniken deutscher Städte'. Foundations of this compi-
235 lation among others are original documents, calculations of city treasurer and year-
236 books. ^{The} Result of the longstanding investigation was the 'Chronik der Stadt München',
237 ~~is~~ three volumes concerning the history of Munich between the years 1157 – 1818. A
238 multitude of flood events along ^{the} river Isar are recorded within 'Chronik der Stadt
239 München'. Each record is furnished with a related city archive reference. ^{the}

241 The early instrumental records in the Bavarian Foreland started in 1826. More than 20
242 historical gauge station records were examined. The data set of EIP/MIP includes
243 about 17.000 flood events of the Bavarian Foreland (MIP = modern instrumental pe-
244 riod). The data were analysed with respect to monthly maxima of the water level.
245 Taking physical structures and vertical erosion of gauge stations into account, the
246 gauge datum has been changed in some cases, sometimes even repeatedly. A high-
247 pass filter was applied to homogenize the data. In the present paper we choose one
248 representative gauge station for each river, each with the longest coherent time series
249 since 1826. They include the time series of Kempten (Iller), Landsberg am Lech
250 (Lech), Landshut (Isar), Wasserburg (Inn) and Burghausen (Salzach). The EIP time
251 series of the river Wertach was merged between overlapping periods, verified by a
252 high Pearson correlation coefficient ($r=0,86$) between the data from the gauge stations
253 Ettringen and Augsburg/Oberhausen. From 1900 onward data were available from the
254 Bavarian Water Authority.

255 Within the IBT all data sets are organized by the following parameters: identification
256 number, event date as accurate as possible (most records are available in daily and

257 monthly resolution), duration of flood, rainfall etc., location with geographic coordi-
258 nates, river relationship, reference and coding concerning hydrological and climato-
259 logical parameters and source text. All data of the Bavarian Foreland have been rec-
260 orded and coded for tambora.org. The activation of the elaborated data base should
261 soon be realized.

262

263 To merge the different data periods, we used the following approach: To compare
264 flood events from written historical sources with flood events measured by water level
265 or discharge during the instrumental period, we used an existing intensity classifica-
266 tion of historical floods which was adapted to the instrumental period. According to
267 the scheme of Sturm et al. (2001), the flood events were classified into three intensity
268 levels. The classification is based on damage reports and descriptions of weather con-
269 ditions if available. If flood events were mentioned in rudimentary descriptions only
270 or there was ~~or~~ little ~~or~~ minor damage, the event was classified as a regional flood (in-
271 tensity Level 1). If damage of water-related structures (e.g. bridges, weirs and mills)
272 or buildings near the rivers have been reported or if there were indicators for long-
273 lasting flooding of farmland, Level 2 was assigned to the flood. The criteria for cata-
274 strophic floods, reported from different river systems, are severe damage or destruc-
275 tion of water-related structures, loss of lives, long-lasting flooding of wide areas and
276 geomorphological changes in the fluvial system; those were classified as Level 3. The
277 selected instrumental data are based on the monthly maxima ~~of~~ water level or dis-
278 charge. The mean values of these measurements plus one, two or three standard devia-
279 tions define the thresholds for the classification of the instrumental data (in case of
280 water level data high-pass filtered data were used, see methods). According to the ex-
281 perience in working with historical flood information, all floods of intensity Levels
282 1,2 and 3 from the period of documentary evidence were considered, whereas Level 1
283 events from the instrumental period were disregarded. An overlapping period (1826 –
284 1880) between the descriptive and the instrumental periods suggested this procedure.

Supports

285 Samples of rudimentary descriptive flood information have shown that the historical
286 flood information through time traditionally is based on strong events for the most
287 part (cf. Böhm 2011). In Table 3 all flood events used for the merged time series
288 “flood frequencies of the Bavarian Foreland” (cf. figs. 3 and 4) are listed. The data for
289 the time series Bavarian Foreland is derived from 1825 different flood records in total
290 which can be assigned to 584 independent flood events.

291

292

293

294

295

296

297

298

299

300

301

302

303

304
305
306
307
308

Table 3. Synoptic table of data basis 'flood frequencies Bavarian Foreland'. Columns a to c contain all outer-alpine flood events of documentary evidences until 1880, segmented after intensity levels. Columns d to e contain all floods derived from instrumental periods until 2008 for one representative gauge per river. EIP = Early Instrumental Period, MIP = Modern Instrumental Period.

River	a) Level 1	b) Level 2	c) Level 3	d) EIP/MIP level 2	e) EIP/MIP level 3
Iller	32	53	15	45	13
Wertach	37	79	20	66	16
Lech	101	159	80	78	38
Isar	88	101	29	55	18
Salzach	154	113	78	56	22
Inn	79	82	63	48	7

309
310

To expand the ^{the} data basis as wide as possible we have applied a methodical practice we have named "Non Critical Approach" (NCA) (cf. Böhm 2011). The NCA is a procedure especially designed for extraordinary hydrological events. Within the range of historical climatic data, flood information has an exceptional position. Common threads connecting available flood information are damages, ~~which have led to burdens on former neighbors~~. The main argument for the NCA is based on the reasonable assumption that historical flood reports - due to the particular burden~~s~~ - contain more objective information than other descriptions of climatic events. In the center of this approach stands the tradition of the gist of 'flood event' through time. Starting point of this approach was the fundamental question if anonymous sources in general may be regarded as verified sources (cf. e.g. Augsburgischer Anonyme Chronik von 1368 bis 1406. In: Die Chroniken der schwäbischen Städte. Augsburg, Band 1. Leipzig 1865). According to a rigorous ^{from} interpretation of source criticism all of the (environmental-related) information ~~of~~ this source would have to be discarded. Based on the NCA we use all available sources and information about flood events of the outer-alpine river sections concerning the period of documentary evidences. Avoiding classical source criticism the NCA contributes to increase acquisition of information and reduces the thinning of relevant information during times of limited flood documentation. This approach minimizes the loss of original written records concerning historical flood information due to anthropogenic or natural calamities.

? I do not understand.

331

To verify the NCA various stress tests have been trialed. Glaser et al. (2002) state that a spatial criterion for the distribution pattern of weather-climatic causes can be implied by sufficient data density. Within the scope of the NCA a spatiotemporal/synoptical criterion has been consulted to verify historical data. All superior flood events of the Bavarian Foreland have been visualized by spatiotemporal flood distribution pattern with the assistance of geographic information systems. Therefore plausible (spatiotemporal/synoptical) evidence for the validity of flood information can be adduced. Further confirmation was given by cross-comparison with verified records, e.g. HISKLID (cf. Böhm 2011).

341
342
343
344

Environmental psychological aspects provide ~~✓~~ further backing for the NCA. In brief damaging flood events have an exceptional position in cultural history and the transfer of information through time based on primal fear ~~still contains the gist~~. A more detailed description of the NCA is to be found in Böhm (2011). ^{remains}


345 **4 Methods**

346

347 In Fig. 2 the monthly maxima of water levels, exemplified by the gauge ^{at} Ettringen/Wertach (35km south of Augsburg) before (a) and after (b) high-pass filtering are depicted. Fig. 2a shows the deepening of the riverbed due to anthropogenic encroachment into the river system. The process of riverbed deepening started around 1860. Since the administration didn't change the datum of the measuring device for a time, but extended the measuring sticks of the gauges into the negative range instead, a total riverbed incision of more than 6 m within 30 years is documented. Furthermore, first counter-measures (like lateral water-buildings) obviously occurred around 1870. Around 1885 the incision seems to have stopped. After a short data gap around 1895, two datum changes can be identified. The high-pass filtering seems to be the most suitable method to address these different changes.


358

avoid contractions
did not



359

360



361

362 **Figure 2a,b.** Monthly maxima of water level (in cm) 1826 – 1937 at the gauge Ettringen/Wertach (35km south of Augsburg). a) Original data from official records, b)
363
364 high-pass filtered data
365

366 In order to be able to properly understand the long-term development of flood events
367 in the Bavarian Foreland, z-transformed 31-year running flood frequencies have been
368 calculated in several studies (e. g. Glaser 2008, Glaser & Stangel 2003b, Böhm &
369 Wetzel 2006, Sturm et al. 2001, Schmoecker-Fackel & Naef 2010). The 31-year time
370 step is derived from the standard reference period of the World Meteorological Or-
371 ganization. This time segment is an established tool to identify the linkage of climatic
372 coherence of time series and exhibits significant changes in flood frequencies. Alt-
373 hough this measure results in a comparatively poor filtration effect it still meets the
374 needs of various geoscientific approaches to define climatological phases
375 (Schönwiese 1992).

376 Due to their geomorphological shapes the catchment areas of the investigated rivers
377 have been divided in an inner- and outer-alpine region. Only the outer-alpine regions
378 (see Fig. 1 dashed line) have been considered for the present paper. To reveal the
379 flood sensitivity of the entire Bavarian Foreland, all flood events of the outer alpine
380 region have been merged into one overall time series. The highest classification ac-
381 cording to damage reports has been counted whereas local events caused by i.e. flash-
382 floods have not been counted. Hence only mesoscale hydrological events have been
383 incorporated into the present analysis.

384 In order to integrate different historical sources and locations referring to one meteor-
385 ological event, a time-window including a maximum number of seven days before
386 and after a designated event has been introduced. Therewith the varying duration of
387 hydro-meteorological sequences (from the genesis of synoptic disturbances to the
388 termination of flood waves) as well as the blur of historical information can be con-
389 sidered. For example, the summer flood of the year 1501, being one of the biggest
390 floods since the beginning of written records (quantitative evidence in terms of a flood
391 mark is located at the “Fischmarkt” in Passau, Bavaria), has been recorded more than
392 150 times in the IBT, yet it is counted only once in the 31-year running flood frequen-
393 cy of the Bavarian Foreland.

394 The determination of flood-rich and flood-poor periods are based on a polynomial
395 function of the 5th degree for the running flood frequencies (see black graph in Fig.
396 4). This method has been adapted from Glaser et al. (2004) who used polynomial
397 functions to visualize long-term development of climatic elements. Using this func-
398 tion the inhomogeneity of the number of cases could best be confronted. Different da-
399 tabases and data densities (e. g. 14th/15th century - turn of the 15th to 16th century - be-
400 ginning of the instrumental period) were thus considered as far as possible. This
401 method does not claim precision for the beginning and the end of the defined periods,
402 but compared to a multitude of other methods and due to the changing data density
403 over time it is the highest-performance method. Different methodical approaches with
404 the aid of quantiles as medians or percentiles could not achieve satisfactory defini-
405 tions for the generated time series and its comparability. The determined periods
406 should come over as the results of a sensitivity analysis. The fixing of the threshold
407 based on a polynomial function of the fifth degree coincides with the fracture points
408 of the t-test analyses (cf. Fig. 5), so the method is provided by statistical measure.

409 In general, the intervals above/below the function graph are defined as flood-
410 rich/flood-poor periods. Due to the changing data basis over the entire time series, it
411 was necessary to interpolate in some cases (compare e.g. flood-rich period #2). With
412 respect to the differing data density as a function of time and not as an increasing fre-
413 quency in general the values in Fig.4 have been z-transformed. So the weak data den-
414 sity until the beginning of the 16th century is denoted by values beneath the zero line.
415 Therewith the under- and over-representative availability of data has been taken into
416 account.

417 For merging the single outer-alpine time series all flood events which took place
418 isochronally (under consideration of the above mentioned time window) have been
419 counted only once. Virtually all counted events can be affirmed with a plurality of
420 flood events which occurred isochronally within in the investigation area as well as in
421 a (Central-) European context.

422

423 **4.1 Statistical significance**

424

425 The time series for the Bavarian Foreland has been submitted to a two-sided t-test
426 which can identify fracture points within the time series (cf. Glaser & Stangl 2003b).
427 The fracture points reveal differences between the means of sliding flood frequencies.
428 The differences shown by estimators above the threshold are expected to detect signif-
429 icant coherence between superior framework conditions like variations of large-scale
430 atmospheric circulation and consequential variability of flood-poor and -rich periods.
431 The α -level of 0.05 (solid line) is depicted in Fig. 5. Most of the detected fracture
432 points coincide with changes in atmospheric conditions (see next section). The flood
433 frequency time series of the Bavarian Foreland has additionally been correlated with
434 reconstructed NAO index-values (cf. Luterbacher et al. 2002a) as well as with moving
435 mean sunspot numbers (data after Hoyt & Schatten 1997). Significant correlations
436 between these quantities include the consideration of inherent autocorrelations (cf.
437 Werner 2002).

438


439

440 **5 Results**

441

442 In Fig. 3 all single time series of the examined catchment areas are depicted. Taking
443 comparability into account all axes have the same scale of values. The itemized time
444 series of the rivers Iller (A), Wertach (B), Lech (C), Isar (D), Salzach (E) and Inn (F)
445 are the foundation of the overall time series Bavarian Foreland depicted in Fig. 4.
446


447



448

449

450



451

452

453

454

455

456

457

458

459

460

461

462

463

Figure 3. 31-year running flood frequencies of the Bavarian Foreland Rivers (red graph). Right ordinate: dark columns show the annual flood frequencies. A Iller, B Wertach, C Lech, D Isar, E Salzach and F Inn.


In Fig. 4 the merged flood history of the Bavarian Foreland between 1300 AD and 2010 is depicted. The data for that time series is derived from 1825 different flood records of the itemized time series (c.f. Fig. 3) which are assigned to 584 independent flood events. Based on a polynomial function of the 5th degree (see red line), 9 flood-rich and 8 flood-poor periods can be identified. ~~The mentioned~~ in the methods chapter, due to changing data density over time it is the highest-performance method and the determined periods constitute the results of a sensitivity analysis. Particularly the

As identified

464 rising data density after the mid-15th century must be seen in a context of the inven-
465 tion of letterpress, among other social reasons and changes. From 1826 onward meas-
466 ured data ~~in~~ daily resolution are available.

467 ~~at a~~
468 Single flood-rich periods will be discussed below. In order to prove significant chang-
469 es within the time series of the 31-year running flood-frequencies of the Bavarian
470 Foreland, we have depicted the estimators of the t-test in Fig. 5 (cf. Glaser & Stangl
471 2003b). In this context, changing climatic parameters like the NAO index and the sun-
472 spot numbers will be briefly approached (a more extensive discussion will follow sub-
473 sequently).

474



475

476 **Figure 4.** 31-year running z-transformed flood frequencies of the Bavarian Foreland. Grey
477 bars label flood-rich periods #1 to #9, black graph: polynomial of 5th degree.

478

479 **5.1 Flood-rich period #1: 1300 – 1335**

480 The first flood-rich period, although based on ~~low~~ low data density, can be associated
481 with changes in the atmospheric framework. Within the investigation area at least 16
482 records could be extracted from compilations like Alexandre (1987), different chroni-
483 cles (cf. e.g. Zillner 1885, Schnurrer 1823) and the Augsburger Urkundenbuch Nr.
484 264 (cf. Gross 1967). Despite small data density significant changes of climatic pa-
485 rameters at the beginning and the end of this flood sensitive period can be stated and
486 should not be withheld.


487 Wanner et al. (2000) ~~are~~ ^e dating the onset of the Little Ice Age at the beginning of the
488 14th century, based on Miller et al. (2012), triggered by multiple volcanic eruptions.
489 Period #1 coincides with the beginning of LIATE3, ~~the~~ ^{the} first period of advancing glacier
490 tongues during the Little Ice Age, additionally enhanced by the Wolf solar activity
491 minimum (1282 – 1342, cf. Glaser 2008). Following the sunspot numbers after
492 Usoskin et al. (2004), the first flood-rich period ~~minutely~~ coincides with a period of
493 sunspot minima. According to Fig. 5, the t-test value shows significant changes with
494 the beginning and the end of Period #1. The estimator declines during the flood-active

495
496
497
498
499
500
501
502
503
504
505
506
507
508

identified flood-rich phases
phases (this can be verified for most of them). While LIATE3 approaches a climax, the flood frequencies decline considering a time lag due to mass input into the alpine glaciers (cf. Wanner et al. 2000). Most of the significant fracture points in Fig. 5 coincide with the beginning and the end of the flood-rich or flood-poor periods in Fig. 4. A further qualitative confirmation for particular climatic circumstances during that period is provided by Lamb (1980).

5.2 Flood-rich period # 2: 1370 – 1450

The increase of flood frequencies is accompanied by significant t-test estimators. Despite the absence of significant estimators, the end of period #2 coincides once more with changing climate conditions characterized by the beginning of the Spoerer Minimum, another sunspot minimum between the years 1450 and 1534 (cf. Glaser 2008).




509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531

Figure 5. Differences of sliding means by 31-year running t-test estimator of flood frequencies of the Bavarian Foreland, threshold value for the two-sided t-test is 2.00 (see red line). Grey bars label flood-rich periods #1 to #9.

5.3 Flood-rich period # 3: 1470 – 1525

The transitional period between the flood-poor period #2 and the flood-rich period #3 coincides with an obviously rising estimator (significant at the α -level 0.05) as well as with the end of a distinct period of negative temperature anomalies in the Alps (cf. Wanner et al. 2000). The end of period #3 is once again marked by highly significant t-test estimator values. The maximum value is accompanied by the end of the Spoerer Minimum, again an indication for changing climate conditions which affect the flood frequencies in the Bavarian Foreland.

From 1500 onward we can use reconstructed NAO-Index (NAOI) values (cf. Luterbacher et al. 2002). The end of period #3 (= first grey bar in Fig. 6) is accompanied by an obvious declining NAOI for annual (Fig. 6) as well as summer seasonal values (Fig. 8). Due to the impact of precipitation to the entire hydrological year on flood progress, the full-year development of the NAO may reflect mean weather conditions. In general, however, the correlation between weather conditions and the NAOI is more significant during winter than during the warmer half of the year.



532
 533 **Figure 6.** Z-transformed time series of annual 31-year running flood frequencies in the Bavar-
 534 ian Foreland (blue) and of annual 31-year running NAOI (red) (NAOI data after
 535 Luterbacher et al. 2002). Grey bars label the flood-rich periods # 3 to # 9 based on
 536 the full-year development.

537
 538 **5.4 Flood-rich period # 4: 1555 – 1590**

539 Between the years 1550 and 1700 the z-transformed NAOI-values are generally char-
 540 acterized by low index-values. Within this period, variations in flood frequencies can
 541 be identified (see Fig. 6) with increasing and declining values ~~being~~ accompanied by
 542 rising and falling NAOI-values, respectively, for the whole year as well as for the sea-
 543 sons, ~~the~~ winter (not shown) and summer (cf. Fig. 8). The t-test estimator indicates sig-
 544 nificant values (> 2.00 at a α -level of 0.05) at the beginning and at the end of period
 545 #4 (cf. Fig. 5). The flood frequency maximum coincides with a distinct period of neg-
 546 ative temperature anomalies that lead into LIATE2.

547
 548 **5.5 Flood-rich period # 5: 1615 – 1665**

549 Period #5 is again accompanied by significant t-test estimators for the beginning as
 550 well as for the end (see Fig. 5). The transition period between period #5 and the
 551 Maunder Minimum (1645 – 1715, cf. Schönwiese 2008) is accompanied by a con-
 552 spicuous behavior of the t-test estimator, a possible evidence for unsettled atmospher-
 553 ic conditions. Period #5 took place during LIATE 2 (1570 – 1640, cf. Wanner et al.
 554 2000); the retreat of the glacier tongues (peak around 1650) coincides with declining
 555 flood frequencies. Period #5 also fell into a period of declining sunspot numbers
 556 which have been directly observed for the first time since 1610 (cf. Fig 7). The abso-
 557 lute minima of sunspots in 1660 turned into the Late Maunder Minimum and they
 558 were accompanied by the absolute flood frequency minimum from 1500 till today.
 559 With the beginning of immediate observation of sunspots we can refer to a distinct
 560 correlation between the 31-year moving sunspot numbers and flood-rich and flood-

561 poor periods until 1930. The end of LIATE 2 coincides with the beginning of the
562 Maunder Minimum, compare Fig. 7 and Wanner et al. (2000).

563
564 The flood frequencies are rising while the mean NAOI-values for the complete year
565 are declining; the flood frequency peak coincides with minimal NAOI-values. The
566 following decline of the flood frequencies is accompanied by rising NAOI-values (see
567 Figs. 6 and 8). The development of the winter NAOI reveals a different pattern. The
568 absolute maximum value of flood frequencies coincides with a short-termed increase
569 of the winter NAOI. This could indicate wet winter conditions resulting in vast water
570 retention which could favor summer floods during snow melt.

571 572 **5.6 Flood-rich period # 6: 1730 – 1780**

573 Period #6 is once again marked by significant t-test estimators during increasing flood
574 frequencies; the end of the period is marked by a distinct evolution of the estimator
575 values. The t-test estimators in general will not show significant variations for the
576 next 80 years (cf. Fig. 5). Period #6 is accompanied by a noticeable development of
577 the NAOI-values. The year-round development shows a parallel increase of flood fre-
578 quencies and NAOI-values, but during the flood frequencies' peak the NAOI-values
579 decline in a distinct way (cf. Fig. 6). The development of the Little Ice Age in terms
580 of glacier tongue movements shows a slight stagnation.

581
582 The following flood-poor years until 1820 are characterized by a next sunspot mini-
583 mum, the Dalton Minimum (1790 – 1830, cf. Schönwiese 2008), again accompanied
584 by relatively low flood frequencies (see Figs. 3 and 4). The end of the Dalton Mini-
585 mum and the beginning of the next flood-rich period is accompanied by extreme flood
586 frequencies (cf. Fig. 4) as well as extraordinary t-test estimators (cf. Fig. 5). Reasons
587 for that will be explained below.

588 589 **5.7 Flood-rich period # 7: 1820 – 1870**

590 Period #7 is marked by a changing database (transition between documentary period
591 and early instrumental period). In addition, it represents the beginning of the transition
592 period between the end of the Little Ice Age and the beginning of the Modern Climat-
593 ic Optimum as well as the beginning of (systematic) anthropogenic interference into
594 the natural river systems. Thus, period #7 falls into a section of different overlapping
595 trends. The t-test estimators reveal this transition at the beginning of the period
596 (unique t-test estimator values, see Fig. 5). The end of the period, based on t-test esti-
597 mators, can be interpreted as inhomogeneous climatic conditions during the transition
598 to the Modern Climatic Optimum. The increasing flood-frequencies of period #7 co-
599 incide with increasing sunspot numbers (cf. Fig. 7).

600
601 The development between LIATE1 (1810 – 1850) and period #7 requires a particular
602 reflection, as LIATE1 and the flood-rich period end at about the same time, whereas
603 the preceding LIATEs (# 3 and #2) did not end at the same time like the correspond-
604 ing flood-rich periods. The maxima of the preceding LIATEs fall into intervals of low
605 flood frequencies framed by the end and the beginning of flood-rich periods. A no-
606 ticeably aligned development between the NAOI-values and the flood frequencies can
607 be highlighted for the whole year as well as for the summer and winter seasons (cf.
608 Figs. 6 and 8). The following years are characterized by both declining flood frequen-

609 cies and NAOI values. In this context we can generally identify an increasing impact
610 and significance of the summer NAOI for the development of flood frequencies in the
611 Bavarian Foreland (discussion see below).

612

613 **5.8 Flood-rich period # 8: 1910 – 1955**

614 Period # 8 is once more framed by significant t-test estimators with a final significant
615 estimator-value within the time series (see Fig. 5). Until 1930 similarities (or correla-
616 tions) between flood frequencies and sunspot numbers do exist. After that the time
617 series seem to be decoupled, as depicted in Fig. 7. The beginning of period #8 coin-
618 cides with rising NAOI-values, the amplitude of these values in general shows a par-
619 allel development (cf. Figs. 6 and 8).

620

621 **5.9 Flood-rich period # 9: 1980 – ?**

622 A last flood-rich period starts in 1980 intersecting with the end of the time series. Due
623 to the use of 31-year running quantities, its validity remains unclear.

624

625

626 **5.10 The flood frequencies of the Bavarian Foreland accompanied by different**
627 **climatic conditions in detail**

628

629 **5.10.1 The correlation of flood frequencies and sunspot numbers**

630 The meaning of varying solar activity for the climatic conditions is currently a matter
631 of fierce discussion (cf. Shindell et al. 2001, Feulner & Rahmstorf 2011, Feulner
632 2011). Generally the variability of solar activity can affect ~~the~~ large-scale atmospheric
633 circulation including climate parameters like temperature, precipitation and transpira-
634 tion (cf. Endlicher & Gerstengarbe 2009). Despite the uncertainty in which way solar
635 activity could have an effect on climate development within the research area, it can
636 be shown that solar trends (of different signs) coincide with changing climatic condi-
637 tions affecting the flood frequencies in the Bavarian Foreland. Within these relation-
638 ships, one certain conspicuousness must be emphasized, but not discussed in detail at
639 this point. One has to distinguish between direct observations or reconstructions based
640 on proxy data. Both periods differ with respect to the sign of the correlation between
641 flood frequencies and solar activity. In general, before direct observations started,
642 high flood frequencies coincided with reduced solar activity, while then flood-rich
643 periods coincided with increased solar activity. The best fit between the 31-year run-
644 ning frequencies is achieved with direct observations starting in 1610 and ending
645 around 1930 (cf. Fig. 7). The year 1930 occupies a special position which will be dis-
646 cussed later ~~in~~. Considering the four sun-spot minima encompassed within the inves-
647 tigation period, we can denote the following:

648

649 The Wolf Minimum (~ 1280 – 1340) coincides with the first calculated flood-rich pe-
650 riod. This temporal correspondence between high flood activity and low sunspot
651 numbers is unique within the total time series (cf. Fig. 4).


652

653 The duration of the Spoerer Minimum differs among a number of authors. According
654 to Glaser (2008), the Spoerer Minimum takes place between the years 1450 and 1534,
655 whereas Schönwiese (2008) identifies the years 1400 – 1510. According to the first
656 definition of the Spoerer Minimum, another sunspot minimum coincides with a flood-
657 rich period (here period #3). Keeping with the sunspot data of Usoskin et al. (2004),
658 the flood frequency maximum of period #3 also corresponds to the low sunspot num-
659 bers of the Spoerer Minimum.

660

661

662



663
664 **Figure 7.** Z-transformed time series of annual 31-year running flood frequencies in the Bavarian
665 Foreland (blue) and “sunspot activity” (red) since 1610 (Data for sunspot activity
666 modified after Hoyt & Schatten 1997). Grey bars label the flood-rich periods # 5 to #
667 9 based on the full-year development.
668

669 With the beginning of the Maunder Minimum period the indication of the relationship
670 changes. As depicted in Fig. 7 running flood frequencies and sunspot numbers are
671 varying in a similar way. The course of both graphs during the Dalton Minimum
672 (1790 –1830) again shows a similar development. These data indicate a significant
673 trend, between the years 1610 and 1995 the Pearson correlation coefficient amounts to
674 0.62, between 1700 and 1930 to 0.7. Autocorrelation could be achieved by using the
675 effective sample size based on a method by Werner (2002), which regards the statisti-
676 cal persistence of database-inherent autocorrelations ($\alpha = 0.01$). After 1930, the rela-
677 tionship of sunspot numbers and flood frequencies seems to be decoupled. A compari-
678 son of global temperatures and GAR (Greater Alpine Region) temperatures shows a
679 temperature leap over the calculated neutral-point temperature development since the
680 beginning of the early instrumental period (cf. Auer et al. 2007).
681


682 Not all flood-rich and flood-poor periods can be connected to sunspot minima or max-
683 ima, but all sunspot extremes can be connected to changes of overall climatic condi-
684 tions irrespective of its trends (cf. Böhm 2011)
685

687 5.10.2 Correlations of flood frequencies and NAOI-values in detail

688
689 Is it possible to associate (in a statistical sense) the variability of the NAO and the
690 flood history of the Bavarian Foreland? Due to oscillations of barometric pressure be-
691 tween the ^{ice} Iceland cyclone and Azores anticyclone weather conditions of the
692 ~~investigated area can be affected in various manners~~ ^{Atlantic can affect W. European climate in various ways}. Based on the reconstruction of
693 the NAOI by Luterbacher et al. (2002), the time series of the flood frequencies (31-
694 year running mean values) and of the NAOI (likewise 31-year running mean values)
695 can be compared from 1500 AD onward (cf. Figs. 6 and 8). The NAO is one of the
696 dominating teleconnection patterns regulating the regional characteristics of many

697
698
699
700
701

climatic parameters. The NAO also reveals seasonal variations (in context of the climatic seasonal cycle).



702
703
704
705
706
707

Figure 8. Z-transformed time series of 31-year running flood frequencies in the Bavarian Foreland (blue) and NAOI values (red) for the meteorological summer (JJA) (NAOI data after Luterbacher et al. 2002). Grey bars label the flood-rich periods # 3 to # 9.

708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723

The annual and the summer time series of flood frequencies and ^{are} of NAOI-values have been compared. The confrontation of the annual valuables ~~is~~ depicted in Fig. 6, the comparison for the summer seasonal in Fig. 8. The particular importance and the general dominance of summer floods in the Bavarian Foreland ~~were~~ ^{have} already ~~accosted~~ ^{been discussed}. The winter flood occurrences are mainly relevant in terms of the retention potential of the alpine catchment areas which supports the development of floods in general. In spite of the general decline of the NAO importance during summer, significant correlations of the NAO (-index) and the time series of summer floods can be recognized. In a first step 100-year intervals have been considered. For the centuries 1500 – 1599 ($r = 0.78$, $\alpha = 0.01$) and 1900 – 1999 ($r = 0.65$, $\alpha = 0.01$) significant ^{positive?} correlations occur (once again verified by the calculated persistence). Shifting the time-interval by 50 years, a significant correlation can be observed for the period 1650 – 1749 ($r = 0.8$). Furthermore, the years from 1830 to 1999 exhibit another highly significant correlation coefficient of $r = 0.8$ ($\alpha = 0.01$). The changing sign in the relationship around 1820 (cf. Fig. 8) coincides with the important transition period between the Little Ice Age and the Modern Climate Optimum.

724
725
726
727

5.11 The flood frequencies of the Bavarian Foreland in confrontation with selected flood frequencies of Central Europe ^{? toward.}

728
729

This confrontation is limited to the period between 1500 and 1900. The limitations result from weak data density in general before 1500 and in a multitude of anthropo-

730 genic overprints of the river systems around the beginning of the 20th century. The
 731 comparison will be limited to the Lower Rhine and Middle Rhine (cf. Glaser 2008),
 732 the Vlatva (an Elbe tributary) and the Czech Elbe itself (cf. Brazdil 1998). The con-
 733 frontation is depicted in Table 4. Due to the decadal visualization of beginnings and
 734 endings of the marked periods they underlie a certain blur. Similarities for all time se-
 735 ries can be particularly highlighted for the second half of the 16th century. In general
 736 an unexpected similarity can be ~~stranded~~ ^{seen} between the time series of the Bavarian
 737 Foreland and the Lower Rhine, except the years 1790 until 1819. Good accordance
 738 between the Bavarian Foreland can be revealed for the first and seventh and eighth
 739 decade of the 16th century. During the 17th century only the sixth decade shows good
 740 accordance. A further good accordance can be highlighted for the end of the 18th and
 741 beginning of the 19th century. Reasons for this variable behavior are founded in the
 742 variability of general synopsis and resulting weather conditions. In that context the
 743 above mentioned NAO is playing a vital role. For a further understanding of the vari-
 744 ability between the confronted time series meteorological aspects must be consulted.

745
 746 **Table 4.** Confrontation of selected flood frequencies. Lower Rhine (RHI), Middle Rhine
 747 (RHm), Czech Elbe (ELBcz), Vlatava (VLA) and Bavarian Foreland (BF). Due to the decadal
 748 visualization beginnings and endings of the marked periods underlie a certain blur. Data al-
 749 tered according to Glaser (2008) and Brazdil (1998).

	RHI	RHm	BF	ELBcz	VLA	
1500						1500
1510						1510
1520						1520
1530						1530
1540						1540
1550						1550
1560						1560
1570						1570
1580						1580
1590						1590
1600						1600
1610						1610
1620						1620
1630						1630
1640						1640
1650						1650
1660						1660
1670						1670
1680						1680
1690						1690
1700						1700
1710						1710

3f.

1720					1720
1730					1730
1740					1740
1750					1750
1760					1760
1770					1770
1780					1780
1790					1790
1800					1800
1810					1810
1820					1820
1830					1830
1840					1840
1850					1850
1860					1860
1870					1870
1880					1880
1890					1890
1900					1900

752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777

6 Conclusions

The flood history of the Bavarian Foreland ^{can} ~~could~~ be analyzed in a statistical way from the beginning of the 14th century onward (based on the existing data). The flood history of the entire Bavarian Foreland was compiled and analyzed, based on both documentary and instrumental data. We could identify statistical correlations between the flood frequencies and conditions of the atmospheric framework up to ~~today~~ ^{the present day} based on different climate proxies and historical observations and transmissions.

The investigation period commences amid the latest climatic depression of the Subatlantic stage (2.5 ka – present) ~~until today~~. Despite the reduced availability of data at the beginning of the time series, including the absence of reconstructions of pressure fields, temperature and precipitation (until 1499), significant changes in the correlation between climate conditions and flood frequencies can be identified. Virtually each shift in the flood frequencies' trend (towards flood-rich or flood-poor periods) coincides with significant fracture points within the time series (according to two-sided t-tests). These fracture points provide indications of changing atmospheric conditions which may affect ~~the~~ flood development ^{in general}.

The NAO is of particular importance for the climatic conditions in the Bavarian Foreland as well as in the Alps in general. However, according to Casty et al. (2005), the NAO alone cannot explain the very sophisticated climatic events of the Greater Alpine Region (GAR). The present work describes multiple mechanisms with influences to major atmospheric conditions. With a view towards the changing directions of the flood frequencies the shifts themselves within multiple climatic factors are playing an

778 important role in the climatic circumstances of flood development. In this context, the
779 dominant role of the summer NAO from 1830 onward is conspicuous. This atmos-
780 pheric parameter coincides with the beginning of the transition period between Little
781 Ice Age and Modern Climate Optimum. Until the end of the time series, high correla-
782 tion coefficients do exist. For this special period a statistical unique coherence within
783 the present work can be emphasized.

784
785 Another ~~significant~~ influence on flood frequency development might come from so-
786 lar activity. Despite marginal changes of global radiation, the correlation between
787 flood frequencies and sunspot numbers from 1610 until 1930 is rather high. For the
788 Maunder Minimum, the solar radiation was ~~markedly~~ reduced by about 0.24 % (com-
789 pared with the present mean value, cf. Lean & Rind 1998). This causes a cooling ef-
790 fect of 0.5° C at the most for the northern hemisphere. Seemingly, the slight change of
791 solar radiation ~~should~~ ^{could} lead to a significant alteration of atmospheric fluxes, particu-
792 larly concerning the moisture content of air masses. Nevertheless, the high correlation
793 between flood frequency and solar activity cannot explain the mechanism of action
794 from the solar surface through the atmospheric ~~steps~~ ^{system} towards the surface. To explain
795 this mechanism of action further investigations are necessary.

796
797 After ~~the year~~ 1930, the natural relationships seem to be superimposed by an increas-
798 ing anthropogenic influence on the climatic conditions. In this context a decoupling of
799 a retrograde signal could be revealed. That assumption will be indicated by trespass- ^{record}
800 ing the threshold of the average temperature deviation for the GAR in 1930 (cf.
801 ZAMG 2011).

802
803 The current report emphasizes the importance of long time series. The complexity of
804 northern hemispheric (or even global) circulation dynamics as well as the range of
805 natural climate variability is, however, only partially understood.

806
807

808 **References**

809

810 **Alexandré, P. (1987):** *Le Climat au Europe au Moyen age.* – Paris 1987.

811

812 **Auer I., Böhm R., Jurkovic A., Lipa W., Orlik A., Potzmann R., Schöner W., Ungers-**
813 **böck M., Matulla C., Briffa K., Jones P.D., Efthymiadis D., Brunetti M., Nanni**
814 **T., Maugeri M., Mercalli L., Mestre O., Moisselin J.M., Begert M., Müller-**
815 **Westermeier G., Kveton V., Bochnicek O., Stastny P., Lapin M., Szalai S.,**
816 **Szentimrey T., Cegnar T., Dolinar M., Gajic-Capka M., Zaninovic K.,**
817 **Majstorovic Z. & E. Nieplova (2007):** HISTALP – historical instrumental clima-
818 **tological surface time series of the greater Alpine region 1760 – 2003.** – In: *Int J*
819 **Climatol** 27. 17-46.

820

821 **Böhm, O. (2011):** *Flood History of the Bavarian Foreland – The summer floods in*
822 *context to the climatic history of Central Europe.* – Dissertation (language Ger-
823 *man), Universität Augsburg.*

824 <http://opus.bibliothek.uni-augsburg.de/opus4/frontdoor/index/index/docId/2261>

825

826 **Böhm, O. & K.-F. Wetzel (2006):** Flood history of the Danube tributaries Lech and
827 **Isar in the Alpine foreland of Germany.** – In: *Hydrological Sciences*, 51 (5), Oc-
828 **tober 2006, Special Issue: Historical Hydrology.** 784-798.

829

830 **Börngen, M. & G. Tetzlaff [Hrsg.] (2000):** Weikinn, C. – *Quellentexte zur Witte-*
831 *rungsgeschichte Europas von der Zeitenwende bis zum Jahre 1850, Hydro-*
832 *graphie, Bd. 5 (1751 – 1800).*

833

834 **Börngen, M. & G. Tetzlaff [Hrsg.] (2002):** Weikinn, C. – *Quellentexte zur Witte-*
835 *rungsgeschichte Europas von der Zeitenwende bis zum Jahre 1850, Hydro-*
836 *graphie, Bd. 6 (1801 – 1850).*

837

838 **Brázdil, R., Dobrovolný, P., Elleder, L., Kakos, V., Kotyza, O., Květoň, V., Macková,**
839 **J., Müller, M., Štekl, J., Tolasz, R. & H. Valášek (2005):** *Historical and Recent*
840 *Floods in the Czech Republic.* – Masaryk University and Czech Hydrometeorog-
841 *ical Institute. Brno, Prague.*

842

843 **Brázdil, R. (1998):** *The history of floods on the rivers Elbe and Vlatava in Bohemia.* –
844 **Pörtge, K.-H. & M. Deutsch [Hrsg.]:** *Erfurter Geographische Studien - Aktuelle*
845 *und historische Hochwasserereignisse.* 93-108.

846

847 **Casty, C., Wanner, H., Luterbacher, J., Esper, J. & R. Böhm (2005):** *Temperature and*
848 *precipitaion variability in the european Alps since 1500.* – In: *International Jour-*
849 *nal of Climatology* Volume 25, Issue 14. 1855-1880.

850

851 **Deutsch, M. & K.-H. Pörtge.(2001):** *Deutsch, M. & K.-H. Pörtge (2001):* *Histo-ri-cal*
852 *floods in Thuringia on the example of the flood in Langensalza on the 15th June*
853 *1815.* – In: *Hlavinková, P. & J. Munzar [Hrsg.]:* *Nature and so-ciety in regional*
854 *context, 4th Moravian Geographical Conference CONGEO `01, Brno/ Czech*
855 *Republic.* 16-23.

- 856 Deutsch, M. & K.-H. Pörtge (2002): Erfassung und Bewertung historischer Hochwas-
857 serereignisse in Thüringen am Beispiel der Gera, In: Fiedler, F., Nestmann, F. &
858 M. Kohler [Hrsg.]: Naturkatastrophen in Mittelgebirgsre-gionen, Proceedings
859 zum Symposium am 11. und 12. Oktober 1999 in Karlsruhe. 185-212.
860
- 861 Deutsch, M., Böhner, J., Pörtge, K.-H. & K. Tilman Rost (2004): Untersuchung histo-
862 rischer Hochwasserereignisse in Thüringen – dargestellt am Beispiel der Werra.
863 – In: Zeitschrift für Geomorphologie, Supplement Vol 135. Borntraeger Berlin,
864 Stuttgart August 2004. 11-23.
865
- 866 Die Chroniken der schwäbischen Städte (1865 – 1928): Augsburg, Bd. 1-9. – Histori-
867 sche Kommission der Bayerischen Akademie der Wissenschaften [Hrsg.] (1862
868 – 1968): Die Chroniken der deutschen Städte. – 37 Bände. Hirzel Leipzig,
869 Perthes Stuttgart/Gotha, Schünemann Bremen.
870
- 871 Endlicher, W. & F.-W. Gerstengarbe [Hrsg.] (2009): Der Klimawandel – Einblicke,
872 Rückblicke und Ausblicke. –
873 <http://edoc.hu-berlin.de/miscellanies/klimawandel> (03.03.2011)
874
- 875 Feulner, G. (2011), Are the most recent estimates for Maunder Minimum solar irradi-
876 ance in agreement with temperature reconstructions?, *Geophys. Res. Lett.*, 38,
877 L16706, doi:10.1029/2011GL048529.
878
- 879 Feulner, G. and Rahmstorf, S. (2010): On the effect of a new grand minimum of solar
880 activity on the future climate on Earth GEOPHYSICAL RESEARCH LET-
881 TERS, VOL. 37, L05707, doi:10.1029/2010GL042710, 2010
882
- 883 Fliri, F. (1998): Naturchronik von Tirol. – Innsbruck.
884
- 885 Glaser, R. (2008): Klimageschichte Mitteleuropas, 1200 Jahre Wetter, Klima, Kata-
886 strophen. – 2. Auflage. Darmstadt.
887
- 888 Glaser, R., Beck, Ch. & H. Stangl (2004a): Zur Temperatur- und Hochwasserentwick-
889 lung der letzten 1000 Jahre in Deutschland. – DWD Klimastatusbericht2003. 55-
890 67.
891
- 892 Glaser, R. & H. Stangl (2003a): Floods in central Europe since 1300. – In: V.R.
893 Thorndycraft, G. Benito, M. Barriendos & M.C. Llasat [Hrsg.]: Palaeofloods,
894 Historical Floods and Climatic Variability: Applications in Flood Risk Assess-
895 ment (Proceedings of the PHEFRA Workshop, Barcelona, 16-19th October,
896 2002). 93-98.
897
- 898 Glaser, R. & H. Stangel (2003b): Historical Floods in the Dutch Rhine Delta. – In:
899 Natural Hazards and Earth System Sciences, Issue 3. 605-613.
900
- 901 Glaser, R., Jacobeit, J., Deutsch, M. & H. Stangel (2002): Hochwasser als historisches
902 Phänomen. – Rundgespräche der Kommission für Ökologie der bay. Akademie

- 903 der Wissenschaft, Bd. 24 Hochwasser und Ökologie – Katastrophe oder Chan-
904 ce? 15-30.
905
- 906 Groos, W. (1967): Beiträge zur Topographie von Alt-Augsburg. – 21. Bericht der Na-
907 turforschenden Gesellschaft Augsburg.
908
- 909 Hoyt, D. V. & K. H. Schatten (1997): Group Sunspot Numbers: A New Solar Activity
910 Reconstruction. – In: Solar Physics 181 (1998). 491-512.
911
- 912 Jacobeit, J., Beck, Ch., Philipp A. (1998): Annual to Decadal Variability in Cli-
913 mate in Europe. – Würzburger Geographische Manuskripte, Heft 43.
914
- 915 Kiss-Laszlovszky: 14th-16th-century Danube floods and long-term water-level
916 changes in archaeological and sedimentary evidence in the western and central
917 Carpathian Basin: An overview with documentary comparison. Journal of Envi-
918 ronmental Geography 2013 6/3-4, 1-11
919
- 920 Lamb, H. H. (1980): Weather and Climate Patterns of the Little Ice Age. – In: Das
921 Klima. Analysen und Modelle, Geschichte und Zukunft. Springer, Berlin, Hei-
922 delberg, New York. 149-160.
923
- 924 Lean, J. and Rind, D.: Climate forcing by changing solar radiation, J. Climate, 11,
925 3069–3094, 1998.
926
- 927 Luterbacher, J., Xoplaki, E., Dietrich, D., Jones, P.D., Davies T.D., Portis, D., Gonza-
928 lez-Rouco, J.F., von Storch, H., Gyalistras, D., Casty, C. & H. Wanner (2002):
929 Extending North Atlantic Oscillation reconstructions back to 1500. – In: Atmos-
930 pheric Science Letters.
931
- 932 Miller G. H., Geirsdóttir, Á., Zhong, Y., Larsen, D. J., Otto-Bliesner, B. L., Holland,
933 M. M., Bailey, D. A., Refsnider, K. A., Lehman, S. J., Southon, J. R., Anderson,
934 C., Björnsson, H. & T., Thordarson, T. (2012): Abrupt onset of the Little Ice
935 Age triggered by volcanism and sustained by sea-ice/ocean feedbacks. – GEO-
936 PHYSICAL RESEARCH LETTERS; VOL. 39, L02708, 5 PP., 2012
937 doi:10.1029/2011GL050168
938
- 939 Miltzer, S. (1998): Klima - Umwelt - Mensch. Band I – III. – Abschlußbericht zum
940 DFG-Projekt MI-493, Leipzig.
941
- 942 Mudelsee, M., Börngen, M., Tetzlaff, G. & U. Grünewald (2004): Extreme floods in
943 central Europe over the past 500 years: Role of cyclone pathway “Zugstrasse
944 Vb”. – In: JOURNAL OF GEOPHYSICAL RESEARCH, VOL. 109.
945 doi:10.1029/2004JD005034
946
- 947 Pfister, Ch. (1984): Klimageschichte der Schweiz 1525 – 1860. – Haupt,
948 Bern/Stuttgart.
949

- 950 Pfister, C. (1996): Häufig, selten oder nie. Zur Wiederkehrperiode der grossräumigen
951 Überschwemmungen im Schweizer Alpenraum seit 1500. – Jb. d. Geogr. Ges.
952 Bern, Bd. 59. 139-149.
953
- 954 Pfister, Ch. (1999): Wetternachhersage. – Haupt Bern/Stuttgart.
955
- 956 Rohr, Ch. (2008): Extreme Naturereignisse im Ostalpenraum. Naturerfahrung im
957 Spätmittelalter und am Beginn der Neuzeit. – Reihe: Umwelthistorische For-
958 schungen 4. Böhlau Köln 2007.
959
- 960 Rohr, Ch. (2013): Floods of the upper Danube River and its tributaries and their im-
961 pact on Urban economies (c. 1350–1600): The Examples of the towns of
962 Krems/Stein and Wels (Austria). – Environment and History 19/2 (2013), 133-
963 148.
964
- 965 Schmoecker-Fackel, P. & Naef, F. (2010): Changes in flood frequencies in Switzer-
966 land since 1500 Hydrol. Earth Syst. Sci., 14, 1581–1594, 2010
967
- 968 Schnurrer, F. (1823): Die Krankheiten des Menschen-Geschlechts historisch und geo-
969 graphische betrachtet. Chronik der Seuchen in Verbindung mit den gleichzeiti-
970 gen Vorgängen in der physischen Welt und in der Geschichte der Menschen.
971 Erster Theil vom Anfang der Geschichte bis in die Mitte des fünfzehnten Jahr-
972 hunderts. – Thübingen 1823, bei Christian Friedrich Osiander.
973
- 974 Schönwiese, Ch.-D. (2008): Klimatologie. – 3. Auflage, UTB Stuttgart.
975
- 976 Schönwiese, Ch.-D. (1992): Praktische Statistik für Meteorologen und Geowissen-
977 schaftler. – Borntraeger, Berlin, 1992.
978
- 979 Shindell, D. T., Schmidt, G.A., Mann, M. E, Rind, D. & A., Waple (2001): Solar
980 Forcing of Regional Climate Change During the Maunder Minimum. –Science 7
981 December 2001: Vol. 294 no. 5549 pp. 2149-2152 DOI:
982 10.1126/science.1064363
983
- 984 Sonklar, K., Edler von Innstaedten (1883): Von den Überschwemmungen. – A. Hart-
985 leben's Verlag, Wien, Pest, Leipzig.
986
- 987 Stahleder, H. (1995 – 2005): Chronik der Stadt München. – 3 Bände, Dölling und Ga-
988 litz Verlag, München.
989
- 990 Sturm, K., Glaser, R., Jacobeit, J., Deutsch, M., Brázdil, R., Pfister, C., Luterbacher, J.
991 & H. Wanner (2001): Hochwasser in Mitteleuropa seit 1500 und ihre Beziehung
992 zur atmosphärischen Zirkulation. – In: Petermanns Geographische Mitteilungen,
993 145. 14-27.
994
- 995 Usoskin, I. G., Schüssler, M., Solanki, S. K., & Mursula, K. (2004): Solar activity
996 over the last 1150 years: does it correlate with climate? – In: Proceedings of the
997 13th Cambridge Workshop on Cool Stars, Stellar Systems and the Sun, held 5-9

- 998 July, 2004 in Hamburg, Germany. Edited by F. Favata, G.A.J. Hussain, and B.
999 Battrick. ESA SP-560, European Space Agency, 2005., p.19
1000
1001 van Bebbber, W. J. (1891): Die Zugstraßen der barometrischen Minima. – In: Meteor.
1002 Z. 361-366.
1003
1004 Weikinn, C. (1958): Quellentexte zur Witterungsgeschichte Europas von der Zei-
1005 tenwende bis zum Jahre 1850. – Band I. Akademieverlag, Berlin.
1006
1007 Weikinn, C. (1960): Quellentexte zur Witterungsgeschichte Europas von der Zei-
1008 tenwende bis zum Jahre 1850. – Band II. Akademieverlag, Berlin.
1009
1010 Weikinn, C. (1961): Quellentexte zur Witterungsgeschichte Europas von der Zei-
1011 tenwende bis zum Jahre 1850. – Band III. Akademieverlag, Berlin.
1012
1013 Weikinn, C. (1963): Quellentexte zur Witterungsgeschichte Europas von der Zei-
1014 tenwende bis zum Jahre 1850. – Band IV. Akademieverlag, Berlin.
1015
1016 Wetter, O., Pfister C, Weingartner R, Luterbacher J, Reist T & J. Trösch (2011): The
1017 largest floods in the high Rhine basin since 1268 assessed from documentary
1018 and instrumental evidence. – Hydrol Sci J 56(5):733-758
1019
1020 Zentralanstalt für Meteorologie und Geodynamik (ZAMG): Lufttemperatur – Ent-
1021 wicklung in Etappen, available at: <http://www.zamg.ac.at/cms/de/klima/informationsportal-klimawandel/klimavergangenheit/neoklima/lufttemperatur> (last
1022 access: 23 June 2014), 2014.
1023
1024
1025 Zillner, F. von (1885): Geschichte der Stadt Salzburg. – Druck von J. Dellache, Salz-
1026 burg.
1027