



HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty, A. M. Kinoshita, and T. S. Hogue

Colorado School of Mines Civil and Environmental Engineering, 1500 Illinois Street, Golden, CO 80401, USA

Received: 14 May 2014 – Accepted: 23 June 2014 – Published: 4 July 2014

Correspondence to: T. S. Hogue (thogue@mines.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

The current work evaluates the spatial and temporal variability in snow after a large forest fire in northern California with Moderate Resolution Imaging Spectroradiometer (MODIS) snow covered area and grain size (MODSCAG) algorithm. MODIS MOD10A1 fractional snow covered area and MODSCAG fractional snow cover products are utilized to detect spatial and temporal changes in snowpack after the 2007 Moonlight Fire and an unburned basin, Grizzly Ridge, for water years (WY) 2002–2012. Estimates of canopy adjusted and non-adjusted MODSCAG fractional snow covered area (fSCA) are smoothed and interpolated to provide a continuous timeseries of daily basin average snow extent over the two basins. The removal of overstory canopy by wildfire exposes more snow cover; however, elemental pixel comparisons and statistical analysis show that the MOD10A1 product has a tendency to overestimate snow coverage pre-fire, muting the effects of wildfire. The MODSCAG algorithm better distinguishes sub-pixel snow coverage in forested areas and is highly correlated to soil burn severity after the fire. Annual MODSCAG fSCA estimates show statistically significant increased fSCA in the Moonlight Fire study area after the fire (WY 2008–2011; $P < 0.01$) compared to pre-fire averages and the control basin. After the fire, the number of days exceeding a pre-fire high snow cover threshold increased by 81%. Canopy reduction increases exposed viewable snow area and the amount of solar radiation that reaches the snowpack leading to earlier basin average melt-out dates compared to the nearby unburned basin. There is also a significant increase in MODSCAG fSCA post-fire regardless of slope or burn severity. Alteration of regional snow cover has significant implications for both short and long-term water supplies for downstream communities and resource managers.

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



1 Introduction

The last several decades have been marked by distinct increases in large-wildfire frequency as well as fire duration and season across the western US (Westerling et al., 2006). Soil and vegetation change after fire result in increased flooding, mass-wasting, increased runoff intensities, long-term changes in the energy and water budget, and increased air pollutants (Swanson, 1981; Kattelman et al., 1983; Stednick, 1996; Webb et al., 2012). Storm runoff also liberates atmospherically deposited contaminants and mobilizes particulate-bound constituents, degrading post-fire water quality (Stein et al., 2012; Burke et al., 2013). Vegetation recovery significantly controls long-term hydrologic conditions and elevated discharged has been observed for nearly ten years post-fire (Kinoshita and Hogue, 2011). Similarly, forest canopy considerably influences snowpack properties and snowmelt response (Faria et al., 2000). Given the dependency of the Western US on snowpack and mountain runoff for water supply (NRCS, 2012) and the assumption of stationarity, under which water reservoir systems are designed and managed (Milly et al., 2008), minimal forest structure alterations will have critical implications for regional and state water resources and management.

Field-based studies have found that disturbance in forest structure considerably impacts snow accumulation and melt properties, altering water yield from snow dominated basins (Kattelman et al., 1983; Stednick, 1996; Faria et al., 2000; Stephens et al., 2012; Webb et al., 2012). Post-fire changes in snowpack energy balance include increased radiation exposure, decreased snow albedo due to surface alterations from charred soils, dust, or vegetation, and changes in soil temperature (Painter et al., 2007; Burles and Boon, 2011; Ebel et al., 2012; Gleason et al., 2013; Harpold et al., 2013). The opposing effects of increased snow accumulation but increased snow ablation have been documented at the plot scale for the first year following a wildfire (Gleason et al., 2013; Harpold et al., 2013). Plot-scale studies generally reported significant increases in snow accumulation in burned areas compared to nearby control plots due to the lack of canopy interception (Burles and Boon, 2011; Harpold et al.,

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2013). Decreased canopy cover reduces snow interception, increases solar radiation exposure, and alters sublimation of the exposed snowpack (Faria et al., 2000; Varhola et al., 2010; Harpold et al., 2013). Harpold et al. (2013) showed winter season ablation reduced snowpack depths by 50 % prior to melt and a 10 % reduction in snow water equivalent in burned areas the first year after fire. Gleason et al. (2013) showed a 40 % decrease in snow albedo accompanied by a 200 % increase in net shortwave radiation in burned forest plots compared to unburned forests. However, effects are undocumented at the watershed scale and there is a need for additional studies on snow accumulation and melt variability from forest cover change (Varhola et al., 2010).

Remote sensing products, including NASA Moderate Resolution Imaging Spectroradiometer (MODIS) MOD10A1 and MODIS Snow Covered Area (SCA) and Grain size (MODSCAG), a spectral mixing product, provide the spatial and temporal resolution necessary for monitoring large-scale wildfires that often impact inaccessible and ungaged snow-dominated basins. To our knowledge, no study has investigated pre-fire and post-fire snow cover change using satellite imagery. The current study facilitates identification of remote sensing tools capable of detecting spatial and temporal changes in post fire snowpack through application of MODIS MOD10A1 and MODSCAG fractional snow covered area (fSCA) products to the 2007 Moonlight Fire in northern Sierra Nevada, California. Specifically, the objectives of our work are to: (1) understand spatial and temporal variability of pre- and post-fire fSCA with MODIS (MOD10A1 and MODSCAG) products, (2) compare MOD10A1 and MODSCAG products in pre- and post-fire conditions to determine the better indicator of SCA, (3) investigate the influence of aspect, burn severity, and general climate patterns on post-fire snow behavior (using fSCA as a proxy), and (4) evaluate post-fire recovery patterns in a snow-dominated basin over several years.

2 Study areas

2.1 Moonlight fire

There is a statistically significant ($P = 0.05$) increase in total area burned in the Sierra Nevada from the 1980s to the present. The 1980s decadal average of burned area increased from 300 to 900 km² in the current decade (Wildland Fire Incidents, 2013). The Moonlight Fire burned over 250 km² (27 370 ha) in the Plumas National Forest (about 190 km north of Sacramento) from 3–15 September 2007 on the eastern side of the northern Sierra Nevada divide (Fig. 1). Since the late 1800s, this was the first major wildfire recorded in this area (California Department of Forestry and Fire Protection, 2012). Steep terrain and high winds caused a mosaic of soil burn severities resulting in concentrated areas of high surrounded by moderate to low/unburned areas (USDA Forest Service RSAC, 2007; Fig. 1). Pre-fire vegetation consisted of mostly evergreen forest (90%) with some riparian and shrub/scrub areas (Fry et al., 2013; Table 1). The slope aspects within Moonlight Fire are evenly distributed, with a dominant south facing slope, followed closely by west, north, and east (Table 1). The Moonlight Fire burn area has an elevation range of 1090–2290 m and receives on average 680 mm of precipitation a year, the majority of which falls in the winter months as snow (Table 1).

2.2 Grizzly Ridge

To evaluate the fire signal relative to regional climate variability a complimentary regional control basin, Grizzly Ridge, was chosen for comparison. The Grizzly Ridge area has not burned within the last 100 years of record (California Department of Forestry and Fire Protection, 2012). It is 150 km² (14 800 ha) approximately 24 km south of Moonlight Fire on the same side of the divide in the Sierra Nevada (Fig. 1). Vegetation within the Grizzly Ridge area is comprised of mostly evergreen forest (80%) and shrub/scrub in the lower elevations (Fry et al., 2013; Table 1). The slope aspects exhibits similar patterns as Moonlight Fire, although Grizzly Ridge has roughly 10%

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



more south facing slopes (Table 1). The Grizzly Ridge area has an elevation range of 1300–2320 m and receives an annual basin average of 880 mm of precipitation.

3 Methods

MODIS MOD10A1 and MODSCAG products were gathered for both areas, Moonlight Fire and Grizzly Ridge, from 1 October 2001 to 30 September 2012 (water year – WY – 2002–2012). Both products only identify areas covered by snow, not snowpack depth – a longer snow season will distinguish more fSCA, but not depth changes or snow water equivalent. Annual and monthly precipitation and maximum and minimum temperatures for Moonlight Fire and Grizzly Ridge were estimated from the Parameter-elevation Regressions on Independent Slopes Model (PRISM) climate data set (Daly, 1994, 1997, 2002). Conterminous US products are downloaded from the PRISM Climate Group (<http://www.prism.oregonstate.edu/>) and the monthly 4 km pixels are extracted within Moonlight Fire and Grizzly Ridge and averaged over both domains for WY 2002–2012.

3.1 Remote sensing products

3.1.1 MODIS MOD10A1

The Terra MODIS SCA product (MOD10A1) provides atmospherically corrected daily fractional snow cover at 500 m spatial resolution based on the normalized difference snow index (NDSI). The preprocessed MODIS product includes spectral thresholds that mask and screen for clouds and low reflectance surfaces such as water (Salomonson and Appel, 2004). To account for snow in densely vegetated areas Klein et al. (1998) developed a method that uses a combined snow reflectance model and canopy reflectance model to map more snow in forested areas using normalized NDSI and the normalized difference vegetation index (NDVI; Klein et al., 1998). The NDVI normalizes

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the reflectances in the near-infrared and visible (red) wavelengths to differentiate vegetation where there is chlorophyll absorption of red light for photosynthesis and reflection of near-infrared light (Tucker, 1979):

$$\text{NDVI} = \frac{R_{\text{NIR}} - R_{\text{VIS}}}{R_{\text{NIR}} + R_{\text{VIS}}} \quad (1)$$

5 where R_{NIR} is near-infrared reflectance and R_{VIS} is red reflectance in the visible spectrum. The NDSI is evaluated as (Dozier, 1989):

$$\text{NDSI} = \frac{R_{\text{VIS}} - R_{\text{SWIR}}}{R_{\text{VIS}} + R_{\text{SWIR}}} \quad (2)$$

10 where, R represents spectral reflectances in the visible and shortwave infrared bands. The vegetation correction is used to map snow when $\text{NDSI} < 0.4$ and $\text{NDVI} > 0.1$.

The newest publicly available version [005] of MODIS fractional snow covered area, MOD10A1 is a daily, 500 m product, available from 2000 to the present (Hall et al., 2006). MOD10A1 fSCA is based on an empirical snow mapping algorithm developed from a linear regression between binary Landsat Thematic Mapper snow cover and
15 MODIS NDSI (Salomonson and Appel, 2004; Hall et al., 1995):

$$\text{fSCA} = -0.01 + 1.45\text{NDSI}. \quad (3)$$

This algorithm is used to map fractional snow cover and performs relatively well in the winter months in mountainous regions compared to other remote sensing products and
20 ground-based observations (Maurer et al., 2003; Pu et al., 2007).

3.1.2 MODSCAG

MODSCAG is derived from a physically-based algorithm which uses a multispectral mixing analysis to identify sub-pixel snow covered area and grain size (Painter et al., 2009). The MODSCAG model has been validated over the Sierra Nevada, Rocky

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



3.1.3 Canopy adjustment

Forest canopy obstructs the view of the ground by MODIS, causing underestimates of snow cover in dense forests (Raleigh et al., 2013). Hence, forest cover density data is used to indicate snow cover masked by canopy and improve MODSCAG estimates of viewable snow cover (Molotch and Margulis, 2008):

$$fSCA_{Adj} = \frac{fSCA_{Ob}}{1 - fVeg} \quad (5)$$

where $fSCA_{Ob}$ is the observed MODSCAG fSCA and $fVeg$ is the annual density of forest cover or the fraction of vegetation estimated using a combination of the MODIS percent tree cover product (MOD44B) and the MODSCAG fraction of vegetation (Painter et al., 2009; DiMiceli et al., 2011). The canopy adjusted fSCA (Eq. 5) assumes that the distribution of snow under a canopy is equivalent to viewable open areas between trees or in clearings. This assumption that spatial distribution of snow in viewable gaps can be interpolated to nearby canopied forests is not as reliable during the accumulation and melt periods (Raleigh et al., 2013). A rigorous correction to improve estimations of snow under canopy using optical sensors remains an area of active research for remote sensing in forested terrains, and is outside the scope of this study. In the current study, MODSCAG fSCA is adjusted for canopy using Eq. (5), while MOD10A1 SCA uses the linear NDSI algorithm and is not scaled by forest cover density.

3.2 Spatial and temporal analysis

3.2.1 Basin fSCA interpolation

Temporal analysis for WY 2002–2012 uses daily basin averaged MODSCAG fSCA for both Moonlight Fire and Grizzly Ridge. The daily data initially has gaps and errors from cloud cover, sensor viewing geometry, or imperfections in the retrieval algorithm. A combination of noise filtering, snow/cloud discrimination, and interpolation and

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

smoothing improves the MODSCAG daily snow cover timeseries (Dozier et al., 2008). Dozier et al. (2008) view the snow data as a space-time cube, which can be filtered, smoothed, and interpolated. In the current study, the space-time cube is filtered to remove cloudy or noisy values; the remaining data is used to interpolate and smooth gaps within the cube.

Filtering consists of several steps: (1) a two-dimensional adaptive Wiener filter (Matlab wiener2 function) is used to identify noise and data dropouts in all seven land reflectance bands, where the Boolean variable is set to 1 for raw fractional snow-covered area that is 0, (2) quality flags from the MOD09 product are used to identify snow-covered pixels as cloudy. False positives and false negatives are identified from MODSCAG snow cover (fSCA) and grain size (r) processing. Then thresholds (false positives: $fSCA > 0.6 \wedge r \geq 100 \mu\text{m}$ and false negatives: $fSCA > 0.6 \wedge r \leq 100 \mu\text{m}$) are used to reduce misidentification, (3) to correct for values obscured by MODIS scan angles (the primary source of error), the time dimension of the space-time cube is interpolated using a cubic smoothing spline (Matlab csaps function). The current study uses 16 days (representing a MODIS viewing angle cycle) for the limits of integration; the smoothing parameter is adaptive and varies spatially depending on the extent of cloud cover or missing data. The weight varies from 0 to 1 and is based on the viewing angle (determined from the corresponding MOD09GA) such that the near-nadir views have the greatest weights. If the cubic smoothing spline yields unrealistic values from gaps in data, the smoothed fSCA values are interpolated using a piecewise interpolant; and (4) after steps 1–3, the whole cube is smoothed with a Gaussian filter, providing a continuous data stream of snow covered area.

3.2.2 Elemental pixel comparison

Differencing maps for each gridded fSCA product, MOD10A1 and MODSCAG, are developed by taking the difference between winter (January–March) pre-fire average fSCA (WY 2002–2007) and post-fire average fSCA (WY 2008–2012); the domain includes 1099 pixels. The difference maps ($\Delta fSCA$) are used to detect spatial changes

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



in snow cover after the fire. An elemental pixel comparison (EPC) between MODSCAG fSCA and MOD10A1 fSCA is evaluated using a least-squares linear regression analysis of individual pre- and post-fire winter pixels. EPC is also used to investigate temporal changes in snow cover based on corresponding basin attributes including burn severity and slope aspect. Gridded daily fSCA is disaggregated over each domain by slope aspects (north, south, east and west) derived from a USGS National Elevation Dataset (NED) 30 m Digital Elevation Model (DEM). Daily basin average estimates are then produced for each slope aspect for WY 2002 to 2012 for Grizzly Ridge and Moonlight Fire. For Moonlight Fire, daily fSCA was also disaggregated to match a 30 m soil burn severity map for EPC (USDA Forest Service RSAC, 2007). A time series of basin averaged fSCA is made based on each burn severity (i.e., high, moderate, and low-unburned) from WY 2002 to 2012 for statistical analyses.

3.3 Statistical analysis

3.3.1 MODSCAG cumulative distribution function

Annual cumulative distribution functions (CDFs) are developed using daily basin averaged fSCA for both Moonlight Fire and Grizzly Ridge to investigate annual shifts in snow cover after fire. Fractional SCA cumulative distribution functions are similar to flow duration curves, which are used to investigate annual changes in flow regimes due to forest disturbance (Lane et al., 2006; Brown et al., 2005). Fractional SCA CDFs are used to determine the probability that a specific basin averaged fSCA will be equaled or exceeded during a given time period. Exceedance probabilities derived from the pre-fire fSCA duration curves are used to establish high and low thresholds for analysis. High snow cover days are defined based on the pre-fire long-term CDFs with an exceedance probability of 10 % or less.

During the beginning and end of the snow season, as MODSCAG and MOD10A1 pixels approach an fSCA value of 15 % (very low fractional snow covered area), there is increased uncertainty and larger errors in positively identifying snow (Rittger et al.,

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2013). This study uses an exceedance probability of 70 % (representing 10 % basin average snow cover) to identify a low SCA melt-out threshold and reduce error from misidentification of snow. A basin average of 10 % fSCA contains enough individual pixels with relatively high fSCA to provide confident basin average estimates. This 70 % exceedance probability threshold also corresponds to the most widely used definition of low flow as derived from flow duration curves (70–99 %; Smakhtin, 2001).

To quantify the change from pre-fire to post-fire, a two-sample Kolmogorov–Smirnov (K–S) test is used to compare the distributions of pre- and post-fire fSCA CDFs. The K–S null hypothesis is that the pre- and post-fire fSCA CDFs are from the same continuous distribution at $\alpha = 0.01$ (Massey, 1951), where the K–S test statistic is the maximum vertical distance between the two curves being evaluated (Cowpertwait et al., 2013).

3.3.2 Analysis of variance

An Analysis of Variance (ANOVA) is used to determine the statistical significance of temporal changes in snow cover after fire. Daily basin averaged fSCA estimates are separated annually based on the water year, excluding summer months (July to September), and by basin attributes (burn severity and slope aspect). The fSCA is then evaluated for statistical differences from the pre-fire period and compared to the control domain (Grizzly Ridge). The null hypothesis that the mean of each post-fire annual fSCA (WY 2008–2012) is similar to the pre-fire annual mean (WY 2002–2007) is tested at $\alpha = 0.01$.

4 Results

4.1 MODSCAG and MOD10A1 comparison

Non-canopy adjusted MODSCAG and MOD10A1 differencing maps for Moonlight Fire show a distinct difference in fSCA after the fire (Fig. 2). Generally, the spatial pattern of

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the increased fSCA for both products follow the high soil burn severity in the Moonlight Fire. Higher soil burn severity near the center of the domain results in reduced canopy cover and more visible snow and snow covered area. An EPC and linear regression of Δ fSCA and soil burn severity shows a stronger correlation of non-canopy adjusted MODSCAG Δ fSCA to soil burn severity ($r = 0.56$) than MOD10A1 Δ fSCA ($r = 0.43$). Non-canopy adjusted MODSCAG has a basin average increase in fSCA of 0.3 (Fig. 2, right panel) after the fire whereas MOD10A1 displays smaller differences throughout the burned domain and increases, on average by 0.2 (Fig. 2, left panel). For the MODSCAG product, 44 % of the Moonlight Fire domain exhibited Δ fSCA values of least 0.3, while MOD10A1 has 21 % of the domain with values of 0.3 or higher.

The least-squared linear regression analysis of MOD10A1 fSCA and MODSCAG fSCA established from the EPC shows a distinct difference between pre- and post-fire correlation (Fig. 3). MOD10A1 tends to produce higher estimates of fSCA compared to MODSCAG across the entire domain pre- and post-fire. MOD10A1 is biased high compared to MODSCAG, but the pre-fire linear correlation between the two products is relatively high ($r = 0.85$). After the fire there is an increase in variability and the linear relationship between MOD10A1 and MODSCAG decreases ($r = 0.69$). The linear regression line is also higher post-fire (Fig. 3). The upward shift in the regression line in the MODSCAG direction is consistent with the increase in visible fSCA (Fig. 2). Decreases in the correlation coefficient after the fire are most likely due to differences in the amount of increased fSCA identified by each product.

Product assessment studies have shown that MOD10A1 fSCA overestimates snow cover in densely vegetated areas (Rittger et al., 2013). These results are consistent with our linear regression analysis. This can be attributed to the MOD10A1 snow-mapping algorithm and NDVI threshold indices (Klein et al., 1998) that are used to identify snow in forested areas. NDVI is a greenness index based on surface reflectances and does not differentiate vegetation types. Therefore, the current NDVI threshold (> 0.1) increases mapped snow cover in areas with shrubs and grasses the same as forested areas. Reduced canopy cover from wildfire should lead to increased

average difference in canopy-adjusted and non-adjusted fSCA ensembles is approximately 0.30 for both Grizzly Ridge and Moonlight, while after the fire the difference is decreased in the Moonlight Fire, on average, to 0.18. The non-adjusted MODSCAG fSCA values show a significant increase in basin averaged fSCA (or exposed snow cover) after the Moonlight Fire in 2007 ($P < 0.01$) due to the stand replacing fire (Fig. 4). MODSCAG fSCA increased, but the canopy adjustment has no statistically significant increase in annual fSCA. However, exposed areas with increased viewable fSCA exhibit altered accumulation and melt behavior due to changes in the snowpack energy budget and are further analyzed with both canopy adjusted and non-adjusted fSCA.

4.3 MODSCAG cumulative distribution functions

Annual CDFs of basin averaged non-canopy adjusted and canopy adjusted MODSCAG fSCA for both Moonlight Fire and Grizzly Ridge highlight shifts in snow cover after the fire (Fig. 5). The spread in the pre-fire (Fig. 5; black) cumulative distribution functions are attributed to snow season climate variability. For post-fire water years 2008–2011 the cumulative distribution functions are statistically different from the pre-fire curve ($P < 0.01$), and the null hypothesis is rejected. However, WY 2012 falls within the pre-fire distributions and is not statically different. The K–S statistic indicates post-fire non-adjusted fSCA distributions are elevated, on average, by 40 % compared to pre-fire non-adjusted curves. The canopy adjusted fSCA curves are not as sensitive, but still increase by 14 % after the fire. The distribution of the post-fire curves in Moonlight is generally higher compared to Grizzly Ridge and is especially apparent using the non-adjusted fSCA (Fig. 5a). The shape of the fSCA curves significantly change after the fire due to the upward shift in inflection points. This shifting distribution indicates a higher post-fire probability that the basin will have larger areas of exposed snow coverage.

Using the thresholds established from the cumulative distribution functions, the consecutive number of high snow cover days with respect to the length of snow season are shown for Moonlight Fire (Fig. 6a and b) and Grizzly Ridge (Fig. 6c and d). Post-fire,

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



there are more days with high snow cover in Moonlight Fire than pre-fire and compared to Grizzly Ridge for both canopy adjusted (Fig. 6c and d) and non-canopy adjusted fSCA values (Fig. 6a and b). On average, there were 13 days that exceeded the high snow cover threshold in the Moonlight Fire before the fire, whereas after the fire there are on average 70 days classified as high snow cover. Temporal distributions highlight daily basin averaged SCA patterns throughout each year for both canopy adjusted and non-adjusted (Fig. 6, right panel). Larger fSCA patterns are noticeable during winter months – December (12) through April (5) – after the fire. The canopy adjusted fSCA plots (Fig. 6b and d) have larger values relative to the non-canopy adjusted due to the linear scaling based on the vegetation fraction (Fig. 6a and c); and is congruent with the annual cumulative distribution functions (Fig. 5).

4.4 ANOVA

An ANOVA of non-adjusted MODSCAG fSCA shows that post-fire annual basin averaged fSCA for WYs 2008–2011 are significantly higher than pre-fire averages in the Moonlight basin at $\alpha=0.01$ ($P < 0.01$; Fig. 7). For the pre-fire years (WY 2002–2007), both Moonlight Fire and Grizzly Ridge follow similar annual basin averaged fSCA trends ($r = 0.92$). Before the fire the Moonlight Fire area had, on average, 17 % less basin averaged fSCA than Grizzly Ridge. After the fire, however, the Moonlight Fire area had an average of 26 % more fSCA than Grizzly Ridge. The Moonlight Fire and Grizzly Ridge domains are also sensitive to winter precipitation, including amount of precipitation and duration of the snow season. Total precipitation as well as the length of snow season in Moonlight Fire and Grizzly Ridge were above average in WY 2005 (Table 2) and yielded more fSCA; while WY 2007 was dry and resulted in less basin averaged fSCA (Fig. 7). For the Moonlight Fire, WY 2012 lies within the pre-fire interval and is similar to the pre-fire average, but may be climate induced. Annual precipitation in WY 2012 is 380 mm (Moonlight Fire) and 520 mm (Grizzly Ridge), which corresponds to the lower fSCA. Annual basin average fSCA estimates in Grizzly Ridge note only one (WY 2011) statically significant increase in fSCA during the post-fire period of WY 2008–2012,

which is attributed to the larger than average annual precipitation and length of snow season (1200 mm).

After the fire, there are significantly higher annual basin averaged fSCA estimates based on slope aspect and soil burn severity (bold values denote statistical significance; Table 3). Regardless of slope aspect and burn severity, statistically significant increases in fSCA for Moonlight Fire are observed from WY 2008 to 2011 ($P < 0.01$). WY 2012 in all aspects and burn severity is not significantly different than pre-fire fSCA values, but is still relatively high considering that it also received the lowest amount of total precipitation in the 11 year study period. Generally, the high soil burn severity areas within the Moonlight Fire domain have slightly larger annual average fSCA values than moderate and low-unburned (Table 3).

4.5 Annual melt-out dates

Annual melt-out dates are estimated for Grizzly Ridge and Moonlight Fire based on the 70 % exceedance (10 % basin averaged fSCA) threshold established from the canopy adjusted MODSCAG fSCA cumulative distribution functions. At 10 % coverage, the domain will have lost the vast majority of its snowpack due to melt. Annual melt-out dates for Grizzly Ridge and Moonlight Fire are compared for pre-fire and post-fire years (Fig. 8). Although the melt-out dates are variable from year to year based on annual snow conditions, Grizzly Ridge and Moonlight Fire melt-out dates are relatively similar pre-fire, where it is observed that Moonlight typically melts out an average of 1.5 days after Grizzly Ridge and ranges from -0.5 to 7 days with a standard deviation of 3 days (Fig. 8b).

The average long-term pre-fire difference in melt-out dates (1.5 days) between Moonlight Fire and the control basin, Grizzly Ridge, are used to estimate the expected melt-out day for WY 2008–2012 assuming no fire (Fig. 8a; red solid diamonds). With the fire, the observed annual difference in melt-out dates between Moonlight Fire and Grizzly Ridge show an average decrease of 7.5 days and more variability post-fire, with a standard deviation of 11 days (Fig. 8b). Thus relative to pre-fire averages, Moonlight melts

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fSCA ensembles in both basins followed similar trends ($r = 0.96$), but there is a notable increase from non-canopy adjusted MODSCAG fSCA in Moonlight Fire as compared to Grizzly Ridge of 26 %, post-fire.

A decomposition of fSCA in the Moonlight Fire area based on slope aspect and soil burn severity using the EPC is employed to investigate the influence of each attribute. Results show statistically significant increases in fSCA from WY 2008 to 2011 regardless of slope aspect and soil burn severity because of acute changes in vegetation structure and the resulting exposure of more snow cover. Water year 2012 is the only year after the fire that does not show statistically significant changes in fSCA compared to average pre-fire conditions and are attributed to the lowest recorded precipitation in the 11 year study period. Compared to the pre-fire low precipitation year (WY 2007), which received slightly more precipitation than WY 2012, and WY 2012 in Grizzly Ridge, fSCA is still increased by nearly 20 % in Moonlight Fire.

In this study, it was beneficial to investigate MODSCAG fSCA estimates adjusted for canopy cover using Eq. (5) and non-adjusted estimates. Using the two estimates, there is a recognizable change in fSCA due to the reduced vegetation fraction which is apparent as post-fire fSCA ensembles increase and begin to approach the canopy adjusted values. This analysis identifies the importance in incorporating dynamic vegetation fractions when using the canopy adjustment. Static vegetation fractions are likely to result in large overestimates of fSCA after fire, as a result of unnecessary linear scaling of fSCA.

Cumulative distribution functions of canopy and non-canopy adjusted basin averaged MODSCAG fSCA are developed for Moonlight Fire and Grizzly Ridge to investigate post-fire shifts in snow cover and establish high snow cover and melt out thresholds. Using the K–S test, we note that annual post-fire fSCA distribution (WY 2008–2011) is elevated up to 40 % compared to the long-term pre-fire distribution, and are significantly different at $\alpha = 0.01$. This represents a higher probability of high fSCA values across the Moonlight Fire. Before the fire, the 10 % exceedance threshold (defined as high snow cover) corresponded to an average snow coverage of 33 % across the domain

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



using non-canopy adjusted fSCA estimates, and 60 % coverage using the adjusted fSCA values. Using these values as thresholds, it was determined that after the fire, there is an average 81 % increase in the number of high snow coverage days (i.e., days exhibiting higher than 33 % snow coverage or higher than 60 % snow coverage using the non-canopy adjusted and canopy adjusted fSCA estimates, respectively) compared to pre-fire conditions and the control basin. Significant changes in the number of days with high snow coverage from elevated annual fSCA cumulative distribution functions compared to both pre-fire conditions, and the control basin are a consequence of the fire and the removal of forest vegetation. It is likely that the increase in fSCA is directly related to additional exposure of the snow surface that was once hidden by forest canopy.

Significant changes in fSCA over the Moonlight Fire domain influence basin melt out dates. Based on the 70 % exceedance probability threshold established from the cumulative distribution functions, the differences in melt out dates between Moonlight Fire and Grizzly Ridge are similar before the fire, only differing on average by 1.5 days. After the fire, for WY 2008–2011, the entire Moonlight Fire domain melts out, on average, 9 days earlier compared to pre-fire conditions with some years melting out up to 23 days early. The significant increases in exposed snow area from reductions in forest canopy cover increase the amount of solar radiation that reaches the snowpack. Early melt due to changes in the snowpack energy balance is consistent with smaller scale field-based studies by Gleason et al. (2013) and Harpold et al. (2013). Changes in melt-out dates can have significant implications for water resource managers in the western US who rely on the mountain snowpack for a majority of their water supply (Bales et al., 2006). Snowpack melt out dates are also correlated to forest types and species present in the Sierra Nevada (Barbour et al., 2002), and may therefore influence vegetation types during the recovery or regeneration period.

According to this study, there is very little evidence of canopy recovery from WY 2008–2012 over the Moonlight Fire domain to pre-fire conditions as compared to the control basin, Grizzly Ridge. Basin averaged fSCA and melt out dates for WY 2012 fall

within pre-fire averages, but this apparent return or recovery to pre-fire values is partly influenced by climate; as WY 2012 had a low annual basin averaged fSCA because of lower than normal precipitation totals. The sustained post-fire increase in remotely sensed fSCA in Moonlight Fire and earlier melt-out dates is a function of canopy loss.

5 Similar to previous post-fire ecosystem studies, recovery is not expected until there is full canopy regeneration or until the system reaches a new equilibrium (Meixner and Wohlgemuth, 2003; Kinoshita and Hogue, 2011).

6 Conclusions

Continuous mapping of mountainous snow at 500 m resolution using remote sensing techniques has been seldom applied to answer forest disturbance related hydrologic questions. Long term analysis identified distinct differences in the pre- and post-fire snow cover and total visible snow over the burned domain (Moonlight Fire) when compared to a control basin (Grizzly Ridge). The changes in snow coverage and melt-out dates from WY 2002 to 2012 in the Moonlight Fire are attributed to the removal of vegetation after fire and are driven by corresponding changes in the snowpack energy balance. Specific key findings of this study include

MODSCAG's spectral mixing algorithm better identifies snow cover in forested areas and is better correlated to soil burn severity compared to MOD10A1. MODSCAG is ultimately better suited to identify changes in snow cover due to reductions in canopy cover after a wildfire.

There is significantly more basin averaged fSCA ($P < 0.01$) after fire due to reduction of canopy cover and therefore increased viewable snow area.

There are significant increases in the total number of high snow cover days after fire, based on pre- and post-fire cumulative distribution functions.

25 Using the relative difference in melt-out dates between Moonlight Fire and Grizzly Ridge, the Moonlight Fire domain melts out, on average, 9 days earlier after the fire.

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



There is minimal spatial or temporal recovery of canopy and snow cover 5 years after the fire.

Climate change and increasing wildfire frequency and size have the potential to highly alter mountain snowpacks. The release of advanced snow mapping products provides a tool for improved application of remote sensing data to better understand hazards such as fire and offers a unique opportunity for future long-term monitoring and research. The successful application of MODSCAG to the Moonlight Fire burn area provides the first watershed-scale analyses of snow cover and snowmelt detection after a large forest fire.

The shifts in the spatial and temporal distribution of snow throughout the year have significant implications for snow accumulation and melt patterns. This study advocates the application of remote sensing products, such as MODSCAG, due its active and continuous spectral mixing analysis, which can contribute additional insight of regional post-fire snowpack and recovery studies. Remote sensing application improves our understanding and prediction of snowmelt behavior and is crucial for water resources and management, especially in regions that are highly dependent on snowpack and subject to frequent and acute forest disturbance.

Acknowledgements. Special thanks to Thomas Painter and his colleges on the snow hydrology team at NASA JPL for the management and distribution of the MODSCAG product. Support for this research was provided by an NSF RAPID Grant (#EAR1361454) as well as an NSF Hydrologic Sciences Program CAREER Grant (#EAR0846662).

References

- Bales, R. C., Molotch, N. P., Painter, T. H., Dettinger, M. D., Rice, R., and Dozier, J.: Mountain hydrology of the western United States, *Water Resour. Res.*, 42, W08432, doi:10.1029/2005WR004387, 2006.
- Barbour, M., Kelley, E., Maloney, P., Rizzo, D., Royce, E., and Fites-Kaufmann, J.: Present and past old-growth forests of the Lake Tahoe Basin, Sierra Nevada, US, *J. Veg. Sci.*, 13, 461–472, doi:10.1111/j.1654-1103.2002.tb02073.x, 2002.

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Brown, A. E., Zhang, L., McMahon, T. A., Western, A. W., and Vertessy, R. A.: A review of paired catchment studies for determining changes in water yield resulting from alterations in vegetation, *J. Hydrol.*, 310, 28–61, doi:10.1016/j.jhydrol.2004.12.010, 2005.
- Burke, M. P., Hogue, T. S., Ferreira, M., Mendez, C. B., Navarro, B., Lopez, S., and Jay, J. A.: The effect of wildfire on soil mercury concentrations in Southern California watersheds, *Water Air Soil Poll.*, 212, 369–385, doi:10.1007/s11270-010-0351-y, 2010a.
- Burke, M. P., Hogue, T. S., Ferreira, M., Mendez, C. B., Navarro, B., Lopez, S., and Jay, J. A.: The effect of wildfire on soil mercury concentrations in Southern California watersheds, *Water Air Soil Poll.*, 212, 369–385, doi:10.1007/s11270-010-0351-y, 2010b.
- Burke, M. P., Hogue, T. S., Kinoshita, A. M., Barco, J., Wessel, C., and Stein, E. D.: Pre- and post-fire pollutant loads in an urban fringe watershed in Southern California, *Environ. Monit. Assess.*, 185, 10131–10145, doi:10.1007/s10661-013-3318-9, 2013.
- Burles, K. and Boon, S.: Snowmelt energy balance in a burned forest plot, Crowsnest Pass, Alberta, Canada, *Hydrol. Process.*, 25, 3012–3029, doi:10.1002/hyp.8067, 2011.
- Cowpervait, P., Ocio, D., Collazos, G., de Cos, O., and Stocker, C.: Regionalised spatiotemporal rainfall and temperature models for flood studies in the Basque Country, Spain, *Hydrol. Earth Syst. Sci.*, 17, 479–494, doi:10.5194/hess-17-479-2013, 2013.
- Daly, C., Neilson, R. P., and Phillips, D. L.: A statistical topographic model for mapping climatological precipitation over mountainous terrain, *J. Appl. Meteorol.*, 33, 140–158, doi:10.1175/1520-0450(1994)033<0140:astmfm>2.0.co;2, 1994.
- Daly, C., Taylor, G., and Gibson, W.: The PRISM approach to mapping precipitation and temperature, 10th Conference on Applied Climatology, American Meteorological Society, Reno, NV, 10–12, 1997.
- Daly, C., Gibson, W. P., Taylor, G. H., Johnson, G. L., and Pasteris, P.: A knowledge-based approach to the statistical mapping of climate, *Clim. Res.*, 22, 99–113, doi:10.3354/cr022099, 2002.
- DiMiceli, C. M., Carroll, M. L., Sohlberg, R. A., Huang, C., Hansen, M. C., and Townshend, J. R. G.: Annual Global Automated MODIS Vegetation Continuous Fields (MOD44B) at 250 m Spatial Resolution for Data Years Beginning Day 65, 2000–2010, Collection 5 Percent Tree Cover, University of Maryland, College Park, 2011.
- Dozier, J.: Spectral signature of alpine snow cover from the Landsat Thematic Mapper, *Remote Sens. Environ.*, 28, 9–22, doi:10.1016/0034-4257(89)90101-6, 1989.

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Dozier, J., Painter, T. H., Rittger, K., and Frew, J. E.: Time-space continuity of daily maps of fractional snow cover and albedo from MODIS, *Adv. Water Resour.*, 31, 1515–1526, doi:10.1016/j.advwatres.2008.08.011, 2008.

Ebel, B. A., Hinckley, E. S., and Martin, D. A.: Soil-water dynamics and unsaturated storage during snowmelt following wildfire, *Hydrol. Earth Syst. Sci.*, 16, 1401–1417, doi:10.5194/hess-16-1401-2012, 2012.

Essery, R., Pomeroy, J., Parviainen, J., and Storck, P.: Sublimation of snow from coniferous forests in a climate model, *J. Climate*, 16, 1855–1864, doi:10.1175/1520-0442(2003)016<1855:sosfcf>2.0.co;2, 2003.

Faria, D. A., Pomeroy, J. W., and Essery, R. L. H.: Effect of covariance between ablation and snow water equivalent on depletion of snow-covered area in a forest, *Hydrol. Process.*, 14, 2683–2695, doi:10.1002/1099-1085(20001030)14:15<2683::aid-hyp86>3.0.co;2-n, 2000.

Gleason, K. E., Nolin, A. W., and Roth, T. R.: Charred forests increase snowmelt: effects of burned woody debris and incoming solar radiation on snow ablation, *Geophys. Res. Lett.*, 40, 4654–4661, doi:10.1002/grl.50896, 2013.

Hall, D. K., Riggs, G. A., and Salomonson, V. V.: Development of methods for mapping global snow cover using Moderate Resolution Imaging Spectroradiometer data, *Remote Sens. Environ.*, 54, 127–140, doi:10.1016/0034-4257(95)00137-p, 1995.

Harpold, A. A., Biederman, J. A., Condon, K., Merino, M., Korgaonkar, Y., Nan, T., Sloat, L. L., Ross, M., and Brooks, P. D.: Changes in snow accumulation and ablation following the Las Conchas Forest Fire, New Mexico, USA, *Ecohydrology*, 7, 440–452, doi:10.1002/eco.1363, 2013.

Kattelman, R. C., Berg, N. H., and Rector, J.: The potential for increasing streamflow from Sierra-Nevada watersheds, *Water Resour. Bull.*, 19, 395–402, 1983.

Kinoshita, A. M. and Hogue, T. S.: Spatial and temporal controls on post-fire hydrologic recovery in Southern California watersheds, *Catena*, 87, 240–252, doi:10.1016/j.catena.2011.06.005, 2011.

Klein, A. G., Hall, D. K., and Riggs, G. A.: Improving snow cover mapping in forests through the use of a canopy reflectance model, *Hydrol. Process.*, 12, 1723–1744, doi:10.1002/(sici)1099-1085(199808/09)12:10/11<1723::aid-hyp691>3.0.co;2-2, 1998.

Lane, P. N. J., Sheridan, G. J., and Noske, P. J.: Changes in sediment loads and discharge from small mountain-catchments following wild-fire in south eastern Australia, *J. Hydrol.*, 331, 495–510, doi:10.1016/j.jhydrol.2006.05.035, 2006.

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Smakhtin, V. U.: Low flow hydrology: a review, *J. Hydrol.*, 240, 147–186, doi:10.1016/s0022-1694(00)00340-1, 2001.

Stednick, J. D.: Monitoring the effects of timber harvest on annual water yield, *J. Hydrol.*, 176, 79–95, doi:10.1016/0022-1694(95)02780-7, 1996.

5 Stephens, S. L., Collins, B. M., and Roller, G.: Fuel treatment longevity in a Sierra Nevada mixed conifer forest, *Forest Ecol. Manage.*, 285, 204–212, doi:10.1016/j.foreco.2012.08.030, 2012.

Swanson, F. J.: Fire and Geomorphic Processes. in: Proceedings, Fire regimes and ecosystems conference, Honolulu, HI, 11–15 December 1979, Gen. Tech. Rep. WO-23, USDA, Washington, D.C., 401–420, 1981.

10 Tucker, C. J.: Red and photographic infrared linear combinations for monitoring vegetation, *Remote Sens. Environ.*, 8, 127–150, 1979.

USDA Forest Service Remote Sensing Applications Center (RSAC): Moonlight Fire occurring on the Plumas National Forest – 2007, available at: <http://edc.usgs.gov>, US Geol. Surv., Sioux Falls, South Dakota USA, 2007.

15 Varhola, A., Coops, N. C., Weiler, M., and Moore, R. D.: Forest canopy effects on snow accumulation and ablation: an integrative review of empirical results, *J. Hydrol.*, 392, 219–233, doi:10.1016/j.jhydrol.2010.08.009, 2010.

Webb, A. A., Kathuria, A., and Turner, L.: Longer-term changes in streamflow following logging and mixed species eucalypt forest regeneration: the Karuah experiment, *J. Hydrol.*, 464, 412–422, doi:10.1016/j.jhydrol.2012.07.034, 2012.

20 Westerling, A. L., Hidalgo, H. G., Cayan, D. R., and Swetnam, T. W.: Warming and earlier spring increase western US forest wildfire activity, *Science*, 313, 940–943, doi:10.1126/science.1128834, 2006.

25 Wildland Fire Incidents: US Historic Fire Perimeters. Geospatial Data Presentation Form: vector digital data, available at: www.geomac.gov, last access: 1 October 2013.

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

Table 1. Domain attributes for Moonlight Fire and Grizzly Ridge.

Domain Attributes	2007 Moonlight Fire	Grizzly Ridge
Area [ha]	27 370	14 800
Elevation Range [m]	1090–2290	1300–2320
Average Annual Precipitation [mm]	680	880
NLCD Land Cover		
Evergreen Forest	89 %	78 %
Shrub/Scrub	9 %	21 %
*Misc.	2 %	1 %
Soil Burn Severity		
High	37 %	N/A
Moderate	18 %	N/A
Low – Unburned	45 %	N/A
Slope Aspect		
North	21 %	17 %
South	33 %	42 %
East	20 %	16 %
West	26 %	25 %

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 2. Length of snow season compared to total winter precipitation for Moonlight Fire and Grizzly Ridge. Post-fire years are in italic.

Moonlight Fire	Length of Snow Season [Days]	Total Winter Precipitation [mm]
WY 2002	100	610
WY 2003	120	890
WY 2004	90	760
WY 2005	160	710
WY 2006	140	1000
WY 2007	60	410
WY 2008	<i>120</i>	<i>450</i>
WY 2009	<i>130</i>	<i>560</i>
WY 2010	<i>140</i>	<i>590</i>
WY 2011	<i>170</i>	<i>820</i>
WY 2012	<i>100</i>	<i>380</i>

Grizzly Ridge	Length of Snow Season [Days]	Total Winter Precipitation [mm]
WY 2002	90	780
WY 2003	120	970
WY 2004	110	800
WY 2005	150	990
WY 2006	130	1300
WY 2007	60	560
WY 2008	140	610
WY 2009	140	790
WY 2010	160	870
WY 2011	180	1200
WY 2012	60	520

Table 3. ANOVA results based on basin attributes for Moonlight Fire. Bold font denotes statistical significance ($P < 0.01$), post-fire years are in italic.

Slope Aspect	South [fSCA]	North [fSCA]	West [fSCA]	East [fSCA]
WY 2002	0.13	0.12	0.12	0.13
WY 2003	0.12	0.10	0.11	0.13
WY 2004	0.11	0.10	0.10	0.11
WY 2005	0.18	0.17	0.17	0.19
WY 2006	0.16	0.15	0.15	0.17
WY 2007	0.08	0.08	0.08	0.09
Pre-Fire Average	0.13	0.12	0.12	0.14
WY 2008	<i>0.22</i>	<i>0.22</i>	<i>0.21</i>	<i>0.23</i>
WY 2009	<i>0.23</i>	<i>0.22</i>	<i>0.21</i>	<i>0.23</i>
WY 2010	<i>0.29</i>	<i>0.28</i>	<i>0.27</i>	<i>0.29</i>
WY 2011	<i>0.29</i>	<i>0.29</i>	<i>0.27</i>	<i>0.32</i>
WY 2012	<i>0.15</i>	<i>0.16</i>	<i>0.14</i>	<i>0.16</i>
Soil Burn Severity	High [fSCA]	Moderate [fSCA]	Low-Unburned [fSCA]	
WY 2002	0.11	0.13	0.15	
WY 2003	0.09	0.12	0.14	
WY 2004	0.09	0.11	0.12	
WY 2005	0.16	0.18	0.20	
WY 2006	0.14	0.16	0.18	
WY 2007	0.07	0.08	0.10	
Pre-Fire Average	0.11	0.13	0.15	
WY 2008	<i>0.23</i>	<i>0.22</i>	<i>0.21</i>	
WY 2009	<i>0.24</i>	<i>0.22</i>	<i>0.21</i>	
WY 2010	<i>0.30</i>	<i>0.28</i>	<i>0.27</i>	
WY 2011	<i>0.30</i>	<i>0.29</i>	<i>0.27</i>	
WY 2012	<i>0.15</i>	<i>0.15</i>	<i>0.15</i>	

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 1. Map of Moonlight Fire with soil burn severity and control basin, Grizzly Ridge.

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.




Figure 2. Pre- and post-fire MOD10A1 fSCA (left panel) and non-canopy adjusted MODSCAG fSCA (right panel) difference maps for winter (January–March) over the Moonlight Fire. Each image contains 1099 pixels.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.




Figure 3. Least-squared linear regression analysis of MOD10A1 and non-adjusted MODSCAG over the Moonlight Fire pre- (black circles) and post-fire (red diamonds).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.




Figure 4. Timeseries of monthly precipitation totals, minimum and maximum temperatures and daily basin averaged MODSCAG fSCA for Moonlight Fire **(a)** and Grizzly Ridge **(b)** for WY 2002 to 2012.

[Title Page](#)

Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.




Figure 5. Annual cumulative frequency curves of daily basin averaged non-canopy adjusted MODSCAG fSCA for Moonlight Fire (a) and Grizzly Ridge (c) and canopy adjusted MODSCAG fSCA for Moonlight Fire (b) and Grizzly Ridge (d). Black lines with black circles represent extreme pre-fire fSCA years (highest and lowest annual curves) and red circles represent post-fire annual curves.

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.




Figure 6. Temporal trends in snow cover of the consecutive number of high snow cover days (pre-fire exceedance probability $\leq 10\%$; [black and red lines]) with respect to the length of snow season (exceedance probability $\geq 70\%$; [black and red crosses]) for Grizzly Ridge (**c** and **d**) and the Moonlight Fire (**a** and **b**). Color maps show annual daily basin averaged fSCA patterns. (**b**) and (**d**) are canopy adjusted MODSCAG fSCA.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

HESSD

11, 7513–7549, 2014

Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.




Figure 7. Basin averaged ANOVA results for Moonlight Fire (left panel) and Grizzly Ridge (right panel) (99 % confidence interval). The post-fire years are shaded for Moonlight Fire.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Application of MODIS snow cover products: wildfire impacts on snow and melt in the Sierra Nevada

P. D. Micheletty et al.




Figure 8. Basin averaged snow cover melt-out dates for Moonlight Fire and Grizzly Ridge (a). Relative difference in melt-out dates (Moonlight Fire – Grizzly Ridge) from the Moonlight Fire and Grizzly Ridge (b).

[Title Page](#)

[Abstract](#) | [Introduction](#)

[Conclusions](#) | [References](#)

[Tables](#) | [Figures](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)

[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

