



**Recharge estimation
and soil moisture
dynamics in a
Mediterranean karst
aquifer**

F. Ries et al.

**Recharge estimation and soil moisture
dynamics in a Mediterranean karst aquifer**

F. Ries^{1,2}, J. Lange¹, S. Schmidt², H. Puhmann³, and M. Sauter²

¹Chair of Hydrology, University of Freiburg, Freiburg, Germany

²Geoscience Center, Applied Geology, University of Göttingen, Göttingen, Germany

³Forest Research Institute Baden-Württemberg, Freiburg, Germany

Received: 3 July 2014 – Accepted: 14 July 2014 – Published: 28 July 2014

Correspondence to: F. Ries (fabian.ries@hydrology.uni-freiburg.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

Knowledge of soil moisture dynamics in the unsaturated soil zone provides valuable information on the temporal and spatial variability of groundwater recharge. This is especially true for the Mediterranean region, where a substantial fraction of long-term groundwater recharge is expected to occur during high magnitude precipitation events of above-average wet winters. To elucidate process understanding of infiltration processes during these extreme events, a monitoring network of precipitation gauges, meteorological stations, and soil moisture plots was installed in an area with a steep climatic gradient in the Jordan Valley region. In three soil moisture plots, Hydrus-1D was used to simulate water movement in the unsaturated soil zone with soil hydraulic parameters estimated by the Shuffled Complex Evolution Metropolis algorithm. To generalize our results, we modified soil depth and rainfall input to simulate the effect of the pronounced climatic gradient and soil depth variability on percolation fluxes and applied the calibrated model to a time series with 62 years of meteorological data.

Soil moisture measurements showed a pronounced seasonality and suggested rapid infiltration during heavy rainstorms. Hydrus-1D successfully simulated short and long-term soil moisture patterns, with the majority of simulated deep percolation occurring during a few intensive rainfall events. Temperature drops in a nearby groundwater well were observed synchronously with simulated percolation pulses, indicating rapid groundwater recharge mechanisms. The 62 year model run yielded annual percolation fluxes of up to 66 % of precipitation depths during wet years and of 0 % during dry years. Furthermore, a dependence of recharge on the temporal rainfall distribution could be shown. Strong correlations between depth of recharge and soil depth were also observed.

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



1 Introduction

In the Mediterranean region, groundwater is the main source for domestic and agricultural water supplies (EUWI, 2007). Knowledge on the quantity of groundwater recharge is a prerequisite for sustainable water resources planning and effective water use.

5 Small-scale differences in climate, geology, land use, topography and soil properties cause a high spatial and temporal variability of groundwater recharge making the assessment and predictions of recharge a challenge (e.g. Zagana et al., 2007). Karst areas are important in this respect, because during high intensity winter storms precipitation may rapidly infiltrate into exposed karst surfaces and induce high recharge rates (De Vries and Simmers, 2002), which are common in the Mediterranean area (Ford and Williams, 2007). A rapidly increasing water demand in the last decades has led to a widespread overexploitation of groundwater resources (EUWI, 2007). Furthermore, the Mediterranean region has been identified as a “hot spot” of current and future climate change (Giorgi, 2006; IPCC, 2013), imposing additional pressure on its limited water resources. Hence, more insights into processes of aquifer replenishment in Mediterranean karst regions are of vital importance.

A large variety of methods suitable for estimating recharge rates were developed in the last decades (De Vries and Simmers, 2002; Scanlon et al., 2002). Infiltration, percolation and recharge quantities in Mediterranean karst have mainly been approached from two sides: on the one hand, hydrologists and geomorphologists characterized the surface water balance on small plots by sprinkling experiments or by runoff/sediment measurements during natural rainstorms (e.g. Cerdà, 1998; Lavee et al., 1998). Large-scale experiments also included tracers and facilitated statements on runoff generation processes (e.g. Lange et al., 2003). However, these studies quantified infiltration by the difference between artificial/natural rainfall and measured overland flow but did not differentiate between recharge and evapotranspiration. On the other hand, hydrogeologists frequently assessed average recharge rates of entire karst catchments from spring discharge measurements or hydraulic head data. Methods include

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



knowledge (GIS)-based mapping (Andreo et al., 2008), multiple linear regression (Al-
loca et al., 2014), conceptual models (e.g. Hartmann et al., 2013a), coupled water-
balance groundwater models (Sheffer et al., 2010), and chloride mass balances (Marei
et al., 2010; Schmidt et al., 2013). However, these studies treat karst systems as units,
including both the unsaturated and the saturated zones, and are limited in temporal and
spatial resolution. Studies on cave drips (Gregory et al., 2009; Arbel et al., 2010) pro-
vided insights into the deeper unsaturated zone in terms of water storage, spatial vari-
ability of percolation and flow paths. Their data was also used to incorporate variability
in recharge modelling (Hartmann et al., 2012). However, it was difficult to distinguish
between processes in the unsaturated soil zone and in the underlying epikarst, and
uncertainty remains regarding the representativeness of cave drip data with respect to
infiltration processes. This is mainly due to the facts that the contributing areas of cave
drips are unknown and caves might have developed their own hydraulic environments.
Therefore cave drips are not necessarily representative for the bulk karst vadose zone
(Lange et al., 2010).

Only limited knowledge on recharge dynamics is available for the carbonate aquifer
system shared between the West Bank and Israel. First recharge estimates were based
on long-term spring discharge and groundwater well abstraction data (Goldschmidt
and Jacobs, 1958). Later, groundwater flow models were used to establish empiri-
cal rainfall–recharge relationships (Baida and Burstein, 1970; Guttman and Zukerman,
1995; Zukerman, 1999). Average recharge rates were assessed by a simple water bal-
ance approach (Hughes et al., 2008) and by a chloride mass balance (Marei et al.,
2010). Scheffer et al. (2010) coupled a water budget model with a groundwater flow
model for the entire western part of the carbonate aquifer and used spring discharge
and groundwater level data for calibration. They reported recharge rates ranging be-
tween 9% and 40% of annual rainfall and showed that the temporal distribution of
rainfall within the winter season had considerable effects on overall recharge rates.

Observations of soil moisture may offer unique insights into near surface hydrologi-
cal processes, because water fluxes are susceptible to conditions and properties of

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

the vadose soil zone across several scales (Vereecken et al., 2008). Yet, soil moisture is rarely measured in semi-arid areas and is seldom used for recharge estimation purposes. Scott et al. (2000) exemplified the potential of soil moisture time series to calibrate Hydrus-1D soil hydraulic parameters in southeastern Arizona. Their results demonstrated the high inter-annual variability of water fluxes in these environments where considerable percolation only occurs during above average wet years.

The objective of this study is to investigate the spatial and temporal variability of percolation, and hence groundwater recharge rates, for an Eastern Mediterranean carbonate aquifer. We use continuously recorded soil moisture data to calibrate one-dimensional water flow models (Hydrus-1D) with the Shuffled Complex Evolution Metropolis (SCEM) algorithm. The calibrated models are then used to assess spatial and temporal patterns of soil water percolation in a Mediterranean karst area, which is characterized by strong climatic gradients and variable soil depths.

2 Study area

Our study area is located on the western margin of the Jordan Rift Valley 25 km north-east of Jerusalem (Fig. 1). Precipitation shows a pronounced seasonality with cold fronts (mainly Cyprus lows) carrying moisture from the Mediterranean Sea during winter season from October to April (Goldreich, 2003). The topographic gradient from the mountain range (highest elevation: 1016 m a.s.l.) in the west to the Jordan Valley in the east results in a strong precipitation gradient. Long-term average annual precipitation decreases from 532 mm in Jerusalem (810 m a.s.l.) to 156 mm in Jericho (290 m b.s.l.) (Morin et al., 2009).

Outcropping geological formations consist of carbonate rocks of Cretaceous age (Begin, 1975). They are composed of fractured and highly permeable layers of limestone and dolomite alternating with marl and chalk layers of low permeability, often considered partial aquicludes (Weiss and Gvirtzman, 2007). In the southeast, Senonian Chalks form outcrops of low hydraulic conductivity (Rofe and Raffety, 1963). Soil

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



parent material consists of residual clay minerals from carbonate rock weathering and from the aeolian input of dust (silt and clay fraction) originating from the Sahara desert (Yaalon, 1997). Predominant soil types are Terra Rossa and Rendzina, both characterized by high clay contents. Rendzina soils contain carbonate in the soil matrix, are thinner and still show recent development, whereas Terra Rossa soils were formed under past climatic conditions (Shapiro, 2006). As a result of irregular carbonate weathering and heterogeneous topography, soil depth is highly variable. On the slopes, outcropping massive bedrock and loose rock fragments of different sizes alternate with soil pockets of variable dimensions, shapes, and depths (Fig. 2). Dissolution cracks and karst fissures are often filled with eroded soil material. In valley bottoms, fine textured alluvial soils (Vertisols) with soil depths up to several meters have developed. Shallow Brown Lithosols and loessial Arid Brown Soils dominate in the eastern, low-lying areas receiving less rainfall (Shapiro, 2006). In general, soils in the region have significantly been transformed by human activities such as land cultivation, terracing, and deforestation during the last 5000 years (Yaalon, 1997).

On hillslopes, annual plants and Mediterranean shrubs (predominantly *Sarcopoterium spinosum*) are the dominant vegetation types. They are used for extensive grazing by goats and sheep. South-facing slopes show a lower vegetation density and a higher proportion of bare soil and outcrops than north-facing slopes, where the presence of biogenic crusts was reported (Kutiel et al., 1998). Minor land use types consist of scattered built-up areas, olive plantations on terraced land and rainfed or partly irrigated agricultural land (annual and perennial crops, herbs and vegetables) in valley bottoms.

3 Material and methods

3.1 Hydrometeorological measurements

To capture the spatial variation of rainfall along the strong climatic gradient, we installed a rain gauge network consisting of 14 tipping buckets (RG3-M) connected to a HOBO pendant event data logger (Onset Computer Corporation) (Fig. 1). All gauges were calibrated before employment, maintained, and cleaned twice a year before and after the rainfall season. Temperature was measured at four climatic stations (Thies GmbH and Onset Computer Corporation). Additional rainfall and climatic data was obtained from the Israel Meteorological Service database (<http://www.data.gov.il/ims>) for long-term analysis. Additionally, groundwater levels and temperatures were recorded in a nearby well (Mini-Diver, Eijkelkamp).

3.2 Soil moisture measurements

Seven soil moisture plots were installed, each equipped with four capacitance soil moisture sensors (5TM/5TE, Decagon Devices Inc.). We paid attention that the plots did not receive lateral surplus water from upslope overland flow by placing them distant from rock outcrops and at locations with minimum slope. To minimize disturbance, we inserted the sensors vertically into the upslope wall of manually dug soil pits (depth between 35 and 80 cm). After installation, we refilled the pits with the parent soil material and compacted approximately to pre-disturbance bulk density. The probes were connected to data loggers (EM50, Decagon Devices Inc.), which were sealed by plastic bags and buried in the soil to avoid vandalism. We used the internal calibration function for mineral soils with a measurement accuracy of 4 % of the volumetric water content (VWC). The measurement interval was set at ten minutes. Further information on the performance of the employed sensors can be found in Kizito et al. (2008). Due to instrument malfunction and vandalism, we obtained continuous data of our entire

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



measurement period (October 2011 to May 2013) from only three locations (SM1–SM3). Characteristics of the plots are summarized in Table 1.

The dielectric permittivity of water changes with temperature (e.g. Wraith and Or, 1999). Hence, measurement techniques of soil moisture based on the difference of dielectric permittivity between water and soil matrix are affected by this phenomenon. In our case, soil temperature was highly variable and changed by up to 20 °C within 24 h due to a strong radiation input and partly uncovered soil. We corrected our soil moisture data applying multiple linear regressions against soil temperature as described by Cobos and Campell (2007).

3.3 Modelling of the soil zone

Water balance at the plot scale in absence of surface runoff can be described with:

$$\frac{ds}{dt} = P - E_a - L \quad \text{with} \quad E_a = E_i + E_s + E_t, \quad (1)$$

where ds/dt is the storage change over time, P is the precipitation, L is the percolation at the profile bottom and E_a is the evapotranspiration per time interval. E_a is composed of the terms E_i (evaporation of intercepted precipitation), E_s (soil evaporation) and E_t (plant transpiration).

For our three soil moisture plots, soil water content and water fluxes were simulated on a daily basis with Hydrus-1D (version 4.16; Šimůnek et al., 2013) for a period of 32 months. Hydrus-1D solves the Richards equation numerically for water transport in variable saturated media. Matric potential dependent water retention and hydraulic conductivity were calculated using the Mualem/van-Genuchten soil hydraulic model (van Genuchten, 1980). To reduce the effect of non-linearity of the hydraulic conductivity function close to saturated conditions, an air entry value of –2 cm as suggested by Vogel et al. (2001) was used. Interception by the plant canopy was calculated by an empirical equation including the leaf area index and daily precipitation values (see Šimůnek et al., 2013 for more details). Potential evapotranspiration was calculated

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by the Hargreaves-equation (Hargreaves and Samani, 1985). Originally developed for a lysimeter station in California, this method adequately reproduced potential evapotranspiration under semi-arid climates (Jensen et al., 1997; Weiß and Menzel, 2008). Potential evapotranspiration was split into potential evaporation from the soil surface and potential transpiration from plants according to Beer's law based on the time variable surface cover fraction. Both fluxes were reduced to actual values based on a root water uptake model (Feddes et al., 1978) applying plant parameters for grass and an energy balance surface evaporation model (Camillo and Gurney, 1986). In our study area, vegetation cover shows a strong seasonality due to the restricted water availability during the dry season. To account for this, time dependent plant growth data was implemented into the model with intra-annual variation of surface cover fraction. According to field observations, the start of the growing season was set to mid November and the maximum vegetation density was assumed for February/March shortly after the largest monthly precipitation amounts were observed. The depth from which plants took up water was controlled by a root distribution function. An exponential decrease of root density with soil depth was assumed, observed at the study sites and often reported for the Mediterranean region (e.g. De Rosnay and Polcher, 1998; De Baets et al., 2008). Temporal variations of rooting depth and root density were disregarded. With these components, Hydrus-1D continuously computed water content and water fluxes at user defined observation points (here: depths of the soil moisture probes) and at the lower profile boundary. Model input data, selected parameter values and their ranges, and the corresponding data sources and calculation methods are summarized in Table 2.

3.4 Calibration procedure, uncertainty analysis and parameter sensitivity

An increase of clay content and bulk density with depth was observed at all profiles and the individual probes in various depths at our plots differed noticeably. As a result, a particular soil material with singular soil hydraulic properties was assigned for each soil moisture probe. Observed soil moisture data from two winter and one summer season

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



(October 2011 to April 2013) were used for calibration of Hydrus-1D. We individually determined soil hydraulic parameters for every soil material by inverse modelling using the Shuffled Complex Evolution Metropolis optimization algorithm (SCEM; Vrugt et al., 2003) and the Kling–Gupta efficiency (KGE; Gupta et al., 2009) in a modified version
5 from Kling et al. (2012) as the objective function:

$$\text{KGE} = 1 - \sqrt{(r - 1)^2 + (\alpha - 1)^2 + (\beta - 1)^2} \quad (2)$$

with:

$$r = \frac{\text{Cov}_{\text{so}}}{\sigma_s \cdot \sigma_o}, \quad \alpha = \frac{\mu_s}{\mu_o} \quad \text{and} \quad \beta = \frac{\sigma_s / \mu_s}{\sigma_o / \mu_o},$$

where r is the correlation coefficient between simulated and observed VWC (Cov_{so} is the covariance between simulated and observed VWC), α is a dimensionless measure for the bias (μ_s and μ_o are the mean simulated and observed VWC) and β is a dimensionless measure for variability (σ_s and σ_o are the standard deviations of simulated and observed VWC). SCEM is widely used to efficiently solve global optimization problems
15 (e.g. Vrugt et al., 2005; Schoups et al., 2005; Feyen, 2007; Hartmann et al., 2012) and to find optimal model parameter sets. As algorithmic parameters for SCEM, 24 complexes/parallel sequences were selected (equal to the number of parameters to be optimized), the population size was set to 144 and the number of accepted draws to infer posterior distribution was set to 1000. The SCEM routine was run until the scale reduction score (SR), a convergence criterion defined by Gelman and Rubin (1992), was fulfilled. As proposed by Vrugt et al. (2003), a SR value of 1.2 was chosen, indicating that the Markov chain had converged to a stationary posterior distribution for all parameters. Predicted soil moisture ranges were used for parameter uncertainty
20 assessment. They were determined by running Hydrus-1D with 1000 parameter sets obtained through the SCEM algorithm after reaching convergence.

Recharge estimation
and soil moisture
dynamics in a
Mediterranean karst
aquifer

F. Ries et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.5 Spatial and temporal extrapolation of percolation

To extrapolate our point measurements of soil water balance, we varied soil depth and climatic input parameters (precipitation and temperature) over ranges observed in our study area. We used the calibrated soil hydraulic parameters of our deepest (1 m) soil moisture plot (SM-1), which had sensor at 10, 25, 40 and 80 cm. Moreover, we assumed that the rooting depth was limited to the soil depth with no changes in the vertical root distribution or plant surface cover fraction. We cut off the profile according to the simulated soil depth, which reduced the number of independent soil layers when the depths fell below 60, 32.5 and 17.5 cm. For soil thicknesses exceeding 1 m, we extended the bottom layer. To simulate the range of climatic conditions with elevations between 400 and 1000 m a.s.l., we modified rainfall and air temperature according to calculated mean annual gradients based on observed rainfall and climatic data. We had three seasons of measured climate data, which we analysed separately due to seasonal differences in cumulative rainfall amount and distribution.

Using a 62 year record of rainfall and temperature (1951–2013) available for Jerusalem (Israel Meteorological Service – www.data.gov.il/ims), we assessed the annual variability of water balance components at the location of our three soil moisture plots. Rainfall and temperature data from Jerusalem station were corrected for elevation differences between the Jerusalem station (810 m a.s.l.) and the three plots based on calculated elevation gradients.

4 Results

4.1 Hydrometeorological conditions

The three years of high resolution measurements of precipitation and meteorological parameters revealed considerable interannual variability and a strong elevation gradient, especially in terms of rainfall. Mean annual precipitation at the Kafr Malek

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



station (810 m a.s.l.) situated close to the Mediterranean Sea–Dead Sea water divide was 526 mm (380–650 mm), while mean annual rainfall at the Auja Village station (270 m b.s.l.) in the Jordan Valley accounted for 106 mm (97–120 mm) leading to seasonal rainfall gradients between 6.4 to 7.2% per 100 m elevation difference (Fig. 3).

5 Mean annual temperature was 7 °C higher at Auja Village whereas relative humidity, wind speed, and net solar radiation were slightly higher at the more elevated station. Stations from the Israel Meteorological Service with long-term records at locations nearby showed similar characteristics.

4.2 Soil moisture dynamics

10 Observed soil moisture at all soil profiles (Fig. 4) showed a strong seasonality where the annual course can be divided into distinct phases. At the beginning of the rainy season, the previously dry soil profile was stepwise wetting up starting from upper to lower sensors. During rainfall events with high amounts and intensities, the soil moisture data showed rapid infiltration of water into the deeper portions of the profile. Particularly at
15 plot SM-1, saturated conditions started from the bottom probe close to the soil–bedrock interface, where these conditions persisted for several hours up to two days. During the strongest rainfall events also upper soil layers reached saturation, however for much shorter periods (Fig. 4b). At the end of the rainy season, the soil was drying out within
20 a few weeks and the soil moisture content further declined at a low rate during the whole dry summer period.

4.3 Modelling of the soil zone

4.3.1 Parameter optimization, uncertainty analysis and model validation

25 Soil hydraulic parameters were optimized for the three soil moisture plots individually, using the Shuffled Complex Evolution Metropolis algorithm. Between 20 000 and 36 000 model runs were conducted until the convergence criterion was fulfilled. The

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

calibrated parameter sets used for further assessment of the plot scale soil water balance, are given in Table 3, and their distributions are illustrated in Fig. 5. All models were generally able to reproduce the observed temporal soil moisture patterns with KGE values between 0.82 and 0.94 (Fig. 6). However, differences in predictive capacities at distinct water content levels could be observed, which vary between the single plots (Figs. 6 and 7). In general, the model tended to overestimate water contents close to saturated conditions except for deeper sections at plot SM-1 where an underestimation of simulated water contents was observed.

Parameter uncertainty was assessed by simulation of water contents using parameter sets obtained with SCEM after fulfilling the convergence criterion. The 95 % soil moisture confidence interval showed a narrow band around the optimum model (Fig. 8 exemplary for plot SM-1). At all sensors the difference between simulated volumetric water content for the best parameter set and the 95 % confidence interval remained below 4 %, i.e. less than the measurement error of the sensors.

Water temperature in a groundwater well near soil moisture plot SM-3 (cf. Fig. 1) indicated five distinct recharge events lowering the mean groundwater temperature from 19 °C by 0.7–4 °C (Fig. 7). These events coincided with the main peaks of modelled percolation from the soil moisture monitoring sites. During these events, mean daily air temperature was less than 6 °C.

4.3.2 Plot scale water balance

Modelled fluxes of the various water balance components showed high temporal variability (Fig. 8) and considerable differences in annual values between single years (Table 4). Evaporation and transpiration started shortly after the first rainfall events of the winter season when the water content in the upper soil layer began to increase. Percolation from the bottom of the soil zone only started after cumulative rainfall during winter season exceeded a certain threshold. This threshold was found to be ca. 240 mm at plot SM-1, 200 mm at plot SM-2, and 150 mm at plot SM-3. The relative proportion of interception, soil evaporation, and transpiration was highly variable during

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the winter season, depending on the length of dry spells between rainfall events and ceased within few weeks after the last rainfall events of the winter season. Mean overall losses through evapotranspiration and interception accounted for 73 % of rainfall. Values slightly above 100 % for the dry year 2010/11 resulted from elevated moisture conditions at the beginning of the simulation period. Percolation strongly varied from negligible amounts during the dry year 2010/11 to values ranging between 28 % and 45 % of cumulative rainfall during 2011/12 and 2012/13, respectively. The largest proportion of percolation was calculated during a few strong rainstorms. On all three plots, more than 50 % of the total percolation of the three years simulation period occurred within a time period of five to ten days.

4.3.3 Spatial extrapolation of deep percolation

During the hydrological year 2010/11, cumulative rainfall was below average with totals ranging between 275 and 425 mm (Fig. 9) and a maximum daily amount below 50 mm. In this season with below average rainfall amounts, percolation was only simulated for soils with depths up to 60 and 110 cm, respectively. Modelled percolation increased to a maximum proportion of 40 % for shallow soils with depths of 10 cm receiving the highest rainfall input. For the following above average wet year 2011/12, seasonal rainfall ranged between 450 and 725 mm. Then simulated percolation rates reached up to 69 % of rainfall and declined to values close to 0 % only under conditions of lowest rainfall amount and soil depths greater than 160 cm. The third simulated year can be regarded as a year with average rainfall conditions (sums of 400 to 600 mm). Percentages of percolation were comparable to the previous year although cumulative rainfall was considerably less. This could be attributed to higher rainfall intensities during 2012/13 when daily rainfall amounts exceeded two times 80 mm and four days of rainfall accounted for almost 50 % of the seasonal amount.

4.3.4 Temporal extrapolation of deep percolation

Modelling water balance components for 62 years (1951–2013) resulted in strong differences of simulated seasonal percolation reflecting the high variability of rainfall input (Fig. 10). Mean annual rainfall was calculated for the three plots to range between 408 and 537 mm (standard deviation: 128–168 mm) and mean percolation fluxes between 82 and 150 mm (standard deviation: 93–141 mm). Percolation at the three plots varied between 0 and 66 % of cumulative seasonal rainfall with an average between 16 and 24 %. Other seasonal fluxes varied much less during the simulation period. The coefficient of determination between seasonal sums of simulated percolation and rainfall ranged between 0.82 and 0.88 on the three plots.

5 Discussion

5.1 Soil moisture dynamics

The observed seasonal dynamics of soil moisture, dominated by short wetting phases during and a rapid decrease after the rainfall season, were comparable with those reported in other studies in the Mediterranean region (Cantón et al., 2010; Ruiz-Sinoga et al., 2011). At all soil moisture plots, our soil moisture data suggested fast infiltration into deeper sections of the soil profile during rainfall events with high intensities and amounts (e.g. plot SM-1 in Fig. 4b). The time lag between the reaction of the uppermost and the lowermost probe was often less than two hours despite high clay contents of the soils. These fast reactions might be an indicator for concentrated infiltration and preferential flow within the vadose soil zone as reported for the Mediterranean by Cerdá et al. (1998), Öhrström et al. (2002) and Van Schaik et al. (2008). Nevertheless, larger macropores were rarely observed in the field and superficial desiccation cracks that appeared during dry summer months closed soon after the beginning of the rainfall season. Moreover, a successive propagation of the wetting front was observed at the

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

soil hydraulic model based on a unimodal pore-size distribution may not be able to represent the heterogeneous pore structure of our clay-rich soil (Durner, 1994). Moreover, persistent saturated conditions during major rainstorms as discussed in the previous section could not be simulated, as a percolation impounding soil–rock interface was not implemented in the model and a free drainage had to be assumed.

Simulated mean evapotranspiration at our plots over the three-years simulation period accounted for 73 % of rainfall, i.e. very close to the long-term average calculated by Schmidt et al. (2014) for the same area. Our values also fall into the range of Cantón et al. (2010), who derived annual effective evapotranspiration rates of more than 64 % of annual rainfall based on eddy covariance measurements in southeastern semi-arid Spain. Our simulated percolation rates ranged between 0 and 45 % of precipitation (arithmetic mean: 28 %) indicating strong inter-annual variability and a strong dependency on depth and temporal distribution of precipitation. During the entire three year period, more than 50 % of overall percolation fluxes occurred during less than 10 days of strong rainfall. These findings are supported by the response of groundwater temperatures observed in a nearby well indicating the arrival of groundwater recharge at the water table (Fig. 8). Tracer experiments in a similar setting demonstrated that percolating water can pass the vadose soil and the epikarst at flow velocities of up to 4.3 m h^{-1} (Lange et al., 2010). Regarding the initiation of percolation at the basis of the soil profiles, we found seasonal rainfall thresholds of ca. 150 mm for the shallow and 240 mm for the deep soil moisture plots. Cave drip studies in the region (Arbel et al., 2010; Lange et al., 2010; Sheffer et al., 2011) measured similar thresholds for the initiation of percolation through the epikarst (100 to 220 mm).

In contrast to humid environments, lateral subsurface flow on rocky semi-arid hillslopes rarely develops, since they consist of individual soil pockets that are poorly connected due to frequently outcropping bedrocks. Soil moisture seldom exceeds field capacity given that evapotranspiration exceeds precipitation depth throughout most of the year (Puigdefabregas et al., 1998). Furthermore, highly permeable bedrocks favour the development of vertical structural pathways in karst areas (dolines, sinkholes) inducing

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

concentrated infiltration from the soil zone (Williams, 1983). We can therefore conclude that one-dimensional modelling of the soil water balance is a reasonable approach to understand percolation and recharge.

Nevertheless, we cannot exclude that frequently outcropping bedrock may affect water redistribution by surface runoff or by preferential infiltration along the soil–rock interface. The importance of these effects on percolation rates and groundwater recharge on the regional scale is subject to current research. During heavy storm events, overland flow generation cannot be excluded (Lange et al., 2003), but surface runoff typically accounts for only a few percent of annual rainfall (Gunkel and Lange, 2012). A second limitation of our investigations of plot scale percolation fluxes is the assumption of an identical vegetation cover at the single sites along the climatic gradient and a constant vegetation cycle throughout years of different seasonal rainfall depths. Although different plant species and vegetation cycles may alter soil moisture conditions prior to rainfall events, we could show that the event rainfall amount is the main factor that influences percolation rates.

5.3 Spatial and temporal extrapolation of deep percolation

Water balance modelling for variable soil depths and rainfall gradients revealed considerable differences for the three winter seasons. During the very dry year 2010/11, soil moisture exceeded field capacity only at locations with relatively shallow soils. During the wet years of 2011/12 and 2012/13, field capacity was exceeded several times at all plots and soils even reached saturation during strong rainfall events. This may lead to substantial percolation and groundwater recharge to local aquifers. These findings are in good agreement with discharge measurements at Auja spring, a large karst spring in the Jordan Valley, where 7 and 8 million m³ were measured for the winter seasons 2011/12 and 2012/13 respectively, but only 0.5 million m³ for the 2010/11 season (Schmidt et al., 2014).

A high temporal variability in percolation fluxes is also apparent from the long-term modelling of water balance components (Fig. 11). For the 62 year simulation period,

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

we calculated seasonal percolation rates between 0 and 66 % (average: 20 to 28 %) for our plots. The highest value was modelled for the extremely wet winter season 1991/92 (five times the mean annual percolation of 150 mm). For a slightly shorter time period, Schmidt et al. (2014) calculated an average recharge rate of 33 % for the Auja spring catchment with a conceptual reservoir model. They found that recharge of only five individual years accounted for one third of total recharge of the 45 years period. In our study seven individual years provided one third of the total recharge. Furthermore, we compared seasonal percolation of our sites with recharge estimations from perched aquifers feeding small karst springs (Weiss and Gvirtzman, 2007) and the entire carbonate aquifer (Guttman und Zukerman, 1995) (Fig. 11). Although our results plotted within the range of these large-scale recharge estimates, we want to emphasize that our calculations display point percolation fluxes. Even in years with below average rainfall, a certain rise in the groundwater table and spring flow can be observed (EXACT, 1998; Schmidt et al., 2014). Then recharge presumably occurs on areas with strongly developed epikarst and shallow or missing soil cover.

Our long-term point calculations suggest substantial differences in percolation fluxes between years of similar rainfall depths. Simulated percolation for plot SM-1 during the seasons 1976/77 and 2004/05 accounted for 16 and 35 % of seasonal rainfall, respectively, although both seasons had very similar above average rainfall (578 and 569 mm). These results are in line with findings of Sheffer et al. (2010) and Abusaada (2011) about the importance of temporal rainfall distribution on groundwater recharge.

5.4 Implications for recharge in Mediterranean karst areas

The steep climatic gradient, the hydraulic properties and characteristics of the carbonate rocks, the heterogeneous soil cover and a high temporal variability of precipitation on event and seasonal scales are dominating hydrological characteristics in our study area. Similar settings can be found across the entire Mediterranean region. Despite recent advances in the determination of groundwater recharge in karst areas, the assessment of the spatial and temporal distribution of recharge is still a challenge. Modelling

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

approaches including hydrochemical and isotopic data (Hartmann et al., 2013b) require additional information from springs (time series of discharge and water chemistry) for model parameter estimation, which are rarely available. Although simulated percolation fluxes from plot-scale soil moisture measurements cannot be directly transferred to regional, i.e. catchment scale, they can provide insight into the various processes responsible for the temporal and spatial variability of groundwater recharge as well as information on the relative importance of different process parameters.

6 Conclusions

This study contributes to the assessment of percolation rates based on soil moisture measurements along a steep climatic gradient in a Mediterranean karst area. We showed that soil moisture measurements together with numerical modelling of the water flow in the unsaturated soil zone are powerful tools to investigate mechanisms and characteristics determining the strong annual variability in the percolation fluxes and the strong dependency on soil thickness, temporal distribution of rainfall and precipitation depth. Although our calculations are based on plot scale measurements, the results closely match long-term observations and the patterns of event and seasonal variability, and reflect the thresholds for the initiation of groundwater recharge reported by other studies in the same region based on different approaches. Our results suggest that groundwater recharge is most prominent when single rainfall events are strong enough to exceed field capacity of soil pockets over a wide range of soil depths. Furthermore, our data revealed that the temporal distribution of rainfall had a strong effect on event and seasonal recharge amounts.

We therefore corroborate the statement of De Vries and Simmers (2002) about the dependence of groundwater recharge in semi-(arid) areas on high intensity rainfall events. The use of empirical rainfall–recharge relationships can lead to large errors, since recharge rates are sensitive with respect to highly variable rainfall distributions and characteristics, which are most probably affected by predicted climate change in

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Begin, Z. B.: The geology of the Jericho sheet, Geological Survey of Israel, Bulletin No. 67, Jerusalem, 1975.
- Booltink, H. W. G. and Bouma, J.: Physical and morphological characterization of bypass flow in a well-structured clay soil, *Soil Sci. Soc. Am. J.*, 55, 1249–1254, 1991.
- 5 Camillo, P. J. and Gurney, R. J.: A resistance parameter for bare-soil evaporation models, *Soil Sci.*, 141, 742–744, 1986.
- Cantón, Y., Villagarcía, L., Moro, M. J., Serrano-Ortíz, P., Were, A., Alcalá, F. J., Kowalski, A. S., Solé-Benet, A., Lázaro, R., and Domingo, F.: Temporal dynamics of soil water balance components in a karst range in southeastern Spain: estimation of potential recharge, *Hydrolog. Sci. J.*, 55, 737–753, 2010.
- 10 Cerdà, A.: Effect of climate on surface flow along a climatological gradient in Israel: a field rainfall simulation approach, *J. Arid Environ.*, 38, 145–159, 1998.
- Cerdà, A., Schnabel, S., Ceballos, A., and Gomez-Amelia, D.: Soil hydrological response under simulated rainfall in the Dehesa land system (Extremadura, SW Spain) under drought conditions, *Earth Surf. Proc. Land.*, 23, 195–209, 1998.
- 15 Cobos, D. R. and Campbell, C.: Correcting temperature sensitivity of ECH2O soil moisture sensors, Application Note #800-755-2751, Decagon Devices, Pullman, WA, 2007.
- Cousin, I., Nicoulaud, B., and Coutadeur, C.: Influence of rock fragments on the water retention and water percolation in a calcareous soil, *Catena*, 53, 97–114, 2003.
- 20 Dan, J.: The distribution and origin of Nari and other lime crusts in Israel, *Israel J. Earth Sci.*, 26, 68–83, 1977.
- De Baets, S., Poesen, J., Reubens, B., Wemans, K., De Baerdemaeker, J., and Muys, B.: Root tensile strength and root distribution of typical Mediterranean plant species and their contribution to soil shear strength, *Plant Soil*, 305, 207–226, 2008.
- 25 de Rosnay, P. and Polcher, J.: Modelling root water uptake in a complex land surface scheme coupled to a GCM, *Hydrol. Earth Syst. Sci.*, 2, 239–255, doi:10.5194/hess-2-239-1998, 1998.
- De Vries, J. J. and Simmers, I.: Groundwater recharge: an overview of processes and challenges, *Hydrogeol. J.*, 10, 5–17, 2002.
- 30 Durner, W.: Hydraulic conductivity estimation for soils with heterogeneous pore structure, *Water Resour. Res.*, 30, 211–223, 1994.
- EUWI – European Water Initiative: Mediterranean groundwater report, technical report on groundwater management in the Mediterranean and the Water Framework Directive,

Produced by the Mediterranean Groundwater Working Group, available at: www.semide.net/initiatives/medeuwi/JP/GroundWater (last access: 9 April 2014), 2007.

EXACT – Executive Action Team: Overview of Middle East Water Resources – Water Resources of Palestinian, Jordanian, and Israeli Interest, compiled by the US Geological Survey for the Executive Action Team, United States Geological Survey, Washington, 44 pp., ISBN: 0-607-91785-7, 1998.

Feddes, R. A., Kowalik, P. J., and Zaradny, H.: Simulation of field water use and crop yield, Simulation Monographs, PUDOC, Wageningen, 189 pp., 1978.

Feyen, L., Vrugt, J. A., Nualláin, B. Ó., van der Knijff, J., and De Roo, A.: Parameter optimisation and uncertainty assessment for large-scale streamflow simulation with the LISFLOOD model, *J. Hydrol.*, 332, 276–289, 2007.

Ford, D. C. and Williams, P. W.: Karst Hydrogeology and Geomorphology, Wiley, Chichester, 2007.

Gelman, A. and Rubin, D. B.: Inference from iterative simulation using multiple sequences, *Stat. Sci.*, 7, 457–472, 1992.

Giorgi, F.: Climate change hot-spots, *Geophys. Res. Lett.*, 33, L08707, doi:10.1029/2006GL025734, 2006.

Giorgi, F. and Lionello, P.: Climate change projections for the Mediterranean region, *Global Planet. Chang.*, 63, 90–104, 2008.

Goldreich, Y.: The climate of Israel: observation research and application, Kluwer Academic Press, London, 270 pp., 2003.

Goldschmidt, M. J. and Jacobs, M.: Precipitation over and replenishment of the Yarqon and Nahal Hatteninim underground catchments, Hydrological Paper 3, Hydrological Service of Israel, Jerusalem, 1958.

Gregory, L., Wilcox, B. P., Shade, B., Munster, C., Owens, K., and Veni, G.: Large-scale rainfall simulation over shallow caves on karst shrublands, *Ecohydrology*, 2, 72–80, 2009.

Gunkel, A. and Lange, J.: New insights into the natural variability of water resources in the Lower Jordan River Basin, *Water Resour. Manag.*, 26, 963–980, 2012.

Gupta, H. V., Kling, H., Yilmaz, K. K., and Martinez, G. F.: Decomposition of the mean squared error and NSE performance criteria: implications for improving hydrological modelling, *J. Hydrol.*, 377, 80–91, 2009.

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Guttman, Y. and Zukerman, C. H.: Flow model in the Eastern Basin of the Judea and Samaria hills, unpublished report in Hebrew, TAHAL Consulting Engineers Ltd. 01/95/66, Tel Aviv, Israel, 1995.
- Hargreaves, G. H. and Samani, Z. A.: Reference crop evapotranspiration from temperature, *Appl. Eng. Agr.* 1, 96–99, 1985.
- Hartmann, A., Lange, J., Weiler, M., Arbel, Y., and Greenbaum, N.: A new approach to model the spatial and temporal variability of recharge to karst aquifers, *Hydrol. Earth Syst. Sci.*, 16, 2219–2231, doi:10.5194/hess-16-2219-2012, 2012.
- Hartmann, A., Wagener, T., Rimmer, A., Lange, J., Brielmann, H., and Weiler, M.: Testing the realism of model structures to identify karst system processes using water quality and quantity signatures, *Water Resour. Res.* 49, 3345–3358, 2013a.
- Hartmann, A., Weiler, M., Wagener, T., Lange, J., Kralik, M., Humer, F., Mizyed, N., Rimmer, A., Barberá, J. A., Andreo, B., Butscher, C., and Huggenberger, P.: Process-based karst modelling to relate hydrodynamic and hydrochemical characteristics to system properties, *Hydrol. Earth Syst. Sci.*, 17, 3305–3321, doi:10.5194/hess-17-3305-2013, 2013b.
- Hughes, A. G., Mansour, M. M., and Robins, N. S.: Evaluation of distributed recharge in an upland semi-arid karst system: the West Bank Mountain Aquifer, Middle East, *Hydrogeol. J.*, 16, 845–854, 2008.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, 1535 pp., 2013.
- IMS – Israel Meteorological Service: online database of the Israel Meteorological Service, Bet Dagan, available at: <http://data.gov.il/ims/3> (last access: 1 March 2014), 2014.
- Jensen, D. T., Hargreaves, G. H., Temesgen, B., and Allen, R. G.: Computation of ETo under nonideal conditions, *J. Irrig. Drain. Eng.*, 123, 394–400, 1997.
- Keshet, N. and Mimran, Y.: Landuse mapping – Ramallah area, unpublished report in Hebrew, Geological Survey of Israel, GSI/14/93, Jerusalem, 1993.
- Kizito, F., Campbell, C. S., Campbell, G. S., Cobos, D. R., Teare, B. L., Carter, B., and Hopmans, J. W.: Frequency, electrical conductivity and temperature analysis of a low-cost capacitance soil moisture sensor, *J. Hydrol.*, 352, 367–378, 2008.
- Kling, H., Fuchs, M., and Paulin, M.: Runoff conditions in the upper Danube basin under an ensemble of climate change scenarios, *J. Hydrol.*, 424, 264–277, 2012.

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Kutiél, P., Lavee, H., and Ackermann, O.: Spatial distribution of soil surface coverage on north and south facing hillslopes along a Mediterranean to extreme arid climatic gradient, *Geomorphology*, 23, 245–256, 1998.
- Lange, J., Greenbaum, N., Husary, S., Ghanem, M., Leibundgut, C., and Schick, A. P.: Runoff generation from successive simulated rainfalls on a rocky, semi-arid, Mediterranean hillslope, *Hydrol. Process.*, 17, 279–296, 2003.
- Lange, J., Arbel, Y., Grodek, T., and Greenbaum, N.: Water percolation process studies in a Mediterranean karst area, *Hydrol. Process.*, 24, 1866–1879, 2010.
- Lavee, H., Imeson, A. C., and Sarah, P.: The impact of climate change on geomorphology and desertification along a Mediterranean-arid transect, *Land Degrad. Dev.*, 9, 407–422, 1998.
- Marei, A., Khayat, S., Weise, S., Ghannam, S., Sbaih, M., and Geyer, S.: Estimating groundwater recharge using the chloride mass-balance method in the West Bank, Palestine, *Hydrolog. Sci. J.*, 55, 780–791, 2010.
- Morin, E., Jacoby, Y., Navon, S., and Bet-Halachmi, E.: Towards flash-flood prediction in the dry Dead Sea region utilizing radar rainfall information, *Adv. Water Resour.*, 32, 1066–1076, 2009.
- Öhrström, P., Persson, M., Albergel, J., Zante, P., Nasri, S., Berndtsson, R., and Olsson, J.: Field-scale variation of preferential flow as indicated from dye coverage, *J. Hydrol.*, 257, 164–173, 2002.
- Puigdefabregas, J., del Barrio, G., Boer, M. M., Gutiérrez, L., and Solé, A.: Differential responses of hillslope and channel elements to rainfall events in a semi-arid area, *Geomorphology*, 23, 337–351, 1998.
- Reiser, H. and Kutiél, H.: The dependence of annual total on the number of rain spells and their yield in the Mediterranean, *Geogr. Ann. A*, 94, 285–299, 2012.
- Rofe and Raffety Consulting Engineers: Jerusalem and District Water Supply: Geological and Hydrological Report, Report to the Central Water Authority of the Hashemite Kingdom of Jordan, Rofe and Raffety Consulting Engineers, London, 1963.
- Ruiz-Sinoga, J. D., Martínez-Murillo, J. F., Gabarrón-Galeote, M. A., and García-Marín, R.: The effects of soil moisture variability on the vegetation pattern in Mediterranean abandoned fields (Southern Spain), *Catena*, 85, 1–11, 2011.
- Samuels, R., Smiatek, G., Krichak, S., Kunstmann, H., and Alpert, P.: Extreme value indicators in highly resolved climate change simulations for the Jordan River area, *J. Geophys. Res. Atmos.*, 116, D24123, doi:10.1029/2011JD016322, 2011.

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Scanlon, B. R., Healy, R. W., and Cook, P. G.: Choosing appropriate techniques for quantifying groundwater recharge, *Hydrogeol. J.*, 10, 18–39, 2002.
- Schmidt, S., Geyer, T., Marei, A., Guttman, J., and Sauter, M.: Quantification of longterm wastewater impacts on karst groundwater resources in a semi-arid environment by chloride mass balance methods, *J. Hydrol.* 502, 177–190, 2013.
- Schmidt, S., Geyer, T., Guttman, J., Marei, A., Ries, F., and Sauter, M.: Characterisation and modelling of conduit restricted karst aquifers – example of the Auja spring, Jordan Valley, *J. Hydrol.*, 511, 750–763, 2014.
- Schoups, G., Hopmans, J. W., Young, C. A., Vrugt, J. A., and Wallender, W. W.: Multi-criteria optimization of a regional spatially-distributed subsurface water flow model, *J. Hydrol.*, 311, 20–48, 2005.
- Scott, R. L., Shuttleworth, W. J., Keefer, T. O., and Warrick, A. W.: Modeling multiyear observations of soil moisture recharge in the semiarid American Southwest, *Water Resour. Res.*, 36, 2233–2247, 2000.
- Shapiro, M. B.: Soils of Israel, *Eurasian Soil Sci.*, 39, 1170–1175, 2006.
- Sheffer, N. A., Dafny, E., Gvirtzman, H., Navon, S., Frumkin, A., and Morin, E.: Hydrometeorological daily recharge assessment model (DREAM) for the Western Mountain Aquifer, Israel: model application and effects of temporal patterns, *Water Resour. Res.*, 46, W05510, doi:10.1029/2008WR007607, 2010.
- Sheffer, N. A., Cohen, M., Morin, E., Grodek, T., Gimburg, A., Magal, E., Gvirtzman, H., Nied, M., Isele, D., and Frumkin, A.: Integrated cave drip monitoring for epikarst recharge estimation in a dry Mediterranean area, Sif Cave, Israel, *Hydrol. Process.*, 25, 2837–2845, 2011.
- Šimůnek, J., Šejna, M., Saito, H., Sakai, M., and van Genuchten, M. Th.: The Hydrus-1D software package for simulating the movement of water, heat, and multiple solutes in variably saturated media, version 4.16, HYDRUS Software Series 3, Department of Environmental Sciences, University of California Riverside, Riverside, California, USA, 340 pp., 2013.
- Sohrt, J., Ries, F., Sauter, M., and Lange, J.: Significance of preferential flow at the rock soil interface in a semi-arid karst environment, *Catena*, 123, 1–10, 2014.
- van Genuchten, M. Th.: A closed-form equation for predicting the hydraulic conductivity of unsaturated soils, *Soil Sci. Soc. Am. J.*, 44, 892–898, 1980.

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Van Schaik, N. L. M. B., Schnabel, S., and Jetten, V. G.: The influence of preferential flow on hillslope hydrology in a semi-arid watershed (in the Spanish Dehesas), *Hydrol. Process.*, 22, 3844–3855, 2008.
- Vereecken, H., Huisman, J. A., Bogena, H., Vanderborght, J., Vrugt, J. A., and Hopmans, J. W.: On the value of soil moisture measurements in vadose zone hydrology: a review, *Water Resour. Res.*, 44, W00D06, doi:10.1029/2008WR006829, 2008.
- Vogel, T., van Genuchten, M. T., and Cislserova, M.: Effect of the shape of the soil hydraulic functions near saturation on variably-saturated flow predictions, *Adv. Water Resour.*, 24, 133–144, 2001.
- Vrugt, J. A., Gupta, H. V., Bouten, W., and Sorooshian, S.: A shuffled complex evolution metropolis algorithm for optimization and uncertainty assessment of hydrologic model parameters, *Water Resour. Res.*, 39, W01017, doi:10.1029/2002WR001642, 2003.
- Vrugt, J. A., Diks, C. G., Gupta, H. V., Bouten, W., and Verstraten, J. M.: Improved treatment of uncertainty in hydrologic modeling: combining the strengths of global optimization and data assimilation, *Water Resour. Res.*, 41, W01017, doi:10.1029/2004WR003059, 2005.
- Weiss, M. and Gvirtzman, H.: Estimating ground water recharge using flow models of perched karstic aquifers, *Ground Water*, 45, 761–773, 2007.
- Weiβ, M. and Menzel, L.: A global comparison of four potential evapotranspiration equations and their relevance to stream flow modeling in semi-arid environments, *Adv. Geosci.*, 18, 15–23, 2008, <http://www.adv-geosci.net/18/15/2008/>.
- Williams, P. W.: The role of the subcutaneous zone in karst hydrology, *J. Hydrol.*, 61, 45–67, 1983.
- Wraith, J. M. and Or, D.: Temperature effects on soil bulk dielectric permittivity measured by time domain reflectometry: a physical model, *Water Resour. Res.*, 35, 371–383, 1999.
- Yaalon, D. H.: Soils in the Mediterranean region: what makes them different?, *Catena*, 28, 157–169, 1997.
- Zagana, E., Kuells, C., Udluft, P., and Constantinou, C.: Methods of groundwater recharge estimation in eastern Mediterranean – a water balance model application in Greece, Cyprus and Jordan, *Hydrol. Process.*, 21, 2405–2414, 2007.
- Zukerman, C. H.: Yarqon-Tanninim-Beer Sheva Basin, Flow Model Update, unpublished report in Hebrew, TAHAL Consulting Engineers Ltd. 6759/700/133, Tel Aviv, Israel, 1999.

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Table 1. Soil moisture plot characteristics.

Plot	Elevation (m a.s.l.)	Average annual rainfall ^a (mm)	Soil depth (cm)	Sensor depths (cm)	Vegetation	Description
SM-1	810	526	100	10, 25, 40, 80	Mediterranean shrubs; annual plants	Sand: 20 % Silt: 40 % Clay: 40 %
SM-2	660	340 ^b	50	5, 10, 20, 35	Annual plants	Sand: 32 % Silt: 33 % Clay: 35 %
SM-3	440	351	60	5, 10, 20, 35	Annual plants	Sand: 46 % Silt: 24 % Clay: 30 %

^a Mean rainfall based on three winter seasons (2010–2013).

^b Rainfall at plot SM-2 is estimated by inverse distance weighted interpolation with elevation as additional predictor.

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Parameters and value ranges for Hydrus-1D modelling.

Parameter	Value/Range	Unit	Source/calculation method	
Soil hydraulic parameter				
Θ_r	Residual soil water content ^b	0–0.3	$\text{m}^3 \text{m}^{-3}$	Calibrated ^a
Θ_s	Saturated soil water content ^b	0.3–0.6	$\text{m}^3 \text{m}^{-3}$	Calibrated ^a
α	Van Genuchten parameter related to air entry suction	0.0001–0.1	mm^{-1}	Calibrated ^a
n	Van Genuchten parameter related to pore size distribution	1.01–3	–	Calibrated ^a
K_s	Saturated hydraulic conductivity	5–10 000	mm day^{-1}	Calibrated ^a
L	Van Genuchten parameter related to tortuosity	–2–2	–	Calibrated ^a
Meteorological parameter				
P	Daily precipitation		mm	Measured time series ^c
T_{\max}	Daily maximum temperature		$^{\circ}\text{C}$	Measured time series ^d
T_{\min}	Daily minimum temperature		$^{\circ}\text{C}$	Measured time series ^d
R_a	Extraterrestrial solar radiation (for Hargreaves equation only)		MJ m^{-2}	Calculated according to Allen et al. (1998)
Vegetation parameter				
D_r	Rooting depth	0.5–1	m	Estimated based on field observations
SCF	Surface Cover Fraction	0.1–1	m/m	Estimated based on field observations
LAI	Leaf Area Index		m/m	Calculated according to Šimůnek (2013)
P_0	Fedde's parameter	–100	mm	Hydrus-1D internal database (grass)
P_{opt}	Fedde's parameter	–250	mm	Hydrus-1D internal database (grass)
$P_{2\text{H}}$	Fedde's parameter	–3000	mm	Hydrus-1D internal database (grass)
$P_{2\text{L}}$	Fedde's parameter	–10 000	mm	Hydrus-1D internal database (grass)
P_3	Fedde's parameter	–80 000	mm	Hydrus-1D internal database (grass)
$r_{2\text{H}}$	Fedde's parameter	5	mm day^{-1}	Hydrus-1D internal database (grass)
$r_{2\text{L}}$	Fedde's parameter	1	mm day^{-1}	Hydrus-1D internal database (grass)
α_i	Interception constant	1	mm	Estimated
D_s	Depth of soil profile	0.5–1	m	Measured at experimental plots

^a Parameter calibrated for each soil material with SCEM algorithm and Kling–Gupta efficiency as optimization criterion.

^b The upper parameter limit of Θ_r and the lower parameter limit of Θ_s were obtained from the lowest respectively highest measured volumetric soil moisture value of each layer in the respective soil moisture plot.

^c Rainfall at plot SM-2 is estimated by inverse distance weighted interpolation with elevation as additional predictor.

^d Maximum and minimum daily air temperature at the soil moisture plots is estimated by calculation of an elevation-temperature gradient based on meteorological stations in the Jordan Valley and the mountains.

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

Table 3. SCEM optimized hydraulic parameter sets for the different plots and probe depths.

Plot	Layer	Θ_r ($\text{m}^3 \text{m}^{-3}$)	Θ_s ($\text{m}^3 \text{m}^{-3}$)	α (1mm^{-1})	n (–)	K_s (mm day^{-1})	L (–)	KGE (–)
SM-1	1 (–10 cm)	0.01	0.41	0.004	1.23	427	2.0	0.91
	2 (–25 cm)	0.12	0.49	0.026	1.30	8159	–2.0	0.94
	3 (–40 cm)	0.11	0.59	0.018	1.54	9468	–2.0	0.90
	4 (–80 cm)	0.10	0.59	0.028	1.36	8732	0.1	0.82
SM-2	1 (–5 cm)	0.00	0.49	0.041	1.18	126	–2.0	0.89
	2 (–10 cm)	0.05	0.40	0.002	1.23	5094	0.6	0.90
	3 (–18 cm)	0.12	0.59	0.012	1.37	9288	2.0	0.87
	4 (–55 cm)	0.13	0.51	0.013	1.43	2679	1.0	0.90
SM-3	1 (–5 cm)	0.00	0.60	0.008	1.23	482	–2.0	0.91
	2 (–10 cm)	0.00	0.56	0.004	1.23	9908	–1.2	0.92
	3 (–20 cm)	0.05	0.46	0.003	1.22	9976	1.2	0.91
	4 (–35 cm)	0.11	0.60	0.001	1.66	5751	2.0	0.94

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Table 4. Cumulative sums of the simulated water balance components in mm and % for the three consecutive hydrological years 2010–2013.

Plot	Year	Rainfall	Interception		Evaporation		Transpiration		Bottom flux	
		(mm)	(mm)	(%)	(mm)	(%)	(mm)	(%)	(mm)	(%)
SM-1	2010/11	381	62	16	99	26	209	55	13	3
	2011/12	650	59	9	93	14	209	32	294	45
	2012/13 ^a	547	39	7	102	19	179	33	224	41
SM-2	2010/11	248	53	21	81	33	117	47	0	0
	2011/12	418	55	13	89	21	159	48	118	28
	2012/13 ^a	346	33	10	84	24	127	37	101	29
SM-3	2010/11	237	47	20	119	50	84	35	2	1
	2011/12	436	53	12	120	27	130	30	135	31
	2012/13 ^a	380	30	8	111	29	105	28	125	33

^a The hydrological year 2012/13 was modelled until 30 April 2013.

HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.




Figure 1. Study area with location of meteorological stations, rain gauges, soil moisture plots (SM-1, SM-2, SM-3) and isohyets of long-term average annual rainfall (≥ 20 years) according to data from ANTEA (1998). Coordinates in the detailed map are in Palestinian Grid format.

Title Page

Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



HESSD

11, 8803–8844, 2014

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.



Figure 2. Typical hillslopes in the study area. The image shows the plain of Ein Samia with semi-arid climatic conditions, where the valley bottom is used for partly irrigated agriculture and the hillslopes are used as extensive grazing land for goats and sheep.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.




Figure 3. Correlation between average annual rainfall and station elevation for the individual hydrological years during the observation period 2010–2013.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.




Figure 4. Observed volumetric soil moisture at different depths of the three experimental plots during the complete monitoring period (a) and details on the winter season 2011/12 for plot SM-1 (b).

[Title Page](#)
[Abstract](#) [Introduction](#)
[Conclusions](#) [References](#)
[Tables](#) [Figures](#)
[◀](#) [▶](#)
[◀](#) [▶](#)
[Back](#) [Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.




Figure 5. Cumulative distribution functions of parameters following 1000 model runs after convergence. The line colours correspond to the various soil depths (see Fig. 6). The limits of the x-axis represent the selected parameter ranges.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.




Figure 6. Measured against simulated volumetric soil water content for the three soil moisture plots (SM-1, SM-2, SM-3) at individual depths for the simulation period. KGE values are shown in brackets in the legend.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.




Figure 7. Time series of rainfall (a), simulated and observed volumetric water content for soil moisture plot SM-1 (b–e), Hydrus-1D simulated percolation (f) and water temperature in a nearby groundwater well (g). The grey shaded area represents the 95 % confidence interval of soil moisture based on model parameter sets obtained using SCEM after fulfilment of the convergence criterion.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Figure 8. Simulated daily water fluxes at the single soil moisture plots for the simulation period 2010–2013.

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.




Figure 9. Simulated percolation vs. soil depth and rainfall amounts along the climatic gradient for three consecutive winter seasons with different rainfall depths and distribution patterns. Simulations were based on calibrated soil hydraulic properties of plots SM-1. The grey shaded areas display rainfall depths, which have not been reached in the study area within altitudes of 400 to 1000 m a.s.l. according to calculated rainfall gradients. The points represent the plot scale simulated percolation fluxes using optimal parameter sets for the single plots SM-1, SM-2 and SM-3.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.




Figure 10. Seasonal sums of simulated water balance components for the period 1951 to 2013 using the calibrated soil hydraulic parameters of the various plots. Rainfall and temperature data were obtained from the nearby Jerusalem central station (<http://www.data.gov.il/ims>) and corrected for the single locations by applying a simple elevation gradient-based correction factor.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Recharge estimation and soil moisture dynamics in a Mediterranean karst aquifer

F. Ries et al.




Figure 11. Simulated seasonal percolation at the plot scale (SM-1, SM-2, SM-3) for the period 1951–2013 in comparison with rainfall–recharge relationships for the carbonate aquifer (Guttman and Zukerman, 1995) and three small karst springs emerging from local perched aquifers (Weiss and Gvirtzman, 2007).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)