

Interactive comment on “On inclusion of water resource management in Earth System models – Part 1: Problem definition and representation of water demand” by A. Nazemi and H. S. Wheater

A. Nazemi and H. S. Wheater

ali.nazemi@usask.ca

Received and published: 4 November 2014

We greatly appreciate Jan Polcher for his positive, constructive and thoughtful comments, which led to substantial improvements in the revised version of our manuscript. In the following, the issues raised are addressed point-by-point in the order they are asked. Jan Polcher’s comments are numbered; our reply to each comment is shown immediately below the comment.

1- 8242, 19 : for me the first attempts to include routing in LSMs (I prefer the Land Surface Model term so that in Earth System modelling the land is at the same level as

C4875

the ocean and models. Who would dare speak about an ocean or atmosphere scheme ?) are: Miller et al, 1994 J. Clim, Hagemann and Dümenil, 1998, Climate Dynamics, Oki and Sud, 1998, Earth Interactions.

Many thanks for your comment. We changed LSSs to LSMs throughout the paper. We have also included the early works you have reminded. Please see the attached draft revisions (lines 69 to 72).

2- 8245, 7 : In the list of possible effects of irrigation and water usage on the climate system, the impact on ocean circulation should be mentioned. This is of particular concern for closed oceans and the polar environment where a change in fresh water input can modify the oceanic circulations and thus feedback on continental rainfall. A recent review of literature showed a few nice examples for the Mediterranean : E. J. Rohling and H. L. Bryden (1992) Man-induced salinity and temperature increases in western Mediterranean deep water. J. Geophys. Res., 97(C7), 11191–11198 M. 11, C3403–C3410, 2014. Vargas-Yañez et al. (2010) Climate change in The Western Mediterranean Sea (1900-2008). Journal of Marine Systems 82(2010) 171-176. N. Skliris and A. Lascaratos (2004) Impacts of the Nile River damming on the thermohaline circulation and water mass characteristics of the Mediterranean Sea. Journal of Marine Systems 52(1–4), 121–143.

Many thanks for the heads-up on this important issue. We added few sentences regarding this and included the references in the text. Please see the attached draft revisions (lines 150 to 156). We believe that these issues are more related to the effect of water resource management on water quality rather than water quantity. As we look at water resource management as a water quantity problem (please see the attached draft revisions, lines 162 to 164), we do not follow this issue further up in the paper.

3- 8245, 15 : A recent study which shows (from one specific model !) the regions where irrigation triggers an atmospheric feedback in the water cycle and those where rainfall is not affected : Guimberteau et al. (2012) Global effect of irrigation and its impact on

C4876

the onset of the Indian summer monsoon, *Climate Dynamics*, Volume 39, Issue 6, pp. 1329-1348.

Many thanks for introducing this valuable contribution. We included the reference where you suggested. We have also used the reference to elaborate our discussion in Section 5.1. Please see the attached draft revisions (line 146 as well as lines 428 to 431, 598 to 600 and 608 to 614).

4- 8245, 24 : I would already write in the abstract that this review paper is in line with GEWEX's ambition to strengthen activities on human-water interactions and raise the awareness on this issue.

We modify the text based on your suggestion. Please see the attached draft revisions (lines 22 to 26).

5- 8246, 9 : Yes, there are still fundamental obstacles to include water resources in large scale models, but I would say that it does not matter if this is on-line or off-line. The nature of the coupling to the atmosphere is not affected by irrigation as it is only evaporation and the surface energy balance which are changed. I would say that in the "water conserving approach" to irrigation, we have to deal with the fundamental problem that man is also modifying the transport of water and tapping non renewable water sources which are outside of the climate system.

Many thanks for this extremely constructive comment. We revised the text to include this very important point in the text and discuss it further in the discussion section. Please see the attached draft revisions (lines 170 to 177).

6- 8252, 20 : In the this discussion of the usage of ETP one has to take into account that LSMs define potential evaporation in a quite different way from FAO, Penman-Monteith or others. Thus using simply the FAO guidelines for estimation irrigation needs will induce inconsistencies at various time scales with the evaporation estimated by the model. This is of particular concern for water stressed surfaces which is the case

C4877

when we expect irrigation to occur. This problem is limited to LSMs which resolve the diurnal cycle and does not occur in GHMs which use anyway some empirical estimates of ETP for evaporation. This issue is well documented in : Milly, P. C. D.: Potential evaporation and soil moisture in general circulation models (1992), *J. Climate*, 5, 209–226. Barella-Ortiz, A., et al. (2013) Potential evaporation estimation through an unstressed surface energy balance and its sensitivity to climate change, *Hydrol. Earth Syst. Sci.*, 17, 4625-4639.

Many thanks for this very useful comment. We made some revisions in the discussion to reflect the difference between calculation of potential evaporation in LSMs and GHMs and incorporated the references indicated in the text. Please see the attached draft revisions (lines 365 to 384).

7- 8254, 5 : Using the extra information available in LSMs we can now do better and the concerns raised here are behind us. The irrigation need can be estimated using potential transpiration. This is the transpiration which would occur should the plan not be water stressed. If this is implemented together with a sub-grid soil moisture division (i.e. bare soils and non-crop PFTs have different soil moisture reservoirs) then the irrigation taken from the water reservoirs optimises photosynthesis and is only evaporated by the crops and not used by other surface types. Furthermore the potential transpiration takes into account the CO₂ fertilisation, the adaptation of the plants to the climatic conditions or crop growth cycles as far as the LSM represents them. This is now present in ORCHIDEE and documented by Guimberteau et al. (see above for the full reference). The next step in the uncertainty is whether the irrigation is sprinkled on the crop, and thus induces some interception loss, or localized and limited to soil moisture processes. But this far beyond the current state of our models and would require knowledge on the irrigation techniques used in each region of the world.

Many thanks for the valuable discussion. We used the first part of your discussion in the next paragraph, where we discuss the potential transpiration algorithms. Please see the attached draft revisions (lines 425 to 449). We incorporated the second part

C4878

of you comment in the discussion related to data uncertainty in Section 6. Please see the attached draft revisions (lines 778 to 780).

8- 8254, 19 : This evolution toward potential transpiration is partially explained in this paragraph but does not address the issue of having to treat separately in the grid box the irrigated vegetation from the rest. Most LSMs today define multiple plant functional types (PFTs) in each grid box and can thus distinguish the various water needs. But as long as all PFTs share the same soil moisture reservoir this does not help. Irrigation will increase the soil moisture of all PFTs and thus reduce water stress for forests as well as crops and in particular increase bare soil evaporation. Thus too much water will be used for irrigation and the evaporation increase overestimated.

Many thanks for the follow up discussion. We merged your discussions in this comment and the previous one and revised the related text accordingly. Please see the attached draft revisions (lines 442 to 449).

9- 8254, 21: The projection of irrigative demand is closely linked to the infrastructures which can be put into place to adducts water to the area where farming occurs. There is some pioneering work being done by economists and which is able to predict which regions can be irrigated and how the irrigation can be sustained in a changing climate. The modelling is purely based on the economical cost of bringing the water from the regions where it is available (generally mountains because of the amount of rainfall and the available potential energy) to those where the farming occurs (sedimentary plains and urbanized areas). The thesis of Hypatia Nassopoulos: <http://halshs.archivesouvertes.fr/pastel-00838516/>, her papers and more generally the group at CIRED are at the forefront of this research. I know the thesis is in French and I am not sure if the part on the model to predict dam operations and water adduction has been published. But Hypatia Nassopoulos can be contacted.

Many thanks for the heads-up on Hypatia's work. We were not aware of her work. We found one of her papers and presentations online, which could help

C4879

us to write a short entry on her work and incorporate it at the end of Section 3.4. The reference are as following: Nassopoulos, H., Dumas, P. and Hallegatte, S.: Adaptation to an uncertain climate change: cost benefit analysis and robust decision making for dam dimensioning, *Climatic change*, 114(3-4), 497-508, doi: 10.1007/s10584-012-0423-7, 2012. Nassopoulos, H., Dumas, P. and Hallegatte, S.: Climate change, precipitation and water management infrastructures, presented at: Water in Africa: Hydro-Pessimism or Hydro-Optimism, 2-3 October 2008, Porto, Portugal, available at: <http://www.slideshare.net/water.in.africa/hypatia-nassopoulos-ppt-presentation>. Please see the attached draft revisions (lines 471 to 475).

10- 8259, 7: In this discussion of the irrigation-induced (or irrigation-displaced) rainfall the rôle of the conservation of water needs to be taken into account. For a model which limits irrigation by the available water stabilising feedbacks can be envisioned. Should irrigation for instance displace rainfall into the neighbouring valley/catchment, then the originally irrigated farmland cannot be sustained as the basin total rainfall might become to low to support the activity. This is perhaps far fetched, but it is a process which can limit irrigation and is not available to parametrisations which do not close the water balance. Thus I would classify these studies into the general topic of surface/atmosphere feedback studies where the surface energy balance perturbation is irrigation. As far as I could verify, none of the studies referred to in Table 3 include the feedbacks generated by water conservation.

Many thanks for the discussion. We merged your discussions in this comment and the next one and revised the related text accordingly to highlight the limitations in current online studies analyzing the irrigation-induced precipitation. Please see the attached draft revisions (lines 620 to 630).

11- 8259, 28 : The studies presented here on the surface/atmosphere interactions are all analysed on the simple scheme of whether evaporation increase can favour moisture convergence or on the contrary reduce it. This has to be linked in some way to the wealth of literature where deforestation (or more academic perturbations) and

C4880

its impact on evaporation are discussed. But I feel there is a recent evolution which is being missed here. It is now accepted that landscape contrasts (transitions between wet and cool and dry and hot areas) are critical in generating rainfall. Irrigation has a huge effect on this type of mechanisms as it creates sharp contrasts in evaporation and surface temperature. But models are known to be limited in their ability to generate the atmospheric perturbations caused by these processes and thus sensitivity experiments have to be analysed with caution. I would suggest that the authors take a look at this part of the literature of which I only highlight 2 recent publications : Taylor (2009) Feedbacks on convection from an African wetland, GRL, VOL. 37, L05406 (These African wetlands are just naturally irrigated areas !) Taylor et al. (2012) Afternoon rain more likely over drier soils, Nature, 489, 423–426.

Many thanks for the follow up discussion. We merged your discussions in this comment and the previous one and revised the related text accordingly to highlight the limitations in current online studies analyzing the irrigation-induced precipitation. Please see the attached draft revisions (lines 620 to 630). Please note that we only incorporate the references indicated, as the text and the reference list are already quite long (i.e., 276 references).

12- 8260, 18: Some LSMs have included irrigation in all of their studies as it simply was available in the model and provided more realistic river discharge values on many of the basins considered. One of these cases are the studies performed by Thanh Ngo-Duc during his thesis. When validating his atmospheric forcing over large basins, looking at the water exchanges between continents and oceans or validating ORCHIDEE with GRACE, the irrigation parametrisation of de Rosnay et al. was used but its impact not specifically analysed. The references are : Ngo-Duc T. et al. (2005) 53 years forcing data set for land-surface models, J. Geophys. Res., 110:D06 116 Ngo-Duc, T. et al. (2005): Effects of land water storage on global mean sea level over the past 50 years. Geophysical Research Letters, 32:L09704 Ngo-Duc, T. et al. (2006): Validation of the land water storage simulated by ORCHIDEE with the GRACE data, role of the routing

C4881

scheme. Water Resources Research, 43(4):W04427.

Many thanks for the heads-up on these references. We incorporated them in the related discussion for offline simulations. Please see the attached draft revisions (lines 652 to 655).

13- 8264, 17 : I believe that in this section the authors mix different aspects of spatial resolution. First there is the spatial resolution needed to represent properly the irrigation processes. This can be achieved either by running the LSM at a higher resolution than the atmospheric component or by obtaining a higher effective resolution at the surface by using tiling approaches. As I pointed out above, if the crop PFTs have their own soil moisture reservoir the impact of irrigation on their evaporation can be quite well represented. The second issue is the adequate resolution to represent the impact of increased evaporation and surface flux contrasts onto the atmospheric processes. For this problem, I do not know of any study as it is probably strongly regionally and seasonally dependent. But this issue of resolution is not independent of the complexity of the parametrisation of irrigation. As the resolution of the surface increases more processes need to be included in order to ensure water conservation within the model as else not enough water is available in each grid box to sustain the enhanced evaporation.

Many thanks for this discussion. We elaborate our discussion by incorporated this into the related text. Please see the modified text in the track changed file attached (lines 733 to 749).

14- 8266, 4 : The uncertainty of the demand linked to the potential evaporation is not that much of an issue as long as the same assumption is used for reference evaporation (or ETP) in the calculation of the crop evaporation and the irrigation demand. If the GHM uses Priestley-Taylor then the FAO guideline has to be re-interpreted accordingly. For the LSM more options are available as ETP or potential transpiration consistent with the surface energy balance can be derived in the model (but significantly different

C4882

from Penman-Monteith as pointed out above). Thus if the consistency of the model is preserved, then the uncertainty of the irrigative demand linked to ETP is the same as that of the evaporation.

Many thanks for the discussion. We merged your discussions in this comment and the next one and revised the related text accordingly to better highlight the main sources of uncertainty in current irrigation demand algorithms. Please see the attached draft revisions (lines 795 to 804).

15- 8266, 4 : To me the largest uncertainty in the parametrisations currently available is the limitation of irrigation by the available water. If the irrigation is limited by the water available within the grid box then we are hindered by our ability to describe water transports and the role played by humans and our lack of geological water used in some regions.

Many thanks for the discussion. We merged your discussions in this comment and the previous one and revised the related text accordingly to better highlight the main sources of uncertainty in current irrigation demand algorithms. Please see the attached draft revisions (lines 795 to 804).

16- 8293, Table 1 : de Rosnay et al. was implemented globally and only analysed over the Indian Peninsula. So it should probably move to table 2.

Many thanks for the heads-up on this. We moved the reference to Table 2.

17- 8295, Table 3 : Guimberteau et al. 2013 is missing.

Many thanks for introducing this reference. The reference is now incorporated in Table 3.

Please also note the supplement to this comment:

<http://www.hydrol-earth-syst-sci-discuss.net/11/C4875/2014/hessd-11-C4875-2014-supplement.pdf>

C4883

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., 11, 8239, 2014.

C4884