Hydrol. Earth Syst. Sci. Discuss., 11, C5635–C5636, 2014 www.hydrol-earth-syst-sci-discuss.net/11/C5635/2014/ © Author(s) 2014. This work is distributed under the Creative Commons Attribute 3.0 License.





Interactive Comment

Interactive comment on "From runoff to rainfall: inverse rainfall–runoff modelling in a high temporal resolution" by M. Herrnegger et al.

D. Brochart

david.brochart@irstea.fr

Received and published: 11 December 2014

Mathew,

I was not talking about the spatial heterogeneity of the catchment. What I meant is that if the catchment is large enough, you don't necessarily see the effect of the rainfall in the discharge at the outlet in the same time step, because of the time lag. Even a lumped model takes this lag into account. So if you wait until you see the discharge increase, it is generally too late: the rainfall that generated this discharge happened before. Regarding the second point, consider this simple fact: it is raining but the discharge continues to decrease - just at a slower rate than if it didn't rain. This is perfectly valid, not a simulation error nor a discharge measurement error. Can your





algorithm infer this rainfall?

Regards,

David.

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., 11, 13259, 2014.

HESSD

11, C5635–C5636, 2014

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

