

## **Response to Anonymous Referee #1's comments on "Analyses of Uncertainties and Scaling of Groundwater Level Fluctuations"**

**General comments:** *This paper deals with uncertainties in groundwater level in unconfined aquifers due to temporal variations of hydrological processes. It derives the head covariance function for 1-D transient flow in a bounded unconfined aquifer with random recharge as well as random initial and boundary conditions. Associated time-dependent spectral densities are also derived, allowing to investigate the existence of temporal scaling of groundwater level fluctuations. The topic of the note lies within the aims and scope of Hydrology and Earth System Sciences and is a valuable addition to the existing literature. The paper is well-written and concise, and deals with a topic of considerable interest. The mathematical derivations are accurate.*

**Response:** Thank you for the reviewer's positive comment on our study.

**Specific comments:** *Specific suggestions to improve the quality of the paper are listed below.*

1. *The authors should mention specific applications of their results to real cases, to help the paper convey a take-home message.*

**Response:** This is an excellent comment. The analytical solutions for the head variances derived in this study provide a way to quantify the uncertainty in the groundwater levels calculated with analytical and numerical solutions with uncertain recharge, source/sink, and boundary conditions. The spectrum relationship among the head, recharge and boundary conditions obtained in this study can help one to improve spectrum analysis for a groundwater level time series and to remove the effects of boundary conditions. Specific applications of the results obtained in this study are to help one to identify and quantify the sources of uncertainties in the system he/she studied. We added these in lines 375-379 of the revised manuscript.

2. *I suggest to add a schematic of the system investigated for the sake of clarity. This will help clarifying the meaning of the quantity  $M$ , defined at line 152 as the average saturated thickness of the aquifer. Since  $h$  is random,  $M$  should incorporate an element of randomness.*

**Response:** As suggested by the reviewer, we added a schematic of the system studied, i.e., the new Figure 1 in the revised manuscript. Please note that it is assumed in this study that the fluctuation of the head is relatively small as compared to the aquifer thickness and the unconfined flow equation was linearized and thus the average saturated thickness of the aquifer ( $M$ ) was assumed to be constant.

3. *A key assumption in the analysis is that  $W(t)$ ,  $Q(t)$ , and  $H(t)$  are uncorrelated (see line 137). Given the geometrical setup, this assumption is not warranted. The paper could benefit from discussing this issue, and, specifically, realistic conditions for the validity of the assumption.*

**Response:** This is again an excellent comment. In general,  $W(t)$ ,  $Q(t)$ , and  $H(t)$  should be correlated. It is possible to consider the relationship among  $W(t)$ ,  $Q(t)$ , and  $H(t)$  by assuming some theoretical correlation functions but the problem is that 1) it is unclear

what kind of correlation exists among these variable, 2) there is little observed or measured data to support any type of the correlation assumed, and 3) simple analytical solutions would be difficult to derive when considering such a correlation. Therefore, we studied the case in which such correlation is weak or no correlation in order to derive some simple analytical solutions. We believe this is an important first step towards solving this complex problem and more research is needed in this direction, especially about the correlation among the recharge, flux, and boundary conditions. We hope to relax this assumption in our future study.

4. (a) *Temporal scaling of groundwater level fluctuations is shown to exist at intermediate and late times, and to be dominated by the effect of random recharge as opposed to that of random boundary conditions. Why? Is this valid only for the specific parameters examined?* (b) *When spectra associated with one random effect at a time are examined, different scalings ( $1/f$ ,  $1/f^2$ ) are found. Why does this happen?*

**Response:**

- a) We think the reason that the temporal scaling of groundwater level is dominated by the random recharge is that the areal recharge occurs over entire the aquifer and thus affect the groundwater level everywhere in the aquifer while the boundary conditions affect a relatively small area near the boundaries in most aquifers. The specific parameters used in this study are typical for a real aquifer. The effect of the boundary conditions or the area of the influence by the boundary conditions would be enhanced in a more permeable aquifer. However, in most aquifers areal recharge should be the dominating force affecting groundwater level fluctuations and its scaling.
- b) We do not totally understand this comment. That the groundwater level fluctuates as a  $1/f$  noise only at  $x'=0$  under the random flux boundary, and it fluctuates as a  $1/f^2$  noise at most locations only under random recharge when the characteristic time scale ( $t_c$ ) is large, i.e.,  $t_c > 400$  days .

Minor points:

1. *Check keywords.*

**Response:** We revised the keywords as: Uncertainty of groundwater levels; Temporal scaling; Random source/sink; Random initial and boundary conditions.

2. *Check line 75.*

**Response:** We checked but didn't find any problem in this line.

3. *Check equation (12).*

**Response:** We checked the Eq. (12) in our original submission (now is Eq. 11 in the revision) and didn't find any error. However, there are two typos in Eq. (8) and (9): one is that the Eqs. (8) and (9) are actually one equation, and the other is in the first term on the right hand of the equation. We corrected these typos. The correct Eq. (8) was given in the revision.

4. *Check Line 173, 'is' is missing.*

**Response:** We added it.

5. *Line 175, in 'decay' a 's' is missing.*

**Response:** We added it.

6. Line 204, check if 'the' is missing.

**Response:** We added "the " before "most aquifers".

7. Line 223, check if 'on' is missing.

**Response:** We deleted this sentence based on Reviewer #2's comments.

8. Check the sentence at lines 346-347.

**Response:** We added 's' after 'curve' and 'location', respectively.

## **Response to Anonymous Referee #2's comments on "Analyses of Uncertainties and Scaling of Groundwater Level Fluctuations"**

### **General comments:**

1. *Liang and Zhang present an extension of previous work on the impact of temporal variations in hydrological processes on the uncertainty in groundwater level fluctuations. The authors derived analytical solutions for variance, covariance and spectrum of groundwater levels under random boundary conditions and a random source/sink, which is something new and interesting to the hydrological community.*

**Response:** Thank you for the positive comment by the reviewer on our study.

2. *The current work strongly relies on previous work of the same authors in that field, it can be interpreted as extension taking into account random boundary conditions and a random source/sink for a bounded groundwater flow model. Therefore, a technical note instead of a full research article appears to me the more appropriate form for publication.*

**Response:** We changed the type of this manuscript to a technical note.

3. *In this line, the section on Results and Discussion, especially section 3.1 could be shortened (specific comments later). Most of the publication is well-written, parts (in particular section 2 and 3.1) need to be improved in language, ideally by a native speaker.*

**Response:** We shortened the section 3.1 significantly based on the reviewer's specific suggestions and had a native speaker to edit and improve the writing of this revision.

4. *Figures and tables are in a good shape. However, the number of figures can be reduced by combining Figures 1+2 and Figures 3+4. I highly recommend to prepare an additional figure illustrating the one-dimensional groundwater flow model, including the nomenclature of the relevant processes (time-dependent source/sink, initial conditions, boundary conditions,...) for improving readability.*

**Response:** Thank you for your suggestions. We combined Figures 1 and 2 as the new Figure 2 and combine Figures 3 and 4 as the new Figure 3. We added a new Figure 1 to illustrate the conceptual model studied in response to both this and another reviewer's comment.

### **Specific comments:**

#### *Introduction*

1. *p.3 l.1: What do you mean with "inherently erroneous"? The sentence could be misinterpreted.*

**Response:** To avoid misinterpretation, we replaced this sentence with "It is obvious that errors always exit in the groundwater levels calculated or simulated with analytical or numerical solutions. " in lines 51-52 of the revised manuscript.

2. *p.3 l.6-7: Please specify the sentence "The uncertainties in model parameters were investigated." (e.g. Which parameters? How?).*

**Response:** We rewritten this sentence as "The uncertainties in the model parameters (e.g., hydraulic conductivity, recharge rate, evapotranspiration, and river conductance) were

investigated based on generalized likelihood uncertainty estimation and Bayesian methods (Nowak et al., 2010; Neuman et al., 2012; Rojas et al., 2008; Rojas et al., 2010)” in lines 56-60 of the revised manuscript.

3. *p.3 l.11: Specify "Little attention" (Who?).*

**Response:** We added corresponding citations, i.e., “Little attention has been given to the uncertainties in groundwater level due to temporal variations of hydrological processes, e.g., recharge, evapotranspiration, discharge to a river, and river stage (Bloomfield and Little, 2010; Zhang and Schilling, 2004; Schilling and Zhang, 2012; Liang and Zhang, 2013a; Zhu et al., 2012)” in lines 63-67 of the revised manuscript.

#### *Formulation and solutions*

4. *p.5 l6-7: Please elaborate more on the simplification of setting  $H_0(x)$  to the steady state solution to the one-dimensional transient groundwater flow equation. Why is that an appropriate assumption?*

**Response:** The initial condition has to be specified in order to solve the mathematical model. For a practical problem, the initial condition can be set based on real measurements or aquifer condition. Our study is theoretical with on real measurements and logical initial condition is a relative steady head distribution that is reached in an aquifer after a rainfall or during a wet season. The steady-state solution to this model was often adopted as the initial condition in previous research (Liang and Zhang, 2012, 2013a, b). Thus we set the initial condition  $H_0(x)$  to be the steady-state solution to the one-dimensional groundwater flow equation. We elaborated this simplification in lines 116-124 of the revised manuscript.

5. *p.5 l.11-12: please explain this step on more detail.*

**Response:** There were some typos in the original manuscript. We corrected them in lines 125 – 126 in the revised manuscript, i.e.,

“Thus, the mean and perturbation of  $H_0(x)$  can be written as,  
 $\langle H_0(x) \rangle = h_0 + 0.5 \langle W_0 \rangle (L^2 - x^2) / T$  and  $H_0'(x) = 0.5 W_0' (L^2 - x^2) / T$ , respectively.”

6. *p.6 l.11-12: Give a justification for the assumption of uncorrelated functions. How realistic is that assumption?*

**Response:** This assumption may not be realistic. In general,  $W(t)$ ,  $Q(t)$ , and  $H(t)$  should be correlated. It is possible to consider the relationship among  $W(t)$ ,  $Q(t)$ , and  $H(t)$  by assuming some theoretical correlation functions but the problem is that 1) it is unclear what kind of correlation exists among these variable, 2) there is little observed data to support the type of the correlation assumed, and 3) simple analytical solutions would be difficulty to derive when considering the correlation . Therefore, we studied the case in which such correlation is weak in order to derive some simple analytical solutions. We believe this is an important first step towards solving this complex problem. and it also beyond the scopes of this paper and hope to relax this assumption in our future study.

7. *There are remarkable differences in the style and language of sections 3.1. and 3.2. To me, section 3.1 is much to circumstantial, where section 3.2 is more compact and to the point of interest. Therefore, section 3.1 should be shortened and adapted in style to that of section 3.2. Steps for improving the readability might be:*

- *reducing doubling of explanations (e.g. p.9 l. 14/15) for all cases discussed*
- *not announcing the content of figures (e.g. p.9 l. 17 – p.14 l. 2) for all cases discussed. You may announce the visualization of results in Figure 1 at the beginning of the section and then directly refer to the Figure of interest, without repeating "The dimensionless standard deviation ... was presented in Figure 1..."*
- *Shortening aspects which can obviously be seen in the figure (e.g. p. 12. l. 3-6) "Similar to ...".*
- *Use a more compact description of the results (e.g. entire page 10).*

**Response:** Thank you for the detailed comments to improve our manuscript. Based on your good suggestions we shortened section 3.1 significantly to make it more compact in the revised manuscript.

#### *Conclusions*

8. *p.17 l.10: What is a "typical aquifer studied"? Please formulate in a more generally way. If it is referred to the previous discussed example, please specify. (In general the conclusions drawn should be understandable without knowing details from the previous sections)*

**Response:** We added a note after “typical aquifer studied” to make this conclusion more clear in lines 376-378 in the revised manuscript, i.e., “In the typical sandy aquifer studied (with the length of aquifer at the direction of water flow  $L=100\text{m}$ , the average saturated thickness  $M=10\text{m}$ , hydraulic conductivity  $K=1\text{m/day}$ , and specific yield  $S_Y=0.25$ ).”

9. *p.17 l.15: In both brackets it is stated "low frequencies".*

**Response:** We replaced first “low frequencies” with “high frequencies” in line 367 of the revised manuscript.

#### *Figures and Tables*

10. *Figure 1: The Figure is in general well constructed to show the different impacts of the processes. The readability of the figure and caption text could be improved by:*
- *specifying the difference in the rows before ": (a) and (b)..." (e.g. by stating in the caption "for different combinations of ... (four rows) ")*
  - *write the specific case to the figures (e.g.  $_{2W6=0}$  to Fig. 1c, etc.)*
  - *The range of time values in Fig. 1b and 1d is different to those  $t$  values in Fig. 1f and 1h., where only the second range (those of 1f and 1h.) is state in the caption.*

**Response:** We rewritten this figure caption and added specific text in each graphs according to your suggestion. We think it is clearer in revised manuscript.

11. *Figure 2: There is the same problem with values of  $t$  in Figure 2b and the caption.*

**Response:** We revised it.

12. *Figure 2 should be combined with Figure 1, being sub-figures 1i and 1j.*

**Response:** We did. Please see our response to your comment 4.

13. *Figure 4: Analogously to Figure 2, this Figure should be combined with Figure 3.*

**Response:** We did. Please see our response to your comment 4.

### Technical Corrections

14. *The text needs language improvements, ideally by a native speaker, in particular section 2 and 3.1.*

**Response:** We had a native speaker to edit the manuscript. Please see our response to your comments 3.

15. *p.3 l.21: typo: "temporospatial"*

**Response:** The word "temporospatial" should be allowed

16. *p.9 l. 13: new paragraph starting.*

**Response:** We corrected it.

17. *p.9 l.17: typo in  $\sigma'_h$*

**Response:** We corrected it.

18. *p.11 l. 6-8: Please rephrase. The sentence ("Unlike ...") is hardly understandable.*

**Response:** We rephrased the sentences in lines 222-226.

19. *p.12 l.16: typo "setting $\sigma$ "*

**Response:** We corrected it.

*Technical note:*

**Analyses of Uncertainties and Scaling  
of Groundwater Level Fluctuations**

Xiuyu Liang<sup>a,b</sup>, You-Kuan Zhang<sup>a\*,b,c</sup>

<sup>a</sup>Center for Hydrosociences Research, Nanjing University,

Nanjing, Jiangsu 210093, P.R. China

<sup>b</sup>School of Earth Sciences and Engineering, Nanjing University,

Nanjing, Jiangsu 210093, P.R. China

<sup>c</sup>Department of Geoscience, University of Iowa,

Iowa City, Iowa 52242, USA

\*Corresponding author

Phone: +86 25-83594485, Fax: +86 25-83596670,

Email: ykzhang@nju.edu.cn

*Resubmitted to Hydrology and Earth System Sciences*

*May, 2015*



## Abstract

Analytical solutions for the variance, covariance, and spectrum of groundwater level,  $h(x, t)$ , in an unconfined aquifer described by a linearized Boussinesq equation with random source/sink and initial and boundary conditions were derived. It was found that in a typical aquifer the error in  $h(x, t)$  in early time is mainly caused by the random initial condition and the error reduces as time progresses to reach a constant error in later time. The duration during which the effect of the random initial condition is significant may last a few hundred days in most aquifers. The constant error in  $h(x, t)$  in later time is due to the combined effects of the uncertainties in the source/sink and flux boundary: the closer to the flux boundary, the larger the error. The error caused by the uncertain head boundary is limited in a narrow zone near the boundary and remains more or less constant over time. The aquifer system behaves as a low-pass filter which filters out high-frequency noises and keeps low-frequency variations. Temporal scaling of groundwater level fluctuations exists in most part of a low permeable aquifer whose horizontal length is much larger than its thickness caused by the temporal fluctuations of areal source/sink.

**Key words:** Uncertainty of groundwater levels; Temporal scaling; Random source/sink; Random initial and boundary conditions.

## 1. Introduction

Groundwater level or hydraulic head ( $h$ ) is the main driving force for water flow and advective contaminant transport in aquifers and thus the most important variable studied in groundwater hydrology and its applications. Knowledge about  $h$  is critical in dealing with groundwater-related environmental problems, such as over-pumping, subsidence, sea water intrusion, and contamination. One often found that the data about groundwater level is limited or unavailable in a hydrogeological investigation. In such cases the groundwater level distribution and its temporal variation are usually obtained with an analytical or numerical solution to a groundwater flow model.

It is obvious that errors always exist in the groundwater levels calculated or simulated with analytical or numerical solutions. The main sources of errors include the simplification or approximation in a conceptual model and the uncertainties in the model parameters. Problems in conceptualization or model structure were dealt with by many researchers (Neuman, 2003; Rojas et al., 2010; Ye et al., 2008; Rojas et al., 2008; Refsgaard et al., 2007; Zeng et al., 2013). The uncertainties in the model parameters (e.g., hydraulic conductivity, recharge rate, evapotranspiration, and river conductance) were investigated based on generalized likelihood uncertainty estimation and Bayesian methods (Nowak et al., 2010; Neuman et al., 2012; Rojas et al., 2008; Rojas et al., 2010). The uncertainty in groundwater level has been one of the main research topics in stochastic subsurface hydrology for more than three decades. Most of these studies were focused on the spatial variability of groundwater

level due to aquifers' heterogeneity (Dagan, 1989;Gelhar, 1993;Zhang, 2002). Little attention has been given to the uncertainties in groundwater level due to temporal variations of hydrological processes, e.g., recharge, evapotranspiration, discharge to a river, and river stage (Bloomfield and Little, 2010;Zhang and Schilling, 2004;Schilling and Zhang, 2012;Liang and Zhang, 2013a;Zhu et al., 2012).

Uncertainties of groundwater level fluctuations have been studied by Zhang and Li (2005, 2006) and most recently by Liang and Zhang (2013a). Based on a linear reservoir model with a white noise or temporally-correlated recharge process, Zhang and Li (2005, 2006) derived the variance and covariance of  $h(t)$  by considering only a random source or sink process assuming deterministic initial and boundary conditions. Liang and Zhang (2013a) extended the studies of Zhang and Li (2005, 2006) and carried out non-stationary spectral analysis and Monte Carlo simulations using a linearized Boussinesq equation, and investigated the temporospatial variations of groundwater level. However, the only random process considered by Liang and Zhang (2013a) is the source/sink. Temporal scaling of groundwater levels discovered first by Zhang and Schilling (2004) was verified in several studies (Zhang and Li, 2005, 2006; Bloomfield and Little, 2010; Zhang and Yang, 2010; Zhu et al., 2012; Schilling and Zhang, 2012). However, we do not know the effect of random boundary conditions on temporal scaling of groundwater levels.

In this study we extended above-mentioned work by considering the groundwater flow in a bounded aquifer described by a linearized Boussinesq equation with a random source/sink as well as random initial and boundary

conditions since the latter processes are known with uncertainties. The objectives of this study are 1) to derive analytical solutions for the covariance, variance and spectrum of groundwater level, and 2) to investigate the individual and combined effects of these random processes on uncertainties and scaling of  $h(x, t)$ . In the following we will first present the formulation and analytical solutions, then discuss the results, and finally draw some conclusions.

## 2. Formulation and Solutions

Under the Dupuit assumption, the one-dimensional transient groundwater flow in an unconfined aquifer near a river (Fig. 1) can be approximated with the linearized Boussinesq equation (Bear, 1972) with the initial and boundary conditions, i.e.,

$$T \frac{\partial^2 h}{\partial x^2} + W(t) = S_y \frac{\partial h}{\partial t} \quad (1a)$$

$$h(x, t)|_{t=0} = H_0(x); \quad T \frac{\partial h}{\partial x} \Big|_{x=0} = Q(t); \quad h(x, t)|_{x=L} = H(t) \quad (1b)$$

where  $T$  [L/T] is the transmissivity,  $h$  [L] is the hydraulic head or groundwater level above the bottom of the aquifer which is assumed to be horizontal,  $W(t)$  [L/T] is the time-dependent source/sink term representing areal recharge or evapotranspiration,  $S_y$  is the specific yield,  $H_0(x)$  [L] is the initial condition,  $Q(t)$  [L<sup>2</sup>/T] is the time-dependent flux at the left boundary,  $H(t)$  [L] is the time-dependent water level at the right boundary,  $L$  [L] is distance from the left to the right boundary,  $x$  [L] is the coordinate, and  $t$  [T] is time. In this study the initial head  $H_0(x)$  is taken to be a spatially random variable, and the source/sink,  $W(t)$ , the flux to the left boundary,  $Q(t)$ , and the head at the right boundary,  $H(t)$ , are all taken to be temporally random

processes and spatially deterministic. The parameters  $T$  and  $S_Y$  are taken to be constant.

The groundwater level,  $h(x, t)$ , the three random processes,  $W(t)$ ,  $Q(t)$ , and  $H(t)$ , and the random variable,  $H_0(x)$ , are expressed in terms of their respective ensemble means plus small perturbations,

$$h(x, t) = \langle h(x, t) \rangle + h'(x, t) \quad (2a)$$

$$W(t) = \langle W(t) \rangle + W'(t); \quad Q(t) = \langle Q(t) \rangle + Q'(t) \quad (2b)$$

$$H(t) = \langle H(t) \rangle + H'(t); \quad H_0(x) = \langle H_0(x) \rangle + H_0'(x) \quad (2c)$$

where  $\langle \rangle$  stands for ensemble average and  $'$  for perturbation. The initial condition  $H_0(x)$  in (1) can be any function. For the conceptualization of the groundwater flow presented in Fig. 1, the steady-state condition can be reached in this aquifer after a rainfall or during a wet season. Thus the steady-state solution to this model were often adopted as initial condition in previous research (Liang and Zhang, 2012, 2013a, b). Thus, in this study, we set initial condition  $H_0(x)$  to be the steady-state solution to the one-dimensional groundwater flow equation, i.e.,  $H_0(x) = h_0 + 0.5W_0(L^2 - x^2)/T$ , where  $h_0$  [L] is the constant groundwater level at the right boundary and  $W_0$  [L/T] is the spatially constant recharge rate (Liang and Zhang, 2012). Since  $h_0$  is taken to be constant, the source of the uncertainty in the initial head  $H_0(x)$  is due to random  $W_0$  only. Thus, the mean and perturbation of  $H_0(x)$  can be written as,  $\langle H_0(x) \rangle = h_0 + 0.5\langle W_0 \rangle(L^2 - x^2)/T$  and  $H_0'(x) = 0.5W_0'(L^2 - x^2)/T$ , respectively. By substituting Eq. (2),  $\langle H_0(x) \rangle$ , and  $H_0'(x)$  into Eq. (1) and taking expectation, one obtains the mean flow equation with the mean initial and boundary conditions as

$$T \frac{\partial^2 \langle h \rangle}{\partial x^2} + \langle W \rangle = S_Y \frac{\partial \langle h \rangle}{\partial t} \quad (3a)$$

$$\langle h(x,0) \rangle = h_0 + \frac{\langle W_0 \rangle}{2T} (L^2 - x^2); \quad T \frac{\partial \langle h \rangle}{\partial x} \Big|_{x=0} = \langle Q \rangle; \quad \langle h(L,t) \rangle = \langle H(t) \rangle \quad (3b)$$

Subtracting Eq. (3) from (1) leads to the following perturbation equation with the initial and boundary conditions

$$T \frac{\partial^2 h'}{\partial x^2} + W' = S_y \frac{\partial h'}{\partial t} \quad (4a)$$

$$h'(x,0) = \frac{W_0'}{2T} (L^2 - x^2); \quad T \frac{\partial h'}{\partial x} \Big|_{x=0} = Q'; \quad h'(L,t) = H'(t) \quad (4b)$$

The analytical solution to Eq. (4) can be derived with integral-transform methods (Ozisik, 1968) given by

$$h' = \frac{2}{L} \sum_{n=0}^{\infty} e^{-\beta b_n^2 t} \cos(b_n x) \left[ \frac{(-1)^n}{b_n^3 T} W_0' + \beta \int_0^t e^{\beta b_n^2 \xi} \left[ \frac{(-1)^n}{T b_n} W'(\xi) - \frac{Q'(\xi)}{T} + H'(\xi) (-1)^n b_n \right] d\xi \right] \quad (5)$$

where  $\beta = T / S_y$ ,  $b_n = (2n+1)\pi / (2L)$ . Using Eq. (5), the temporal covariance of the groundwater level fluctuations can be derived as

$$\begin{aligned} C_{hh}(x, t_1; x, t_2) &= E[h'(x, t_1) h'(x, t_2)] \\ &= \frac{4}{L^2} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} e^{-\beta(b_m^2 t_1 + b_n^2 t_2)} \cos(b_m x) \cos(b_n x) \left[ \frac{(-1)^{m+n}}{T^2 b_m^3 b_n^3} \sigma_{W_0}^2 \right. \\ &\quad \left. + \beta^2 \int_0^{t_1} \int_0^{t_2} e^{\beta(b_m^2 \xi + b_n^2 \rho)} \left[ \frac{(-1)^{n+m}}{T^2 b_m b_n} C_{WW}(\xi, \rho) + \frac{C_{QQ}(\xi, \rho)}{T^2} + C_{HH}(\xi, \rho) (-1)^{m+n} b_m b_n \right] d\xi d\rho \right] \end{aligned} \quad (6)$$

in which  $\sigma_{W_0}^2$  is the variance of  $W_0$ , and  $C_{WW}(\xi, \rho)$ ,  $C_{QQ}(\xi, \rho)$  and  $C_{HH}(\xi, \rho)$  are the temporal auto-covariance of  $W(t)$ , of  $Q(t)$ , and  $H(t)$ , respectively. We assume that  $W(t)$ ,  $Q(t)$ , and  $H(t)$  are uncorrelated in order to simplify our analyses. It is shown in Eq. (6) that the head covariance depends on the variance of  $W_0$  and the covariances of  $W(t)$ ,  $Q(t)$ , and  $H(t)$  and this equation can be evaluated for any random  $W(t)$ ,  $Q(t)$ , and  $H(t)$ . We assume that these processes are white noises as employed in previous

studies (Gelhar, 1993; Hantush and Marino, 1994; Liang and Zhang, 2013a). More

realistic randomness of these processes will be considered in future studies.

Following Gelhar (1993, p.34), we express the spectra of  $W(t)$ ,  $Q(t)$ , and  $H(t)$  as

$S_{WW} = \sigma_W^2 \lambda_W / \pi$ ,  $S_{QQ} = \sigma_Q^2 \lambda_Q / \pi$ , and  $S_{HH} = \sigma_H^2 \lambda_H / \pi$ , respectively, where  $\sigma_W^2$ ,

$\sigma_Q^2$ , and  $\sigma_H^2$  are the variances and  $\lambda_W$ ,  $\lambda_Q$ , and  $\lambda_H$  are the correlation time

intervals of these three processes, respectively. The corresponding covariance of

$W(t)$ ,  $Q(t)$  and  $H(t)$  are  $C_{WW}(\xi, \rho) = 2\sigma_W^2 \lambda_W \delta(\xi - \rho)$ ,  $C_{QQ}(\xi, \rho) = 2\sigma_Q^2 \lambda_Q \delta(\xi - \rho)$ ,

and  $C_{HH}(\xi, \rho) = 2\sigma_H^2 \lambda_H \delta(\xi - \rho)$ . Substituting these covariance into (6) and taking

integration, one obtain analytical solution of head covariance

$$C_{hh}(x', t', \tau') = \frac{4\beta L^2}{T^2} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \cos(b'_m x') \cos(b'_n x') \left\{ e^{-\left[ (b_m'^2 + b_n'^2) t' + (b_n'^2 - b_m'^2) \frac{\tau'}{2} \right]} \frac{L^2 (-1)^{m+n} \sigma_{W_0}^2}{\beta b_m'^3 b_n'^3} \right. \\ \left. + 2 \frac{(e^{-b_m'^2 \tau'} - e^{-2b_m'^2 t'})}{(b_m'^2 + b_n'^2)} \left[ \frac{(-1)^{m+n} \sigma_W^2 \lambda_W}{b'_m b'_n} + \frac{\sigma_Q^2 \lambda_Q}{L^2} + \frac{(-1)^{m+n} b'_m b'_n T^2 \sigma_H^2 \lambda_H}{L^4} \right] \right\} \quad (7)$$

where  $\tau' = t'_2 - t'_1$  and  $t' = (t'_2 + t'_1) / 2$ . The analytical solution for the head variance can

be obtain by setting  $\tau' = 0$

$$\sigma_h^2(x', t') = \frac{4\beta L^2}{T^2} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \cos(b'_m x') \cos(b'_n x') \left\{ e^{-(b_m'^2 + b_n'^2) t'} \frac{L^2 (-1)^{m+n} \sigma_{W_0}^2}{\beta b_m'^3 b_n'^3} + \right. \\ \left. 2 \frac{1 - e^{-2b_m'^2 t'}}{(b_m'^2 + b_n'^2)} \left[ \frac{(-1)^{m+n} \sigma_W^2 \lambda_W}{b'_m b'_n} + \frac{\sigma_Q^2 \lambda_Q}{L^2} + \frac{(-1)^{m+n} b'_m b'_n T^2 \sigma_H^2 \lambda_H}{L^4} \right] \right\} \quad (8)$$

where

$$t' = \frac{t}{t_c}; \quad x' = \frac{x}{L}; \quad t_c = \frac{L^2}{\beta}; \quad b'_n = \frac{(2n+1)\pi}{2}$$

in which  $t_c (= S_Y L^2 / (KM)) [1/T]$  is a characteristic timescale (Gelhar, 1993) where

the transmissivity ( $T$ ) is replaced by the product of the hydraulic conductivity ( $K$ ) and

the average saturated thickness ( $M$ ) of the aquifer. The characteristic timescale ( $t_c$ ) is

an important parameter and its value for most shallow aquifers is usually larger than 100 day since the horizontal extent of a shallow aquifer is usually much larger than its thickness. For instance, the value of  $t_c$  is 250 days for a sandy aquifer with  $L=100\text{m}$ ,  $M=10\text{m}$ ,  $K=1\text{m/day}$ , and  $S_Y=0.25$ .

The spectral density of  $h(x, t)$  can't be derived by ordinary Fourier transform since the head covariance and variance depend on time  $t'$  and thus  $h(x, t)$  are temporally non-stationary as shown in Eqs. (7) and (8). Priestley (1981) defined the spectral density of non-stationary processes (Wigner spectrum) as the Fourier transform of time-dependent auto-covariance with fixed reference time  $t$  and derived time-dependent spectral density. In order to obtain the spectrum of  $h(x, t)$ , we applied Priestley's method and obtained the time-dependent spectral density (Priestley, 1981; Zhang and Li, 2005; Liang and Zhang, 2013a), i.e.,

$$\begin{aligned}
 S_{hh}(x, t, \omega) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} C_{hh}(x, t, \tau) e^{-i\omega\tau} d\tau \\
 &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \cos(b_m x) \cos(b_n x) \frac{2t_c (b_n^2 - b_m^2) e^{-\beta(b_m^2 + b_n^2)t}}{\beta^2 (b_n^2 - b_m^2)^2 / 4 + \omega^2} \frac{(-1)^{m+n} \sigma_{W_0}^2}{\pi T^2 b_m^3 b_n^3} + \\
 &\quad \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \cos(b_m x) \cos(b_n x) \frac{8\beta b_m^2}{t_c (b_m^2 + b_n^2)} \frac{1}{\beta^2 b_m^4 + \omega^2} \left[ \frac{(-1)^{m+n} S_{WW}}{T^2 b_m b_n} + \frac{S_{QQ}}{T^2} + (-1)^{m+n} b_m b_n S_{HH} \right]
 \end{aligned} \tag{9}$$

where  $\omega$  is angular frequency and  $\omega = 2\pi f$ ,  $f$  is frequency, and  $i = \sqrt{-1}$ . It is seen in Eq. (9) that the spectrum  $S_{hh}$  is dependent on not only frequency and locations but also time  $t$ . The time-dependent term (i.e., first term) in Eq. (9) is caused by the random initial condition and is proportional to  $e^{-\beta(b_m^2 + b_n^2)t}$  which decays quickly with  $t$ . We evaluated the first term in the Eq. (9) by setting  $t=0$  and found that it is much smaller than the second term in Eq. (9). We thus ignored the first term and evaluated the spectrum using the approximation,



$$S_{hh}(x', \omega) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{8\beta b_m'^2 \cos(b_m' x') \cos(b_n' x')}{t_c (b_m'^2 + b_n'^2) (\beta^2 b_m'^4 / L^4 + \omega^2)} \left[ \frac{(-1)^{m+n} S_{WW} L^2}{T^2 b_m' b_n'} + \frac{S_{QQ}}{T^2} + \frac{(-1)^{m+n} b_m' b_n' S_{HH}}{L^2} \right] \quad (10)$$

### 3. Results and Discussion

#### 3.1 Variance of groundwater levels

The general expression of the head variance in Eq. (8) depends on the variances of the four random processes,  $\sigma_{W_0}^2$ ,  $\sigma_W^2$ ,  $\sigma_Q^2$ , and  $\sigma_H^2$ . In the following we will study their individual and combined effects on the head variation and focus our attention only on the variance of  $h(x, t)$ . The dimensionless standard deviation of  $h(x, t)$ ,  $\sigma'_h$ , or the square root of the dimensionless variance ( $\sigma'^2_h$ ) as a function of the dimensionless time ( $t'$ ) were evaluated and presented in the left column of Fig. 2 at fixed dimensionless locations ( $x'$ ). The  $\sigma'_h$  as a function of  $x'$  was evaluated and presented in the right column of Fig. 2 at fixed  $t'$ .

We first evaluate the effect of the random initial condition due to the random term,  $W_0$ , by setting  $\sigma_W^2 = \sigma_Q^2 = \sigma_H^2 = 0$ . In this case the dimensionless variance in Eq. (8) reduces to

$$\sigma'^2_h(x', t') = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(-1)^{m+n}}{b_m'^3 b_n'^3} \cos(b_m' x') \cos(b_n' x') e^{-(b_m'^2 + b_n'^2) t'} \quad (11)$$

where  $\sigma'^2_h = \sigma_h^2 T^2 / (4L^4 \sigma_{W_0}^2)$ . The changes of the  $\sigma'_h$  with  $x'$  and  $t'$  were presented in Fig 2a and 2b, respectively. It is shown in Fig. 2a that for a fixed location the  $\sigma'_h$  is at its maximum at  $t'=0$  and decreases with time gradually to a negligible number at  $t'=1.0$ . This means that the error in  $h(x, t)$  predicted by an analytical or numerical solution due to the uncertain initial condition is significant at

early time, especially near a flux boundary. The time duration during which the effect of the uncertain initial condition is significant depends on the value of the characteristic timescale ( $t_c$ ) since  $t' = t/t_c$ . In the most aquifers this duration may last many days. In the typical aquifer studied the effect of the uncertainty in initial condition on  $h(x, t)$  is significant during first 250 days ( $t' = 1.0$ ). This duration should be relatively short, however, in a more permeable aquifer whose horizontal extent ( $L$ ) is relatively smaller than its thickness ( $M$ ). It is seen in Fig. 2b that for a fixed time, the  $\sigma'_h$  is the largest at the left flux boundary ( $x' = 0.0$ ) and becomes zero at the right constant head boundary ( $x' = 1.0$ ) since the right boundary is deterministic. This means that the error in  $h(x, t)$  predicted by an analytical or numerical solution due to the uncertain initial condition is significant almost everywhere in the aquifer: the further away from a constant head boundary, the larger the error.

We then consider the uncertainty in the areal source/sink term ( $W$ ) by setting

$\sigma_{w_0}^2 = \sigma_Q^2 = \sigma_H^2 = 0$ . In this case the dimensionless variance in Eq. (8) reduces to

$$\sigma_h'^2(x', t') = 2 \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} c_o(b'_m x') c_o(b'_n x') \frac{(1 - e^{-2b_m'^2 t'}) (-1)^{m+n}}{(b_m'^2 + b_n'^2) b'_m b'_n} \quad (12)$$

where  $\sigma_h'^2 = \sigma_h^2 TS_Y / (4L^2 \sigma_W^2 \lambda_W)$ . The changes of the  $\sigma'_h$  with  $x'$  and  $t'$  were presented in Fig 2c and 2d, respectively. It is noticed in Fig. 2c that at a fixed location, the  $\sigma'_h$  is zero initially, gradually increases as time goes, and approaches a constant limit at later time. This means that the error in  $h(x, t)$  due to an source/sink is at its minimum at early time and increases with time to approach a constant limit at later time: the closer to the left flux boundary, the larger the limit. For a fixed time the  $\sigma'_h$  decreases smoothly from the left to the right boundary (Fig. 2d). The error in  $h(x,$

$t$ ) due to the uncertainty in the source/sink is significant almost everywhere in the aquifer: the further away from the constant head boundary, the larger the error, similar to the previous case with the random initial condition (Fig. 2b).

Thirdly, we investigate the effect of the left random flux boundary by setting  $\sigma_{W_0}^2 = \sigma_W^2 = \sigma_H^2 = 0$  in Eq. (8). In this case the dimensionless head variance is given by

$$\sigma_h'^2(x', t') = 2 \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \cos(b'_m x') \cos(b'_n x') \frac{1 - e^{-2b_m'^2 t'}}{b_m'^2 + b_n'^2} \quad (13)$$

where  $\sigma_h'^2 = \sigma_h^2 TS_Y / (4\sigma_Q^2 \lambda_Q)$ . The changes of the  $\sigma_h'$  with  $x'$  and  $t'$  were presented in Fig 2e and 2f, respectively. At any location the  $\sigma_h'$  in Fig. 2e or the error in  $h(x, t)$  due to an uncertain flux boundary is at its minimum at early time and increases quickly with time to approach a constant limit: the closer to the left flux boundary, the larger the limit. At any time the  $\sigma_h'$  in Fig. 2f or the error in the head due to the uncertain flux boundary is at its maximum at the left boundary but decreases quickly away from the boundary to become insignificant for  $x' > 0.8$ .

Fourthly, we investigated the effect of the random head boundary by setting  $\sigma_{W_0}^2 = \sigma_W^2 = \sigma_Q^2 = 0$  in Eq. (8). The dimensionless head variance in this case is given by

$$\sigma_h'^2(x', t') = 2 \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \cos(b'_m x') \cos(b'_n x') \frac{(-1)^{m+n} b'_m b'_n (1 - e^{-2b_m'^2 t'})}{(b_m'^2 + b_n'^2)} \quad (14)$$

where  $\sigma_h'^2 = \sigma_h^2 L^2 S_Y / (4T\sigma_H^2 \lambda_H)$ . The changes of this  $\sigma_h'$  with  $x'$  and  $t'$  were presented in Fig 2g and 2h, respectively. It seen in Fig. 2g that at any location the  $\sigma_h'$  or the error in  $h(x, t)$  due to the random head boundary increases with time

quickly to approach a constant limit: the closer to the uncertain head boundary, the larger the error. The spatial variation of  $\sigma'_h$  can be clearly observed in Fig. 2h for fixed  $t'$ . At any time  $\sigma'_h$  is at its maximum at the right boundary ( $x'=1$ ) where the head is uncertain, decreases quickly away from the boundary. The error in  $h(x, t)$  due to the uncertain head boundary is limited in a narrow zone near the boundary ( $x'>0.8$ ) (Fig. 2h).

Finally, we consider the combined effects of the uncertainties from all four sources, i.e., the initial condition, sources, and flux and head boundaries. The head variance in Eq. (8) is written in the dimensionless form as

$$\sigma'^2_h(x', t') = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \cos(b'_m x') \cos(b'_n x') \left\{ e^{-(b'^2_m + b'^2_n)t'} \frac{(-1)^{m+n} \sigma'^2_{w_0}}{b'^3_m b'^3_n} + 2 \frac{1 - e^{-2b'^2_m t'}}{(b'^2_m + b'^2_n)} \left[ \frac{(-1)^{m+n}}{b'_m b'_n} + \sigma'^2_Q + (-1)^{m+n} b'_m b'_n \sigma'^2_H \right] \right\} \quad (15)$$

where

$$\sigma'^2_h = \frac{\sigma^2_h T S_Y}{4 L^2 \sigma^2_w \lambda_w}; \quad \sigma'^2_{w_0} = \frac{L^2 S_Y \sigma^2_{w_0}}{T \sigma^2_w \lambda_w}; \quad \sigma'^2_Q = \frac{\sigma^2_Q \lambda_Q}{L^2 \sigma^2_w \lambda_w}; \quad \sigma'^2_H = \frac{T^2 \sigma^2_H \lambda_H}{L^4 \sigma^2_w \lambda_w}$$

The dimensionless variances,  $\sigma'^2_{w_0}$ ,  $\sigma'^2_Q$  and  $\sigma'^2_H$ , need to be specified in order to evaluate the dimensionless  $\sigma'^2_h(x', t')$  in Eq. (15). For the typical aquifer mentioned above with  $L=100\text{m}$ ,  $T=10 \text{ m}^2/\text{day}$  (or  $K=1\text{m}/\text{day}$  and  $M=10\text{m}$ ) and  $S_Y=0.25$ , we set  $\sigma^2_{w_0}/(\sigma^2_w \lambda_w) = 10^{-1}$ ,  $\sigma^2_Q \lambda_Q/(\sigma^2_w \lambda_w) = 10^3$ ,  $\sigma^2_H \lambda_H/(\sigma^2_w \lambda_w) = 10^4$  and obtain  $\sigma'^2_{w_0} = 25$ ,  $\sigma'^2_Q = 0.1$  and  $\sigma'^2_H = 0.01$ .

The changes of this  $\sigma'_h$  with  $x'$  and  $t'$  were presented in Fig 2i and 2j, respectively. It is observed in Fig. 2i that at any location the  $\sigma'_h$  is at its maximum

due to the uncertainty in the initial condition, gradually decreases as time goes, and approaches a constant limit at later time ( $t' > 0.6$ ) which is due to the combined effects of the uncertain source/sink and flux and head boundaries. This means that the error in the head in early time is significant if the initial condition is uncertain and reduces as time goes to reach a constant limit. The error in head in later time is determined by the uncertainties in the source/sink, flux and head boundaries. It can be observed in Fig. 2j that  $\sigma'_h$  is relatively larger near both boundaries. The values of  $\sigma'_h$  at the two boundaries are equivalent ( $\sim 1.3$ ) at early time, say  $t' = 0.01$  (the top curve in Fig. 2j) and it reduces slowly away from the flux boundary but quickly away from the head boundary. As time progresses, the  $\sigma'_h$  near the head boundary stays more or less the same but reduces significantly in most part of the aquifer. This means that in early time the error in  $h(x, t)$  in most part of the aquifer is mainly caused by the initial condition and at later time it is due to the combined effects of the uncertain areal source/sink and flux boundary. The effect of the uncertain head boundary on  $h(x, t)$  doesn't change with time significantly but is limited in a narrow zone near the boundary.

### 3.2 Spectrum of groundwater levels

We first evaluated  $S_{hh}$  in Eq. (10) due to the effect of the white noise flux boundary only by setting  $S_{QQ} \neq 0$ ,  $S_{WW} = 0$ , and  $S_{HH} = 0$ . The dimensionless spectrum  $S_{hh}/S_{QQ}$  as a function of the frequency ( $f$ ) was evaluated and presented in the log-log plot (Fig. 3a-3c) for three values of  $t_c$  (40, 400, and 4,000 days) since the value of  $t_c$  is 250 days for a sandy aquifer as we mentioned above and at the six

locations ( $x' = 0.0, 0.2, 0.4, 0.6, 0.8, \text{ and } 0.9$ ). The spectrum  $S_{hh}/S_{QQ}$  in Fig. 3a is more or less horizontal (i.e., white noise) at low frequencies and decrease gradually as  $f$  increases, indicating that an aquifer acts as a low-bass filter that filter signals at high frequencies and keep signals at low frequencies. The aquifer has significantly dampened the fluctuations of the groundwater level. The spectrum varies with the location  $x'$ : the smaller the value of  $x'$  or the closer to the left flux boundary ( $x'=0$ ), the larger the spectrum (Fig. 3a-3c). All spectra in Fig. 3a are not a straight line in the log-log plot, meaning that the temporal scaling of  $h(x, t)$  doesn't exist in the range of  $f = 10^{-3} \sim 10^0$  when  $t_c=40$  days. As  $t_c$  increases to 400 and 4000 days, however, the spectrum at  $x'=0$  become a straight line (the top curve in Fig. 3b and 3c) or has a power-law relation with  $f$ , i.e.,  $S_{hh}/S_{QQ} \propto 1/f$ , since its slope is approximately one. The fluctuations of  $h(0, t)$  is a pink noise due to the white noise fluctuations flux boundary when the characteristic timescale ( $t_c$ ) is large which means that the aquifer is relatively less permeable and/or has a much larger horizontal length than its thickness.

Secondly, the spectrum  $S_{hh}/S_{HH}$  due to the sole effect of the random head boundary was evaluated by setting  $S_{HH} \neq 0$ ,  $S_{ww} = 0$ , and  $S_{QQ} = 0$  in Eq. (10) for the same three values of  $t_c$  and six locations and presented in Fig. 3d-3f as a function of  $f$ . It is shown that similar to Fig. 3a-3c, the spectrum decreases as  $f$  increases but different from Fig. 3a-3c, the spectrum is larger at  $x'=0.9$  near the right boundary (the top curves in Fig. 3d-3f) than that  $x'=0.0$  (the bottom curves). Furthermore, none of the spectra are a straight line in the log-log plot, indicating that the temporal

scaling of groundwater level fluctuations doesn't exist in the case of the white noise head boundary.

Thirdly, the spectrum  $S_{hh}/S_{ww}$  due the effect of the white noise recharge only was evaluated by setting  $S_{ww} \neq 0$ ,  $S_{qq} = 0$ , and  $S_{HH} = 0$  in Eq. (10) for the same values of  $t_c$  and  $x'$  and presented in Fig. 3g-3i as a function of  $f$ . It is shown that when  $t_c=40$  day the spectrum in Fig. 3g is horizontal at low frequencies and become a straight line at high frequencies: the closer to the right head boundary, the later it approaches a straight line (Fig. 3h). As  $t_c$  increases to 400 and 4000 days, the slope of the spectrum at all locations except at  $x'=0.9$  approaches to a straight line with a slope of 2 (Fig. 3h and 3i), indicating a temporal scaling of  $h(x, t)$ . The fluctuations of groundwater level is a Brownian motion, i.e.,  $S \propto 1/f^2$ , when  $t_c \geq 4000$  day or in a relatively less permeable and/or has a much larger horizontal length than its thickness.

Finally, the head spectrum due to the combined effect of all three random sources (the white noise recharge, and flux and head boundaries) was evaluated, i.e.,  $S_{ww} \neq 0$ ,  $S_{qq} \neq 0$ , and  $S_{HH} \neq 0$  in Eq. (10). The spectrum of  $S_{hh}/S_{ww}$  as a function of  $f$  was presented in Fig. 3j-3l for the same values of  $t_c$  and  $x'$  where  $S_{qq}/S_{ww} = 1000$  and  $S_{HH}/S_{ww} = 10000$  which are same with the values using in previous section. It is noticed that the general patterns of  $S_{hh}/S_{ww}$  in the combined case is similar to the case under the random source/sink only (Fig. 3g-3i) except at  $x'=0.0$  and  $0.9$  (the dashed and dotted curves in Fig. 3j, respectively) due to the strong effects of the boundary conditions at these two locations. At  $t_c=4000$

day, the spectra at all locations except  $x'=0.0$  (Fig. 3l) are similar to those in Fig. 3i, indicating the dominating effect of the random areal source/sink. The spectrum at  $x'=0$  in this case is also a straight line (the dashed curve in Fig. 3l) but with a different slope due to the effect of the random flux boundary which is similar to the top straight line in Fig. 3c. Above results provide a theoretical explanation as why temporal scaling exists in the observed groundwater level fluctuations (Zhang and Schilling, 2004; Bloomfield and Little, 2010; Zhu et al., 2012). We thus conclude that temporal scaling of  $h(x, t)$  may indeed exist in real aquifers due to the strong effect of the areal source/sink.

#### 4. Conclusions

In this study the effects of random source/sink, and initial and boundary conditions on the uncertainty and temporal scaling of the groundwater level,  $h(x, t)$  were investigated. The analytical solutions for the variance, covariance and spectrum of  $h(x, t)$  in an unconfined aquifer described by a linearized Boussinesq equation with white noise source/sink, and initial and boundary conditions were derived. The standard deviations of  $h(x, t)$  for various cases were evaluated. Based on the results, the following conclusions can be drawn.

1. The error in  $h(x, t)$  due to a random initial condition is significant at early time, especially near a flux boundary. The duration during which the effect is significant may last a few hundred days in most aquifers;

2. The error in  $h(x, t)$  due to a random areal source/sink is significant in most part of an aquifer: the closer to a flux boundary, the larger the error;



3. The errors in  $h(x, t)$  due to random flux and head boundaries are significant near the boundaries: the closer to the boundaries, the larger the errors. The random flux boundary may affect the head over a larger region near the boundary than the random head boundary;

4. In the typical sandy aquifer studied (with the length of aquifer at the direction of water flow  $L=100\text{m}$ , the average saturated thickness  $M=10\text{m}$ , hydraulic conductivity  $K=1\text{m/day}$ , and specific yield  $S_y=0.25$ ) the error in  $h(x, t)$  in early time is mainly caused by an uncertain initial condition and the error reduces as time goes to reach a constant error in later time. The constant error in  $h(x, t)$  is mainly due to the combined effects of uncertain source/sink and boundaries;

5. The aquifer system behaves as a low-pass filter which filter the short-term (high frequencies) fluctuations and keep the long-term (low frequencies) fluctuations;

6. Temporal scaling of groundwater level fluctuations may indeed exist in most part of a low permeable aquifer whose horizontal length is much larger than its thickness caused by the temporal fluctuations of areal source/sink.

Finally, it is pointed out that the analyses carried out in this study is under the assumptions that the processes,  $W(t)$ ,  $Q(t)$ , and  $H(t)$  are uncorrelated white noises. In reality, they may be correlated and spatially varied. We plan to relax those constraints and study more realistic cases in the near future. It is also noted that the analytical solutions for head variances derived in this study provide a way to identify and quantify the uncertainty. The spectrum relationship obtained among the head,

recharge and boundary conditions can help one to improve spectrum analysis for a groundwater level time series and removed the effects of the boundary conditions.

## Acknowledgment

This study was partially supported with the research grants from the National Nature Science Foundation of China (NSFC-41272260; NSFC-41330314; NSFC-41302180), the Natural Science Foundation of Jiangsu Province (SBK201341336) and from the National Key project “Water Pollution Control in the Huai River Basin” of China (2012ZX07204-001, 2012ZX07204-003).

## References

- Bear, J.: Dynamics of fluids in porous media, Environmental science series (New York, 1972- ), American Elsevier Pub. Co., New York., xvii, 764 p. pp., 1972.
- Beven, K., and Binley, A.: The Future of Distributed Models - Model Calibration and Uncertainty Prediction, Hydrol Process, 6, 279-298, 1992.
- Bloomfield, J. P., and Little, M. A.: Robust evidence for random fractal scaling of groundwater levels in unconfined aquifers, J Hydrol, 393, 362-369, DOI 10.1016/j.jhydrol.2010.08.031, 2010.
- Dagan, G.: Flow and transport in porous formations, Springer-Verlag, Berlin ; New York, xvii, 465 p. pp., 1989.
- Gelhar, L. W.: Stochastic subsurface hydrology, Prentice-Hall, Englewood Cliffs, N.J., x, 390 p. pp., 1993.
- Hantush, M. M., and Marino, M. A.: One-Dimensional Stochastic-Analysis in Leaky Aquifers Subject to Random Leakage, Water Resour Res, 30, 549-558, Doi 10.1029/93wr02887, 1994.
- Liang, X. Y., and Zhang, Y. K.: A new analytical method for groundwater recharge and discharge estimation, J Hydrol, 450, 17-24, DOI 10.1016/j.jhydrol.2012.05.036, 2012.
- Liang, X. Y., and Zhang, Y. K.: Temporal and spatial variation and scaling of groundwater levels in a bounded unconfined aquifer, J Hydrol, 479, 139-145, DOI 10.1016/j.jhydrol.2012.11.044, 2013a.
- Liang, X. Y., and Zhang, Y. K.: Analytic solutions to transient groundwater flow under time-dependent sources in a heterogeneous aquifer bounded by fluctuating river stage, Adv Water Resour, 58, 1-9, DOI 10.1016/j.advwatres.2013.03.010, 2013b.
- Neuman, S. P.: Maximum likelihood Bayesian averaging of uncertain model predictions, Stoch Env Res Risk A, 17, 291-305, 10.1007/s00477-003-0151-7, 2003.
- Neuman, S. P., Xue, L., Ye, M., and Lu, D.: Bayesian analysis of data-worth considering model and parameter uncertainties, Advances in Water Resources, 36, 75-85, 10.1016/j.advwatres.2011.02.007, 2012.


- Nowak, W., de Barros, F. P. J., and Rubin, Y.: Bayesian geostatistical design: Task-driven optimal site investigation when the geostatistical model is uncertain, *Water Resour Res*, 46, Artn W03535 Doi 10.1029/2009wr008312, 2010.
- Priestley, M. B.: *Spectral Analysis and Time Series*, Academic Press, San Diego, 1981.
- Refsgaard, J. C., van der Sluijs, J. P., Hojberg, A. L., and Vanrolleghem, P. A.: Uncertainty in the environmental modelling process - A framework and guidance, *Environmental Modelling & Software*, 22, 1543-1556, 10.1016/j.envsoft.2007.02.004, 2007.
- Rojas, R., Feyen, L., and Dassargues, A.: Conceptual model uncertainty in groundwater modeling: Combining generalized likelihood uncertainty estimation and Bayesian model averaging, *Water Resources Research*, 44, 10.1029/2008wr006908, 2008.
- Rojas, R., Feyen, L., Batelaan, O., and Dassargues, A.: On the value of conditioning data to reduce conceptual model uncertainty in groundwater modeling, *Water Resources Research*, 46, 20, 10.1029/2009wr008822, 2010.
- Schilling, K. E., and Zhang, Y. K.: Temporal Scaling of Groundwater Level Fluctuations Near a Stream, *Ground Water*, 50, 59-67, DOI 10.1111/j.1745-6584.2011.00804.x, 2012.
- Vrugt, J. A., Gupta, H. V., Bouten, W., and Sorooshian, S.: A Shuffled Complex Evolution Metropolis algorithm for optimization and uncertainty assessment of hydrologic model parameters, *Water Resources Research*, 39, 10.1029/2002wr001642, 2003.
- Ye, M., Meyer, P. D., and Neuman, S. P.: On model selection criteria in multimodel analysis, *Water Resources Research*, 44, 10.1029/2008wr006803, 2008.
- Zeng, X. K., Wang, D., Wu, J. C., and Chen, X.: Reliability Analysis of the Groundwater Conceptual Model, *Human and Ecological Risk Assessment*, 19, 515-525, 10.1080/10807039.2012.713822, 2013.
- Zhang, D.: *Stochastic methods for flow in porous media : coping with uncertainties*, Academic, San Diego, Calif. London, xiv, 350 p. pp., 2002.
- Zhang, Y. K., and Schilling, K.: Temporal scaling of hydraulic head and river base flow and its implication for groundwater recharge, *Water Resour Res*, 40, -, Artn W03504 Doi 10.1029/2003wr002094, 2004.
- Zhang, Y. K., and Li, Z. W.: Temporal scaling of hydraulic head fluctuations: Nonstationary spectral analyses and numerical simulations, *Water Resources Research*, 41, 10.1029/2004wr003797, 2005.
- Zhang, Y. K., and Li, Z. W.: Effect of temporally correlated recharge on fluctuations of groundwater levels, *Water Resources Research*, 42, 10.1029/2005wr004828, 2006.
- Zhang, Y. K., and Yang, X. Y.: Effects of variations of river stage and hydraulic conductivity on temporal scaling of groundwater levels: numerical simulations, *Stoch Env Res Risk A*, 24, 1043-1052, DOI 10.1007/s00477-010-0437-5, 2010.
- Zhu, J. T., Young, M. H., and Osterberg, J.: Impacts of riparian zone plant water use on temporal scaling of groundwater systems, *Hydrol Process*, 26, 1352-1360, Doi 10.1002/Hyp.8241, 2012.

## Figure captions

**Figure 1** A schematic of the unconfined aquifer studied where  $W(t)$  is the random time-dependent source/sink,  $H_0(x)$  is the random initial condition,  $Q(t)$  is the random time-dependent flux at the left boundary,  $H(t)$  is the random time-dependent water level at the right boundary,  $L$  is distance from the left to the right boundary, and  $h(x, t)$  is the random groundwater level in the aquifer.

**Figure 2** The graphs on the left column are the standard deviation ( $\sigma'_h$ ) of groundwater level ( $h(x, t)$ ) versus the dimensionless time ( $t'$ ) at the dimensionless locations  $x'=0.0, 0.2, 0.4, 0.6$ , and  $0.8$ . The graphs on the right column are  $\sigma'_h$  versus  $x'$  for the different  $t'$ : b) and d) are for  $t'=0.0, 0.2, 0.4, 0.6$  and  $0.8$ , f) and h) are for  $t'=0.01, 0.1$ , and  $1.0$ , and j) is for  $t'=0.01, 0.2, 0.4, 0.6$  and  $0.8$ . Also, a) and b) are based on Eq.(11) where  $\sigma_W^2 = \sigma_Q^2 = \sigma_H^2 = 0$ ; c) and d) are based on Eq. (12) where  $\sigma_{W_0}^2 = \sigma_Q^2 = \sigma_H^2 = 0$ ; e) and f) are based on Eq. (13) where  $\sigma_{W_0}^2 = \sigma_W^2 = \sigma_H^2 = 0$ ; g) and h) are based on Eq. (14) where  $\sigma_{W_0}^2 = \sigma_W^2 = \sigma_Q^2 = 0$ ; i) and j) are based on Eq.(15) where  $\sigma_{W_0}^2 \neq \sigma_W^2 \neq \sigma_Q^2 \neq \sigma_H^2 \neq 0$ .

**Figure 3** The dimensionless power spectrum versus frequency ( $f$ ) at the dimensionless locations  $x'=0.0, 0.2, 0.4, 0.6, 0.8$ , and  $0.9$ . The graphs on the left column are for  $t_c = 40$  day, the graphs on the middle column are for  $t_c = 400$  day, and the graphs on the right column are for  $t_c = 4000$  day. The graphs on the first row are the dimensionless spectrum  $S_{hh}/S_{QQ}$  when  $S_{WW}=0$ ,  $S_{HH}=0$ , and  $S_{QQ} \neq 0$  in Eq. (10), the graphs on the second row is  $S_{hh}/S_{HH}$  when  $S_{WW}=0$ ,  $S_{QQ}=0$ , and  $S_{HH} \neq 0$ , the graphs on the third row are  $S_{hh}/S_{WW}$  when  $S_{QQ}=0$ ,  $S_{HH}=0$ , and  $S_{WW} \neq 0$ , and the graphs on the bottom row is  $S_{hh}/S_{WW}$  when  $S_{QQ} \neq 0$ ,  $S_{HH} \neq 0$ , and  $S_{WW} \neq 0$ .



**Figure 1**





Figure 2



**Figure 3**