

1 **Editor Decision: Reconsider after major revisions** (21 Dec 2015) by Efrat Morin

2 **Comments to the Author:**

3 The manuscript deals with an important topic and the basic of this study has a good scientific  
4 flow. The reviewers however have major comments that must be addressed and improved,  
5 such as shortening the manuscript and provide clarifications as well as others, as detailed in  
6 the reviewers reports. I ask the authors to carefully read those reports and address those  
7 comments in the revised manuscript.

8 **General answer**

9 We report below our detailed answers to all questions and comments. The too long part 7.4  
10 was reduced from 3.5 to 2 pages. The ancient Figures 14, 16b and 16c were deleted, and the  
11 discussion on the relative importance of the catchment area and the channel system in  
12 controlling catchment response as well. However, we added all the additional available  
13 information on the description of the watershed, the land use and the forms of erosion  
14 occurring in this basin (new Fig. 2 and 3). The quantification of changes in forcings on  
15 different sources of erosion is now commented in conclusion.

16 We warmly thank the editor and the reviewers who helped us to improve the paper. A  
17 sentence was added in the acknowledgements.

18  
19 **Detailed answers to Review #1**

20 This manuscript provides a detailed assessment of changes in the hydrological response of the  
21 Wadi Abd, a 2480 km<sup>2</sup> catchment in Algeria, based on a record of precipitation, flow and  
22 sediment load extending back over 40 years. Particular emphasis is placed on sediment load,  
23 which, looking more generally, has received much less attention than flow. The findings  
24 reported are important since:

- 25 (a) Their emphasis is on sediment load;
- 26 (b) There have been few studies of this nature undertaken on catchments in semi-arid  
27 areas and particularly catchments in North Africa;
- 28 (c) The catchment has evidenced a shift from perennial flow to intermittent flow, which  
29 again has received little attention to date
- 30 (d) The lengthy record (40 years) is also an important feature of the study.

31 (e) The study catchment has not experienced other major changes over the past 40 years  
32 due to human impact (e.g. major land use change) and therefore provides a good  
33 dataset for evaluating the impact of climate change.

34 The study builds on a previous paper by the authors published in 2007. This paper dealt with  
35 the 22 year record from 1973 to 1995. The current paper covers the 40 year period from 1970-  
36 2010 and therefore represents an important update on the previous publication.


37 Overall, I would suggest that the manuscript is rather ‘heavy going’ and too long and that the  
38 authors should focus on reporting the changes that have occurred in selected key indicators of  
39 changes in the behaviour of the catchment. This is what readers are likely to find most useful.  
40 I have, for example, indicated below that the discussion of changes in the parameters a and b  
41 of the rating curves should be curtailed. I would recommend suggest that the ms should be  
42 shortened by about 15-20%. This would strengthen its message. The results presented should  
43 be of general interest to readers of HESS.

44 The authors warmly thank the reviewer for his suggestion. The revised version will be  
45 shortened (especially in the discussion on (a, b) changes) and will more clearly report the  
46 changes in the behaviour of the catchment as well as the associated time scales.

47 I have little further to say in terms of the analysis and results presented. However, four e  
48 issues need further attention.


49 I. Since the ms places emphasis on changes in the sediment yield of the study catchment,  
50 section 2.1 should provide a much fuller and clearer description of the landscape of the  
51 catchment and more particularly the main erosion processes and sediment sources and  
52 their likely relative importance. Is the catchment surface or the channel system the main  
53 sediment source. Are there gullies etc etc. Some of this information may be presented in  
54 the 2007 paper but I do not think that it is acceptable that the reader has to search out  
55 another paper to find key information.

56 More detailed information that were neither presented in 2007 nor in the submitted  
57 manuscript are proposed in the revised version of the paper: maps of geology, slope and  
58 vegetation cover (new Fig. 2) and typical examples of linear erosion processes within the  
59 catchment (new Fig. 3). A text is also added: “In the plain, sheet (interrill) and rill  
60 erosion dominates (Fig. 3 b, f). Gully erosion is mainly restricted to the mountainous  
61 regions of Frenda and Tiaret in the North (Fig. 3 c, d and Fig. 2c), while some mid-slope  
62 areas are gullying (Fig. 3 a, e).”




63 (a)

64




65 (b)

66



67 (c)

68



69 (d)

70 (new) Fig. 2 The Wadi Abd catchment area. (a) Rain and hydrometric stations including HS1  
 71 at Takhmaret and HS2 at Ain Hamara, (b) Geology, (c) Slopes from the Digital Elevation  
 72 Model of North Algeria, (d) Vegetation cover from Landsat ETM+ data of 2009

73

74


75

76



77

78



82

83 (new) Fig. 3. Linear erosion forms in the Wadi Abd basin. (a) and (e) Gully erosion (depth: 30-50  
 84 cm), (c) and (d) Gully erosion (depth: 50-200 cm), (b) and (f) Sheet (interrill) and rill erosion

85 II. The authors rightly stress that their study is important because they are able to look at the  
86 impact of climate change without having to distinguish this from other human impacts on  
87 the catchment. I think that there is a need to provide an explicit discussion of the lack of  
88 change in other drivers of the hydrological response of the catchment. Has there really  
89 been no land use change? Have there been changes in livestock densities? Has the  
90 construction of small reservoirs etc caused changes in the flow regime? This discussion  
91 should be linked back to I above.

92 We are not able to prove that the changes are only due to climate change. If it was the  
93 case, the paper would be entitled: "Effects of climate change on sediment load in the  
94 Wadi Abd".

95 Livestock density is very low (see the new Fig. 2d). This information was added in the  
96 following sentence of the conclusion: "The lack of data on land use and land cover  
97 changes over 40 years does not allow us to isolate the factors directly related to climate  
98 change from those related to other anthropogenic activities, but this question was not in  
99 the scope of the paper. The small population, the low coverage of pasture (see Fig. 2d), of  
100 cultivated areas and vegetation (43 %) in the basin and the small volume of reservoirs  
101 (nominally 2.3% of the annual discharge, but silted up to 70%) make us think that in this  
102 system the effects of climate change dominate anthropogenic effects."

103 III. I am concerned that the authors prefer to lump all the sediment data together to produce a  
104 rating curve for the entire period which is then used to calculate annual sediment loads.  
105 This seems unacceptable when they claim that they are looking at a non-stationary system  
106 demonstrating the effects of climate change. This issue needs to be explicitly discussed.

107 The comparison detailed p.10464 (lines 10-25) between two strategies of rating curve  
108 application from one side, and the long-term changes in (a, b) regime which becomes  
109 visible with values averaged over 15 years from the other side, rise maybe a main lesson  
110 of this study: the non-stationarity of the system occurs at two time scales. The revised  
111 version states more clearly the different time scales involved in the change of the basin's  
112 behaviour. We add the following paragraph in the conclusion: "The rapid change in  
113 sediment regime which is instantaneously driven by the changing flow regime should be  
114 distinguished from the slow change in the concentration-flow relationship. The change in  
115 flow regime can be precisely dated in May-July 1986 (with 49 consecutive dry days),  
116 while the change in the C-Q relationship needs averaging over 15 years of a, b and

117 specific sediment yield to become evident. Such inertial effect may be attributed to the  
118 time for the basin soil properties (such as humidity) or vegetation to adapt to the new  
119 climate conditions. It likely depends amongst other factors on the underground water  
120 storage, and thus on basin lithology and infiltration history. On the Wadi Abd basin, the  
121 time needed for the flow regime to change after the dryness settlement in early 1970's  
122 (see Fig. 4) is around 15 years.”

123 Anyway, the new figure 5 (previous Fig. 3) shows that the use of one single relationship  
124 over 1970-2010 is suitable to reproduce the variations of daily C from those of daily Q.

125 IV. I found the discussion and interpretation of the parameters a and b in the sediment rating  
126 curve unconvincing and contrived. I would strongly suggest that this part of the  
127 manuscript should be deleted or at least greatly reduced. I do not believe that these two  
128 parameters provide information on the relative importance of the catchment surface and  
129 the channel system in controlling catchment response. The authors suggest that the  
130 changes in a and b reflect increased erosion of the watershed and decrease in the erosive  
131 power of the river. I would see the changes in a and b and simply reflecting the shift of  
132 the river regime from perennial flow to intermittent flow. As a result of this change flows  
133 will decline to zero and sediment concentrations associated with lower flows during  
134 storm events will increase. Parameter a represents the concentration associated with a  
135 flow of 1. Since flows decline to zero after storms, high concentrations can be recorded at  
136 low flows on the rising limb of the hydrograph when flow resumes during storm events.  
137 Therefore parameter a will increase. There is no need to invoke a increase in erosion  
138 within the watershed. If parameter a increases the slope of the line (i.e. b) will inevitably  
139 decrease. Again there is no need to invoke a change in the erosive power of the river. The  
140 authors refer to the work of (Wang et al. 2008) on the Yangtze below the Three Gorges  
141 Dam as providing further evidence of the potential to use the two parameters to identify  
142 changes in the relative contribution of the basin surface and the river. Dam construction is  
143 a special case, which is very different from the response of a more natural catchment. If a  
144 dam is constructed upstream, the suspended sediment load and concentration are likely to  
145 decline due to sediment trapping by the dam. As a result sediment concentrations for a  
146 given flow will decline and a would decrease. If a decreases, b is likely to increase and  
147 this will also reflect scour of the channel below the dam during high flows due to the low  
148 sediment load (i.e. Kondolf's hungry water) which will reduce any decline of sediment  
149 concentrations at high flows.

150 Part 7.4 was 3.5 pages long, it was curtailed and reduced to 2 pages. Figures 14, 16b and  
151 16c were removed. The discussion on the relative importance of the catchment surface  
152 and the channel system in controlling catchment response was deleted, as well as the  
153 reference to Wang et al (2008). The conclusion was also reduced on the interpretation of  
154 a and b values, and the reference to Trimble (1999) was removed.

155 My remaining comments relate, firstly, to the need for revision to deal with the use of terms  
156 which are likely to be unclear to native English speakers and which probably reflect terms  
157 used in France and, secondly, to other improvements in the text and minor queries. The terms  
158 that need attention are:

159 (a) RUPTURE This term is used very frequently in the ms. to refer to break points or  
160 change points in the double mass curve and similar contexts. Its use with this meaning  
161 will not be familiar to readers of the ms. 'Shift' is a word that might be appropriate in  
162 some places.

163 [Thanks. It was corrected.](#)

164 (b) The term SPECIFIC DEGRADATION, although commonly used in French  
165 (degradation spécifique) will not be familiar to most readers. If the term 'sediment  
166 yield' is used for the total load i.e.  $t \text{ year}^{-1}$ , then the term specific sediment yield should  
167 be used to refer to  $t \text{ km}^{-2} \text{ year}^{-1}$ .

168 [Done](#)

169 (c) SEDIMENT WASH-DOWN (page 10460 line 3) It is not clear what you mean by the  
170 ratio of sediment wash-down to river discharge. What is sediment wash-down. What is  
171 the ratio? Is it effectively load/discharge i.e. concentration?

172 [Yes, it is the ratio load/discharge i.e. concentration \( \$10.7 \text{ g L}^{-1}\$  over 22 years in  
173 AO2007\). The correction is done.](#)

174 (d) AGGLOMERATIONS Page 10461 line 25. What are agglomerations?

175 ["Cities" \(sorry, we let the French word in the submitted version\). Corrected.](#)

176 (e) GAPS On line 11 page 10471 you use this word when you really mean 'differences'

177 [Corrected](#)

178 (f) UNDERGROUND WATER LAYERS You refer to this on line 6 page 10472. I think  
179 you probably mean 'underground storage'.

180 Corrected

181 (g) PARTICLES not PARTICULES (page 10481 line 15).

182 Thanks. Corrected

183 There are many other instances of poor English which need to be dealt with. The ms needs  
184 careful editing by a native English speaker with expertise in hydrology. It may be necessary to  
185 make use of commercial scientific editing service. Some suggested corrections and comments  
186 are listed below.

187 1) Page 10458 line 14. 'scatter' of what?? "of the C-Q pairs" (added)

188 2) Page 10459 'patterns involved in' change this to 'factors controlling' Thanks. Done.

189 3) Page 10460 line 26 Change to 'relationship between sediment load and runoff over...'  
190 Thanks. Done.

191 4) Page 10463 lines 1-3. It is not clear how the mean daily values are obtained from the  
192 instantaneous values. There is a need to specify key procedures used and not to expect the  
193 reader to search out other papers to find this information.

194 The text was changed into: "From these 9076 coincident instantaneous data measured  
195 during 1213 days, average arithmetic values were calculated per day so as to obtain 1213  
196 pairs of "mean daily" (C, Q) values. The resulting "mean daily Q" differs from the (true)  
197 daily discharge obtained from the averaging of 24h of continuous instant Q."

198 5) Page 10464. Lines 10-25. The argument here seems counterintuitive and questionable. I do  
199 not think it is acceptable to use a rating relationship developed for the 40 years of data  
200 when the system is clearly not stationary. You are identifying important changes but then  
201 apparently ignoring them by using a lumped rating relationship.

202 See issue III above.


203 6) Page 19465 line 3. 'to understand better..' Avoid the spilt infinitive. Thanks. Done.

204 7) Page 10468 line 3. Avoid using the words 'sediment delivery' since they are often used to  
205 refer to the processes operating between sediment mobilisation in the catchment and the  
206 sediment load at the catchment outlet i.e. conveyance losses and storage. You are referring  
207 to sediment output, sediment load or sediment yield. Thanks. We changed to "sediment  
208 load" or "sediment yield" all along the paper.



- 209 8) Page 10469 lines 1-3. It is not clear why you need daily values of concentration and  
210 discharge to calculate average values of SPM\*. If is this is load -weighted mean (i.e.  
211 Load/Discharge) you need to make this clear. You also need to make it clear whether the  
212 daily concentration data are measured data or extrapolated values obtained using the rating  
213 curve relationship. You're right. SPM\* is the load weighted mean and can be obtained  
214 from daily values but also from weekly, monthly, seasonally or directly yearly values of Q  
215 and Qs, since  $SPM^* = Q_s/Q$ . The lines are changed into: "The average value of SPM\*  
216 calculated over the period 1970-2010 is 12.3 g.L<sup>-1</sup>. The 40 annual values of SPM\*  
217 calculated for each year from measured discharges and concentrations estimated using the  
218 rating curve (3) vary between 2.5 g.L<sup>-1</sup> and 50.2 g.L<sup>-1</sup>"
- 219 9) Page 10469 lines 24 and 25. The wording needs improving. I think that you mean  
220 '...showed that, over 22 years, 71% of the variance of the annual SPM values .....was  
221 accounted for by the annual discharge.. and 73% by the 95th percentile ... ' Thank you.  
222 Corrected as suggested.
- 223 10) Page 10472 line 5 '...runs into...' Line 14 reword '...and the water level in the aquifers  
224 will be lowered.' Line 26 '...suspended sediment loads were greater in the autumn during  
225 the 2000s..'Lines 27-28 reword to ' shifted from.....to a regime with one dominant  
226 season in the 2000s'. Thanks. Done.
- 227 11) Page 10475 Line 9 'the Maghreb' Line 10 '...correlated with..' Line 14. Do you mean  
228 '...on the highest discharges than on the average discharge'. Line 27 '...of 0.226 when  
229 correlated with ...' Thanks. Done.
- 230 12) Page 10476 line 2 'is responsible for..' Lines 6-7 'consequences for'. Done. Line 24. It is  
231 not clear what is meant by 'averaged per decade'. How can a single value of average flow  
232 provide information on the date of the first flood? We added and corrected the first  
233 sentence following: "The analysis of the time series of daily flows enables to determine the  
234 start of the first summer flood. The average daily flow per decade suddenly increases the  
235 day at which the first summer flood occurred, at least once in the decade. By observing  
236 these decadal averaged daily flows, there is no ambiguity on the start of the earlier flood by  
237 decade."
- 238 13) Page 10477 Line 1 'starts on average on the 6th September...' Line 4 ' the first flood of  
239 summer'. Line 8 'was observable'. Line 10 'gave the following results for...'. Done. Line  
240 24. Give reference for NQAA. 2 references were added (Hansen et al., 2010; GISTEMP

241 Team, 2015) as requested by NASA. By error, the previous Figure 13 (which now  
242 becomes Fig. 15) showed only 22 points corresponding to 1973-1995. A new Fig. is  
243 provided in the revised version and the  $r^2$  value (0.388) is changed accordingly in the text:



244  
245 [Ref. added:](#)

246 [GISTEMP Team: GISS Surface Temperature Analysis \(GISTEMP\). NASA Goddard Institute](#)  
247 [for Space Studies. Dataset accessed 2015-12-08 at http://data.giss.nasa.gov/gistemp/](#)  
248 [Hansen, J., Ruedy, R., Sato, M., and Lo, K.: Global surface temperature change, \*Rev.\*](#)  
249 [Geophysics, 48, RG4004, doi:10.1029/2010RG000345, 2010](#)

250  
251 14) Page 10478 Line 3. Please check the Langbein and Schumm paper. I think the  
252 relationship was with annual effective rainfall rather than annual rainfall. The two are  
253 different. You're right. We changed the text into: "Many authors studied the variations of  
254 sediment load per unit of catchment area against annual rainfall (e.g. Summerfield and  
255 Hulton, 1994) or effective rainfall (e.g. Langbein and Schumm, 1958)" and added the  
256 following reference:

257 [Summerfield, M.A., Hulton, N.J.: Natural controls of fluvial denudation rates in major](#)  
258 [world drainage basins, \*J. Geophys. Res.\*, 99 B7, 13871-13883, 1994.](#)

259 Line 18 'fluvial sediment rather than 'riverine sediment'. Done.

260 Line 23. It is not clear what is meant by ‘parameter evolutions’ The text was changed into:  
261 “Although the river regime shift clearly impacted several parameters between the two  
262 periods”

263 15) Page 10481 line 3 ‘after’ rather than ‘posterior to’ Done.

264 16) Page 10482 Line 6. ‘favour’ and ‘as a sediment source..’ Line 16 ‘correlation with...’  
265 Done.

266 17) Page 10483 line 2 ‘estimation’? line 5 ‘coefficient of determination’. Lines 6 and 7.  
267 ‘..established for only the last decade did not provide a reliable estimate of the soild  
268 discharge....’ Line 17 reword ‘...rating curves, water discharge must be recorded at  
269 frequent intervals, although measurements of ...’ Done.

270 Line 23 meaning of ‘advance’ not clear. The text was changed into: “rainfall moved  
271 forward during the late warm season and the watershed of Wadi Abd experienced a  
272 significant change”

273 Line 24 ‘..increased variability at both the inter-annual and intra-annual levels. Done.

274 18) Page 10484 lines 9-12. As noted above you need to provide some indication of the likely  
275 relative importance of the catchment surface and the channel system as sediment sources.  
276 The discussion on the sediment sources (catchment surface or channel system) was deleted  
277 in the new version of the paper. Line 18 . What are INSTANT models?? “instant” was  
278 deleted. Line 23 Should it be ‘marl’? Yes. Correction done.

279 19) Page 10485 Line 1 wording needs correction. ‘..makes us think that...’ Line 20 ‘due to..’  
280 Done.

281 20) Page 10486 line 2 ‘..in the Maghreb..’ Done.

282 21) Page 10496 Table 2 avoid the term specific degradation - refer to specific sediment yield.  
283 Degradation could involve chemical weathering. Thanks. Correction done.

284 22) Page 10497 Table 3 title. Delete ‘sediment delivery’ and insert ‘sediment output’ or  
285 ‘sediment yield’. Done.

286 23) Page 10499 Figure 2 title. ‘...mean annual temperatures...’ Done.

287 24) Page 10502 Figure 5 title. See 22 above. Done.

288 25) Figure 6 title. Refer to ‘flowing water’ not ‘running water’ . Running water comes out of  
289 a tap! Thanks. Done.

- 290 26) Page 10504 Figure 7 y axis. The units cannot be t. It must be a multiple of t. The unit  
291 should be tonnes not tons. [Sorry, the unit is 10<sup>6</sup> metric tons.](#)
- 292 27) Page 10505 Figure 8 title. Find an alternative term to ‘sediment delivery parameters’. As  
293 indicated above, do not use ‘specific degradation’ in the title and on the y axis Bb.  
294 “[sediment delivery](#)” was corrected into “[sediment load](#)”, and “[specific degradation](#)” by  
295 “[specific sediment yield](#)”
- 296 28) Page 10506 figure 9. Change labelling of y axis in Fig. 9C to ‘Seasonal contribution to  
297 mean annual sediment load (%) (9 year average). [Done.](#)
- 298 29) Figure 13 title. Provide a reference to GSS [Done \(see also comment 14 above\)](#)
- 299 30) Page 10513 The key should refer to  $r^2$  and not  $R^2$  . You are plotting simple bivariate  
300 relationships.  $R^2$  is used for multiple correlation. [Done on every figure.](#)

301  
302

303

### [Answers to Review #2](#)

304 The authors didn’t present a thorough critical study of the solid matters measurements. In  
305 order to carry out a work on the impact of climate changes on this phenomenon, such a study  
306 is a much needed. What does the portion of suspended matter represent in the total sediment  
307 transport? What does the soils erosion represent in these values measured at the basin mouth?  
308 Remini Boualem showed, in Algeria, that banks ablation is a significant proportion of in the  
309 sediment transport in Algeria, what about the banks ablation in your work? In order to be able  
310 to study the relationship between Climate Change and the sediment transport. This  
311 phenomenon has connection with the extreme events which also are synonyms of climate  
312 change. Also, the vegetational cover evolution that has a predominant role in the soils erosion  
313 is not mentioned. The climate changes directly affect this evolution and the spatial and  
314 temporal evolution of sediment transport as a result. In my view, the authors should first deal  
315 with those aspects prior to undertaking the statistic study of the measurements series which  
316 were not criticized rigorously. The coefficients obtained along the paper have no physical  
317 meaning without extending the intervening phenomena that interact in the phenomenon  
318 accountable for the production of those solid matters at the mouth. All the parties: erosion,  
319 ablation... must be evaluated in order to take account only of the party which can be in  
320 connection with the change in pluviometric and hydrometric mechanisms.


321 [Answers:](#)

322 [1. The solid matter measurements are questioned by the reviewer. As explained in the paper,](#)  
323 [the river discharge and suspended sediment concentration measurements were performed and](#)  
324 [provided by the ANRH, the National Agency of Algeria in charge of Hydraulic Resources.](#)  
325 [The protocol which is described in § 2.2 is the same in all rivers from Algeria and thus for all](#)  
326 [papers on water resources in Algeria \(including the works by Remini Boualem or the](#)  
327 [references he referred to in his papers – one reference to Remini’s paper was added in the](#)  
328 [revised version\). The protocol can be questioned, as we did in the last part of our paper: in the](#)  
329 [case of an intermittent river, more frequent measurements \(and continuous automatic](#)  
330 [measurements if possible\) are suggested in the future.](#)

331 2. “What does the soils erosion (and the bank erosion) represent in these values measured at  
332 the basin mouth?”: The origin of the particles transported at the hydrologic station is unknown  
333 and can’t be determined precisely from the available data set. Additional measurements  
334 would be requested. This question is beyond the scope of the paper.

335 3. “This phenomenon has connection with the extreme events which also are synonyms of  
336 climate change”: For sure, sediment flux is mainly driven by the extreme precipitation  
337 episodes. However, we don’t agree that extreme events are “synonyms” of climate change.  
338 Extreme events have occurred before climate has started to change. In this paper, the entire  
339 data set of rainfall – including the extreme events – is considered so as to provide a general  
340 view of the hydrology and sediment transport dynamics in the basin. A future study could be  
341 restricted on extreme events, but a general glance is a first mandatory step before extracting  
342 parts of the full time series of measurements.

343 4. “the vegetational cover evolution that has a predominant role in the soils erosion is not  
344 mentioned”: The available information on the vegetation cover are given in § 2.1. An  
345 additional figure of the vegetation cover in 2009 is added in the revised version (see figure  
346 below). The question of its evolution between 1970 and 2010 goes far from the scope of the  
347 paper. This paper focuses on temporal changes of sediment dynamics in a river turning from  
348 perennial to ephemeral, which is very rarely reported in the literature (see the general  
349 comment of the reviewer #1). A future and complementary study of the vegetation cover  
350 history and its impact on the hydrologic behavior of the basin can be envisaged using  
351 historical satellite data and a numerical model (such as WEPP, EUROSEM or SWAT). A  
352 paragraph was added to address this question in the conclusion (see below “General  
353 comments”).



354

355

(New) Fig. 2d added: Vegetation cover of the Abd basin in 2009

356 5. “The coefficients obtained along the paper have no physical meaning without extending the  
357 intervening phenomena that interact in the phenomenon accountable for the production of  
358 those solid matters at the mouth”: The scientific literature on the rating curve coefficients  
359 values and their variability is rich, because the question of their meaning is still pending. We  
360 agree that these values should be related to the processes of sediment production (and  
361 transport, see next comment). The dataset faces us with a non linear answer of a system on  
362 changing forcing. This study aims at describing the dynamics of this very complex system.  
363 The coefficients are discussed on the basis of the available information but their physical  
364 meaning is not fully solved by this paper. By the way, is this question solved in any paper?  
365 We agree with the reviewer that this study opens avenues on potential future studies on  
366 different aspects of sediment dynamics in the basin.

367 6. “All the parties: erosion, ablation... must be evaluated in order to take account only of the  
368 party which can be in connection with the change in pluviometric and hydrometric  
369 mechanisms”: Because the suffix “metric” refers to measurements and not to processes or  
370 mechanisms, this request is not clear to us. Shall we understand “... in connection with the  
371 changes in rainfall and river discharge”? In that case, if we consider the changes from a  
372 decade to another at the scale of one century (see for example the alternation of 20 wet years  
373 in the 50s and 60s, followed by a severe drought in the 70s and 80s, fig. 4, § 4.2 and 7.1.1),  
374 which period could be considered as a reference? This paper emphasizes the fact that a period  
375 of 40 years of measurements is not long enough to define a reference behavior. That’s why  
376 we don’t discuss the mean value of specific suspended sediment yield, for example, in this  
377 paper.

378

379

### **General comment and answer to reviewer 2**

380 This paper is based on the complementary study of variability of water and suspended  
381 sediment discharge at one gauging station, considering some related climatologic changes (in  
382 temperature, and rainfall intensities and periods of occurrence). We thank the anonymous  
383 reviewer #2 who is much concerned by the soil cover, erosion process and by the origin of the  
384 suspended particles, i.e. by a pedological study of the catchment. A paragraph is added in the  
385 conclusion of the revised version: “The quantification of forcing changes on sediment sources  
386 (raindrop erosion, sheet erosion, rill erosion, gully erosion, stream channel erosion) may be  
387 investigated in situ (e.g. Poesen et al., 2003) and/or estimated using a numerical model of the  
388 hydrologic and sedimentological functioning of the basin, such as WEPP (Nearing et al.,  
389 1989), EUROSEM (Morgan et al., 1998) or SWAT (Neitsch et al., 2011). Such a model could  
390 help us to test hypothesis and quantify or at least estimate the effects of different forcing  
391 changes (temperature, runoff, vegetation etc) in future studies.”

392 Ref. added in the revised version:

393 Morgan R.P.C., Quinton, J.N., Smith R.E., Govers, G., Poesen, J.W.A., Auerswald, K., Chisci  
394 G., Torri, D., and Styczen, M.E.: The European Soil Erosion Model (EUROSEM): A  
395 dynamic approach for predicting sediment transport from fields and small catchments,  
396 Earth Surf. Process. Landforms, 23, 527-544, 1998.

397 Neitsch, S.L., Arnold, J.G., Kiniry J.R., and Williams, J.R.: Soil and Water Assessment Tool  
398 – Theoretical Documentation version 2009. Texas Water Inst. Techn. Report n°406,  
399 Texas A&M University, College Station, 2011.

400 Poesen J., Nachtergaele, J., Verstraeten, G., and Valentin, C.: Gully erosion and  
401 environmental change: importance and research needs, Catena, 50, 91-133, 2003.

402 Remini B., Leduc C., and Hallouche, W.: Evolution des grands barrages en régions arides:  
403 quelques exemples algériens, Sécheresse, 20 (1), 96-103, 2009.

404

405

Marked-up manuscript version

406 **Recent changes in climate, hydrology and sediment load in**  
407 **the Wadi Abd, Algeria (1970-2010)**

408

409 **Mohammed Achite<sup>1</sup>, Sylvain Ouillon<sup>2,3</sup>**

410 [1]{Laboratoire Eau-Environnement, Université Hassiba Ben Bouali, BP 151, Hay Es-Salem,  
411 Chlef, Algeria}

412 [2]{Université de Toulouse, UPS (OMP), IRD, UMR 5566 LEGOS, 14 avenue E. Belin,  
413 31400 Toulouse, France}

414 [3]{University of Science and Technology of Hanoi, Dept. Water-Environment-  
415 Oceanography, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam}

416 Correspondence to: Sylvain Ouillon, ([sylvain.ouillon@legos.obs-mip.fr](mailto:sylvain.ouillon@legos.obs-mip.fr))

417

418 **Abstract**

419 Here we investigate the changes of temperature, precipitation, river runoff and sediment  
420 transport in the Wadi Abd in NW Algeria over a time series of 40 hydrological years (1970-  
421 2010). Temperature increased and precipitation decreased with the reduction in rainfall being  
422 relatively higher during the rainy season. A shift towards an earlier onset of first rains during  
423 summer was also found with cascading effects on hydrology (hydrological regimes,  
424 vegetation etc) and thus on erosion and sediment yield. During the 1980s, the flow regime  
425 shifted from perennial to intermittent with an amplification of the variations of discharge and  
426 a modification of the sediment regime with higher and more irregular suspended particulate  
427 flux. Sediment flux was shown to almost double every decade from 1970s to 2000s. The  
428 sediment regime shifted from two equivalent seasons of sediment **yield** (spring and autumn)  
429 to a single major season regime. In 2000s, autumn produced over 4 times more sediment than  
430 spring. The enhanced scatter **of the C-Q pairs** denotes an increase of hysteresis phenomena in  
431 the Wadi Abd that is probably related to the change in the hydrologic regime. **The increased**  
432 **erosion of the watershed is accompanied by a decrease in the coefficient b of its rating curves**  
433 **and a decrease in the erosive power of the river.** At the end of the period, due to the

434 irregularity of the discharge, the ability of a rating curve to derive suspended sediment  
435 concentration from river discharge was poor.

436 **Keywords:** water erosion; suspended sediment concentration; sediment transport; rating  
437 curve; hydroclimatology; wadi; intermittent river; Algeria

438

## 439 **1 Introduction**

440 Fluvial and estuarine suspended sediment fluxes have been dramatically changing under the  
441 combined effects of anthropogenic activities and climate change. On a global scale, recent  
442 changes showed a trend towards increasing land erosion and decreasing fluxes to coastal  
443 waters (Walling and Fang, 2003; Vörösmarty et al, 2003; Wang et al., 2006). The sediment  
444 flux trapped in regulated basins with reservoirs is higher than 50% (Vörösmarty et al., 2003).  
445 Locally, it can reach more than 60% after the impoundment of one single dam like on the Red  
446 River (Vinh et al., 2014), and more than 80% on rivers with many dams (86% on the Yellow  
447 River, Wang et al., 2007; >95% on the Ebro river, Durand et al., 2002). Other engineering  
448 activities (meander cutoffs, river-training structures, bank revetments, soil erosion controls)  
449 also affect significantly the sediment fluxes and can participate to shift from a transport-  
450 limited system to a supply-limited system, like on the Missouri-Mississippi (Meade and  
451 Moody, 2010).

452 With increasing temperature and evaporation, climate change tends to accelerate the water  
453 cycle and modify hydrologic regimes (Bates et al., 2008). Precipitation intensities and the  
454 frequency of extreme events are projected to increase under climate change, leading to more  
455 frequent flood events of higher magnitude that will, in turn, affect patterns of erosion and  
456 deposition within river basins (Tucker and Slingerland, 1997; Pruski and Nearing, 2002;  
457 Tockner and Stanford, 2002; Coulthard et al., 2012). Recent studies focused on the impact of  
458 climate change on sediment transport (e.g. Gomez et al., 2009; Hancock, 2009; Walling,  
459 2009; Hancock and Coulthard, 2011; Knight and Harrison, 2013; Lu et al., 2013). Syvitski  
460 (2003) showed on an example that sediment transport may increase due to the increasing  
461 discharge or decrease because of the enhanced temperature. Studies compared the trends in  
462 hydrological and sediment time-series to the land use changes (Wang et al., 2007; Memariam  
463 et al., 2012; Gao et al., 2012). Climate projections are consistent on warming and acceleration  
464 of the water cycle (IPCC, 2013) but they remain to be defined on sediment transport where  
465 projections shows a high uncertainty (Shrestha et al., 2013; Lu et al., 2013). This is in part due



466 to the fact that climate affects many **factors controlling** sediment yield, such as surface  
467 moisture availability, weathering processes and rates, and the nature of the riparian vegetation  
468 (Nanson et al., 2002).

469 While sediment transport is well-documented in perennial rivers in humid or temperate  
470 climates, its study in semiarid areas is still fragmentary due to the difficulty of sampling  
471 during flashfloods. Amongst the factors favoring erosion (slope, nature of rocks, relief,  
472 climate, human activities), climate is recognized to be the main factor in semi-arid  
473 mediterranean areas of Algeria which experience short and intense rain episodes, high  
474 evaporating power of wind, prolonged droughts and freezing and thawing cycles (Touaibia,  
475 2010; Houyou et al., 2014). Erosion is extremely active and the **average concentration** is at  
476 least one order of magnitude higher than at global scale (Achite and Ouillon, 2007). One of  
477 the main impacts of this high erosion is the rapid silting up of reservoirs (up to 2 to 5% per  
478 year, Kassoul et al., 1997; **Remini et al., 2009**; Touaibia, 2010) with important consequences  
479 on water resources management in a region where 85% of rain evaporates (Benhamiche et al.,  
480 2014). The high temporal variability and recent changes in forcings mean that it is necessary  
481 to study sediment dynamics in such environments over time-periods of several decades in  
482 order to document and understand the changes in sediment regime.

483 In this context, this paper extends to cover a 40-year period (1970-2010) the analysis of  
484 sediment transport changes of a wadi already studied over a 22-year period (1973-1995 in  
485 Achite and Ouillon, 2007 hereafter referred as AO2007). The hydrologic gauging station is  
486 located upstream from a dam and is not affected by any major management. This river sub-  
487 basin is also particularly suitable for such study because its hydrologic regime was shown to  
488 have drastically changed between the 1970s and the 1980s. Precipitation decreased and  
489 became more irregular, the flow regime shifted from perennial to intermittent with 26% of dry  
490 days in average in 1990-1995, amplified variations of discharge, and a modified sediment  
491 regime with higher and more irregular suspended particulate flux, 4.7 times higher over 1985-  
492 1995 than over 1973-1985. AO2007, showing the advantage of working over 22 years of  
493 measurement, however, stressed the difficulty of defining a reference period, and the need to  
494 extend the study period longer. The objectives of this additional study are to 1) describe the  
495 precipitation, discharge and sediment flux variability of the Wadi Abd basin over a 40-years  
496 period, 2) detect the **shift** if any in temperature, runoff and sediment yield, 3) determine the  
497 relationship **between sediment load and runoff over** the last 40 years, 4) detect when a shift  
498 occurred in the runoff-sediment load relationship, 5) analyze the possible causes of the change

499 in flow regime and its consequences on suspended sediment discharge, 6) assess the use of  
500 rating curves and the physical signification of its parameters when a river is experiencing a  
501 transition and turns from a perennial regime to an intermittent regime.

## 502 **2 Study area: the Wadi Abd**

### 503 **2.1 General information**

504 The Wadi Abd, located in the North-Western of Algeria, is a tributary of the Wadi Cheliff, the  
505 major river of Algeria (Fig. 1). The length of the Wadi Abd's main stream is 118 km, its basin  
506 area is 2480 km<sup>2</sup> and the drainage density is 3.70 km km<sup>-2</sup> (Fig. 2a). The Wadi Abd supplies  
507 downstream the Sidi Mohamed Benaouda (SMB) reservoir which basin area is 4900 km<sup>2</sup>. The  
508 Wadi Abd catchment area is formed of erodible sedimentary rocks from Upper Jurassic  
509 (45.9% of its surface), Middle Jurassic (20.2%) and Pliocene (7.4%) (Fig. 2b). Soft bottom  
510 sedimentary deposits from the Quaternary cover 13% of the basin along the wadi (Tescult  
511 International, 2004).

512 The climate is Mediterranean and characterized by a dry season from April to  
513 August/September, and a wet season from September to March. The hydraulic deficit is very  
514 high. Annual precipitation is 264 mm on average while the mean evapotranspiration over the  
515 SMB basin is 1525 mm (Tescult International, 2004).

516 ~~The main physical, geological, topographical and vegetation characteristics of the river and~~  
517 ~~watershed, and a location map as well, are provided in AO2007. Seven hill reservoirs were~~  
518 ~~built in the Wadi Abd basin from 1986 to 2004 for agriculture (irrigation, livestock watering)~~  
519 ~~or for fire fighting measures. Their total cumulated capacity is 0.88 hm<sup>3</sup>, representing 2.3 %~~  
520 ~~of the yearly averaged discharge at Ain Hamara station. These small reservoirs are now silted~~  
521 ~~up to 70% of their volume.~~

522 The watershed mainly consists of steep slopes (Fig. 2c) with very sparse vegetation or bare  
523 soil (Fig. 2d). The main land use is natural environment (73%; 17% of forests + 56 % of scrub  
524 and bare steppe soils), cultivated lands cover about 26% and cities 0.4%. ~~Seven hill reservoirs~~  
525 ~~were built in the Wadi Abd basin from 1986 to 2004 for agriculture (irrigation, livestock~~  
526 ~~watering) or for fire fighting measures. Their total cumulated capacity is 0.88 hm<sup>3</sup>,~~  
527 ~~representing 2.3 % of the yearly averaged discharge at Ain Hamara station. These small~~  
528 ~~reservoirs are now silted up to 70% of their volume.~~

529 123.000 inhabitants were living in the Wadi Abd basin in 2008 (average density: 49  
530 inhabitants/km<sup>2</sup>), 44% of them living in the city of Takhmaret. The Wadi Abd is thus little  
531 influenced by human activities, in view of its extensive surface that is subject to severe  
532 natural erosion.

533 In the plain, sheet (interrill) and rill erosion dominates (Fig. 3 b, f). Gully erosion is mainly  
534 restricted to the mountainous regions of Frenda and Tiaret in the North (Fig. 3 c, d and Fig.  
535 2c), while some mid-slope areas are gullying (Fig. 3 a, e).

## 536 2.2 Data

537 Long-term series of temperature measured at 3 stations in Algeria were extracted from  
538 CRUTEM4 (Jones et al., 2012; Osborn and Jones, 2014). These stations are located at Chlef  
539 (36.20°N, 1.30°E - 1951-2011), Miliana (36.30°N, 2.20°E - 1922-2011) and Dar El Beida  
540 (36.70°N, 3.30°E - 1856-2011). The annual average temperatures were calculated for each  
541 station from the 12 monthly averages. 20 missing monthly data over 480 did not allow us to  
542 exactly calculate mean measured yearly temperature at Chlef, the nearest station from the  
543 Wadi Abd. In order to estimate the change per decade at Chlef either at the yearly or seasonal  
544 scale, the 20 lacking values were extrapolated from the monthly temperatures measured at  
545 Miliana and Dar El Beida using the relationships between the monthly average temperatures  
546 at Chlef and Miliana, and Chlef and Dar El Beida. Such relationships established at the annual  
547 scale are shown on Fig. 42. The resulting estimates of temperature at Chlef at seasonal and  
548 yearly scales allowed us to estimate changes by decade over the period 1970-2010.

549 Rainfall and hydrometric records were provided by the National Agency of Hydraulic  
550 Resources (ANRH). Time series of rainfall data are available at 6 stations within the basin  
551 (see Fig. 2a): S1 Ain Kermes (altitude: 1162 m), S2 Rosfa (960 m), S3 Sidi Youcef (1100 m),  
552 S4 Tiricine (1070 m), S5 Takhmaret (655 m) and S6 Ain Hamara (288 m). 9076 coincident  
553 instantaneous measurements of water discharge (namely Q, in m<sup>3</sup> s<sup>-1</sup>) and suspended sediment  
554 concentrations (C, in g L<sup>-1</sup>) were recorded at the Ain Hamara gauging station between  
555 September 1970 and August 2010. Water depths were measured continuously and a  
556 calibration between water level and discharge was regularly performed from velocity profiles.  
557 Concentrations derived from water samples taken at one or two points, after filtration on pre-  
558 weighed Whatman Glass Fibre Filters (GFF) filters, oven-dried and weighed again following  
559 the protocol described by A02007 and Megnounif et al. (2013). From these 9076 coincident  
560 instantaneous data measured during 1213 days, average arithmetic values were calculated per

561 day so as to obtain 1213 pairs of “mean daily” (C, Q) values. The resulting “mean daily Q”  
562 differs from the (true) daily discharge obtained from the averaging of 24h of continuous  
563 instant Q.

564 The Atlantic Multidecadal Oscillation (AMO) index is an index of North Atlantic  
565 temperatures. The monthly unsmoothed values used in this study were calculated by NOAA,  
566 Earth System Research Laboratory, Physical Sciences Division/ESRL/PSD1  
567 (<http://www.esrl.noaa.gov/psd/data/timeseries/AMO/>).

### 568 **3 Models and Methods**

#### 569 **3.1 Trends**

570 The analysis of trends was conducted following a method fully described by Stahl et al.  
571 (2010) and Déry et al. (2005) for river runoff. The Kendall-Theil Robust Line furnishes a  
572 linear equation from a time-series of  $n$  measurements such as

$$573 \quad y = m t + b \quad (1)$$

574 where  $t$  is time (year),  $y$  denotes the hydrological parameter (precipitation, river discharge,  
575 sediment discharge), and  $m$  is the magnitude of the trend over this period.  $m$  is calculated as  
576 the median of all slopes  $m_k$  of consecutive pairs of values:

$$577 \quad m_k = \frac{y_j - y_i}{t_j - t_i} \quad (2)$$

578 where  $k = [1, n(n-1)/2]$ ,  $i = [1, n-1]$ ,  $j = [2, n]$ . This slope is often referred to as the Sen slope (Sen,  
579 1968). The significance of this trend at a level  $p$  was calculated following Ziegler et al.  
580 (2003).

#### 581 **3.2 Rating curves**

582 C and Q measurements were used to define rating curves that estimate C from measured  
583 values of Q, according to a common approach (e.g. Walling, 1977; Asselman, 2000; El Mahi  
584 et al., 2012; Tebbi et al., 2012; Louamri et al., 2013). The most suitable model is a power law  
585 of the type  $C = aQ^b$  for which the coefficients (a, b) determined empirically account for the  
586 effectiveness of erosion and transport. In this paper, the rating curve established from the  
587 1213 daily averages of C and Q data available for the period 1970-2010 enabled the  
588 estimation of C then  $Q_s$  ( $Q_s = CxQ$ ) for the whole period from the measured daily Q values.

589 Considering the change in hydrologic regime during the study period, we wondered if the

590 estimate of C and  $Q_s$  per sub-periods like decades could be better adapted than globally. We  
591 therefore applied the 4 rating curves established for the 4 decades to the time series of daily Q  
592 to obtain daily C and then daily  $Q_s$ . This method (B) enabled us to compare the estimated  
593 solid discharge with the value provided by the global relationship established from 40-years  
594 of data (method A). The average error for daily  $Q_s$  values was 51% using method A and  
595 42.1% using method B. However, the cumulative flux of suspended matter over the 1213 days  
596 for which daily data are available was over-estimated by 3.1% using method A while it was  
597 under-estimated by 5% using method B. A comparison of the estimates by these two methods  
598 showed that method B is not reliable for high discharge during the last decade because of an  
599 increase in scattering of the C, Q pairs. The relationship obtained over the last decade (2000-  
600 10) lead to an under-estimation of  $Q_s$  of 23% over the 314 days for which daily C and Q are  
601 known. In contrast, the global algorithm from method A led to an under-estimation of the  
602 same cumulated  $Q_s$  by only 3.5% over the same period. The relationship established over 40  
603 years was therefore used for this study.

604 It should be noted that although method A provides some daily solid discharges from the  
605 1213 daily Q values with a high error (the average error being 51%) it enabled the  
606 reconstruction of good trends of  $Q_s$  values over more than 7 orders of magnitude (Fig. 53).  
607 However, the temporal variability of the coefficients a, b of the rating curves calculated over  
608 years or decades will be discussed in light of the variability of the forcings and their sediment  
609 transport consequences, to [understand better](#) their physical meaning.

### 610 **3.3 Average loads**

611 In order to analyze the temporal variability of suspended sediment flux, we use the average  
612 concentration resulting from the ratio between the solid and the liquid flow rate, denoted  
613 (SPM\*), which can be defined for any integration period (day, month, season, year).

### 614 **3.4 Study of breaks: double mass curve**

615 Double-mass curves were used to determine long term trends and changes in the hydro-  
616 sedimentary regime (Searcy and Hardison, 1960; Walling, 1997; 2006).

## 617 **4 Interannual variations of temperature, precipitation, river discharge and** 618 **flow regime**

619 The statistics of hydrological parameters at Ain Hamara gauging station over 1970-2010 are  
620 reported in Table 1.

### 621 **4.1 Temperature**

622 Temperature in Northern Algeria at the three stations of Chlef, Miliana and Dar El Beida  
623 increased from the 1970s onwards (Fig. 64). On average, temperature was higher at Chlef  
624 (between 17.5°C and 20.3°C) than at Dar El Beida (15-18.5°C) and Miliana (14.5-18.5°C). In  
625 average, temperature at Chlef increased by 0.96°C between 1970-85 and 1985-2010, and by  
626 1.17°C from the 1970s to the 2000s (Table 2). The increase was, on average, 0.87°C between  
627 the 1970s and the 1980s which is more than four times the difference between the 1980s and  
628 the 1990s (+0.19°C) and the 1990s and the 2000s (+0.12°C). As has been shown on a global  
629 scale, the decade of the 2000s was the warmest (IPCC, 2013).

### 630 **4.2 Precipitation**

631 Annual precipitation at Ain Hamara station was highly irregular, varying between 165 mm yr<sup>-1</sup>  
632 and 506 mm yr<sup>-1</sup> (Table 1, Fig. 75). Mean annual precipitation (*P*) was 264 mm, with a  
633 coefficient of variation (CV) of 27% between 1970–71 and 2009–10. The interannual  
634 variations of *P* (Fig. 75) showed trends towards a decrease of rainfall (-1.86 mm yr<sup>-1</sup> on  
635 average over 40 years, *p* < 0.05). *P* decreased by 15 % (from 310 to 264 mm) between 1970s  
636 and 2000s if we consider the values averaged over decades (Table 2). However, a more  
637 precise analysis shows that rainfall greatly decreased from 1970s to the next decade (from 310  
638 to 231 mm, -25%), then slightly increased in the two following decades (average of 250 mm  
639 yr<sup>-1</sup> in 1990s and 264 mm yr<sup>-1</sup> in 2000s, see Table 2).

640 The average precipitation over the 6 rainfall gauging stations within the basin was 273 mm yr<sup>-1</sup>  
641 <sup>1</sup>. Their interannual variations were consistent and showed a similar variation to Ain Hamara  
642 station. Amongst decades, the coefficient of variation varied between 12% and 20%. Five out  
643 of 6 stations show a decrease in precipitation between 1970-1985 and 1985-2010, the average  
644 deficit being equal to 3.7 %.

### 645 **4.3 River discharge and flow regime**

646 The mean annual discharge at the Ain Hamara gauging station was  $1.18 \text{ m}^3 \text{ s}^{-1}$  over the 40-  
647 year period of observation (Table 1). The interannual variability of yearly averaged values of  
648 discharge (CV=44.4%, see Table 1) was higher than that of yearly precipitation. Yearly  
649 averaged values of  $Q$  showed a trend towards an increase of river flow ( $+11.3 \text{ L s}^{-1} \text{ yr}^{-1}$  on  
650 average over 40 years,  $p < 0.01$ ; Fig. 75). The averaged values over decades decreased  
651 between the 1970s and the 1980s, then increased (Table 2). Globally, they increased by 25%  
652 (from  $1.16$  to  $1.45 \text{ m}^3 \text{ s}^{-1}$ ) between 1970-80 and 2000-2010.

653 The detailed analysis of the daily river discharge shows that the river was perennial in the  
654 1970s and then became intermittent during the 1980s (Fig. 86). The driest year occurred in  
655 1993-94 with 117 days of fully dry river. On Fig. 86, the very low river discharges (around  
656  $0.01 \text{ m}^3 \text{ s}^{-1}$ ) were not considered as days of dry river.

657 The “wet discharge”, denoted  $Q_w$ , i.e. the yearly average river discharge of the days of  
658 running river (and not calculated over the full year) was also calculated (Table 2). Over the  
659 40-year period when  $Q$  increased by 25%,  $Q_w$  averaged over 10-years increased by more than  
660 35% from 1970-80 to 2000-2010 (from  $1.16$  to  $1.57 \text{ m}^3 \text{ s}^{-1}$ ).

661  $Q$  and  $Q_w$  increased as did the number of dry days (and consequently the intra-annual  
662 variability) and their intra-decade variability (Table 2). Two indicators of intra-annual  
663 discharge variability are shown in Fig. 75:  $Q_{98}$ , the 98th percentile of annual flows calculated  
664 from daily discharge and the standard deviation of daily discharge within each year ( $\sigma_Q$ ).  $Q_{98}$   
665 increased from an average of  $4.37 \text{ m}^3 \text{ s}^{-1}$  over the period 1970-80 to  $13.94 \text{ m}^3 \text{ s}^{-1}$  over the  
666 period 2000-2010, a factor 3.2 increase.  $Q_{98}$  is also a good indicator of changes in sediment  
667 transport as it occurs during the highest flood events that occur each year.

### 668 **4.4 Summary: changes of hydrologic forcings**

669 These results indicate that four significant changes occurred during the 40-year period in the  
670 Wadi Abd basin (Table 2): (1) an increase of temperature at Chlef by  $1.17^\circ\text{C}$  between the  
671 1970s and the 2000s; (2) a decrease in precipitation of 15% over 4 decades; (3) an increase in  
672 average annual flow of 25 % over the same period, or 35% if we consider only the days when  
673 the river is not dry; (4) a change in the flow regime, from a perennial regime to intermittent  
674 regime. The pivotal year after from which the river experiences dry weeks is the hydrological

675 year 1985/86 with 49 days with no flow. This number increased in the following years (Fig.  
676 86).

## 677 5 Interannual variation of sediment load

### 678 5.1 Rating curve

679 The rating curve obtained from 1213 pairs of daily averages gave:

$$680 C = 2.270 Q^{0.647} \quad (3)$$

681 where C is expressed in  $\text{g L}^{-1}$  and Q in  $\text{m}^3 \text{ s}^{-1}$ . 43% of the variations of C are explained by  
682 those of Q ( $r^2=0.431$ ). The rating curve obtained between Q and  $Q_s$  shows a much higher  
683 determination coefficient ( $r^2=0.831$ ) but is biased since  $Q_s = C \times Q$ . Nevertheless, both  
684 relationships give estimates of  $Q_s$  values from Q with less than 1% difference which is less  
685 than the uncertainty of  $Q_s$ .

### 686 5.2 Yearly sediment fluxes and concentrations

#### 687 Decadal variability of $Q_s$

688  $Q_s$  increased from 180 to 1130  $10^3$  tons per year between the 1970s and the 2000s (Table 2).  
689 The increase from one decade to the next is remarkably regular: +85% between the 1970s and  
690 the 80s, + 84% between the 80s and the 90s, +84% between the 90s and the 2000s and is  
691 statistically significant ( $+19.7 \text{ } 10^3 \text{ t yr}^{-1}$  in average,  $p < 0.05$ ). Specific sediment yield follows  
692 the same trend increasing from  $72 \text{ t km}^{-2} \text{ yr}^{-1}$  in the 1970s to  $455 \text{ t km}^{-2} \text{ yr}^{-1}$  in the 2000s.

#### 693 Variability of mean annual load SPM\*

694 The average value of SPM\* calculated over the period 1970-2010 ~~from daily concentration~~  
695 ~~and discharge~~ is  $12.3 \text{ g L}^{-1}$ . The 40 annual values of SPM\* calculated for each year ~~from~~  
696 ~~measured discharges and concentrations estimated using the rating curve (3)~~ vary between 2.5  
697  $\text{g L}^{-1}$  and  $50.2 \text{ g L}^{-1}$  (Tables 1, 2). Their interannual variation was smaller than that of solid  
698 discharge because annual SPM\* is the ratio of the annual  $Q_s$  to the annual Q (which increased  
699 less than  $Q_s$ ). The variability of SPM\* was thus smaller than that of annual  $Q_s$  (CV=86.0%  
700 instead of 123.3% over 40 years).

#### 701 Analysis of break points

702 The double mass plot enabled us to identify changes in the sediment response of the stream  
703 (Fig. 97). A major break occurred in 1985-86. A secondary break was noticed in 1991-92, but



704 the entire period 1985-86/2009-10 may be considered as a single period (with the relationship  
705 « cumulated  $Q_s$  » = 0.021 « cumulated  $Q$  » - 9.417,  $r^2=0.989$ ). The period 1985-86/1991-92  
706 may thus be considered as a transient event towards a new regime.

707 The response of sediment flow to various constraints (changes in precipitation, hydrology,  
708 plant, agricultural practices etc.) differs clearly from that of discharge from the year 1985-86  
709 onwards. This break corresponds to the first year of dry river over a long period in summer  
710 (49 days). This initiates a phase of intermittent flow regime. The averaged parameters for the  
711 two periods 1970-1985 and 1985-2010 were added to the tables, in addition to average values  
712 throughout the full study period and values for decades to illustrate the dynamics of the  
713 hydrological and hydro-sedimentary change.

### 714 **5.3 High dependency of the solid discharge on Q variability**

715 The variability of  $Q$  and  $Q_s$  or  $SPM^*$  at different time scales were compared. AO2007 showed  
716 that, over 22 years, 71% of the variance of the annual  $SPM^*$  values was accounted for by the  
717 annual discharge and 73% by the 95<sup>th</sup> percentile of daily discharge within the given year  $Q_{95}$ .  
718 This means that  $SPM^*$  was mainly driven by the 10 to 15 highest daily discharges in a year  
719 suggesting a strong correlation between yearly  $Q_s$  and the discharge variability. Finally, they  
720 showed a remarkable linearity between  $SPM^*$  and the standard deviation of the daily  
721 discharge per year ( $\sigma_Q$ ).

722 Yearly  $SPM^*$  and yearly  $\sigma_Q$  still showed a strong linearity over 40-years ( $r^2=0.956$ , Fig.  
723 108a). A higher correlation was obtained between yearly  $Q_s$  or  $SSY$ , the specific sediment  
724 yield, and yearly  $\sigma_Q$  ( $r^2=0.991$ , Fig. 108b). This is one of the most important conclusions from  
725 this river where the solid discharge depends on discharge following a rating curve: the yearly  
726 solid discharge is more closely dependent on the discharge variability than on discharge  
727 values.

### 728 **6 Variation of the seasonality of climatic and hydrological parameters**

729 The yearly values of temperature at Chlef increased on average but the monthly averages  
730 showed high discrepancies. Temperature from March to November increased with a  
731 maximum of increase in June (+3.30°C on average between the 1970s and 2000s), it remained  
732 quite constant in December and February and decreased by 0.98°C in January over the same  
733 period. Considering the average values per season, winter values (Dec-Feb) decreased by  
734 0.33°C between the 1970s and the 2000s, while spring values (Mar-May) increased by

735 1.66°C, summer values (Jun-Aug) by 2.22°C and fall values (Sep-Nov) by 1.29°C. In  
736 summary, annual temperature differences increased with minimum temperatures down  
737 slightly and maximum temperatures rising sharply. The increase was most marked in July-  
738 August.

739 Averaged seasonal values of P, Q and  $Q_s$  for each decade are given in absolute values and in  
740 percent of the yearly values in Table 3. The seasonal relative contribution of P, Q and  $Q_s$   
741 centered and averaged over 9 consecutive years are presented in Figure 119. The monthly  
742 values of P, Q and  $Q_s$  per decade over 40-years also clearly illustrate the absolute changes in  
743 intensity and in seasonality of the river regime (Fig. 1240). The main conclusions of the  
744 analysis of T, P, Q and  $Q_s$  variations are the following:

745 • Rainfall decreased in spring and increased in autumn. Precipitation in autumn increased  
746 from 22 to 30 % at the expense of spring rains (decreasing from 41% to 29%). It is  
747 striking to note that for the decade 2000-2010 precipitation was the same in autumn and  
748 in spring (78 mm) while for the decade 1970-1980 spring rainfall was 87% higher than in  
749 fall (128.2 mm vs. 68.5 mm; see Table 3 & Fig. 119a).

750 Average monthly rainfall from six weather stations in the river basin for 1970-1985 and  
751 1985-2010 (Fig. 1344) illustrates the changes. Two marked seasons typical of a  
752 Mediterranean climate are present (a dry season and a rainy season) but the following  
753 changes are observable: (1) differences between seasons decrease, as indicated by the CV  
754 of monthly rainfall from 57.3 % in 1970-85 to 45.9 % in 1985-2000. There is a decrease  
755 of spring rains (March-May) and at the beginning of the cold season (November-  
756 December) and the strengthening of rain in the warm season (July-October) and in winter  
757 (January-February) ; (2) advancement of the rainy season as evidenced by precipitation in  
758 October and November; (3) spreading of the rainy season over 9 months (September-  
759 May) for 1985-2010 from previously 7 or 8 months (from October or November  
760 onwards, according to the criteria that are defined for the rainy season) ; (4) increased  
761 regularity of rainy season precipitation.

762 • Proportionally, flow decreased in all seasons from winter to summer and increased  
763 dramatically in autumn from just over a quarter (27.3%) of the flow delivered over the  
764 decade 1970-1980 to more than one half (52.5%) over the period 2000-2010 (Table 3 and  
765 Fig. 119b). Flow decreased in summer and the river became dry for much of the summer.  
766 Over the last decade, it is striking to see the difference between the average flow rates in  
767 fall and spring: the fall rate is almost three times that of spring with almost the same  
768 rainfall. This trend is evident over the 40 year period (Fig 119b).

- 769 • These results point towards a change in runoff as defined by the ratio Q/P. Considering  
770 the whole basin area, the river discharge at Ain Hamara station averaged over 40-years  
771 corresponds to a water depth of 15 mm yr<sup>-1</sup>, while the average precipitation is 264 mm yr<sup>-1</sup>.  
772 For comparison, on average 85% of rain in this region evaporates and the remaining  
773 15% runs into surface waters or infiltrate into underground storage (Sari, 2009, quoted by  
774 Benhamiche et al., 2014). On the Wadi Abd, Q/P averages 5.7%. We calculated the value  
775 of Q/P averaged over 3 consecutive years and over 3 consecutive months (centered) and  
776 then took the average per decade (Fig. 1412). It appears that the Q/P ratio remains  
777 constant during the months from December to April (around 4.4% in average), it  
778 increased slightly in November and May during the decade 2000-2010 and it increased  
779 significantly from September to November. In other words, runoff increased, rain  
780 decreased slightly and the temperature (and therefore ETP) increased. As a consequence,  
781 infiltration will decrease and the water level in the aquifers will be lowered. Moreover,  
782 Q/P, which was very high in July and August in 1970s, has nearly halved since 1980s.
- 783 • In absolute value, solid discharge has been increasing in all seasons over 4 decades, but  
784 more so in the fall than in the other seasons (Table 3 and Fig. 1240c). During autumn, it  
785 more than doubled from one decade to another (x 2.07 in the 1980s vs. 1970s, x 2.17  
786 from 1980s to 1990s, and x 2.88 from 1990s to 2000s). During the other seasons, it  
787 doubled or tripled within 30 years, between 1970s and 2000s. The average annual load  
788 was multiplied per 1.84 from one decade to another (1.8 10<sup>6</sup> tons during the 1970s, 3.34  
789 in 1980s, 6.14 in 1990s and 11.30 in 2000s, see Table 2). While during the 1970s the  
790 Wadi Abd had two major periods of roughly equivalent sediment discharge in the fall and  
791 spring, suspended sediment loads were greater in the autumn during the 2000s (> 70%).  
792 The Wadi shifted from a regime with two equivalent seasons of sediment production to a  
793 regime with one dominant season in the 2000s. Autumn produced over 4 times more  
794 sediment than spring in 2000s (Table 3, Fig. 119c). This phenomenon does not seem to  
795 be due to some exceptional floods because the trend is observable over 4 consecutive  
796 decades (Fig. 119c).

## 797 **7 Discussion**

### 798 **7.1 Interannual variations**

#### 799 Hydrology and climate change over 40 years

800 Temperature increased rapidly between the 1970s and 1980s (+0.88°C on average at Chlef).  
801 The increases were lower during the following decades (1980s to the 2000s). An increase in  
802 temperature of 1.6°C between 1977-1979 and 2000-2006 was noted by Dahmani and Meddi  
803 (2009) for the Wadi Fekan basin in West Algeria and Bakreti et al. (2013) also showed a  
804 significant trend of increasing temperature in spring by 0.0183 °C per year in the Tafna basin  
805 in West Algeria over the same period. However, temperature did not increase so fast during  
806 the whole 20th century (Fig. 64) and as mentioned by IPCC (2013), “trends based on short  
807 records are very sensitive to the beginning and end dates and do not in general reflect long-  
808 term climate trends.” The longest available time series of temperature in Algeria was  
809 measured at Dar El Beida near Algiers. At this station, average temperature increased by 0.62  
810 °C between 1850-1900 (29 yearly values available) and 2003-2012 (Fig. 64), while it  
811 increased between 1880 and 2012 by 0.85°C globally (IPCC, 2013).

812 A global trend towards an increasing temperature and increasing dryness in Algeria from the  
813 1970s onwards has already been described (Meddi and Meddi, 2009). Over the period 1923-  
814 2006 North Algeria experienced an alternation of wet periods (1923-1939, 1947-1973) and  
815 dry periods (1939-1946 and from 1974 onwards) (Benhamiche et al., 2014). Over 70 years in  
816 the Wadi Fekan, Dahmani and Meddi (2009) showed that the period 1943-1960 was rather  
817 wet, that 1960-1975 was average, and that the period 1975 onwards (up to the end of their  
818 data set in 2004) was dry and of an exceptional long duration. Using three different statistical  
819 tests (Pettitt, Lee Heghinian and Hubert), Meddi and Meddi (2007) shown that a **shift** was  
820 observed between 1973 and 1980 over most of the rain gauges in Algeria. In North-West  
821 Algeria, a **shift** was noticed in 1973 in winter rainfall and between 1974 and 1980 in spring  
822 rainfall, both of them being responsible of the yearly rainfall deficit (Meddi and Talia, 2008).  
823 From the rainfall dataset at the Ain Hamara station between 1968 and 2007, Hallouz et al.  
824 (2013) showed that the **break** in annual rainfall occurred in 1976 and calculated a deficit of  
825 19% between 1968-1976 (304 mm yr<sup>-1</sup>) and 1976-2007 (247 mm yr<sup>-1</sup>). At the stations Ponteba  
826 and Rechaiga, near to the Abd basin, the trends of decreasing total precipitation and of  
827 increasing mean length of dry spells were amongst the 5 highest in the Maghreb area over the  
828 22 stations considered by Trambly et al. (2013, see their Fig. 86).

829 As a consequence of the decrease of rainfall after the 1970s **break** which was observed in  
830 most basins of Western Algeria, river discharges were generally seen to decrease as well.  
831 Meddi and Hubert (2003) showed that the decrease in river discharge varied between -37%  
832 and -70% from the Eastern Algeria to the Western Algeria. Over the Mecta basin in North-

833 West Algeria, runoff was estimated to be 28-36% lower in 1976-2002 as compared to 1949-  
834 1976 (Meddi et al., 2009). Over the Tafna basin also in North-West Algeria, Ghenim and  
835 Megnounif (2013a, 2013b) showed that the decrease of precipitation by 29% on average over  
836 the basin (especially in winter and spring) after the **break point** was accompanied by a  
837 decrease of 60 % in river flow.

838 In this context, the Wadi Abd had a different behavior since the river discharge increased. The  
839 counter-intuitive increase of runoff with decreasing rainfall has also been observed in Sahel  
840 and is referred to as « the Sahelian paradox » (see Mahé and Paturel, 2009; Mahé et al., 2012).  
841 A closer look at the seasonal variations of the different parameters shows that Q decreased in  
842 winter and spring but that Q/P increased in autumn when rainfall increased. Overall Q  
843 increased. The decrease of rainfall in spring and its low level in summer may have lead to a  
844 change in vegetation cover which would in turn decrease **ef**-infiltration. However, although  
845 studying the vegetation dynamics of the basin goes beyond the scope of this study, this aspect  
846 could be investigated in the future using satellite data, for example.

847 What is the influence of large-scale circulation indices?

848 Changes in precipitation are derived from atmospheric-oceanic signals (Milliman et al., 2008;  
849 Giuntoli et al., 2013). Low frequency fluctuations related to climate change are modulated  
850 with higher frequency interannual fluctuations, such as ENSO (El Niño Southern Oscillation),  
851 NAO (North Atlantic Oscillation), AMO (Atlantic Multidecadal Oscillation) or MO  
852 (Mediterranean Oscillation). Trambly et al (2013) showed that the precipitation amounts and  
853 the number of dry days **over the Maghreb** were significantly **correlated with** the MO and  
854 NAO patterns. MO and NAO showed positive trends from the 1970s onwards which could  
855 explain the trend towards decreasing frontal conditions over the Mediterranean basin and thus  
856 increasing droughts.

857 Interannual influence by the Austral oscillation ENSO over Algeria was shown to be higher in  
858 North-West Algeria **on the highest discharges than on the average discharge**. The maximum Q  
859 seems to be smaller during El Niño and higher during La Niña in North-West Algeria (Ward  
860 et al., 2014). Average discharge is less influenced by ENSO than the maximum yearly  
861 discharge (Ward et al., 2014). The frequency of extreme rainfall events shows the highest  
862 correlation with the Mediterranean Oscillation Index in Algiers and with the Southern  
863 Oscillation Index in Oran (Taibi et al., 2014).

864 In this study, no significant correlation was established between a series of hydrological  
865 parameters in the Wadi Abd and the Southern Oscillation Index. The average of AMO per  
866 hydrologic year was calculated from its monthly values. AMO has increased from 1970s to  
867 the 2000s, with negative values up to 1993-94, then positive afterwards (except in 1996-97).  
868 Its decadal average was -0.25 in the 1970s, -0.12 in the 1980s, 0.0 in the 1990s and 0.18 in the  
869 2000s. AMO and the discharge variability of the Wadi Abd within the year increased  
870 coincidentally. The yearly AMO values have a coefficient of determination of 0.226 when  
871 correlated with the standard deviation of daily river discharges within the year, a proxy for the  
872 variability of daily discharge. However, this information does not allow us to conclude that  
873 the Atlantic Multidecadal Oscillation is responsible for hydrological changes in the Wadi Abd  
874 basin.

#### 875 Break point in 1985-86: change of flow regime

876 The several weeks of dry river for the first time in 1985-86 (49 days) can be considered as a  
877 threshold effect, which marks the start of a new flow regime. The appearance of a dry regime  
878 is a break, a fully nonlinear phenomenon. It has strong consequences for water infiltration and  
879 groundwater recharge, on seasonality, intensity and type of floods, and in turn, on erosion and  
880 sediment transport. 1985 is also a pivotal year for recent climate change as evidenced by the  
881 rapid increase in global mean temperature anomaly of air from that year until 1993 (Fig 1 in  
882 Lockwood and Fröhlich, 2007). The hypothesis of a temporary warming caused by dust  
883 emitted during the eruption of Mount Pinatubo had been advanced to explain the warming  
884 since 1985, but climate scientists later recognized that the temperature anomaly has been  
885 increasing since 1993, reaching about 0.6°C by 2007 compared to the global average  
886 temperature calculated for the period 1951-1980 (Lockwood and Fröhlich, 2007).

887 This threshold is coincident with hydrological shifts in the Tafna basin in North-West Algeria.  
888 Bakreti et al. (2013) analyzed the baseflow and baseflow index of five of its sub-basins  
889 between 1976 and 2006 and evidenced ruptures of the baseflow index between 1984 and 1990  
890 depending on the sub-basin, in 1984, 1985 and 1990 in the mountains, and in 1985 and 1986  
891 in the plain. These changes in flow regimes of the Tafna basin were likely caused by shifts in  
892 rainfall late 1970s in the Mounts of Tlemcen and early 1980s in the plains (Ghenim and  
893 Megnounif, 2013a).

#### 894 Shift of the onset of the first summer flood

895 The analysis of the time series of daily flows enables to determine the start of the first  
896 summer flood. The average daily flow per decade suddenly increases the day at which the  
897 first summer flood occurred, at least once in the decade. By observing these decadal averaged  
898 daily flows, there is no ambiguity on the start of the earlier flood by decade:

899 - in 1970-80, the first flood starts on ~~average~~ the 6th September with an average 4-days  
900 discharge (6-9 September) of  $1.59 \text{ m}^3 \text{ s}^{-1}$ , while it was on average  $0.58 \text{ m}^3 \text{ s}^{-1}$  over the four  
901 previous days,

902 - in 2000-2010, the first flood of summer starts on August 8 with an average 4-days discharge  
903 (8-11 August) of  $2.03 \text{ m}^3 \text{ s}^{-1}$ , while it was on average  $0.03 \text{ m}^3 \text{ s}^{-1}$  from 4 to 7 August.

904 During the 2000s, the first flood in summer started close to one month before that of the  
905 1970s and the magnitude was 27% higher. It can be asked if this trend was observable over  
906 the 40-year period or only between two specific decades. The analysis of mean dates and  
907 discharges of the first flood in late dry season gave the following results for the intermediate  
908 decades:

909 - 1980-1990: the first flood started in average on August 31 with a 4-days average discharge  
910 (August 31-September 3) of  $2.69 \text{ m}^3 \text{ s}^{-1}$ , while the average rate over the four previous days  
911 was  $0.13 \text{ m}^3 \text{ s}^{-1}$

912 - 1990-2000: the first flood started in average on August 22 with a 4-days average discharge  
913 (August 22-25) of  $7.67 \text{ m}^3 \text{ s}^{-1}$ , while the average rate over the four previous days was  
914  $0.33 \text{ m}^3 \text{ s}^{-1}$ . The existence of a precursor peak on August 17, which was not observed in  
915 previous decades, was also observed.

916 It therefore appears that the date of the first flood advanced by about ten days each decade  
917 over the previous 40 years. The shift in the onset of the first flood in summer probably has  
918 important consequences on flow and erosion rates.

## 919 **7.2 Relationships between several parameters and sediment yield**

### 920 Temperature and sediment yield

921 The curve showing annual suspended load versus global air temperature anomaly (base period  
922 1951-1980) calculated by hydrological year from monthly data provided by NOAA (Hansen  
923 et al., 2010; GISTEMP Team, 2015) shows a correlation between the sediment yield and  
924 ongoing climate change ( $r^2=0.388$ , Fig. 1513).

925 Precipitation and sediment yield

926 Many authors studied the variations of sediment load per unit of catchment area against  
927 annual rainfall (e.g. Summerfield and Hulton, 1994) or effective rainfall (e.g. Langbein and  
928 Schumm, 1958). On the Wadi Abd, annual rainfall was 310 mm yr<sup>-1</sup> in the 1970s, fell sharply  
929 in the 1980s then slightly increased over the following decades to between 231 and 264 mm  
930 yr<sup>-1</sup>. Meanwhile, yearly sediment concentration and suspended sediment discharge have  
931 increased. The comparison of their respective variations shows a lack of correlation between  
932 precipitation and annual sediment yield ( $r^2 < 0.1$  regardless of the type of regression  
933 considered). Regarding the relationship between precipitation and erosion, if there are  
934 correlations between their spatial variations reported in the literature (though with a strong  
935 scatter, see Riebe et al., 2001), our study shows that the temporal variations of precipitation  
936 and sediment yield are not correlated in the Wadi Abd. This may be due to the change of flow  
937 regime within the study period.

938 Runoff and sediment yield

939 Although runoff was noted to have a limited impact on the distribution of sediment yield at  
940 regional or global scales by Aalto et al. (2006), Syvitski and Milliman (2007), Vanmaercke et  
941 al (2014), the temporal variability in precipitation, runoff (or discharge) and consecutive  
942 vegetation cover was shown to be locally the main impact on fluvial sediment load (see the  
943 review of Vanmaercke et al. 2014, p. 360). Our results confirm that, on the Wadi Abd, the  
944 yearly suspended sediment load was highly correlated with discharge (Q mean or its highest  
945 percentiles) and to its intra-annual fluctuation with the highest percentiles of variability of  
946 discharges, and especially to its highest values (Q<sub>98</sub>), the best correlation being obtained with  
947 the standard deviation of daily discharge within the year (Fig. 108). Climate change alters the  
948 hydrology of a river basin such as the Wadi Abd. Although the river regime shift clearly  
949 impacted several parameters between the two periods, the relationship between yearly  
950 sediment load and discharge variability did not change over the study period.

951 ~~Climate change alters the hydrology of a river basin such as the Wadi Abd. We show that~~  
952 ~~suspended sediment yield was highly correlated with discharge (Q mean or its highest~~  
953 ~~percentiles) and to its intra-annual fluctuation (Fig. 8).~~ We cannot conclude on the exact  
954 origin of the regime change but note that it occurred when dry periods started, annual  
955 precipitation timing shifted and runoff increased.



956 **7.3 On the use of double-mass curves to determine the climate change and**  
957 **anthropogenic influences**

958 Double-mass curves are often used to determine the impact of developments such as dams on  
959 sediment discharge (e.g. Lu et al., 2013). Our findings warn about extrapolations that could be  
960 wrongly made to quantify the impact of a development by extending the double mass curves.  
961 Indeed, this study shows that the double-mass curve can change its slope (here increasing)  
962 when the flow regime change is driven by seasonal temporal variation in precipitation and  
963 runoff that isn't linked to any specific anthropogenic activity (such as a dam impoundment)  
964 within the basin.

965 **7.4 Physical meaning of rating parameters a & b**

966 Interannual variation of (a, b)

967 Since  $C = a Q^b$ , with  $b \neq 0$ ,  $C(1) = a$ . a thus represents the sediment concentration when the  
968 river discharge is  $1 \text{ m}^3 \text{ s}^{-1}$ , and b reflects the sensitivity of concentration to discharge  
969 variation. The general formula  $\ln C = \ln (aQ^b)$  provides:

970 
$$dC/C = b dQ/Q \tag{4a}$$

971 
$$b = dC/dQ Q/C = 1/a dC/dQ Q^{(1-b)} \tag{4b}$$

972 thus b varies almost like 1/a (Asselman 2000). Many papers discuss the physical meaning of  
973 the rating parameters a and b (see AO2007) and try to connect their values to physiographical  
974 characteristics, vegetation cover or hydro-meteorological forcing. ~~A study such as the present  
975 one on a single basin avoids physiographical variations and enables the analysis of the  
976 dependence of a and b on the hydro-meteorological forcing and, if data are available, on  
977 vegetation cover and land use.~~

978 ~~Over 40 years the variations of (a, b) averaged over one or several years can be instructive  
979 (Fig. 14). In the present study, we only considered the yearly (a, b) values corresponding to  
980 yearly C-Q rating curves with  $r^2 > 0.21$  (or to yearly Q-Q<sub>s</sub> rating curves with  $r^2 > 0.71$  in our  
981 dataset. Based on this criterion, 3 yearly (a, b) values over 40 for the hydrological years  
982 1972/73, 1975/76 and 2009/10 were removed from the present analysis).~~

983 ~~At yearly scale, a explains 58% or 66% of the variance of b depending on the considered  
984 relationship, respectively (Fig. 14):~~

985 ~~$$b = -0.294 \ln a + 0.912 \tag{5a}$$~~

986  ~~$\ln b = -0.188 a + 0.042$  (5b)~~

987 ~~Relationship 5a is close to the one established for 138 flood episodes between 1973 and 1995~~  
 988 ~~( $b = -0.311 \ln a + 1.066$ ) (AO2007).~~ The river's regime change is accompanied by a change  
 989 in the (a, b) pairs of rating curves defined for multi-year periods such that a increases and b  
 990 decreases (Table 2), following:

991  $b = -0.294 \ln a + 0.912$  ( $r^2=0.582$ ) (5a)

992  $\ln b = -0.188 a + 0.042$  ( $r^2=0.649$ ) (5b)

993 Equation (5a) is ~~also~~ very similar to that presented by Iadanza and Napolitano (2006) for the  
 994 Tiber River after the construction of a dam ( $b = -0.3815 \ln a + 0.7794$ ,  $r^2=0.992$ ). Before the  
 995 construction of this dam, another relationship ( ~~$b = -0.4457 \ln a + 0.9615$ ,  $r^2 = 0.991$ )~~  
 996 corresponded to more than 3 times higher sediment yields. Asselman (2000) suggested to  
 997 interpret regression lines in a  $\ln a - b$  graph as different sediment transport regimes.

998 On the Wadi Abd, the change in sediment transport regime is not evident from the yearly (a,  
 999 b) values but it becomes clearly observable when considering a and b values averaged over  
 1000 moving periods of several years. The best correlations were obtained for running averages  
 1001 over 15 years named  $a_{15}$  and  $b_{15}$  (N=25, from 1970-1985 to 1995-2010, see Fig. 165). The  
 1002 available data set does not allow us to determine if results obtained from averaging over  
 1003 longer periods would perform best.

1004 The time evolution of the moving average pair ( $a_{15}$ ,  $b_{15}$ ) clearly shows a first relationship with  
 1005 the values dominated by the pre-1985 regime (8 values from 1970-1985 to 1977-1991),  
 1006 another one for the values predominantly after 1990 (12 values from 1983-1997 to 1995-  
 1007 2010), both with  $a_{15}$  increasing and  $b_{15}$  decreasing, and a transitional regime centered on the  
 1008 period 1985-1990 (Fig. 165). ~~They gave:~~

1009  ~~$b_{15} = -0.178 a_{15} + 1.043$  ( $r^2=0.960$ , N=8)~~ (6)

1010 ~~from 1970-1985 to 1977-1991 (average values calculated with mainly pre-1985 data), with  $a_{15}$~~   
 1011 ~~increasing and  $b_{15}$  decreasing, and~~

1012  ~~$b_{15} = -0.126 a_{15} + 1.021$  ( $r^2=0.982$ , N=12)~~ (7)

1013 ~~from 1983-1997 to 1995-2010 (mainly post-1990).~~ During the transition period centered over  
 1014 1985-1990,  $b_{15}$  was almost constant (between 0.72 and 0.74) while  $a_{15}$  was increasing from  
 1015 2.01 to 2.34. During the period 1985-1991 the yearly values of b varied very little (between  
 1016 0.653 and 0.672) while yearly a increased significantly from 1.81 in 1985-86 to 3.23 in 1990-

1017 91. Higher a and lower b values are in the literature typical of highly arid river basins, such as  
1018 the ephemeral Nahal Eshtemoa in Israel, where  $a=16.98$  and  $b=0.43$  (Alexandrov et al., 2003).  
1019 ~~The changes of moving average ( $a_{15}$ ,  $b_{15}$ ) pairs are associated to given river basin and types of~~  
1020 ~~transported particles. Their change depends only on the hydro-meteorological and vegetation~~  
1021 ~~forcings for a given physiography (slope, geology). As the break points ruptures were~~  
1022 coincident, it is possible to analyze the change of ( $a_{15}$ ,  $b_{15}$ ) in terms of shift of hydrological  
1023 regime ~~The two relationships before and after the transition can be considered as signatures of~~  
1024 ~~the dominant hydrological regime. However, if the new hydrological regime was immediate~~  
1025 from 1985 onwards, the change in the C-Q relationship was only evidenced in the Wadi Abd  
1026 at mid-term, considering 15-years average values ~~We can thus conclude that a true change in~~  
1027 ~~sediment transport regime occurred on the Wadi Abd basin, with. Within this change of~~  
1028 ~~regime, the rating parameters showed a clear tendency to evolve towards increasing values of~~  
1029 ~~a and decreasing values of b (Table 2). High a and low b are in the literature typical of A~~  
1030 ~~similar trend is generally considered in the literature as typical of an increasing aridity. One~~  
1031 ~~example of highly arid river basins, such as was given by Alexandrov et al (2003) for the case~~  
1032 ~~of the ephemeral Nahal Eshtemoa in Israel, where  $a=16.98$  and  $b=0.43$  (Alexandrov et al.,~~  
1033 ~~2003).~~

1034 ~~Studying the impact of the Three Gorges dam over the Yantgze River, Wang et al. (2008)~~  
1035 ~~showed that a decreased and b increased after dam impoundment and associate this change to~~  
1036 ~~a decrease in sediment supply from the basin (with a) and an increase of the erosive power of~~  
1037 ~~the river (with b) which scours the river downstream of the dam. Considering the two main~~  
1038 ~~sediment sources (the basin and the river bed) and applying the same reasoning, we could~~  
1039 ~~infer that the erosive power of the river decreased while the erosion of the basin increased.~~  
1040 ~~This means that in the Wadi Abd climate change would favour the river basin as a sediment~~  
1041 ~~source to the detriment of the bed.~~

#### 1042 Parameters that explain a (or b)

1043 ~~The value of a (or b, which is deduced from a) varies with the hydro-meteorological and~~  
1044 ~~vegetation forcing. The annual average liquid flow explains only 63.3% of the variance of a.~~  
1045 The coefficient of determination between a and specific sediment yield (SSY) is low at the  
1046 annual scale but higher when we consider the moving averages of a and SSY over 15-years.  
1047 The specific sediment yield explained 95.2% of the variance in the interannual scale (Fig.  
1048 176a), much more than the average river flow did ( $r^2= 0.839$ ; see Fig. 16b), following:

1049  $a_{15} = 6.104 \cdot 10^{-3} \text{ SSY}_{15} + 1.117 \quad (r^2=0.952)$  (86)

1050  $b_{15}$  showed a lower correlation with the SSY (Fig. 16e  $r^2=0.853$ ) than  $a_{15}$  did.

1051 In summary, ~~our results indicate that~~ the moving average of  $a$  is strongly correlated to specific  
1052 ~~sediment yield~~ over the same moving period of 15 years, and ~~that~~ the moving average of  $b$  can  
1053 be deduced from  $a$  over the same period using a relationship which is given per flow regime,  
1054 either perennial or intermittent. ~~During the transition to an intermittent regime,  $b$  remained~~  
1055 ~~almost constant while  $a$  increased. The coincident increase of  $a$  and specific sediment yield is~~  
1056 ~~consistent with the previous hypothesis of increasing erosion within the basin as a~~  
1057 ~~consequence of the regime change.~~

#### 1058 Validity range of rating curves

1059 The estimation of sediment yield from flow measurements and a rating curve is still  
1060 acceptable throughout the study period (Fig. 53). However, it should be noted that the pairs  
1061 ( $C$ ,  $Q$ ) become increasingly scattered with time around the best-fit curve. The coefficient of  
1062 determination has decreased from one decade to another over 40 years, from 0.57 to 0.38  
1063 (Table 2). ~~As already explained, the rating curve established for only the last decade did not~~  
1064 ~~provide a reliable estimate of the solid discharge, leading to an error of 23% as compared to~~  
1065 ~~the calculation from measurements.~~

1066 Intermittent flows induce a stronger dependency of river behavior on antecedent wetness  
1067 (Beven, 2002) and antecedent weathering, i.e. a strong dependency on memory through  
1068 threshold and hysteresis effects. With increasing memory effects, coincident values of  $C$  and  
1069  $Q$  become less dependent on each other and the rating curves less suitable to model their  
1070 relation. The study of sediment dynamics in the Wadi Abd will thus likely require in the  
1071 future a more appropriate method than rating curves, such as the study of each individual  
1072 flood, like Megnounif et al. (2013) did in the Wadi Sebdou. This finding may have  
1073 consequences on water management as well. When dealing with rating curves, ~~water~~  
1074 ~~discharge must be recorded at frequent intervals, although measurements of concentration can~~  
1075 ~~be sparser.~~ When rating curves cannot be applied, river discharge and sediment concentration  
1076 should be both frequently and simultaneously measured.

## 1077 **8 Conclusions**

1078 Over the last 40 years, in response to climate change which resulted in an increase in  
1079 temperature of around 1.1°C between the 1970s and 2000s years, ~~rainfall moved forward~~  
1080 ~~during the late warm season and the watershed of Wadi Abd experienced a significant change~~

1081 in the flow regime of the river and an increased [variability at both the inter-annual and intra-](#)  
1082 [annual levels](#). These changes ultimately lead to a dramatic and continuous increase in  
1083 sediment load over 4 decades (in average 84% more every decade as compared to the  
1084 previous one).

1085 The main result of our analysis is the shift of the onset of the first summer flood that occurred  
1086 1 month earlier in the 2000s than in the 1970s. This shift is likely responsible for the  
1087 cascading effects on the hydrological regime of the Wadi Abd. In particular, earlier floods  
1088 during the warmer season have higher evaporation which limits the groundwater storage. A  
1089 parallel study of seasonal changes in vegetation cover is needed to provide additional  
1090 information.

1091 The increase in erosion of the watershed (coefficient a) is accompanied by a decrease in the  
1092 coefficient b ~~that seems to be associated with a decrease in the erosive power of the river. But~~  
1093 ~~this interpretation is still questionable since the sediment fluxes do not reflect erosion~~  
1094 ~~processes only, but also sediment storage (Trimble, 1999)~~. The traditional rating curves  
1095 approach which was applicable when the river was perennial is now less adapted to model the  
1096 behavior of the river (Table 2). This could be explained by a more pronounced hysteresis  
1097 phenomenon, which is consistent with the change of hydrological regime in the dry season  
1098 thereby limiting the utility of rating curves to model C-Q relationships. ~~Indeed, system~~  
1099 ~~memory related to hysteresis phenomena cannot be accounted for by instant models such as~~  
1100 ~~rating curves~~. Other methods such as that proposed by Megnounif et al. (2013) are probably  
1101 better adapted to understand future sediment dynamics of the Wadi Abd.

1102 ~~The rapid change in sediment regime which is instantaneously driven by the changing flow~~  
1103 ~~regime should be distinguished from the slow change in the concentration-flow relationship.~~  
1104 ~~The change in flow regime can be precisely dated in May-July 1986 (with 49 consecutive dry~~  
1105 ~~days), while the change in the C-Q relationship needs averaging over 15 years of a, b and~~  
1106 ~~specific sediment yield to become evident. Such inertial effect may be attributed to the time~~  
1107 ~~for the basin soil properties (such as humidity) or vegetation to adapt to the new climate~~  
1108 ~~conditions. It likely depends amongst other factors on the underground water storage, and~~  
1109 ~~thus on basin lithology and infiltration history. On the Wadi Abd basin, the time needed for~~  
1110 ~~the flow regime to change after the dryness settlement in early 1970's (see Fig. 6) is estimated~~  
1111 ~~around 15 years in this study.~~

1112 This present analysis only includes hydrological parameters. Management programs that  
1113 were conducted to fight erosion in Algeria from 1960s until 1990s by reforestation and setting  
1114 up banks over cultivated marl and clay areas proved to be little or no efficiency (Touaibia,  
1115 2010). Human activities may have influenced the hydrological regime change and increased  
1116 erosion, in particular through firewood cutting during economically difficult periods (1990s),  
1117 however the shift was shown to occur earlier. The lack of data on land use and land cover  
1118 changes over 40 years does not allow us to isolate the factors directly related to climate  
1119 change from those related to other anthropogenic activities, ~~but this question was not in the~~  
1120 ~~scope of the paper~~. However, the small population, the low coverage of pasture (see Fig. 2d),  
1121 of cultivated areas and vegetation (43 %) in the basin and the small volume of reservoirs  
1122 (nominally 2.3% of the annual discharge, ~~but~~ and silted up to 70%) make us think that in this  
1123 system the effects of climate change dominate anthropogenic effects. The quantification of  
1124 forcing changes on sediment sources (raindrop erosion, sheet erosion, rill erosion, gully  
1125 erosion, stream channel erosion) may be investigated in situ (e.g. Poesen et al., 2003) and/or  
1126 estimated using a numerical model of the hydrologic and sedimentological functioning of the  
1127 basin, such as WEPP (Nearing et al., 1989), EUROSEM (Morgan et al., 1998) or SWAT  
1128 (Neitsch et al., 2011). Such a model could help us to test hypothesis and quantify or at least  
1129 estimate the effects of different forcing changes (temperature, runoff, vegetation etc) in future  
1130 studies.

1131 It is important to emphasize that it is impossible to define long-term hydrological averages in  
1132 the context of a changing flow regime. Our analysis is based on the shift from a perennial  
1133 regime to an intermittent one. The example of the Wadi Abd shows that the difficulty is  
1134 challenging with regard to sediment transport in suspension, since changes of flux cannot be  
1135 counted as a fraction but can reach an order of magnitude.

1136 Changes in flow regime in relation to climate change can be investigated using climate  
1137 models. Das et al. (2013) using 16 climate projections showed that more intense floods of a  
1138 return period of 2-50 years should occur in the Sierra Nevada, regardless of the rainfall  
1139 variation. The recent changes in the Wadi Abd show that extreme events with increasing  
1140 variability already occur in the basin. Over Algeria, an increase of 1-2°C in temperature could  
1141 induce a reduction of 10% in precipitation before the end of the 21<sup>st</sup> century (Benhamiche et  
1142 al., 2014) with unknown consequences on erosion and sediment transport. Lu et al. (2013)  
1143 calculated the impact on sediment loads of every 1% change in precipitation or river  
1144 discharge in large Chinese rivers. Such a calculation has no meaning in our basin since the

1145 rainfall and discharge were not monotonic (severe decrease in the 1970s then slight increase  
1146 during 30 years) while the sediment loads have always increased. The difficulty of forecasting  
1147 climate change-driven impacts on sediment yield [due to](#) non linear effects has been  
1148 underlined by geomorphologists (see Goudie, 2006; Jerolmack and Paola, 2010; Coulthard et  
1149 al., 2012; Knight and Harrison, 2013). The present study illustrates that the change of flow  
1150 regime induce a fully non linear effect between river discharge and sediment yield. This needs  
1151 be considered in forecasts especially in small river basins in semi-arid areas.

1152 Changes in erosion and sediment transport under new climate constraints will induce changes  
1153 on the middle to long term that decision-makers must integrate into water resources  
1154 management, habitat status, agricultural adaptation (O’Neal et al., 2005), landscape evolution  
1155 (Temme and Veldkamp, 2009) as well as in many other environmental adaptations (Ouillon,  
1156 1998). We thus encourage the local adaptation of sampling strategies and measurements to  
1157 take into account changing in flow regimes. Furthermore, due to the uncertainty of water  
1158 resources and erosion [in the Maghreb](#) (Taabni and El Jihad, 2012) and in the Mediterranean  
1159 basin (Nunes et al., 2008), we also encourage the development of studies on long-term  
1160 sediment transport in North African basins, in connection with changes in forcing factors.

## 1161 **Acknowledgements**

1162 The authors warmly thank Miss Abda Leila from the Hydrometry office, Mr Ould Lamara  
1163 Arezki from the Climatology Office of the Agence Nationale des Ressources Hydrauliques  
1164 (ANRH) in Alger, Mr Boudalia Mohamed from the Hydrometry Office of the ANRH in Oran,  
1165 and all the staff of ANRH who participated to the field measurements at Ain Hamara station.  
1166 They also acknowledge Mr Abderrezak Kamel Toubal for his help in drawing [Figures 1 and](#)  
1167 [2. Two anonymous reviewers are warmly thanked for their reviews and comments on](#)  
1168 [previous versions of this paper. The editor, Efrat Morin, is gratefully acknowledged.](#) Emma  
1169 Rochelle-Newall, a native English speaker, is warmly thanked for English corrections.

1170

1171 **References**

- 1172 Aalto, R., Dunne, T., and Guyot, J.L.: Geomorphic controls on Andean denudation rates, The  
1173 Journal of Geology, 114, 85-99, 2006.
- 1174 Achite, M., and Ouillon, S.: Suspended sediment transport in a semiarid watershed, Wadi  
1175 Abd, Algeria (1973-1995), J. Hydrol., 343 (3-4), 187-202, 2007.
- 1176 Alexandrov, Y., Laronne, J.B., and Reid, I.: Suspended sediment concentration and its  
1177 variation with water discharge in a dryland ephemeral channel, northern Negev, Israel,  
1178 J. Arid Envir., 53, 73-84, 2003.
- 1179 Asselman, N.E.M.: Fitting and interpretation of sediment rating curves, J. Hydrol., 234, 228-  
1180 248, 2000.
- 1181 Bakreti, A., Braud, I., Leblois, E., and Benali, A.: Analyse conjointe des régimes  
1182 pluviométriques et hydrologiques dans le bassin de la Tafna (Algérie Occidentale),  
1183 Hydrol. Sci. J., 58 (1), 133-151, 2013.
- 1184 Bates, B.C., Kundzewicz Z.W., Wu S., and Palutikof J.P., Eds.: Climate Change and Water.  
1185 Technical Paper of the Intergovernmental Panel on Climate Change, IPCC Secretariat,  
1186 Geneva, 210 pp., 2008
- 1187 Benhamiche, N., Madani, K., and Laignel, B.: Chapter 12 Impact of climate change on water  
1188 resources in Algeria, in Behnassi M. et al. (eds), *Vulnerability of agriculture, water and*  
1189 *fisheries to climate change: toward sustainable adaptation strategies*, Springer, 193-  
1190 205, doi:10.1007/978-94-017-8962-2\_12, 2014.
- 1191 Beven, K.: Runoff generation in semi-arid areas. In Bull L.J., Kirkby M.J. (ed), 2002. Dryland  
1192 rivers: hydrology and geomorphology of semi-arid channels, Wiley, London, pp 57-  
1193 106, 2002.
- 1194 Coulthard, T.J., Ramirez, J., Fowler, H.J., and Glenis, V.: Using the UKCP09 probabilistic  
1195 scenarios to model the amplified impact of climate change on river basin sediment  
1196 yield, Hydrol. Earth Syst. Sci., 16, 4401-4416, 2012.
- 1197 Dahmani, A., and Meddi, M.: Climate variability and its impact on water resources in the  
1198 catchment area of the Wadi Fekan Wilaya of Mascara (West Algeria), Eur. J. Sci. Res.,  
1199 36 (3), 458-472, 2009.
- 1200 Das, T., Maurer, E.P., Pierce, D.W., Dettinger, M.D., and Cayan, D.R.: Increases in flood  
1201 magnitudes in California under warming climates, J. Hydrol., 501, 101-110, 2013.



1202 Déry, S.J., Stieglitz, M., McKenna, E.C., and Wood, E.F.: Characteristics and trends of river  
1203 discharge into Hudson, James, and Ungava bays, 1964–2000, *J. Climate*, 18, 2540–  
1204 2557, 2005.

1205 Durand, N., Fiandrino, A., Fraunie, P., Ouillon, S., Forget, P., and Naudin, J.J.: Suspended  
1206 matter dispersion in the Ebro ROFI: an integrated approach, *Cont. Shelf Res.*, 22, 267–  
1207 284, 2002.

1208 El Mahi, A., Meddi, M., and Bravard, J.P.: Analyse du transport solide en suspension dans le  
1209 bassin versant de l'Oued El Hammam (Algérie du Nord), *Hydrol. Sci. J.*, 57 (8), 1642–  
1210 1661, 2012.

1211 Gao, Z.L., Fu, Y.L., Li, Y.H., Liu, J.X., Chen, N., and Zhang, X.P.: Trends of streamflow,  
1212 sediment load and their dynamic relation for the catchments in the middle reaches of the  
1213 Yellow River over the past five decades, *Hydrol. Earth Syst. Sciences*, 16, 3219–3231,  
1214 2012.

1215 Ghenim, A.N., and Megnounif, A.: Analyse des précipitations dans le Nord-Ouest algérien,  
1216 *Sécheresse*, 24, 107–114, 2013a.

1217 Ghenim, A.N., and Megnounif, A.: Ampleur de la sécheresse dans le bassin d'alimentation du  
1218 barrage Meffrouche (Nord-Ouest de l'Algérie), *Physio-Géo*, 7, 35–49, 2013b.

1219 [GISTEMP Team: GISS Surface Temperature Analysis \(GISTEMP\). NASA Goddard Institute](#)  
1220 [for Space Studies. Dataset accessed 2015-12-08 at http://data.giss.nasa.gov/gistemp/](#)

1221 Giuntoli, I., Renard, B., Vidal, J.P., and Bard, A.: Low flows in France and their relationship  
1222 to large-scale climate indices, *J. Hydrol.*, 482, 105–118, 2013.

1223 Gomez, B., Cui, Y., Kettner, A.J., Peacock, D.H., and Syvitski, J.P.M.: Simulating changes to  
1224 the sediment transport regime of the Waipaoa River, New Zealand, driven by climate  
1225 change in the twenty-first century, *Global Planet. Change*, 67, 153–166, 2009.

1226 Goudie, A.S.: Global warming and fluvial geomorphology, *Geomorphology*, 79, 384–394,  
1227 2006.

1228 Hallouz, F., Meddi, M., and Mahé, G.: Modification du régime hydroclimatique dans le bassin  
1229 de l'oued Mina (Nord-Ouest d'Algérie), *Revue Sci. Eau*, 26 (1), 33–38, 2013.

1230 Hancock, G.R.: A catchment scale assessment of increased rainfall and storm intensity on  
1231 erosion and sediment transport for Northern Australia, *Geoderma*, 152, 350–360, 2009.

1232 Hancock, G.R., and Coulthard, T.J.: Channel movement and erosion response to rainfall  
1233 variability in southeast Australia, *Hydrol. Process.*, 26, 663–673, 2011.

1234 [Hansen, J., Ruedy, R., Sato, M., and Lo, K.: Global surface temperature change, \*Rev.\*](#)  
1235 [Geophysics](#), 48, RG4004, doi:10.1029/2010RG000345, 2010.

1236 Houyou Z., Biolders, C.L., Benhorma, H.A., Dellal A., and Boutemdjet, A.: Evidence of  
1237 strong land degradation by wind erosion as a result of rainfed cropping in the Algerian  
1238 steppe: a case study at Laghouat, *Land Degrad. Develop.*, doi:10.1002/ldr.2295, 2014.

1239 Iadanza, C., and Napolitano, F.: Sediment transport time series in the Tiber River, *Physics*  
1240 *Chem. Earth*, 31, 1212-1227, 2006.

1241 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I  
1242 to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change;  
1243 Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y.  
1244 Xia, V. Bex and P.M. Midgley (eds.). Cambridge University Press, Cambridge, United  
1245 Kingdom and New York, NY, USA, 1535 pp., 2013.

1246 Jerolmack, D.J., and Paola, C.: Shredding of environmental signals by sediment transport,  
1247 *Geophys. Res. Lett.*, 37, L19401, doi:10.1029/2010GL044638, 2010.

1248 Jones, P.D., Lister, D.H., Osborn, T.J., Harpham, C., Salmon, M., and Morice, C.P.:  
1249 Hemispheric and large-scale land surface air temperature variations: an extensive  
1250 revision and an update to 2010. *J. Geophys. Res.*, 117, D05127. doi:  
1251 [10.1029/2011JD017139](https://doi.org/10.1029/2011JD017139), 2012.

1252 Kassoul, M., Abdelgader, A., and Belorgey, M.: Caractérisation de la sédimentation des  
1253 barrages en Algérie. *Rev. Sci. Eau*, 3, 339-358, 1997.

1254 Knight, J., and Harrison, S.: The impacts of climate change on terrestrial Earth surface  
1255 systems, *Nature Climate Change*, 3, 24-29, 2013.

1256 Langbein, W.B., and Schumm, S.A.: Yield of sediment in relation to mean annual  
1257 precipitation, *Trans. Amer. Geophys. Union*, 39, 1076-1084, 1958.

1258 Lockwood, M., and Fröhlich, C.: Recent oppositely directed trends in solar climate forcings  
1259 and the global mean surface air temperature, *Proc. R. Soc. A*,  
1260 doi:10.1098/rspa.2007.1880, 2007.

1261 Louamri, A., Mebarki, A., and Laignel, B.: Variabilité interannuelle et intra-annuelle des  
1262 transports solides de l'Oued Bouhamdane, à l'amont du barrage Hammam Debagh  
1263 (Algérie orientale), *Hydrol. Sci. J.*, 58:7, 1559-1572, 2013.

1264 Lu, X.X., Ran, L.S., Liu, S., Jiang, T., Zhang, S.R., and Wang, J.J.: Sediment loads response  
1265 to climate change: A preliminary study of eight large Chinese rivers, *Int. J. Sed. Res.*,  
1266 28, 1-14, 2013.

1267 Mahé, G., and Paturel, J.E.: 1896-2006 Sahelian annual rainfall variability and runoff increase  
1268 of Sahelain Rivers, *C.R. Geoscience*, 314, 538-546, 2009.

- 1269 Mahé, G., Lienou, G., Descroix, L., Bamba, F., Paturol, J.E., Laraque, A., Meddi, M.,  
 1270 Habaieb, H., Adeaga, O., Dieulin, C., Chahnez Kotti, F., and Khomsi, K.: The rivers of  
 1271 Africa: witness of climate change and human impact on the environment, *Hydrol. Proc.*,  
 1272 27, 2105-2114, 2012.
- 1273 Meade, R.H., and Moody, J.A.: Causes for the decline of suspended-sediment discharge in the  
 1274 Mississippi River system, 1940-2007, *Hydrol. Proc.*, 24, 35-49, 2010.
- 1275 Meddi, M., and Hubert, P.: Impact de la modification du régime pluviométrique sur les  
 1276 ressources en eau du nord-ouest de l'Algérie, in: *Hydrology of the Mediterranean and*  
 1277 *Semiarid Regions*, IAHS Publ. n°278, 229-235, 2003.
- 1278 Meddi, H., and Meddi, M.: Variabilité spatiale et temporelle des précipitations du Nord-Ouest  
 1279 de l'Algérie, *Geographica Technica* n°2/2007, 49-55, 2007.
- 1280 Meddi, H., and Meddi, M.: Variabilité des précipitations annuelles du Nord-Ouest de  
 1281 l'Algérie, *Sécheresse*, 20 (1), 57-65, 2009.
- 1282 Meddi, M., and Talia, A.: Pluviometric regime evolution in the North of Algeria. *Arab Gulf J.*  
 1283 *Sci. Res.*, 26 (3), 152-162, 2008.
- 1284 Meddi, M., Talia, A., and Martin, C.: Evolution récente des conditions climatiques et des  
 1285 écoulements sur le bassin versant de la Macta (nord-ouest de l'Algérie), *Physio-Géo*, 3,  
 1286 61-84, 2009.
- 1287 Megnounif, A., Terfous, A., and Ouillon, S.: A graphical method to study suspended sediment  
 1288 dynamics during flood events in the Wadi Sebdo, NW Algeria (1973-2004), *J. Hydrol.*,  
 1289 497, 24-36, 2013.
- 1290 Memariam, H., Balasundram, S.K., Talib, J.B., Sood, A.M., and Abbaspour, K.C.: Trend  
 1291 analysis of water discharge and sediment load during the past three decades of  
 1292 development in the Langat basin, Malaysia, *Hydrol. Sci. J.*, 57 (6), 1207-1222, 2012.
- 1293 Milliman, J.D., Farnsworth, K.L., Jones, P.D., Xu, K.H., and Smith, L.C.: Climatic and  
 1294 anthropogenic factors affecting river discharge to the global ocean, 1951-2000, *Global*  
 1295 *Planet. Change*, 62, 187-194, 2008.
- 1296 [Morgan R.P.C., Quinton, J.N., Smith R.E., Govers, G., Poesen, J.W.A., Auerswald, K., Chisci](#)  
 1297 [G., Torri, D., and Styczen, M.E.: The European Soil Erosion Model \(EUROSEM\): A](#)  
 1298 [dynamic approach for predicting sediment transport from fields and small catchments,](#)  
 1299 [Earth Surf. Process. Landforms](#), 23, 527-544, 1998.
- 1300 Nanson, G.C., Tooth, S., and Knighton, A.D.: A global perspective on dryland rivers:  
 1301 perceptions, misconceptions and distinctions. In Bull L.J., Kirkby M.J. (ed), 2002.

1302 Dryland rivers: hydrology and geomorphology of semi-arid channels, Wiley, London.  
 1303 pp 17-54, 2002.

1304 [Neitsch, S.L., Arnold, J.G., Kiniry J.R., and Williams, J.R.: Soil and Water Assessment Tool](#)  
 1305 [– Theoretical Documentation version 2009. Texas Water Inst. Techn. Report n°406,](#)  
 1306 [Texas A&M University, College Station, 2011.](#)

1307 Nunes, J.P., Seixas, J., and Pacheco, N.R.: Vulnerability of water resources, vegetation  
 1308 productivity and soil erosion to climate change in Mediterranean watersheds, *Hydrol.*  
 1309 *Process.*, 22, 3115–3134, 2008.

1310 Oliver, J.E.: Monthly precipitation distribution: a comparative index, *Prof. Geogr.*, 32 (3),  
 1311 300–309, 1980.

1312 O’Neal, M.R., Nearing, M.A., Vining, R.C., Southworth, J., and Pfeifer, R.A.: Climate  
 1313 change impacts on soil erosion in Midwest United States with changes in crop  
 1314 management, *Catena*, 61, 165–184, 2005.

1315 Osborn, T.J., and Jones, P.D.: The CRUTEM4 land-surface air temperature dataset:  
 1316 construction, previous versions and dissemination via Google Earth, *Earth System*  
 1317 *Science Data* 6, 61-68, 2014.

1318 Ouillon, S.: Erosion and sediment transport: width and stakes, *La Houille Blanche*, 53 (2), 52-  
 1319 58, 1998.

1320 Paturel, J.E., Ouedraogo, M., Servat, E., Mahé, G., Dezetter, A., and Boyer, J.F.: The concept  
 1321 of rainfall and stream flow normals in West and Central Africa in a context of climatic  
 1322 variability, *Hydrol. Sci. J.*, 48 (1), 125-137, 2003.

1323 [Poesen J., Nachtergaele, J., Verstraeten, G., and Valentin, C.: Gully erosion and](#)  
 1324 [environmental change: importance and research needs, \*Catena\*, 50, 91-133, 2003.](#)

1325 Pruski, F.F., and Nearing, M.A.: Climate-induced changes in erosion during the 21st century  
 1326 for eight U.S. locations, *Water Resour. Res.*, 38, 1298, doi:10.1029/2001WR000493,  
 1327 2002.

1328 [Remini B., Leduc C., and Hallouche, W.: Evolution des grands barrages en régions arides:](#)  
 1329 [quelques exemples algériens, \*Sécheresse\*, 20 \(1\), 96-103, 2009.](#)

1330 Riebe, C.S., Kirchner, J.W., Granger, D.E., and Finkel, R.C.: Minimal climatic control on  
 1331 erosion rates in the Sierra Nevada, California, *Geology*, 29, 447-450, 2001.

1332 Sari, A.: Hydrologie de surface, cours d’initiation, Editions Distribution Houma, 155 pp,  
 1333 2009.

1334 Searcy, J.K., and Hardison, C.H.: Double-mass curves, USGS Water-Supply Paper 1541-B,  
 1335 1960.

- 1336 Sen, P. K.: Estimates of regression coefficients based on Kendall's tau, *J. Am. Stat. Assoc.*,  
 1337 63, 1379–1389, 1968.
- 1338 Shrestha, B., Babel, M.S., Maskey, S., van Griensven, A., Uhlenbrook, S., Green, A., and  
 1339 Akkharath, I. Impact of climate change on sediment yield in the Mekong River basin: a  
 1340 case study of the Nam Ou basin, Lao PDR, *Hydrol. Earth System Sciences*, 17, 1-20,  
 1341 2013.
- 1342 Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L.M., van Lanen, H.A.J., Sauquet, E., Demuth,  
 1343 S., Fendekova, M., and Jodar, J.: Streamflow trends in Europe: Evidence from a dataset  
 1344 of near-natural catchments, *Hydrol. Earth Syst. Sci.*, 14, 2367–2382, 2010.
- 1345 [Summerfield, M.A., Hulton, N.J.: Natural controls of fluvial denudation rates in major world  
 1346 drainage basins, \*J. Geophys. Res.\*, 99 B7, 13871-13883, 1994.](#)
- 1347 Syvitski, J.P.M.: Supply and flux of sediment along hydrological pathways: research for the  
 1348 21<sup>st</sup> century, *Global Planet. Change*, 39, 1-11, 2003.
- 1349 Syvitski, J.P.M., and Milliman, J.D.: Geology, geography, and humans battle for dominance  
 1350 over the delivery of fluvial sediment to the coastal ocean, *The Journal of Geology*, 115,  
 1351 1-19, 2007.
- 1352 Taabni, M., and El Jihad, M.D.: Eau et changement climatique au Maghreb: quelles stratégies  
 1353 d'adaptation?, *Les Cahiers d'Outre-Mer*, 260, 493-518, 2012.
- 1354 Taibi, S., Meddi, M., Mahé, G., and Assani, A.: Variability of annual and extreme rainfall  
 1355 over Northern Algeria and relationship with teleconnections patterns, *Proc. of the  
 1356 Mediterranean meeting on "Monitoring, modeling and early warning of extreme events  
 1357 triggered by heavy rainfalls"*, Univ. of Calabria, Cosenza, Italy, June 26-28, 2014.
- 1358 Tebbi, F.Z., Dridi, H., and Morris, G.L.: Optimization of cumulative trapped sediment curve  
 1359 for an arid zone reservoir: Foum El Kherza (Biskra, Algeria), *Hydrol. Sci. J.*, 57 (7),  
 1360 1368-1377, 2012.
- 1361 Temme, A.J.A.M., and Veldkamp, A.: Multi-process Late Quaternary landscape evolution  
 1362 modelling reveals lags in climate response over small spatial scales, *Earth Surf. Proc.  
 1363 Land.*, 34, 573–589, 2009.
- 1364 Tecsalt International: Etude de la protection des bassins versants de l'ensemble des barrages  
 1365 en études d'avant projet détaillé (lots n°1 à 4) et des bassins versants des barrages en  
 1366 exploitation (lots n° 5 et 6), Phase I description du milieu, lot n°5: barrages Gargar et  
 1367 S.M. Ben-Aouda, Ministère des Ressources en Eau, Algérie, 2004.
- 1368 Tockner, K., and Stanford, J.A.: Riverine flood plains: present state and future trends,  
 1369 *Environ. Conserv.*, 29, 308–330, 2002.

1370 Touaibia, B.: Problématique de l'érosion et du transport solide en Algérie septentrionale,  
1371 Sécheresse, 21 (1e), 1-6, 2010.

1372 Tramblay, Y., El Adlouni, S., and Servat, E.: Trends and variability in extreme precipitation  
1373 indices over Maghreb countries. *Natural Hazards Earth Syst. Sci.*, 13, 3235-3248, 2013.

1374 ~~Trimble, S.W.: Decreased rates of alluvial sediment storage in the Coon Creek basin,  
1375 Wisconsin, 1975-93, *Science*, 285, 1244-1246, 1999.~~

1376 Tucker, G.E., and Slingerland, R.: Drainage basin responses to climate change, *Water Resour.*  
1377 *Res.*, 33, 2031-2047, 1997.

1378 Vanmaercke, M., Poersen, J., Broeckx, J., and Nyssen, J.: Sediment yield in Africa, *Earth-*  
1379 *Science Rev.*, 136, 350-368, 2014.

1380 Vinh, V.D., Ouillon, S., Thanh, T.D., and Chu, L.V.: Impact of the Hoa Binh dam (Vietnam)  
1381 on water and sediment budgets in the Red River basin and delta, *Hydrol. Earth System*  
1382 *Science*, 18, 3987-4005, 2014.

1383 Vörösmarty, C.J., Meybeck, M., Fekete, B., Sharma, K., Green, P., and Syvitski, J.P.M.:  
1384 Anthropogenic sediment retention: major global impact from registered river  
1385 impoundments, *Global Planet. Change*, 39, 169-190, 2003.

1386 Walling, D.E.: Assessing the accuracy of suspended sediment rating curves for a small basin,  
1387 *Water Res. Res.*, 13 (3), 531-538, 1977.

1388 Walling, D.E.: The response of sediment yields to environmental change, *IAHS Publ.* 245,  
1389 77-89, 1997.

1390 Walling, D.E.: Human impact on land-ocean sediment transfer by the world's rivers,  
1391 *Geomorphology*, 79, 192-216, 2006.

1392 Walling, D.E.: The impact of Global Change on Erosion and Sediment Transport by Rivers:  
1393 Current progress and Future Challenges, *The United Nations World Water*  
1394 *Development Report 3*, UNESCO, Paris, 26 p., 2009.

1395 Walling, D.E., and Fang, D.: Recent trends in the suspended sediment loads of the world's  
1396 rivers. *Global Planet. Change*, 39, 111-126, 2003.

1397 Wang, H., Yang, Z., Saito, Y., Liu, J., and Sun, X.: Interannual and seasonal variation of the  
1398 Huanghe (Yellow River) water discharge over the past 50 years: connections to impacts  
1399 from ENSO events and dams, *Global Planet. Change*, 50, 212-225, 2006.

1400 Wang, H.J., Yang, Z.S., Saito, Y., Liu, J.P., Sun, X.X., and Wang, Y.: Stepwise decreases of  
1401 the Huanghe (Yellow River) sediment load (1950-2005): Impacts of climate change and  
1402 human activities, *Global Planet. Change*, 57 (3-4), 331-354, 2007.

- 1403 ~~Wang, H.J., Yang, Z.S., Wang, Y., Saito, Y., and Liu, J.P.: Reconstruction of sediment flux~~  
1404 ~~from the Changjiang (Yangtze River) to the sea since the 1860s, J. Hydrology, 349,~~  
1405 ~~318-332, 2008.~~
- 1406 Ward, P.J., Eisner, S., Flörke, M., Dettinger, M.D., and Kummu, M.: Annual flood  
1407 sensitivities to El Niño-Southern Oscillation at the global scale, Hydrol. Earth Syst. Sci.,  
1408 18, 47-66, 2014.
- 1409 Ziegler, A.D., Sheffield, J., Maurer, E.P., Nijssen, B., Wood, E.F., and Lettenmaier, D.P.:  
1410 Detection of intensification in global- and continental-scale hydrological cycles:  
1411 Temporal scale of evaluation. J. Climate, 16, 535–547, 2003.

1412 **Table 1.** General statistics of the yearly averages of hydrologic parameters of the Wadi Abd  
 1413 at Ain Hamara gauging station over 1970-2010 (Note: T at Chlef was estimated from  
 1414 measurements at Dar El Beida and Miliana for 20 months over 480)

1415

statistic value	T (Chlef) °C	P mm yr <sup>-1</sup>	Q m <sup>3</sup> s <sup>-1</sup>	Q <sub>w</sub> m <sup>3</sup> s <sup>-1</sup>	M 10 <sup>3</sup> t yr <sup>-1</sup>	SPM* g L <sup>-1</sup>
Mean	19.09	264	1.18	1.29	564	12.3
Min	17.52	165	0.37	0.46	33.1	2.56
(Year)	1971-72	1999-00	1992-93	1983-84	1992-93	1975-76
Max	20.32	506	2.19	2.98	3266	50.25
(Year)	1989-90	1995-96	1994-95	1994-95	2007-08	2007-08
standard deviation	0.69	71.2	0.52	0.59	696	10.6
CV (%)		27.0	44.4	45.6	123.3	86.0

1416

1417



**Table 2.** General statistics of the averages of hydrologic parameters of the Wadi Abd at Ain Hamara gauging station per decade and significant period over 1970-2010 (Note: T at Chlef was estimated from measurements at Dar El Beida and/or Miliana for 20 months with missing values over 480)

Period	T at Chlef	NRD, average yearly number of rainy days	P, yearly precipitation		NDD, average yearly number of dry days (Q=0)	Q, yearly discharge		Q <sub>w</sub> , yearly discharge of wet days		Q <sub>s</sub> , yearly sediment load		Q <sub>98</sub> , average of yearly values		SSY, average specific sed. yield (t km <sup>-2</sup> yr <sup>-1</sup> )	SPM*		Rating curve parameters			
	Average (°C)		Average (mm)	CV (%)		Average (m <sup>3</sup> s <sup>-1</sup> )	CV (%)	Average (m <sup>3</sup> s <sup>-1</sup> )	CV (%)	Average (10 <sup>3</sup> tons yr <sup>-1</sup> )	CV (%)	Average (m <sup>3</sup> s <sup>-1</sup> )	CV (%)		Average (g L <sup>-1</sup> )	CV (%)	a	b	R <sup>2</sup>	N
1970-2010	19.09		264.10	27.0	28.3	1.18	44.4	1.29	45.7	564	123.3	9.18	78.6	227.6	12.3	86.0	2.270	0.647	0.431	1213
1970-1980	18.32		310.53	19.4	1.2	1.16	32.9	1.16	32.9	180	78.8	4.37	66.9	72.7	4.54	47.9	1.021	0.890	0.573	240
1980-1990	19.19		231.23	16.8	24.1	0.98	36.8	1.07	41.5	334	91.7	7.39	68.0	134.5	9.93	57.0	2.049	0.649	0.449	316
1990-2000	19.37		250.42	40.5	59.9	1.13	55.1	1.34	55.2	614	98.3	11.03	88.5	247.5	14.36	69.2	2.753	0.659	0.418	343
2000-2010	19.49		264.22	19.7	28.1	1.45	43.3	1.57	42.2	1130	90.3	13.94	44.5	455.6	20.55	68.7	4.440	0.412	0.384	324
1970-1985	18.51		284.34	23.1	0.8	1.02	37.8	1.02	38.2	159	78.9	4.13	58.8	64.2	5.16	58.9	1.213	0.818	0.519	346
1985-2010	19.47		251.96	29.0	44.8	1.28	45.1	1.45	43.7	808	97.0	12.21	61.1	325.6	16.65	67.4	2.974	0.576	0.415	867

1 **Table 3.** Variation of precipitation, water discharge and sediment **yield** averaged per season  
 2 over each decade  
 3

	Precipitation (mm)				Water discharge (m <sup>3</sup> s <sup>-1</sup> )				Sediment <b>yield</b> (10 <sup>3</sup> tons)			
	autumn	winter	spring	summer	autumn	winter	spring	summer	autumn	winter	spring	summer
1970-1980	68.5	102.6	128.2	11.2	3.79	4.15	4.22	1.75	62.2	43.7	66.1	8.4
1980-1990	56.0	94.4	70.7	10.1	3.45	3.86	3.25	1.19	128.8	61.0	97.2	46.8
1990-2000	67.0	81.1	86.9	15.5	5.58	2.98	3.33	1.62	279.1	57.8	130.9	146.0
2000-2010	78.6	98.4	77.7	9.5	9.13	4.05	3.18	1.05	804.9	94.4	195.3	35.4

	Precipitation (%)				Water discharge (%)				Sediment <b>yield</b> (%)			
	autumn	winter	spring	summer	autumn	winter	spring	summer	autumn	winter	spring	summer
1970-1980	22.1	33.0	41.3	3.6	27.3	29.8	30.3	12.6	34.5	24.2	36.6	4.7
1980-1990	24.2	40.8	30.6	4.4	29.4	32.8	27.7	10.1	38.6	18.3	29.1	14.0
1990-2000	26.7	32.4	34.7	6.2	41.3	22.1	24.6	12.0	45.5	9.4	21.3	23.8
2000-2010	29.7	37.3	29.4	3.6	52.5	23.2	18.2	6.1	71.2	8.4	17.3	3.1

4

## Figure Captions

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

**Fig. 1.** Location of the Wadi Abd sub-basin within the Mina and Cheliff basins, and the other main basins of Algeria

**Fig. 2.** The Wadi Abd catchment area. (a) Rain and hydrometric stations including HS1 at Takhmaret and HS2 at Ain Hamara, (b) Geology, (c) Slopes from the Digital Elevation Model of North Algeria, (d) Vegetation cover from Landsat ETM+ data of 2009

**Fig. 3.** Linear erosion forms in the Wadi Abd basin. (a) and (e) Gullying (depth: 30-50 cm, width < 1 m), (c) and (d) Gully erosion (depth: 50-200 cm), (b) and (f) Interrill and rill erosion

**Fig. 42** Relationships between mean annual temperatures at the three stations of Dar El Beida, Miliana and Chlef (from CRUTEM4)

**Fig. 53** Comparison between estimates of  $Q_s$  obtained from  $Q$  and the global rating curve, and measured  $Q_s$

**Fig. 64** Interannual variations of mean yearly temperature (calculated from September to August monthly temperatures) at three stations in northern Algeria: Dar El Beida, Miliana, Chlef (from measurements of CRUTEM4 only, extrapolated values are not shown)

1 **Fig. 75.** Interannual variations of annual precipitation, water discharge and sediment yield at  
2 Ain Hamara station  
3  
4 **Fig. 86.** Variation of hydrological regime with annual % of time of flowing water,  $Q_{98}$   
5 (amongst daily discharges, per year) and annual standard deviation of daily river  
6 discharge  
7  
8 **Fig. 97** Double mass plot of sediment yield versus water flow  
9  
10 **Fig. 108** Yearly average of related sediment load parameters vs intra-annual variability of  
11 daily river discharge, characterized by their annual standard deviation. (a) SPM\*, (b)  
12 Specific Sediment Yield  
13  
14 **Fig. 119** Trends of the seasonal indexes of precipitation (a), discharge (b) and (c) sediment  
15 discharge in the Wadi Abd basin.  
16  
17 **Fig. 1210** Monthly values of precipitation (a), Q (b) and Qs (c) averaged over decades in the  
18 Wadi Abd basin.  
19  
20 **Fig. 1311** Monthly values of precipitation averaged over 6 stations, for the two periods 1970-  
21 1985 and 1985-2010.  
22  
23 **Fig. 1412** Monthly values of the ratio Q/P averaged over decades  
24

1 **Fig. 1513** Variations of SPM\* against the global mean temperature anomaly (from [GISTEMP](#)  
2 [Team, 2015](#))

3

4 ~~**Fig. 14** Relationship between yearly values of the rating curve parameters (a, b)~~

5

6 **Fig. 165** Relationship between the rating curves parameters averaged over 15 years

7

8 **Fig. 176** Relationship between the rating curve parameters  $a$  averaged over 15 years and the  
9 averaged values of ~~specific sediment yield or river discharge~~ over 15 years