

1 **Editor Decision: Reconsider after major revisions** (29 Feb 2016) by Efrat Morin

2 Comments to the Author:

3 I have read the revised paper, and examined the author response to the two reviewers
4 comments. Although the paper was improved it still needs more revisions.

5 General answer: We thank the reviewer#1 who agreed that the paper can be now accepted for
6 publication and whose help was really important to improve the paper from its initial version.
7 We also warmly thank the editor for his support to make the paper more concise and easier to
8 read. Detailed answers are given point by point below.

9 Additional corrections: Please note that the third column of the former Table 2 was deleted
10 since it was empty, and that the 16 seasonal averages of water discharge (in $\text{m}^3 \text{s}^{-1}$) were
11 corrected in Table 3 (all of them had to be divided by 3).

12 1) In the first round, reviewer #1 pointed out that the paper is too long and "heavy". I
13 am afraid the revised version still has this problem. The authors should put more efforts
14 in make the paper much more concise. See the comment below in this regard.

15 Detailed answer: Paragraphs were shortened all along the paper, as suggested. Section 4.4 was
16 deleted. Globally, the paper was reduced by 5.8%, from 8734 to 8230 words (sections 1 to 8).

17 2) Often, the authors describe in details the changes from decade to decade of different
18 variables, which is a repeat of the data presented for example in Table 2. I do not see
19 the point of this, as all info is in the table. The authors should concisely point only to
20 important features. This will help shortening the paper.

21 Detailed answer: Done, thank you for the suggestion. Paragraphs were shortened, especially
22 from sections 4 to 8 (conclusion). See below the details on the marked manuscript.

23 3) In view of one of reviewer #2 comments, I propose to add information on the
24 sediment load measurement technique, its reliability, and some details on what is
25 actually measured.

26 Detailed answer: More details are given in section 2.2. The former description:
27 “Concentrations derived from water samples taken at one or two points, after filtration on pre-
28 weighed Whatman Glass Fibre Filters (GFF) filters, oven-dried and weighed again following
29 the protocol described by A02007 and Megnounif et al. (2013).” was replaced by “Water
30 samples were filtered on pre-weighed Whatman Glass Fiber Filters (GFF), oven-dried at
31 105°C for 24h, and weighed again to determine the concentration. This method, used by
32 ANRH at all hydrologic stations in Algeria, underestimates the suspended load as compared

33 to its value averaged over the cross section under low turbulence (i.e. at low flow) since water
 34 is sampling near the surface (Touat, 1989). During floods, which transport most of the
 35 sediment load, turbulence is high enough to homogenize suspension load. While this
 36 underestimation may slightly affect the budget, it doesn't severely affect the time variability
 37 of suspended matter which is analyzed in this paper." The reference Touat (1989) was also
 38 added.

39 4) I would like to ask the authors to confirm the significance of all their statistical tests
 40 (p-value is always reported as lower than the selected significance level, i.e. all trends
 41 are significant). I cannot be sure, but it seems that the signal of the trend of precipitation
 42 for example presented in figure 7 is much smaller than the "noise", i.e., precipitation
 43 variability. Please recheck the p-value for this and other statistical tests presented.

44 Detailed answer: The trends and the p-values were double-checked and confirmed.

45 5) L 107 - add "potential" before evaporation.

46 Detailed answer: Done, thank you.

47 6) Some language editing is needed.

Detailed answer: Sorry for our mistakes. The paper was corrected by one English-native
 colleague. All the corrections are reported below.

Report #1

| | | | | |
|--|-------------------|---------------------|-----------|-----------------------|
| Submitted | on | 19 | Feb | 2016 |
| Anonymous Referee #1 | | | | |
| Anonymous during peer-review: Yes No | | | | |
| Anonymous in acknowledgements of published article: Yes No | | | | |
| Recommendation to the Editor | | | | |
| 1) | Scientific | Significance | Excellent | Good Fair Poor |
| Does the manuscript represent a substantial contribution to scientific progress within the scope of this journal (substantial new concepts, ideas, methods, or data)? | | | | |
| 2) | Scientific | Quality | Excellent | Good Fair Poor |
| Are the scientific approach and applied methods valid? Are the results discussed in an appropriate and balanced way (consideration of related work, including appropriate references)? | | | | |

3) Presentation Quality Excellent **Good** Fair Poor
Are the scientific results and conclusions presented in a clear, concise, and well structured way (number and quality of figures/tables, appropriate use of English language)?

For final publication, the manuscript should be

accepted as is

accepted subject to **technical corrections**

accepted subject to **minor revisions**

reconsidered after **major revisions**

I would like to review the revised paper

I would NOT be willing to review the revised paper

rejected

Please note that this rating only refers to this version of the manuscript!

Suggestions for revision or reasons for rejection (will be published if the paper is accepted for final publication)

48

49

50

Marked-up manuscript version

51 **Recent changes in climate, hydrology and sediment load in**
52 **the Wadi Abd, Algeria (1970-2010)**

53

54 **Mohammed Achite¹, Sylvain Ouillon^{2,3}**

55 [1]{Laboratoire Eau-Environnement, Université Hassiba Ben Bouali, BP 151, Hay Es-Salem,
56 Chlef, Algeria}

57 [2]{Université de Toulouse, UPS (OMP), IRD, UMR 5566 LEGOS, 14 avenue E. Belin,
58 31400 Toulouse, France}

59 [3]{University of Science and Technology of Hanoi, Dept. Water-Environment-
60 Oceanography, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam}

61 Correspondence to: Sylvain Ouillon, (sylvain.ouillon@legos.obs-mip.fr)

62

63 **Abstract**

64 Here we investigate the changes of temperature, precipitation, river runoff and sediment
65 transport in the Wadi Abd in NW Algeria over a time series of 40 hydrological years (1970-
66 2010). Temperature increased and precipitation decreased with the reduction in rainfall being
67 relatively higher during the rainy season. A shift towards an earlier onset of first rains during
68 summer was also found with cascading effects on hydrology (hydrological regimes,
69 vegetation, etc.) and thus on erosion and sediment yield. During the 1980s, the flow regime
70 shifted from perennial to intermittent with an amplification of the variations of discharge and
71 a modification of the sediment regime with higher and more irregular suspended particulate
72 flux. Sediment flux was shown to almost double every decade from 1970s to 2000s. The
73 sediment regime shifted from two equivalent seasons of sediment yield (spring and autumn)
74 to a single major season regime. In 2000s, autumn produced over 4 times more sediment than
75 spring. The enhanced scatter of the C-Q pairs denotes an increase of hysteresis phenomena in
76 the Wadi Abd that is probably related to the change in the hydrologic regime. At the end of
77 the period, due to ~~the~~ irregularity of the discharge, the ability of a rating curve to derive
78 suspended sediment concentration from river discharge was poor.

79 **Keywords:** water erosion; suspended sediment concentration; sediment transport; rating
80 curve; hydroclimatology; wadi; intermittent river; Algeria

81

82 1 Introduction

83 Fluvial and estuarine suspended sediment fluxes ~~are have been dramatically~~ changing
84 ~~dramatically~~ under the combined effects of anthropogenic activities and climate change. On a
85 global scale, recent changes showed a trend towards increasing land erosion and decreasing
86 fluxes to coastal waters (Walling and Fang, 2003; Vörösmarty et al, 2003; Wang et al., 2006).
87 The sediment flux trapped in regulated basins with reservoirs is higher than 50% (Vörösmarty
88 et al., 2003). Locally, it can reach more than 60% after the impoundment of one single dam
89 like on the Red River (Vinh et al., 2014), and more than 80% on rivers with many dams (86%
90 on the Yellow River, Wang et al., 2007; >95% on the Ebro ~~R~~river, Durand et al., 2002). Other
91 engineering activities (meander cutoffs, river-training structures, bank revetments, soil
92 erosion controls) also ~~affect~~ significantly affect ~~the~~ sediment fluxes and can participate to ~~the~~
93 shift from a transport-limited system to a supply-limited system, like on the Missouri-
94 Mississippi (Meade and Moody, 2010).

95 ~~Climate change, through~~With increasing temperatures and evaporation, ~~climate change~~ tends
96 to accelerate the water cycle and modify hydrologic regimes (Bates et al., 2008). Precipitation
97 intensities and the frequency of extreme events are projected to increase under climate
98 change, leading to more frequent flood events of higher magnitude that will, in turn, affect
99 patterns of erosion and deposition within river basins (Tucker and Slingerland, 1997; Pruski
100 and Nearing, 2002; Tockner and Stanford, 2002; Coulthard et al., 2012). Recent studies
101 focused on the impact of climate change on sediment transport (e.g. Gomez et al., 2009;
102 Hancock, 2009; Walling, 2009; Hancock and Coulthard, 2011; Knight and Harrison, 2013; Lu
103 et al., 2013). Syvitski (2003) showed on an example that sediment transport may increase due
104 to the increasing discharge or decrease because of the enhanced temperature. Studies ~~have~~
105 compared ~~the~~ trends in hydrological and sediment time-series to ~~the~~ land use changes (Wang
106 et al., 2007; Memariam et al., 2012; Gao et al., 2012). Climate projections are consistent
107 ~~regarding~~on warming and ~~the~~ acceleration of the water cycle (IPCC, 2013) ~~however, but~~ they
108 remain to be defined on sediment transport where projections shows a high uncertainty
109 (Shrestha et al., 2013; Lu et al., 2013). This is in part due to the fact that climate affects many

110 factors controlling sediment yield, such as surface moisture availability, weathering processes
111 and rates, and the nature of ~~the~~ riparian vegetation (Nanson et al., 2002).

112 While sediment transport is well-documented in perennial rivers in humid or temperate
113 climates, its study in semiarid areas is still fragmentary due to the difficulty of sampling
114 during flashfloods. Amongst the factors favoring erosion (slope, nature of rocks, relief,
115 climate, human activities), climate is recognized to be the main factor in semi-arid
116 mediterranean areas of Algeria which experience short and intense rain episodes, high
117 evaporating power of wind, prolonged droughts and freezing and thawing cycles (Touaibia,
118 2010; Houyou et al., 2014). Erosion is extremely active and the average concentration is at
119 least one order of magnitude higher than ~~at the~~ global ~~scale~~average (Achite and Ouillon,
120 2007). One of the main impacts of this high erosion is the rapid silting up of reservoirs (up to
121 2 to 5% per year, Kassoul et al., 1997; Remini et al., 2009; Touaibia, 2010) with important
122 consequences on water resources management in a region where 85% of rain evaporates
123 (Benhamiche et al., 2014). The high temporal variability and recent changes in forcings mean
124 that it is necessary to study sediment dynamics in such environments over time-periods of
125 several decades in order to document and understand the changes in sediment regime.

126 Achite and Ouillon (2007 hereafter referred as AO2007) analysed sediment transport changes
127 in the Wadi Abd, an Algerian wadi over a 22-year period (1973-1995). Here we extend this
128 analysis of sediment transport changes to cover a 40-year period (1970-2010). ~~In this context,~~
129 ~~this paper extends to cover a 40-year period (1970-2010) the analysis of sediment transport~~
130 ~~changes of a wadi already studied over a 22-year period (1973-1995 in Achite and Ouillon,~~
131 ~~2007 hereafter referred as AO2007).~~ The hydrologic gauging station is located upstream from
132 a dam and is not affected by any major management. This river sub-basin is also particularly
133 suitable for such study because its hydrologic regime was shown to have drastically changed
134 between the 1970s and the 1980s. Precipitation decreased and became more irregular, ~~and~~ the
135 flow regime shifted from perennial to intermittent with 26% ~~of~~ dry days ~~on~~ average in 1990-
136 1995. ~~Variations of discharge were amplified variations of discharge,~~ and a modified
137 sediment regime ~~occurred~~ with a higher and more irregular suspended particulate flux, ~~that~~
138 ~~was~~ 4.7 times higher over the period 1985-1995 than over 1973-1985. AO2007, showing the
139 advantage of working over 22 years of measurement, however, stressed the difficulty of
140 defining a reference period, and the need to extend the study ~~over a longer period longer of~~
141 ~~time~~. The objectives of this additional study are to 1) describe ~~the~~ precipitation, discharge and
142 sediment flux variability of the Wadi Abd basin over a 40-years period; 2) detect the shift, if

143 any, in temperature, runoff and sediment yield;; 3) determine the relationship between
144 sediment load and runoff over the last 40 years;; 4) detect when a shift occurred in the runoff-
145 sediment load relationship;; 5) analyze the possible causes of the change in flow regime and
146 its consequences on suspended sediment discharge;; 6) assess the use of rating curves and the
147 physical signification of its parameters when a river is experiencing a transition and ~~turns~~
148 ~~shifts~~ from a perennial regime to an intermittent regime.

149 **2 Study area: the Wadi Abd**

150 **2.1 General information**

151 The Wadi Abd, located in ~~the North-Western of~~ Algeria, is a tributary of the Wadi Cheliff,
152 the ~~principal~~major river of Algeria (Fig. 1). The length of the Wadi Abd's main stream is 118
153 km, ~~its the~~ basin area is 2480 km² and the drainage density is 3.70 km km⁻² (Fig. 2a). The
154 Wadi Abd supplies the downstream ~~the~~Sidi Mohamed Benaouda (SMB) reservoir which ~~has~~
155 a basin area ~~isof~~ 4900 km². The Wadi Abd catchment area is formed of erodible sedimentary
156 rocks from the Upper Jurassic (45.9% of its surface), Middle Jurassic (20.2%) and Pliocene
157 (7.4%) (Fig. 2b). Soft bottom sedimentary deposits from the Quaternary cover 13% of the
158 basin along the wadi (Tescult International, 2004).

159 The climate is Mediterranean and is characterized by a dry season from April to
160 August/September, and a wet season from September to March. The hydraulic deficit is very
161 high. Annual precipitation is 264 mm on average while the mean ~~potential~~ evapotranspiration
162 over the SMB basin is 1525 mm (Tescult International, 2004).

163 The watershed mainly consists of steep slopes (Fig. 2c) with very sparse vegetation or bare
164 soil (Fig. 2d). The main land use is natural environment (73%; 17% of forests + 56 % of scrub
165 and bare steppe soils), cultivated lands cover about 26% and cities 0.4%. Seven hill reservoirs
166 were built in the Wadi Abd basin from 1986 to 2004 for agriculture (irrigation ~~and~~, livestock
167 watering) or for fire fighting measures. Their total cumulated capacity is 0.88 hm³,
168 representing 2.3 % of the yearly averaged discharge at Ain Hamara station. These small
169 reservoirs are now silted up to 70% of their volume.

170 123.000 inhabitants were living in the Wadi Abd basin in 2008 (average density: 49
171 inhabitants km⁻²), 44% of them living in the city of Takhmaret. The Wadi Abd is thus little
172 influenced by human activities, in view of its extensive surface that is subject to severe
173 natural erosion.

174 In the plain, sheet (interrill) and rill erosion dominates (Fig. 3 b, f). Gully erosion is mainly
175 restricted to the mountainous regions of Freneda and Tiaret in the North (Fig. 3 c, d and Fig.
176 2c), while some mid-slope areas are gullying (Fig. 3 a, e).

177 **2.2 Data**

178 Long-term series of temperature measured at 3 stations in Algeria were extracted from
179 CRUTEM4 (Jones et al., 2012; Osborn and Jones, 2014). These stations are located at Chlef
180 (36.20°N, 1.30°E - 1951-2011), Miliana (36.30°N, 2.20°E - 1922-2011) and Dar El Beida
181 (36.70°N, 3.30°E - 1856-2011). ~~The a~~Annual average temperatures were calculated for each
182 station from the 12 monthly averages. ~~The 20 missing monthly data (out of over 480) did not~~
183 ~~allow us to exactly calculate mean measured yearly temperature~~ at Chlef, the nearest station
184 from the Wadi Abd. ~~In order to estimate the change per decade at Chlef either at the yearly or~~
185 ~~seasonal scale, the 20 lacking values~~ were extrapolated from the monthly temperatures
186 measured at Miliana and Dar El Beida using the relationships between the monthly average
187 temperatures at Chlef and Miliana, and Chlef and Dar El Beida. ~~Such relationships~~
188 ~~established at the annual scale are shown on~~ (Fig. 4). The resulting estimates of temperature at
189 Chlef ~~at on~~ seasonal and yearly scales allowed us to estimate changes by decade over the
190 period 1970-2010.

191 Rainfall and hydrometric records were provided by the National Agency of Hydraulic
192 Resources (ANRH). Time series of rainfall data are available at 6 stations within the basin
193 (see Fig. 2a): S1 Ain Kermes (altitude: 1162 m), S2 Rosfa (960 m), S3 Sidi Youcef (1100 m),
194 S4 Tiricine (1070 m), S5 Takhmaret (655 m) and S6 Ain Hamara (288 m). 9076 coincident
195 instantaneous measurements of water discharge (namely Q , in $\text{m}^3 \text{s}^{-1}$) and suspended sediment
196 concentrations (C , in g L^{-1}) were recorded at the Ain Hamara gauging station between
197 September 1970 and August 2010. Water depths were measured continuously and a
198 calibration between water level and discharge was regularly performed from velocity profiles.
199 ~~During flow measurements, water was manually sampled once or twice using a 1 L dip at the~~
200 ~~edge of the wadi. The number of samples was adapted to the flow regime. During baseflow~~
201 ~~samples were collected every other day, whereas, during floods samples were collected at~~
202 ~~higher rates (up to one every 30 min). Water samples were filtered on pre-weighed Whatman~~
203 ~~Glass Fiber Filters (GFF) filters, oven-dried at 105°C for 24h, and weighed again to determine~~
204 ~~the concentration following the protocol described by A02007 and Megnounif et al. (2013).~~
205 ~~This method, used by ANRH at all hydrologic stations in Algeria, underestimates the~~

206 suspended load as compared to its value averaged over the cross section under low turbulence
207 (i.e. at low flow) since water is sampling near the surface (Touat, 1989). During floods, which
208 transport most of the sediment load, turbulence is high enough to homogenize suspension
209 load. While this underestimation may slightly affect the budget, it doesn't severely affect the
210 time variability of suspended matter which is analyzed in this paper. From ~~these~~ 9076
211 coincident instantaneous data measured during 1213 days, average arithmetic values were
212 calculated per day so as to obtain 1213 pairs of "mean daily" (C, Q) values. The resulting
213 "mean daily Q" differs from the (true) daily discharge obtained from the averaging of 24h of
214 continuous instant Q.

215 The Atlantic Multidecadal Oscillation (AMO) index is an index of North Atlantic
216 temperatures. The monthly unsmoothed values used in this study were calculated by NOAA,
217 Earth System Research Laboratory, Physical Sciences Division/ESRL/PSD1
218 (<http://www.esrl.noaa.gov/psd/data/timeseries/AMO/>).

219 **3 Models and Methods**

220 **3.1 Trends**

221 The analysis of trends was conducted following a method fully described by Stahl et al.
222 (2010) and Déry et al. (2005) for river runoff. The Kendall-Theil Robust Line furnishes a
223 linear equation from a time-series of n measurements such as

$$224 \quad y = m t + b \quad (1)$$

225 where t is time (year), y denotes the hydrological parameter (precipitation, river discharge,
226 sediment discharge), and m is the magnitude of the trend over this period. m is calculated as
227 the median of all slopes m_k of consecutive pairs of values:

$$228 \quad m_k = \frac{y_j - y_i}{t_j - t_i} \quad (2)$$

229 where $k = [1, n(n-1)/2]$, $i = [1, n-1]$, $j = [2, n]$. This slope is often referred to as the Sen slope (Sen,
230 1968). The significance of this trend at a level p was calculated following Ziegler et al.
231 (2003).

232 **3.2 Rating curves**

233 C and Q measurements were used to define rating curves that estimate C from measured
234 values of Q, according to a common approach (e.g. Walling, 1977; Asselman, 2000; El Mahi

235 et al., 2012; Tebbi et al., 2012; Louamri et al., 2013). The most suitable model is a power law
236 of the type $C=aQ^b$ for which the coefficients (a, b) determined empirically account for the
237 effectiveness of erosion and transport. ~~In this paper,~~ the rating curve established from the
238 1213 daily averages of C and Q data ~~available for the period 1970-2010~~ enabled the estimation
239 of C then Q_s ($Q_s = C \times Q$) for the whole period 1970-2010 from the measured daily Q values.

240 Considering the change in hydrologic regime during the study period, we wondered if the
241 estimate of C and Q_s per sub-periods ~~such as~~ like decades ~~would be~~ more appropriate ~~better~~
242 ~~adapted than globally~~. We therefore applied the 4 rating curves established for the 4 decades
243 to the time series of daily Q to obtain daily C and then daily Q_s . This method (B) enabled us
244 to compare the estimated solid discharge with the value provided by the global relationship
245 established from 40-years of data (method A). The average error for daily Q_s values was 51%
246 using method A and 42.1% using method B. However, the cumulative flux of suspended
247 matter over the 1213 days for which daily data are available was over-estimated by 3.1%
248 using method A while it was under-estimated by 5% using method B. A comparison of the
249 estimates by these two methods showed that method B is not reliable ~~at~~ for high discharge
250 during the last decade because of an increase in scattering of the C, Q pairs. The relationship
251 obtained over the last decade (2000-10) lead to an under-estimation of Q_s of 23% over the 314
252 days for which daily C and Q are known. In contrast, the global algorithm from method A led
253 to an under-estimation of the same cumulated Q_s by only 3.5% over the same period. The
254 relationship established over 40 years was therefore used for this study.

255 It should be noted that although method A provides some daily solid discharges from the
256 1213 daily Q values with a high error (the average error being 51%), it enabled the
257 reconstruction of good trends of Q_s values over more than 7 orders of magnitude (Fig. 5).
258 ~~However,~~ ~~†~~The temporal variability of the coefficients a ~~and~~, b of the rating curves calculated
259 over years or decades will be discussed in light of the variability of the forcings and their
260 ~~consequences on~~ sediment transport ~~consequences,~~ in order to ~~better~~ understand ~~better~~ their
261 physical meaning.

262 3.3 Average loads

263 In order to analyze the temporal variability of suspended sediment flux, we use the average
264 concentration resulting from the ratio between the solid and the liquid flow rate, denoted as
265 ~~(~~SPM*~~)~~, which can be defined for any integration period (day, month, season, year).

266 **3.4 Study of breaks: double mass curve**

267 Double-mass curves were used to determine long term trends and changes in the hydro-
268 sedimentary regime (Searcy and Hardison, 1960; Walling, 1997; 2006).

269 **4 Interannual variations of temperature, precipitation, river discharge and**
270 **flow regime**

271 The statistics of hydrological parameters at Ain Hamara gauging station over the period 1970-
272 2010 are reported in Table 1.

273 **4.1 Temperature**

274 Temperature in Northern Algeria at the three stations of Chlef, Miliana and Dar El Beida
275 increased from the 1970s onwards (Fig. 6), with higher values. On average, temperature was
276 higher at Chlef (between 17.5°C and 20.3°C) than at Dar El Beida (15-18.5°C) and Miliana
277 (14.5-18.5°C). In average, temperature at Chlef increased by 0.96°C between 1970-85 and
278 1985-2010, and by 1.17°C from the 1970s to the 2000s (Table 2). The increase was, on
279 average, 0.87°C between the 1970s and the 1980s which is more than four times the
280 difference between the 1980s and the 1990s (+0.19°C) and the 1990s and the 2000s
281 (+0.12°C). As has been shown on a global scale, the decade of the 2000s was the warmest
282 (IPCC, 2013).

283 **4.2 Precipitation**

284 Annual precipitation at Ain Hamara station was highly irregular, varying between 165 mm yr⁻¹
285 and 506 mm yr⁻¹ (Table 1, Fig. 7). Mean annual precipitation (*P*) was 264 mm, with a
286 coefficient of variation (CV) of 27% between 1970-71 and 2009-10. The interannual
287 variations of *P* (Fig. 7) showed trends towards a decrease of rainfall (-1.86 mm yr⁻¹ on
288 average over 40 years, *p* < 0.05). *P* decreased by 15 % (from 310 to 264 mm) between the
289 1970s and 2000s if we consider the values averaged over decades (Table 2). However, a
290 more precise analysis shows that rainfall greatly decreased from the 1970s to the next decade
291 (from 310 to 231 mm, -25%), then slightly increased in the two following decades (average of
292 250 mm yr⁻¹ in 1990s and 264 mm yr⁻¹ in 2000s, see Table 2).

293 The average precipitation over the 6 rainfall gauging stations within the basin was 273 mm yr⁻¹
294 ¹, with consistent variations as compared to. Their interannual variations were consistent and
295 showed a similar variation to Ain Hamara station. Amongst decades, the coefficient of

296 ~~variation varied between 12% and 20%.~~ Five out of 6 stations show a decrease in precipitation
297 between 1970-1985 and 1985-2010, the average deficit being equal to 3.7 %.

298 **4.3 River discharge and flow regime**

299 The ~~mean~~ annual discharge ~~at the Ain Hamara gauging station~~ was $1.18 \text{ m}^3 \text{ s}^{-1}$ on average
300 ~~over the 40 year period of observation (Table 1).~~ and exhibited a higher ~~The~~ interannual
301 variability ~~of yearly averaged values of discharge~~ (CV=44.4%, ~~see Table 1~~) ~~was higher~~ than
302 ~~that of annual~~ yearly precipitation (Table 1). Yearly ~~averaged values of Q~~ values showed a
303 trend towards an increase of river flow ($+11.3 \text{ L s}^{-1} \text{ yr}^{-1}$ on average over 40 years, $p < 0.01$;
304 Fig. 7), ~~with decreasing decadal values.~~ ~~The averaged values over decades decreased~~ between
305 the 1970s and the 1980s, then ~~increasing values afterwards, similar to P~~ (Table 2). ~~Globally,~~
306 ~~they increased by 25% (from 1.16 to 1.45 m³ s⁻¹) between 1970s-80 and 2000s-2010.~~

307 ~~The~~ detailed analysis of ~~the~~ daily river discharge shows that the river was perennial in the
308 1970s and then became intermittent during the 1980s (Fig. 8). The driest year occurred in
309 1993-94 with 117 days of fully dry river. On Fig. 8, ~~the~~ very low river discharges (around
310 $0.01 \text{ m}^3 \text{ s}^{-1}$) were not considered as days of dry river.

311 ~~The “wet discharge”, denoted Q_w , i.e. the yearly average river discharge of the days of~~
312 ~~running river (and not calculated over the full year) was also calculated (Table 2).~~ From 1970s
313 to 2000s, ~~Over the 40 year period~~ when Q averaged over 10-years increased by 25%, the wet
314 discharge Q_w (i.e. the yearly average discharge of the days of running river) ~~averaged over 10-~~
315 ~~years~~ increased by more than 35% ~~from 1970-80 to 2000-2010~~ (Table 2) ~~from 1.16 to 1.57 m³~~
316 ~~s⁻¹).~~

317 ~~Q and Q_w increased as did the number of dry days (and consequently the intra-annual~~
318 ~~variability) and their intra-decade variability (Table 2).~~ Two indicators of intra-annual
319 discharge variability are shown in Fig. 7: Q_{98} , the 98th percentile of annual flows calculated
320 from daily discharge and the standard deviation of daily discharge within each year (σ_Q). Q_{98}
321 increased by a factor 3.2 ~~between from an average of 4.37 m³ s⁻¹ over the period 1970-80~~
322 ~~and to 13.94 m³ s⁻¹ over the period 2000-2010 (Table 2), a factor 3.2 increase.~~ Q_{98} is ~~also~~ a
323 good indicator of changes in sediment transport as it occurs during the highest flood events
324 that occur each year.

325 **4.4 Summary: changes of hydrologic forcings**

326 These results indicate that four significant changes occurred during the 40 year period in the
327 Wadi Abd basin (Table 2): (1) an increase of temperature at Chlef by 1.17°C between the
328 1970s and the 2000s; (2) a decrease in precipitation of 15% over 4 decades; (3) an increase in
329 average annual flow of 25 % over the same period, or 35% if we consider only the days when
330 the river is not dry; (4) a change in the flow regime, from a perennial regime to intermittent
331 regime. The pivotal year after from which the river experiences dry weeks is the hydrological
332 year 1985/86 with 49 days with no flow. This number increased in the following years (Fig.
333 8).

334 **5 Interannual variation of sediment load**

335 **5.1 Rating curve**

336 The rating curve obtained from 1213 pairs of daily averages gave:

$$337 C = 2.270 Q^{0.647} \quad (r^2=0.431) \quad (3)$$

338 where C is expressed in g L⁻¹ and Q in m³ s⁻¹. 43% of the variations of C are explained by
339 those of Q (~~r²=0.431~~). The rating curve obtained between Q and Q_s shows a much higher
340 ~~determination~~ coefficient of determination (r²=0.831) but is biased since Q_s = C x Q.
341 Nevertheless, both relationships give estimates of Q_s values from Q with less than 1%
342 difference which is less than the uncertainty of Q_s.

343 **5.2 Yearly sediment fluxes and concentrations**

344 Decadal variability of Q_s

345 Q_s increased from 180 to 1130 x10³ tons per year between the 1970s and the 2000s (Table 2).
346 The increase from one decade to the next is remarkably regular: +85% between the 1970s and
347 the 80s, + 84% between the 80s and the 90s, +84% between the 90s and the 2000s and is
348 statistically significant (+19.7 10³ t yr⁻¹ in average, p < 0.05). Specific sediment yield follows
349 the same trend ~~increasing from 72 t km⁻² yr⁻¹ in the 1970s to 455 t km⁻² yr⁻¹ in the 2000s~~
350 (Table 2).

351 Variability of mean annual load SPM*

352 The average value of SPM* ~~calculated~~ over the period 1970-2010 is 12.3 g L⁻¹, with annual
353 values comprised ~~. The 40 annual values of SPM* calculated for each year from measured~~

354 ~~discharges and concentrations estimated using the rating curve (3) vary~~ between 2.5 g L⁻¹ and
355 50.2 g L⁻¹ (Tables 1, 2). Their interannual variation was smaller than that of solid discharge
356 because annual SPM* is the ratio of the annual Q_s to the annual Q (which increased less than
357 Q_s). ~~The variability of SPM* was thus smaller than that of annual Q_s (CV=86.0% instead of~~
358 ~~123.3% over 40 years).~~

359 Analysis of break points

360 The double mass plot enabled us to identify changes in the sediment response of the stream
361 (Fig. 9). A major break occurred in 1985-86. A secondary break was noticed in 1991-92, but
362 the entire period 1985-86/2009-10 may be considered as a single period (with the relationship
363 « cumulated Q_s » = 0.021 « cumulated Q » - 9.417, r²=0.989). The period 1985-86/1991-92
364 may thus be considered as a transient event towards a new regime.

365 The response of sediment flow to various constraints (~~changes in precipitation, hydrology,~~
366 ~~plant, agricultural practices etc.~~) differs clearly from that of discharge from the year 1985-86
367 onwards. This break corresponds to the first year of dry river over a long period in summer
368 (49 days). This initiates a phase of intermittent flow regime. The averaged parameters for the
369 two periods 1970-1985 and 1985-2010 were added to the tables, in addition to average values
370 throughout the full study period and values for decades to illustrate the dynamics of the
371 hydrological and hydro-sedimentary change.

372 **5.3 High dependency of the solid discharge on Q variability**

373 The variability of Q and Q_s or SPM* at different time scales were compared. AO2007 showed
374 that, over 22 years, 71% of the variance of the annual SPM* values was accounted for by ~~the~~
375 annual discharge and 73% by the 95th percentile of daily discharge within the given year Q₉₅.
376 This means that SPM* was mainly driven by the 10 to 15 highest daily discharges in a year
377 suggesting a strong correlation between yearly Q_s and the discharge variability. Finally, they
378 showed a remarkable linearity between SPM* and the standard deviation of the daily
379 discharge per year (σ_Q).

380 Yearly SPM* and yearly σ_Q still showed a strong linearity over 40-years ~~period~~ (r²=0.956,
381 Fig. 10a). A higher correlation was obtained between yearly Q_s or SSY, the specific sediment
382 yield, and yearly σ_Q (r²=0.991, Fig. 10b). ~~In conclusion, for this river, This is one of the most~~
383 ~~important conclusions from this river where the solid discharge depends on discharge~~

384 ~~following a rating curve:~~ the yearly solid discharge is more closely dependent on the
385 discharge variability than on discharge values.

386 **6 Variation of the seasonality of climatic and hydrological parameters**

387 The yearly values of temperature at Chlef ~~generally~~ increased ~~on average~~ but the monthly
388 averages showed high discrepancies. Temperature from March to November increased with a
389 maximum of increase in June (+3.30°C on average between the 1970s and 2000s), it remained
390 quite constant in December and February and decreased by 0.98°C in January over the same
391 period. Considering the average values per season, winter values (Dec-Feb) decreased by
392 0.33°C between the 1970s and the 2000s, while spring values (Mar-May) increased by
393 1.66°C, summer values (Jun-Aug) by 2.22°C and fall values (Sep-Nov) by 1.29°C. In
394 summary, annual temperature differences increased with minimum temperatures down
395 slightly and maximum temperatures rising sharply. The increase was most marked in July-
396 August.

397 Averaged seasonal values of P, Q and Q_s for each decade are given in absolute values and in
398 percent of the yearly values in Table 3. The seasonal relative contribution of P, Q and Q_s
399 centered and averaged over 9 consecutive years are presented in Figure 11. The monthly
400 values of P, Q and Q_s per decade over 40-years also clearly illustrate the absolute changes in
401 intensity and in seasonality of the river regime (Fig. 12). The main conclusions of the analysis
402 of ~~T~~, P, Q and Q_s ~~seasonal~~ variations are the following:

- 403 • Rainfall decreased in spring and increased in autumn. Precipitation in autumn increased
404 from 22 to 30 % at the expense of spring rains (decreasing from 41% to 29%). ~~It is~~
405 ~~striking to note that~~ for the decade 2000-2010 precipitation was the same in autumn and
406 in spring (78 mm) while for the decade 1970-1980 spring rainfall was 87% higher than in
407 fall (~~128.2 mm vs. 68.5 mm~~; see Table 3 & Fig. 11a).

408 Average monthly rainfall from six weather stations in the river basin for 1970-1985 and
409 1985-2010 (Fig. 13) illustrates the changes. Two marked seasons typical of a
410 Mediterranean climate are present (a dry season and a rainy season) but the following
411 changes are observable: (1) differences between seasons decrease, as indicated by the CV
412 of monthly rainfall from 57.3 % in 1970-85 to 45.9 % in 1985-2000. There is a decrease
413 of spring rains (~~March-May~~) and at the beginning of the cold season (~~November-~~
414 ~~December~~) and the strengthening of rain in the warm season (~~July-October~~) and in winter
415 (~~January-February~~); (2) advancement of the rainy season as evidenced by precipitation in
416 ~~September and October and November~~; (3) spreading of the rainy season over 9 months

417 (September-May) for 1985-2010 from previously 7 or 8 months (from October or
 418 November onwards, ~~according to the criteria that are defined for the rainy season~~) ; (4)
 419 increased regularity of rainy season precipitation.

- 420 • Proportionally, flow decreased ~~in all seasons~~ from winter to summer and increased
 421 dramatically in autumn from just over a quarter (27.3%) of the flow delivered over ~~the~~
 422 ~~decade~~ 1970-1980 to more than one half (52.5%) over ~~the period~~ 2000-2010 (Table 3 and
 423 Fig. 11b). Flow decreased in summer and the river became dry for much of the summer.
 424 Over the last decade, it is striking to see the difference between the average flow rates in
 425 fall and spring: the fall rate is almost three times that of spring with almost the same
 426 rainfall. This trend is evident over the 40 year period (Fig 11b).
- 427 • These results point towards a change in runoff as defined by the ratio Q/P. Considering
 428 the whole basin area, the river discharge at Ain Hamara station averaged over 40-years
 429 corresponds to a water depth of 15 mm yr⁻¹, while the average precipitation is 264 mm yr⁻¹.
 430 For comparison, on average 85% of rain in this region evaporates and the remaining
 431 15% runs into surface waters or infiltrates into underground storage (Sari, 2009, quoted
 432 by Benhamiche et al., 2014). On the Wadi Abd, Q/P averages 5.7%. We calculated the
 433 value of Q/P averaged over 3 consecutive years and over 3 consecutive months (centered)
 434 and then took the average per decade (Fig. 14). It appears that the Q/P ratio remains
 435 constant during the months from December to April (around 4.4% in average), it
 436 increased slightly in November and May during the decade 2000-2010 and it increased
 437 significantly from September to November. In other words, runoff increased, rain
 438 decreased slightly and the temperature (and therefore ETP) increased. As a consequence,
 439 infiltration will decrease and the water level in the aquifers will be lowered. ~~Moreover,~~
 440 ~~Q/P, which was very high in July and August in the 1970s, has nearly halved since the~~
 441 ~~1980s.~~
- 442 • In absolute values, solid discharge has been increasing in all seasons over 4 decades, but
 443 more so in the fall than in the other seasons (Table 3 and Fig. 12c). During autumn, it
 444 more than doubled from one decade to another ~~(x 2.07 in the 1980s vs. 1970s, x 2.17~~
 445 ~~from 1980s to 1990s, and x 2.88 from 1990s to 2000s).~~ During the other seasons, it
 446 doubled or tripled ~~within 30 years,~~ between the 1970s and 2000s. ~~The average annual load~~
 447 ~~was multiplied per 1.84 from one decade to another (1.8 10⁶ tons during the 1970s, 3.34~~
 448 ~~in 1980s, 6.14 in 1990s and 11.30 in 2000s,~~ see Table 2). While during the 1970s the
 449 Wadi Abd had two major periods of roughly equivalent sediment discharge in the fall and
 450 spring, suspended sediment loads were greater in the autumn during the 2000s (> 70%).

451 The Wadi shifted from a regime with two equivalent seasons of sediment production to a
452 regime with one dominant season in the 2000s. Autumn produced over 4 times more
453 sediment than spring in the 2000s (Table 3, Fig. 11c). This phenomenon does not seem to
454 be due to some exceptional floods because the trend is observable over 4 consecutive
455 decades (Fig. 11c).

456 7 Discussion

457 7.1 Interannual variations

458 Hydrology and climate change over 40 years

459 Temperature increased rapidly between the 1970s and 1980s (+0.878°C on average at Chlef).
460 The increases were lower during the three following decades ~~(1980s to the 2000s)~~. An
461 increase in temperature of 1.6°C between 1977-1979 and 2000-2006 was noted by Dahmani
462 and Meddi (2009) for the Wadi Fekan basin in West Algeria and Bakreti et al. (2013) also
463 showed a significant trend of increasing temperature in spring by 0.0183 °C per year in the
464 Tafna basin in West Algeria over the same period. However, temperature ~~has~~ not increased
465 ~~as rapidly over the so fast during the whole~~ 20th century (Fig. 6) and as mentioned by IPCC
466 (2013), “trends based on short records are very sensitive to the beginning and end dates and
467 do not in general reflect long-term climate trends.” The longest available time series of
468 temperature in Algeria was measured at Dar El Beida near Algiers. At this station, average
469 temperature increased by 0.62 °C between 1850-1900 (29 yearly values available) and 2003-
470 2012 (Fig. 6), while it increased between 1880 and 2012 by 0.85°C globally (IPCC, 2013).

471 A global trend towards ~~an~~-increasing temperatures and increasing dryness in Algeria from the
472 1970s onwards has already been described (Meddi and Meddi, 2009). Over the period 1923-
473 2006 North Algeria experienced an alternation of wet periods (1923-1939, 1947-1973) and
474 dry periods (1939-1946 and from 1974 onwards) (Benhamiche et al., 2014). Over 70 years in
475 the Wadi Fekan, Dahmani and Meddi (2009) showed that the period 1943-1960 was rather
476 wet, that 1960-1975 was average, and that the period 1975 onwards (up to the end of their
477 data set in 2004) was dry and of an exceptional long duration. Using three different statistical
478 tests (Pettitt, Lee Heghinian and Hubert), Meddi and Meddi (2007) shown that a shift was
479 observed between 1973 and 1980 over most of the rain gauges in Algeria. In North-West
480 Algeria, a shift was noticed in 1973 in winter rainfall and between 1974 and 1980 in spring
481 rainfall, both of them being responsible of the yearly rainfall deficit (Meddi and Talia, 2008).
482 From the rainfall dataset at the Ain Hamara station between 1968 and 2007, Hallouz et al.

483 (2013) showed that the break in annual rainfall occurred in 1976 and calculated a deficit of
484 19% between 1968-1976 (304 mm yr⁻¹) and 1976-2007 (247 mm yr⁻¹). At the stations Ponteba
485 and Rechaiga, near to the Abd basin, the trends of decreasing total precipitation and of
486 increasing mean length of dry spells were amongst the 5 highest in the Maghreb area over the
487 22 stations considered by Trambly et al. (2013, see their Fig. 8).

488 As a consequence of the decrease of rainfall after the 1970s break which was observed in
489 most basins of Western Algeria, river discharges were generally seen to decrease as well.
490 Meddi and Hubert (2003) showed that the decrease in river discharge varied between -37%
491 and -70% from ~~the~~ Eastern Algeria to ~~the~~ Western Algeria. ~~Over~~In the Mecta basin in North-
492 West Algeria, runoff was estimated to be 28-36% lower in 1976-2002 as compared to 1949-
493 1976 (Meddi et al., 2009). ~~Over~~In the Tafna basin, also in North-West Algeria, Ghenim and
494 Megnounif (2013a, 2013b) showed that the decrease ~~in~~ precipitation ~~after the break point~~
495 ~~was, on average, by~~ 29% ~~on average~~ over the ~~whole~~ basin (especially in winter and spring)
496 ~~after the break point and~~ was accompanied by a decrease of 60 % in river flow.

497 In ~~contrast~~~~this context~~, the Wadi Abd ~~had a behaved~~ differently ~~behavior since then~~ that river
498 discharge increased. The counter-intuitive increase of runoff with decreasing rainfall has also
499 been observed in ~~the~~ Sahel and is referred to as « the Sahelian paradox » (see Mahé and
500 Paturel, 2009; Mahé et al., 2012). A closer look at the seasonal variations of the different
501 parameters shows that Q decreased in winter and spring but that Q/P increased in autumn
502 when rainfall increased. Overall Q increased. The decrease of rainfall in spring and its low
503 level in summer may have lead to a change in vegetation cover which would in turn decrease
504 infiltration. However, although studying the vegetation dynamics of the basin goes beyond the
505 scope of this study, this aspect could be investigated in the future using satellite data, for
506 example.

507 What is the influence of large-scale circulation indices?

508 Changes in precipitation are derived from atmospheric-oceanic signals (Milliman et al., 2008;
509 Giuntoli et al., 2013). Low frequency fluctuations related to climate change are modulated
510 with higher frequency interannual fluctuations, such as ENSO (El Niño Southern Oscillation),
511 NAO (North Atlantic Oscillation), AMO (Atlantic Multidecadal Oscillation) or MO
512 (Mediterranean Oscillation). Trambly et al (2013) showed that the precipitation amounts and
513 the number of dry days over the Maghreb were significantly correlated with the MO and
514 NAO patterns. MO and NAO showed positive trends from the 1970s onwards which could

515 explain the trend towards decreasing frontal conditions over the Mediterranean basin and thus
516 increasing droughts.

517 Interannual influence by the Austral oscillation ENSO over Algeria was shown to be higher in
518 North-West Algeria on the highest discharges than on the average discharge. The maximum Q
519 seems to be smaller during El Niño and higher during La Niña in North-West Algeria (Ward
520 et al., 2014). ~~Average discharge is less influenced by ENSO than the maximum yearly~~
521 ~~discharge (Ward et al., 2014).~~The frequency of extreme rainfall events shows the highest
522 correlation with the Mediterranean Oscillation Index in Algiers and with the Southern
523 Oscillation Index in Oran (Taibi et al., 2014).

524 In this study, no significant correlation was established between a series of hydrological
525 parameters in the Wadi Abd and the Southern Oscillation Index. The average of AMO per
526 hydrologic year was calculated from its monthly values. AMO has increased from 1970s to
527 the 2000s, with negative values up to 1993-94, then positive ~~values thereafterwards~~ (except in
528 1996-97). Its decadal average was -0.25 in the 1970s, -0.12 in the 1980s, 0.0 in the 1990s and
529 0.18 in the 2000s. AMO and the discharge variability of the Wadi Abd within the year
530 increased coincidentally. The yearly AMO values have a coefficient of determination of 0.226
531 when correlated with the standard deviation of daily river discharges within the year, a proxy
532 for the variability of daily discharge. However, this information does not allow us to conclude
533 that the Atlantic Multidecadal Oscillation is responsible for hydrological changes in the Wadi
534 Abd basin.

535 Break point in 1985-86: change of flow regime

536 The several weeks of dry river for the first time in 1985-86 (49 days) can be considered as a
537 threshold effect, which marks the start of a new flow regime. The appearance of a dry regime
538 is a break, a fully nonlinear phenomenon. It has strong consequences for water infiltration and
539 groundwater recharge, on ~~the~~ seasonality, intensity and type of floods, and in turn, on erosion
540 and sediment transport. 1985 is also a pivotal year for recent climate change as evidenced by
541 the rapid increase in global mean ~~air~~ temperature anomaly ~~of air~~ from that year until 1993
542 (Fig 1 in Lockwood and Fröhlich, 2007). The hypothesis of a temporary warming caused by
543 dust emitted during the eruption of Mount Pinatubo had been advanced to explain the
544 warming since 1985, but climate scientists later recognized that the temperature anomaly has
545 been increasing since 1993, reaching about 0.6°C by 2007 compared to the global average
546 temperature calculated for the period 1951-1980 (Lockwood and Fröhlich, 2007).

547 This threshold is coincident with hydrological shifts in the Tafna basin in North-West Algeria.
548 Bakreti et al. (2013) analyzed the baseflow and baseflow index of five of its sub-basins
549 between 1976 and 2006 and ~~found evidence of~~ ruptures of the baseflow index between 1984
550 and 1990 depending on the sub-basin, in 1984, 1985 and 1990 in the mountains, and in 1985
551 and 1986 in the plain. These changes in flow regimes ~~in~~ the Tafna basin were likely caused
552 by shifts in rainfall late 1970s in the Mounts of Tlemcen and early 1980s in the plains
553 (Ghenim and Megnounif, 2013a).

554 Shift of the onset of the first summer flood

555 The analysis of the time series of daily flows enables ~~the determination of~~ ~~to determine the~~
556 start of the first summer flood. The average daily flow per decade suddenly increases the day
557 at which the first summer flood occurred, at least once in the decade. By observing these
558 decadal averaged daily flows, there is no ambiguity on the start of the ~~earliest~~ flood by
559 decade:

560 - in 1970-80, the first flood starts on the 6th September with an average 4-days discharge (6-9
561 ~~September~~) of $1.59 \text{ m}^3 \text{ s}^{-1}$, while it was on average $0.58 \text{ m}^3 \text{ s}^{-1}$ over the four previous days,

562 - in 2000-2010, the first flood of summer starts on August 8 with an average 4-days discharge
563 (8-11 ~~August~~) of $2.03 \text{ m}^3 \text{ s}^{-1}$, while it was on average $0.03 \text{ m}^3 \text{ s}^{-1}$ from 4 to 7 August.

564 During the 2000s, the first flood in summer started close to one month before that of the
565 1970s and the magnitude was 27% higher. It can be asked if this trend was observable over
566 the 40-year period or only between two specific decades. The analysis of mean dates and
567 discharges of the first flood in late dry season gave the following results for the intermediate
568 decades:

569 - 1980-1990: the first flood started ~~on~~ in average on August 31 with a 4-days average discharge
570 (~~August~~ 31-~~September~~ 3) of $2.69 \text{ m}^3 \text{ s}^{-1}$, while the average rate over the four previous days
571 was $0.13 \text{ m}^3 \text{ s}^{-1}$

572 - 1990-2000: the first flood started in average on August 22 with a 4-days average discharge
573 (~~August~~ 22-25) of $7.67 \text{ m}^3 \text{ s}^{-1}$, while the average rate over the four previous days was
574 $0.33 \text{ m}^3 \text{ s}^{-1}$. The existence of a precursor peak on August 17, which was not observed in
575 previous decades, was also observed.

576 It therefore appears that the date of the first flood advanced by about ten days each decade
577 over the previous 40 years. The shift in the onset of the first flood in summer probably has
578 important consequences on flow and erosion rates.

579 **7.2 Relationships between several parameters and sediment yield**

580 Temperature and sediment yield

581 The curve showing annual suspended load versus [the](#) global air temperature anomaly (base
582 period 1951-1980) calculated by hydrological year from monthly data provided by NOAA
583 (Hansen et al., 2010; GISTEMP Team, 2015) shows a correlation between the sediment yield
584 and ongoing climate change ($r^2=0.388$, Fig. 15).

585 Precipitation and sediment yield

586 Many authors studied the variations of sediment load per unit of catchment area against
587 annual rainfall (e.g. Summerfield and Hulton, 1994) or effective rainfall (e.g. Langbein and
588 Schumm, 1958). On the Wadi Abd, annual rainfall ~~fell sharply between was 310 mm yr⁻¹ in~~
589 the 1970s ~~and, fell sharply in~~ the 1980s then slightly increased over the following decades ~~to~~
590 ~~between 231 and 264 mm yr⁻¹~~. Meanwhile, yearly sediment concentration and suspended
591 sediment discharge have increased. The comparison of their respective variations shows a
592 lack of correlation between precipitation and annual sediment yield ($r^2 < 0.1$ regardless of the
593 type of regression considered). Regarding the relationship between precipitation and erosion,
594 if there are correlations between their spatial variations reported in the literature (though with
595 a strong scatter, see Riebe et al., 2001), our study shows that the temporal variations of
596 precipitation and sediment yield are not correlated in the Wadi Abd. This may be due to the
597 change of flow regime within the study period.

598 Runoff and sediment yield

599 Although runoff was noted to have a limited impact on the distribution of sediment yield at
600 regional or global scales by Aalto et al. (2006), Syvitski and Milliman (2007), Vanmaercke et
601 al. (2014), the temporal variability in precipitation, runoff (or discharge) and consecutive
602 vegetation cover was shown to be locally the main impact on fluvial sediment load (see [the](#)
603 ~~review of~~ Vanmaercke et al. 2014, p. 360). ~~Our results confirm that, on~~ the Wadi Abd, the
604 yearly suspended sediment load was highly correlated with discharge (Q mean or its highest
605 percentiles) and to its intra-annual fluctuation (Fig. 10). ~~Climate change alters the hydrology~~
606 ~~of a river basin such as the Wadi Abd.~~ Although the river regime shift clearly impacted

607 several parameters ~~between the two periods~~, the relationship between yearly sediment load
608 and discharge variability did not change over the 40-year study period.

609 ~~We cannot conclude on the exact origin of the regime change but note that it occurred when~~
610 ~~dry periods started, annual precipitation timing shifted and runoff increased.~~

611 **7.3 On the use of double-mass curves to determine the climate change and** 612 **anthropogenic influences**

613 Double-mass curves are often used to determine the impact of developments such as dams on
614 sediment discharge (e.g. Lu et al., 2013). Our findings warn about extrapolations that could be
615 wrongly made to quantify the impact of a development by extending the double mass curves.
616 Indeed, this study shows that the double-mass curve can change its slope (here increasing)
617 when the flow regime change is driven by seasonal temporal variation in precipitation and
618 runoff that isn't linked to any specific anthropogenic activity (such as a dam impoundment)
619 within the basin.

620 **7.4 Physical meaning of rating parameters a & b**

621 Interannual variation of (a, b)

622 Since $C = a Q^b$, with $b \neq 0$, $C(1) = a$. a thus represents the sediment concentration when the
623 river discharge is $1 \text{ m}^3 \text{ s}^{-1}$, and b reflects the sensitivity of concentration to discharge
624 variation. The general formula $\ln C = \ln (aQ^b)$ provides:

$$625 \quad dC/C = b \, dQ/Q \quad (4a)$$

$$626 \quad b = dC/dQ \, Q/C = 1/a \, dC/dQ \, Q^{(1-b)} \quad (4b)$$

627 thus b varies almost like $1/a$ (Asselman, 2000). Many papers discuss the physical meaning of
628 the rating parameters a and b (see AO2007) and try to connect their values to physiographical
629 characteristics, vegetation cover or hydro-meteorological forcing.

630 The river's regime change is accompanied by a change in the (a, b) pairs of rating curves
631 defined for multi-year periods such that a increases and b decreases (Table 2), following:

$$632 \quad b = -0.294 \, \ln a + 0.912 \quad (r^2=0.582) \quad (5a)$$

$$633 \quad \ln b = -0.188 \, a + 0.042 \quad (r^2=0.649) \quad (5b)$$

634 Equation (5a) is very similar to that presented by Iadanza and Napolitano (2006) for the Tiber
635 River after the construction of a dam ($b = -0.3815 \, \ln a + 0.7794$, $r^2=0.992$). Before the

636 construction of this dam, another relationship corresponded to more than 3 times higher
637 sediment yields. Asselman (2000) has suggested to interpreting these regression lines in a Ln a
638 - b graph as different sediment transport regimes.

639 On the Wadi Abd, the change in sediment transport regime is not evident from the yearly (a,
640 b) values but it becomes clearly observable when considering a and b values averaged over
641 moving periods of several years. The best correlations were obtained for running averages
642 over 15 years named a_{15} and b_{15} (N=25, from 1970-1985 to 1995-2010, see Fig. 16). The
643 available data set does not allow us to determine if results obtained from averaging over
644 longer periods would perform best.

645 The time evolution of the moving average pair (a_{15} , b_{15}) clearly shows a first relationship with
646 the values dominated by the pre-1985 regime (8 values from 1970-1985 to 1977-1991),
647 another one for the values predominantly after 1990 (12 values from 1983-1997 to 1995-
648 2010), both with a_{15} increasing and b_{15} decreasing, and a transitional regime centered on the
649 period 1985-1990 (Fig. 16). During the transition period centered over 1985-1990, b_{15} was
650 almost constant (between 0.72 and 0.74) while a_{15} was increasing from 2.01 to 2.34. During
651 the period 1985-1991 the yearly values of b varied very little (between 0.653 and 0.672) while
652 yearly a increased significantly from 1.81 in 1985-86 to 3.23 in 1990-91. Higher a and lower
653 b values are in the literature typical of highly arid river basins, such as the ephemeral Nahal
654 Eshtemoa in Israel, where $a=16.98$ and $b=0.43$ (Alexandrov et al., 2003).

655 As the break points were coincident, it is possible to analyze the change of (a_{15} , b_{15}) in terms
656 of shift of hydrological regime. However, if the new hydrological regime was immediate from
657 1985 onwards, the change in the C-Q relationship was only evidenced in the Wadi Abd at
658 mid-term, considering 15-years average values.

659 Parameters that explain a (or b)

660 The coefficient of determination between a and specific sediment yield (SSY) is low at the
661 annual scale but higher when we consider the moving averages of a and SSY over 15-years.
662 The specific sediment yield explained 95.2% of the variance in the interannual scale (Fig. 17),
663 much more than the average river flow did ($r^2= 0.839$). ,following:

$$664 \quad a_{15} = 6.104 \cdot 10^{-3} \cdot SSY_{15} + 1.117 \quad (r^2 = 0.952) \quad (6)$$

665 b_{15} showed a lower correlation with the SSY ($r^2=0.853$) than a_{15} did.

666 In summary, the moving average of a is strongly correlated to specific sediment yield over the
667 same moving period of 15 years, and the moving average of b can be deduced from a ~~over the~~
668 ~~same period~~ using ~~thea~~ relationship which is given in Fig. 16 as a function of ~~per~~-flow regime,
669 either perennial or intermittent.

670 Validity range of rating curves

671 The estimation of sediment yield from flow measurements and a rating curve is still
672 acceptable throughout the study period (Fig. 5). However, ~~it should be noted that~~ the pairs (C,
673 Q) become increasingly scattered with time around the best-fit curve, ~~as attested by the~~
674 ~~decrease of t.~~The coefficient of determination ~~has decreased~~ from one decade to another ~~over~~
675 ~~40 years, from 0.57 to 0.38~~ (Table 2).

676 Intermittent flows induce a stronger dependency of river behavior on antecedent wetness
677 (Beven, 2002) and antecedent weathering, i.e. a strong dependency on memory through
678 threshold and hysteresis effects. With increasing memory effects, coincident values of C and
679 Q become less dependent on each other and the rating curves less suitable to model their
680 relation. The study of sediment dynamics in the Wadi Abd will thus likely require in the
681 future a more appropriate method than rating curves, such as the study of each individual
682 flood, like Megnounif et al. (2013) did in the Wadi Sebdou. This finding may have
683 consequences on water management as well. When dealing with rating curves, water
684 discharge must be recorded at frequent intervals, although measurements of concentration can
685 be sparser. When rating curves cannot be applied, river discharge and sediment concentration
686 should be both frequently and simultaneously measured.

687 **8 Conclusions**

688 ~~IOver the last 40 years,~~ in response to climate change which resulted in an increase in
689 temperature of around 1.1°C between the 1970s and 2000s years ~~at Chlef,~~ rainfall moved
690 forward during the late warm season and the watershed of Wadi Abd experienced a
691 significant change in ~~the~~-flow regime ~~of the river~~ and an increased variability at both the inter-
692 annual and intra-annual levels. These changes ultimately lead to a dramatic and continuous
693 increase in sediment load over 4 decades (~~ion~~ average 84% more every decade as compared to
694 the previous one).

695 The main result of our analysis is the shift of the onset of the first summer flood that occurred
696 1 month earlier in the 2000s than in the 1970s. This shift is likely responsible for the
697 cascading effects on the hydrological regime of the Wadi Abd. In particular, earlier floods

698 during the warmer season have higher evaporation which limits the groundwater storage. A
699 parallel study of seasonal changes in vegetation cover is needed to provide additional
700 information.

701 The increase in erosion of the watershed (coefficient a) is accompanied by a decrease in the
702 coefficient b. The traditional rating curves approach which was applicable when the river was
703 perennial is now less adapted to model the behavior of the river (Table 2). This could be
704 explained by a more pronounced hysteresis phenomenon, which is consistent with the change
705 of hydrological regime in the dry season thereby limiting the utility of rating curves to model
706 C-Q relationships. ~~Other methods such as that proposed by Megnounif et al. (2013) are~~
707 ~~probably better adapted to understand future sediment dynamics of the Wadi Abd.~~

708 The rapid change in sediment regime which is instantaneously driven by the changing flow
709 regime should be distinguished from the slow change in the concentration-flow relationship.
710 The change in flow regime can be precisely dated in May-July 1986 (with 49 consecutive dry
711 days), while the change in the C-Q relationship ~~requires~~ ~~needs~~ averaging over ~~several~~ ~~15~~ years
712 of a, b and specific sediment yield to become evident. Such inertial effect may be attributed to
713 the time for the basin soil properties (such as humidity) or vegetation to adapt to the new
714 climate conditions. It likely depends, amongst other factors, on ~~the~~ underground water
715 storage, and thus on basin lithology and infiltration history. ~~Over~~ the Wadi Abd basin, the
716 time needed for the flow regime to change after the dryness settlement in early 1970's (see
717 Fig. 6) is estimated ~~as being~~ around 15 years in this study.

718 The present analysis only includes hydrological parameters. Management programs that were
719 conducted to fight erosion in Algeria from 1960s until 1990s by reforestation and setting up
720 banks over cultivated marl and clay areas proved to be little or ~~not efficient~~ (Touaibia,
721 2010). Human activities may have influenced the hydrological regime change and increased
722 erosion, in particular through firewood cutting during economically difficult periods (1990s),
723 however the shift was shown to occur earlier. The lack of data on land use and land cover
724 changes over 40 years does not allow us to isolate the factors directly related to climate
725 change from those related to other anthropogenic activities. However, the small population,
726 the low coverage of pasture (see Fig. 2d), of cultivated areas and vegetation (43 %) in the
727 basin and the small volume of reservoirs (nominally 2.3% of the annual discharge, and silted
728 up to 70%) make us think that in this system the effects of climate change dominate
729 anthropogenic effects. The quantification of forcing changes on sediment sources (raindrop
730 erosion, sheet erosion, rill erosion, gully erosion, stream channel erosion) may be investigated

731 in situ (e.g. Poesen et al., 2003) and/or estimated using a numerical model of the hydrologic
732 and sedimentological functioning of the basin, such as WEPP (Nearing et al., 1989),
733 EUROSEM (Morgan et al., 1998) or SWAT (Neitsch et al., 2011). Such a model could help
734 us to test hypotheses and quantify or at least estimate the effects of different forcing changes
735 (temperature, runoff, vegetation, etc.) in future studies.

736 It is important to emphasize that it is impossible to define long-term hydrological averages in
737 the context of a changing flow regime. ~~Our analysis is based on the shift from a perennial~~
738 ~~regime to an intermittent one.~~ The example of the Wadi Abd shows that the difficulty is
739 challenging with regard to sediment transport in suspension, since changes of flux cannot be
740 counted as a fraction but can reach an order of magnitude.

741 Changes in flow regime in relation to climate change can be investigated using climate
742 models. Das et al. (2013) using 16 climate projections showed that more intense floods of a
743 return period of 2-50 years should occur in the Sierra Nevada, regardless of the rainfall
744 variation. The recent changes in the Wadi Abd show that extreme events with increasing
745 variability already occur in the basin. Over Algeria, an increase of 1-2°C in temperature could
746 induce a reduction of 10% in precipitation before the end of the 21st century (Benhamiche et
747 al., 2014) with unknown consequences on erosion and sediment transport. Lu et al. (2013)
748 calculated the impact on sediment loads of every 1% change in precipitation or river
749 discharge in large Chinese rivers. Such a calculation has no meaning in our basin since the
750 rainfall and discharge were not monotonic (severe decrease in the 1970s then slight increase
751 during 30 years) while the sediment loads have always increased. The difficulty of forecasting
752 climate change-driven impacts on sediment yield due to non linear effects has been
753 underlined by geomorphologists (see Goudie, 2006; Jerolmack and Paola, 2010; Coulthard et
754 al., 2012; Knight and Harrison, 2013). The present study illustrates that the change ~~in~~ flow
755 regime induced a fully non linear effect between river discharge and sediment yield. This
756 needs be considered in forecasts especially in small river basins in semi-arid areas.

757 Changes in erosion and sediment transport under new climate constraints will induce changes
758 ~~over~~ the middle to long term that decision-makers must integrate into water resources
759 management, habitat status, agricultural adaptation (O'Neal et al., 2005), landscape evolution
760 (Temme and Veldkamp, 2009) as well as in many other environmental adaptations (Ouillon,
761 1998). We thus encourage the local adaptation of sampling strategies and measurements to
762 take into account changing in flow regimes. Furthermore, due to the uncertainty of water
763 resources and erosion in the Maghreb (Taabni and El Jihad, 2012) and in the Mediterranean

764 basin (Nunes et al., 2008), we also encourage the development of studies on long-term
765 sediment transport in North African basins, in connection with changes in forcing factors.

766 **Acknowledgements**

767 The authors warmly thank Miss Abda Leila from the Hydrometry office, Mr Ould Lamara
768 Arezki from the Climatology Office of the Agence Nationale des Ressources Hydrauliques
769 (ANRH) in Alger, Mr Boudalia Mohamed from the Hydrometry Office of the ANRH in Oran,
770 and all the staff of ANRH who participated to the field measurements at Ain Hamara station.

771 They also acknowledge Mr Abderrezak Kamel Toubal for his help in drawing Figures 1 and
772 2. Two anonymous reviewers are warmly thanked for their reviews and comments on
773 previous versions of this paper. The editor, Efrat Morin, is gratefully acknowledged. Emma
774 Rochelle-Newall, a native English speaker, is warmly thanked for English corrections.

775

776 **References**

- 777 Aalto, R., Dunne, T., and Guyot, J.L.: Geomorphic controls on Andean denudation rates, The
778 Journal of Geology, 114, 85-99, 2006.
- 779 Achite, M., and Ouillon, S.: Suspended sediment transport in a semiarid watershed, Wadi
780 Abd, Algeria (1973-1995), J. Hydrol., 343 (3-4), 187-202, 2007.
- 781 Alexandrov, Y., Laronne, J.B., and Reid, I.: Suspended sediment concentration and its
782 variation with water discharge in a dryland ephemeral channel, northern Negev, Israel,
783 J. Arid Envir., 53, 73-84, 2003.
- 784 Asselman, N.E.M.: Fitting and interpretation of sediment rating curves, J. Hydrol., 234, 228-
785 248, 2000.
- 786 Bakreti, A., Braud, I., Leblois, E., and Benali, A.: Analyse conjointe des régimes
787 pluviométriques et hydrologiques dans le bassin de la Tafna (Algérie Occidentale),
788 Hydrol. Sci. J., 58 (1), 133-151, 2013.
- 789 Bates, B.C., Kundzewicz Z.W., Wu S., and Palutikof J.P., Eds.: Climate Change and Water.
790 Technical Paper of the Intergovernmental Panel on Climate Change, IPCC Secretariat,
791 Geneva, 210 pp., 2008
- 792 Benhamiche, N., Madani, K., and Laignel, B.: Chapter 12 Impact of climate change on water
793 resources in Algeria, in Behnassi M. et al. (eds), *Vulnerability of agriculture, water and*
794 *fisheries to climate change: toward sustainable adaptation strategies*, Springer, 193-
795 205, doi:10.1007/978-94-017-8962-2_12, 2014.
- 796 Beven, K.: Runoff generation in semi-arid areas. In Bull L.J., Kirkby M.J. (ed), 2002. Dryland
797 rivers: hydrology and geomorphology of semi-arid channels, Wiley, London, pp 57-
798 106, 2002.
- 799 Coulthard, T.J., Ramirez, J., Fowler, H.J., and Glenis, V.: Using the UKCP09 probabilistic
800 scenarios to model the amplified impact of climate change on river basin sediment
801 yield, Hydrol. Earth Syst. Sci., 16, 4401-4416, 2012.
- 802 Dahmani, A., and Meddi, M.: Climate variability and its impact on water resources in the
803 catchment area of the Wadi Fekan Wilaya of Mascara (West Algeria), Eur. J. Sci. Res.,
804 36 (3), 458-472, 2009.
- 805 Das, T., Maurer, E.P., Pierce, D.W., Dettinger, M.D., and Cayan, D.R.: Increases in flood
806 magnitudes in California under warming climates, J. Hydrol., 501, 101-110, 2013.

807 Déry, S.J., Stieglitz, M., McKenna, E.C., and Wood, E.F.: Characteristics and trends of river
808 discharge into Hudson, James, and Ungava bays, 1964–2000, *J. Climate*, 18, 2540–
809 2557, 2005.

810 Durand, N., Fiandrino, A., Fraunie, P., Ouillon, S., Forget, P., and Naudin, J.J.: Suspended
811 matter dispersion in the Ebro ROFI: an integrated approach, *Cont. Shelf Res.*, 22, 267–
812 284, 2002.

813 El Mahi, A., Meddi, M., and Bravard, J.P.: Analyse du transport solide en suspension dans le
814 bassin versant de l'Oued El Hammam (Algérie du Nord), *Hydrol. Sci. J.*, 57 (8), 1642–
815 1661, 2012.

816 Gao, Z.L., Fu, Y.L., Li, Y.H., Liu, J.X., Chen, N., and Zhang, X.P.: Trends of streamflow,
817 sediment load and their dynamic relation for the catchments in the middle reaches of the
818 Yellow River over the past five decades, *Hydrol. Earth Syst. Sciences*, 16, 3219–3231,
819 2012.

820 Ghenim, A.N., and Megnounif, A.: Analyse des précipitations dans le Nord-Ouest algérien,
821 *Sécheresse*, 24, 107–114, 2013a.

822 Ghenim, A.N., and Megnounif, A.: Ampleur de la sécheresse dans le bassin d'alimentation du
823 barrage Meffrouche (Nord-Ouest de l'Algérie), *Physio-Géo*, 7, 35–49, 2013b.

824 GISTEMP Team: GISS Surface Temperature Analysis (GISTEMP). NASA Goddard Institute
825 for Space Studies. Dataset accessed 2015-12-08 at <http://data.giss.nasa.gov/gistemp/>

826 Giuntoli, I., Renard, B., Vidal, J.P., and Bard, A.: Low flows in France and their relationship
827 to large-scale climate indices, *J. Hydrol.*, 482, 105–118, 2013.

828 Gomez, B., Cui, Y., Kettner, A.J., Peacock, D.H., and Syvitski, J.P.M.: Simulating changes to
829 the sediment transport regime of the Waipaoa River, New Zealand, driven by climate
830 change in the twenty-first century, *Global Planet. Change*, 67, 153–166, 2009.

831 Goudie, A.S.: Global warming and fluvial geomorphology, *Geomorphology*, 79, 384–394,
832 2006.

833 Hallouz, F., Meddi, M., and Mahé, G.: Modification du régime hydroclimatique dans le bassin
834 de l'oued Mina (Nord-Ouest d'Algérie), *Revue Sci. Eau*, 26 (1), 33–38, 2013.

835 Hancock, G.R.: A catchment scale assessment of increased rainfall and storm intensity on
836 erosion and sediment transport for Northern Australia, *Geoderma*, 152, 350–360, 2009.

837 Hancock, G.R., and Coulthard, T.J.: Channel movement and erosion response to rainfall
838 variability in southeast Australia, *Hydrol. Process.*, 26, 663–673, 2011.

839 Hansen, J., Ruedy, R., Sato, M., and Lo, K.: Global surface temperature change, *Rev.*
840 *Geophysics*, 48, RG4004, doi:10.1029/2010RG000345, 2010.

841 Houyou Z., Biolders, C.L., Benhorma, H.A., Dellal A., and Boutemdjet, A.: Evidence of
842 strong land degradation by wind erosion as a result of rainfed cropping in the Algerian
843 steppe: a case study at Laghouat, *Land Degrad. Develop.*, doi:10.1002/ldr.2295, 2014.

844 Iadanza, C., and Napolitano, F.: Sediment transport time series in the Tiber River, *Physics*
845 *Chem. Earth*, 31, 1212-1227, 2006.

846 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I
847 to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change;
848 Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y.
849 Xia, V. Bex and P.M. Midgley (eds.). Cambridge University Press, Cambridge, United
850 Kingdom and New York, NY, USA, 1535 pp., 2013.

851 Jerolmack, D.J., and Paola, C.: Shredding of environmental signals by sediment transport,
852 *Geophys. Res. Lett.*, 37, L19401, doi:10.1029/2010GL044638, 2010.

853 Jones, P.D., Lister, D.H., Osborn, T.J., Harpham, C., Salmon, M., and Morice, C.P.:
854 Hemispheric and large-scale land surface air temperature variations: an extensive
855 revision and an update to 2010. *J. Geophys. Res.*, 117, D05127.
856 doi:10.1029/2011JD017139, 2012.

857 Kassoul, M., Abdelgader, A., and Belorgey, M.: Caractérisation de la sédimentation des
858 barrages en Algérie. *Rev. Sci. Eau*, 3, 339-358, 1997.

859 Knight, J., and Harrison, S.: The impacts of climate change on terrestrial Earth surface
860 systems, *Nature Climate Change*, 3, 24-29, 2013.

861 Langbein, W.B., and Schumm, S.A.: Yield of sediment in relation to mean annual
862 precipitation, *Trans. Amer. Geophys. Union*, 39, 1076-1084, 1958.

863 Lockwood, M., and Fröhlich, C.: Recent oppositely directed trends in solar climate forcings
864 and the global mean surface air temperature, *Proc. R. Soc. A*,
865 doi:10.1098/rspa.2007.1880, 2007.

866 Louamri, A., Mebarki, A., and Laignel, B.: Variabilité interannuelle et intra-annuelle des
867 transports solides de l'Oued Bouhamdane, à l'amont du barrage Hammam Debagh
868 (Algérie orientale), *Hydrol. Sci. J.*, 58:7, 1559-1572, 2013.

869 Lu, X.X., Ran, L.S., Liu, S., Jiang, T., Zhang, S.R., and Wang, J.J.: Sediment loads response
870 to climate change: A preliminary study of eight large Chinese rivers, *Int. J. Sed. Res.*,
871 28, 1-14, 2013.

872 Mahé, G., and Paturel, J.E.: 1896-2006 Sahelian annual rainfall variability and runoff increase
873 of Sahelain Rivers, *C.R. Geoscience*, 314, 538-546, 2009.

- 874 Mahé, G., Lienou, G., Descroix, L., Bamba, F., Paturel, J.E., Laraque, A., Meddi, M.,
875 Habaieb, H., Adeaga, O., Dieulin, C., Chahnez Kotti, F., and Khomsi, K.: The rivers of
876 Africa: witness of climate change and human impact on the environment, *Hydrol. Proc.*,
877 27, 2105-2114, 2012.
- 878 Meade, R.H., and Moody, J.A.: Causes for the decline of suspended-sediment discharge in the
879 Mississippi River system, 1940-2007, *Hydrol. Proc.*, 24, 35-49, 2010.
- 880 Meddi, M., and Hubert, P.: Impact de la modification du régime pluviométrique sur les
881 ressources en eau du nord-ouest de l'Algérie, in: *Hydrology of the Mediterranean and*
882 *Semiarid Regions*, IAHS Publ. n°278, 229-235, 2003.
- 883 Meddi, H., and Meddi, M.: Variabilité spatiale et temporelle des précipitations du Nord-Ouest
884 de l'Algérie, *Geographica Technica* n°2/2007, 49-55, 2007.
- 885 Meddi, H., and Meddi, M.: Variabilité des précipitations annuelles du Nord-Ouest de
886 l'Algérie, *Sécheresse*, 20 (1), 57-65, 2009.
- 887 Meddi, M., and Talia, A.: Pluviometric regime evolution in the North of Algeria. *Arab Gulf J.*
888 *Sci. Res.*, 26 (3), 152-162, 2008.
- 889 Meddi, M., Talia, A., and Martin, C.: Evolution récente des conditions climatiques et des
890 écoulements sur le bassin versant de la Macta (nord-ouest de l'Algérie), *Physio-Géo*, 3,
891 61-84, 2009.
- 892 Megnounif, A., Terfous, A., and Ouillon, S.: A graphical method to study suspended sediment
893 dynamics during flood events in the Wadi Sebdo, NW Algeria (1973-2004), *J. Hydrol.*,
894 497, 24-36, 2013.
- 895 Memariam, H., Balasundram, S.K., Talib, J.B., Sood, A.M., and Abbaspour, K.C.: Trend
896 analysis of water discharge and sediment load during the past three decades of
897 development in the Langat basin, Malaysia, *Hydrol. Sci. J.*, 57 (6), 1207-1222, 2012.
- 898 Milliman, J.D., Farnsworth, K.L., Jones, P.D., Xu, K.H., and Smith, L.C.: Climatic and
899 anthropogenic factors affecting river discharge to the global ocean, 1951-2000, *Global*
900 *Planet. Change*, 62, 187-194, 2008.
- 901 Morgan R.P.C., Quinton, J.N., Smith R.E., Govers, G., Poesen, J.W.A., Auerswald, K., Chisci
902 G., Torri, D., and Styczen, M.E.: The European Soil Erosion Model (EUROSEM): A
903 dynamic approach for predicting sediment transport from fields and small catchments,
904 *Earth Surf. Process. Landforms*, 23, 527-544, 1998.
- 905 Nanson, G.C., Tooth, S., and Knighton, A.D.: A global perspective on dryland rivers:
906 perceptions, misconceptions and distinctions. In Bull L.J., Kirkby M.J. (ed), 2002.

907 Dryland rivers: hydrology and geomorphology of semi-arid channels, Wiley, London.
908 pp 17-54, 2002.

909 Neitsch, S.L., Arnold, J.G., Kiniry J.R., and Williams, J.R.: Soil and Water Assessment Tool
910 – Theoretical Documentation version 2009. Texas Water Inst. Techn. Report n°406,
911 Texas A&M University, College Station, 2011.

912 Nunes, J.P., Seixas, J., and Pacheco, N.R.: Vulnerability of water resources, vegetation
913 productivity and soil erosion to climate change in Mediterranean watersheds, *Hydrol.*
914 *Process.*, 22, 3115–3134, 2008.

915 Oliver, J.E.: Monthly precipitation distribution: a comparative index, *Prof. Geogr.*, 32 (3),
916 300–309, 1980.

917 O’Neal, M.R., Nearing, M.A., Vining, R.C., Southworth, J., and Pfeifer, R.A.: Climate
918 change impacts on soil erosion in Midwest United States with changes in crop
919 management, *Catena*, 61, 165–184, 2005.

920 Osborn, T.J., and Jones, P.D.: The CRUTEM4 land-surface air temperature dataset:
921 construction, previous versions and dissemination via Google Earth, *Earth System*
922 *Science Data* 6, 61-68, 2014.

923 Ouillon, S.: Erosion and sediment transport: width and stakes, *La Houille Blanche*, 53 (2), 52-
924 58, 1998.

925 Paturel, J.E., Ouedraogo, M., Servat, E., Mahé, G., Dezetter, A., and Boyer, J.F.: The concept
926 of rainfall and stream flow normals in West and Central Africa in a context of climatic
927 variability, *Hydrol. Sci. J.*, 48 (1), 125-137, 2003.

928 Poesen J., Nachtergaele, J., Verstraeten, G., and Valentin, C.: Gully erosion and
929 environmental change: importance and research needs, *Catena*, 50, 91-133, 2003.

930 Pruski, F.F., and Nearing, M.A.: Climate-induced changes in erosion during the 21st century
931 for eight U.S. locations, *Water Resour. Res.*, 38, 1298, doi:10.1029/2001WR000493,
932 2002.

933 Remini B., Leduc C., and Hallouche, W.: Evolution des grands barrages en régions arides:
934 quelques exemples algériens, *Sécheresse*, 20 (1), 96-103, 2009.

935 Riebe, C.S., Kirchner, J.W., Granger, D.E., and Finkel, R.C.: Minimal climatic control on
936 erosion rates in the Sierra Nevada, California, *Geology*, 29, 447-450, 2001.

937 Sari, A.: Hydrologie de surface, cours d’initiation, Editions Distribution Houma, 155 pp,
938 2009.

939 Searcy, J.K., and Hardison, C.H.: Double-mass curves, USGS Water-Supply Paper 1541-B,
940 1960.

941 Sen, P. K.: Estimates of regression coefficients based on Kendall's tau, *J. Am. Stat. Assoc.*,
942 63, 1379–1389, 1968.

943 Shrestha, B., Babel, M.S., Maskey, S., van Griensven, A., Uhlenbrook, S., Green, A., and
944 Akkharath, I. Impact of climate change on sediment yield in the Mekong River basin: a
945 case study of the Nam Ou basin, Lao PDR, *Hydrol. Earth System Sciences*, 17, 1-20,
946 2013.

947 Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L.M., van Lanen, H.A.J., Sauquet, E., Demuth,
948 S., Fendekova, M., and Jodar, J.: Streamflow trends in Europe: Evidence from a dataset
949 of near-natural catchments, *Hydrol. Earth Syst. Sci.*, 14, 2367–2382, 2010.

950 Summerfield, M.A., Hulton, N.J.: Natural controls of fluvial denudation rates in major world
951 drainage basins, *J. Geophys. Res.*, 99 B7, 13871-13883, 1994.

952 Syvitski, J.P.M.: Supply and flux of sediment along hydrological pathways: research for the
953 21st century, *Global Planet. Change*, 39, 1-11, 2003.

954 Syvitski, J.P.M., and Milliman, J.D.: Geology, geography, and humans battle for dominance
955 over the delivery of fluvial sediment to the coastal ocean, *The Journal of Geology*, 115,
956 1-19, 2007.

957 Taabni, M., and El Jihad, M.D.: Eau et changement climatique au Maghreb: quelles stratégies
958 d'adaptation?, *Les Cahiers d'Outre-Mer*, 260, 493-518, 2012.

959 Taibi, S., Meddi, M., Mahé, G., and Assani, A.: Variability of annual and extreme rainfall
960 over Northern Algeria and relationship with teleconnections patterns, *Proc. of the*
961 *Mediterranean meeting on "Monitoring, modeling and early warning of extreme events*
962 *triggered by heavy rainfalls"*, Univ. of Calabria, Cosenza, Italy, June 26-28, 2014.

963 Tebbi, F.Z., Dridi, H., and Morris, G.L.: Optimization of cumulative trapped sediment curve
964 for an arid zone reservoir: Foum El Kherza (Biskra, Algeria), *Hydrol. Sci. J.*, 57 (7),
965 1368-1377, 2012.

966 Temme, A.J.A.M., and Veldkamp, A.: Multi-process Late Quaternary landscape evolution
967 modelling reveals lags in climate response over small spatial scales, *Earth Surf. Proc.*
968 *Land.*, 34, 573–589, 2009.

969 Tecsalt International: Etude de la protection des bassins versants de l'ensemble des barrages
970 en études d'avant projet détaillé (lots n°1 à 4) et des bassins versants des barrages en
971 exploitation (lots n° 5 et 6), Phase I description du milieu, lot n°5: barrages Gargar et
972 S.M. Ben-Aouda, Ministère des Ressources en Eau, Algérie, 2004.

973 Tockner, K., and Stanford, J.A.: Riverine flood plains: present state and future trends,
974 *Environ. Conserv.*, 29, 308–330, 2002.

- 975 Touaibia, B.: Problématique de l'érosion et du transport solide en Algérie septentrionale,
976 Sécheresse, 21 (1e), 1-6, 2010.
- 977 [Touat, S.: Contrôle de la représentativité de l'échantillon des transports solides en suspension.](#)
978 [Revue Eaux et Sols d'Algérie, 3, 48-54, 1989.](#)
- 979 Trambly, Y., El Adlouni, S., and Servat, E.: Trends and variability in extreme precipitation
980 indices over Maghreb countries. *Natural Hazards Earth Syst. Sci.*, 13, 3235-3248, 2013.
- 981 Tucker, G.E., and Slingerland, R.: Drainage basin responses to climate change, *Water Resour.*
982 *Res.*, 33, 2031–2047, 1997.
- 983 Vanmaercke, M., Poersen, J., Broeckx, J., and Nyssen, J.: Sediment yield in Africa, *Earth-*
984 *Science Rev.*, 136, 350-368, 2014.
- 985 Vinh, V.D., Ouillon, S., Thanh, T.D., and Chu, L.V.: Impact of the Hoa Binh dam (Vietnam)
986 on water and sediment budgets in the Red River basin and delta, *Hydrol. Earth System*
987 *Science*, 18, 3987-4005, 2014.
- 988 Vörösmarty, C.J., Meybeck, M., Fekete, B., Sharma, K., Green, P., and Syvitski, J.P.M.:
989 Anthropogenic sediment retention: major global impact from registered river
990 impoundments, *Global Planet. Change*, 39, 169-190, 2003.
- 991 Walling, D.E.: Assessing the accuracy of suspended sediment rating curves for a small basin,
992 *Water Res. Res.*, 13 (3), 531-538, 1977.
- 993 Walling, D.E.: The response of sediment yields to environmental change, *IAHS Publ.* 245,
994 77-89, 1997.
- 995 Walling, D.E.: Human impact on land-ocean sediment transfer by the world's rivers,
996 *Geomorphology*, 79, 192-216, 2006.
- 997 Walling, D.E.: The impact of Global Change on Erosion and Sediment Transport by Rivers:
998 Current progress and Future Challenges, *The United Nations World Water*
999 *Development Report 3*, UNESCO, Paris, 26 p., 2009.
- 1000 Walling, D.E., and Fang, D.: Recent trends in the suspended sediment loads of the world's
1001 rivers. *Global Planet. Change*, 39, 111–126, 2003.
- 1002 Wang, H., Yang, Z., Saito, Y., Liu, J., and Sun, X.: Interannual and seasonal variation of the
1003 Huanghe (Yellow River) water discharge over the past 50 years: connections to impacts
1004 from ENSO events and dams, *Global Planet. Change*, 50, 212–225, 2006.
- 1005 Wang, H.J., Yang, Z.S., Saito, Y., Liu, J.P., Sun, X.X., and Wang, Y.: Stepwise decreases of
1006 the Huanghe (Yellow River) sediment load (1950-2005): Impacts of climate change and
1007 human activities, *Global Planet. Change*, 57 (3-4), 331-354, 2007.

- 1008 Ward, P.J., Eisner, S., Flörke, M., Dettinger, M.D., and Kummu, M.: Annual flood
1009 sensitivities to El Niño-Southern Oscillation at the global scale, *Hydrol. Earth Syst. Sci.*,
1010 18, 47-66, 2014.
- 1011 Ziegler, A.D., Sheffield, J., Maurer, E.P., Nijssen, B., Wood, E.F., and Lettenmaier, D.P.:
1012 Detection of intensification in global- and continental-scale hydrological cycles:
1013 Temporal scale of evaluation. *J. Climate*, 16, 535–547, 2003.

1014 **Table 1.** General statistics onf the yearly averages of hydrologic parameters fromof the Wadi
 1015 Abd at Ain Hamara gauging station over the period 1970-2010 (Note: T at Chlef was
 1016 estimated from measurements at Dar El Beida and Miliana for 20 months over 480)
 1017

| Sstatistic-value | T (Chlef) °C | P mm yr ⁻¹ | Q m ³ s ⁻¹ | Q _w m ³ s ⁻¹ | M 10 ³ t yr ⁻¹ | SPM* g L ⁻¹ |
|--------------------|-----------------|--------------------------|-------------------------------------|--|---|---------------------------|
| Mean | 19.09 | 264 | 1.18 | 1.29 | 564 | 12.3 |
| Min | 17.52 | 165 | 0.37 | 0.46 | 33.1 | 2.56 |
| (Year) | 1971-72 | 1999-00 | 1992-93 | 1983-84 | 1992-93 | 1975-76 |
| Max | 20.32 | 506 | 2.19 | 2.98 | 3266 | 50.25 |
| (Year) | 1989-90 | 1995-96 | 1994-95 | 1994-95 | 2007-08 | 2007-08 |
| standard deviation | 0.69 | 71.2 | 0.52 | 0.59 | 696 | 10.6 |
| CV (%) | | 27.0 | 44.4 | 45.6 | 123.3 | 86.0 |

1018
 1019

Table 2. General statistics of the averages of hydrologic parameters (averages) from the Wadi Abd at Ain Hamara gauging station per decade and for the entire significant period from 1970-2010 (Note: T at Chlef was estimated from measurements at Dar El Beida and/or Miliana for 20 months with missing values over 480)

| Period | T at Chlef | P, yearly precipitation | | NDD, average yearly number of dry days (Q=0) | Q, yearly discharge | | Q _w , yearly discharge of wet days | | Q _s , yearly sediment load | | Q ₉₈ , average of yearly values | | SSY, average specific sed. yield (t km ⁻² yr ⁻¹) | SPM* | | Rating curve parameters | | | |
|-----------|--------------|-------------------------|--------|--|---|--------|---|--------|--|--------|--|--------|---|------------------------------|--------|-------------------------|-------|----------------|------|
| | Average (°C) | Average (mm) | CV (%) | | Average (m ³ s ⁻¹) | CV (%) | Average (m ³ s ⁻¹) | CV (%) | Average (10 ³ tons yr ⁻¹) | CV (%) | Average (m ³ s ⁻¹) | CV (%) | | Average (g L ⁻¹) | CV (%) | a | b | R ² | N |
| 1970-2010 | 19.09 | 264.10 | 27.0 | 28.3 | 1.18 | 44.4 | 1.29 | 45.7 | 564 | 123.3 | 9.18 | 78.6 | 227.6 | 12.3 | 86.0 | 2.270 | 0.647 | 0.431 | 1213 |
| 1970-1980 | 18.32 | 310.53 | 19.4 | 1.2 | 1.16 | 32.9 | 1.16 | 32.9 | 180 | 78.8 | 4.37 | 66.9 | 72.7 | 4.54 | 47.9 | 1.021 | 0.890 | 0.573 | 240 |
| 1980-1990 | 19.19 | 231.23 | 16.8 | 24.1 | 0.98 | 36.8 | 1.07 | 41.5 | 334 | 91.7 | 7.39 | 68.0 | 134.5 | 9.93 | 57.0 | 2.049 | 0.649 | 0.449 | 316 |
| 1990-2000 | 19.37 | 250.42 | 40.5 | 59.9 | 1.13 | 55.1 | 1.34 | 55.2 | 614 | 98.3 | 11.03 | 88.5 | 247.5 | 14.36 | 69.2 | 2.753 | 0.659 | 0.418 | 343 |
| 2000-2010 | 19.49 | 264.22 | 19.7 | 28.1 | 1.45 | 43.3 | 1.57 | 42.2 | 1130 | 90.3 | 13.94 | 44.5 | 455.6 | 20.55 | 68.7 | 4.440 | 0.412 | 0.384 | 324 |
| 1970-1985 | 18.51 | 284.34 | 23.1 | 0.8 | 1.02 | 37.8 | 1.02 | 38.2 | 159 | 78.9 | 4.13 | 58.8 | 64.2 | 5.16 | 58.9 | 1.213 | 0.818 | 0.519 | 346 |
| 1985-2010 | 19.47 | 251.96 | 29.0 | 44.8 | 1.28 | 45.1 | 1.45 | 43.7 | 808 | 97.0 | 12.21 | 61.1 | 325.6 | 16.65 | 67.4 | 2.974 | 0.576 | 0.415 | 867 |

Table 3. Variation of precipitation, water discharge and sediment yield averaged per season over each decade

| | Precipitation (mm) | | | | Water discharge (m ³ s ⁻¹) | | | | Sediment yield (10 ³ tons) | | | |
|-----------|--------------------|--------|--------|--------|---|--------|--------|--------|---------------------------------------|--------|--------|--------|
| | autumn | winter | spring | summer | autumn | winter | spring | summer | autumn | winter | spring | summer |
| 1970-1980 | 68.5 | 102.6 | 128.2 | 11.2 | 1.26 | 1.38 | 1.41 | 0.58 | 62.2 | 43.7 | 66.1 | 8.4 |
| 1980-1990 | 56.0 | 94.4 | 70.7 | 10.1 | 1.15 | 1.29 | 1.08 | 0.40 | 128.8 | 61.0 | 97.2 | 46.8 |
| 1990-2000 | 67.0 | 81.1 | 86.9 | 15.5 | 1.86 | 0.99 | 1.11 | 0.54 | 279.1 | 57.8 | 130.9 | 146.0 |
| 2000-2010 | 78.6 | 98.4 | 77.7 | 9.5 | 3.04 | 1.35 | 1.06 | 0.35 | 804.9 | 94.4 | 195.3 | 35.4 |

| | Precipitation (%) | | | | Water discharge (%) | | | | Sediment yield (%) | | | |
|-----------|-------------------|--------|--------|--------|---------------------|--------|--------|--------|--------------------|--------|--------|--------|
| | autumn | winter | spring | summer | autumn | winter | spring | summer | autumn | winter | spring | summer |
| 1970-1980 | 22.1 | 33.0 | 41.3 | 3.6 | 27.3 | 29.8 | 30.3 | 12.6 | 34.5 | 24.2 | 36.6 | 4.7 |
| 1980-1990 | 24.2 | 40.8 | 30.6 | 4.4 | 29.4 | 32.8 | 27.7 | 10.1 | 38.6 | 18.3 | 29.1 | 14.0 |
| 1990-2000 | 26.7 | 32.4 | 34.7 | 6.2 | 41.3 | 22.1 | 24.6 | 12.0 | 45.5 | 9.4 | 21.3 | 23.8 |
| 2000-2010 | 29.7 | 37.3 | 29.4 | 3.6 | 52.5 | 23.2 | 18.2 | 6.1 | 71.2 | 8.4 | 17.3 | 3.1 |

Figure Captions

Fig. 1 Location of the Wadi Abd sub-basin within the Mina and Cheliff basins, and the other main basins of Algeria

Fig. 2 The Wadi Abd catchment area. (a) Rain and hydrometric stations including HS1 at Takhmaret and HS2 at Ain Hamara, (b) Geology, (c) Slopes from the Digital Elevation Model of North Algeria, (d) Vegetation cover from Landsat ETM+ data of 2009

Fig. 3 Linear erosion forms in the Wadi Abd basin. (a) and (e) Gullying (depth: 30-50 cm, width < 1 m), (c) and (d) Gully erosion (depth: 50-200 cm), (b) and (f) Interrill and rill erosion

Fig. 4 Relationships between mean annual temperatures at the three stations of Dar El Beida, Miliana and Chlef (from CRUTEM4)

Fig. 5 Comparison between estimates of Q_s obtained from Q and the global rating curve, and measured Q_s

Fig. 6 Interannual variations of mean yearly temperature (calculated from September to August monthly temperatures) at three stations in northern Algeria: Dar El Beida, Miliana, Chlef (from measurements of CRUTEM4 only, extrapolated values are not shown)

Fig. 7 Interannual variations of annual precipitation, water discharge and sediment yield at Ain Hamara station

Fig. 8 Variation of hydrological regime with annual % of time of flowing water, Q_{98} (amongst daily discharges, per year) and annual standard deviation of daily river discharge

Fig. 9 Double mass plot of sediment yield versus water flow

Fig. 10 Yearly average of related sediment load parameters vs intra-annual variability of daily river discharge, characterized by their annual standard deviation. (a) SPM^* , (b) Specific sediment yield

Fig. 11 Trends of the seasonal indexes of precipitation (a), discharge (b) and (c) sediment discharge in the Wadi Abd basin.

Fig. 12 Monthly values of precipitation (a), Q (b) and Q_s (c) averaged over decades in the Wadi Abd basin.

Fig. 13 Monthly values of precipitation averaged over 6 stations, for the two periods: 1970-1985 and 1985-2010.

Fig. 14 Monthly values of the ratio Q/P averaged over decades

Fig. 15 Variations of SPM^* against the global mean temperature anomaly (from GISTEMP Team, 2015)

Fig. 16 Relationship between the rating curves parameters averaged over 15 years

Fig. 17 Relationship between the rating curve parameter a averaged over 15 years and the averaged values of specific sediment yield over 15 years




Fig. 1





Fig. 2





Fig. 2 (continued)




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10a




Fig. 10b




Fig. 11




Figure 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17