

Reviewer 1

While the authors have done a nice job in revision, there is still a lack of a clear central research question or gap being filled. While the presentation has improved, I still miss a clear statement at the end of the introduction motivating what the goal of the study is. P4L29-P5L29 is poorly written and does not address this concern. This is a critical need that must be addressed. This was a weakness in the first submission and it remains a fundamental shortcoming. The current study characterizes some variability and applies a model. These alone (without a strong context and motivation) are likely of limited interest to the readers of HESS where there is an expectation to see advancements of process understanding and fundamental hydrological understanding.

As a general comment, the language gets rather loose throughout the text. The terminology and presentation must be improved and tightened. Too often are blanket statements or generalizations made without clear support. The language is often not precise and rather vague. I have tried to target several examples in the minor comments section of this review, but I am sure there are more. Please improve.

Further, the study needs streamlined (this goes hand-in-hand with the comment regarding the main question). The use of a hydrologic model is a good addition to the manuscript. However, the presentation of the modeling setup (section 2.5) leaves something to be desired. Rather than lengthen an already long manuscript, I would suggest shortening the text in the manuscript and adding an extended appendix or supplementary text section. This will allow you to list equations, have schematics and cover in detail the calibration/validation aspects of the modeling. As it is currently presented, the modeling is not fully described at a level of detail needed to understand the steps to create a working model. Rather than expanding this in the manuscript, I am suggesting to expand it in an appendix. If the central question for the study is to use the chemical data to improve the modeling, then much more is needed with regards to presenting the modeling without considering the tracer data in comparison with the model results considering the tracer data (and more information would be needed on exactly how the data were considered).

In connection with the two previous general comments, I suggest also moving the methods and results text pertaining to the travel time modeling into an appendix or supplementary text section. As they are currently presented, they come across as "extra" and just confuse the flow of the study. The level of detail is underdeveloped. Moving them to a separate text section would really help streamline the presentation of this study. This is something that is needed as the current presentation is overly complicated and gets confusing with all the information jumbled together. This will allow you to focus in on the central research question of the study.

Lastly, from my reading, the discussion section 4.3 appears to be the main discussion section. The other two subsections more or less are just reiterating the results. This section 4.3 (while well written) could do a better job relating to more current studies on mixing assumptions (both temporal and spatial). The authors start into that consideration (at P28L26-30) but do not go further. I suggest reducing the emphasis on sections 4.1 and 4.2 (which cover the empirical and site-specific aspects of the research) and breaking the section 4.3 up into smaller subsections that potentially relate back to central research question(s). Explore the literature on these mixing-storage topics over the past 5 years. Placing findings in that context will significantly raise the level of presentation.

Apologies for harping on the presentation and structuring. I think the study presents a nice set of data. The challenge to the authors is to leverage these data into a fundamental advancement with regards to catchment hydrology. Taking a critical view of the research question and the general manuscript structure (i.e. how the results answer the question and the discussion provides relevance) should help elucidate this aspect.

We thank Referee #1 for the comments which were very useful to improve the paper. We followed the reviewer's comments by strengthening and shortening the manuscript (by 7 pages) now having a stronger focus on process understanding. We rewrote the Introduction including our research

motivation. We further moved the Material and methods sections about MTT estimations and groundwater age modelling to Appendix I and II which also include paragraphs about the objective of our modelling approach, model setup, model's boundary conditions, parameters, and lists equations. We think that the applied groundwater model is fully described now. With our model we were able to get a rough estimate of residence times greater than the limit of stable water isotopes (>5 years). We split the Discussion section and included an additional section 4.4 to further discuss our modelling results with regard to current studies on mixing assumptions.

Minor comments

P1: J. Pferdmenges does not have an affiliation.

Corrected.

P2L1: "smoothed out" is not precise. Adjusted terminology.

Changed to "became less pronounced".

P2L27: Provide here a clear definition of what you mean with "low angle". Also, make it clear that "low angle" is different from "lowland" since that is the focus of the next few sentences.

There does not exist an international standard definition of a mountain (Goudie and Goudie, 2013) and thus, a low-mountainous terrain. The definition of mountain regions is largely arbitrary because multiple criteria can be used to define such areas, e.g. relative relief, threshold altitude (1000 m) etc. (Perry and Taylor, 2009). Following Perry and Taylor (2009) a hilly terrain, which we equate with a low-mountainous terrain, has an altitudinal difference of 50-100 m (over 5 km distance) which applies to our catchment. Thus we included the following: "altitudinal difference of 50–100 m over 5 km distance (Perry and Taylor, 2009)".

P2L27: Remove "very early,"

Followed reviewer's suggestion.

P3L22: Again, see work by van der Velde ([Van der Velde Y, Torfs PJJF, van der Zee SEATM, Uijlenhoet R. 2012. Quantifying catchment-scale mixing and its effects on time-varying travel time distributions. Water Resources Research 48: W06536])

We included the reference in Discussion section 4.4.

P4L10: "Following the way"? This is vague and confusing.

We deleted the whole section.

P4L29-P5L29: This section is poorly written and disconnected from the rest of the introduction. Parts seem like methods. It should be removed since it adds no value to presenting the study. Rather, end the introduction here with a simple statement of the gap in research being targeted and the research question(s) answered. Be concise and clear to motivate the research.

We removed most of this section and rewrote the Introduction including a research motivation.

P8L5: Change "was fallen" to "fall"

Followed reviewer's suggestion.

P8L19: Change "was" to "were"

Followed reviewer's suggestion.

P8L28: "as connecting compartment"? Vague and confusing.

We deleted "as connecting compartment between precipitation and groundwater".

P10L10: I think it should be "connection" rather than "connectivity". Connectivity has taken on special meaning in the literature that is not consistent with how it is necessarily being used here.

We changed it to "connection" throughout the manuscript.

P11L9: Check the significant figures in your numbers.

Edited in Appendix II.

P15L28: Perhaps you mean “proposed”?

Edited accordingly.

P16L4: “higher than further below” is confusing. Re-write.

Edited as follows: “are higher than in the subsoil”.

P18L13-L23: This should be its own results subsection and not presented with the section on stream water variability. Further, perhaps consider taking this and the methods text describing this approach into their own stand-alone appendix or supplementary text section. This would really help streamline the presentation.

We moved the methods text to Appendix I. However, we disagree with the reviewer that the respective lines should be moved to their own results section as this would add additional text to an already long manuscript. We also think that the results are not relevant enough to justify their own subsection. We therefore kept this paragraph under heading 3.3. However, we restructured the Discussion section and added a section on “Water age dynamics”.

P19L12-13: Confusing sentence. Re-write.

We rephrased the sentence as follows: “Due to different water flow paths of groundwater along the studied stream, we expected to find distinguished groundwater isotopic signatures.”

P20L28: Change “:” to “.”

1 Edited accordingly.

2

3 **Reviewer 2**

4 Review of Revision: Exploring water cycle dynamics by sampling multiple stable water isotope pools
5 in a small developed landscape of Germany by Orlowski et al. in HESS

6

7 The authors improved several aspects of the paper with this revision and accounted for three of my
8 main concerns:

9 - They use a rigorous statistical methodology for analysing their stable isotope data

10 - They present the results from transfer function analysis of stable isotope data to determine
11 catchment transit times (no good fit achieved- which is also a result)

12 - And restructured the paper, eventually increasing readability

13 The paper is well written. Yet, I don't think the paper is ready to be published, and still requires some
14 work. Besides the changes and the rework the introduction does not sufficiently frame the research
15 question. The authors outline four objectives: 1. Linkages between water cycle components, 2.
16 investigate transformation of precipitation to soil and groundwater, 3. Analysis of landscape
17 characteristics On soil water isotopic composition and 4. estimate groundwater ages and flow
18 directions... via ... model.

19 While these are definitely interesting questions to ask, the “why” remains unclear. Several times the
20 other use the term “poorly understood” in the introduction, while it is never clear to me, what
21 exactly is poorly understood. For example, the authors claim that groundwater-surface water
22 interactions and mechanism that explain the release of old water to streams fall in this category. I
23 think there exist a plurality of work for both problems; so why not pinpointing on what exactly is not
24 understood? For me, this needs some more attention. The big question: What brings this work
25 beyond the local case study?

26 Moreover, the use of the CMF model seems a little bit weak. As there is no calculation of streamflow
27 age, nor any validation of the modelled ages, it seems somewhat useless to me.

28 What would generally be interesting, seeing the claimed decoupling of precipitation with
29 groundwater and streamflow, is the calculation of the isotopic signal in groundwater recharge. Can

30 the soil profiles, and meteorological data combined with a simple approach be used to show that the
31 soils modify the precipitation isotope signal in a way that it is consistent with the values in stream
32 and groundwater. This would show that the system is not decoupled, but the soils behave in a
33 translator flow way, or at least as a filter for incoming precipitation. This could also underpin the
34 claim in the conclusion that it was shown “that groundwater was predominantly recharged during
35 winter”. I think this interpretation is right, but the proof is not convincingly presented. An alternative
36 could be that a larger groundwater system controls the streamflow.

37 We gratefully acknowledge the comments of the reviewer, which helped us to improve the
38 manuscript. In general, we shortened (by 7 pages) and streamlined the whole manuscript, rewrote
39 the Introduction focusing on our research motivation, and included new studies, e.g. by Hrachowitz
40 et al. (2016) or McDonnell and Beven (2014). By moving the Material and methods section about the
41 MTT estimations and our hydrological model to the Appendix, we were able to describe our
42 approach in more detail (objective of our modelling approach, model setup, model’s boundary
43 conditions, parameters, and equations). As mentioned in the manuscript, we estimated MTTs for the
44 two studied streams (Schwingbach and Vollnkirchener Bach) as well as mean residence times from
45 precipitation to groundwater comprising thirteen groundwater sampling points (therewith covering
46 water fluxes through the vadose zone). The improved discussion and streamlined structure should
47 cover the issues raised by the reviewer.

48

49 **Some minor comments:**

50 Title: “small” is rather subjective here with catchments of >3km² and ~10km²

51 We deleted “small”.

52 Abstract, L19 “decoupled” is a strange word for me in this perspective. As precipitation contributes
53 to groundwater, but with the soils acting as filter.

54 We replaced “decoupled” by “disconnected”.

55 P2L16: origion of? Formation of?

56 We revised the sentence as follows: “The application of stable water isotopes as natural tracers in
57 combination with hydrodynamic methods has been proven to be a valuable tool for studying the
58 origin and formation of recharged water as well as the interrelationship between surface water and
59 groundwater (Blasch and Bryson, 2007)...”

60 P3L23-28: this is not done yet?

61 We deleted and rephrased most of this paragraph.

62 P4L14: I think this sentence is grammatically wrong

63 We deleted “thus” from the sentence.

64 P4L18: Replace “This” with “The enrichment...”

65 Rephrased as follows: “The isotopic enrichment decreases exponentially with depth...”

66 P5L21: What is your hypothesis? What is the question that differentiates this work from a local case
67 study?

68 We shortened and rewrote the Introduction focusing on our research motivation.

69 P10L28ff. How is the model validated? How useful is it to leave out soil processes if they are the only
70 component where stable isotopes change? I am not convinced about the added value of the GW
71 model (and I don’t think it is needed for the work)

72 We added Appendix I and II to account for this. We included a subsection describing the objectives of
73 our groundwater modelling approach as follows: “Stable water isotopes are only a tool to determine
74 the residence time for a few years (McDonnell et al., 2010). In cases of longer residence times and a
75 strong mixing effect, seasonal variation of isotopes vanishes and results in stable flat lines of the
76 isotopic signal. To get a rough estimate of residence times greater than the limit of stable water

77 isotopes (>5 years), we split the water flow path in our catchment in two parts: the flow from
78 precipitation to groundwater, which was calculated via FlowPC and the longer groundwater
79 transport. The simplest method to estimate the residence time of groundwater transport is via the
80 storage-to-input-relation, with the storage as the aquifer size and the input as the groundwater
81 recharge time. However, this method ignores the topographic setting, and water input
82 heterogeneity. In our study we used a simplified groundwater flow model with tracer transport to
83 calculate the groundwater age dynamics. The numerical output of water ages cannot be validated
84 with the given isotope data, since the model is used to fill a residence time gap, where stable water
85 isotopes are not feasible to apply. The model is falsified however, if the residence time is short
86 enough (<5 years) to be calculable via FlowPC. Hence, the results of the groundwater age model
87 should be handled with care and only seen as the order of magnitude of flow time scales.”

88 P15L13ff. Now you mention the stream water in the precipitation results, this is a little bit strange,
89 since the stream results are not presented yet

90 We deleted this paragraph and shifted stream water results to the respective section (3.3).

91 P15L26: Insert “they” before “exhibited”

92 We deleted this sentence and moved the previous sentence to the Discussion section (4.2.1).

93 P15L27-30. These lines seem unnecessary, since repetitive.

94 We deleted this paragraph.

95 P18L20/21: This conclusion only related to Transfer function based methods.

96 We rewrote the sentence as follows: “Therefore, we conclude that transfer function based MTT
97 estimation methods applying stable water isotope data failed for the Schwingbach.”

98 P21L3: Any change for streamflow age information?

99 The model confirms that the stream contains water with different transit times. Therefore, the
100 stream water does not have a discrete age, but a distribution of ages due to variable flow paths,
101 which is also mentioned in the manuscript. MTTs for both streams were further calculated using
102 FlowPC and the groundwater model for different reaches of the Vollnkirchner Bach.

103 P21L5ff.: Since there was no objective in the introduction that relates to precipitation, this discussion
104 is somewhat oversized (but good)

105 In general, we shortened the length of the manuscript by 7 pages from the Abstract to the
106 Conclusions. We have also cut down the length of this section.

107 P23-25: Can you calculate the stable isotope composition of recharge from your data? That would be
108 very helpful. So you have a Hewlett and Hibbert like translator flow for GW recharge? Or is only
109 winter rainfall contributing?

110 We measured spatial isotopic uniformity of the subsurface soil water (chapter 3.2) similar to the
111 groundwater composition throughout the study area. We would not expect different values in the
112 groundwater recharge. Hence, variable source area effects are not detectable via the isotopic
113 composition in our case.

114 P14: How do you define baseflow? I would rather say, that the stream is always groundwater
115 dominated, but you also observe higher flows, which are not really baseflow situations. But again,
116 here the stream is groundwater fed.

117 In our catchment baseflow is defined as discharge $<10 \text{ L}\cdot\text{s}^{-1}$ (Orlowski et al., 2014), which is also
118 defined in the manuscript now.

119 P27-30: They work; they just show 100% groundwater contributions for the observed samples.
120 We rephrased the sentence as follows: “Just by comparing mean precipitation, stream, and
121 groundwater isotopic signatures (Table 1), it is obvious that simple mixing calculations do not work,
122 i.e. showing predominant groundwater contribution.”

123 P28L4: I am not fully happy with the term “disconnected”, as GW levels react instantaneously.

124 We rewrote the sentence and included “isotopically disconnected”.

125 P29L14ff. - So what is new? What is the beyond study area relevance?

126 - Show clearer that it is winter precipitation recharging

127 We have added some key points to the Conclusion section, especially the partitioning of the flow
128 path as an analytical tool.

129 P29L21-22: Delete “old water paradox” as this is repetitive and vague. There is a wide range of
130 processes that are behind this paradox. It would be better to clarify this functioning

131 Followed reviewer’s suggestion.

132

133 **Relevant changes**

134 • Shortened (by 7 pages) and streamlined the whole manuscript

135 • Restructured the manuscript: Revised the Introduction completely and moved the MTT
136 estimation and hydrological modelling methods parts to Appendix I and II

137 • Extended the description about the hydrological model including: Objectives, model setup,
138 boundary conditions, parameters, groundwater age calculations, as well as relevant
139 equations

140 • Included an additional section 4.4 to further discuss our modelling results with regard to
141 current studies on mixing assumptions

142

143 All changes we made to the manuscript can be found in the “Manuscript version showing changes”
144 below.

145

146

Manuscript version showing changes

147

148 **Exploring water cycle dynamics by sampling multiple**
149 **stable water isotope pools in a ~~small~~-developed landscape**
150 **of Germany**

151

152 **N. Orlowski^{1,2}, P. Kraft¹, J. ~~Pferdmenges~~Pferdmenges¹ and L. Breuer^{1,3}**

153 [1]{Institute for Landscape Ecology and Resources Management (ILR), Research Centre for
154 BioSystems, Land Use and Nutrition (~~IFZ~~IFZ), Justus-Liebig-University Giessen, Giessen,
155 Germany}

156 [2]{Global Institute for Water Security, ~~School of Environment and Sustainability~~, University
157 of Saskatchewan, Saskatoon, Canada}

158 [3]{Centre for International Development and Environmental Research, Justus Liebig
159 University Giessen, Germany}

160 Correspondence to: N. Orlowski (Natalie.Orlowski@umwelt.uni-giessen.de)

161

162 **Abstract**

163 A dual stable water isotope ($\delta^2\text{H}$ and $\delta^{18}\text{O}$) study was conducted in the developed (managed)
164 landscape of the Schwingbach catchment (Germany). The two-year weekly to biweekly
165 measurements of precipitation, stream, and groundwater isotopes revealed that surface and
166 groundwater are ~~decoupled~~isotopically disconnected from the annual precipitation cycle but
167 showed bidirectional interactions between each other. Apparently, snowmelt played a
168 fundamental role for groundwater recharge explaining the observed differences to
169 precipitation δ -values.

170 A spatially distributed snapshot sampling of soil water isotopes in two soil depths at 52
171 sampling points across different land uses (arable land, forest, and grassland) revealed that top
172 soil isotopic signatures were similar to the precipitation input signal. Preferential water flow
173 paths occurred under forested soils explaining the isotopic similarities between top and
174 subsoil isotopic signatures. Due to human-impacted agricultural land use (tilling and
175 compression) of arable and grassland soils, water delivery to the deeper soil layers was
176 reduced, resulting in significant different isotopic signatures. However, the land use influence

177 ~~smoothed out~~ became less pronounced with depth and soil water approached groundwater δ -
178 values. Seasonally tracing stable water isotopes through soil profiles showed that the
179 influence of new percolating soil water decreased with depth as no remarkable seasonality in
180 soil isotopic signatures was obvious at depth >0.9 m and constant values were observed
181 through space and time.

182 Since classic isotope evaluation methods such as transfer function based mean transit time
183 ~~evaluation~~ calculations failed, we established a hydrological model to estimate spatially
184 distributed groundwater ages and flow directions within the Vollnkirchener Bach
185 subcatchment. Our model revealed that complex age dynamics exist within the subcatchment
186 and that much of the runoff must have been stored ~~in the catchment~~ for much longer than event
187 water-

188 (average water age is 16 years). Tracing stable water isotopes through the water cycle in
189 combination with ~~our~~ hydrological model was valuable for determining interactions between
190 different water cycle components and unravelling age dynamics within the study area. This
191 knowledge can further improve catchment specific process understanding of developed,
192 human-impacted landscapes.

193 **Introduction**

194 The application of stable water isotopes as natural tracers in combination with hydrodynamic
195 methods has been proven to be a valuable tool for studying the origin, and formation, ~~and of~~
196 recharged water as well as the interrelationship between surface water and groundwater
197 (~~Blasch and Bryson, 2007; Goni, 2006~~), ~~partitioning evaporation and transpiration (Phillips~~
198 ~~and Gregg, 2003; Rothfuss et al., 2010, 2012; Wang and Yakir, 2000)~~ (Blasch and Bryson,
199 2007), ~~partitioning evaporation and transpiration (Wang and Yakir, 2000)~~, and further mixing
200 processes between various water sources (~~Aggarwal et al., 2007; Clark and Fritz, 1997c;~~
201 ~~Kendall and Coplen, 2001; Wu et al., 2012)~~ (Clark and Fritz, 1997c). Particularly in
202 catchment hydrology, stable water isotopes play a major role since they can be utilised for
203 hydrograph separations (~~Buttle, 2006; Hoeg et al., 2000; Ladouche et al., 2001; Munyaneza et~~
204 ~~al., 2012~~), ~~to calculate the mean transit time (McGuire et al., 2002, 2005; Rodgers et al.,~~
205 ~~2005b)~~, ~~to investigate water flow paths (Barthold et al., 2011; Goller et al., 2005; Rodgers et~~
206 ~~al., 2005a)~~, or ~~to improve hydrological model simulations (Birkel et al., 2010; Koivusalo et~~
207 ~~al., 1999; Liebinger et al., 2007; Rodgers et al., 2005b)~~. However, spatio-temporal sources
208 ~~of stream water in low angle, developed catchments are still poorly understood.~~ (Buttle, 2006),

209 [to calculate the mean transit time \(McGuire and McDonnell, 2006\), to investigate water flow](#)
210 [paths \(Barthold et al., 2011\), or to improve hydrological model simulations \(Windhorst et al.,](#)
211 [2014\). However, most of our current understanding is resulting from studies in forested](#)
212 [catchments. Spatio-temporal studies of stream water in low angle, developed, agricultural](#)
213 [dominated, and managed catchments are less abundant.](#) This is partly caused by damped
214 stream water isotopic signatures excluding traditional hydrograph separations [in low-relief](#)
215 [catchments](#) (Klaus et al., 2015). Unlike the distinct watershed components found in steeper
216 headwater counterparts, lowland areas often exhibit a complex groundwater–surface water
217 interaction (Klaus et al., 2015). ~~This interaction between groundwater and surface water~~
218 ~~remains poorly understood in many catchments throughout the world but process~~
219 ~~understanding is fundamental to effectively manage the quantity and quality of water~~
220 ~~resources (Ivkovic, 2009). Sklash and Farvolden (1979) showed very early, Sklash and~~
221 [Farvolden \(1979\) showed](#) that groundwater plays an important role as a generating factor for
222 storm and snowmelt runoff processes. In many catchments, streamflow responds promptly to
223 rainfall inputs but variations in passive tracers such as water isotopes are often strongly
224 damped (Kirchner, 2003). ~~This indicates that storm runoff in these catchments is dominated~~
225 ~~mostly by “old water” (Buttle, 1994; Neal and Rosier, 1990; Sklash, 1990)~~ [This indicates that](#)
226 [storm runoff in these catchments is dominated mostly by “old water” \(Buttle, 1994; Neal and](#)
227 [Rosier, 1990; Sklash, 1990\).](#) However, not all “old water” is the same (Kirchner, 2003). This
228 catchment behaviour was described by Kirchner (2003) as the old water paradox. Thus, there
229 is evidence of complex age dynamics within catchments and ~~that~~ much of the runoff is stored
230 in the catchment for much longer than event water (Rinaldo et al., 2015). Still, some of the
231 physical processes controlling the release of “old water” from catchments are poorly
232 understood, roughly modelled, and the observed data do not suggest a common catchment
233 behaviour (Botter et al., 2010). [However, old water paradox behaviour was observed in many](#)
234 [catchments worldwide but it may have the strongest effect in agriculturally managed](#)
235 [catchments, where surprisingly only small changes in stream chemistry have been observed](#)
236 [\(Hrachowitz et al., 2016\).](#)

237 ~~Moreover, due to human-induced alterations of river systems (e.g. channelisation of~~
238 ~~streambeds or draining) (O’Driscoll et al., 2010), water fluxes in developed (managed)~~
239 ~~landscapes can be especially diverse. Almost all European river systems were already~~
240 ~~substantially modified by humans before river ecology research developed (Klapper, 1990;~~
241 ~~Allan, 2004). Through changes in land use, land cover and irrigation, agriculture has~~

242 ~~substantially modified the hydrological cycle in terms of both water quality and quantity~~
243 ~~(Gordon et al., 2010) as well as altered the functioning of aquatic ecosystem processes (Pierce~~
244 ~~et al., 2012; Rockström et al., 2014). This complex character of developed, agricultural~~
245 ~~dominated catchments is often disregarded and established research approaches often failed to~~
246 ~~fully capture agro ecosystem functioning at multiple scales (Orlowski et al., 2014). Since~~
247 ~~agricultural land use (arable land, permanent crops, and grassland) is the most dominant land~~
248 ~~use in Europe (UNEP, 2002), there exists a pressing need to understand biogeochemical~~
249 ~~fluxes (e.g. nitrogen compounds or pesticides) coupled with water fluxes in these managed~~
250 ~~landscapes (Orlowski et al., 2014) and to figure out a way to embed this landscape~~
251 ~~heterogeneity or the consequence of the heterogeneity into models (McDonnell et al., 2007).~~

252 Moreover, almost all European river systems were already substantially modified by humans
253 before river ecology research developed (Allan, 2004). Through changes in land use, land
254 cover, irrigation, and draining, agriculture has substantially modified the water cycle in terms
255 of both quality and quantity (Gordon et al., 2010) as well as hydrological functioning (Pierce
256 et al., 2012). Hrachowitz et al. (2016) recently stated the need for a stronger linkage between
257 catchment-scale hydrological and water quality communities. Further, McDonnell et al.
258 (2007) concluded that we need to figure out a way to embed landscape heterogeneity or the
259 consequence of the heterogeneity (i.e. of agricultural dominated and managed catchments)
260 into models as current generation catchment-scale hydrological and water quality models are
261 poorly linked (Hrachowitz et al., 2016).

262 One way to better understand ~~the relationship between precipitation, stream, soil, and~~
263 ~~groundwater, is detailed knowledge about the isotopic composition of the different~~catchment
264 behaviour and the interaction among the various water sources (surface, subsurface, and
265 groundwater) and their variation in space and time is a detailed knowledge about their
266 isotopic composition. In principal, isotopic signatures of precipitation are altered by
267 temperature, amount (or rainout), continental, altitudinal, and seasonal effects. ~~They are~~
268 ~~mainly influenced by prevailing atmospheric conditions during rainfall and snowfall causing a~~
269 ~~depletion of isotopes (Araguás-Araguás et al., 2000; Blasch and Bryson, 2007; Clark and~~
270 ~~Fritz, 1997c; Gat, 1996; Rohde, 1998). The input signal becomes more pronounced in snow-~~
271 ~~dominated systems where snowfall and snowmelt are depleted in heavy stable water isotopes~~
272 ~~relative to rainfall (Maule et al., 1994; O'Driscoll et al., 2005). Stream water isotopic~~
273 signatures can reflect precipitation isotopic composition and moreover, ~~depend~~dependent on

274 discharge variations be affected by seasonally variable contributions of different water
275 sources such as bidirectional water exchange with the groundwater body during baseflow, or
276 high event-water contributions during stormflow (Genereux and Hooper, 1998; Koeniger et
277 al., 2009). ~~Following the way of precipitation over the unsaturated zone to the groundwater,~~
278 ~~the process of infiltration in itself is known to be a non-fractionating process (Gonfiantini et~~
279 ~~al., 1998), except for mixing between different water pools (e.g. moving and standing water)~~
280 ~~(Gat, 1996). However, precipitation falling on vegetated areas is intercepted by plants and re-~~
281 ~~evaporated thus isotopically fractionated.~~ Precipitation falling on vegetated areas is intercepted
282 by plants and re-evaporated isotopically fractionated. The remaining throughfall infiltrates
283 slower and can be affected by evaporation resulting in an enrichment of heavy isotopes,
284 particularly in the upper soil layers (Gonfiantini et al., 1998; Kendall and Caldwell, 1998). In
285 the soil, specific isotopic profiles develop, characterized by an evaporative layer near the
286 surface ~~especially under arid and semi-arid climate. This decreases exponentially with depth~~
287 ~~(Zimmermann et al., 1968).~~ The isotopic enrichment decreases exponentially with depth,
288 representing a balance between the upward convective flux and the downward diffusion of the
289 evaporative signature (Barnes and Allison, 1988). In humid and semi-humid areas, this
290 exponential decrease is generally interrupted by the precipitation isotopic signal. Hence, the
291 combination of the evaporation effect and the precipitation isotopic signature determine the
292 isotope profile in the soil (Song et al., 2011). Once soil water reaches the saturated zone, this
293 isotope information is finally transferred to the groundwater (Song et al., 2011). Soil water
294 can therefore be seen as a link between precipitation and groundwater, and the dynamics of
295 isotopic composition in soil water are indicative of the processes of precipitation infiltration,
296 evaporation of soil water, and recharge to groundwater (Blasch and Bryson, 2007; Song et al.,
297 2011).

298 To compare different water sources on the catchment-scale, a local meteoric water (LMWL)
299 line is developed and evaporation water lines (EWLs) are used. We started our research with
300 results obtained through an earlier study in the managed Schwingbach catchment that implied
301 a high responsiveness of the system to precipitation inputs indicated by very fast rises in
302 discharge and groundwater head levels They represent the linear relationship between $\delta^2\text{H}$ and
303 $\delta^{18}\text{O}$ of meteoric waters (Cooper, 1998) in contrast to the global meteoric water line
304 (GMWL), which describes the world-wide average stable isotopic composition in
305 precipitation (Craig, 1961a). ~~Thus, the comparison of stable isotope data for stream, soil, or~~

306 groundwater samples relative to the global or local meteoric water lines can provide general
307 understandings on water cycle processes at specific research sites (Song et al., 2011).

308 Identifying the origin of water vapour sources and moisture recycling (Gat et al., 2001; Lai
309 and Ehleringer, 2011), the deuterium excess (d excess), defined by Dansgaard (1964) as
310 $d = \delta^2\text{H} - 8 \times \delta^{18}\text{O}$ can be used, since the d excess mainly depends on the mean relative
311 humidity of the air masses formed above the ocean surface (Zhang et al., 2013). In addition,
312 the d excess reflects the prevailing conditions during evolution, interaction, or mixing of air
313 masses en route to the precipitation site (Fröhlich et al., 2002).

314 To capture spatial landscape heterogeneity, but to keep data acquisition simple, stable water
315 isotope data were coupled with hydrodynamic data from a previous study by Orłowski et al.
316 (2014) in the developed Schwingbach catchment (Germany) to unravel water flow paths and
317 interactions between different water cycle components. Results obtained through this earlier
318 study imply that the Schwingbach catchment is highly responsive, indicated by fast runoff
319 responses to precipitation inputs (Orłowski et al., 2014). Moreover, groundwater reacted
320 almost as quickly as streamflow to precipitation events with raising head levels. Thus, the
321 catchment exhibited “old water” paradox like behaviour (Kirchner, 2003). We further showed
322 that streamflow was predominantly generated in the catchment headwater area and that
323 gaining and losing stream reaches occurred in parallel along the studied stream affected by the
324 underlying geology.

325 Thus, stable water isotopes in combination with hydrodynamic data of a two-year monitoring
326 period (July 2011 to July 2013) were utilised to explore spatio-temporal isotopic variations,
327 unravel linkages between the different water cycle components, investigate the
328 transformations from precipitation to soil and groundwater, and analyse the effect of small-
329 scale landscape characteristics (i.e. soil physical properties, topographic wetness index (TWI),
330 distance to stream, and vegetation cover) on soil water isotopic composition. Further, stable
331 water isotope data was utilized to estimate groundwater ages and flow directions in the
332 Vollnkirchener Bach subcatchment via an hydrological model setup based on the findings of
333 Orłowski et al. (2014).

334 (Orłowski et al., 2014). However, as there was only a negligible influence of the precipitation
335 input signal on the stable water isotopic composition in streams, our initial data set showed
336 evidence for complex age dynamics within the catchment. Nevertheless, a rapid flow response
337 to a precipitation input may also be mistaken (as conceptualized in the vast majority of

338 [catchment-scale conceptual hydrological models](#)) as the actual input signal already reaching
339 [the stream, while in reality it is the remainder of past input signals that slowly travelled](#)
340 [through the system \(Hrachowitz et al., 2016\). The observable hydrological response therefore](#)
341 [acts at different time scales than the tracer response \(Hrachowitz et al., 2016\) as described by](#)
342 [the celerity vs. velocity concept \(McDonnell and Beven, 2014\). The observed patterns in our](#)
343 [catchment therefore inspired us to use a combined approach of hydrodynamic data analyses,](#)
344 [stable water isotope investigations, and data-driven hydrological modelling to determine](#)
345 [catchment dynamics \(response times and groundwater age patterns\) and unravel water flow](#)
346 [paths at multiple spatial scales. This work should further improve our knowledge on](#)
347 [hydrological flow paths in developed, human-impacted catchments.](#)

348 **Materials and methods**

349 **Study area**

350 [The research was carried out in the Schwingbach catchment](#) (50°30'4.23"N, 8°33'2.82"E)
351 (Germany) (Fig. 1a). The Schwingbach and its main tributary the Vollnkirchener Bach are
352 low-mountainous creeks (~~Fig. having an altitudinal difference of 50–100 m over 5 km distance~~
353 [\(Perry and Taylor, 2009\) \(Fig. 1c\)](#) with an altered physical structure of the stream system
354 (channelled stream reaches, pipes, drainage systems, fishponds). The ~~whole~~-Schwingbach
355 catchment ~~encompasses an area of (9.6 km², with an altitude range)~~ [ranges](#) from 233–415 m
356 a.s.l. [in altitude](#). The Vollnkirchener Bach tributary is ~~about 4.7-~~ km in length and drains a 3.7
357 km² subcatchment area (Fig. 1c), ~~which ranges in elevation with elevations~~
358 a.s.l. Almost 46% of the overall Schwingbach catchment is forested, which slightly exceeds
359 agricultural land use (35%) (Fig. 1c). Grassland (10%) is mainly distributed along streams
360 and smaller meadow orchards are located around the villages.

361 ~~The Schwingbach~~ main catchment is underlain by argillaceous shale in the northern parts,
362 serving as aquicludes (~~Mazor, 2003~~). Graywacke zones with lydite in the central, as well as
363 limestone, quartzite, and sandstone regions in the headwater area provide aquifers with large
364 storage capacities (~~Choi, 1997; Mazor, 2003~~) (~~Fig. (Fig. 1f)~~). Loess covers Paleozoic bedrock
365 at north- and east bounded hillsides (Fig. 1f). Streambeds consists of sand and debris covered
366 by loam and some larger rocks (Lauer et al., 2013). Many downstream sections of both creeks
367 are framed by armor stones (Orlowski et al., 2014). ~~Figure 1e shows that the~~ [The](#) dominant
368 soil types in the overall study area are Stagnosols (41%) and mostly forested Cambisols

369 (38%). Stagnic Luvisols with thick loess layers ~~are under agricultural use. The same is true~~
370 ~~for~~, Regosol, Luvisols, and Anthrosols, ~~which encompass an area of 7%. are found under~~
371 ~~agricultural use and~~ Gleysols ~~are found predominantly~~ under grassland ~~sites~~ along the creeks.

372

373

[Figure 1 near here]

374

375 The climate ~~in the study area~~ is classified as temperate with a mean annual temperature of
376 8.2°C. An annual precipitation sum of 633 mm (for the hydrological year 1 November 2012
377 to 31 October 2013) was measured at the catchment's climate station (site 13, Fig. 1b). The
378 year 2012 to 2013 was an average hydrometeorological year. For comparison, the climate
379 station Giessen/Wettenberg (25 km N of the catchment) operated by the German
380 Meteorological Service (DWD, 2014) records a mean annual temperature of 9.6 °C and a
381 mean annual precipitation sum of 666 ~~± ±~~103 mm for the period 1980 ~~to~~ 2010. Discharge
382 peaks from December to April (measured by the use of RBC-flumes with maximum peak
383 flow of 114 L s⁻¹, Eijkelkamp Agrisearch Equipment, Giesbeek, NL) and low flows occur
384 from July until November. Substantial snowmelt peaks were observed during December 2012
385 and February 2013. Furthermore, May 2013 was an exceptional wet month characterised by
386 discharge of 2–3 mm d⁻¹. A ~~more~~ detailed description of runoff characteristics, ~~especially for~~
387 ~~the Vollnkirchener Bach~~ is given ~~in a previous study~~ by Orłowski et al. (2014).

388 **Monitoring network and water isotope sampling**

389 ~~The monitoring network consists of an automated~~ climate station (site 13, Fig. 1 b–c)
390 (Campbell Scientific Inc., AQ5, UK; equipped with a CR1000 data logger ~~collecting air~~
391 ~~temperature at 2 m height, wind speed and direction, relative humidity, and solar radiation~~),
392 three tipping buckets, ~~and~~ 15 precipitation collectors, six stream water sampling points, and
393 22 piezometers (Fig. 1 b–c). Precipitation data were corrected according to Xia (2006).

394 Two stream water sampling points (sites 13 and 18) in the Vollnkirchener Bach are installed
395 with trapezium shaped RBC-flumes for gauging discharge (Eijkelkamp Agrisearch
396 Equipment, Giesbeek, NL), and a V-weir is located at sampling point 64. RBC-flumes and
397 V-weir are equipped with Mini-Divers® (Eigenbrodt Inc. & Co. KG, Königsmoor, DE) for
398 automatically recording water levels ~~and deriving continuous discharge data through the~~

399 ~~given stage discharge relationships (Eijkelkamp, 2014).~~ Discharge at the remaining stream
400 sampling points was manually measured applying the salt dilution method (WTW-cond340i,
401 WTW, Weilheim, DE), ~~which can be precise to $\pm 5\%$ (Day, 1976; Moore, 2004). The 22~~
402 ~~piezometers (Fig. 1b) situated between the conjunction of the Schwingbach with the~~
403 ~~Vollnkirchener Bach and the upper RBC flume of the Vollnkirchener Bach (site 18) are made~~
404 ~~from perforated PVC tubes sealed with a bentonite clay). The 22 piezometers (Fig. 1b) are~~
405 made from perforated PVC tubes sealed with bentonite at the upper part of the tube to prevent
406 contamination by surface water. For monitoring shallow groundwater levels, either combined
407 water level/temperature loggers (Odyssey Data Flow System, Christchurch, NZ) or Mini-
408 Diver® water level loggers (Eigenbrodt Inc. & Co. KG, Königsmoor, DE) are installed.
409 Accuracy of Mini-Diver® is ± 5 mm and for Odyssey data logger ± 1 mm. For calibration
410 purposes, groundwater levels are additionally measured manually via an electric contact
411 gauge.

412 Stable water isotope samples of rainfall, stream-, and groundwater were taken ~~over a two year~~
413 ~~observation period (from July 2011 to July 2013) approximately~~ on weekly intervals, ~~except~~
414 ~~for the winter period.~~ In winter 2012 ~~to~~ 2013, snow core samples over the entire snow depth
415 of < 0.15 m were collected in tightly sealed jars at same sites as open rainfall was sampled.
416 We sampled shortly after snow ~~was fallen~~ fall because sublimation, recrystallization, partial
417 melting, rainfall on snow, and redistribution by wind can alter the ~~primary~~ isotopic
418 composition ~~of the snowfall~~ (Clark and Fritz, 1997b). Samples were melted overnight
419 following Kendall and Caldwell (1998); and analysed for their isotopic composition. Open
420 rainfall was collected in self-constructed samplers. ~~Each collector was made from a 1 L glass~~
421 ~~bottle prepared with a circular funnel of 0.10 m in diameter. Funnels were covered with a~~
422 ~~mosquito net to keep out leaves, insects, or windblown debris. Bottles were placed in PVC~~
423 ~~tubes to avoid heating, screwed to wooden pales, and installed 1 m above ground. To avoid~~
424 ~~sample evaporation, a table tennis ball was placed into each funnel and two layers of small~~
425 ~~plastic balls were inserted into the glass bottles (Windhorst et al., 2013).~~

426 ~~Stream water samples were taken as grab samples at six locations — three sampling points at~~
427 ~~each stream (Vollnkirchener Bach sites: 13, 18, and 94; Schwingbach sites: 11, 19, and 64)~~
428 ~~(Fig. 1b c). To account for possible spatial variation in groundwater, grab samples were~~
429 ~~collected from 17 piezometers (Fig. 1b). Since spatial variations between the piezometers~~
430 ~~under meadow was small, the amount of sampled piezometers was reduced to three as per~~

431 [Windhorst et al. \(2013\)](#). Grab samples of stream water were taken at six locations, three
432 [sampling points at each stream \(Fig. 1b–c\)](#). Since spatial isotopic variations of groundwater
433 [among piezometers under meadow were small, samples were collected at three out of eight](#)
434 sampling points under meadow (sites 1, 6, and 21), five under the arable field (sites 25–29),
435 and four [besidenext to](#) the Vollnkirchener Bach (sites 24, 31, 32, and 35) (Fig. 1b).
436 Additionally, a drainage pipe (site 15) located ~226 m downstream of site 18 was sampled.
437 According to IAEA standard procedures, all samples were filled and stored in 2 mL brown
438 glass vials, sealed with a solid lid, and wrapped up with Parafilm® ([Mook, 2001](#)).

439 **Isotopic soil sampling**

440 **Spatial variability**

441 In order to analyse the effect of small-scale characteristics such as distance to stream, TWI,
442 and land use on soil isotopic signatures ~~as connecting compartment between precipitation and~~
443 ~~groundwater~~, we sampled a snapshot of 52 points evenly distributed over a 200 m grid around
444 the Vollnkirchener Bach (Fig. 1d). Soil samples were taken at four consecutive rainless days
445 (1 to 4 November 2011) at altitudes of 235–294 m a.s.l.. Sampling sites were selected via a
446 stratified, GIS-based sampling plan (ArcGIS, Arc Map 10.2.1, Esri, California, USA),
447 including three classes of topographic wetness indices (TWIs: 4.4–6.5; 6.5–7.7; 7.7–18.4),
448 two different distances to stream (0–121 m, 121–250 m), and three land ~~use units~~ ~~suses~~ (arable
449 land, forest, and grassland), with each class containing the same number of sampling points.
450 Samples were collected at depths of 0.2 m and 0.5 m. Gravimetric water content was
451 measured according to DIN-ISO 11465 by drying soils for 24 h at 110 °C. Soil pH was
452 analysed following DIN-ISO 10390 on 1:1 soil-water-mixture with a handheld pH-meter
453 (WTW cond340i, WTW Inc., DE). Bulk density was determined according to DIN-ISO
454 11272, and soil texture by finger testing ([Whitefield, 2004](#)).

455 **Seasonal isotope soil profiling and isotope analysis**

456 ~~In order to trace the seasonal development of stable water isotopes from rainfall to~~
457 ~~groundwater, seven soil profiles were taken in the dry summer season (28 August 2011),~~
458 ~~seven in the wet winter period (28 March 2013), and two profiles in the transitional season~~
459 spring (24 April 2013) under different vegetation cover (arable land and grassland) (Fig. 1d).
460 Soil was sampled utilising a hand-auger (Eijkelkamp Agrisearch Equipment BV, Giesbeek,

461 ~~DE). Samples were taken~~ from the soil surface to 2 m depth. Samples were collected in
462 greater detail near the soil surface since this area is known to have the greatest isotopic
463 variability (Barnes and Allison, 1988; ~~Hsieh et al., 1998; Zimmermann et al., 1968~~).

464 ~~Soil samples were stored in amber glass tubes, sealed with Parafilm®, and kept frozen until~~
465 ~~water extraction (Orlowski et al., 2013).~~ Soil water was extracted cryogenically with 180 min
466 extraction duration, a vacuum threshold of 0.3 Pa, and an extraction temperature of 90°C
467 following Orlowski et al. (2013). Isotopic signatures of $\delta^{18}\text{O}$ and $\delta^2\text{H}$ were analysed via off-
468 axis integrated cavity output spectroscopy (OA-ICOS) (DLT-100, Los Gatos Research Inc.,
469 Mountain View, ~~CA~~, USA). Within each isotope analysis three calibrated stable water isotope
470 standards of different water isotope ratios were included (LGR working standard number 1, 3,
471 and 5; Los Gatos Research Inc., CA, US). After every fifth sample the LGR working
472 standards are measured. For each sample, six sequential 900 μL aliquot of a water sample are
473 injected into the analyser. Then, the first three measurements are discarded. The remaining are
474 averaged and corrected for per mil scale linearity following the IAEA laser spreadsheet
475 template (Newman et al., 2009). Following this IAEA standard procedure allows for drift and
476 memory corrections. Isotopic ratios are reported in per mil (‰) relative to Vienna Standard
477 Mean Ocean Water (VSMOW) (Craig, 1961b). Accuracy of analyses was 0.6‰ for $\delta^2\text{H}$ and
478 0.2‰ for $\delta^{18}\text{O}$ (LGR, 2013). Leaf water extracts typically contain a high fraction of organic
479 contaminations ~~(West et al., 2010),~~ which might lead to spectral interferences when using
480 isotope ratio infrared absorption spectroscopy ~~(Leen et al., 2012),~~ causing erroneous isotope
481 values (Schultz et al., 2011). ~~Therefore, isotopic data of plant water extracts are usually~~
482 ~~checked for spectral interferences using the Spectral Contamination Identifier (LWIA-SCI)~~
483 ~~post processing software (Los Gatos Research Inc.). However, for soil water extracts no~~
484 ~~evidence for such interferences have been observed so far~~ However, for soil water extracts
485 there exists no need to check or correct such data (Schultz et al., 2011; Zhao et al., 2011).
486 ~~Thus, there exists no need to check or correct such data.~~

487 Mean transit time estimation

488 ~~To understand the connectivity between the different water cycle components in the~~
489 ~~Schwingbach catchment, the mean transit times (MTT) for the Vollnkirchner Bach (sites 13,~~
490 ~~18, and 94) and the Schwingbach (sites 11, 19, and 64) were calculated using FlowPC~~
491 ~~(Małozzewski and Zuber, 1996). Different models (dispersion model with different dispersion~~
492 ~~parameters $D_p = 0.05, 0.4,$ and $0.8,$ exponential model, exponential-piston-flow model, linear~~

493 ~~model, and linear piston flow model) were compared for their results (sigma as goodness of~~
494 ~~fit and model efficiencies (ME)). A model efficiency $ME = 1$ indicates an ideal fit of the~~
495 ~~model to the concentrations observed, while $ME = 0$ indicates that the model fits the data no~~
496 ~~better than a horizontal line through the mean concentration observed (Maloszewski and~~
497 ~~Zuber, 2002). The same is true for sigma.~~

498 ~~For calculations with FlowPC, weekly averages of precipitation and stream water isotopic~~
499 ~~signatures were calculated. We also bias corrected the precipitation input data with two~~
500 ~~different approaches: the mean precipitation value was subtracted from every single~~
501 ~~precipitation value and then divided by the standard deviation of precipitation isotopic~~
502 ~~signatures. Afterwards, this value was subtracted from the weekly precipitation values (bias1).~~
503 ~~For the second approach, the difference of the mean stream water isotopic value and the mean~~
504 ~~precipitation value was calculated and also subtracted from the weekly precipitation values~~
505 ~~(bias2).~~

506 **Hydrological model setup**

507 ~~To estimate the age dynamics of the groundwater body in the Vollnkirchener Bach~~
508 ~~subcatchment, a hydrological model was established on the basis of the conceptual model~~
509 ~~presented by Orłowski et al. (2014). For this purpose the *Catchment Modelling Framework*~~
510 ~~(CMF) by Kraft et al. (2011) was used. CMF is a modular framework for hydrological~~
511 ~~modelling based on the concept of finite volume method by Qu and Duffy (2007). CMF is~~
512 ~~applicable for simulating one- to three-dimensional water fluxes but also advective transport~~
513 ~~of stable water isotopes (^{18}O and ^2H). Thus, it is especially suitable for our tracer study and~~
514 ~~can be used to study the origin (Windhorst et al., 2014) and age of water.~~

515 ~~The generated model is a highly simplified representation of the Vollnkirchener Bach~~
516 ~~subcatchment's groundwater body. The subcatchment is divided into 353 polygonal shaped~~
517 ~~cells ranging from 101.7–38940.1 m², manually adjusted on the basis of land use, soil types,~~
518 ~~and contour lines following Qu and Duffy (2007) and Windhorst et al. (2014). Each cell~~
519 ~~contains two layers, one comprises a water storage. The upper layer, representing the~~
520 ~~groundwater body, is generated based on soil depth measurements and reaches down to 20 m~~
521 ~~below the surface. Due to the fact that groundwater depth was not measured, the layer-~~
522 ~~thickness is a rough estimation. The second layer (20–40 m below the surface) represents the~~
523 ~~bedrock. The main fresh water input is the groundwater recharge, which is a constant value~~

524 ~~over time for each cell. It is calculated as the difference between rainfall, evapotranspiration,~~
525 ~~and the change in stored water. Precipitation and evapotranspiration values are calculated~~
526 ~~using a fully distributed 3D model established through CMF with a one year simulation~~
527 ~~period of the same subcatchment. The change in stored water is set to zero since a steady state~~
528 ~~is simulated (see below) and therefore the water content in the system is stable.~~

529 ~~Besides the groundwater recharge, a combined sewer overflow (site 38) is considered as an~~
530 ~~additional water input based on findings of Orłowski et al. (2014). Moreover, there are two~~
531 ~~water outlets in the two lowest cells for efficient draining. Both cells are located in the very~~
532 ~~north of the subcatchment. The compartments within the system are linked by a series of~~
533 ~~flow accounting equations: Richards equation for percolation, Darcy equation for lateral~~
534 ~~subsurface flow, Neumann boundary condition for input of fresh water (groundwater~~
535 ~~recharge, pipe source), and constant Dirichlet boundary conditions representing the system~~
536 ~~outlets.~~

537 ~~For estimating the groundwater age, a virtual tracer is used. It is modelled as a radioactive~~
538 ~~decay tracer with a fixed concentration at the input to the system. From the modelled~~
539 ~~concentration of the tracer in each cell, the mean age of the water for this cell is derived.~~

540 ~~**Model assumptions:**~~ The saturated hydraulic conductivity of the groundwater body is set to
541 ~~0.1007 m d⁻¹, as measured in the study area. For the~~ bedrock compartment there is no data
542 available. However, expecting a high rate of joints, preliminary testing revealed that a
543 saturated hydraulic conductivity of 0.25 m d⁻¹ seemed to be a realistic estimation (based on
544 field measurements).

545 To understand the connection between the different water cycle components in the
546 Schwingbach catchment, mean transit times (MTT) for both streams as well as mean
547 residence times (MRT) from precipitation to groundwater were calculated using FlowPC
548 (Maloszewski and Zuber, 2002). See Appendix I for details about the applied method.

549 **Model-based groundwater age dynamics**

550 To estimate the age dynamics of the groundwater body in the Vollnkirchener Bach
551 subcatchment, a hydrological model was established on the basis of the conceptual model
552 presented by Orłowski et al. (2014) and the isotopic measurements presented here. Appendix
553 II outlines the modelling concept, model set up, and its parameterization.

554 **Statistical analyses**

555 For statistical analyses, we used IBM SPSS Statistics (Version 22, SPSS Inc., Chicago, IL,
556 US) and R (version Rx64 3.2.2). The R package igraph was utilized for plotting (Csardi and
557 Nepusz, 2006). Studying temporal and spatial variations in meteoric and groundwater, isotope
558 data were tested for normal distribution. Subsequently, t-tests or Multivariate Analyses of
559 Variances (MANOVAs) were applied and Tukey-HSD tests were run to determine which
560 groups were significantly different ($p \leq 0.05$). Event mean values of isotopes in
561 precipitation, stream, and groundwater were calculated when no spatial variation was
562 observed. [Regression analyses were run to determine the effect of small-scale characteristics
563 such as distance to stream, TWI, and land use on soil isotopic signatures.](#)

564 We used a topology inference network map (Kolaczyk, 2014) in combination with a principal
565 component analysis (~~Jolliffe, 2002) to show $\delta^{18}\text{O}$ isotope relationships between surface and
566 groundwater sampling points to show $\delta^{18}\text{O}$ isotope relationships between surface and
567 groundwater sampling points.~~ To explore the sensitivity of missing data, we used both the
568 complete isotope time series and randomly selected 80% of the whole data sets. Overall, the
569 cluster relationships of the surface and groundwater sampling points are largely similar for
570 both whole and subsets of isotope data sets, despite some differences of the exact cluster
571 centroid locations. We therefore decided to use randomly selected 80% of the isotope time
572 series to illustrate our results. In the network map, each node of the network represents an
573 isotope sampling point. The locations of the nodes are based on the first two components
574 (PC1 and PC2). The correlations between isotope time series are represented by the edges
575 connecting nodes. The thickness of edges characterizes the strength of the correlations. The p-
576 values of correlations are approximated by using the F-distributions and mid-ranks are used
577 for the ties (Hollander et al., 2013). Only statistically significant connections ($p < 0.05$) are
578 shown ~~in the network diagram. Basic background information related to graph theory can be
579 found in Wallis (2007).~~

580 [To compare different water sources on the catchment-scale, a local meteoric water \(LMWL\)
581 line was developed and evaporation water lines \(EWLs\) were used.](#) They represent the linear
582 relationship between $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of meteoric waters (Cooper, 1998) in contrast to the global
583 meteoric water line (GMWL), which describes the world-wide average stable isotopic
584 composition in precipitation (Craig, 1961a). [Identifying the origin of water vapour sources](#)

585 [and moisture recycling \(Gat et al., 2001; Lai and Ehleringer, 2011\), the deuterium-excess \(d-](#)
586 [excess\), defined by Dansgaard \(1964\) as \$d = \delta^2\text{H} - 8 \times \delta^{18}\text{O}\$ was used.](#)

587 For comparisons, precipitation isotope data from the closest GNIP (Global Network of
588 Isotopes in Precipitation) station Koblenz (DE; [73.874](#) km SW of the study area, 97 m a.s.l.)
589 was used (IAEA, 2014; Stumpp et al., 2014). For monthly comparisons with Schwingbach d-
590 [excess](#) values, we used a data set from the GNIP station Koblenz that includes 24 values
591 starting from July 2011 to July 2013.

592 **Results**

593 ~~Descriptive statistics of isotopic composition in precipitation, stream, and groundwater are~~
594 ~~shown along with d-excess values in Table 1 and are described in detail in the following:~~

595

596 [\[Table 1 near here\]](#)

597

598 **Variations of precipitation isotopes and d-excess**

599 The $\delta^2\text{H}$ values of all precipitation isotope samples (~~$N = 592$~~) ~~taken throughout the~~
600 ~~observation period (July 2011 to July 2013)~~ ranged from -167.6 to -8.3‰ (Table 1). To
601 examine the spatial isotopic ~~variation~~[variations](#), rainfall was collected at 15 open field site
602 locations throughout the Schwingbach main catchment (Fig. 1b–c) for a 7-month period.
603 ~~However, but~~ no spatial variation could be observed ~~in the Schwingbach catchment.~~ Thus,
604 rainfall was collected at the catchment outlet (site 13) from 23 October 2014 onward ~~and~~
605 ~~event mean δ values were calculated for the previous isotope data.~~

606 ~~.~~ Analysing effects that influence the isotopic composition of precipitation, neither an amount
607 effect nor an altitude effect ~~was~~[were](#) found – not surprisingly, as the greatest altitudinal
608 difference between sampling points was only 101 m. Nevertheless, a slight temperature effect
609 ($R^2 = 0.5$ for $\delta^2\text{H}$ and $R^2 = 0.6$ for $\delta^{18}\text{O}$, respectively) was observed showing enriched
610 isotopic signatures at higher temperatures.

611 [\[Table 1 near here\]](#)

612 Strong temporal variations in precipitation isotopic signatures, as well as pronounced seasonal
613 isotopic effects were measured with greatest isotopic differences occurring between summer

614 and winter. Samples taken in the fall and spring were isotopically similar, however, differed
615 from winter isotopic signature, which were somewhat lighter (Fig. 2). Furthermore, in the
616 winter of 2012–13 snow could be sampled, which decreased the mean winter isotopic values
617 for this period in comparison to the previous winter period (2011–12). ~~No~~The mean $\delta^2\text{H}$
618 isotope values of snow samples were approximately 84‰ lighter than mean precipitation
619 isotopic signatures (Fig. 3). Further, no statistically significant ($p>0.05$) inter-annual variation
620 was detected between the summer periods of 2011 and 2012 (Fig. 2), ~~which could have~~
621 ~~reflected varying local climate conditions (Koeniger et al., 2009).~~

622
623 [Figure 2 near here]

624
625 Examining the influence of moisture recycling on the isotopic compositions of precipitation,
626 the ~~deuterium excess~~ (d-excess) was calculated for each individual rain event at the
627 Schwingbach catchment. ~~For the two year observation period, d~~D-excess values (~~N=108~~)
628 ranged from -7.8‰ to $+19.4\text{‰}$ and averaged $+7.1\text{‰}$ (Fig. 2). In general, 37% of all events
629 were ~~sampled~~sampled in summer periods (21 June to 21/22 September) and showed lower d-
630 excess values in comparison to the 19% winter precipitation events (21/22 December to 19/20
631 March) (Fig. 2). D-excess greater than $+10\text{‰}$ was determined for 22% of all events. ~~As a~~
632 ~~reference the d-excess of the GMWL $d = 10$ is depicted in Figure 2 (solid line).~~ Lowest values
633 corresponded to summer precipitation events ~~with~~where evaporation of the raindrops below
634 the cloud base ~~at mean daily air temperatures between 12–18°C may occur~~. Most of the higher
635 values ($>+10\text{‰}$) appeared in cold seasons (fall/winter) and winter snow samples of the
636 Schwingbach catchment with ~~very~~much depleted δ -values showed highest d-excess ~~values~~
637 (Fig. 2).

638 In comparison with the GNIP station Koblenz (~~years~~ 2011–2013), the mean annual d-excess
639 at the Schwingbach catchment was on average 3.9‰ higher (~~7.1‰ for 2011–12 and 2012–13,~~
640 ~~respectively~~),₂ showing a greater impact of oceanic moisture sources than the further south-
641 west located station Koblenz. The ~~unweighted mean annual d-excess at the GNIP station~~
642 ~~Koblenz was 2.9‰ for July 2011 to June 2012 and 3.6‰ for July 2012 to June 2013, whereas~~
643 ~~the~~ long-term mean d-excess was 4.4‰ for the Koblenz station (1978–2009) (Stumpp et al.,
644 2014). ~~Nevertheless, highest~~Highest d-excesses at the GNIP station matched highest values in
645 the Schwingbach catchment, both occurring in the cold seasons (October to December 2011

646 and November to December 2012). ~~Since no amount effect on the $\delta^2\text{H}$ and $\delta^{18}\text{O}$ values was~~
647 ~~observed in the Schwingbach, also no linear regression of event d-excess with precipitation~~
648 ~~amount was detected.~~

649 ~~The linear relationship of $\delta^2\text{H}$ and $\delta^{18}\text{O}$ content in local precipitation, results in a local~~
650 ~~meteoric water line (LMWL) (Fig. 3), which can be utilised to link the relative contribution of~~
651 ~~seasonal precipitation to ground and surface water sources (Wassenaar et al., 2011). The~~
652 ~~global meteoric water line (GMWL) established by Craig (1961a), and more recently refined~~
653 ~~by Rozanski et al. (1993) is $\delta^2\text{H} = 8.13 \times \delta^{18}\text{O} + 10.8 \text{ ‰}$, provides a valuable benchmark~~
654 ~~against which regional or local waters can be compared (Song et al., 2011). The slope of the~~
655 ~~LMWL of the Schwingbach catchment is well in agreement with the one from the closest~~
656 ~~GNIP station in Koblenz ($\delta^2\text{H} = 7.66 \times \delta^{18}\text{O} + 2.0 \text{ ‰}$; $R^2 = 3$). The slope of the Schwingbach~~
657 ~~LMWL is well in agreement with the one from the GNIP station Koblenz~~
658 ~~($\delta^2\text{H} = 7.66 \times \delta^{18}\text{O} + 2.0 \text{ ‰}$; $R^2 = 0.97$; 1978–2009 (Stumpp et al., 2014)), but is slightly lower in~~
659 ~~comparison to the revised GMWL, showing stronger local evaporation conditions. Since~~
660 ~~evaporation causes a differential increase in $\delta^2\text{H}$ and $\delta^{18}\text{O}$ values of the remaining water, the~~
661 ~~slope for the linear relationship between $\delta^2\text{H}$ and $\delta^{18}\text{O}$ is lower in comparison to the GMWL~~
662 ~~(Rozanski et al., 2001; Wu et al., 2012). The lower intercept (d-excess), dependent on the~~
663 ~~humidity and temperature conditions in the evaporation region (Mook, 2001), nevertheless~~
664 ~~shows that moisture recycling did obviously not play a major role in the study area.~~

665

666

[Figure 3 near here]

667

668 ~~Considering isotope samples of the different water cycle components in comparison with the~~
669 ~~LMWL revealed that mean isotope values of snow samples were for $\delta^2\text{H}$ approximately 84%~~
670 ~~lighter than mean precipitation isotopic signatures (Fig. 3). Stream water isotope samples of~~
671 ~~both creeks (Schwingbach and Vollnkirchener Bach) fell on the LMWL, showing slight~~
672 ~~evaporative enrichment for few samples (Fig. 3). Moreover, isotopic values for stream water~~
673 ~~were almost identical to those found in groundwater (Table 1, Fig. 3).~~

674 Isotopes of soil water

675 Spatial variability

676 Determining the impact of landscape characteristics on soil water isotopic signatures, we
677 found no ~~relationship statistically significant connection~~ between the parameters distance to
678 stream, TWI, soil water content, soil texture, pH, and bulk density with the soil isotopic
679 signatures in ~~two depths (0.2 and 0.5 m), except for land use both soil depths, except for land~~
680 ~~use~~. This was potentially attributed to the small variation in soil textures (mainly clayey silts
681 and loamy sandy silts), bulk densities, and pH values for both soil depths (Table 2). ~~Water~~
682 ~~contents showed the greatest standard deviation within the two soil depths (Table 2),~~
683 ~~however, exhibited no effect on soil water isotopes. Moreover, no tendency of higher TWI~~
684 ~~values with decreasing distance to stream was obvious. Distances to the stream are linked to~~
685 ~~water flow path lengths and were therefore supposed to be a controlling factor. However, no~~
686 ~~impact of different distances to the stream on soil water isotopic signatures could be observed.~~

687

688 [Table 2 near here]

689

690 ~~The mean δ values in the~~ top 0.2 m of the soil profile ~~is~~are higher than ~~further below in the~~
691 ~~subsoil~~, reflecting a stronger impact of precipitation in the topsoil (Table 2, Fig. 4). ~~The~~While
692 ~~the~~ δ -values ~~of top soil for subsoil~~ and precipitation ~~did not vary differed~~ significantly
693 ~~statistically~~ ($p > \leq 0.05$), ~~which is they did~~ not ~~the case~~ for ~~precipitation and subsoil topsoil~~ (Fig.
694 4). Subsoil isotopic values were statistically equal to stream ~~water~~ and groundwater ~~isotopic~~
695 ~~values~~ (Fig. 4).

696

697 [Figure 4 near here]

698

699 Generally, all soil water isotopic values fell on the ~~local meteoric water line~~LMWL,
700 indicating no evaporative enrichment ~~of soil water~~ (Fig. 5). Comparing soil isotopic
701 signatures between different land covers showed generally higher and statistically
702 significantly different δ -values ($p \leq \leq 0.05$) at 0.2 m soil depth under arable land as compared
703 to forests and grasslands. ~~However, all top soil isotopic values reflected precipitation isotopic~~

704 ~~signals (Fig. 5, top).~~ For the lower 0.5 m of the soil column, isotopic signatures under all land
705 ~~use units~~ suses showed statistically similar values; ~~nevertheless, differing significantly from~~
706 ~~precipitation ($p \leq 0.05$) (Fig. 5, bottom).~~

707

708 ~~].~~ [Figure 5 near here]

709

710 Comparing soil water $\delta^2\text{H}$ values between top and subsoil under different land use units
711 showed significant differences ($p \leq 0.05$) under arable and grassland but not under forested
712 sites (Fig. ~~5, capital letters~~ 5).

713 [Figure 5 near here]

714 **Seasonal isotope soil profiling**

715 Isotope compositions of soil water varied seasonally: More depleted soil water was found in
716 the winter and spring (Fig. 6); contrary, soil water was enriched in summer due to evaporation
717 during warmer and drier periods (Darling, 2004). For summer soil profiles in the
718 Vollnkirchener subcatchment, no evidence for evaporation was obvious below 0.4 m soil
719 depth. However, snowmelt isotopic signatures could be traced down to a soil depth of 0.9 m
720 during spring rather than winter, pointing to a depth-translocation of meltwater in the soil,
721 more remarkable for the deeper profile under arable land (Fig. 6, upper left panel).
722 Furthermore, shallow soil water (<0.4 m) showed larger standard deviations with values
723 closer to mean seasonal precipitation inputs (Fig. 6, upper panels). Winter profiles exhibited
724 somewhat greater standard deviations in comparison to summer isotopic soil profiles. The
725 observed seasonal amplitude ~~smoothed out~~ became less pronounced with depth as soil water
726 isotope signals approached groundwater average in >0.9 m depth. Generally, deeper soil
727 water isotope values were relatively constant through time and space.

728

729 [Figure 6 near here]

730

731 **Isotopes of stream water**

732 ~~Analysing spatial~~No statistically significant differences ~~in isotopic compositions~~were found
733 between ~~the~~ Schwingbach (sites 11, 19, and 64) and Vollnkirchener Bach (sites 13, 18, and
734 94) stream water ~~resulted in no statistically significant differences for all sampling points~~
735 (Fig. 7). ~~In general,~~7) with isotope data falling on the LMWL but showing slight evaporative
736 ~~enrichment for few samples (Fig. 3).~~ $\delta^{18}\text{O}$ values varied for the Vollnkirchener Bach by
737 $-8.4\pm 0.4\text{‰}$ and for the Schwingbach by $-8.4\pm 0.6\text{‰}$ ~~over the two-year observation~~
738 ~~period~~(Table 1). Stream water isotopic signatures were by approximately -15‰ in $\delta^2\text{H}$ more
739 ~~depleted than precipitation signatures and similar to groundwater~~ (Table 1).

740

741 [Figure 7 near here]

742

743 ~~Stream water isotopic signatures in the Schwingbach catchment were by approximately~~
744 ~~-15‰ in $\delta^2\text{H}$ more depleted than precipitation signatures (Table 1). However, surface water~~
745 ~~isotopic compositions were similar to groundwaters (Table 1).~~

746 Examining temporal isotopic variations, ~~A~~ damped seasonality (~~less variation~~) of the isotope
747 concentration in stream water ~~in comparison to~~versus precipitation was ~~measured with main~~
748 ~~seasonal differences~~ occurring between summer and winter ~~periods~~(Fig. 7). Most outlying
749 depleted stream water isotopic signatures (e.g. in March 2012 and 2013) ~~could~~can be
750 explained by snowmelt (Fig. 7). However, the outlier at the Schwingbach stream water
751 sampling site 64 (-66.7‰ for $\delta^2\text{H}$) is by 8.5‰ more depleted than the two-year average of
752 Schwingbach stream water (Table 1). Rainfall falling on 24 September 2012 was -31.9‰ for
753 $\delta^2\text{H}$. This period in September was generally characterized by low flow and little rainfall
754 (~~antecedent precipitation index: AP8 was 8mm~~). Thus, little contribution of new water was
755 observed and stream water isotopic signatures were groundwater-dominated. For site 13, the
756 outlier in May 2012 (-44.2‰ for $\delta^2\text{H}$) was by 13.8‰ more enriched than the average stream
757 water isotopic composition of the Vollnkirchener Bach over the two-year observation period
758 (Table 1). A runoff peak at site 13 of ~~0.45215~~ 15 mm d^{-1} and a 2.9 mm rainfall event were
759 recorded on 23 May 2012. ~~Moreover, AP8 was 23.2 mm~~. Thus, this outlier could be explained
760 by precipitation contributing to stream flow causing more enriched isotopic values in stream
761 water, which approached average precipitation δ -values (-43.9 ± 23.4).

762 Calculated MTT for the Schwingbach ranged ~~betweenfrom~~ 52 ~~and~~ 67 weeks and for the
763 Vollnkirchener Bach ~~betweenfrom~~ 47 ~~and~~ 66 weeks, whereby linear and exponential models
764 provided the best fits for all sampling points. However, the calculated output data did not fit
765 the observed values in terms of the quality ~~critierioncriteria~~ sigma and model efficiency
766 (Timbe et al., 2014). ~~Model efficiencies for the Schwingbach sampling points were~~
767 ~~($ME_{Schwingbach}$ -0.1-0.0 and for the Vollnkirchener, $ME_{Vollnkirchener}$ Bach 0.0-0.4. Sigma values;~~
768 ~~sigma~~ for all sampling points ~~were 0.1 for the best fit models, respectively).~~ Even a bias
769 correction of the input data (~~precipitation~~) did not improve the model outputs (~~sigma = 0.1~~).
770 Therefore, we conclude that ~~the application of transfer function based~~ MTT estimation
771 methods ~~based on stable water isotopes failed in the Schwingbach catchment and developed a~~
772 ~~new data driven groundwater model to simulate observed~~ applying stable water isotope data
773 failed for the Schwingbach.

774 **Isotopes of groundwater**

775 ~~Since groundwater head levels responded almost as quickly as streamflow to rainfall events,~~
776 ~~rainfall isotopic signatures were assumed to be rapidly transferred to the groundwater. This~~
777 ~~was likewise underlined by the fact that Orłowski et al. (2014) observed bidirectional water~~
778 ~~interactions between the groundwater body and the stream. Studying groundwater isotopic~~
779 ~~signatures at the downstream section of the Vollnkirchener Bach, almost constant isotopic~~
780 ~~values (Fig. 8, Table 1) throughout the study period were observed ($\delta^2H: -57.6 \pm 1.6\text{‰}$ for~~
781 ~~piezometers under meadow). Most depleted groundwater isotopic values ($\leq -80\text{‰}$ for δ^2H)~~
782 ~~were measured for piezometer 32 during snowmelt events in March and April 2013 as well as~~
783 ~~for piezometer 27 from December 2012 to February 2013. As shown by Orłowski et al.~~
784 ~~(2014) piezometer 32 is highly responsive to rainfall runoff events and groundwater head~~
785 ~~elevations showed significant correlations with mean daily discharge at this site. Further,~~
786 ~~effluent conditions and lowest K_{sat} values ($7-14 \text{ mm h}^{-1}$) were measured in this stream section~~
787 ~~(piezometers 32-35) (Orłowski et al., 2014).~~

788 **In the Schwingbach catchment, groundwater/Isotopes of groundwater**

789 Since groundwater head levels responded almost as quickly as streamflow to rainfall events,
790 rainfall isotopic signatures were assumed to be rapidly transferred to the groundwater. For the
791 piezometers under meadow, almost constant isotopic values (Fig. 8, Table 1) were observed
792 ($\delta^2H: -57.6 \pm 1.6\text{‰}$). Most depleted groundwater isotopic values ($\leq -80\text{‰}$ for δ^2H) were

793 [measured for piezometer 32 during snowmelt events in March and April 2013 as well as for](#)
794 [piezometer 27 from December 2012 to February 2013. Piezometer 32 is highly responsive to](#)
795 [rainfall-runoff events and groundwater head elevations showed significant correlations with](#)
796 [mean daily discharge at this site \(Orlowski et al., 2014\).](#)

797 [Groundwater](#) under meadow differed from mean precipitation values by about -14‰ for $\delta^2\text{H}$
798 showing no evidence of a rapid transfer of rainfall isotopic signatures to the groundwater-
799 [\(Fig. 8\). This was underlined by the results of our MRT estimations which varied between](#)
800 [56–65 weeks for the thirteen considered piezometers. However, the calculated output data did](#)
801 [not fit the observed values showing very low MEs \(ME: \$-0.62\$ – \$-0.09\$ for \$\delta^{18}\text{O}\$ and \$-0.49\$ –](#)
802 [\$0.16\$ for \$\delta^2\text{H}\$; sigma: \$0.08\$ – \$0.15\$ for \$\delta^{18}\text{O}\$ and \$0.62\$ – \$1.11\$ for \$\delta^2\text{H}\$ \).](#)

803

804 [Figure 8 near here]

805

806 Due to different water flow paths of groundwater along the studied stream, [we expected to](#)
807 [find](#) distinguished groundwater isotopic signatures ~~were assumed to be found.~~ In fact, we
808 could identify spatial statistical differences between grassland and arable land groundwater
809 isotopic signatures (Fig. 9). Groundwater isotopic signatures under arable land (sites: 25–29,
810 Fig. 1b) showed more enriched values (Fig. 8) and showed significant correlations ($p < 0.05$)
811 among each other (Fig. 9). Arable land groundwater plotted furthest away from surface water
812 sampling points in our network map, showing no significant correlations to either the
813 Schwingbach or the Vollnkirchener Bach. ~~This hydrological disconnectivity was already~~
814 ~~observed in the study of Orlowski et al. (2014). In general,~~ $\delta^{18}\text{O}$ time series of piezometers
815 along the stream and under the meadow showed ~~elose~~closest relations to surface water
816 sampling points (Fig. 9). We further found high correlations ($R^2 > 0.6$) of $\delta^{18}\text{O}$ time series of
817 piezometers located under the meadow ~~(sites: 3, 6, and 21)~~ among each other. Additionally,
818 $\delta^{18}\text{O}$ values of piezometer 3 correlated significantly ($p < 0.05$) with surface water sampling
819 points 18 and 94 ($R^2 = 0.6$ and 0.8 , respectively) and piezometer 32 with sampling points 13
820 and 64 ($R^2 = 0.8$ and 0.6 , respectively).

821

822 [Figure 9 near here]

823

824 We further observed close relations ($p < 0.05$) among $\delta^{18}\text{O}$ values of Vollnkirchener Bach
825 sampling sites 13, 18, and 94 as well as of Schwingbach sites 11, 19, and 64 along with
826 significant correlations between each other.

827 **4.1.1 Modelled groundwater Groundwater age dynamics**

828 Since MTT calculations failed, we modelled the groundwater age in the Vollnkirchener Bach
829 subcatchment using CMF, involving (Appendix II), applying observed hydrometric as well as
830 stable water isotope data (Fig. 10).

831

832 [Figure 10 near here]

833

834 The maximum age of water is highly variable throughout the subcatchment, which results in a
835 very heterogeneous spatial age distribution. The groundwater in most of the outer cells is very
836 young (0–10 years), whereas the inner cells, which incorporate the
837 VollnkirehnerVollnkirchener Bach, contain older water (>30 years). The oldest water (≥ 55
838 years) can be found in the Northern part of the catchment (Fig. 10, detail view), where the
839 VollnkirehnerVollnkirchener Bach drains into the Schwingbach. The main outlets of the
840 subcatchment (dark red coloured cell and green cell) even reach an age of 100 and 55 years,
841 respectively. This can be explained by the fact that these are it is the lowest cellscell within the
842 subcatchment. Thus, and that water flows from the higher to the lower cells and water from
843 the whole subcatchment accumulates at these cells here. The overall flow path to these
844 cellsthis cell is the longest and as a consequence the groundwater age of these cells in this cell
845 is the highest.

846 In general, six2% of cells contain groundwater that is older than 50 years (, <1.7% of cells),
847 and two cells% reveal ages >70 years (0.6%). In contrast, 47 cells (, 13.3%)% contain water
848 with an age of less than one year, and 52.4% with an age <15 years. Thus, most of the cells
849 contain young to moderately old water (<15 years), while few cells comprise old water (>50
850 years). The average groundwater age in the Vollnkirchener Bach subcatchment is 16 years.
851 RelatingCorrelating the groundwater age toagainst the distance to the stream, we found a
852 linear correlation ($R^2 = 0.3$) with a distinct trend. The water tends to be younger with
853 greater distance to the stream.

854 ~~The modelled main flow direction is towards the Vollnkirchener Bach (Fig. 10).~~ The amount
855 of flowing water depicted by the length of the arrows is generally higher near the stream,
856 whereas in most of the outer cells the amount ~~of water is very low (Fig. 10); is very low (Fig.~~
857 10). The modelled main flow direction is towards the Vollnkirchener Bach but many arrows
858 show flow direction across the stream indicating bidirectional water exchange between the
859 stream and the groundwater body.

860 Discussion

861 Variations of precipitation isotopes and d-excess

862 Analysing effects that influence the precipitation isotopic ~~composition,~~ We found no spatial
863 variation in precipitation isotopes ~~was observed~~ throughout the Schwingbach catchment.
864 Mook et al. (1974) also observed for north-western Europe that precipitation collected over
865 periods of 8 and 24 h from three different locations within 6 km² at the same altitude were
866 consistent within 0.3‰ for δ¹⁸O. ~~Further, no amount or altitude effect for isotopes in~~
867 precipitation was found. However, the observed linear relationship (δ¹⁸O = Further, we detected
868 no amount or altitude effect on isotopes in precipitation. Amount effects generally occur most
869 likely in the tropics or for intense convective rain events and are not a key factor for
870 explaining isotope distributions in German precipitation (Stumpp et al., 2014).

871 The observed linear relationship (δ¹⁸O = 0.44T - 12.05‰) between air temperature and
872 precipitation δ¹⁸O values compares reasonably well with a correlation reported by Yurtsever
873 (1975) based on north Atlantic and European stations from the GNIP network
874 δ¹⁸O = ~~(=)~~ (0.521 ± 0.014)T - (14.96 ± 0.21)‰. The same is true for a correlation found by
875 Rozanski et al. (1982) calculated δ²H = (2.4 ± 0.3)T - (80.5 ± 4.2)‰ (R² = 0.89) at the for the
876 GNIP station Stuttgart, which is located 196 km South of the Schwingbach study area. This
877 relationship is similar to the correlation found for the Schwingbach catchment. Stumpp et al.
878 (2014) analysed long-term precipitation data from meteorological stations across Germany
879 and found that 23 out of 24 tested stations showed a positive long-term temperature trend over
880 time, ~~whereas the precipitation amount was not a key factor for explaining isotope~~
881 ~~distributions or average values in German precipitation. The temperature isotope relationship~~
882 ~~was likewise strongly influenced by seasonality (Stumpp et al., 2014). For the Schwingbach~~
883 ~~catchment, 53% of the events were sampled at mean daily temperatures >10°C, resulting in a~~
884 ~~slight overrepresentation of values measured at warmer temperatures. Nevertheless, the.~~ The

885 observed correspondence between the degree of isotope depletion and the temperature reflects
886 the influence of the temperature effect in the Schwingbach catchment, which mainly appears
887 in continental, middle–high latitudes (Jouzel et al., 1997; Wu et al., 2012). (Jouzel et al.,
888 [1997](#)). Furthermore, the correlation between $\delta^2\text{H}$ in monthly precipitations and local surface
889 air temperature becomes increasingly stronger towards the centre of the continent (Rozanski
890 et al., 1982). [Thus, the observed seasonal differences in precipitation \$\delta\$ -values in the
891 Schwingbach catchment could mainly be attributed to seasonal differences in air temperature
892 and the presence of snow in the winter of 2012–13 \(Fig. 2\).](#)

~~893 Thus, the observed inter-seasonal differences in precipitation δ -values in the Schwingbach
894 catchment could mainly be attributed to seasonal differences in air temperature and water
895 vapour and their effect on evaporation (Schürch et al., 2003) and the presence of snow in the
896 winter of 2012–13 (Fig. 2). This observation is well in agreement with Gat et al. (2001) who
897 stated that for temperate climates the $\delta^{18}\text{O}$ values generally do not vary by more than 1‰
898 inter-annually, and a large part of the spread is caused by variations in the average annual
899 temperature. Moreover, the interior of the continent is obviously far more stable with regard
900 to isotopic inputs than areas under greater influence of Atlantic weather patterns. Perhaps in
901 view of this stability, only few isotope data are available for this region, apart from the
902 general GNIP maps (Bowen and Wilkinson, 2002; Darling, 2004; IAEA, 2014) and recent
903 work (Stumpp et al., 2014), for which this work contributes valuable information.~~

904 Considering ~~d-excess values, it is well known that precipitation~~[Precipitation](#) events
905 originating from oceanic moisture show d-excess values close to +10‰ (Craig, 1961a;
906 Dansgaard, 1964; Wu et al., 2012) and one of the main sources for precipitation in Germany
907 is moisture from the Atlantic Ocean (Stumpp et al., 2014). Lowest values corresponded to
908 summer precipitation events ~~with~~[where](#) evaporation of the [falling](#) raindrops below the cloud
909 base ~~at mean daily air temperatures between 12–18°C occurs~~. Same observations were made
910 by Rozanski et al. (1982) for European GNIP stations. ~~Accordingly, even more negative
911 summer d-excess values were measured at air temperatures around 26–27°C for a study site in
912 Greece (Argiriou and Lykoudis, 2006). Most of the higher values measured in the
913 Schwingbach catchment (>+10‰) appeared in cold seasons (fall/winter) (Fig. 2), similar to d-
914 excess values observed by Wu et al. (2012) for a continental, semi-arid study area in Inner
915 Mongolia (China). Winter snow samples of the Schwingbach catchment with very depleted δ -
916 values showed highest d-excess values, which was again~~[Winter snow samples of the](#)

917 Schwingbach catchment with very depleted δ -values showed highest d-excess values
918 ($>+10\%$), well in agreement with results of Rozanski et al. (1982) for European GNIP
919 stations. Continental precipitation events originating from oceanic moisture can approach d-
920 excess values of $+10\%$ (Wu et al., 2012) (Fig. 2, solid line). Air mass trajectories at
921 intercontinental, southern and eastern regions are suggested to be more stable with less
922 variable moisture sources in these regions compared to sites near the coast (Stumpp et al.,
923 2014). Therefore, rainout histories on the continent itself are more stable (Stumpp et al.,
924 2014). The observed differences in d-excess values between the Schwingbach catchment and
925 the GNIP station Koblenz can be attributed to differences in elevation range and the different
926 regional climatic settings at both sites (Koblenz is located in the relatively warmer Rhine river
927 valley). Further, no amount effect on d-excess could be determined for the Schwingbach
928 catchment, which generally occurs most likely in the tropics (Bony et al., 2008) or for intense
929 convective rain events (Gat et al., 2001) at monsoon dominated sites (Risi et al., 2008).

930 **Isotopes of soil water**

931 **Spatial variability**

932 Determining potential relationships between small scale characteristics such as distance to
933 stream, TWI, and land use on soil isotopic signatures, no tendency of higher TWI values with
934 decreasing distance to stream was obvious. Garvelmann et al. (2012) investigated two
935 hillslopes in a humid 0.9 km^2 catchment in the southern Black Forest (Germany) and found
936 that soil profiles upslope or with a weak affinity for saturation (low TWIs) preserved the
937 precipitation isotopic signal. In our study, the δ -values of top soil and precipitation did not
938 vary significantly statistically (Fig. 4), which is not the case for precipitation and subsoil. A
939 mixing and homogenization of new and old soil water with depth could not clearly be seen in
940 0.5 m soil depth, which would have resulted in a lower standard deviation (Song et al., 2011),
941 but standard deviations of isotopic signatures in top and subsoil were similar (Table 2).
942 Subsoil isotopic values were statistically equal to stream We found no statistically significant
943 connection between the parameters distance to stream, TWI, soil water content, soil texture,
944 pH, and bulk density with the soil isotopic signatures in both soil depths. This was potentially
945 attributed to the small variation in soil textures (mainly clayey silts and loamy sandy silts),
946 bulk densities, and pH values for both soil depths (Table 2). and groundwater isotopic values
947 (Fig. 4) implying that the catchment was under baseflow conditions during the sampling

948 ~~campaign and that capillary rise of groundwater occurred. Nevertheless, the rainfall isotopic~~
949 ~~signal was not directly transferred through the soil to the groundwater body, even so prompt~~
950 ~~groundwater head level raises as a result of rainfall runoff events occurred. This supports the~~
951 ~~assumption of double paradox like catchment behaviour.~~

952 Garvelmann et al. (2012) obtained high resolution $\delta^2\text{H}$ vertical depth profiles of pore water at
953 various points along two fall lines of a pasture hillslope in the ~~southern~~-Black Forest
954 (Germany) by applying the $\text{H}_2\text{O}(\text{liquid})\text{-H}_2\text{O}(\text{vapor})$ equilibration laser spectroscopy method.
955 The authors showed that groundwater was flowing through the soil in the riparian zone
956 (downslope profiles) and dominated streamflow during baseflow conditions. Their
957 comparison indicated that the percentage of pore water soil samples with a very similar
958 stream water $\delta^2\text{H}$ signature is increasing towards the stream channel (Garvelmann et al.,
959 2012). In contrast, we found no such relationship between the distance to stream or TWI and
960 soil isotopic values in the Vollnkirchener Bach subcatchment over various heights above sea
961 level (235–294_m a.s.l.-.) and locations. We attributed this to the gentle, low angle hillslopes
962 and the low subsurface flow contribution in large parts of the catchment.

963 ~~Comparing soil water $\delta^2\text{H}$ values between top and subsoil under different land use units~~
964 ~~showed significant differences ($p \leq 0.05$) under arable and grassland but not under forested~~
965 ~~sites (Fig. 5). This could be explained through the occurrence of vertical preferential flow~~
966 ~~paths and interconnected macropore flow such as continuous root channels or earthworm~~
967 ~~burrows~~In our study, the δ -values of top soil and precipitation did not differ statistically (Fig.
968 4), but for precipitation and subsoil they did. The latter indicates either the influence of
969 evaporation in the topsoil or the mixing with groundwater in the subsoil. However, a mixing
970 and homogenization of new and old soil water with depth could not clearly be seen in 0.5 m
971 soil depth, which would have resulted in a lower standard deviation (Song et al., 2011), but
972 standard deviations of isotopic signatures in top and subsoil were similar (Table 2). Subsoil
973 isotopic values were statistically equal to stream water and groundwater (Fig. 4) implying that
974 capillary rise of groundwater occurred. Overall, the rainfall isotopic signal was not directly
975 transferred through the soil to the groundwater; even so groundwater head level rose promptly
976 after rainfall events. This behaviour reflects the differences of celerity and velocity in the
977 catchment's rainfall-runoff response (McDonnell and Beven, 2014).

978 Soil water $\delta^2\text{H}$ between top and subsoil showed significant differences ($p < 0.05$) under arable
979 and grassland but not under forested sites (Fig. 5). This could be explained through the

980 occurrence of vertical preferential flow paths and interconnected macropore flow (Buttle and
981 McDonald, 2002) characteristic for forested soils (~~Alaoui et al., 2011~~). Alaoui et al. (2011)
982 showed that macropore flow with high interaction with the surrounding soil matrix occurred
983 in forest soils, while macropore flow with low to mixed interaction with the surrounding soil
984 matrix dominates in grassland soils. ~~The authors attributed~~ Seasonal tilling prevents the low
985 efficiency establishment of grassland soil macropores in transporting all water vertically
986 downward to the fine and dense few topsoil layers caused by the land use that limit water flux
987 into the underlying macropores. In general, the upper part of most preferential flow paths
988 under agricultural human impacted soils is restructured annually due to seasonal tillingsites
989 and is regularly done in the Schwingbach catchment, whereas the structure of forest soils,
990 may remain ~~unchanged for years and be~~ uninterrupted throughout the entire soil profile for
991 years (in particular the macropores and biopores) (Alaoui et al., 2011). ~~Considering the bulk~~
992 density in the Schwingbach catchment increasing values from forest (1.10 g cm⁻³) over
993 grassland (1.25 g cm⁻³) to arable land soils (1.41 g cm⁻³) were measured in the top soil. As
994 reported in a study by Price et al. (2010) for North Carolina (USA), soils underlying forest
995 trees generally feature low bulk density in a comparison with soils impacted by human land
996 use. The reduced hydrological connectivity between top and subsoil under arable and
997 grassland observed in the Vollnkirchener Bach subcatchment therefore led to different
998 isotopic signatures (Fig. This is reflected in the bulk density of the soils in the Schwingbach
999 catchment that increases from forests (1.10 g cm⁻³) over grassland (1.25 g cm⁻³) to arable
1000 land (1.41 g cm⁻³) in the top soil. We infer that reduced hydrological connection between top
1001 and subsoil under arable and grassland led to different isotopic signatures (Fig. 5).

1002 ~~Although, vegetation cover has been proven to have often shown~~ an impact on soil water
1003 isotopes (~~Brødersen et al., 2000; Gat, 1996; Li et al., 2007~~)(Gat, 1996), only few data are
1004 available for Central Europe (Darling, 2004). ~~Burger and Seiler (1992)~~ Burger and Seiler
1005 (1992) found that soil water isotopic enrichment under spruce forest in Upper Bavaria was
1006 double that beneath neighbouring arable land. ~~However, but~~ soil ~~water isotopic~~
1007 ~~signatures~~ isotope values were not comparable to groundwater ~~isotope values~~ (~~Burger and~~
1008 ~~Seiler, 1992~~)(Burger and Seiler, 1992). ~~Brødersen et al. (2000) reported the effect of~~
1009 ~~vegetation structure on δ¹⁸O values of rainwater and soil water in the unsaturated zone in~~
1010 ~~southern Germany. In their study,~~ Gehrels et al. (1998) also detected (though only slightly)
1011 heavier isotopic signatures under forested sites in the Netherlands in comparison to non-
1012 forested sites (grassland and heathland). Contrasting, in southern Germany Brodersen et al.

1013 ~~(2000) observed only a negligible effect of~~ throughfall isotopic signatures ~~(of different tree~~
1014 ~~species (spruce and beech) seemed to have a negligible effect~~ on soil water isotopes, since soil
1015 water in the upper layers followed the seasonal trend in the precipitation input and had a very
1016 constant signature in greater depth. ~~In contrast, Gehrels et al. (1998) detected slightly heavier~~
1017 ~~isotopic signatures under forested sites at a field site in the Netherlands in comparison to non-~~
1018 ~~forested sites (grassland and heathland), both showing isotopic signatures comparable to~~
1019 ~~precipitation signals.~~ For the Schwingbach catchment, we conclude that the observed land use
1020 effect in the upper soil column is mainly attributed to different preservation and transmission
1021 of the precipitation input signal. It is most likely not ~~attributed~~attributable to distinguished
1022 throughfall isotopic signatures, impact of evaporation or interception losses, since top soil
1023 water isotopic signals followed the precipitation input signal under all land use units. ~~The~~
1024 ~~precipitation influence smoothed out with depth since soil water isotopes approached~~
1025 ~~groundwater signatures at 0.5 m soil depth.~~

1026 **Seasonal isotope soil profiling**

1027 ~~Soil water was enriched in summer due to evaporation during warmer and drier periods~~
1028 ~~(Darling, 2004).~~ The depth to which soil water isotopes are significantly affected by
1029 evaporation is rarely more than 1–2 m below ground, and often less under temperate climates
1030 (Darling, 2004). In contrast, winter profiles exhibited somewhat greater standard deviations in
1031 comparison to summer isotopic soil profiles, indicative for wetter soils (Fig. 6, lower panels)
1032 and shorter residence times (Thomas et al., 2013). Generally, deeper soil water isotope values
1033 were relatively constant through time and space. Similar findings were made by Foerstel et al.
1034 (1991) on a sandy soil at Juelich, in western Germany ~~and by~~ McConville et al. (2001) under
1035 predominately agriculturally used gley and till soils in Northern Ireland, ~~and~~ Thomas et al.
1036 (2013) ~~likewise observed that soil water isotope samples from shallow soils (≤ 30 cm) were~~
1037 ~~comparable to precipitation isotopic composition, while samples from intermediate soils (40–~~
1038 ~~100 cm) plot near the groundwater average for~~ in a forested catchment ~~located~~ in central
1039 Pennsylvania, USA. Furthermore, Tang and Feng (2001) showed for a sandy loam soil
1040 sampling site in New Hampshire (USA) that the influence of summer precipitation decreased
1041 with increasing depth, and soil at 0.5 m ~~can~~ only receiverreceived water from large
1042 storms. ~~For~~In our summer soil profiles under arable land, precipitation input signals similarly
1043 decreased with depth (Fig. 6, upper left panel). Generally, the replacement of old soil water
1044 with new infiltrating water is dependent on the frequency and intensity of precipitation and

1045 the soil texture, structure, wetness, and water potential of the soil (Li et al., 2007; Tang and
1046 Feng, 2001). ~~It is usually more efficient in a wet year than in a dry year (Tang and Feng,~~
1047 ~~2001). As a result of soil water recharge near the surface~~As a result, the amount of percolating
1048 water decreases with depth and consequently, deeper soil layers have less chance to obtain
1049 new water (Tang and Feng, 2001). ~~Summer and winter profiles show higher water contents in~~
1050 ~~the upper 0.2 m than further down (Fig. 6, lower panels). Furthermore, in~~In the growing
1051 season, the percolation depth is additionally limited by plants' transpiration (Tang and Feng,
1052 2001). For the Schwingbach catchment we conclude that the ~~influence~~percolation of new
1053 ~~percolating~~soil water ~~decreased with depth~~this low as no remarkable seasonality in soil isotopic
1054 signatures was obvious at >0.9 m and constant values were observed through space and time.
1055 Although replications over several years are missing, this result indicates a transit time
1056 through the rooting zone (1m) of approximately one year.

1057 **Linkages between water cycle components**

1058 ~~In general, stream~~Stream water isotopic time series of the Vollnkirchener Bach and
1059 Schwingbach showed ~~(with few exceptions)~~ little deflections through time and, consequently,
1060 provided little insight into time and source-components ~~connectivity. Schürch et al. (2003)~~
1061 ~~likewise observed damped river water isotopic signatures as compared with precipitation~~
1062 ~~isotopic signatures for sampling points of the “Swiss National Network for the Observation of~~
1063 ~~Isotopes in the Water Cycle”.~~ For larger rivers like the Elbe at Torgau in eastern Germany
1064 ~~seasonal isotopic composition varied with an amplitude of 1.5‰ in $\delta^{18}\text{O}$ (Darling, 2004).~~

1065 ~~As described above, MTT calculations did not provide meaningful results. The failure of the~~
1066 ~~MTT estimations is mainly attributed to the little variation in stream water isotopic~~
1067 ~~signatures~~connection. . Just as in the here presented results, Klaus et al. (2015) had difficulties
1068 to apply traditional methods of isotope hydrology (MTT estimation, hydrograph separation) to
1069 their dataset due to the lack of temporal isotopic variation in stream water of a forested low-
1070 mountainous catchment in South Carolina (USA). Furthermore, stable water isotopes can only
1071 be utilised for estimations of younger water (<5 years) ~~(McGuire et al., 2005; Stewart et al.,~~
1072 ~~2010), suggesting that transit times in the Schwingbach catchment are longer than the range~~
1073 ~~used for stable water isotopes.~~

1074 Due to the observed isotopic similarities of stream and groundwater, we ~~assume~~conclude that
1075 groundwater predominantly feeds baseflow: (discharge <10 L·s⁻¹). Even during peak flow

1076 occurring in January 2012, December to April or May 2013, rainfall input did not play a
1077 major role for stream water isotopic composition although fast rainfall-runoff behaviours
1078 were observed by Orłowski et al. (2014). ~~The damped groundwater isotopic signatures~~ Same
1079 observations were made by Jin et al. (2010) ~~for the Red Canyon Creek watershed (Wyoming,~~
1080 ~~USA), indicating good hydraulic connection between surface water and shallow groundwater~~
1081 ~~and by Klaus et al. (2015) for a low mountainous forested watershed in South Carolina~~
1082 ~~(USA), comparable to the Schwingbach catchment. The damped groundwater isotopic~~
1083 ~~signatures, which likewise showed little variation through time, rather~~ seemed to be a mixture
1084 of former lighter precipitation events and snowmelt, since meltwater is known to be depleted
1085 in stable isotopes as compared to ~~the annual mean of~~ precipitation or groundwater (Rohde,
1086 1998); ~~(Figure 3)~~. However, ~~one should be aware that~~ differences in the snow sampling
1087 method (new snow, snow pit layers, meltwater) can affect the isotopic composition (Penna et
1088 al., 2014; Taylor et al., 2001). As groundwater at the observed piezometers in the
1089 Vollnkirchener subcatchment is shallow (Orłowski et al., 2014), the snowmelt signal is
1090 allowed to move rapidly through the soil. Pulses of snowmelt water causing a depletion in
1091 spring and early summer was also observed by other studies (Darling, 2004; Kortelainen and
1092 Karhu, 2004). ~~We therefore assume that groundwater is mainly recharged throughout the~~
1093 ~~winter. Generally, less than 5 to 25% of precipitation infiltrates to the groundwater table in~~
1094 ~~temperate climates; the rest is lost to runoff, evaporation from soils, and transpiration by~~
1095 ~~vegetation (Clark and Fritz, 1997a).~~ ~~We therefore conclude that groundwater is mainly~~
1096 ~~recharged throughout the winter.~~ During spring runoff when soils are saturated, temperatures
1097 are low, and vegetation is inactive, recharge rates are generally highest. In contrast, recharge
1098 is very low during summer when most precipitation is transpired back to the atmosphere
1099 (Clark and Fritz, 1997a). Similarly, O’Driscoll et al. (2005) showed that summer precipitation
1100 does not significantly contribute to recharge in the Spring Creek watershed ~~of central~~
1101 ~~(Pennsylvania, USA)~~ since $\delta^{18}\text{O}$ values in summer precipitation were enriched compared to
1102 mean annual groundwater composition.

1103 Further, Orłowski et al. (2014) showed that influent and effluent conditions (bidirectional
1104 water exchange) occurred simultaneously at different stream sections of the Vollnkirchener
1105 Bach affecting stream and groundwater isotopic compositions, equally. ~~Since groundwater~~
1106 ~~head levels in the Vollnkirchener Bach subcatchment closely followed stream runoff~~
1107 ~~dynamics and responded to stormflow events with rising head levels (Fig. 8), we conclude~~
1108 ~~that bidirectional water exchange between the groundwater body and the Vollnkirchener Bach~~

1109 ~~occurred.~~ Our network map supported this assumption (Fig. 9) as surface water
1110 ~~sampling~~sampling points plotted close to groundwater sampling points (especially to the
1111 sampling points under the meadow and along the stream). ~~However, both water compartments~~
1112 ~~This was also underlined by our groundwater model showing flow directions across the~~
1113 ~~Vollnkirchener Bach. Nevertheless, both stream and groundwater~~ differed significantly from
1114 rainfall isotopic signatures (Table 1). ~~These divergent isotopic signatures but the prompt~~
1115 ~~reaction of the groundwater body to rainfall-runoff events indicate that 'old' groundwater can~~
1116 ~~be released during very short times (Kirchner, 2003).~~ Thus, our catchment showed double
1117 water paradox behaviour as ~~described earlier by~~ per Kirchner (2003) ~~as the~~with fast
1118 ~~releasing~~release of very old water ~~with~~but little variation in tracer concentration. ~~This paradox~~
1119 ~~behaviour could likewise be a reason for the failure of the MTT estimation.~~

1120 Water age dynamics

1121 Our ~~MTT and MRT calculations did not provide meaningful results.~~ Just by comparing mean
1122 precipitation ($\delta^{18}\text{O} = -6.2 \pm 3.1$), stream (e.g. $\delta^{18}\text{O} = -8.4 \pm 0.4$ for the Vollnkirchener Bach),
1123 and groundwater isotopic signatures ($\delta^{18}\text{O} = -8.2 \pm 0.4$ for the meadow) (Table 1), it is obvious
1124 that simple mixing calculations do not work, i.e. showing predominant groundwater
1125 contribution. Same observations were made by Jin et al. (2010) indicating good hydraulic
1126 connectivity between surface water and shallow groundwater. Just as in the here presented
1127 results, Klaus et al. (2015) had difficulties to apply traditional methods of isotope hydrology
1128 (MTT estimation, hydrograph separation) to their dataset due to the lack of temporal isotopic
1129 variation in stream water of a forested low-mountainous catchment in South Carolina (USA).
1130 Furthermore, stable water isotopes can only be utilised for estimations of younger water (<5
1131 years) ~~either~~(Stewart et al., 2010) as they are blind to older contributions (Duvert et al.,
1132 2016). In our catchment, transit times are orders of magnitudes longer than the timescale of
1133 hydrologic response (prompt discharge of old water) (McDonnell et al., 2010) and the range
1134 used for stable water isotopes.

1135 ~~Nevertheless, to still estimate groundwater ages in the Vollnkirchener Bach subcatchment, we~~
1136 ~~established a hydrological model. Our model results suggest that the main groundwater flow~~
1137 ~~direction is towards the stream and the quantity of flowing water is highest near the stream~~
1138 ~~(Fig. 10). This further supports the assumption that stream water is mainly fed by~~
1139 ~~groundwater. Moreover, the simulation underlines the conclusion that the groundwater body~~
1140 ~~and stream water are disconnected from the precipitation cycle, since only 13.3% of cells~~

1141 ~~contained water with an age <1 year. The results of the model reveal a spatially highly~~
1142 ~~heterogeneous age distribution of groundwater throughout the Vollnkirchener Bach~~
1143 ~~subcatchment. The age varies from about two days to more than 100 years with oldest water~~
1144 ~~near the stream. Accurately capturing the transit time of the old water fraction is essential~~
1145 ~~(Duvert et al., 2016) and could previously only be determined via other tracers such as tritium~~
1146 ~~(e.g. Thus, our model provides the opportunity to make use of stable water isotope~~
1147 ~~information along with climate, land use, and soil type data, in combination with a digital~~
1148 ~~elevation map to estimate residence times >5 years. Such long residence times could~~
1149 ~~previously only be determined via other tracers such as tritium (e.g. Michel (1992)). Current~~
1150 ~~studies on mixing assumptions either consider spatial or time-varying MTTs. Heidbüchel et~~
1151 ~~al. (2012) proposed the concept of the master transit time distribution that accounts for the~~
1152 ~~temporal variability of MTT. The time-varying transit time concept of Botter et al. (2011) and~~
1153 ~~van der Velde et al. (2012), was recently reformulated by Harman (2015) so that the storage~~
1154 ~~selection function became a function of the watershed storage and actual time. Instead of~~
1155 ~~quantifying time-variant travel times, our model facilitates the estimation of spatially~~
1156 ~~distributed groundwater ages, which opens up new opportunities to compare groundwater~~
1157 ~~ages from over a range of scales within catchments. It further gives a deeper understanding of~~
1158 ~~the groundwater-surface water connection across the landscape than a classical MTT~~
1159 ~~calculation could provide. Our work complements recent advances in spatially distributed~~
1160 ~~modelling of age distributions through transient groundwater flows (e.g. Gomez and Wilson,~~
1161 ~~2013; Woolfenden and Ginn, 2009). The results of our model reveal a spatially highly~~
1162 ~~heterogeneous age distribution of groundwater throughout the Vollnkirchener Bach~~
1163 ~~subcatchment (ages of 2 days–100 years) with oldest water near the stream. Thus, our model~~
1164 ~~provides the opportunity to make use of stable water isotope information along with climate,~~
1165 ~~land use, and soil type data, in combination with a digital elevation map to estimate residence~~
1166 ~~times >5 years. If stable water isotope information is used alone, it is known to cause a~~
1167 ~~truncation of stream residence time distributions (Stewart et al., 2010). Moreover, our model~~
1168 ~~facilitates the estimation of spatially distributed groundwater ages, which opens up new~~
1169 ~~opportunities to compare groundwater ages from over a range of scales within~~
1170 ~~catchments. Further, our groundwater model suggests that the main groundwater flow direction~~
1171 ~~is towards and across the stream and the quantity of flowing water is highest near the stream~~
1172 ~~(Fig. 10). This further supports the assumption that stream water is mainly fed by older~~
1173 ~~groundwater. Moreover, the simulation underlines the conclusion that the groundwater body~~

1174 and stream water are isotopically disconnected from the precipitation cycle, since only 13% of
1175 cells contained water with and age <1 year.

1176 ~~The observation that gaining and losing stream reaches occur simultaneously along the~~
1177 ~~Vollkirchner Bach could similarly be supported by our model results. However, due to the~~
1178 ~~model assumption of a constant groundwater recharge over the course of a year, no~~
1179 ~~seasonality was simulated. Moreover, model results differ somewhat from the conceptual~~
1180 ~~model of Orłowski et al. (2014). This is due to the fact that the hydrological model only~~
1181 ~~estimates groundwater fluxes but not surface water fluxes. Moreover, no spatial differences in~~
1182 ~~soil properties of the groundwater layer were considered. Nevertheless, as shown by the~~
1183 ~~diverse ages of water in the stream cells and the assumption of spatially gaining conditions,~~
1184 ~~the model confirms that the stream contains water with different transit times. Therefore, the~~
1185 ~~stream water does not have a discrete age, but a distribution of ages due to variable flow paths~~
1186 ~~throughout the subcatchment (Stewart et al., 2010). Heidebüchel et al. (2012) proposed the~~
1187 ~~concept of the master transit time distribution that accounts for temporal variability of MTT.~~
1188 ~~Our model provides a different approach that considers spatial aspects of transit times and~~
1189 ~~gives a much deeper understanding of the groundwater-surface water connectivity across the~~
1190 ~~landscape than a classical MTT calculation could provide.~~

1191 However, our semi-conceptual model approach has also some limitations. During model setup
1192 a series of assumptions and simplifications were made to develop a realistic hydrologic model
1193 without a severe loss in performance. Therefore~~Due to the assumption of a constant~~
1194 groundwater recharge over the course of a year, no seasonality was simulated. Moreover, no
1195 spatial differences in soil properties of the groundwater layer were considered. Further,
1196 several parameters such as the depth of the groundwater body are only rough estimations,
1197 while others like evapotranspiration are based on simulations. Moreover, the groundwater
1198 body is highly simplified since e.g. properties of the simulated aquifer are assumed to be
1199 constant over the subcatchment. ~~However, the complexity of the model is higher than in a~~
1200 ~~simple one-dimensional model (with only one cell and one layer), which results in a better~~
1201 ~~spatial resolution, but lower than in a fully distributed variable-saturated 3D~~
1202 ~~model.~~Nevertheless, as shown by the diverse ages of water in the stream cells and the
1203 assumption of spatially gaining conditions, the model confirms that the stream contains water
1204 with different transit times and supports the assumption that surface and groundwater are
1205 isotopically disconnected from precipitation. Therefore, the stream water does not have a

1206 discrete age, but a distribution of ages due to variable flow paths (Stewart et al., 2010). In
1207 future models a more diverse groundwater body based on small-scale measurements of
1208 aquifer parameters should be implemented. Especially data of saturated hydraulic
1209 conductivity with high spatial resolution, as well as the implementation of a temporal
1210 dynamic groundwater recharge could lead to an enhanced model performance. ~~Nevertheless,~~
1211 ~~our hydrological model enables a good assessment of the groundwater age for the~~
1212 ~~Vollnkirchner Bach subcatchment and supports the assumption that surface and groundwater~~
1213 ~~are disconnected from precipitation.~~

1214 **Conclusions**

1215 Conducting a stable water isotope study in the Schwingbach catchment helped to identify
1216 relationships between precipitation, stream, soil, and groundwater in a developed (managed)
1217 catchment. The close isotopic link between groundwater and the streams revealed that
1218 groundwater controls streamflow. Moreover, it could be shown that groundwater was
1219 predominately recharged during winter but was decoupled from the annual precipitation
1220 cycle. Even so streamflow and groundwater head levels promptly responded to precipitation
1221 inputs, there was no obvious change in their isotopic composition due to rain events ~~(old~~
1222 ~~water paradox behaviour). This was underlined by the fact that no remarkable seasonality in~~
1223 ~~soil isotopic signatures as interface between precipitation and groundwater was obvious at~~
1224 ~~>0.9 m and constant values were observed through space and time.~~

1225 Nevertheless, the lack of temporal variation in stable isotope time series of stream and
1226 groundwater (with few exceptions) limited the application of classical methods of isotope
1227 hydrology, i.e. ~~mean transit time~~ transfer function based MTT estimations. By splitting the
1228 flow path into different compartments (upper and lower vadose zone, groundwater, stream),
1229 we were able to determine, where the water age passes the limit of using stable isotopes for
1230 age calculations. This limit is in the Schwingbach catchment. We therefore set up lower vadose
1231 zone approximately 1–2 m below ground. To estimate the total transit time to the stream, we
1232 set up a hydrological model ~~with CMF to estimate~~ calculating spatially distributed
1233 groundwater ages and flow directions in the Vollnkirchner Bach subcatchment. Our model
1234 result~~results~~ supported the finding that the water in the catchment is >5 years (on average 16
1235 years) and that stream water is mainly fed by groundwater. Our modelling approach was
1236 valuable to overcome the limitations of MTT calculations with traditional methods and/or
1237 models. ~~Thus~~ Further, our dual isotope study in combination with the hydrological model

1238 approach ~~was valuable for determining the connectivity~~enabled the determination of
1239 connection and ~~disconnectivity~~disconnection between different water cycle components.

1240

1241 **Acknowledgements**

1242 The first author acknowledges financial support by the Friedrich-Ebert-Stiftung (Bonn, DE).
1243 Furthermore, this work was supported by the Deutsche Forschungsgemeinschaft under Grant
1244 BR2238/10-1. Thanks also to the student assistants, BSc. and MSc. students for their help
1245 during field sampling campaigns and hydrological modelling. In this regard we like to
1246 acknowledge especially Julia Mechsner, Julia Klöber, and Judith Henkel. We thank Dr.
1247 Christine Stumpp from the Helmholtz Zentrum München for providing us the isotope data
1248 from GNIP station Koblenz.

1249

1250 **References**

- 1251 [Aggarwal, P. K., Gat, J. R. and Froehlich, K. F., Eds.: Isotopes in the Water Cycle: Past,](#)
1252 [Present and Future of a Developing Science, Springer, Dordrecht, Netherlands., 2007.](#)
- 1253 [Alaoui, A., Caduff, U., Gerke, H. H. and Weingartner, R.: Preferential Flow Effects on](#)
1254 [Infiltration and Runoff in Grassland and Forest Soils, Vadose Zone J., 10\(1\), 367,](#)
1255 [doi:10.2136/vzj2010.0076, 2011.](#)
- 1256 [Allan, J. D.: Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems,](#)
1257 [Annu. Rev. Ecol. Evol. Syst., 35, 257–284, 2004.](#)
- 1258 [Araguás-Araguás, L., Froehlich, K. and Rozanski, K.: Deuterium and oxygen-18 isotope](#)
1259 [composition of precipitation and atmospheric moisture, Hydrol. Process., 14\(8\), 1341–1355,](#)
1260 [2000.](#)
- 1261 [Argiriou, A. A. and Lykoudis, S.: Isotopic composition of precipitation in Greece, J. Hydrol.,](#)
1262 [327\(3–4\), 486–495, doi:10.1016/j.jhydrol.2005.11.053, 2006.](#)
- 1263 [Barnes, C. J. and Allison, G. B.: Tracing of water movement in the unsaturated zone using](#)
1264 [stable isotopes of hydrogen and oxygen, J. Hydrol., 100\(1–3\), 143–176, doi:10.1016/0022-](#)
1265 [1694\(88\)90184-9, 1988.](#)
- 1266 [Barthold, F. K., Tyralla, C., Schneider, K., Vaché, K. B., Frede, H. G. and Breuer, L.: How](#)
1267 [many tracers do we need for end-member mixing analysis \(EMMA\)? A sensitivity analysis,](#)
1268 [Water Resour. Res., 47\(8\), doi:10.1029/2011WR010604, 2011.](#)
- 1269 [Birkel, C., Dunn, S. M., Tetzlaff, D. and Soulsby, C.: Assessing the value of high-resolution](#)
1270 [isotope tracer data in the stepwise development of a lumped conceptual rainfall-runoff model,](#)
1271 [Hydrol. Process., 24\(16\), 2335–2348, doi:10.1002/hyp.7763, 2010.](#)
- 1272 [Blasch, K. W. and Bryson, J. R.: Distinguishing Sources of Ground Water Recharge by Using](#)
1273 [\$\delta^2\text{H}\$ and \$\delta^{18}\text{O}\$, Ground Water, 45\(3\), 294–308, doi:10.1111/j.1745-6584.2006.00289.x, 2007.](#)
- 1274 [Bony, S., Risi, C. and Vimeux, F.: Influence of convective processes on the isotopic](#)
1275 [composition \(\$\delta^{18}\text{O}\$ and \$\delta\text{D}\$ \) of precipitation and water vapor in the tropics: 1. Radiative-](#)
1276 [convective equilibrium and Tropical Ocean Global Atmosphere Coupled Ocean Atmosphere](#)
1277 [Response Experiment \(TOGA-COARE\) simulations, J. Geophys. Res. Atmospheres,](#)
1278 [113\(D19\), D19305, doi:10.1029/2008JD009942, 2008.](#)
- 1279 [Botter, G., Bertuzzo, E. and Rinaldo, A.: Transport in the hydrologic response: Travel time](#)
1280 [distributions, soil moisture dynamics, and the old-water paradox, Water Resour. Res., 46\(3\),](#)
1281 [W03514, doi:10.1029/2009WR008371, 2010.](#)
- 1282 [Bowen, G. J. and Wilkinson, B.: Spatial distribution of \$\delta^{18}\text{O}\$ in meteoric precipitation,](#)
1283 [Geology, 30\(4\), 315–318, doi:10.1130/0091-7613\(2002\)030<0315:SDOIM>2.0.CO;2,](#)
1284 [2002.](#)

- 1285 ~~Brodersen, C., Pohl, S., Lindenlaub, M., Leibundgut, C. and Wilpert, K. v: Influence of~~
1286 ~~vegetation structure on isotope content of throughfall and soil water, *Hydrol. Process.*, 14(8),~~
1287 ~~1439–1448, doi:10.1002/1099-1085(20000615)14:8<1439::AID-HYP985>3.0.CO;2-3, 2000.~~
- 1288 ~~Burger, H. M. and Seiler, K. P.: Evaporation from soil water under humid climate conditions~~
1289 ~~and its impact on deuterium and ¹⁸O concentrations in groundwater., edited by International~~
1290 ~~Atomic Energy Agency, pp. 674–678, International Atomic Energy Agency, Vienna, Austria.,~~
1291 ~~1992.~~
- 1292 ~~Buttle, J. M.: Isotope hydrograph separations and rapid delivery of pre-event water from~~
1293 ~~drainage basins, *Prog. Phys. Geogr.*, 18(1), 16–41, doi:10.1177/030913339401800102, 1994.~~
- 1294 ~~Buttle, J. M.: Isotope Hydrograph Separation of Runoff Sources, in *Encyclopedia of*~~
1295 ~~Hydrological Sciences, edited by M. G. Anderson, p. 10:116, John Wiley & Sons, Ltd,~~
1296 ~~Chichester, Great Britain, 2006.~~
- 1297 ~~Buttle, J. M. and McDonald, D. J.: Coupled vertical and lateral preferential flow on a forested~~
1298 ~~slope, *Water Resour. Res.*, 38(5), 18–1, doi:10.1029/2001WR000773, 2002.~~
- 1299 ~~Choi, S. H.: Characteristics of hydrogeologic units of groundwater resources in the Republic~~
1300 ~~of Korea, in *Engineering Geology and the Environment*, vol. 4, edited by P. G. Marinos, G. C.~~
1301 ~~Koukis, G. C. Tsiambaos, and G. C. Stournaras, pp. 1219–1224, CRC Press, Florida, USA.,~~
1302 ~~1997.~~
- 1303 ~~Clark, I. D. and Fritz, P.: Groundwater, in *Environmental Isotopes in Hydrogeology*, p.80,~~
1304 ~~CRC Press, Florida, FL, USA., 1997a.~~
- 1305 ~~Clark, I. D. and Fritz, P.: Methods for Field Sampling, in *Environmental Isotopes in*~~
1306 ~~Hydrogeology, p. 283, CRC Press, Florida, FL, USA., 1997b.~~
- 1307 ~~Clark, I. D. and Fritz, P.: The Environmental Isotopes, in *Environmental Isotopes in*~~
1308 ~~Hydrogeology, pp. 2–34, CRC Press, Florida, FL, USA., 1997c.~~
- 1309 ~~Cooper, L. W.: Isotopic Fractionation in Snow Cover, in *Isotope Tracers in Catchment*~~
1310 ~~Hydrology, edited by C. Kendall and J. J. McDonnell, pp. 119–136, Elsevier, Amsterdam,~~
1311 ~~Netherlands., 1998.~~
- 1312 ~~Craig, H.: Isotopic Variations in Meteoric Waters, *Science*, 133(3465), 1702–1703,~~
1313 ~~doi:10.1126/science.133.3465.1702, 1961a.~~
- 1314 ~~Craig, H.: Standard for reporting concentrations of deuterium and oxygen-18 in natural~~
1315 ~~waters, *Science*, 133(3467), 1833, 1961b.~~
- 1316 ~~Csardi, G. and Nepusz, T.: The igraph software package for complex network research,~~
1317 ~~*Complex Systems* 1695. [online] Available from: <http://igraph.sf.net> (Accessed 24 December~~
1318 ~~2015), 2006.~~
- 1319 ~~Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16(4), 436–468, doi:10.1111/j.2153-~~
1320 ~~3490.1964.tb00181.x, 1964.~~

- 1321 Darling, W. G.: Hydrological factors in the interpretation of stable isotopic proxy data present
1322 and past: a European perspective, *Quat. Sci. Rev.*, 23(7-8), 743-770,
1323 doi:10.1016/j.quascirev.2003.06.016, 2004.
- 1324 Day, T. J.: On the precision of salt dilution gauging, *J. Hydrol.*, 31(3-4), 293-306,
1325 doi:10.1016/0022-1694(76)90130-X, 1976.
- 1326 DWD: Deutscher Wetterdienst Wetter und Klima, Bundesministerium für Verkehr und
1327 digitale Infrastruktur, [online] Available from: <http://dwd.de/> (Accessed 17 February 2014),
1328 2014.
- 1329 Eijkelkamp: RBC flumes, [online] Available from:
1330 [http://en.eijkelkamp.com/products/water/hydrological-research/water-discharge-](http://en.eijkelkamp.com/products/water/hydrological-research/water-discharge-measurements/rbc-flumes.htm)
1331 [measurements/rbc-flumes.htm](http://en.eijkelkamp.com/products/water/hydrological-research/water-discharge-measurements/rbc-flumes.htm) (Accessed 13 January 2014), 2014.
- 1332 Foerstel, H., Frinken, J., Huetzen, H., Lembrich, D. and Puetz, T.: Application of H₂¹⁸O as a
1333 tracer of water flow in soil, *International Atomic Energy Agency, Vienna, Austria*, 1991.
- 1334 Froehlich, K., Gibson, J. J. and Aggarwal, P. K.: Deuterium excess in precipitation and its
1335 climatological significance, in *Study of Environmental Changes Using Isotope Techniques*,
1336 pp. 54-65, C&S Paper Series 13/P, International Atomic Energy Agency, Vienna, Austria.,
1337 2002.
- 1338 Garvelmann, J., Kuells, C. and Weiler, M.: A porewater-based stable isotope approach for the
1339 investigation of subsurface hydrological processes, *Hydrol. Earth Syst. Sci.*, 16(2), 631-640,
1340 doi:10.5194/hess-16-631-2012, 2012.
- 1341 Gat, J. R.: Oxygen and Hydrogen Isotopes in the Hydrologic Cycle, *Annu. Rev. Earth Planet.*
1342 *Sci.*, 24(1), 225-262, doi:10.1146/annurev.earth.24.1.225, 1996.
- 1343 Gat, J., R., Mook, W. G. and Meijer, H. A. J.: Environmental isotopes in the hydrological
1344 cycle: Principles and Applications, edited by W. G. Mook, International Hydrological
1345 Programme; United Nations Educational, Scientific and Cultural Organization and
1346 International Atomic Energy Agency, Paris, France., 2001.
- 1347 Gehrels, J. C., Peeters, J. E. M., de Vries, J. J. and Dekkers, M.: The mechanism of soil-water
1348 movement as inferred from ¹⁸O stable isotope studies, *Hydrol. Sci. J.*, 43(4), 579-594,
1349 doi:10.1080/02626669809492154, 1998.
- 1350 Genreux, D. P. and Hooper, R. P.: Oxygen and Hydrogen Isotopes in Rainfall-Runoff
1351 Studies, in *Isotope Tracers in Catchment Hydrology*, edited by C. Kendall and J. J.
1352 McDonnell, pp. 319-346, Elsevier, Amsterdam, Netherlands., 1998.
- 1353 Goller, R., Wilcke, W., Leng, M. J., Tobschall, H. J., Wagner, K., Valarezo, C. and Zech, W.:
1354 Tracing water paths through small catchments under a tropical montane rain forest in south
1355 Ecuador by an oxygen isotope approach, *J. Hydrol.*, 308(1-4), 67-80,
1356 doi:10.1016/j.jhydrol.2004.10.022, 2005.
- 1357 Gonfiantini, R., Fröhlich, K., Araguás-Araguás, L. and Rozanski, K.: Isotopes in
1358 Groundwater Hydrology, in *Isotope Tracers in Catchment Hydrology*, edited by C. Kendall
1359 and J. J. McDonnell, pp. 203-246, Elsevier, Amsterdam, Netherlands., 1998.

- 1360 Goni, I. B.: Tracing stable isotope values from meteoric water to groundwater in the
1361 southwestern part of the Chad basin, *Hydrogeol. J.*, 14(5), 742–752, doi:10.1007/s10040-005-
1362 0469-y, 2006.
- 1363 Gordon, L. J., Finlayson, C. M. and Falkenmark, M.: Managing water in agriculture for food
1364 production and other ecosystem services, *Agric. Water Manag.*, 97(4), 512–519,
1365 doi:10.1016/j.agwat.2009.03.017, 2010.
- 1366 Heidebuechel, I., Troch, P. A., Lyon, S. W. and Weiler, M.: The master transit time distribution
1367 of variable flow systems, *Water Resour. Res.*, 48(6), W06520, doi:10.1029/2011WR011293,
1368 2012.
- 1369 Hoeg, S., Uhlenbrook, S. and Leibundgut, C.: Hydrograph separation in a mountainous
1370 catchment—combining hydrochemical and isotopic tracers, *Hydrol. Process.*, 14(7), 1199–
1371 1216, doi:10.1002/(SICI)1099-1085(200005)14:7<1199::AID-HYP35>3.0.CO;2-K, 2000.
- 1372 Hollander, M., Wolfe, D. A. and Chieken, E.: *Nonparametric Statistical Methods*, John Wiley
1373 & Sons, New York, NY, USA., 2013.
- 1374 Hsieh, J. C. C., Chadwick, O. A., Kelly, E. F. and Savin, S. M.: Oxygen isotopic composition
1375 of soil water: quantifying evaporation and transpiration, *Geoderma*, 82(1–3), 269–293,
1376 doi:10.1016/S0016-7061(97)00105-5, 1998.
- 1377 IAEA: International Atomic Energy Agency: Water Resources Programme—Global Network
1378 of Isotopes in Precipitation, [online] Available from: [http://www-
1379 naweb.iaea.org/naweb/ih/IHS_resources_gnip.html](http://www-naweb.iaea.org/naweb/ih/IHS_resources_gnip.html) (Accessed 11 August 2014), 2014.
- 1380 Ivkovic, K. M.: A top-down approach to characterise aquifer–river interaction processes, *J.*
1381 *Hydrol.*, 365(3–4), 145–155, doi:10.1016/j.jhydrol.2008.11.021, 2009.
- 1382 Jin, L., Siegel, D. I., Lautz, L. K., Mitchell, M. J., Dahms, D. E. and Mayer, B.: Calcite
1383 precipitation driven by the common ion effect during groundwater–surface water mixing: A
1384 potentially common process in streams with geologic settings containing gypsum, *Geol. Soc.
1385 Am. Bull.*, 122(7/8), B30011.1, doi:10.1130/B30011.1, 2010.
- 1386 Jin, L., Siegel, D. I., Lautz, L. K. and Lu, Z.: Identifying streamflow sources during spring
1387 snowmelt using water chemistry and isotopic composition in semi-arid mountain streams, *J.*
1388 *Hydrol.*, 470–471, 289–301, doi:10.1016/j.jhydrol.2012.09.009, 2012.
- 1389 Jolliffe, I. T.: *Principal Component Analysis*, 2nd ed., Springer Science & Business Media,
1390 New York, NY, USA., 2002.
- 1391 Jouzel, J., Alley, R. B., Cuffey, K. M., Dansgaard, W., Grootes, P., Hoffmann, G., Johnsen, S.
1392 J., Koster, R. D., Peel, D., Shuman, C. A., Stievenard, M., Stuiver, M. and White, J.: Validity
1393 of the temperature reconstruction from water isotopes in ice cores, *J. Geophys. Res.*, 102,
1394 26471–26487, doi:10.1029/97JC01283, 1997.
- 1395 Kendall, C. and Caldwell, E. A.: *Fundamentals of Isotope Geochemistry*, in *Isotope Tracers in
1396 Catchment Hydrology*, edited by C. Kendall and J. J. McDonnell, pp. 51–86, Elsevier,
1397 Amsterdam, Netherlands., 1998.

- 1398 Kendall, C. and Coplen, T. B.: Distribution of oxygen-18 and deuterium in river waters across
1399 the United States, *Hydrol. Process.*, 15, 1363–1393, doi:10.1002/hyp.217, 2001.
- 1400 Kirchner, J. W.: A double paradox in catchment hydrology and geochemistry, *Hydrol.*
1401 *Process.*, 17(4), 871–874, doi:10.1002/hyp.5108, 2003.
- 1402 Klapper, H.: Historical Change of Large Alluvial Rivers: Western Europe, edited by G. E.
1403 Petts, H. Möller, and A. L. Roux, John Wiley and Sons Ltd., Chichester, Great Britain, 1990.
- 1404 Klaus, J., McDonnell, J. J., Jackson, C. R., Du, E. and Griffiths, N. A.: Where does
1405 streamwater come from in low-relief forested watersheds? A dual-isotope approach, *Hydrol.*
1406 *Earth Syst. Sci.*, 19(1), 125–135, 2015.
- 1407 Koeniger, P., Leibundgut, C. and Stiehler, W.: Spatial and temporal characterisation of stable
1408 isotopes in river water as indicators of groundwater contribution and confirmation of
1409 modelling results; a study of the Weser river, Germany, *Isotopes Environ. Health Stud.*, 45(4),
1410 289–302, doi:10.1080/10256010903356953, 2009.
- 1411 Koivusalo, H., Karvonen, T. and Lepistö, A.: A quasi-three-dimensional model for predicting
1412 rainfall-runoff processes in a forested catchment in Southern Finland, *Hydrol. Earth Syst.*
1413 *Sci.*, 4(1), 65–78, doi:10.5194/hess-4-65-2000, 1999.
- 1414 Kolaczyk, E. D.: Statistical Analysis of Network Data with R, Springer Science & Business
1415 Media, New York, NY, USA., 2014.
- 1416 Kortelainen, N. M. and Karhu, J. A.: Regional and seasonal trends in the oxygen and
1417 hydrogen isotope ratios of Finnish groundwaters: a key for mean annual precipitation, *J.*
1418 *Hydrol.*, 285(1–4), 143–157, doi:10.1016/j.jhydrol.2003.08.014, 2004.
- 1419 Kraft, P., Vaché, K. B., Frede, H. G. and Breuer, L.: CMF: A Hydrological Programming
1420 Language Extension For Integrated Catchment Models, *Environ. Model. Softw.*, 26(6), 828–
1421 830, doi:10.1016/j.envsoft.2010.12.009, 2011.
- 1422 Ladouche, B., Probst, A., Viville, D., Idir, S., Baqu'e, D., Loubet, M., Probst, J. L. and
1423 Bariaac, T.: Hydrograph separation using isotopic, chemical and hydrological approaches
1424 (Strengbach catchment, France), *J. Hydrol.*, 242(3), 255–274, 2001.
- 1425 Lai, C. T. and Ehleringer, J. R.: Deuterium excess reveals diurnal sources of water vapor in
1426 forest air, *Oecologia*, 165(1), 213–223, doi:10.1007/s00442-010-1721-2, 2011.
- 1427 Lauer, F., Frede, H. G. and Breuer, L.: Uncertainty assessment of quantifying spatially
1428 concentrated groundwater discharge to small streams by distributed temperature sensing,
1429 *Water Resour. Res.*, 49(1), 400–407, doi:10.1029/2012WR012537, 2013.
- 1430 Leen, J. B., Berman, E. S. F., Liebson, L. and Gupta, M.: Spectral contaminant identifier for
1431 off-axis integrated cavity output spectroscopy measurements of liquid water isotopes, *Rev.*
1432 *Sci. Instrum.*, 83(4), doi:10.1063/1.4704843, 2012.
- 1433 LGR: Los Gatos Research, Greenhouse Gas, isotope and trace gas analyzers, [online]
1434 Available from: <http://www.lgrinc.com/> (Accessed 5 February 2013), 2013.

- 1435 [Liebminger, A., Haberhauer, G., Papesch, W., Heiss, G. and others: Footprints of climate in](#)
1436 [groundwater and precipitation, *Hydrol. Earth Syst. Sci.*, 11\(2\), 785–791, doi:10.5194/hess-11-](#)
1437 [785-2007, 2007.](#)
- 1438 [Li, F., Song, X., Tang, C., Liu, C., Yu, J. and Zhang, W.: Tracing infiltration and recharge](#)
1439 [using stable isotope in Taihang Mt., North China, *Environ. Geol.*, 53\(3\), 687–696,](#)
1440 [doi:10.1007/s00254-007-0683-0, 2007.](#)
- 1441 [Maloszewski, P. and Zuber, A.: Lumped parameter models for interpretation of](#)
1442 [environmental tracer data, in *Manual on Mathematical Models in Isotope Hydrogeology,*](#)
1443 [International Atomic Energy Agency, Vienna, Austria., 1996.](#)
- 1444 [Maloszewski, P. and Zuber, A.: Manual on lumped parameter models used for the](#)
1445 [interpretation of environmental tracer data in groundwaters, in *Use of isotopes for analyses of*](#)
1446 [flow and transport dynamics in groundwater systems, p. 50, International Atomic Energy](#)
1447 [Agency, Vienna, Austria., 2002.](#)
- 1448 [Maule, C. P., Chanaszyk, D. S. and Muehlenbachs, K.: Isotopic determination of snow water](#)
1449 [contribution to soil water and groundwater, *J. Hydrol.*, 155\(1–2\), 73–91, doi:10.1016/0022-](#)
1450 [1694\(94\)90159-7, 1994.](#)
- 1451 [Mazor, E.: Part I. Physical and Geological Concepts, 2. Basic Hydrological Concepts, in](#)
1452 [Chemical and Isotopic Groundwater Hydrology, pp. 15–48, CRC Press, Boca Raton, Florida,](#)
1453 [USA., 2003.](#)
- 1454 [McConville, C., Kalin, R. m., Johnston, H. and McNeill, G. w.: Evaluation of Recharge in a](#)
1455 [Small Temperate Catchment Using Natural and Applied \$\delta^{18}\text{O}\$ Profiles in the Unsaturated](#)
1456 [Zone, *Ground Water*, 39\(4\), 616–623, doi:10.1111/j.1745-6584.2001.tb02349.x, 2001.](#)
- 1457 [McDonnell, J. J., Sivapalan, M., Vaché, K., Dunn, S., Grant, G., Haggerty, R., Hinz, C.,](#)
1458 [Hooper, R., Kirchner, J., Roderick, M. L. and others: Moving beyond heterogeneity and](#)
1459 [process complexity: a new vision for watershed hydrology, *Water Resour. Res.*, 43\(7\),](#)
1460 [W07301, 2007.](#)
- 1461 [McGuire, K. ., DeWalle, D. . and Gburek, W. .: Evaluation of mean residence time in](#)
1462 [subsurface waters using oxygen 18 fluctuations during drought conditions in the mid-](#)
1463 [Appalachians, *J. Hydrol.*, 261\(1–4\), 132–149, doi:10.1016/S0022-1694\(02\)00006-9, 2002.](#)
- 1464 [McGuire, K. J., McDonnell, J. J., Weiler, M., Kendall, C., McGlynn, B. L., Welker, J. M. and](#)
1465 [Seibert, J.: The role of topography on catchment scale water residence time, *Water Resour.*](#)
1466 [Res., 41\(5\), W05002, doi:10.1029/2004WR003657, 2005.](#)
- 1467 [Michel, R. L.: Residence times in river basins as determined by analysis of long-term tritium](#)
1468 [records, *J. Hydrol.*, 130\(1\), 367–378, doi:10.1016/0022-1694\(92\)90117-E, 1992.](#)
- 1469 [Mook, W. G., Ed.: *Environmental Isotopes in the Hydrological Cycle: Principles and*](#)
1470 [Applications, International Hydrological Programme; United Nations Educational, Scientific](#)
1471 [and Cultural Organization and International Atomic Energy Agency, Paris, Vienna., 2001.](#)

- 1472 Mook, W. G., Groeneveld, D. J., Brouwn, A. E. and Ganswijk, A. J. van: Analysis of a run-
1473 off hydrograph by means of natural ^{18}O , in *Isotope techniques in groundwater hydrology*, vol.
1474 1, pp. 159–169, International Atomic Energy Agency, Vienna, Austria., 1974.
- 1475 Moore, R. D.: Introduction to Salt Dilution Gauging for Streamflow Measurement: Part 1,
1476 *Streamline Watershed Manag. Bull.*, 7(4), 20–23, 2004.
- 1477 Munyaneza, O., Wenninger, J. and Uhlenbrook, S.: Identification of runoff generation
1478 processes using hydrometric and tracer methods in a meso-scale catchment in Rwanda,
1479 *Hydrol. Earth Syst. Sci.*, 16(7), 1991–2004, doi:10.5194/hess-16-1991-2012, 2012.
- 1480 Neal, C. and Rosier, P. T. W.: Chemical studies of chloride and stable oxygen isotopes in two
1481 conifer afforested and moorland sites in the British uplands, *J. Hydrol.*, 115(1–4), 269–283,
1482 doi:10.1016/0022-1694(90)90209-G, 1990.
- 1483 O'Driscoll, M., Clinton, S., Jefferson, A., Manda, A. and McMillan, S.: Urbanization Effects
1484 on Watershed Hydrology and In-Stream Processes in the Southern United States, *Water*, 2(3),
1485 605–648, doi:10.3390/w2030605, 2010.
- 1486 O'Driscoll, M. A., DeWalle, D. R., McGuire, K. J. and Gburek, W. J.: Seasonal ^{18}O variations
1487 and groundwater recharge for three landscape types in central Pennsylvania, USA, *J. Hydrol.*,
1488 303(1–4), 108–124, doi:10.1016/j.jhydrol.2004.08.020, 2005.
- 1489 Orłowski, N., Frede, H. G., Brüggemann, N. and Breuer, L.: Validation and application of a
1490 cryogenic vacuum extraction system for soil and plant water extraction for isotope analysis, *J.*
1491 *Sens. Sens. Syst.*, 2(2), 179–193, doi:10.5194/jsss-2-179-2013, 2013.
- 1492 Orłowski, N., Lauer, F., Kraft, P., Frede, H. G. and Breuer, L.: Linking Spatial Patterns of
1493 Groundwater Table Dynamics and Streamflow Generation Processes in a Small Developed
1494 Catchment, *Water*, 6(10), 3085–3117, doi:10.3390/w6103085, 2014.
- 1495 Penna, D., Ahmad, M., Birks, S. J., Bouchaou, L., Brenič, M., Butt, S., Holko, L., Jeelani,
1496 G., Martínez, D. E., Melikadze, G., Shanley, J. B., Sokratov, S. A., Stadnyk, T., Sugimoto, A.
1497 and Vreča, P.: A new method of snowmelt sampling for water stable isotopes, *Hydrol.*
1498 *Process.*, 28(22), 5637–5644, doi:10.1002/hyp.10273, 2014.
- 1499 Phillips, D. L. and Gregg, J. W.: Source partitioning using stable isotopes: coping with too
1500 many sources, *Oecologia*, 136(2), 261–269, doi:10.1007/s00442-003-1218-3, 2003.
- 1501 Pierce, S. C., Kröger, R. and Pezeshki, R.: Managing Artificially Drained Low Gradient
1502 Agricultural Headwaters for Enhanced Ecosystem Functions, *Biology*, 1(3), 794–856,
1503 doi:10.3390/biology1030794, 2012.
- 1504 Price, K., Jackson, C. R. and Parker, A. J.: Variation of surficial soil hydraulic properties
1505 across land uses in the southern Blue Ridge Mountains, North Carolina, USA, *J. Hydrol.*,
1506 383(3–4), 256–268, doi:10.1016/j.jhydrol.2009.12.041, 2010.
- 1507 Qu, Y. and Duffy, C. J.: A semidiscrete finite volume formulation for multiprocess watershed
1508 simulation, *Water Resour. Res.*, 43(8), W08419, doi:10.1029/2006WR005752, 2007.

1509 Rinaldo, A., Benettin, P., Harman, C. J., Hrachowitz, M., McGuire, K. J., van der Velde, Y.,
1510 Bertuzzo, E. and Botter, G.: Storage selection functions: A coherent framework for
1511 quantifying how catchments store and release water and solutes, *Water Resour. Res.*, 51(6),
1512 4840–4847, doi:10.1002/2015WR017273, 2015.

1513 Risi, C., Bony, S., Vimeux, F., Descroix, L., Ibrahim, B., Lebreton, E., Mamadou, I. and
1514 Sultan, B.: What controls the isotopic composition of the African monsoon precipitation?
1515 Insights from event-based precipitation collected during the 2006 AMMA field campaign,
1516 *Geophys. Res. Lett.*, 35(24), L24808, doi:10.1029/2008GL035920, 2008.

1517 Rockström, J., Falkenmark, M., Allan, T., Folke, C., Gordon, L., Jägerskog, A., Kumm, M.,
1518 Lannerstad, M., Meybeck, M., Molden, D., Postel, S., Savenije, H. h. g., Svedin, U., Turton,
1519 A. and Varis, O.: The unfolding water drama in the Anthropocene: towards a resilience-based
1520 perspective on water for global sustainability, *Ecohydrol.*, 7(5), 1249–1261,
1521 doi:10.1002/eco.1562, 2014.

1522 Rodgers, P., Soulsby, C. and Waldron, S.: Stable isotope tracers as diagnostic tools in
1523 upscaling flow path understanding and residence time estimates in a mountainous mesoscale
1524 catchment, *Hydrol. Process.*, 19(11), 2291–2307, doi:10.1002/hyp.5677, 2005a.

1525 Rodgers, P., Soulsby, C., Waldron, S. and Tetzlaff, D.: Using stable isotope tracers to assess
1526 hydrological flow paths, residence times and landscape influences in a nested mesoscale
1527 catchment, *Hydrol. Earth Syst. Sci.*, 9(3), 139–155, 2005b.

1528 Rohde, A.: Snowmelt-Dominated Systems, in *Isotope Tracers in Catchment Hydrology*,
1529 edited by C. Kendall and J. J. McDonnell, pp. 391–433, Elsevier, Amsterdam, Netherlands.,
1530 1998.

1531 Rothfuss, Y., Biron, P., Braud, I., Canale, L., Durand, J. L., Gaudet, J. P., Richard, P.,
1532 Vauclin, M. and Bariac, T.: Partitioning evapotranspiration fluxes into soil evaporation and
1533 plant transpiration using water stable isotopes under controlled conditions, *Hydrol. Process.*,
1534 24(22), 3177–3194, doi:10.1002/hyp.7743, 2010.

1535 Rothfuss, Y., Braud, I., Le Moine, N., Biron, P., Durand, J. L., Vauclin, M. and Bariac, T.:
1536 Factors controlling the isotopic partitioning between soil evaporation and plant transpiration:
1537 Assessment using a multi-objective calibration of SiSPAT-Isotope under controlled
1538 conditions, *J. Hydrol.*, 442–443, 75–88, doi:10.1016/j.jhydrol.2012.03.041, 2012.

1539 Rozanski, K., Sonntag, C. and Münnich, K. O.: Factors controlling stable isotope composition
1540 of European precipitation, *Tellus*, 34(2), 142–150, doi:10.1111/j.2153-3490.1982.tb01801.x,
1541 1982.

1542 Rozanski, K., Araguás-Araguás, L. and Gonfiantini, R.: Isotopic Patterns in Modern Global
1543 Precipitation, in *Climate Change in Continental Isotopic Records*, edited by P. K. Swart, K. C.
1544 Lohmann, J. Mckenzie, and S. Savin, pp. 1–36, American Geophysical Union, Washington,
1545 D. C., US. [online] Available from:
1546 <http://onlinelibrary.wiley.com/doi/10.1029/GM078p0001/summary> (Accessed 13 August
1547 2014), 1993.

1548 Rozanski, K., Froehlich, K. and Mook, W. G.: in *Environmental isotopes in the hydrological*
1549 *cycle: Principles and Applications*, vol. 3, International Hydrological Programme; United

- 1550 Nations Educational, Scientific and Cultural Organization and International Atomic Energy
1551 Agency, Paris, Vienna., 2001.
- 1552 Schultz, N. M., Griffis, T. J., Lee, X. and Baker, J. M.: Identification and correction of
1553 spectral contamination in $^2\text{H}/^1\text{H}$ and $^{18}\text{O}/^{16}\text{O}$ measured in leaf, stem, and soil water, *Rapid*
1554 *Commun. Mass Spectrom.*, 25(21), 3360–3368, doi:10.1002/rem.5236, 2011.
- 1555 Schürch, M., Kozel, R., Schotterer, U. and Tripet, J. P.: Observation of isotopes in the water
1556 eyele the Swiss National Network (NISOT), *Environ. Geol.*, 45(1), 1–11,
1557 doi:10.1007/s00254-003-0843-9, 2003.
- 1558 Sklash, M. G.: Environmental isotope studies of storm and snowmelt runoff generation, in
1559 *Process studies in hillslope hydrology*, edited by M. G. Anderson, pp. 410–435, Wiley, New
1560 York, NY, USA., 1990.
- 1561 Sklash, M. G. and Farvolden, R. N.: The Role Of Groundwater In Storm Runoff, in
1562 *Developments in Water Science*, vol. 12, edited by W. B. and D. A. Stephenson, pp. 45–65,
1563 Elsevier. [online] Available from:
1564 <http://www.sciencedirect.com/science/article/pii/S0167564809700097> (Accessed 8 October
1565 2015), 1979.
- 1566 Song, X., Wang, P., Yu, J., Liu, X., Liu, J. and Yuan, R.: Relationships between precipitation,
1567 soil water and groundwater at Chongling catchment with the typical vegetation cover in the
1568 Taihang mountainous region, China, *Environ. Earth Sci.*, 62(4), 787–796,
1569 doi:10.1007/s12665-010-0566-7, 2011.
- 1570 Stewart, M. K., Morgenstern, U. and McDonnell, J. J.: Truncation of stream residence time:
1571 how the use of stable isotopes has skewed our concept of streamwater age and origin, *Hydrol.*
1572 *Process.*, 24(12), 1646–1659, doi:10.1002/hyp.7576, 2010.
- 1573 Stumpp, C., Klaus, J. and Stiehler, W.: Analysis of long-term stable isotopic composition in
1574 German precipitation, *J. Hydrol.*, 517, 351–361, doi:10.1016/j.jhydrol.2014.05.034, 2014.
- 1575 Tang, K. and Feng, X.: The effect of soil hydrology on the oxygen and hydrogen isotopic
1576 compositions of plants' source water, *Earth Planet. Sci. Lett.*, 185(3–4), 355–367, 2001.
- 1577 Taylor, S., Feng, X., Kirchner, J. W., Osterhuber, R., Klaue, B. and Renshaw, C. E.: Isotopic
1578 evolution of a seasonal snowpack and its melt, *Water Resour. Res.*, 37(3), 759–769,
1579 doi:10.1029/2000WR900341, 2001.
- 1580 Thomas, E. M., Lin, H., Duffy, C. J., Sullivan, P. L., Holmes, G. H., Brantley, S. L. and Jin,
1581 L.: Spatiotemporal Patterns of Water Stable Isotope Compositions at the Shale Hills Critical
1582 Zone Observatory: Linkages to Subsurface Hydrologic Processes, *Vadose Zone J.*, 12(4), 0,
1583 doi:10.2136/vzj2013.01.0029, 2013.
- 1584 Timbe, E., Windhorst, D., Crespo, P., Frede, H. G., Feyen, J. and Breuer, L.: Understanding
1585 uncertainties when inferring mean transit times of water through tracer-based lumped-
1586 parameter models in Andean tropical montane cloud forest catchments, *Hydrol. Earth Syst.*
1587 *Sci.*, 18(4), 1503–1523, doi:10.5194/hess-18-1503-2014, 2014.

1588 [UNEP: United Nations Environment Programme: Global Environment Outlook 3: Past,](#)
1589 [Present and Future Perspectives, Earthscan Publications Ltd, London, United Kingdom,](#)
1590 [2002.](#)

1591 [Wallis, W. D.: A Beginner's Guide to Graph Theory, Birkhäuser Boston, Boston, MA, USA,](#)
1592 [2007.](#)

1593 [Wang, X. F. and Yakir, D.: Using stable isotopes of water in evapotranspiration studies,](#)
1594 [Hydrol. Process., 14\(8\), 1407–1421, doi:10.1002/1099-1085\(20000615\)14:8<1407::AID-](#)
1595 [HYP992>3.0.CO;2-K, 2000.](#)

1596 [Wassenaar, L. I., Athanasopoulos, P. and Hendry, M. J.: Isotope hydrology of precipitation,](#)
1597 [surface and ground waters in the Okanagan Valley, British Columbia, Canada, J. Hydrol.,](#)
1598 [411\(1–2\), 37–48, doi:10.1016/j.jhydrol.2011.09.032, 2011.](#)

1599 [Whitefield, P.: Chapter 3: Soil, in Earth Care Manual: A Permaculture Handbook for Britain](#)
1600 [& Other Temperate Climates, pp. 39–66, Permanent Publications, East Meon, UK., 2004.](#)

1601 [Windhorst, D., Waltz, T., Timbe, E., Frede, H. G. and Breuer, L.: Impact of elevation and](#)
1602 [weather patterns on the isotopic composition of precipitation in a tropical montane rainforest,](#)
1603 [Hydrol. Earth Syst. Sci., 17\(1\), 409–419, doi:10.5194/hess-17-409-2013, 2013.](#)

1604 [Windhorst, D., Kraft, P., Timbe, E., Frede, H. G. and Breuer, L.: Stable water isotope tracing](#)
1605 [through hydrological models for disentangling runoff generation processes at the hillslope](#)
1606 [scale, Hydrol. Earth Syst. Sci., 18\(10\), 4113–4127, doi:10.5194/hess-18-4113-2014, 2014.](#)

1607 [Wu, J., Ding, Y., Ye, B., Yang, Q., Hou, D. and Xue, L.: Stable isotopes in precipitation in](#)
1608 [Xilin River Basin, northern China and their implications, Chin. Geogr. Sci., 22\(5\), 531–540,](#)
1609 [doi:10.1007/s11769-012-0543-z, 2012.](#)

1610 [Xia, Y.: Optimization and uncertainty estimates of WMO regression models for the](#)
1611 [systematic bias adjustment of NLDAS precipitation in the United States, J. Geophys. Res.](#)
1612 [Atmospheres, 111\(D8\), D08102, doi:10.1029/2005JD006188, 2006.](#)

1613 [Yurtsever, Y.: Worldwide survey of stable isotopes in precipitation, International Atomic](#)
1614 [Energy Agency, Vienna, Austria., 1975.](#)

1615 [Zhang, Y., Shen, Y., Chen, Y. and Wang, Y.: Spatial characteristics of surface water and](#)
1616 [groundwater using water stable isotope in the Tarim River Basin, northwestern China,](#)
1617 [Ecohydrol., 6\(6\), 1031–1039, doi:10.1002/eco.1416, 2013.](#)

1618 [Zhao, L., Xiao, H., Zhou, J., Wang, L., Cheng, G., Zhou, M., Yin, L. and McCabe, M. F.:](#)
1619 [Detailed assessment of isotope ratio infrared spectroscopy and isotope ratio mass](#)
1620 [spectrometry for the stable isotope analysis of plant and soil waters, Rapid Commun. Mass](#)
1621 [Spectrom., 25\(20\), 3071–3082, doi:10.1002/rem.5204, 2011.](#)

1622 [Zimmermann, U., Ehhalt, D. and Muennich, K. O.: Soil Water Movement and](#)
1623 [Evapotranspiration: Changes in the Isotopic Composition of the Water., in Isotopes in](#)
1624 [Hydrology., pp. 567–85, International Atomic Energy Agency, Vienna, Austria., 1968.](#)

1625

- 1626 [Alaoui, A., Caduff, U., Gerke, H. H. and Weingartner, R.: Preferential Flow Effects on](#)
1627 [Infiltration and Runoff in Grassland and Forest Soils, *Vadose Zone J.*, 10\(1\), 367,](#)
1628 [doi:10.2136/vzj2010.0076, 2011.](#)
- 1629 [Allan, J. D.: Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems,](#)
1630 [Annu. Rev. Ecol. Evol. Syst., 35, 257–284, doi:10.1146/annurev.ecolsys.35.120202.110122,](#)
1631 [2004.](#)
- 1632 [Barnes, C. J. and Allison, G. B.: Tracing of water movement in the unsaturated zone using](#)
1633 [stable isotopes of hydrogen and oxygen, *J. Hydrol.*, 100\(1–3\), 143–176, doi:10.1016/0022-](#)
1634 [1694\(88\)90184-9, 1988.](#)
- 1635 [Barthold, F. K., Tyralla, C., Schneider, K., Vaché, K. B., Frede, H.-G. and Breuer, L.: How](#)
1636 [many tracers do we need for end member mixing analysis \(EMMA\)? A sensitivity analysis,](#)
1637 [Water Resour. Res., 47\(8\), doi:10.1029/2011WR010604, 2011.](#)
- 1638 [Blasch, K. W. and Bryson, J. R.: Distinguishing Sources of Ground Water Recharge by Using](#)
1639 [\$\delta^2\text{H}\$ and \$\delta^{18}\text{O}\$, *Ground Water*, 45\(3\), 294–308, doi:10.1111/j.1745-6584.2006.00289.x, 2007.](#)
- 1640 [Botter, G., Bertuzzo, E. and Rinaldo, A.: Transport in the hydrologic response: Travel time](#)
1641 [distributions, soil moisture dynamics, and the old water paradox, *Water Resour. Res.*, 46\(3\),](#)
1642 [W03514, doi:10.1029/2009WR008371, 2010.](#)
- 1643 [Botter, G., Bertuzzo, E. and Rinaldo, A.: Catchment residence and travel time distributions:](#)
1644 [The master equation, *Geophys. Res. Lett.*, 38\(11\), L11403, doi:10.1029/2011GL047666,](#)
1645 [2011.](#)
- 1646 [Brodersen, C., Pohl, S., Lindenlaub, M., Leibundgut, C. and Wilpert, K. v: Influence of](#)
1647 [vegetation structure on isotope content of throughfall and soil water, *Hydrol. Process.*, 14\(8\),](#)
1648 [1439–1448, doi:10.1002/1099-1085\(20000615\)14:8<1439::AID-HYP985>3.0.CO;2-3, 2000.](#)
- 1649 [Burger, H. M. and Seiler, K. P.: Evaporation from soil water under humid climate conditions](#)
1650 [and its impact on deuterium and \$^{18}\text{O}\$ concentrations in groundwater, edited by International](#)
1651 [Atomic Energy Agency, International Atomic Energy Agency, Vienna, Austria., 1992.](#)
- 1652 [Buttle, J. M.: Isotope hydrograph separations and rapid delivery of pre-event water from](#)
1653 [drainage basins, *Prog. Phys. Geogr.*, 18\(1\), 16–41, doi:10.1177/030913339401800102, 1994.](#)
- 1654 [Buttle, J. M.: Isotope Hydrograph Separation of Runoff Sources, in *Encyclopedia of*](#)
1655 [Hydrological Sciences, edited by M. G. Anderson, p. 10:116, John Wiley & Sons, Ltd,](#)
1656 [Chichester, Great Britain., 2006.](#)
- 1657 [Buttle, J. M. and McDonald, D. J.: Coupled vertical and lateral preferential flow on a forested](#)
1658 [slope, *Water Resour. Res.*, 38\(5\), 18–1, doi:10.1029/2001WR000773, 2002.](#)
- 1659 [Clark, I. D. and Fritz, P.: Groundwater, in *Environmental Isotopes in Hydrogeology*, p. 80,](#)
1660 [CRC Press, Florida, FL, USA., 1997a.](#)
- 1661 [Clark, I. D. and Fritz, P.: Methods for Field Sampling, in *Environmental Isotopes in*](#)
1662 [Hydrogeology, p. 283, CRC Press, Florida, FL, USA., 1997b.](#)

- 1663 [Clark, I. D. and Fritz, P.: The Environmental Isotopes, in Environmental Isotopes in](#)
1664 [Hydrogeology, pp. 2–34, CRC Press, Florida, FL, USA., 1997c.](#)
- 1665 [Cooper, L. W.: Isotopic Fractionation in Snow Cover, in Isotope Tracers in Catchment](#)
1666 [Hydrology, edited by C. Kendall and J. J. McDonnell, pp. 119–136, Elsevier, Amsterdam,](#)
1667 [Netherlands., 1998.](#)
- 1668 [Craig, H.: Isotopic Variations in Meteoric Waters, Science, 133\(3465\), 1702–1703,](#)
1669 [doi:10.1126/science.133.3465.1702, 1961a.](#)
- 1670 [Craig, H.: Standard for reporting concentrations of deuterium and oxygen-18 in natural](#)
1671 [waters, Science, 133\(3467\), 1833, 1961b.](#)
- 1672 [Csardi, G. and Nepusz, T.: The igraph software package for complex network research,](#)
1673 [Complex Systems 1695. \[online\] Available from: <http://igraph.sf.net> \(Accessed 24 December](#)
1674 [2015\), 2006.](#)
- 1675 [Dansgaard, W.: Stable isotopes in precipitation, Tellus, 16\(4\), 436–468, doi:10.1111/j.2153-](#)
1676 [3490.1964.tb00181.x, 1964.](#)
- 1677 [Darling, W. G.: Hydrological factors in the interpretation of stable isotopic proxy data present](#)
1678 [and past: a European perspective, Quat. Sci. Rev., 23\(7–8\), 743–770,](#)
1679 [doi:10.1016/j.quascirev.2003.06.016, 2004.](#)
- 1680 [Duvert, C., Stewart, M. K., Cendón, D. I. and Raiber, M.: Time series of tritium, stable](#)
1681 [isotopes and chloride reveal short-term variations in groundwater contribution to a stream,](#)
1682 [Hydrol. Earth Syst. Sci., 20\(1\), 257–277, doi:10.5194/hess-20-257-2016, 2016.](#)
- 1683 [DWD: Deutscher Wetterdienst - Wetter und Klima, Bundesministerium für Verkehr und](#)
1684 [digitale Infrastruktur, \[online\] Available from: <http://dwd.de/> \(Accessed 17 February 2014\),](#)
1685 [2014.](#)
- 1686 [Foerstel, H., Frinken, J., Huetzen, H., Lembrich, D. and Puetz, T.: Application of H₂¹⁸O as a](#)
1687 [tracer of water flow in soil, International Atomic Energy Agency, Vienna, Austria., 1991.](#)
- 1688 [Garvelmann, J., Kuells, C. and Weiler, M.: A porewater-based stable isotope approach for the](#)
1689 [investigation of subsurface hydrological processes, Hydrol. Earth Syst. Sci., 16\(2\), 631–640,](#)
1690 [doi:10.5194/hess-16-631-2012, 2012.](#)
- 1691 [Gat, J. R.: Oxygen and Hydrogen Isotopes in the Hydrologic Cycle, Annu. Rev. Earth Planet.](#)
1692 [Sci., 24\(1\), 225–262, doi:10.1146/annurev.earth.24.1.225, 1996.](#)
- 1693 [Gat, J., R., Mook, W. G. and Meijer, H. A. J.: Environmental isotopes in the hydrological](#)
1694 [cycle: Principles and Applications, edited by W. G. Mook, International Hydrological](#)
1695 [Programme; United Nations Educational, Scientific and Cultural Organization and](#)
1696 [International Atomic Energy Agency, Paris, France., 2001.](#)
- 1697 [Gehrels, J. C., Peeters, J. E. M., de Vries, J. J. and Dekkers, M.: The mechanism of soil water](#)
1698 [movement as inferred from ¹⁸O stable isotope studies, Hydrol. Sci. J., 43\(4\), 579–594,](#)
1699 [doi:10.1080/02626669809492154, 1998.](#)

- 1700 [Genereux, D. P. and Hooper, R. P.: Oxygen and Hydrogen Isotopes in Rainfall-Runoff](#)
1701 [Studies, in Isotope Tracers in Catchment Hydrology, edited by C. Kendall and J. J.](#)
1702 [McDonnell, pp. 319–346, Elsevier, Amsterdam, Netherlands., 1998.](#)
- 1703 [Gomez, J. D. and Wilson, J. L.: Age distributions and dynamically changing hydrologic](#)
1704 [systems: Exploring topography-driven flow, Water Resour. Res., 49\(3\), 1503–1522,](#)
1705 [doi:10.1002/wrcr.20127, 2013.](#)
- 1706 [Gonfiantini, R., Fröhlich, K., Araguás-Araguás, L. and Rozanski, K.: Isotopes in](#)
1707 [Groundwater Hydrology, in Isotope Tracers in Catchment Hydrology, edited by C. Kendall](#)
1708 [and J. J. McDonnell, pp. 203–246, Elsevier, Amsterdam, Netherlands., 1998.](#)
- 1709 [Gordon, L. J., Finlayson, C. M. and Falkenmark, M.: Managing water in agriculture for food](#)
1710 [production and other ecosystem services, Agric. Water Manag., 97\(4\), 512–519,](#)
1711 [doi:10.1016/j.agwat.2009.03.017, 2010.](#)
- 1712 [Harman, C. J.: Time-variable transit time distributions and transport: Theory and application](#)
1713 [to storage-dependent transport of chloride in a watershed, Water Resour. Res., 51\(1\), 1–30,](#)
1714 [doi:10.1002/2014WR015707, 2015.](#)
- 1715 [Heidbüchel, I., Troch, P. A., Lyon, S. W. and Weiler, M.: The master transit time distribution](#)
1716 [of variable flow systems, Water Resour. Res., 48\(6\), W06520, doi:10.1029/2011WR011293,](#)
1717 [2012.](#)
- 1718 [Hindmarsh, A. C., Brown, P. N., Grant, K. E., Lee, S. L., Serban, R., Shumaker, D. E. and](#)
1719 [Woodward, C. S.: SUNDIALS: Suite of Nonlinear and Differential/Algebraic Equation](#)
1720 [Solvers, ACM Trans. Math. Softw., 31\(3\), 363–396, doi:10.1145/1089014.1089020, 2005.](#)
- 1721 [Hollander, M., Wolfe, D. A. and Chicken, E.: Nonparametric Statistical Methods, John Wiley](#)
1722 [& Sons, New York, NY, USA., 2013.](#)
- 1723 [Hrachowitz, M., Benettin, P., van Breukelen, B. M., Fovet, O., Howden, N. J. K., Ruiz, L.,](#)
1724 [van der Velde, Y. and Wade, A. J.: Transit times—the link between hydrology and water](#)
1725 [quality at the catchment scale, Wiley Interdiscip. Rev. Water, doi:10.1002/wat2.1155, 2016.](#)
- 1726 [IAEA: International Atomic Energy Agency: Water Resources Programme – Global Network](#)
1727 [of Isotopes in Precipitation, \[online\] Available from: \[http://www-\]\(http://www-naweb.iaea.org/napc/ih/IHS_resources_gnip.html\)](#)
1728 [naweb.iaea.org/napc/ih/IHS_resources_gnip.html \(Accessed 11 August 2014\), 2014.](#)
- 1729 [Jin, L., Siegel, D. I., Lautz, L. K., Mitchell, M. J., Dahms, D. E. and Mayer, B.: Calcite](#)
1730 [precipitation driven by the common ion effect during groundwater–surface-water mixing: A](#)
1731 [potentially common process in streams with geologic settings containing gypsum, Geol. Soc.](#)
1732 [Am. Bull., 122\(7/8\), B30011.1, doi:10.1130/B30011.1, 2010.](#)
- 1733 [Jin, L., Siegel, D. I., Lautz, L. K. and Lu, Z.: Identifying streamflow sources during spring](#)
1734 [snowmelt using water chemistry and isotopic composition in semi-arid mountain streams, J.](#)
1735 [Hydrol., 470–471, 289–301, doi:10.1016/j.jhydrol.2012.09.009, 2012.](#)
- 1736 [Jouzel, J., Alley, R. B., Cuffey, K. M., Dansgaard, W., Grootes, P., Hoffmann, G., Johnsen, S.](#)
1737 [J., Koster, R. D., Peel, D., Shuman, C. A., Stievenard, M., Stuiver, M. and White, J.: Validity](#)

- 1738 [of the temperature reconstruction from water isotopes in ice cores, J. Geophys. Res., 102,](#)
1739 [26471–26487, doi:10.1029/97JC01283, 1997.](#)
- 1740 [Kendall, C. and Caldwell, E. A.: Fundamentals of Isotope Geochemistry, in Isotope Tracers in](#)
1741 [Catchment Hydrology, edited by C. Kendall and J. J. McDonnell, pp. 51–86, Elsevier,](#)
1742 [Amsterdam, Netherlands., 1998.](#)
- 1743 [Kirchner, J. W.: A double paradox in catchment hydrology and geochemistry, Hydrol.](#)
1744 [Process., 17\(4\), 871–874, doi:10.1002/hyp.5108, 2003.](#)
- 1745 [Klaus, J., McDonnell, J. J., Jackson, C. R., Du, E. and Griffiths, N. A.: Where does](#)
1746 [streamwater come from in low-relief forested watersheds? A dual-isotope approach, Hydrol.](#)
1747 [Earth Syst. Sci., 19\(1\), 125–135, 2015.](#)
- 1748 [Koeniger, P., Leibundgut, C. and Stichler, W.: Spatial and temporal characterisation of stable](#)
1749 [isotopes in river water as indicators of groundwater contribution and confirmation of](#)
1750 [modelling results; a study of the Weser river, Germany, Isotopes Environ. Health Stud., 45\(4\),](#)
1751 [289–302, doi:10.1080/10256010903356953, 2009.](#)
- 1752 [Kolaczyk, E. D.: Statistical Analysis of Network Data with R, Springer Science & Business](#)
1753 [Media, New York, NY, USA., 2014.](#)
- 1754 [Kortelainen, N. M. and Karhu, J. A.: Regional and seasonal trends in the oxygen and](#)
1755 [hydrogen isotope ratios of Finnish groundwaters: a key for mean annual precipitation, J.](#)
1756 [Hydrol., 285\(1–4\), 143–157, doi:10.1016/j.jhydrol.2003.08.014, 2004.](#)
- 1757 [Kraft, P., Vaché, K. B., Frede, H.-G. and Breuer, L.: CMF: A Hydrological Programming](#)
1758 [Language Extension For Integrated Catchment Models, Environ. Model. Softw., 26\(6\), 828–](#)
1759 [830, doi:10.1016/j.envsoft.2010.12.009, 2011.](#)
- 1760 [Lai, C.-T. and Ehleringer, J. R.: Deuterium excess reveals diurnal sources of water vapor in](#)
1761 [forest air, Oecologia, 165\(1\), 213–223, doi:10.1007/s00442-010-1721-2, 2011.](#)
- 1762 [Lauer, F., Frede, H.-G. and Breuer, L.: Uncertainty assessment of quantifying spatially](#)
1763 [concentrated groundwater discharge to small streams by distributed temperature sensing,](#)
1764 [Water Resour. Res., 49\(1\), 400–407, doi:10.1029/2012WR012537, 2013.](#)
- 1765 [LGR: Los Gatos Research, Greenhouse Gas, isotope and trace gas analyzers, \[online\]](#)
1766 [Available from: <http://www.lgrinc.com/> \(Accessed 5 February 2013\), 2013.](#)
- 1767 [Li, F., Song, X., Tang, C., Liu, C., Yu, J. and Zhang, W.: Tracing infiltration and recharge](#)
1768 [using stable isotope in Taihang Mt., North China, Environ. Geol., 53\(3\), 687–696,](#)
1769 [doi:10.1007/s00254-007-0683-0, 2007.](#)
- 1770 [Maloszewski, P. and Zuber, A.: Manual on lumped parameter models used for the](#)
1771 [interpretation of environmental tracer data in groundwaters, in Use of isotopes for analyses of](#)
1772 [flow and transport dynamics in groundwater systems, p. 50, International Atomic Energy](#)
1773 [Agency, Vienna, Austria., 2002.](#)

- 1774 [McConville, C., Kalin, R. m., Johnston, H. and McNeill, G. w.: Evaluation of Recharge in a](#)
1775 [Small Temperate Catchment Using Natural and Applied \$\delta^{18}\text{O}\$ Profiles in the Unsaturated](#)
1776 [Zone, *Ground Water*, 39\(4\), 616–623, doi:10.1111/j.1745-6584.2001.tb02349.x, 2001.](#)
- 1777 [McDonnell, J. J. and Beven, K.: Debates—The future of hydrological sciences: A \(common\)](#)
1778 [path forward? A call to action aimed at understanding velocities, celerities and residence time](#)
1779 [distributions of the headwater hydrograph, *Water Resour. Res.*, 50\(6\), 5342–5350,](#)
1780 [doi:10.1002/2013WR015141, 2014.](#)
- 1781 [McDonnell, J. J., Sivapalan, M., Vaché, K., Dunn, S., Grant, G., Haggerty, R., Hinz, C.,](#)
1782 [Hooper, R., Kirchner, J., Roderick, M. L. and others: Moving beyond heterogeneity and](#)
1783 [process complexity: a new vision for watershed hydrology, *Water Resour. Res.*, 43\(7\),](#)
1784 [W07301, 2007.](#)
- 1785 [McDonnell, J. J., McGuire, K., Aggarwal, P., Beven, K. J., Biondi, D., Destouni, G., Dunn,](#)
1786 [S., James, A., Kirchner, J., Kraft, P., Lyon, S., Maloszewski, P., Newman, B., Pfister, L.,](#)
1787 [Rinaldo, A., Rodhe, A., Sayama, T., Seibert, J., Solomon, K., Soulsby, C., Stewart, M.,](#)
1788 [Tetzlaff, D., Tobin, C., Troch, P., Weiler, M., Western, A., Worman, A. and Wrede, S.: How](#)
1789 [old is streamwater? Open questions in catchment transit time conceptualization, modelling](#)
1790 [and analysis, *Hydrol. Process.*, 24\(12\), 1745–1754, 2010.](#)
- 1791 [McGuire, K. J. and McDonnell, J. J.: A review and evaluation of catchment transit time](#)
1792 [modeling, *J. Hydrol.*, 330\(3\), 543–563, 2006.](#)
- 1793 [Michel, R. L.: Residence times in river basins as determined by analysis of long-term tritium](#)
1794 [records, *J. Hydrol.*, 130\(1\), 367–378, doi:10.1016/0022-1694\(92\)90117-E, 1992.](#)
- 1795 [Mook, W. G., Groeneveld, D. J., Brouwn, A. E. and Ganswijk, A. J. van: Analysis of a run-](#)
1796 [off hydrograph by means of natural \$^{18}\text{O}\$, in *Isotope techniques in groundwater hydrology*, vol.](#)
1797 [1, pp. 159–169, International Atomic Energy Agency, Vienna, Austria., 1974.](#)
- 1798 [Neal, C. and Rosier, P. T. W.: Chemical studies of chloride and stable oxygen isotopes in two](#)
1799 [conifer afforested and moorland sites in the British uplands, *J. Hydrol.*, 115\(1–4\), 269–283,](#)
1800 [doi:10.1016/0022-1694\(90\)90209-G, 1990.](#)
- 1801 [O’Driscoll, M. A., DeWalle, D. R., McGuire, K. J. and Gburek, W. J.: Seasonal \$^{18}\text{O}\$ variations](#)
1802 [and groundwater recharge for three landscape types in central Pennsylvania, USA, *J. Hydrol.*,](#)
1803 [303\(1–4\), 108–124, doi:10.1016/j.jhydrol.2004.08.020, 2005.](#)
- 1804 [Orlowski, N., Frede, H.-G., Brüggemann, N. and Breuer, L.: Validation and application of a](#)
1805 [cryogenic vacuum extraction system for soil and plant water extraction for isotope analysis, *J.*](#)
1806 [Sens. Sens. Syst., 2\(2\), 179–193, doi:10.5194/jsss-2-179-2013, 2013.](#)
- 1807 [Orlowski, N., Lauer, F., Kraft, P., Frede, H.-G. and Breuer, L.: Linking Spatial Patterns of](#)
1808 [Groundwater Table Dynamics and Streamflow Generation Processes in a Small Developed](#)
1809 [Catchment, *Water*, 6\(10\), 3085–3117, doi:10.3390/w6103085, 2014.](#)
- 1810 [Penna, D., Ahmad, M., Birks, S. J., Bouchaou, L., Brenčič, M., Butt, S., Holko, L., Jeelani,](#)
1811 [G., Martínez, D. E., Melikadze, G., Shanley, J. B., Sokratov, S. A., Stadnyk, T., Sugimoto, A.](#)
1812 [and Vreča, P.: A new method of snowmelt sampling for water stable isotopes, *Hydrol.*](#)
1813 [Process., 28\(22\), 5637–5644, doi:10.1002/hyp.10273, 2014.](#)

- 1814 [Perry, C. and Taylor, K.: Environmental Sedimentology, p. 36, Blackwell Publishing, Oxford,](#)
1815 [OX, UK, 2009.](#)
- 1816 [Pierce, S. C., Kröger, R. and Pezeshki, R.: Managing Artificially Drained Low-Gradient](#)
1817 [Agricultural Headwaters for Enhanced Ecosystem Functions, Biology, 1\(3\), 794–856,](#)
1818 [doi:10.3390/biology1030794, 2012.](#)
- 1819 [Qu, Y. and Duffy, C. J.: A semidiscrete finite volume formulation for multiprocess watershed](#)
1820 [simulation, Water Resour. Res., 43\(8\), W08419, doi:10.1029/2006WR005752, 2007.](#)
- 1821 [Rinaldo, A., Benettin, P., Harman, C. J., Hrachowitz, M., McGuire, K. J., van der Velde, Y.,](#)
1822 [Bertuzzo, E. and Botter, G.: Storage selection functions: A coherent framework for](#)
1823 [quantifying how catchments store and release water and solutes, Water Resour. Res., 51\(6\),](#)
1824 [4840–4847, doi:10.1002/2015WR017273, 2015.](#)
- 1825 [Rohde, A.: Snowmelt-Dominated Systems, in Isotope Tracers in Catchment Hydrology,](#)
1826 [edited by C. Kendall and J. J. McDonnell, pp. 391–433, Elsevier, Amsterdam, Netherlands.,](#)
1827 [1998.](#)
- 1828 [Rozanski, K., Sonntag, C. and Münnich, K. O.: Factors controlling stable isotope composition](#)
1829 [of European precipitation, Tellus, 34\(2\), 142–150, doi:10.1111/j.2153-3490.1982.tb01801.x,](#)
1830 [1982.](#)
- 1831 [Rozanski, K., Froehlich, K. and Mook, W. G.: in Environmental isotopes in the hydrological](#)
1832 [cycle: Principles and Applications, vol. 3, International Hydrological Programme; United](#)
1833 [Nations Educational, Scientific and Cultural Organization and International Atomic Energy](#)
1834 [Agency, Paris, Vienna., 2001.](#)
- 1835 [Schultz, N. M., Griffis, T. J., Lee, X. and Baker, J. M.: Identification and correction of](#)
1836 [spectral contamination in \$^2\text{H}/^1\text{H}\$ and \$^{18}\text{O}/^{16}\text{O}\$ measured in leaf, stem, and soil water, Rapid](#)
1837 [Commun. Mass Spectrom., 25\(21\), 3360–3368, doi:10.1002/rcm.5236, 2011.](#)
- 1838 [Shuttleworth, W. J. and Wallace, J. S.: Evaporation from sparse crops-an energy combination](#)
1839 [theory, Q. J. R. Meteorol. Soc., 111\(469\), 839–855, doi:10.1002/qj.49711146910, 1985.](#)
- 1840 [Sklash, M. G.: Environmental isotope studies of storm and snowmelt runoff generation, in](#)
1841 [Process studies in hillslope hydrology, edited by M. G. Anderson, pp. 410–435, Wiley, New](#)
1842 [York, NY, USA., 1990.](#)
- 1843 [Sklash, M. G. and Farvolden, R. N.: The Role Of Groundwater In Storm Runoff, in](#)
1844 [Developments in Water Science, vol. 12, edited by W. B. and D. A. Stephenson, pp. 45–65,](#)
1845 [Elsevier, Amsterdam, Netherlands., 1979.](#)
- 1846 [Song, X., Wang, P., Yu, J., Liu, X., Liu, J. and Yuan, R.: Relationships between precipitation,](#)
1847 [soil water and groundwater at Chongling catchment with the typical vegetation cover in the](#)
1848 [Taihang mountainous region, China, Environ. Earth Sci., 62\(4\), 787–796,](#)
1849 [doi:10.1007/s12665-010-0566-7, 2011.](#)
- 1850 [Stewart, M. K., Morgenstern, U. and McDonnell, J. J.: Truncation of stream residence time:](#)
1851 [how the use of stable isotopes has skewed our concept of streamwater age and origin, Hydrol.](#)
1852 [Process., 24\(12\), 1646–1659, doi:10.1002/hyp.7576, 2010.](#)

- 1853 [Stumpp, C., Klaus, J. and Stichler, W.: Analysis of long-term stable isotopic composition in](#)
1854 [German precipitation, *J. Hydrol.*, 517, 351–361, doi:10.1016/j.jhydrol.2014.05.034, 2014.](#)
- 1855 [Tang, K. and Feng, X.: The effect of soil hydrology on the oxygen and hydrogen isotopic](#)
1856 [compositions of plants' source water, *Earth Planet. Sci. Lett.*, 185\(3–4\), 355–367, 2001.](#)
- 1857 [Taylor, S., Feng, X., Kirchner, J. W., Osterhuber, R., Klaue, B. and Renshaw, C. E.: Isotopic](#)
1858 [evolution of a seasonal snowpack and its melt, *Water Resour. Res.*, 37\(3\), 759–769,](#)
1859 [doi:10.1029/2000WR900341, 2001.](#)
- 1860 [Thomas, E. M., Lin, H., Duffy, C. J., Sullivan, P. L., Holmes, G. H., Brantley, S. L. and Jin,](#)
1861 [L.: Spatiotemporal Patterns of Water Stable Isotope Compositions at the Shale Hills Critical](#)
1862 [Zone Observatory: Linkages to Subsurface Hydrologic Processes, *Vadose Zone J.*, 12\(4\), 0,](#)
1863 [doi:10.2136/vzj2013.01.0029, 2013.](#)
- 1864 [Timbe, E., Windhorst, D., Crespo, P., Frede, H.-G., Feyen, J. and Breuer, L.: Understanding](#)
1865 [uncertainties when inferring mean transit times of water trough tracer-based lumped-](#)
1866 [parameter models in Andean tropical montane cloud forest catchments, *Hydrol. Earth Syst.*](#)
1867 [Sci., 18\(4\), 1503–1523, doi:10.5194/hess-18-1503-2014, 2014.](#)
- 1868 [van der Velde, Y., Torfs, P. J. J. F., van der Zee, S. E. a. T. M. and Uijlenhoet, R.:](#)
1869 [Quantifying catchment-scale mixing and its effect on time-varying travel time distributions,](#)
1870 [Water Resour. Res., 48\(6\), W06536, doi:10.1029/2011WR011310, 2012.](#)
- 1871 [Wang, X. F. and Yakir, D.: Using stable isotopes of water in evapotranspiration studies,](#)
1872 [Hydrol. Process., 14\(8\), 1407–1421, doi:10.1002/1099-1085\(20000615\)14:8<1407::AID-](#)
1873 [HYP992>3.0.CO;2-K, 2000.](#)
- 1874 [Windhorst, D., Waltz, T., Timbe, E., Frede, H.-G. and Breuer, L.: Impact of elevation and](#)
1875 [weather patterns on the isotopic composition of precipitation in a tropical montane rainforest,](#)
1876 [Hydrol. Earth Syst. Sci., 17\(1\), 409–419, doi:10.5194/hess-17-409-2013, 2013.](#)
- 1877 [Windhorst, D., Kraft, P., Timbe, E., Frede, H.-G. and Breuer, L.: Stable water isotope tracing](#)
1878 [through hydrological models for disentangling runoff generation processes at the hillslope](#)
1879 [scale, *Hydrol. Earth Syst. Sci.*, 18\(10\), 4113–4127, doi:10.5194/hess-18-4113-2014, 2014.](#)
- 1880 [Woolfenden, L. R. and Ginn, T. R.: Modeled Ground Water Age Distributions, *Ground*](#)
1881 [Water, 47\(4\), 547–557, doi:10.1111/j.1745-6584.2008.00550.x, 2009.](#)
- 1882 [Wu, J., Ding, Y., Ye, B., Yang, Q., Hou, D. and Xue, L.: Stable isotopes in precipitation in](#)
1883 [Xilin River Basin, northern China and their implications, *Chin. Geogr. Sci.*, 22\(5\), 531–540,](#)
1884 [doi:10.1007/s11769-012-0543-z, 2012.](#)
- 1885 [Xia, Y.: Optimization and uncertainty estimates of WMO regression models for the](#)
1886 [systematic bias adjustment of NLDAS precipitation in the United States, *J. Geophys. Res.*](#)
1887 [Atmospheres, 111\(D8\), D08102, doi:10.1029/2005JD006188, 2006.](#)
- 1888 [Yurtsever, Y.: Worldwide survey of stable isotopes in precipitation, International Atomic](#)
1889 [Energy Agency, Vienna, Austria., 1975.](#)

1890 [Zhao, L., Xiao, H., Zhou, J., Wang, L., Cheng, G., Zhou, M., Yin, L. and McCabe, M. F.:](#)
1891 [Detailed assessment of isotope ratio infrared spectroscopy and isotope ratio mass](#)
1892 [spectrometry for the stable isotope analysis of plant and soil waters, Rapid Commun. Mass](#)
1893 [Spectrom., 25\(20\), 3071–3082, doi:10.1002/rcm.5204, 2011.](#)


1894

1895 Table 1. Descriptive statistics of $\delta^2\text{H}$, $\delta^{18}\text{O}$, and d-excess values for precipitation, stream, and
 1896 groundwater over the two-year observation period including all sampling points.

Sample type	Mean \pm SD		Min		Max		D-excess mean \pm SD	N
	$\delta^2\text{H}$	$\delta^{18}\text{O}$	$\delta^2\text{H}$	$\delta^{18}\text{O}$	$\delta^2\text{H}$	$\delta^{18}\text{O}$		
	[‰]	[‰]	[‰]	[‰]	[‰]	[‰]		
Precipitation	-43.9 \pm 23.4	-6.2 \pm 3.1	-167.6	-22.4	-8.3	-1.2	5.9 \pm 5.7	592
Vollnkirchener Bach	-58.0 \pm 2.8	-8.4 \pm 0.4	-66.3	-10.0	-26.9	-6.7	9.0 \pm 2.3	332
Schwingbach	-58.2 \pm 4.3	-8.4 \pm 0.6	-139.7	-18.3	-47.2	-5.9	9.0 \pm 2.2	463
Groundwater meadow	-57.6 \pm 1.6	-8.2 \pm 0.4	-64.9	-9.2	-50.8	-5.7	7.9 \pm 5.5	375
Groundwater arable land	-56.2 \pm 3.7	-8.0 \pm 0.5	-91.6	-12.3	-49.5	-6.8	1.7 \pm 5.0	338
Groundwater along stream	-59.9 \pm 6.8	-8.5 \pm 0.9	-94.5	-13.0	-49.5	-7.0	8.2 \pm 1.5	108

Table 2. Mean and standard deviation for isotopic signatures and soil physical properties in 0.2 m and 0.5 m soil depth (N=52 per depth).

	$\delta^2\text{H}$ [‰]		$\delta^{18}\text{O}$ [‰]		water content [% w/w]		pH		bulk density [g cm ⁻³]	
	0.2 m	0.5 m	0.2 m	0.5 m	0.2 m	0.5 m	0.2 m	0.5 m	0.2 m	0.5 m
Mean±SD	-46.9±8.4	-58.5±8.3	-6.6±1.2	-8.2±1.2	16.8±7.2	16.1±8.3	5.0±1.0	5.3±1.0	1.3±0.2	1.3±0.2




1898

1899


1900 Figure 1. Maps show (a) the location of the Schwingbach catchment in Germany, (b) the main
 1901 monitoring area, (c) the land use, elevation, and instrumentation, (d) the locations of the

1902 snapshot as well as the seasonal soil samplings, (e) soil types, and (f) geology of the
1903 Schwingbach catchment including the Vollnkirchener Bach subcatchment boundaries.
1904



1905
 1906
 1907
 1908
 1909
 1910
 1911


Figure 2. Temporal variation of precipitation amount, isotopic signatures ($\delta^2\text{H}$ and $\delta^{18}\text{O}$) including snow samples (grey striped box), and d-excess values for the study area compared to monthly d-excess values (July 2011 to July 2013) of GNIP station Koblenz with reference d-excess of GMWL ($d=10$; solid black line).



1912

1913

1914 Figure 3. Local Meteoric Water Line for the Schwingbach catchment (LMWL) in comparison
 1915 to GMWL, including comparisons between precipitation, stream water, groundwater, and soil
 1916 water isotopic signatures and the respective EWLs.




1917

1918

1919 Figure 4. Boxplots of $\delta^2\text{H}$ values comparing precipitation, stream, groundwater, and soil
 1920 isotopic composition in 0.2 m and 0.5 m depth ($N=52$ per depth). Different letters indicate
 1921 significant differences ($p \leq 0.05$).

1922




1923

1924

1925 Figure 5. Dual isotope plot of soil water isotopic signatures in 0.2 m and 0.5 m depth
 1926 compared by land use including precipitation isotope data from 19, 21, and 28 October 2011.

1927 Insets: Boxplots comparing $\delta^2\text{H}$ isotopic signatures between different land use units and
 1928 precipitation (small letters) in top and subsoil (capital letters). Different letters indicate

1929 significant differences ($p \leq 0.05$).



1930

1931

1932 Figure 6. Seasonal $\delta^2\text{H}$ profiles of soil water (upper panels) and water content (lower panels)
 1933 for winter (28 March 2013), summer (28 August 2011), and spring (24 April 2013). Error bars
 1934 represent the natural isotopic variation of the replicates taken during each sampling campaign.
 1935 For reference, mean groundwater (grey shaded) and mean seasonal precipitation $\delta^2\text{H}$ values
 1936 are shown (coloured arrows at the top).




Figure 7. Mean daily discharge at the Vollnkirchener Bach (13, 18) and Schwingbach (site 11, 19, and 64) with automatically recorded data (solid lines) and manual discharge measurements (asterisks), temporal variation of $\delta^2\text{H}$ of stream water in the Schwingbach (site 11, 19, and 64) and Vollnkirchener Bach (site 13, 18, and 94) including moving averages (MA) for streamflow isotopes.




Figure 8. Temporal variation of discharge at the Vollnkirchener Bach with automatically recorded data (solid line) and manual discharge measurements (asterisks) (site 18), groundwater

head levels, and $\delta^2\text{H}$ values (coloured dots) for selected piezometers under meadow (site 3 and 21), arable land (site 26, 27, and 28), and beside the Vollnkirchener Bach (site 24 and 32) including moving averages for groundwater isotopes.




Figure 9. Network map of $\delta^{18}\text{O}$ relationships between surface water (SW) and groundwater (GW) sampling points. Yellow circles represent groundwater sampling points on the arable field, light green circles are piezometers located on the grassland close to the conjunction of the Schwingbach with the Vollnkirchener Bach, and dark green circles represent piezometers along the Vollnkirchener Bach. Light blue circles stand for Schwingbach and darker blue circles for Vollnkirchener Bach surface water sampling points. See Figure 1 for an overview of all sampling points. Only statistically significant connections between $\delta^{18}\text{O}$ time series ($p < 0.05$) are shown in the network diagram.




Figure 10. Maps of modelled groundwater ages (colour scheme) and flow directions (white arrows) of (a) the Vollnkirchner Bach subcatchment and (b) detail view of the northern part of the subcatchment. The intensity of flow is depicted by the length of the white arrows.

Appendix I

Mean transit time estimation

We applied a set of five different models to estimate the MTT using the FlowPC software (Maloszewski and Zuber, 2002): dispersion model (with different dispersion parameters $D_p=0.05, 0.4, \text{ and } 0.8$), exponential model, exponential-piston-flow model, linear model, and linear-piston-flow model. We evaluated these results using two goodness of fit criteria, i.e. sigma (σ) and model efficiency (ME) following Maloszewski and Zuber (2002):

$$\sigma = \frac{\sqrt{\sum (c_{mi} - c_{oi})^2}}{m} \quad (1)$$

$$ME = 1 - \frac{\sum (c_{mi} - c_{oi})^2}{\sum (c_{oi} - \bar{c}_o)^2} \quad (2)$$

Where:

- c_{mi} : The i-th model result
- c_{oi} : The i-th observed result
- \bar{c}_o : The arithmetic mean of all observations

A model efficiency $ME=1$ indicates an ideal fit of the model to the concentrations observed, while $ME=0$ indicates that the model fits the data no better than a horizontal line through the mean observed concentration (Maloszewski and Zuber, 2002). The same is true for sigma. For calculations with FlowPC, weekly averages of precipitation and stream water isotopic signatures are calculated. We firstly calculated the MTT from precipitation to the streams for three sampling points in the Vollnkirchener Bach (sites 13, 18 and 94) and three points in the Schwingbach (sites 11, 19 and 64). For the second set of simulations, the mean residence time from precipitation to groundwater comprising thirteen groundwater sampling points was determined. We also bias-corrected the precipitation input data with two different approaches: the mean precipitation value is subtracted from every single precipitation value and then divided by the standard deviation of precipitation isotopic signatures. Afterwards, this value is subtracted from the weekly precipitation values (bias1). For the second approach, the difference of the mean stream water isotopic value and the mean precipitation value is calculated and also subtracted from the weekly precipitation values (bias2).

Appendix II

Model-based groundwater age dynamics

Objective:

Stable water isotopes are only a tool to determine the residence time for a few years (McDonnell et al., 2010). In cases of longer residence times and a strong mixing effect, seasonal variation of isotopes vanishes and results in stable flat lines of the isotopic signal. To get a rough estimate of residence times greater than the limit of stable water isotopes (>5 years), we split the water flow path in our catchment in two parts: the flow from precipitation to groundwater, which was calculated via FlowPC and the longer groundwater transport. The simplest method to estimate the residence time of groundwater transport is via the storage-to-input-relation, with the storage as the aquifer size and the input as the groundwater recharge time. However, this method ignores the topographic setting, and water input heterogeneity. In our study we used a simplified groundwater flow model with tracer transport to calculate the groundwater age dynamics. The numerical output of water ages cannot be validated with the given isotope data, since the model is used to fill a residence time gap, where stable water isotopes are not feasible to apply. The model is falsified however, if the residence time is short enough (<5 years) to be calculable via FlowPC. Hence, the results of the groundwater age model should be handled with care and only seen as the order of magnitude of flow time scales.

Model setup:

We set up a tailored hydrological model for the Vollnkirchener Bach subcatchment using the *Catchment Modelling Framework (CMF)* by Kraft et al. (2011). CMF is a modular framework for hydrological modelling based on the concept of finite volume method by Qu and Duffy (2007). CMF is applicable for simulating one- to three-dimensional water fluxes but also advective transport of stable water isotopes (^{18}O and ^2H). Thus, it is especially suitable for our tracer study and can be used to study the origin (Windhorst et al., 2014) and age of water. The generated model is a highly simplified representation of the Vollnkirchener Bach subcatchment's groundwater body. The subcatchment is divided into 353 polygonal-shaped cells ranging from 100–40'000 m² in size based on land use, soil type, and topography. The model is vertically

divided in two compartments, the upper soft rock aquifer, and the lower bed rock aquifer, referred to as upper and lower layer from now onwards.

The layers of each cell are connected using a mass conservative Darcy approach with a finite volume discretization. The water storage dynamic of one layer in one cell i of the groundwater model is given as:

$$\frac{dV_{i,s}}{dt} = R_i - S_i - \sum_{j=1}^{N_i} \left(K_s \frac{\Psi_{i,s} - \Psi_{j,s}}{d_{ij}} A_{ij,s} \right) \quad (3)$$

$$\frac{dV_{i,b}}{dt} = S_i - \sum_{j=1}^{N_i} \left(K_b \frac{\Psi_{i,b} - \Psi_{j,b}}{d_{ij}} A_{ij,b} \right)$$

Where:

- V_j : The water volume stored by the layer in m^3 in cell I for soft rock (s) and bedrock (b), respectively
- R_j : The groundwater recharge rate in $m^3 \cdot d^{-1}$
- S_j : the percolation from the soft rock to the bedrock aquifer, calculated by the gradient and geometric mean conductivity between the layers: $S_i = \sqrt{K_s K_b} \frac{\Psi_{i,s} - \Psi_{i,b}}{d_{sb}} A_i$, where d_{sb} is the distance between the layers and A_i is the cell area
- N_j : Number of adjacent cells to cell i
- K : Saturated hydraulic conductivity in $m \cdot d^{-1}$ for soft rock (s) and bedrock (b), respectively
- Ψ : Water head in the current cell i and the neighbour cell j in m for soft rock (s) and bedrock (b), respectively
- d_{ij} : The distance between the current cell i and the neighbour cell j in m
- $A_{i,j,x}$: The wetted area of the joint layer boundary in m^2 between cells i and j in layer x

The volume head relation is linearized as $\Psi = \phi \frac{V}{A}$, with ϕ being the fillable porosity and A the cell area. The resulting ordinary differential equation system is integrated using the CVODE

solver by Hindmarsh et al. (2005), an error controlled Krylov-Newton multistep implicit solver with an adaptive order of 1–5 according to stability constraints.

Boundary conditions:

The upper boundary condition of the groundwater system – the mean groundwater recharge – is modelled applying a Richard’s equation based model using measured rainfall data (2011–2013) and calculated evapotranspiration with the Shuttleworth-Wallace method (Shuttleworth and Wallace, 1985) including land cover and climate data. To retrieve long-term steady state conditions, the groundwater recharge is averaged and used as constant flow Neumann boundary condition. The total outflow is calibrated against measured outflow data; hence, the unsaturated model’s role is mainly to account for spatial heterogeneity of groundwater recharge. As an additional input, a combined sewer overflow (site 38, Fig. 1b) is considered based on findings of Orłowski et al. (2014). Moreover, there are two water outlets in the two lowest cells for efficient draining, reflecting measured groundwater flow directions throughout most of the year at piezometers 1–6 (Fig. 1b). Both cells are located in the very north of the subcatchment and their outlets are modelled as constant head Dirichlet boundary condition.

Parameters:

The saturated hydraulic conductivity of the groundwater body is set to 0.1007 m d⁻¹, as measured in the study area. For the lower bedrock compartment there is no data available. However, expecting a high rate of joints, preliminary testing revealed that a saturated hydraulic conductivity of 0.25 m d⁻¹ seemed to be a realistic estimation (based on field measurements).

Water Age:

To calculate the water age in each cell, a virtual tracer flows through the system using advective transport. To calculate the water age from the tracer that enters the system with a unity concentration by groundwater recharge, a linear decay is used to reduce the tracer concentration with time:

$$\frac{dX_{i,s}}{dt} = 1 \frac{u}{m^3} R_i - S_i[X]_{i,s} - \sum_{j=1}^{N_i} \left([X]_{i,s} K_s \frac{\Psi_{i,s} - \Psi_{j,s}}{d_{ij}} A_{ij,s} \right) - r X_{i,s} \quad (4)$$

$$\frac{dX_{i,b}}{dt} = S_i[X]_{i,s} - \sum_{j=1}^{N_i} \left([X]_{i,b} K_b \frac{\Psi_{i,b} - \Psi_{j,b}}{d_{ij}} A_{ij,b} \right) - rX_{i,b}$$

$$t_{ix} = \frac{\ln[X]_{ix}}{r}$$

Where:

- $X_{i,x}$: Amount of virtual tracer in layer x in cell i in virtual unit u
- $1 \frac{u}{m^3} R_i$: Tracer input with groundwater recharge R with unity concentration
- $[X]_{i,x}$: Concentration of tracer in layer x of cell i in $u m^{-3}$
- r : Arbitrary chosen decay constant, for water age calculation in d^{-1} . Rounding errors occur due to low concentrations when r is set to a high value. We found a good numerical performance with values between 10^{-6} – $10^{-9} d^{-1}$
- t_{ix} : Water age in days in layer x in cell i

To ensure long term steady state conditions, the model is run for 2000 years. However, after 300 years of model run time, steady state is reached.