

## 1 Reply to referee

2  
3 In the following please find the corrections and comments to the referee's response.

4  
5 I think the paper clearly improved from the last version, I think it can be published if the following aspects are  
6 commented (for my first concern) and changed (for #2 and #3).

7 I think there are some points in the modelling that could lead to several rounds of arguing, but I leave this here.

8 I would just appreciate if the authors could comment on the following concerns regarding the model setup.  
9

10 1) The model still does not convince me, as I see it as a pure model exercise. The authors stated that the model  
11 can simulate isotope transport, so why not showing the resulting stable isotope in the stream? I am aware that  
12 the length of the time series is with the current travel times not long enough, but you could repeat the same  
13 input time series and see if this somewhat fits the observation.

14 **The model is only used for the groundwater part and not for the vadose zone of the transport path. The**  
15 **isotopic dynamics in the groundwater are already fairly small and show similarities to the stream water close to**  
16 **the isotope analyzer's precision. This small signal-to-noise ratio led us to the decision of not including "real"**  
17 **isotope tracer data in the model. It is instead driven by realistic made-up-data (as the reviewer's suggestion of**  
18 **repeated time series). We are using a virtual tracer to estimate travel time (cf. eq. 4). Although, Windhorst et**  
19 **al. 2014 showed that direct modelling of isotope transport is feasible with the chosen software framework**  
20 **(CMF). To clarify our decision, we added the following sentence to L1069: "To avoid errors in transit time**  
21 **calculations from small differences between the isotopic signal in groundwater and stream water, we are**  
22 **tracing the transit time of groundwater and not the real isotopic values in this study."**

23 Currently, I am a little but concerned that a change in model parameters will have huge impacts on the  
24 calculated travel times.

25 **Yes, it has (see the disclaimer in L1058ff). We think that reporting the results of a complete**  
26 **sensitivity/uncertainty analysis of our model would be out of scope for an "Appendix II" model which was setup**  
27 **to get an estimate of transit time ranges. However, we have tested the model with different parameters, albeit**  
28 **in a rather exploring then rigid manner. It is possible to get much longer transit times with different**  
29 **parameters. The reported values are rather on the faster side.**

30 Furthermore, I was wondering if it is possible to also model the streamflow age, and not only the groundwater  
31 age?

32 **The mean streamflow age could be calculated for the groundwater contributing to each stream segment,**  
33 **weighted with the relative contribution to the stream. In this way, we would superimpose the uncertainties of**  
34 **the groundwater age with the uncertainties of the spatial distribution of groundwater to the stream. However,**  
35 **providing such a number would not improve the statements or conclusions we made.**

36 The last point, not unimportant, how is the model treating celerities versus velocities as it was mentioned  
37 several times in the manuscript?

38 **As a Darcy's law based groundwater model, the pressure wave can travel much faster than the water itself. The**  
39 **velocity of the water is the velocity of a single water molecule; whereas, the celerity is the propagation velocity**  
40 **of the pressure signal. It would be possible to provide a graph together with a rather lengthy explanation (~1**  
41 **page) on how velocity and celerity differ but we see this beyond the scope of this article.**  
42

43 2) A point that needs to be improved: You state several times that mixing calculation did not work, bc you  
44 yielded 100% groundwater contribution (L598/599). For me this statement is wrong, as already mentioned in  
45 my previous review. Why is it not working? It is, 100% is a clear result. It just shows that there is no  
46 precipitation water /new water in streamflow. I cannot see how the claim "not working" can be justified. Either  
47 change or elaborate more.

48 **We edited our statements throughout the manuscript which now read as follows:"did not provide a good fit**  
49 **between the observed and calculated data".**  
50

51 3) Other comments that can be improved with minor revisions:

52 P2L46: delete "further"

53 **Deleted**

54 P2L52: "low angle". As commented by me in an earlier review, and again by the other fellow reviewer this time.

55 What is low angle, you define it later as low mountain range and give elevation changes page 5 L126. But if you  
56 call it angle, that it should be related to the slope (i.e. mean, median slope of the catchment). But low

57 mountain and low angle are completely different things. If you have mountains, than you cannot have low  
58 angle, which mean to me that the catchments is more or less flat.  
59 We removed the term low angle as it seems to be confusing. Instead we added to P5L129 “with an average  
60 slope of 8.0%”.  
61 P3L90: add “partly” between “is” and “intercepted”  
62 Edited as suggested by the reviewer.  
63 P7L194: You are using elevation/altitude/height inconsistent. Furthermore, it must be elevation at this  
64 instance. See: McVicar, T. R., & Körner, C. (2013). On the use of elevation, altitude, and height in the ecological  
65 and climatological literature. *Oecologia*, 171(2), 335-337.  
66 Corrected throughout the manuscript following reviewer’s suggestion.  
67 P8L222. The Newman et al. reference is not in the literature list. Please add, please also recheck the  
68 consistency between text and reference list  
69 We added the Newman et al. (2009) reference to the list which we carefully checked again.  
70 L233: I think the current terminology also refers to piezometer samples as travel time, not as residence times..  
71 Changed accordingly.  
72 L281-283. Please rewrite the sentence  
73 We have split the sentence into two and rewritten them.  
74 L291: replace “could be” by “was”  
75 Edited as suggested by the reviewer.  
76 L292: What is unclear to me is if there was not any snow in 2011-2012 or if it was not sampled  
77 The snow was unfortunately not sampled in winter 2011-2012. We added the following to the sentence:  
78 “where no snow sampling could be conducted”.  
79 L294: change “Further” to “Furthermore”  
80 Edited as suggested by the reviewer.  
81 L300 “sampl ed” to “sampled”  
82 Edited as suggested by the reviewer.  
83 L301: “lower...events”. I don’t understand what this sentence should mean.  
84 We have split the sentence into two and rewritten them.  
85 L358-360: Rewrite  
86 We rewrote these sentences.  
87 L375: Contrary to the earlier findings of 100% Groundwater, you can see here the rainfall influence on  
88 streamflow. How big is the contribution of new water to streamflow, can you explain the inconsistency with  
89 the findings that groundwater contributes 100%, and how would this effect your general interpretation.  
90 As we do not have sequential data from storm events but generally only grab sample data, we cannot perform  
91 a hydrograph separation only with the one data point that we are describing in this section. We also do not  
92 think that this finding contradicts with our general statement that the stream water is mainly recharged by  
93 groundwater. As already mentioned in the manuscript “this outlier could be explained by precipitation  
94 contributing to stream flow causing more enriched isotopic values in stream water, which approached average  
95 precipitation  $\delta$ -values ( $-43.9 \pm 23.4$ )”. We mention several times in the manuscript that “groundwater  
96 predominantly feeds baseflow”.  
97 L379: I see it somewhat useless to report the results of a bad fit. I recommend deleting 379-381, and just report  
98 that no good fit was achieved. A numeric result might mislead.  
99 Edited as suggested by the reviewer.  
100 L383: delete “Even a”  
101 Edited as suggested by the reviewer.  
102 L389: I would not write that it was assumed that the signal directly transfers. Just write what you see.  
103 We deleted this statement.  
104 L398: Again, I would not report the ages  
105 Edited as suggested by the reviewer.  
106 L420: can you also extend this to streamflow  
107 See comment further up.  
108 L565: But you got insights to the source of streamflow. The source is groundwater.  
109 We deleted the following: “and, consequently, provided little insight into time and source-components  
110 connection”.  
111 L576: “allowed” to “able”  
112 Edited as suggested by the reviewer.

113 L598: For me this is not obvious at all, if it would be obvious, you would not achieve any result  
114 We edited the sentence as follows: “Just by comparing mean precipitation, stream, and groundwater isotopic  
115 signatures (Table 1), one could expect that simple mixing calculations would not work, i.e. showing  
116 predominant groundwater contribution.”  
117 L1058: Does this statement hold, under the recent developments following the work of Botter (2010, 2011).  
118 The work of Botter represents a promising new theory to model water transport in catchments. However, the  
119 mentioned papers do not deal with the question how these models can be validated using tracer data. The  
120 articles provide no ready-to-apply method to overcome the effect of smoothed out seasonal variation of the  
121 isotopic composition. We cannot identify any contradiction of the statement from this work, but have cited  
122 Duvert et al. 2016 in section 4.4, who report this statement very recently.  
123 L1059: You don’t have a flat line....  
124 We deleted “flat line” and revised the sentence: “in barely varying isotopic signals”.  
125

126 **Exploring water cycle dynamics by sampling multiple**  
127 **stable water isotope pools in a developed landscape of**  
128 **Germany**

129

130 **N. Orlowski<sup>1,2</sup>, P. Kraft<sup>1</sup>, J. Pferdmenges<sup>1</sup> and L. Breuer<sup>1,3</sup>**

131 [1]{Institute for Landscape Ecology and Resources Management (ILR), Research Centre for  
132 BioSystems, Land Use and Nutrition (iFZ), Justus Liebig University Giessen, Giessen,  
133 Germany}

134 [2]{Global Institute for Water Security, University of Saskatchewan, Saskatoon, Canada}

135 [3]{Centre for International Development and Environmental Research, Justus Liebig  
136 University Giessen, Germany}

137 Correspondence to: N. Orlowski (Natalie.Orlowski@umwelt.uni-giessen.de)

138

139 **Abstract**

140 A dual stable water isotope ( $\delta^2\text{H}$  and  $\delta^{18}\text{O}$ ) study was conducted in the developed (managed)  
141 landscape of the Schwingbach catchment (Germany). The two-year weekly to biweekly  
142 measurements of precipitation, stream, and groundwater isotopes revealed that surface and  
143 groundwater are isotopically disconnected from the annual precipitation cycle but showed  
144 bidirectional interactions between each other. Apparently, snowmelt played a fundamental role  
145 for groundwater recharge explaining the observed differences to precipitation  $\delta$ -values.

146 A spatially distributed snapshot sampling of soil water isotopes in two soil depths at 52  
147 sampling points across different land uses (arable land, forest, and grassland) revealed that top  
148 soil isotopic signatures were similar to the precipitation input signal. Preferential water flow  
149 paths occurred under forested soils explaining the isotopic similarities between top and subsoil  
150 isotopic signatures. Due to human-impacted agricultural land use (tilling and compression) of  
151 arable and grassland soils, water delivery to the deeper soil layers was reduced, resulting in  
152 significant different isotopic signatures. However, the land use influence became less  
153 pronounced with depth and soil water approached groundwater  $\delta$ -values. Seasonally tracing  
154 stable water isotopes through soil profiles showed that the influence of new percolating soil

155 water decreased with depth as no remarkable seasonality in soil isotopic signatures was obvious  
156 at depth >0.9 m and constant values were observed through space and time. Since classic  
157 isotope evaluation methods such as transfer function based mean transit time calculations  
158 failed did not provide a good fit between the observed and calculated data, we established a  
159 hydrological model to estimate spatially distributed groundwater ages and flow directions  
160 within the Vollnkirchener Bach subcatchment. Our model revealed that complex age dynamics  
161 exist within the subcatchment and that much of the runoff must have been stored for much longer  
162 than event water (average water age is 16 years). Tracing stable water isotopes through the  
163 water cycle in combination with our hydrological model was valuable for determining  
164 interactions between different water cycle components and unravelling age dynamics within  
165 the study area. This knowledge can further improve catchment specific process understanding  
166 of developed, human-impacted landscapes.

## 167 **1 Introduction**

168 The application of stable water isotopes as natural tracers in combination with hydrodynamic  
169 methods has been proven to be a valuable tool for studying the origin and formation of  
170 recharged water as well as the interrelationship between surface water and groundwater  
171 (Blasch and Bryson, 2007), partitioning evaporation and transpiration (Wang and Yakir,  
172 2000), and ~~further~~ mixing processes between various water sources (Clark and Fritz, 1997c).  
173 Particularly in catchment hydrology, stable water isotopes play a major role since they can be  
174 utilised for hydrograph separations (Buttle, 2006), to calculate the mean transit time (McGuire  
175 and McDonnell, 2006), to investigate water flow paths (Barthold et al., 2011), or to improve  
176 hydrological model simulations (Windhorst et al., 2014). However, most of our current  
177 understanding is resulting from studies in forested catchments. Spatio-temporal studies of  
178 stream water in ~~low angle~~, developed, agricultural dominated, and managed catchments are  
179 less abundant. This is partly caused by damped stream water isotopic signatures excluding  
180 traditional hydrograph separations in low-relief catchments (Klaus et al., 2015). Unlike the  
181 distinct watershed components found in steeper headwater counterparts, lowland areas often  
182 exhibit a complex groundwater–surface water interaction (Klaus et al., 2015). Sklash and  
183 Farvolden (1979) showed that groundwater plays an important role as a generating factor for  
184 storm and snowmelt runoff processes. In many catchments, streamflow responds promptly to  
185 rainfall inputs but variations in passive tracers such as water isotopes are often strongly  
186 damped (Kirchner, 2003). This indicates that storm runoff in these catchments is dominated  
187 mostly by “old water” (Buttle, 1994; Neal and Rosier, 1990; Sklash, 1990). However, not all

188 “old water” is the same (Kirchner, 2003). This catchment behaviour was described by  
189 Kirchner (2003) as the old water paradox. Thus, there is evidence of complex age dynamics  
190 within catchments and much of the runoff is stored in the catchment for much longer than  
191 event water (Rinaldo et al., 2015). Still, some of the physical processes controlling the release  
192 of “old water” from catchments are poorly understood, roughly modelled, and the observed  
193 data do not suggest a common catchment behaviour (Botter et al., 2010). However, old water  
194 paradox behaviour was observed in many catchments worldwide but it may have the strongest  
195 effect in agriculturally managed catchments, where surprisingly only small changes in stream  
196 chemistry have been observed (Hrachowitz et al., 2016).

197 Moreover, almost all European river systems were already substantially modified by humans  
198 before river ecology research developed (Allan, 2004). Through changes in land use, land  
199 cover, irrigation, and draining, agriculture has substantially modified the water cycle in terms  
200 of both quality and quantity (Gordon et al., 2010) as well as hydrological functioning (Pierce et  
201 al., 2012). Hrachowitz et al. (2016) recently stated the need for a stronger linkage between  
202 catchment-scale hydrological and water quality communities. Further, McDonnell et al. (2007)  
203 concluded that we need to figure out a way to embed landscape heterogeneity or the  
204 consequence of the heterogeneity (i.e. of agricultural dominated and managed catchments) into  
205 models as current generation catchment-scale hydrological and water quality models are poorly  
206 linked (Hrachowitz et al., 2016).

207 One way to better understand catchment behaviour and the interaction among the various water  
208 sources (surface, subsurface, and groundwater) and their variation in space and time is a detailed  
209 knowledge about their isotopic composition. In principal, isotopic signatures of precipitation  
210 are altered by temperature, amount (or rainout), continental, altitudinal, and seasonal effects.  
211 Stream water isotopic signatures can reflect precipitation isotopic composition and moreover,  
212 dependent on discharge variations be affected by seasonally variable contributions of different  
213 water sources such as bidirectional water exchange with the groundwater body during baseflow,  
214 or high event-water contributions during stormflow (Genereux and Hooper, 1998; Koeniger et  
215 al., 2009). Precipitation falling on vegetated areas is partly intercepted by plants and re-  
216 evaporated isotopically fractionated. The remaining throughfall infiltrates slower and can be  
217 affected by evaporation resulting in an enrichment of heavy isotopes, particularly in the upper  
218 soil layers (Gonfiantini et al., 1998; Kendall and Caldwell, 1998). In the soil, specific isotopic  
219 profiles develop, characterized by an evaporative layer near the surface. The isotopic

220 enrichment decreases exponentially with depth, representing a balance between the upward  
221 convective flux and the downward diffusion of the evaporative signature (Barnes and Allison,  
222 1988). In humid and semi-humid areas, this exponential decrease is generally interrupted by the  
223 precipitation isotopic signal. Hence, the combination of the evaporation effect and the  
224 precipitation isotopic signature determine the isotope profile in the soil (Song et al., 2011).  
225 Once soil water reaches the saturated zone, this isotope information is finally transferred to the  
226 groundwater (Song et al., 2011). Soil water can therefore be seen as a link between precipitation  
227 and groundwater, and the dynamics of isotopic composition in soil water are indicative of the  
228 processes of precipitation infiltration, evaporation of soil water, and recharge to groundwater  
229 (Blasch and Bryson, 2007; Song et al., 2011).

230 We started our research with results obtained through an earlier study in the managed  
231 Schwingbach catchment that implied a high responsiveness of the system to precipitation inputs  
232 indicated by very fast rises in discharge and groundwater head levels (Orlowski et al., 2014).  
233 However, as there was only a negligible influence of the precipitation input signal on the stable  
234 water isotopic composition in streams, our initial data set showed evidence for complex age  
235 dynamics within the catchment. Nevertheless, a rapid flow response to a precipitation input may  
236 also be mistaken (as conceptualized in the vast majority of catchment-scale conceptual  
237 hydrological models) as the actual input signal already reaching the stream, while in reality it  
238 is the remainder of past input signals that slowly travelled through the system (Hrachowitz et  
239 al., 2016). The observable hydrological response therefore acts at different time scales than the  
240 tracer response (Hrachowitz et al., 2016) as described by the celerity vs. velocity concept  
241 (McDonnell and Beven, 2014). The observed patterns in our catchment therefore inspired us to  
242 use a combined approach of hydrodynamic data analyses, stable water isotope investigations,  
243 and data-driven hydrological modelling to determine catchment dynamics (response times and  
244 groundwater age patterns) and unravel water flow paths at multiple spatial scales. This work  
245 should further improve our knowledge on hydrological flow paths in developed, human-  
246 impacted catchments.

## 247 **2 Materials and methods**

### 248 **2.1 Study area**

249 The research was carried out in the Schwingbach catchment (50°30'4.23"N, 8°33'2.82"E)  
250 (Germany) (Fig. 1a). The Schwingbach and its main tributary the Vollnkirchener Bach are low-

251 mountainous creeks having an altitudinal difference of 50–100 m over 5 km distance (Perry and  
252 Taylor, 2009) (Fig. 1c) with an altered physical structure of the stream system (channelled  
253 stream reaches, pipes, drainage systems, fishponds). The Schwingbach catchment (9.6 km<sup>2</sup>)  
254 ranges from 233–415 m a.s.l. ~~in altitude with an average slope of 8.0%~~. The Vollnkirchener  
255 Bach tributary is 4.7 km in length and drains a 3.7 km<sup>2</sup> subcatchment area (Fig. 1c), with  
256 elevations from 235–351 m a.s.l. Almost 46% of the overall Schwingbach catchment is  
257 forested, which slightly exceeds agricultural land use (35%) (Fig. 1c). Grassland (10%) is  
258 mainly distributed along streams and smaller meadow orchards are located around the villages.

259 The Schwingbach main catchment is underlain by argillaceous shale in the northern parts,  
260 serving as aquicludes. Graywacke zones with lydite in the central, as well as limestone, quartzite,  
261 and sandstone regions in the headwater area provide aquifers with large storage capacities (Fig.  
262 1f). Loess covers Paleozoic bedrock at north- and east bounded hillsides (Fig. 1f). Streambeds  
263 consists of sand and debris covered by loam and some larger rocks (Lauer et al., 2013). Many  
264 downstream sections of both creeks are framed by armor stones (Orlowski et al., 2014). The  
265 dominant soil types in the overall study area are Stagnosols (41%) and mostly forested  
266 Cambisols (38%). Stagnic Luvisols with thick loess layers, Regosol, Luvisols, and Anthrosols  
267 are found under agricultural use and Gleysols under grassland along the creeks.

268 [Figure 1 near here]

269 The climate is classified as temperate with a mean annual temperature of 8.2°C. An annual  
270 precipitation sum of 633 mm (for the hydrological year 1 November 2012 to 31 October 2013)  
271 was measured at the catchment's climate station (site 13, Fig. 1b). The year 2012 to 2013 was  
272 an average hydrometeorological year. For comparison, the climate station Giessen/Wettenberg  
273 (25 km N of the catchment) operated by the German Meteorological Service (DWD, 2014)  
274 records a mean annual temperature of 9.6 °C and a mean annual precipitation sum of 666±103  
275 mm for the period 1980–2010. Discharge peaks from December to April (measured by the use  
276 of RBC-flumes with maximum peak flow of 114 L s<sup>-1</sup>, Eijkelkamp Agrisearch Equipment,  
277 Giesbeek, NL) and low flows occur from July until November. Substantial snowmelt peaks  
278 were observed during December 2012 and February 2013. Furthermore, May 2013 was an  
279 exceptional wet month characterised by discharge of 2–3 mm d<sup>-1</sup>. A detailed description of  
280 runoff characteristics is given by Orlowski et al. (2014).



## 281 **2.2 Monitoring network and water isotope sampling**

282 The monitoring network consists of an automated climate station (site 13, Fig. 1 b–c) (Campbell  
283 Scientific Inc., AQ5, UK; equipped with a CR1000 data logger), three tipping buckets, and 15  
284 precipitation collectors, six stream water sampling points, and 22 piezometers (Fig. 1 b–c).  
285 Precipitation data were corrected according to Xia (2006).

286 Two stream water sampling points (sites 13 and 18) in the Vollnkirchener Bach are installed  
287 with trapezium shaped RBC-flumes for gauging discharge (Eijkelkamp Agrisearch Equipment,  
288 Giesbeek, NL), and a V-weir is located at sampling point 64. RBC-flumes and V-weir are  
289 equipped with Mini-Divers® (Eigenbrodt Inc. & Co. KG, Königsmoor, DE) for automatically  
290 recording water levels. Discharge at the remaining stream sampling points was manually  
291 measured applying the salt dilution method (WTW-cond340i, WTW, Weilheim, DE). The 22  
292 piezometers (Fig. 1b) are made from perforated PVC tubes sealed with bentonite at the upper  
293 part of the tube to prevent contamination by surface water. For monitoring shallow groundwater  
294 levels, either combined water level/temperature loggers (Odyssey Data Flow System,  
295 Christchurch, NZ) or Mini-Diver® water level loggers (Eigenbrodt Inc. & Co. KG,  
296 Königsmoor, DE) are installed. Accuracy of Mini-Diver® is  $\pm 5$  mm and for Odyssey data  
297 logger  $\pm 1$  mm. For calibration purposes, groundwater levels are additionally measured  
298 manually via an electric contact gauge.

299 Stable water isotope samples of rainfall, stream-, and groundwater were taken from July 2011  
300 to July 2013 on weekly intervals. In winter 2012–2013, snow core samples over the entire snow  
301 depth of  $<0.15$  m were collected in tightly sealed jars at same sites as open rainfall was sampled.  
302 We sampled shortly after snow fall because sublimation, recrystallization, partial melting,  
303 rainfall on snow, and redistribution by wind can alter the isotopic composition (Clark and Fritz,  
304 1997b). Samples were melted overnight following Kendall and Caldwell (1998) and analysed  
305 for their isotopic composition. Open rainfall was collected in self-constructed samplers as per  
306 Windhorst et al. (2013). Grab samples of stream water were taken at six locations, three  
307 sampling points at each stream (Fig. 1b–c). Since spatial isotopic variations of groundwater  
308 among piezometers under meadow were small, samples were collected at three out of eight  
309 sampling points under meadow (sites 1, 6, and 21), five under the arable field (sites 25–29), and  
310 four next to the Vollnkirchener Bach (sites 24, 31, 32, and 35) (Fig. 1b). Additionally, a  
311 drainage pipe (site 15) located  $\sim 226$  m downstream of site 18 was sampled. According to IAEA

312 standard procedures, all samples were filled and stored in 2 mL brown glass vials, sealed with  
313 a solid lid, and wrapped up with Parafilm®.

## 314 **2.3 Isotopic soil sampling**

### 315 **2.3.1 Spatial variability**

316 In order to analyse the effect of small-scale characteristics such as distance to stream, TWI, and  
317 land use on soil isotopic signatures, we sampled a snapshot of 52 points evenly distributed over  
318 a 200 m grid around the Vollnkirchener Bach (Fig. 1d). Soil samples were taken at four  
319 consecutive rainless days (1 to 4 November 2011) at altitudeselevations of 235–294 m a.s.l..  
320 Sampling sites were selected via a stratified, GIS-based sampling plan (ArcGIS, Arc Map  
321 10.2.1, Esri, California, USA), including three classes of topographic wetness indices (TWIs:  
322 4.4–6.5; 6.5–7.7; 7.7–18.4), two different distances to stream (0–121 m, 121–250 m), and three  
323 land uses (arable land, forest, and grassland), with each class containing the same number of  
324 sampling points. Samples were collected at depths of 0.2 m and 0.5 m. Gravimetric water  
325 content was measured according to DIN-ISO 11465 by drying soils for 24 h at 110 °C. Soil pH  
326 was analysed following DIN-ISO 10390 on 1:1 soil-water-mixture with a handheld pH-meter  
327 (WTW cond340i, WTW Inc., DE). Bulk density was determined according to DIN-ISO 11272,  
328 and soil texture by finger testing.

### 329 **2.3.2 Seasonal isotope soil profiling and isotope analysis**

330 In order to trace the seasonal development of stable water isotopes from rainfall to groundwater,  
331 seven soil profiles were taken in the dry summer season (28 August 2011), seven in the wet  
332 winter period (28 March 2013), and two profiles in spring (24 April 2013) under different  
333 vegetation cover (arable land and grassland) (Fig. 1d). Soil was sampled utilising a hand-auger  
334 (Eijkelpkamp Agrisearch Equipment BV, Giesbeek, DE) from the soil surface to 2 m depth.  
335 Samples were collected in greater detail near the soil surface since this area is known to have  
336 the greatest isotopic variability (Barnes and Allison, 1988).

337 Soil samples were stored in amber glass tubes, sealed with Parafilm®, and kept frozen until  
338 water extraction. Soil water was extracted cryogenically with 180 min extraction duration, a  
339 vacuum threshold of 0.3 Pa, and an extraction temperature of 90°C following Orłowski et al.  
340 (2013). Isotopic signatures of  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  were analysed via off-axis integrated cavity output  
341 spectroscopy (OA-ICOS) (DLT-100, Los Gatos Research Inc., Mountain View, USA). Within

342 each isotope analysis three calibrated stable water isotope standards of different water isotope  
343 ratios were included (LGR working standard number 1, 3, and 5; Los Gatos Research Inc., CA,  
344 US). After every fifth sample the LGR working standards are measured. For each sample, six  
345 sequential 900  $\mu\text{L}$  aliquot of a water sample are injected into the analyser. Then, the first three  
346 measurements are discarded. The remaining are averaged and corrected for per mil scale  
347 linearity following the IAEA laser spreadsheet template (~~Newman et al., 2009~~); (Newman et al.,  
348 2009). Following this IAEA standard procedure allows for drift and memory corrections.  
349 Isotopic ratios are reported in per mil (‰) relative to Vienna Standard Mean Ocean Water  
350 (VSMOW) (Craig, 1961b). Accuracy of analyses was 0.6‰ for  $\delta^2\text{H}$  and 0.2‰ for  $\delta^{18}\text{O}$  (LGR,  
351 2013). Leaf water extracts typically contain a high fraction of organic contaminations, which  
352 might lead to spectral interferences when using isotope ratio infrared absorption spectroscopy,  
353 causing erroneous isotope values (Schultz et al., 2011). However, for soil water extracts there  
354 exists no need to check or correct such data (Schultz et al., 2011; Zhao et al., 2011).

#### 355 **2.4 Mean transit time estimation**

356 To understand the connection between the different water cycle components in the  
357 Schwingbach catchment, mean transit times (MTT) for both streams as well as ~~mean residence~~  
358 ~~times (MRT)~~ from precipitation to groundwater were calculated using FlowPC (Maloszewski  
359 and Zuber, 2002). See Appendix I for details about the applied method.

#### 360 **2.5 Model-based groundwater age dynamics**

361 To estimate the age dynamics of the groundwater body in the Vollnkirchener Bach  
362 subcatchment, a hydrological model was established on the basis of the conceptual model  
363 presented by Orłowski et al. (2014) and the isotopic measurements presented here. Appendix  
364 II outlines the modelling concept, model set up, and its parameterization.

#### 365 **2.6 Statistical analyses**

366 For statistical analyses, we used IBM SPSS Statistics (Version 22, SPSS Inc., Chicago, IL, US)  
367 and R (version Rx64 3.2.2). The R package igraph was utilized for plotting (Csardi and Nepusz,  
368 2006). Studying temporal and spatial variations in meteoric and groundwater, isotope data were  
369 tested for normal distribution. Subsequently, t-tests or Multivariate Analyses of Variances  
370 (MANOVAs) were applied and Tukey-HSD tests were run to determine which groups were

371 significantly different ( $p \leq 0.05$ ). Event mean values of isotopes in precipitation, stream, and  
372 groundwater were calculated when no spatial variation was observed. Regression analyses were  
373 run to determine the effect of small-scale characteristics such as distance to stream, TWI, and  
374 land use on soil isotopic signatures.

375 We used a topology inference network map (Kolaczyk, 2014) in combination with a principal  
376 component analysis to show  $\delta^{18}\text{O}$  isotope relationships between surface and groundwater  
377 sampling points. To explore the sensitivity of missing data, we used both the complete isotope  
378 time series and randomly selected 80% of the whole data sets. Overall, the cluster relationships  
379 of the surface and groundwater sampling points are largely similar for both whole and subsets  
380 of isotope data sets, despite some differences of the exact cluster centroid locations. We  
381 therefore decided to use randomly selected 80% of the isotope time series to illustrate our  
382 results. In the network map, each node of the network represents an isotope sampling point.  
383 The locations of the nodes are based on the first two components (PC1 and PC2). The  
384 correlations between isotope time series are represented by the edges connecting nodes. The  
385 thickness of edges characterizes the strength of the correlations. The p-values of correlations  
386 are approximated by using the F-distributions and mid-ranks are used for the ties (Hollander et  
387 al., 2013). Only statistically significant connections ( $p < 0.05$ ) are shown.

388 To compare different water sources on the catchment-scale, a local meteoric water (LMWL)  
389 line was developed and evaporation water lines (EWLs) were used. They represent the linear  
390 relationship between  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  of meteoric waters (Cooper, 1998) in contrast to the global  
391 meteoric water line (GMWL), which describes the world-wide average stable isotopic  
392 composition in precipitation (Craig, 1961a). Identifying the origin of water vapour sources and  
393 moisture recycling (Gat et al., 2001; Lai and Ehleringer, 2011), the deuterium-excess (d-  
394 excess), defined by Dansgaard (1964) as  $d = \delta^2\text{H} - 8 \times \delta^{18}\text{O}$  was used.

395 For comparisons, precipitation isotope data from the closest GNIP (Global Network of Isotopes  
396 in Precipitation) station Koblenz (DE; 74 km SW of the study area, 97 m a.s.l.) was used (IAEA,  
397 2014; Stumpp et al., 2014). For monthly comparisons with Schwingbach d-excess values, we  
398 used a data set from the GNIP station Koblenz that includes 24 values starting from July 2011  
399 to July 2013.

## 400 3 Results

### 401 3.1 Variations of precipitation isotopes and d-excess

402 The  $\delta^2\text{H}$  values of all precipitation isotope samples ranged from  $-167.6$  to  $-8.3\text{‰}$  (Table 1). To  
403 examine the spatial isotopic variations, rainfall was collected at 15 open field site locations  
404 throughout the Schwingbach main catchment (Fig. 1b–c) for a 7-month period, but no spatial  
405 variation could be observed. Thus, rainfall was collected at the catchment outlet (site 13) from  
406 23 October 2014 onward. ~~Analysing effects that influence the isotopic composition of~~  
407 ~~precipitation, We could~~ neither identify an amount effect nor an altitude effect ~~were found—~~  
408 ~~surprisingly, as the~~ in our precipitation isotope data. The greatest altitudinal difference between  
409 sampling points was also only 101 m. Nevertheless, a slight temperature effect ( $R^2=0.5$  for  $\delta^2\text{H}$   
410 and  $R^2=0.6$  for  $\delta^{18}\text{O}$ , respectively) was observed showing enriched isotopic signatures at higher  
411 temperatures.

412 [Table 1 near here]

413 Strong temporal variations in precipitation isotopic signatures, as well as pronounced seasonal  
414 isotopic effects were measured with greatest isotopic differences occurring between summer  
415 and winter. Samples taken in the fall and spring were isotopically similar, however, differed  
416 from winter isotopic signature, which were somewhat lighter (Fig. 2). Furthermore, in the  
417 winter of 2012–13 snow ~~could be~~ was sampled, which decreased the mean winter isotopic values  
418 for this period in comparison to the previous winter period (2011–12) ~~;) where no snow sampling~~  
419 ~~could be conducted.~~ The mean  $\delta^2\text{H}$  isotope values of snow samples were approximately 84‰  
420 lighter than mean precipitation isotopic signatures (Fig. 3). ~~Further~~ Furthermore, no statistically  
421 significant ( $p>0.05$ ) inter-annual variation was detected between the summer periods of 2011  
422 and 2012 (Fig. 2).

423 [Figure 2 near here]

424 Examining the influence of moisture recycling on the isotopic compositions of precipitation,  
425 the d-excess was calculated for each individual rain event at the Schwingbach catchment. D-  
426 excess values ranged from  $-7.8\text{‰}$  to  $+19.4\text{‰}$  and averaged  $+7.1\text{‰}$  (Fig. 2). In general, 37% of  
427 all events were ~~sampled~~ sampled in summer periods (21 June to 21/22 September) ~~and~~. These  
428 summer events showed lower d-excess values in comparison to the 19% winter precipitation  
429 events (21/22 December to 19/20 March) (Fig. 2). D-excess greater than  $+10\text{‰}$  was determined  
430 for 22% of all events. Lowest values corresponded to summer precipitation events where

431 evaporation of the raindrops below the cloud base may occur. Most of the higher values  
432 (>+10‰) appeared in cold seasons (fall/winter) and winter snow samples of the Schwingbach  
433 catchment with much depleted  $\delta$ -values showed highest d-excess (Fig. 2).

434 In comparison with the GNIP station Koblenz (2011–2013), the mean annual d-excess at the  
435 Schwingbach catchment was on average 3.9‰ higher, showing a greater impact of oceanic  
436 moisture sources than the further south-west located station Koblenz. The long-term mean d-  
437 excess was 4.4‰ for the Koblenz station (1978–2009) (Stumpp et al., 2014). Highest d-  
438 excesses at the GNIP station matched highest values in the Schwingbach catchment, both  
439 occurring in the cold seasons (October to December 2011 and November to December 2012).

440 The linear relationship of  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  content in local precipitation, results in a local meteoric  
441 water line (LMWL) (Fig. 3). The slope of the Schwingbach LMWL is well in agreement with  
442 the one from the GNIP station Koblenz ( $\delta^2\text{H}=7.66\times\delta^{18}\text{O}+2.0\text{‰}$ ;  $R^2=0.97$ ; 1978–2009 (Stumpp  
443 et al., 2014)), but is slightly lower in comparison to the GMWL, showing stronger local  
444 evaporation conditions. Since evaporation causes a differential increase in  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  values  
445 of the remaining water, the slope for the linear relationship between  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  is lower in  
446 comparison to the GMWL (Rozanski et al., 2001; Wu et al., 2012).

447 [Figure 3 near here]

## 448 **3.2 Isotopes of soil water**

### 449 **3.2.1 Spatial variability**

450 Determining the impact of landscape characteristics on soil water isotopic signatures, we found  
451 no statistically significant connection between the parameters distance to stream, TWI, soil  
452 water content, soil texture, pH, and bulk density with the soil isotopic signatures in both soil  
453 depths, except for land use.

454 [Table 2 near here]

455 The mean  $\delta$ -values in the top 0.2 m of the soil profile are higher than in the subsoil, reflecting  
456 a stronger impact of precipitation in the topsoil (Table 2, Fig. 4). While the  $\delta$ -values for subsoil  
457 and precipitation differed significantly ( $p\leq 0.05$ ), they did not for topsoil (Fig. 4). Subsoil  
458 isotopic values were statistically equal to stream water and groundwater (Fig. 4).

459 [Figure 4 near here]

460 Generally, all soil water isotopic values fell on the LMWL, indicating no evaporative  
461 enrichment (Fig. 5). Comparing soil isotopic signatures between different land covers showed  
462 generally higher and statistically significantly different  $\delta$ -values ( $p \leq 0.05$ ) at 0.2 m soil depth  
463 under arable land as compared to forests and grasslands. For the lower 0.5 m of the soil column,  
464 isotopic signatures under all land uses showed statistically similar values. Comparing soil water  
465  $\delta^2\text{H}$  values between top and subsoil under different land use units showed significant  
466 differences ( $p \leq 0.05$ ) under arable and grassland but not under forested sites (Fig. 5).

467 [Figure 5 near here]

### 468 3.2.2 Seasonal isotope soil profiling

469 Isotope compositions of soil water varied seasonally: More depleted soil water was found in  
470 the winter and spring (Fig. 6); contrary, soil water was enriched in summer due to evaporation  
471 during warmer and drier periods (Darling, 2004). For summer soil profiles in the Vollnkirchener  
472 subcatchment, no evidence for evaporation was obvious below 0.4 m soil depth. However,  
473 snowmelt isotopic signatures could be traced down to a soil depth of 0.9 m during spring rather  
474 than winter, pointing to a depth-translocation of meltwater in the soil, more remarkable for the  
475 deeper profile under arable land (Fig. 6, upper left panel). Furthermore, shallow soil water (<0.4  
476 m) showed larger standard deviations with values closer to mean seasonal precipitation inputs  
477 (Fig. 6, upper panels). Winter profiles exhibited somewhat greater standard deviations in  
478 comparison to summer isotopic soil profiles. The observed seasonal amplitude became less  
479 pronounced with depth as soil water isotope signals approached groundwater average in >0.9 m  
480 depth. Generally, deeper soil water isotope values were relatively constant through time and  
481 space.

482 [Figure 6 near here]

### 483 3.3 Isotopes of stream water

484 No statistically significant differences were found between the Schwingbach and  
485 Vollnkirchener Bach stream water (Fig. 7) ~~with~~. All stream water isotope ~~data fallingsamples~~  
486 fell on the LMWL ~~but showing slight evaporative enrichment~~ except for few evaporatively  
487 enriched samples (Fig. 3).  $\delta^{18}\text{O}$  values varied for the Vollnkirchener Bach by  $-8.4 \pm 0.4\%$  and  
488 for the Schwingbach by  $-8.4 \pm 0.6\%$  (Table 1). Stream water isotopic signatures were by

489 approximately  $-15\%$  in  $\delta^2\text{H}$  more depleted than precipitation signatures and similar to  
490 groundwater (Table 1).

491 [Figure 7 near here]

492 A damped seasonality of the isotope concentration in stream water versus precipitation was  
493 occurring between summer and winter (Fig. 7). Most outlying depleted stream water isotopic  
494 signatures (e.g. in March 2012 and 2013) can be explained by snowmelt (Fig. 7). However, the  
495 outlier at the Schwingbach stream water sampling site 64 ( $-66.7\%$  for  $\delta^2\text{H}$ ) is by  $8.5\%$  more  
496 depleted than the two-year average of Schwingbach stream water (Table 1). Rainfall falling on  
497 24 September 2012 was  $-31.9\%$  for  $\delta^2\text{H}$ . This period in September was generally characterized  
498 by low flow and little rainfall. Thus, little contribution of new water was observed and stream  
499 water isotopic signatures were groundwater-dominated. For site 13, the outlier in May 2012  
500 ( $-44.2\%$  for  $\delta^2\text{H}$ ) was by  $13.8\%$  more enriched than the average stream water isotopic  
501 composition of the Vollnkirchener Bach over the two-year observation period (Table 1). A  
502 runoff peak at site 13 of  $0.15 \text{ mm d}^{-1}$  and a  $2.9 \text{ mm}$  rainfall event were recorded on 23 May  
503 2012. Thus, this outlier could be explained by precipitation contributing to stream flow causing  
504 more enriched isotopic values in stream water, which approached average precipitation  $\delta$ -values  
505 ( $-43.9 \pm 23.4$ ).

506 ~~Calculated MTT calculations for the Schwingbach ranged from 52–67 weeks and for the~~  
507 ~~Vollnkirchener Bach from 47–66 weeks, whereby linear and exponential models provided the~~  
508 ~~best fits for all sampling points. However, the calculated output data did not provide a good fit~~  
509 ~~the observed values in terms of the quality criteria sigma and model efficiency (Timbe et al.,~~  
510 ~~2014) ( $ME_{\text{Schwingbach}} -0.1-0.0$ ,  $ME_{\text{Vollnkirchener Bach}} 0.0-0.4$ ; sigma for all sampling points 0.1).~~  
511 ~~Even a bias correction of the input data did not improve the model outputs (sigma=0.1).~~  
512 ~~Therefore, we conclude that transfer function based MTT estimation methods applying stable~~  
513 ~~water isotope data failed for the Schwingbach.~~

### 514 3.4 Isotopes of groundwater

515 ~~Since groundwater head levels responded almost as quickly as streamflow to rainfall events,~~  
516 ~~rainfall isotopic signatures were assumed to be rapidly transferred to the groundwater.~~ For the  
517 piezometers under meadow, almost constant isotopic values (Fig. 8, Table 1) were observed  
518 ( $\delta^2\text{H}$ :  $-57.6 \pm 1.6\%$ ). Most depleted groundwater isotopic values ( $< -80\%$  for  $\delta^2\text{H}$ ) were  
519 measured for piezometer 32 during snowmelt events in March and April 2013 as well as for



520 piezometer 27 from December 2012 to February 2013. Piezometer 32 is highly responsive to  
521 rainfall-runoff events and groundwater head elevations showed significant correlations with  
522 mean daily discharge at this site (Orlowski et al., 2014).

523 Groundwater under meadow differed from mean precipitation values by about  $-14\%$  for  $\delta^2\text{H}$   
524 showing no evidence of a rapid transfer of rainfall isotopic signatures to the groundwater (Fig.  
525 8). ~~This was underlined by~~For the ~~results of our MRTMTT~~ estimations ~~which varied between~~  
526 ~~56–65 weeks for of~~ the thirteen ~~considered~~ piezometers. ~~However~~, the calculated output data  
527 did not fit the observed values showing very low MEs (ME:  $-0.62$ – $-0.09$  for  $\delta^{18}\text{O}$  and  $-0.49$ –  
528  $0.16$  for  $\delta^2\text{H}$ ; sigma:  $0.08$ – $0.15$  for  $\delta^{18}\text{O}$  and  $0.62$ – $1.11$  for  $\delta^2\text{H}$ ).

529 [Figure 8 near here]

530 Due to different water flow paths of groundwater along the studied stream, we expected to find  
531 distinguished groundwater isotopic signatures. In fact, we could identify spatial statistical  
532 differences between grassland and arable land groundwater isotopic signatures (Fig. 9).  
533 Groundwater isotopic signatures under arable land (sites: 25–29, Fig. 1b) showed more  
534 enriched values (Fig. 8) and showed significant correlations ( $p < 0.05$ ) among each other (Fig.  
535 9). Arable land groundwater plotted furthest away from surface water sampling points in our  
536 network map showing no significant correlations to either the Schwingbach or the  
537 Vollnkirchener Bach.  $\delta^{18}\text{O}$  time series of piezometers along the stream and under the meadow  
538 showed closest relations to surface water sampling points (Fig. 9). We further found high  
539 correlations ( $R^2 > 0.6$ ) of  $\delta^{18}\text{O}$  time series of piezometers located under the meadow among each  
540 other. Additionally,  $\delta^{18}\text{O}$  values of piezometer 3 correlated significantly ( $p < 0.05$ ) with surface  
541 water sampling points 18 and 94 ( $R^2 = 0.6$  and  $0.8$ , respectively) and piezometer 32 with  
542 sampling points 13 and 64 ( $R^2 = 0.8$  and  $0.6$ , respectively).

543 [Figure 9 near here]

544 We further observed close relations ( $p < 0.05$ ) among  $\delta^{18}\text{O}$  values of Vollnkirchener Bach  
545 sampling sites 13, 18, and 94 as well as of Schwingbach sites 11, 19, and 64 along with  
546 significant correlations between each other.

### 547 3.5 Groundwater age dynamics

548 Since MTT calculations ~~failed~~did not provide a good fit between the observed and calculated  
549 output data, we modelled the groundwater age in the Vollnkirchener Bach subcatchment using

550 CMF (Appendix II), applying observed hydrometric as well as stable water isotope data (Fig.  
551 10).

552 [Figure 10 near here]

553 The maximum age of water is highly variable throughout the subcatchment, which results in a  
554 heterogeneous spatial age distribution. The groundwater in most of the outer cells is young (0–  
555 10 years), whereas the inner cells, which incorporate the Vollnkirchener Bach, contain older  
556 water (>30 years). The oldest water ( $\geq 55$  years) can be found in the Northern part of the  
557 catchment (Fig. 10, detail view), where the Vollnkirchener Bach drains into the Schwingbach.  
558 The main outlets of the subcatchment (dark red coloured cell and green cell) even reach an age  
559 of 100 and 55 years, respectively. This can be explained by the fact that it is the lowest cell  
560 within the subcatchment and that water accumulates here. The overall flow path to this cell is  
561 the longest and as a consequence the groundwater age in this cell is the highest.

562 In general, 2% of cells contain groundwater that is older than 50 years, <1% reveal ages >70  
563 years, 13% contain water with an age of less than one year, and 52% with an age <15 years.  
564 Thus, most of the cells contain young to moderately old water (<15 years), while few cells  
565 comprise old water (>50 years). The average groundwater age in the Vollnkirchener Bach  
566 subcatchment is 16 years. Correlating the groundwater age against the distance to the stream,  
567 we found a linear correlation ( $R^2=0.3$ ) with a distinct trend. The water tends to be younger with  
568 greater distance to the stream.

569 The amount of flowing water depicted by the length of the arrows is generally higher near the  
570 stream, whereas in most of the outer cells the amount is very low (Fig. 10). The modelled main  
571 flow direction is towards the Vollnkirchener Bach but many arrows show flow direction across  
572 the stream indicating bidirectional water exchange between the stream and the groundwater  
573 body.

## 574 **4 Discussion**

### 575 **4.1 Variations of precipitation isotopes and d-excess**

576 We found no spatial variation in precipitation isotopes throughout the Schwingbach catchment.  
577 Mook et al. (1974) also observed for north-western Europe that precipitation collected over  
578 periods of 8 and 24 h from three different locations within 6 km<sup>2</sup> at the same altitude/elevation  
579 were consistent within 0.3‰ for  $\delta^{18}\text{O}$ . Further, we detected no amount or altitude effect on

580 isotopes in precipitation. Amount effects generally occur most likely in the tropics or for intense  
581 convective rain events and are not a key factor for explaining isotope distributions in German  
582 precipitation (Stumpp et al., 2014).

583 The observed linear relationship ( $\delta^{18}\text{O}=0.44T-12.05\text{‰}$ ) between air temperature and  
584 precipitation  $\delta^{18}\text{O}$  values compares reasonably well with a correlation reported by Yurtsever  
585 (1975) based on north Atlantic and European stations from the GNIP network  
586  $\delta^{18}\text{O}=(0.521\pm 0.014)T-(14.96\pm 0.21)\text{‰}$ . The same is true for a correlation found by Rozanski et  
587 al. (1982) for the GNIP station Stuttgart, 196 km South of the Schwingbach. Stumpp et al.  
588 (2014) analysed long-term precipitation data from meteorological stations across Germany and  
589 found that 23 out of 24 tested stations showed a positive long-term temperature trend over time.  
590 The observed correspondence between the degree of isotope depletion and the temperature  
591 reflects the influence of the temperature effect in the Schwingbach catchment, which mainly  
592 appears in continental, middle–high latitudes (Jouzel et al., 1997). Furthermore, the correlation  
593 between  $\delta^2\text{H}$  in monthly precipitations and local surface air temperature becomes increasingly  
594 stronger towards the centre of the continent (Rozanski et al., 1982). Thus, the observed seasonal  
595 differences in precipitation  $\delta$ -values in the Schwingbach catchment could mainly be attributed  
596 to seasonal differences in air temperature and the presence of snow in the winter of 2012–13  
597 (Fig. 2).

598 Precipitation events originating from oceanic moisture show d-excess values close to +10‰  
599 (Craig, 1961a; Dansgaard, 1964; Wu et al., 2012) and one of the main sources for precipitation  
600 in Germany is moisture from the Atlantic Ocean (Stumpp et al., 2014). Lowest values  
601 corresponded to summer precipitation events where evaporation of the falling raindrops below  
602 the cloud base occurs. Same observations were made by Rozanski et al. (1982) for European  
603 GNIP stations. Winter snow samples of the Schwingbach catchment with very depleted  $\delta$ -  
604 values showed highest d-excess values ( $>+10\text{‰}$ ), well in agreement with results of Rozanski et  
605 al. (1982) for European GNIP stations. The observed differences in d-excess values between  
606 the Schwingbach catchment and the GNIP station Koblenz can be attributed to differences in  
607 elevation range and the different regional climatic settings at both sites (Koblenz is located in  
608 the relatively warmer Rhine river valley).

## 609 4.2 Isotopes of soil water

### 610 4.2.1 Spatial variability

611 We found no statistically significant connection between the parameters distance to stream,  
612 TWI, soil water content, soil texture, pH, and bulk density with the soil isotopic signatures in  
613 both soil depths. This was potentially attributed to the small variation in soil textures (mainly  
614 clayey silts and loamy sandy silts), bulk densities, and pH values for both soil depths (Table 2).  
615 Garvelmann et al. (2012) obtained high resolution  $\delta^2\text{H}$  vertical depth profiles of pore water at  
616 various points along two fall lines of a pasture hillslope in the Black Forest (Germany) by  
617 applying the  $\text{H}_2\text{O}(\text{liquid})\text{--H}_2\text{O}(\text{vapor})$  equilibration laser spectroscopy method. The authors  
618 showed that groundwater was flowing through the soil in the riparian zone (downslope profiles)  
619 and dominated streamflow during baseflow conditions. Their comparison indicated that the  
620 percentage of pore water soil samples with a very similar stream water  $\delta^2\text{H}$  signature is  
621 increasing towards the stream channel (Garvelmann et al., 2012). In contrast, we found no such  
622 relationship between the distance to stream or TWI and soil isotopic values in the  
623 Vollnkirchener Bach subcatchment over various heightselevations (235–294 m a.s.l.) and  
624 locations. We attributed this to the gentle, ~~low-angle~~ hillslopes and the low subsurface flow  
625 contribution in large parts of the catchment.

626 In our study, the  $\delta$ -values of top soil and precipitation did not differ statistically (Fig. 4), but for  
627 precipitation and subsoil they did. The latter indicates either the influence of evaporation in the  
628 topsoil or the mixing with groundwater in the subsoil. However, a mixing and homogenization  
629 of new and old soil water with depth could not clearly be seen in 0.5 m soil depth, which would  
630 have resulted in a lower standard deviation (Song et al., 2011), but standard deviations of  
631 isotopic signatures in top and subsoil were similar (Table 2). Subsoil isotopic values were  
632 statistically equal to stream water and groundwater (Fig. 4) implying that capillary rise of  
633 groundwater occurred. Overall, the rainfall isotopic signal was not directly transferred through  
634 the soil to the groundwater; even so groundwater head level rose promptly after rainfall events.  
635 This behaviour reflects the differences of celerity and velocity in the catchment's rainfall-runoff  
636 response (McDonnell and Beven, 2014).

637 Soil water  $\delta^2\text{H}$  between top and subsoil showed significant differences ( $p \leq 0.05$ ) under arable  
638 and grassland but not under forested sites (Fig. 5). This could be explained through the  
639 occurrence of vertical preferential flow paths and interconnected macropore flow (Buttle and

640 McDonald, 2002) characteristic for forested soils. Alaoui et al. (2011) showed that macropore  
641 flow with high interaction with the surrounding soil matrix occurred in forest soils, while  
642 macropore flow with low to mixed interaction with the surrounding soil matrix dominates in  
643 grassland soils. Seasonal tilling prevents the establishment of preferential flow paths under  
644 agricultural sites and is regularly done in the Schwingbach catchment, whereas the structure of  
645 forest soils, may remain uninterrupted throughout the entire soil profile for years (in particular  
646 the macropores and biopores) (Alaoui et al., 2011). This is reflected in the bulk density of the  
647 soils in the Schwingbach catchment that increases from forests ( $1.10 \text{ g cm}^{-3}$ ) over grassland  
648 ( $1.25 \text{ g cm}^{-3}$ ) to arable land ( $1.41 \text{ g cm}^{-3}$ ) in the top soil. We infer that reduced hydrological  
649 connection between top and subsoil under arable and grassland led to different isotopic  
650 signatures (Fig. 5).

651 Although, vegetation cover has often shown an impact on soil water isotopes (Gat, 1996), only  
652 few data are available for Central Europe (Darling, 2004). Burger and Seiler (1992) found that  
653 soil water isotopic enrichment under spruce forest in Upper Bavaria was double that beneath  
654 neighbouring arable land but soil isotope values were not comparable to groundwater (Burger  
655 and Seiler, 1992). Gehrels et al. (1998) also detected (though only slightly) heavier isotopic  
656 signatures under forested sites in the Netherlands in comparison to non-forested sites (grassland  
657 and heathland). Contrasting, in southern Germany Brodersen et al. (2000) observed only a  
658 negligible effect of throughfall isotopic signatures (of spruce and beech) on soil water isotopes,  
659 since soil water in the upper layers followed the seasonal trend in the precipitation input and  
660 had a very constant signature in greater depth. For the Schwingbach catchment we conclude  
661 that the observed land use effect in the upper soil column is mainly attributed to different  
662 preservation and transmission of the precipitation input signal. It is most likely not attributable  
663 to distinguished throughfall isotopic signatures, impact of evaporation or interception losses,  
664 since top soil water isotopic signals followed the precipitation input signal under all land use  
665 units.

#### 666 **4.2.2 Seasonal isotope soil profiling**

667 Soil water was enriched in summer due to evaporation during warmer and drier periods. The  
668 depth to which soil water isotopes are significantly affected by evaporation is rarely more than  
669 1–2 m below ground, and often less under temperate climates (Darling, 2004). In contrast,  
670 winter profiles exhibited somewhat greater standard deviations in comparison to summer  
671 isotopic soil profiles, indicative for wetter soils (Fig. 6, lower panels) and shorter residence

672 times (Thomas et al., 2013). Generally, deeper soil water isotope values were relatively constant  
673 through time and space. Similar findings were made by Foerstel et al. (1991) on a sandy soil in  
674 western Germany, McConville et al. (2001) under predominately agriculturally used gley and  
675 till soils in Northern Ireland, and Thomas et al. (2013) in a forested catchment in central  
676 Pennsylvania, USA. Furthermore, Tang and Feng (2001) showed for a sandy loam in New  
677 Hampshire (USA) that the influence of summer precipitation decreased with increasing depth,  
678 and soils at 0.5 m only received water from large storms. In our summer soil profiles under  
679 arable land, precipitation input signals similarly decreased with depth (Fig. 6, upper left panel).  
680 Generally, the replacement of old soil water with new infiltrating water is dependent on the  
681 frequency and intensity of precipitation and the soil texture, structure, wetness, and water  
682 potential of the soil (Li et al., 2007; Tang and Feng, 2001). As a result, the amount of percolating  
683 water decreases with depth and consequently, deeper soil layers have less chance to obtain new  
684 water (Tang and Feng, 2001). In the growing season, the percolation depth is additionally  
685 limited by plants' transpiration (Tang and Feng, 2001). For the Schwingbach catchment we  
686 conclude that the percolation of new soil water is low as no remarkable seasonality in soil  
687 isotopic signatures was obvious at >0.9 m and constant values were observed through space  
688 and time. Although replications over several years are missing, this result indicates a transit  
689 time through the rooting zone (1m) of approximately one year.

### 690 **4.3 Linkages between water cycle components**

691 Stream water isotopic time series of the Vollnkirchener Bach and Schwingbach showed little  
692 deflections through time ~~and, consequently, provided little insight into time and source~~  
693 ~~components connection.~~ Due to the observed isotopic similarities of stream and groundwater,  
694 we conclude that groundwater predominantly feeds baseflow (discharge <10 L·s<sup>-1</sup>). Even  
695 during peak flow occurring in January 2012, December to April or May 2013, rainfall input did  
696 not play a major role for stream water isotopic composition although fast rainfall-runoff  
697 behaviours were observed by Orłowski et al. (2014). The damped groundwater isotopic  
698 signatures seemed to be a mixture of former lighter precipitation events and snowmelt, since  
699 meltwater is known to be depleted in stable isotopes as compared to precipitation or  
700 groundwater (Rohde, 1998) (Figure 3). However, differences in the snow sampling method  
701 (new snow, snow pit layers, meltwater) can affect the isotopic composition (Penna et al., 2014;  
702 Taylor et al., 2001). As groundwater at the observed piezometers in the Vollnkirchener  
703 subcatchment is shallow (Orłowski et al., 2014), the snowmelt signal is allowed to move

704 rapidly through the soil. Pulses of snowmelt water causing a depletion in spring and early  
705 summer was also observed by other studies (Darling, 2004; Kortelainen and Karhu, 2004). We  
706 therefore conclude that groundwater is mainly recharged throughout the winter. During spring  
707 runoff when soils are saturated, temperatures are low, and vegetation is inactive, recharge rates  
708 are generally highest. In contrast, recharge is very low during summer when most precipitation  
709 is transpired back to the atmosphere (Clark and Fritz, 1997a). Similarly, O'Driscoll et al. (2005)  
710 showed that summer precipitation does not significantly contribute to recharge in the Spring  
711 Creek watershed (Pennsylvania, USA) since  $\delta^{18}\text{O}$  values in summer precipitation were enriched  
712 compared to mean annual groundwater composition.

713 Further, Orłowski et al. (2014) showed that influent and effluent conditions (bidirectional water  
714 exchange) occurred simultaneously at different stream sections of the Vollnkirchener Bach  
715 affecting stream and groundwater isotopic compositions, equally. Our network map supported  
716 this assumption (Fig. 9) as surface water sampling points plotted close to groundwater sampling  
717 points (especially to the sampling points under the meadow and along the stream). This was  
718 also underlined by our groundwater model showing flow directions across the Vollnkirchener  
719 Bach. Nevertheless, both stream and groundwater differed significantly from rainfall isotopic  
720 signatures (Table 1). Thus, our catchment showed double water paradox behaviour as per  
721 Kirchner (2003) with fast release of very old water but little variation in tracer concentration.

#### 722 **4.4 Water age dynamics**

723 Our MTT ~~and MRT~~ calculations did not provide ~~meaningful results: a good fit between the~~  
724 ~~observed and calculated data.~~ Just by comparing mean precipitation, stream, and groundwater  
725 isotopic signatures (Table 1), ~~it is obvious one could expect~~ that simple mixing calculations  
726 ~~would~~ not work ~~to derive MTTs~~, i.e. showing predominant groundwater contribution. Same  
727 observations were made by Jin et al. (2010) indicating good hydraulic connectivity between  
728 surface water and shallow groundwater. Just as in the here presented results, Klaus et al. (2015)  
729 had difficulties to apply traditional methods of isotope hydrology (MTT estimation, hydrograph  
730 separation) to their dataset due to the lack of temporal isotopic variation in stream water of a  
731 forested low-mountainous catchment in South Carolina (USA). Furthermore, stable water  
732 isotopes can only be utilised for estimations of younger water (<5 years) (Stewart et al., 2010)  
733 as they are blind to older contributions (Duvert et al., 2016). In our catchment, transit times are  
734 orders of magnitudes longer than the timescale of hydrologic response (prompt discharge of old  
735 water) (McDonnell et al., 2010) and the range used for stable water isotopes.

736 Accurately capturing the transit time of the old water fraction is essential (Duvert et al., 2016)  
737 and could previously only be determined via other tracers such as tritium (e.g. Michel (1992)).  
738 Current studies on mixing assumptions either consider spatial or time-varying MTTs.  
739 Heidebüchel et al. (2012) proposed the concept of the master transit time distribution that  
740 accounts for the temporal variability of MTT. The time-varying transit time concept of Botter  
741 et al. (2011) and van der Velde et al. (2012), was recently reformulated by Harman (2015) so  
742 that the storage selection function became a function of the watershed storage and actual time.  
743 Instead of quantifying time-variant travel times, our model facilitates the estimation of spatially  
744 distributed groundwater ages, which opens up new opportunities to compare groundwater ages  
745 from over a range of scales within catchments. It further gives a deeper understanding of the  
746 groundwater-surface water connection across the landscape than a classical MTT calculation  
747 could provide. Our work complements recent advances in spatially distributed modelling of age  
748 distributions through transient groundwater flows (e.g. Gomez and Wilson, 2013; Woolfenden  
749 and Ginn, 2009). The results of our model reveal a spatially highly heterogeneous age  
750 distribution of groundwater throughout the Vollnkirchener Bach subcatchment (ages of 2 days–  
751 100 years) with oldest water near the stream. Thus, our model provides the opportunity to make  
752 use of stable water isotope information along with climate, land use, and soil type data, in  
753 combination with a digital elevation map to estimate residence times >5 years. If stable water  
754 isotope information is used alone, it is known to cause a truncation of stream residence time  
755 distributions (Stewart et al., 2010). Further, our groundwater model suggests that the main  
756 groundwater flow direction is towards and across the stream and the quantity of flowing water  
757 is highest near the stream (Fig. 10). This further supports the assumption that stream water is  
758 mainly fed by older groundwater. Moreover, the simulation underlines the conclusion that the  
759 groundwater body and stream water are isotopically disconnected from the precipitation cycle,  
760 since only 13% of cells contained water with and age <1 year.

761 However, our semi-conceptual model approach has also some limitations. During model setup  
762 a series of assumptions and simplifications were made to develop a realistic hydrologic model  
763 without a severe loss in performance. Due to the assumption of a constant groundwater recharge  
764 over the course of a year, no seasonality was simulated. Moreover, no spatial differences in soil  
765 properties of the groundwater layer were considered. Further, several parameters such as the  
766 depth of the groundwater body are only rough estimations, while others like evapotranspiration  
767 are based on simulations. Moreover, the groundwater body is highly simplified since e.g.  
768 properties of the simulated aquifer are assumed to be constant over the subcatchment.



769 Nevertheless, as shown by the diverse ages of water in the stream cells and the assumption of  
770 spatially gaining conditions, the model confirms that the stream contains water with different  
771 transit times and supports the assumption that surface and groundwater are isotopically  
772 disconnected from precipitation. Therefore, the stream water does not have a discrete age, but  
773 a distribution of ages due to variable flow paths (Stewart et al., 2010). In future models a more  
774 diverse groundwater body based on small-scale measurements of aquifer parameters should be  
775 implemented. Especially data of saturated hydraulic conductivity with high spatial resolution,  
776 as well as the implementation of a temporal dynamic groundwater recharge could lead to an  
777 enhanced model performance.

## 778 **5 Conclusions**

779 Conducting a stable water isotope study in the Schwingbach catchment helped to identify  
780 relationships between precipitation, stream, soil, and groundwater in a developed (managed)  
781 catchment. The close isotopic link between groundwater and the streams revealed that  
782 groundwater controls streamflow. Moreover, it could be shown that groundwater was  
783 predominately recharged during winter but was decoupled from the annual precipitation cycle.  
784 Even so streamflow and groundwater head levels promptly responded to precipitation inputs,  
785 there was no obvious change in their isotopic composition due to rain events.

786 Nevertheless, the lack of temporal variation in stable isotope time series of stream and  
787 groundwater limited the application of classical methods of isotope hydrology, i.e. transfer  
788 function based MTT estimations. By splitting the flow path into different compartments (upper  
789 and lower vadose zone, groundwater, stream), we were able to determine, where the water age  
790 passes the limit of using stable isotopes for age calculations. This limit is in the lower vadose  
791 zone approximately 1–2 m below ground. To estimate the total transit time to the stream, we  
792 set up a hydrological model calculating spatially distributed groundwater ages and flow  
793 directions in the Vollnkirchener Bach subcatchment. Our model results supported the finding  
794 that the water in the catchment is >5 years (on average 16 years) and that stream water is mainly  
795 fed by groundwater. Our modelling approach was valuable to overcome the limitations of MTT  
796 calculations with traditional methods and/or models. Further, our dual isotope study in  
797 combination with the hydrological model approach enabled the determination of connection  
798 and disconnection between different water cycle components.

## 799 **Acknowledgements**

800 The first author acknowledges financial support by the Friedrich-Ebert-Stiftung (Bonn, DE).  
801 Furthermore, this work was supported by the Deutsche Forschungsgemeinschaft under Grant  
802 BR2238/10-1. Thanks also to the student assistants, BSc. and MSc. students for their help  
803 during field sampling campaigns and hydrological modelling. In this regard we like to  
804 acknowledge especially Julia Mechsner, Julia Klöber, and Judith Henkel. We thank Dr.  
805 Christine Stumpp from the Helmholtz Zentrum München for providing us the isotope data from  
806 GNIP station Koblenz.  
807

808 **References**

- 809 ~~Alaoui, A., Caduff, U., Gerke, H. H. and Weingartner, R.: Preferential Flow Effects on~~  
810 ~~Infiltration and Runoff in Grassland and Forest Soils, *Vadose Zone J.*, 10(1), 367,~~  
811 ~~doi:10.2136/vzj2010.0076, 2011.~~
- 812 ~~Allan, J. D.: Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems,~~  
813 ~~*Annu. Rev. Ecol. Evol. Syst.*, 35, 257–284, doi:10.1146/annurev.ecolsys.35.120202.110122,~~  
814 ~~2004.~~
- 815 ~~Barnes, C. J. and Allison, G. B.: Tracing of water movement in the unsaturated zone using~~  
816 ~~stable isotopes of hydrogen and oxygen, *J. Hydrol.*, 100(1–3), 143–176, doi:10.1016/0022-~~  
817 ~~1694(88)90184-9, 1988.~~
- 818 ~~Barthold, F. K., Tyralla, C., Schneider, K., Vaché, K. B., Frede, H. G. and Breuer, L.: How~~  
819 ~~many tracers do we need for end member mixing analysis (EMMA)? A sensitivity analysis,~~  
820 ~~*Water Resour. Res.*, 47(8), doi:10.1029/2011WR010604, 2011.~~
- 821 ~~Blasch, K. W. and Bryson, J. R.: Distinguishing Sources of Ground Water Recharge by Using~~  
822  ~~$\delta^2\text{H}$  and  $\delta^{18}\text{O}$ , *Ground Water*, 45(3), 294–308, doi:10.1111/j.1745-6584.2006.00289.x, 2007.~~
- 823 ~~Botter, G., Bertuzzo, E. and Rinaldo, A.: Transport in the hydrologic response: Travel time~~  
824 ~~distributions, soil moisture dynamics, and the old water paradox, *Water Resour. Res.*, 46(3),~~  
825 ~~W03514, doi:10.1029/2009WR008371, 2010.~~
- 826 ~~Botter, G., Bertuzzo, E. and Rinaldo, A.: Catchment residence and travel time distributions:~~  
827 ~~The master equation, *Geophys. Res. Lett.*, 38(11), L11403, doi:10.1029/2011GL047666, 2011.~~
- 828 ~~Brodersen, C., Pohl, S., Lindenlaub, M., Leibundgut, C. and Wilpert, K. v: Influence of~~  
829 ~~vegetation structure on isotope content of throughfall and soil water, *Hydrol. Process.*, 14(8),~~  
830 ~~1439–1448, doi:10.1002/1099-1085(20000615)14:8<1439::AID-HYP985>3.0.CO;2-3, 2000.~~
- 831 ~~Burger, H. M. and Seiler, K. P.: Evaporation from soil water under humid climate conditions~~  
832 ~~and its impact on deuterium and  $^{18}\text{O}$  concentrations in groundwater, edited by International~~  
833 ~~Atomic Energy Agency, International Atomic Energy Agency, Vienna, Austria., 1992.~~
- 834 ~~Buttle, J. M.: Isotope hydrograph separations and rapid delivery of pre-event water from~~  
835 ~~drainage basins, *Prog. Phys. Geogr.*, 18(1), 16–41, doi:10.1177/030913339401800102, 1994.~~
- 836 ~~Buttle, J. M.: Isotope Hydrograph Separation of Runoff Sources, in *Encyclopedia of*~~  
837 ~~*Hydrological Sciences*, edited by M. G. Anderson, p. 10:116, John Wiley & Sons, Ltd,~~  
838 ~~Chichester, Great Britain., 2006.~~
- 839 ~~Buttle, J. M. and McDonald, D. J.: Coupled vertical and lateral preferential flow on a forested~~  
840 ~~slope, *Water Resour. Res.*, 38(5), 18–1, doi:10.1029/2001WR000773, 2002.~~
- 841 ~~Clark, I. D. and Fritz, P.: Groundwater, in *Environmental Isotopes in Hydrogeology*, p. 80, CRC~~  
842 ~~Press, Florida, FL, USA., 1997a.~~
- 843 ~~Clark, I. D. and Fritz, P.: Methods for Field Sampling, in *Environmental Isotopes in*~~  
844 ~~*Hydrogeology*, p. 283, CRC Press, Florida, FL, USA., 1997b.~~

845 Clark, I. D. and Fritz, P.: The Environmental Isotopes, in Environmental Isotopes in  
846 Hydrogeology, pp. 2–34, CRC Press, Florida, FL, USA., 1997c.

847 Cooper, L. W.: Isotopic Fractionation in Snow Cover, in Isotope Tracers in Catchment  
848 Hydrology, edited by C. Kendall and J. J. McDonnell, pp. 119–136, Elsevier, Amsterdam,  
849 Netherlands., 1998.

850 Craig, H.: Isotopic Variations in Meteoric Waters, *Science*, 133(3465), 1702–1703,  
851 doi:10.1126/science.133.3465.1702, 1961a.

852 Craig, H.: Standard for reporting concentrations of deuterium and oxygen-18 in natural waters,  
853 *Science*, 133(3467), 1833, 1961b.

854 Csardi, G. and Nepusz, T.: The igraph software package for complex network research,  
855 *Complex Systems* 1695. [online] Available from: <http://igraph.sf.net> (Accessed 24 December  
856 2015), 2006.

857 Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16(4), 436–468, doi:10.1111/j.2153-  
858 3490.1964.tb00181.x, 1964.

859 Darling, W. G.: Hydrological factors in the interpretation of stable isotopic proxy data present  
860 and past: a European perspective, *Quat. Sci. Rev.*, 23(7–8), 743–770,  
861 doi:10.1016/j.quascirev.2003.06.016, 2004.

862 Duvert, C., Stewart, M. K., Cendón, D. I. and Raiber, M.: Time series of tritium, stable isotopes  
863 and chloride reveal short-term variations in groundwater contribution to a stream, *Hydrol. Earth  
864 Syst. Sci.*, 20(1), 257–277, doi:10.5194/hess-20-257-2016, 2016.

865 DWD: Deutscher Wetterdienst—Wetter und Klima, Bundesministerium für Verkehr und  
866 digitale Infrastruktur, [online] Available from: <http://dwd.de/> (Accessed 17 February 2014),  
867 2014.

868 Foerstel, H., Frinken, J., Huetzen, H., Lembrich, D. and Puetz, T.: Application of H<sub>2</sub><sup>18</sup>O as a  
869 tracer of water flow in soil, International Atomic Energy Agency, Vienna, Austria., 1991.

870 Garvelmann, J., Kuells, C. and Weiler, M.: A porewater-based stable isotope approach for the  
871 investigation of subsurface hydrological processes, *Hydrol. Earth Syst. Sci.*, 16(2), 631–640,  
872 doi:10.5194/hess-16-631-2012, 2012.

873 Gat, J. R.: Oxygen and Hydrogen Isotopes in the Hydrologic Cycle, *Annu. Rev. Earth Planet.  
874 Sci.*, 24(1), 225–262, doi:10.1146/annurev.earth.24.1.225, 1996.

875 Gat, J., R., Mook, W. G. and Meijer, H. A. J.: Environmental isotopes in the hydrological cycle:  
876 Principles and Applications, edited by W. G. Mook, International Hydrological Programme;  
877 United Nations Educational, Scientific and Cultural Organization and International Atomic  
878 Energy Agency, Paris, France., 2001.

879 Gehrels, J. C., Peeters, J. E. M., de Vries, J. J. and Dekkers, M.: The mechanism of soil water  
880 movement as inferred from <sup>18</sup>O stable isotope studies, *Hydrol. Sci. J.*, 43(4), 579–594,  
881 doi:10.1080/02626669809492154, 1998.

882 Genereux, D. P. and Hooper, R. P.: Oxygen and Hydrogen Isotopes in Rainfall–Runoff Studies,  
883 in *Isotope Tracers in Catchment Hydrology*, edited by C. Kendall and J. J. McDonnell, pp. 319–  
884 346, Elsevier, Amsterdam, Netherlands., 1998.

885 Gomez, J. D. and Wilson, J. L.: Age distributions and dynamically changing hydrologic  
886 systems: Exploring topography driven flow, *Water Resour. Res.*, 49(3), 1503–1522,  
887 doi:10.1002/wrer.20127, 2013.

888 Gonfiantini, R., Fröhlich, K., Araguás-Araguás, L. and Rozanski, K.: Isotopes in Groundwater  
889 Hydrology, in *Isotope Tracers in Catchment Hydrology*, edited by C. Kendall and J. J.  
890 McDonnell, pp. 203–246, Elsevier, Amsterdam, Netherlands., 1998.

891 Gordon, L. J., Finlayson, C. M. and Falkenmark, M.: Managing water in agriculture for food  
892 production and other ecosystem services, *Agric. Water Manag.*, 97(4), 512–519,  
893 doi:10.1016/j.agwat.2009.03.017, 2010.

894 Harman, C. J.: Time-variable transit time distributions and transport: Theory and application to  
895 storage-dependent transport of chloride in a watershed, *Water Resour. Res.*, 51(1), 1–30,  
896 doi:10.1002/2014WR015707, 2015.

897 Heidebüchel, I., Troch, P. A., Lyon, S. W. and Weiler, M.: The master transit time distribution  
898 of variable flow systems, *Water Resour. Res.*, 48(6), W06520, doi:10.1029/2011WR011293,  
899 2012.

900 Hindmarsh, A. C., Brown, P. N., Grant, K. E., Lee, S. L., Serban, R., Shumaker, D. E. and  
901 Woodward, C. S.: SUNDIALS: Suite of Nonlinear and Differential/Algebraic Equation  
902 Solvers, *ACM Trans. Math. Softw.*, 31(3), 363–396, doi:10.1145/1089014.1089020, 2005.

903 Hollander, M., Wolfe, D. A. and Chickén, E.: *Nonparametric Statistical Methods*, John Wiley  
904 & Sons, New York, NY, USA., 2013.

905 Hrachowitz, M., Benettin, P., van Breukelen, B. M., Fovet, O., Howden, N. J. K., Ruiz, L., van  
906 der Velde, Y. and Wade, A. J.: Transit times—the link between hydrology and water quality at  
907 the catchment scale, *Wiley Interdiscip. Rev. Water*, doi:10.1002/wat2.1155, 2016.

908 IAEA: International Atomic Energy Agency: Water Resources Programme—Global Network  
909 of Isotopes in Precipitation, [online] Available from: [http://www-](http://www-naweb.iaea.org/naweb/ih/IHS_resources_gnip.html)  
910 [naweb.iaea.org/naweb/ih/IHS\\_resources\\_gnip.html](http://www-naweb.iaea.org/naweb/ih/IHS_resources_gnip.html) (Accessed 11 August 2014), 2014.

911 Jin, L., Siegel, D. I., Lautz, L. K., Mitchell, M. J., Dahms, D. E. and Mayer, B.: Calcite  
912 precipitation driven by the common ion effect during groundwater surface water mixing: A  
913 potentially common process in streams with geologic settings containing gypsum, *Geol. Soc.  
914 Am. Bull.*, 122(7/8), B30011.1, doi:10.1130/B30011.1, 2010.

915 Jin, L., Siegel, D. I., Lautz, L. K. and Lu, Z.: Identifying streamflow sources during spring  
916 snowmelt using water chemistry and isotopic composition in semi-arid mountain streams, *J.  
917 Hydrol.*, 470–471, 289–301, doi:10.1016/j.jhydrol.2012.09.009, 2012.

918 Jouzel, J., Alley, R. B., Cuffey, K. M., Dansgaard, W., Grootes, P., Hoffmann, G., Johnsen, S.  
919 J., Koster, R. D., Peel, D., Shuman, C. A., Stievenard, M., Stuiver, M. and White, J.: Validity

920 of the temperature reconstruction from water isotopes in ice cores, *J. Geophys. Res.*, 102,  
921 26471–26487, doi:10.1029/97JC01283, 1997.

922 Kendall, C. and Caldwell, E. A.: Fundamentals of Isotope Geochemistry, in *Isotope Tracers in*  
923 *Catchment Hydrology*, edited by C. Kendall and J. J. McDonnell, pp. 51–86, Elsevier,  
924 Amsterdam, Netherlands., 1998.

925 Kirchner, J. W.: A double paradox in catchment hydrology and geochemistry, *Hydrol. Process.*,  
926 17(4), 871–874, doi:10.1002/hyp.5108, 2003.

927 Klaus, J., McDonnell, J. J., Jackson, C. R., Du, E. and Griffiths, N. A.: Where does streamwater  
928 come from in low relief forested watersheds? A dual isotope approach, *Hydrol. Earth Syst.*  
929 *Sci.*, 19(1), 125–135, 2015.

930 Koeniger, P., Leibundgut, C. and Stiehler, W.: Spatial and temporal characterisation of stable  
931 isotopes in river water as indicators of groundwater contribution and confirmation of modelling  
932 results; a study of the Weser river, Germany, *Isotopes Environ. Health Stud.*, 45(4), 289–302,  
933 doi:10.1080/10256010903356953, 2009.

934 Kolaczyk, E. D.: *Statistical Analysis of Network Data with R*, Springer Science & Business  
935 Media, New York, NY, USA., 2014.

936 Kortelainen, N. M. and Karhu, J. A.: Regional and seasonal trends in the oxygen and hydrogen  
937 isotope ratios of Finnish groundwaters: a key for mean annual precipitation, *J. Hydrol.*, 285(1–  
938 4), 143–157, doi:10.1016/j.jhydrol.2003.08.014, 2004.

939 Kraft, P., Vaché, K. B., Frede, H. G. and Breuer, L.: CMF: A Hydrological Programming  
940 Language Extension For Integrated Catchment Models, *Environ. Model. Softw.*, 26(6), 828–  
941 830, doi:10.1016/j.envsoft.2010.12.009, 2011.

942 Lai, C. T. and Ehleringer, J. R.: Deuterium excess reveals diurnal sources of water vapor in  
943 forest air, *Oecologia*, 165(1), 213–223, doi:10.1007/s00442-010-1721-2, 2011.

944 Lauer, F., Frede, H. G. and Breuer, L.: Uncertainty assessment of quantifying spatially  
945 concentrated groundwater discharge to small streams by distributed temperature sensing, *Water*  
946 *Resour. Res.*, 49(1), 400–407, doi:10.1029/2012WR012537, 2013.

947 LGR: Los Gatos Research, Greenhouse Gas, isotope and trace gas analyzers, [online] Available  
948 from: <http://www.lgrinc.com/> (Accessed 5 February 2013), 2013.

949 Li, F., Song, X., Tang, C., Liu, C., Yu, J. and Zhang, W.: Tracing infiltration and recharge using  
950 stable isotope in Taihang Mt., North China, *Environ. Geol.*, 53(3), 687–696,  
951 doi:10.1007/s00254-007-0683-0, 2007.

952 Maloszewski, P. and Zuber, A.: Manual on lumped parameter models used for the interpretation  
953 of environmental tracer data in groundwaters, in *Use of isotopes for analyses of flow and*  
954 *transport dynamics in groundwater systems*, p. 50, International Atomic Energy Agency,  
955 Vienna, Austria., 2002.

956 McConville, C., Kalin, R. m., Johnston, H. and McNeill, G. w.: Evaluation of Recharge in a  
957 Small Temperate Catchment Using Natural and Applied  $\delta^{18}\text{O}$  Profiles in the Unsaturated Zone,  
958 *Ground Water*, 39(4), 616–623, doi:10.1111/j.1745-6584.2001.tb02349.x, 2001.

959 McDonnell, J. J. and Beven, K.: Debates—The future of hydrological sciences: A (common)  
960 path forward? A call to action aimed at understanding velocities, celerities and residence time  
961 distributions of the headwater hydrograph, *Water Resour. Res.*, 50(6), 5342–5350,  
962 doi:10.1002/2013WR015141, 2014.

963 McDonnell, J. J., Sivapalan, M., Vaché, K., Dunn, S., Grant, G., Haggerty, R., Hinz, C., Hooper,  
964 R., Kirchner, J., Roderick, M. L. and others: Moving beyond heterogeneity and process  
965 complexity: a new vision for watershed hydrology, *Water Resour. Res.*, 43(7), W07301, 2007.

966 McDonnell, J. J., McGuire, K., Aggarwal, P., Beven, K. J., Biondi, D., Destouni, G., Dunn, S.,  
967 James, A., Kirchner, J., Kraft, P., Lyon, S., Maloszewski, P., Newman, B., Pfister, L., Rinaldo,  
968 A., Rodhe, A., Sayama, T., Seibert, J., Solomon, K., Soulsby, C., Stewart, M., Tetzlaff, D.,  
969 Tobin, C., Troch, P., Weiler, M., Western, A., Worman, A. and Wrede, S.: How old is  
970 streamwater? Open questions in catchment transit time conceptualization, modelling and  
971 analysis, *Hydrol. Process.*, 24(12), 1745–1754, 2010.

972 McGuire, K. J. and McDonnell, J. J.: A review and evaluation of catchment transit time  
973 modeling, *J. Hydrol.*, 330(3), 543–563, 2006.

974 Michel, R. L.: Residence times in river basins as determined by analysis of long term tritium  
975 records, *J. Hydrol.*, 130(1), 367–378, doi:10.1016/0022-1694(92)90117-E, 1992.

976 Mook, W. G., Groeneveld, D. J., Brouwn, A. E. and Ganswijk, A. J. van: Analysis of a run-off  
977 hydrograph by means of natural  $^{18}\text{O}$ , in *Isotope techniques in groundwater hydrology*, vol. 1,  
978 pp. 159–169, International Atomic Energy Agency, Vienna, Austria., 1974.

979 Neal, C. and Rosier, P. T. W.: Chemical studies of chloride and stable oxygen isotopes in two  
980 conifer afforested and moorland sites in the British uplands, *J. Hydrol.*, 115(1–4), 269–283,  
981 doi:10.1016/0022-1694(90)90209-G, 1990.

982 O’Driscoll, M. A., DeWalle, D. R., McGuire, K. J. and Gburek, W. J.: Seasonal  $^{18}\text{O}$  variations  
983 and groundwater recharge for three landscape types in central Pennsylvania, USA, *J. Hydrol.*,  
984 303(1–4), 108–124, doi:10.1016/j.jhydrol.2004.08.020, 2005.

985 Orłowski, N., Frede, H. G., Brüggemann, N. and Breuer, L.: Validation and application of a  
986 cryogenic vacuum extraction system for soil and plant water extraction for isotope analysis, *J.*  
987 *Sens. Sens. Syst.*, 2(2), 179–193, doi:10.5194/jsss-2-179-2013, 2013.

988 Orłowski, N., Lauer, F., Kraft, P., Frede, H. G. and Breuer, L.: Linking Spatial Patterns of  
989 Groundwater Table Dynamics and Streamflow Generation Processes in a Small Developed  
990 Catchment, *Water*, 6(10), 3085–3117, doi:10.3390/w6103085, 2014.

991 Penna, D., Ahmad, M., Birks, S. J., Bouchaou, L., Brenčič, M., Butt, S., Holko, L., Jeelani, G.,  
992 Martínez, D. E., Melikadze, G., Shanley, J. B., Sokratov, S. A., Stadnyk, T., Sugimoto, A. and  
993 Vreča, P.: A new method of snowmelt sampling for water stable isotopes, *Hydrol. Process.*,  
994 28(22), 5637–5644, doi:10.1002/hyp.10273, 2014.



- 995 Perry, C. and Taylor, K.: *Environmental Sedimentology*, p. 36, Blackwell Publishing, Oxford,  
996 OX, UK, 2009.
- 997 Pierce, S. C., Kröger, R. and Pezeshki, R.: *Managing Artificially Drained Low Gradient*  
998 *Agricultural Headwaters for Enhanced Ecosystem Functions*, *Biology*, 1(3), 794–856,  
999 doi:10.3390/biology1030794, 2012.
- 1000 Qu, Y. and Duffy, C. J.: *A semidiscrete finite volume formulation for multiprocess watershed*  
1001 *simulation*, *Water Resour. Res.*, 43(8), W08419, doi:10.1029/2006WR005752, 2007.
- 1002 Rinaldo, A., Benettin, P., Harman, C. J., Hrachowitz, M., McGuire, K. J., van der Velde, Y.,  
1003 Bertuzzo, E. and Botter, G.: *Storage selection functions: A coherent framework for quantifying*  
1004 *how catchments store and release water and solutes*, *Water Resour. Res.*, 51(6), 4840–4847,  
1005 doi:10.1002/2015WR017273, 2015.
- 1006 Rohde, A.: *Snowmelt-Dominated Systems*, in *Isotope Tracers in Catchment Hydrology*, edited  
1007 by C. Kendall and J. J. McDonnell, pp. 391–433, Elsevier, Amsterdam, Netherlands., 1998.
- 1008 Rozanski, K., Sonntag, C. and Münnich, K. O.: *Factors controlling stable isotope composition*  
1009 *of European precipitation*, *Tellus*, 34(2), 142–150, doi:10.1111/j.2153-3490.1982.tb01801.x,  
1010 1982.
- 1011 Rozanski, K., Froehlich, K. and Mook, W. G.: in *Environmental isotopes in the hydrological*  
1012 *cycle: Principles and Applications*, vol. 3, International Hydrological Programme; United  
1013 Nations Educational, Scientific and Cultural Organization and International Atomic Energy  
1014 Agency, Paris, Vienna., 2001.
- 1015 Schultz, N. M., Griffis, T. J., Lee, X. and Baker, J. M.: *Identification and correction of spectral*  
1016 *contamination in  $^2\text{H}/^1\text{H}$  and  $^{18}\text{O}/^{16}\text{O}$  measured in leaf, stem, and soil water*, *Rapid Commun.*  
1017 *Mass Spectrom.*, 25(21), 3360–3368, doi:10.1002/rem.5236, 2011.
- 1018 Shuttleworth, W. J. and Wallace, J. S.: *Evaporation from sparse crops—an energy combination*  
1019 *theory*, *Q. J. R. Meteorol. Soc.*, 111(469), 839–855, doi:10.1002/qj.49711146910, 1985.
- 1020 Sklash, M. G.: *Environmental isotope studies of storm and snowmelt runoff generation*, in  
1021 *Process studies in hillslope hydrology*, edited by M. G. Anderson, pp. 410–435, Wiley, New  
1022 York, NY, USA., 1990.
- 1023 Sklash, M. G. and Farvolden, R. N.: *The Role Of Groundwater In Storm Runoff*, in  
1024 *Developments in Water Science*, vol. 12, edited by W. B. and D. A. Stephenson, pp. 45–65,  
1025 Elsevier, Amsterdam, Netherlands., 1979.
- 1026 Song, X., Wang, P., Yu, J., Liu, X., Liu, J. and Yuan, R.: *Relationships between precipitation,*  
1027 *soil water and groundwater at Chongling catchment with the typical vegetation cover in the*  
1028 *Taihang mountainous region, China*, *Environ. Earth Sci.*, 62(4), 787–796, doi:10.1007/s12665-  
1029 010-0566-7, 2011.
- 1030 Stewart, M. K., Morgenstern, U. and McDonnell, J. J.: *Truncation of stream residence time:*  
1031 *how the use of stable isotopes has skewed our concept of streamwater age and origin*, *Hydrol.*  
1032 *Process.*, 24(12), 1646–1659, doi:10.1002/hyp.7576, 2010.



1033 Stumpp, C., Klaus, J. and Stichler, W.: Analysis of long-term stable isotopic composition in  
1034 German precipitation, *J. Hydrol.*, 517, 351–361, doi:10.1016/j.jhydrol.2014.05.034, 2014.

1035 Tang, K. and Feng, X.: The effect of soil hydrology on the oxygen and hydrogen isotopic  
1036 compositions of plants' source water, *Earth Planet. Sci. Lett.*, 185(3–4), 355–367, 2001.

1037 Taylor, S., Feng, X., Kirchner, J. W., Osterhuber, R., Klaue, B. and Renshaw, C. E.: Isotopic  
1038 evolution of a seasonal snowpack and its melt, *Water Resour. Res.*, 37(3), 759–769,  
1039 doi:10.1029/2000WR900341, 2001.

1040 Thomas, E. M., Lin, H., Duffy, C. J., Sullivan, P. L., Holmes, G. H., Brantley, S. L. and Jin, L.:  
1041 Spatiotemporal Patterns of Water Stable Isotope Compositions at the Shale Hills Critical Zone  
1042 Observatory: Linkages to Subsurface Hydrologic Processes, *Vadose Zone J.*, 12(4), 0,  
1043 doi:10.2136/vzj2013.01.0029, 2013.

1044 Timbe, E., Windhorst, D., Crespo, P., Frede, H.-G., Feyen, J. and Breuer, L.: Understanding  
1045 uncertainties when inferring mean transit times of water trough tracer-based lumped-parameter  
1046 models in Andean tropical montane cloud forest catchments, *Hydrol. Earth Syst. Sci.*, 18(4),  
1047 1503–1523, doi:10.5194/hess-18-1503-2014, 2014.

1048 van der Velde, Y., Torfs, P. J. J. F., van der Zee, S. E. a. T. M. and Uijlenhoet, R.: Quantifying  
1049 catchment-scale mixing and its effect on time-varying travel time distributions, *Water Resour.*  
1050 *Res.*, 48(6), W06536, doi:10.1029/2011WR011310, 2012.

1051 Wang, X. F. and Yakir, D.: Using stable isotopes of water in evapotranspiration studies, *Hydrol.*  
1052 *Process.*, 14(8), 1407–1421, doi:10.1002/1099-1085(20000615)14:8<1407::AID-  
1053 HYP992>3.0.CO;2-K, 2000.

1054 Windhorst, D., Waltz, T., Timbe, E., Frede, H.-G. and Breuer, L.: Impact of elevation and  
1055 weather patterns on the isotopic composition of precipitation in a tropical montane rainforest,  
1056 *Hydrol. Earth Syst. Sci.*, 17(1), 409–419, doi:10.5194/hess-17-409-2013, 2013.

1057 Windhorst, D., Kraft, P., Timbe, E., Frede, H.-G. and Breuer, L.: Stable water isotope tracing  
1058 through hydrological models for disentangling runoff generation processes at the hillslope  
1059 scale, *Hydrol. Earth Syst. Sci.*, 18(10), 4113–4127, doi:10.5194/hess-18-4113-2014, 2014.

1060 Woolfenden, L. R. and Ginn, T. R.: Modeled Ground Water Age Distributions, *Ground Water*,  
1061 47(4), 547–557, doi:10.1111/j.1745-6584.2008.00550.x, 2009.

1062 Wu, J., Ding, Y., Ye, B., Yang, Q., Hou, D. and Xue, L.: Stable isotopes in precipitation in  
1063 Xilin River Basin, northern China and their implications, *Chin. Geogr. Sci.*, 22(5), 531–540,  
1064 doi:10.1007/s11769-012-0543-z, 2012.

1065 Xia, Y.: Optimization and uncertainty estimates of WMO regression models for the systematic  
1066 bias adjustment of NLDAS precipitation in the United States, *J. Geophys. Res. Atmospheres*,  
1067 111(D8), D08102, doi:10.1029/2005JD006188, 2006.

1068 Yurtsever, Y.: Worldwide survey of stable isotopes in precipitation, International Atomic  
1069 Energy Agency, Vienna, Austria., 1975.

1070 [Zhao, L., Xiao, H., Zhou, J., Wang, L., Cheng, G., Zhou, M., Yin, L. and McCabe, M. F.: Detailed assessment of isotope ratio infrared spectroscopy and isotope ratio mass spectrometry for the stable isotope analysis of plant and soil waters, \*Rapid Commun. Mass Spectrom.\*, 25\(20\), 3071–3082, doi:10.1002/rcm.5204, 2011.](#)

1071

1072

1073

1074 [Alaoui, A., Caduff, U., Gerke, H. H. and Weingartner, R.: Preferential Flow Effects on Infiltration and Runoff in Grassland and Forest Soils, \*Vadose Zone J.\*, 10\(1\), 367, doi:10.2136/vzj2010.0076, 2011.](#)

1075

1076

1077 [Allan, J. D.: Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems, \*Annu. Rev. Ecol. Evol. Syst.\*, 35, 257–284, doi:10.1146/annurev.ecolsys.35.120202.110122, 2004.](#)

1078

1079

1080 [Barnes, C. J. and Allison, G. B.: Tracing of water movement in the unsaturated zone using stable isotopes of hydrogen and oxygen, \*J. Hydrol.\*, 100\(1–3\), 143–176, doi:10.1016/0022-1694\(88\)90184-9, 1988.](#)

1081

1082

1083 [Barthold, F. K., Tyralla, C., Schneider, K., Vaché, K. B., Frede, H.-G. and Breuer, L.: How many tracers do we need for end member mixing analysis \(EMMA\)? A sensitivity analysis, \*Water Resour. Res.\*, 47\(8\), doi:10.1029/2011WR010604, 2011.](#)

1084

1085

1086 [Blasch, K. W. and Bryson, J. R.: Distinguishing Sources of Ground Water Recharge by Using  \$\delta^2\text{H}\$  and  \$\delta^{18}\text{O}\$ , \*Ground Water\*, 45\(3\), 294–308, doi:10.1111/j.1745-6584.2006.00289.x, 2007.](#)

1087

1088 [Botter, G., Bertuzzo, E. and Rinaldo, A.: Transport in the hydrologic response: Travel time distributions, soil moisture dynamics, and the old water paradox, \*Water Resour. Res.\*, 46\(3\), W03514, doi:10.1029/2009WR008371, 2010.](#)

1089

1090

1091 [Botter, G., Bertuzzo, E. and Rinaldo, A.: Catchment residence and travel time distributions: The master equation, \*Geophys. Res. Lett.\*, 38\(11\), L11403, doi:10.1029/2011GL047666, 2011.](#)

1092

1093 [Brodersen, C., Pohl, S., Lindenlaub, M., Leibundgut, C. and Wilpert, K. v: Influence of vegetation structure on isotope content of throughfall and soil water, \*Hydrol. Process.\*, 14\(8\), 1439–1448, doi:10.1002/1099-1085\(20000615\)14:8<1439::AID-HYP985>3.0.CO;2-3, 2000.](#)

1094

1095

1096 [Burger, H. M. and Seiler, K. P.: Evaporation from soil water under humid climate conditions and its impact on deuterium and  \$^{18}\text{O}\$  concentrations in groundwater, edited by International Atomic Energy Agency, International Atomic Energy Agency, Vienna, Austria., 1992.](#)

1097

1098

1099 [Buttle, J. M.: Isotope hydrograph separations and rapid delivery of pre-event water from drainage basins, \*Prog. Phys. Geogr.\*, 18\(1\), 16–41, doi:10.1177/030913339401800102, 1994.](#)

1100

1101 [Buttle, J. M.: Isotope Hydrograph Separation of Runoff Sources, in \*Encyclopedia of Hydrological Sciences\*, edited by M. G. Anderson, p. 10:116, John Wiley & Sons, Ltd, Chichester, Great Britain., 2006.](#)

1102

1103

1104 [Buttle, J. M. and McDonald, D. J.: Coupled vertical and lateral preferential flow on a forested slope, \*Water Resour. Res.\*, 38\(5\), 18–1, doi:10.1029/2001WR000773, 2002.](#)

1105

1106 [Clark, I. D. and Fritz, P.: Groundwater, in \*Environmental Isotopes in Hydrogeology\*, p. 80, CRC Press, Florida, FL, USA., 1997a.](#)

1107

1108 [Clark, I. D. and Fritz, P.: Methods for Field Sampling, in \*Environmental Isotopes in Hydrogeology\*, p. 283, CRC Press, Florida, FL, USA., 1997b.](#)

1109

1110 [Clark, I. D. and Fritz, P.: The Environmental Isotopes, in \*Environmental Isotopes in Hydrogeology\*, pp. 2–34, CRC Press, Florida, FL, USA., 1997c.](#)

1111

- 1112 Cooper, L. W.: Isotopic Fractionation in Snow Cover, in Isotope Tracers in Catchment  
1113 Hydrology, edited by C. Kendall and J. J. McDonnell, pp. 119–136, Elsevier, Amsterdam,  
1114 Netherlands., 1998.
- 1115 Craig, H.: Isotopic Variations in Meteoric Waters, Science, 133(3465), 1702–1703,  
1116 doi:10.1126/science.133.3465.1702, 1961a.
- 1117 Craig, H.: Standard for reporting concentrations of deuterium and oxygen-18 in natural waters,  
1118 Science, 133(3467), 1833, 1961b.
- 1119 Csardi, G. and Nepusz, T.: The igraph software package for complex network research,  
1120 Complex Systems 1695. [online] Available from: <http://igraph.sf.net> (Accessed 24 December  
1121 2015), 2006.
- 1122 Dansgaard, W.: Stable isotopes in precipitation, Tellus, 16(4), 436–468, doi:10.1111/j.2153-  
1123 3490.1964.tb00181.x, 1964.
- 1124 Darling, W. G.: Hydrological factors in the interpretation of stable isotopic proxy data present  
1125 and past: a European perspective, Quat. Sci. Rev., 23(7–8), 743–770,  
1126 doi:10.1016/j.quascirev.2003.06.016, 2004.
- 1127 Duvert, C., Stewart, M. K., Cendón, D. I. and Raiber, M.: Time series of tritium, stable isotopes  
1128 and chloride reveal short-term variations in groundwater contribution to a stream, Hydrol. Earth  
1129 Syst. Sci., 20(1), 257–277, doi:10.5194/hess-20-257-2016, 2016.
- 1130 DWD: Deutscher Wetterdienst - Wetter und Klima, Bundesministerium für Verkehr und  
1131 digitale Infrastruktur, [online] Available from: <http://dwd.de/> (Accessed 17 February 2014),  
1132 2014.
- 1133 Foerstel, H., Frinken, J., Huetzen, H., Lembrich, D. and Puetz, T.: Application of H<sub>2</sub><sup>18</sup>O as a  
1134 tracer of water flow in soil, International Atomic Energy Agency, Vienna, Austria., 1991.
- 1135 Garvelmann, J., Kuells, C. and Weiler, M.: A porewater-based stable isotope approach for the  
1136 investigation of subsurface hydrological processes, Hydrol. Earth Syst. Sci., 16(2), 631–640,  
1137 doi:10.5194/hess-16-631-2012, 2012.
- 1138 Gat, J. R.: Oxygen and Hydrogen Isotopes in the Hydrologic Cycle, Annu. Rev. Earth Planet.  
1139 Sci., 24(1), 225–262, doi:10.1146/annurev.earth.24.1.225, 1996.
- 1140 Gat, J., R., Mook, W. G. and Meijer, H. A. J.: Environmental isotopes in the hydrological cycle:  
1141 Principles and Applications, edited by W. G. Mook, International Hydrological Programme;  
1142 United Nations Educational, Scientific and Cultural Organization and International Atomic  
1143 Energy Agency, Paris, France., 2001.
- 1144 Gehrels, J. C., Peeters, J. E. M., de Vries, J. J. and Dekkers, M.: The mechanism of soil water  
1145 movement as inferred from <sup>18</sup>O stable isotope studies, Hydrol. Sci. J., 43(4), 579–594,  
1146 doi:10.1080/02626669809492154, 1998.
- 1147 Genereux, D. P. and Hooper, R. P.: Oxygen and Hydrogen Isotopes in Rainfall-Runoff Studies,  
1148 in Isotope Tracers in Catchment Hydrology, edited by C. Kendall and J. J. McDonnell, pp. 319–  
1149 346, Elsevier, Amsterdam, Netherlands., 1998.
- 1150 Gomez, J. D. and Wilson, J. L.: Age distributions and dynamically changing hydrologic  
1151 systems: Exploring topography-driven flow, Water Resour. Res., 49(3), 1503–1522,  
1152 doi:10.1002/wrcr.20127, 2013.
- 1153 Gonfiantini, R., Fröhlich, K., Araguás-Araguás, L. and Rozanski, K.: Isotopes in Groundwater  
1154 Hydrology, in Isotope Tracers in Catchment Hydrology, edited by C. Kendall and J. J.

- 1155 McDonnell, pp. 203–246, Elsevier, Amsterdam, Netherlands., 1998.
- 1156 Gordon, L. J., Finlayson, C. M. and Falkenmark, M.: Managing water in agriculture for food  
1157 production and other ecosystem services, Agric. Water Manag., 97(4), 512–519,  
1158 doi:10.1016/j.agwat.2009.03.017, 2010.
- 1159 Harman, C. J.: Time-variable transit time distributions and transport: Theory and application to  
1160 storage-dependent transport of chloride in a watershed, Water Resour. Res., 51(1), 1–30,  
1161 doi:10.1002/2014WR015707, 2015.
- 1162 Heidbüchel, I., Troch, P. A., Lyon, S. W. and Weiler, M.: The master transit time distribution  
1163 of variable flow systems, Water Resour. Res., 48(6), W06520, doi:10.1029/2011WR011293,  
1164 2012.
- 1165 Hindmarsh, A. C., Brown, P. N., Grant, K. E., Lee, S. L., Serban, R., Shumaker, D. E. and  
1166 Woodward, C. S.: SUNDIALS: Suite of Nonlinear and Differential/Algebraic Equation  
1167 Solvers, ACM Trans. Math. Softw., 31(3), 363–396, doi:10.1145/1089014.1089020, 2005.
- 1168 Hollander, M., Wolfe, D. A. and Chicken, E.: Nonparametric Statistical Methods, John Wiley  
1169 & Sons, New York, NY, USA., 2013.
- 1170 Hrachowitz, M., Benettin, P., van Breukelen, B. M., Fovet, O., Howden, N. J. K., Ruiz, L., van  
1171 der Velde, Y. and Wade, A. J.: Transit times—the link between hydrology and water quality at  
1172 the catchment scale, Wiley Interdiscip. Rev. Water, doi:10.1002/wat2.1155, 2016.
- 1173 IAEA: International Atomic Energy Agency: Water Resources Programme – Global Network  
1174 of Isotopes in Precipitation, [online] Available from: [http://www-](http://www-naweb.iaea.org/napc/ih/IHS_resources_gnip.html)  
1175 naweb.iaea.org/napc/ih/IHS\_resources\_gnip.html (Accessed 11 August 2014), 2014.
- 1176 Jin, L., Siegel, D. I., Lautz, L. K., Mitchell, M. J., Dahms, D. E. and Mayer, B.: Calcite  
1177 precipitation driven by the common ion effect during groundwater–surface-water mixing: A  
1178 potentially common process in streams with geologic settings containing gypsum, Geol. Soc.  
1179 Am. Bull., 122(7/8), B30011.1, doi:10.1130/B30011.1, 2010.
- 1180 Jin, L., Siegel, D. I., Lautz, L. K. and Lu, Z.: Identifying streamflow sources during spring  
1181 snowmelt using water chemistry and isotopic composition in semi-arid mountain streams, J.  
1182 Hydrol., 470–471, 289–301, doi:10.1016/j.jhydrol.2012.09.009, 2012.
- 1183 Jouzel, J., Alley, R. B., Cuffey, K. M., Dansgaard, W., Grootes, P., Hoffmann, G., Johnsen, S.  
1184 J., Koster, R. D., Peel, D., Shuman, C. A., Stievenard, M., Stuiver, M. and White, J.: Validity  
1185 of the temperature reconstruction from water isotopes in ice cores, J. Geophys. Res., 102,  
1186 26471–26487, doi:10.1029/97JC01283, 1997.
- 1187 Kendall, C. and Caldwell, E. A.: Fundamentals of Isotope Geochemistry, in Isotope Tracers in  
1188 Catchment Hydrology, edited by C. Kendall and J. J. McDonnell, pp. 51–86, Elsevier,  
1189 Amsterdam, Netherlands., 1998.
- 1190 Kirchner, J. W.: A double paradox in catchment hydrology and geochemistry, Hydrol. Process.,  
1191 17(4), 871–874, doi:10.1002/hyp.5108, 2003.
- 1192 Klaus, J., McDonnell, J. J., Jackson, C. R., Du, E. and Griffiths, N. A.: Where does streamwater  
1193 come from in low-relief forested watersheds? A dual-isotope approach, Hydrol. Earth Syst.  
1194 Sci., 19(1), 125–135, 2015.
- 1195 Koeniger, P., Leibundgut, C. and Stichler, W.: Spatial and temporal characterisation of stable  
1196 isotopes in river water as indicators of groundwater contribution and confirmation of modelling  
1197 results; a study of the Weser river, Germany, Isotopes Environ. Health Stud., 45(4), 289–302,

1198 [doi:10.1080/10256010903356953](https://doi.org/10.1080/10256010903356953), 2009.

1199 [Kolaczyk, E. D.: Statistical Analysis of Network Data with R, Springer Science & Business](#)  
1200 [Media, New York, NY, USA., 2014.](#)

1201 [Kortelainen, N. M. and Karhu, J. A.: Regional and seasonal trends in the oxygen and hydrogen](#)  
1202 [isotope ratios of Finnish groundwaters: a key for mean annual precipitation, J. Hydrol., 285\(1–](#)  
1203 [4\), 143–157, doi:10.1016/j.jhydrol.2003.08.014, 2004.](#)

1204 [Kraft, P., Vaché, K. B., Frede, H.-G. and Breuer, L.: CMF: A Hydrological Programming](#)  
1205 [Language Extension For Integrated Catchment Models, Environ. Model. Softw., 26\(6\), 828–](#)  
1206 [830, doi:10.1016/j.envsoft.2010.12.009, 2011.](#)

1207 [Lai, C.-T. and Ehleringer, J. R.: Deuterium excess reveals diurnal sources of water vapor in](#)  
1208 [forest air, Oecologia, 165\(1\), 213–223, doi:10.1007/s00442-010-1721-2, 2011.](#)

1209 [Lauer, F., Frede, H.-G. and Breuer, L.: Uncertainty assessment of quantifying spatially](#)  
1210 [concentrated groundwater discharge to small streams by distributed temperature sensing, Water](#)  
1211 [Resour. Res., 49\(1\), 400–407, doi:10.1029/2012WR012537, 2013.](#)

1212 [LGR: Los Gatos Research, Greenhouse Gas, isotope and trace gas analyzers, \[online\] Available](#)  
1213 [from: <http://www.lgrinc.com/> \(Accessed 5 February 2013\), 2013.](#)

1214 [Li, F., Song, X., Tang, C., Liu, C., Yu, J. and Zhang, W.: Tracing infiltration and recharge using](#)  
1215 [stable isotope in Taihang Mt., North China, Environ. Geol., 53\(3\), 687–696,](#)  
1216 [doi:10.1007/s00254-007-0683-0, 2007.](#)

1217 [Maloszewski, P. and Zuber, A.: Manual on lumped parameter models used for the interpretation](#)  
1218 [of environmental tracer data in groundwaters, in Use of isotopes for analyses of flow and](#)  
1219 [transport dynamics in groundwater systems, p. 50, International Atomic Energy Agency,](#)  
1220 [Vienna, Austria., 2002.](#)

1221 [McConville, C., Kalin, R. m., Johnston, H. and McNeill, G. w.: Evaluation of Recharge in a](#)  
1222 [Small Temperate Catchment Using Natural and Applied  \$\delta^{18}\text{O}\$  Profiles in the Unsaturated Zone,](#)  
1223 [Ground Water, 39\(4\), 616–623, doi:10.1111/j.1745-6584.2001.tb02349.x, 2001.](#)

1224 [McDonnell, J. J. and Beven, K.: Debates—The future of hydrological sciences: A \(common\)](#)  
1225 [path forward? A call to action aimed at understanding velocities, celerities and residence time](#)  
1226 [distributions of the headwater hydrograph, Water Resour. Res., 50\(6\), 5342–5350,](#)  
1227 [doi:10.1002/2013WR015141, 2014.](#)

1228 [McDonnell, J. J., Sivapalan, M., Vaché, K., Dunn, S., Grant, G., Haggerty, R., Hinz, C., Hooper,](#)  
1229 [R., Kirchner, J., Roderick, M. L. and others: Moving beyond heterogeneity and process](#)  
1230 [complexity: a new vision for watershed hydrology, Water Resour. Res., 43\(7\), W07301, 2007.](#)

1231 [McDonnell, J. J., McGuire, K., Aggarwal, P., Beven, K. J., Biondi, D., Destouni, G., Dunn, S.,](#)  
1232 [James, A., Kirchner, J., Kraft, P., Lyon, S., Maloszewski, P., Newman, B., Pfister, L., Rinaldo,](#)  
1233 [A., Rodhe, A., Sayama, T., Seibert, J., Solomon, K., Soulsby, C., Stewart, M., Tetzlaff, D.,](#)  
1234 [Tobin, C., Troch, P., Weiler, M., Western, A., Worman, A. and Wrede, S.: How old is](#)  
1235 [streamwater? Open questions in catchment transit time conceptualization, modelling and](#)  
1236 [analysis, Hydrol. Process., 24\(12\), 1745–1754, 2010.](#)

1237 [McGuire, K. J. and McDonnell, J. J.: A review and evaluation of catchment transit time](#)  
1238 [modeling, J. Hydrol., 330\(3\), 543–563, 2006.](#)

1239 [Michel, R. L.: Residence times in river basins as determined by analysis of long-term tritium](#)  
1240 [records, J. Hydrol., 130\(1\), 367–378, doi:10.1016/0022-1694\(92\)90117-E, 1992.](#)



1241 [Mook, W. G., Groeneveld, D. J., Brouwn, A. E. and Ganswijk, A. J. van: Analysis of a run-off](#)  
1242 [hydrograph by means of natural <sup>18</sup>O, in Isotope techniques in groundwater hydrology, vol. 1,](#)  
1243 [pp. 159–169, International Atomic Energy Agency, Vienna, Austria., 1974.](#)

1244 [Neal, C. and Rosier, P. T. W.: Chemical studies of chloride and stable oxygen isotopes in two](#)  
1245 [conifer afforested and moorland sites in the British uplands, J. Hydrol., 115\(1–4\), 269–283,](#)  
1246 [doi:10.1016/0022-1694\(90\)90209-G, 1990.](#)

1247 [Newman, B., Tanweer, A. and Kurtas, T.: IAEA Standard Operating Procedure for the Liquid-](#)  
1248 [Water Stable Isotope Analyser, Laser Proced. IAEA Water Resour. Programme, 2009.](#)

1249 [O’Driscoll, M. A., DeWalle, D. R., McGuire, K. J. and Gburek, W. J.: Seasonal <sup>18</sup>O variations](#)  
1250 [and groundwater recharge for three landscape types in central Pennsylvania, USA, J. Hydrol.,](#)  
1251 [303\(1–4\), 108–124, doi:10.1016/j.jhydrol.2004.08.020, 2005.](#)

1252 [Orlowski, N., Frede, H.-G., Brüggemann, N. and Breuer, L.: Validation and application of a](#)  
1253 [cryogenic vacuum extraction system for soil and plant water extraction for isotope analysis, J.](#)  
1254 [Sens. Sens. Syst., 2\(2\), 179–193, doi:10.5194/jsss-2-179-2013, 2013.](#)

1255 [Orlowski, N., Lauer, F., Kraft, P., Frede, H.-G. and Breuer, L.: Linking Spatial Patterns of](#)  
1256 [Groundwater Table Dynamics and Streamflow Generation Processes in a Small Developed](#)  
1257 [Catchment, Water, 6\(10\), 3085–3117, doi:10.3390/w6103085, 2014.](#)

1258 [Penna, D., Ahmad, M., Birks, S. J., Bouchaou, L., Brenčič, M., Butt, S., Holko, L., Jeelani, G.,](#)  
1259 [Martínez, D. E., Melikadze, G., Shanley, J. B., Sokratov, S. A., Stadnyk, T., Sugimoto, A. and](#)  
1260 [Vreča, P.: A new method of snowmelt sampling for water stable isotopes, Hydrol. Process.,](#)  
1261 [28\(22\), 5637–5644, doi:10.1002/hyp.10273, 2014.](#)

1262 [Perry, C. and Taylor, K.: Environmental Sedimentology, p. 36, Blackwell Publishing, Oxford,](#)  
1263 [OX, UK, 2009.](#)

1264 [Pierce, S. C., Kröger, R. and Pezeshki, R.: Managing Artificially Drained Low-Gradient](#)  
1265 [Agricultural Headwaters for Enhanced Ecosystem Functions, Biology, 1\(3\), 794–856,](#)  
1266 [doi:10.3390/biology1030794, 2012.](#)

1267 [Qu, Y. and Duffy, C. J.: A semidiscrete finite volume formulation for multiprocess watershed](#)  
1268 [simulation, Water Resour. Res., 43\(8\), W08419, doi:10.1029/2006WR005752, 2007.](#)

1269 [Rinaldo, A., Benettin, P., Harman, C. J., Hrachowitz, M., McGuire, K. J., van der Velde, Y.,](#)  
1270 [Bertuzzo, E. and Botter, G.: Storage selection functions: A coherent framework for quantifying](#)  
1271 [how catchments store and release water and solutes, Water Resour. Res., 51\(6\), 4840–4847,](#)  
1272 [doi:10.1002/2015WR017273, 2015.](#)

1273 [Rohde, A.: Snowmelt-Dominated Systems, in Isotope Tracers in Catchment Hydrology, edited](#)  
1274 [by C. Kendall and J. J. McDonnell, pp. 391–433, Elsevier, Amsterdam, Netherlands., 1998.](#)

1275 [Rozanski, K., Sonntag, C. and Münnich, K. O.: Factors controlling stable isotope composition](#)  
1276 [of European precipitation, Tellus, 34\(2\), 142–150, doi:10.1111/j.2153-3490.1982.tb01801.x,](#)  
1277 [1982.](#)

1278 [Rozanski, K., Froehlich, K. and Mook, W. G.: in Environmental isotopes in the hydrological](#)  
1279 [cycle: Principles and Applications, vol. 3, International Hydrological Programme; United](#)  
1280 [Nations Educational, Scientific and Cultural Organization and International Atomic Energy](#)  
1281 [Agency, Paris, Vienna., 2001.](#)

1282 [Schultz, N. M., Griffis, T. J., Lee, X. and Baker, J. M.: Identification and correction of spectral](#)  
1283 [contamination in <sup>2</sup>H/<sup>1</sup>H and <sup>18</sup>O/<sup>16</sup>O measured in leaf, stem, and soil water, Rapid Commun.](#)

1284 [Mass Spectrom., 25\(21\), 3360–3368, doi:10.1002/rcm.5236, 2011.](#)

1285 [Shuttleworth, W. J. and Wallace, J. S.: Evaporation from sparse crops-an energy combination](#)  
1286 [theory, Q. J. R. Meteorol. Soc., 111\(469\), 839–855, doi:10.1002/qj.49711146910, 1985.](#)

1287 [Sklash, M. G.: Environmental isotope studies of storm and snowmelt runoff generation, in](#)  
1288 [Process studies in hillslope hydrology, edited by M. G. Anderson, pp. 410–435, Wiley, New](#)  
1289 [York, NY, USA., 1990.](#)

1290 [Sklash, M. G. and Farvolden, R. N.: The Role Of Groundwater In Storm Runoff, in](#)  
1291 [Developments in Water Science, vol. 12, edited by W. B. and D. A. Stephenson, pp. 45–65,](#)  
1292 [Elsevier, Amsterdam, Netherlands., 1979.](#)

1293 [Song, X., Wang, P., Yu, J., Liu, X., Liu, J. and Yuan, R.: Relationships between precipitation,](#)  
1294 [soil water and groundwater at Chongling catchment with the typical vegetation cover in the](#)  
1295 [Taihang mountainous region, China, Environ. Earth Sci., 62\(4\), 787–796, doi:10.1007/s12665-](#)  
1296 [010-0566-7, 2011.](#)

1297 [Stewart, M. K., Morgenstern, U. and McDonnell, J. J.: Truncation of stream residence time:](#)  
1298 [how the use of stable isotopes has skewed our concept of streamwater age and origin, Hydrol.](#)  
1299 [Process., 24\(12\), 1646–1659, doi:10.1002/hyp.7576, 2010.](#)

1300 [Stumpp, C., Klaus, J. and Stichler, W.: Analysis of long-term stable isotopic composition in](#)  
1301 [German precipitation, J. Hydrol., 517, 351–361, doi:10.1016/j.jhydrol.2014.05.034, 2014.](#)

1302 [Tang, K. and Feng, X.: The effect of soil hydrology on the oxygen and hydrogen isotopic](#)  
1303 [compositions of plants' source water, Earth Planet. Sci. Lett., 185\(3–4\), 355–367, 2001.](#)

1304 [Taylor, S., Feng, X., Kirchner, J. W., Osterhuber, R., Klaue, B. and Renshaw, C. E.: Isotopic](#)  
1305 [evolution of a seasonal snowpack and its melt, Water Resour. Res., 37\(3\), 759–769,](#)  
1306 [doi:10.1029/2000WR900341, 2001.](#)

1307 [Thomas, E. M., Lin, H., Duffy, C. J., Sullivan, P. L., Holmes, G. H., Brantley, S. L. and Jin, L.:](#)  
1308 [Spatiotemporal Patterns of Water Stable Isotope Compositions at the Shale Hills Critical Zone](#)  
1309 [Observatory: Linkages to Subsurface Hydrologic Processes, Vadose Zone J., 12\(4\), 0,](#)  
1310 [doi:10.2136/vzj2013.01.0029, 2013.](#)

1311 [Timbe, E., Windhorst, D., Crespo, P., Frede, H.-G., Feyen, J. and Breuer, L.: Understanding](#)  
1312 [uncertainties when inferring mean transit times of water trough tracer-based lumped-parameter](#)  
1313 [models in Andean tropical montane cloud forest catchments, Hydrol. Earth Syst. Sci., 18\(4\),](#)  
1314 [1503–1523, doi:10.5194/hess-18-1503-2014, 2014.](#)

1315 [van der Velde, Y., Torfs, P. J. J. F., van der Zee, S. E. a. T. M. and Uijlenhoet, R.: Quantifying](#)  
1316 [catchment-scale mixing and its effect on time-varying travel time distributions, Water Resour.](#)  
1317 [Res., 48\(6\), W06536, doi:10.1029/2011WR011310, 2012.](#)

1318 [Wang, X. F. and Yakir, D.: Using stable isotopes of water in evapotranspiration studies, Hydrol.](#)  
1319 [Process., 14\(8\), 1407–1421, doi:10.1002/1099-1085\(20000615\)14:8<1407::AID-](#)  
1320 [HYP992>3.0.CO;2-K, 2000.](#)

1321 [Windhorst, D., Waltz, T., Timbe, E., Frede, H.-G. and Breuer, L.: Impact of elevation and](#)  
1322 [weather patterns on the isotopic composition of precipitation in a tropical montane rainforest,](#)  
1323 [Hydrol. Earth Syst. Sci., 17\(1\), 409–419, doi:10.5194/hess-17-409-2013, 2013.](#)

1324 [Windhorst, D., Kraft, P., Timbe, E., Frede, H.-G. and Breuer, L.: Stable water isotope tracing](#)  
1325 [through hydrological models for disentangling runoff generation processes at the hillslope](#)  
1326 [scale, Hydrol. Earth Syst. Sci., 18\(10\), 4113–4127, doi:10.5194/hess-18-4113-2014, 2014.](#)

1327 [Woolfenden, L. R. and Ginn, T. R.: Modeled Ground Water Age Distributions, Ground Water, 47\(4\), 547–557, doi:10.1111/j.1745-6584.2008.00550.x, 2009.](#)  
1328  
1329 [Wu, J., Ding, Y., Ye, B., Yang, Q., Hou, D. and Xue, L.: Stable isotopes in precipitation in](#)  
1330 [Xilin River Basin, northern China and their implications, Chin. Geogr. Sci., 22\(5\), 531–540,](#)  
1331 [doi:10.1007/s11769-012-0543-z, 2012.](#)  
1332 [Xia, Y.: Optimization and uncertainty estimates of WMO regression models for the systematic](#)  
1333 [bias adjustment of NLDAS precipitation in the United States, J. Geophys. Res. Atmospheres,](#)  
1334 [111\(D8\), D08102, doi:10.1029/2005JD006188, 2006.](#)  
1335 [Yurtsever, Y.: Worldwide survey of stable isotopes in precipitation, International Atomic](#)  
1336 [Energy Agency, Vienna, Austria., 1975.](#)  
1337 [Zhao, L., Xiao, H., Zhou, J., Wang, L., Cheng, G., Zhou, M., Yin, L. and McCabe, M. F.:](#)  
1338 [Detailed assessment of isotope ratio infrared spectroscopy and isotope ratio mass spectrometry](#)  
1339 [for the stable isotope analysis of plant and soil waters, Rapid Commun. Mass Spectrom., 25\(20\),](#)  
1340 [3071–3082, doi:10.1002/rcm.5204, 2011.](#)  
1341



1342 Table 1. Descriptive statistics of  $\delta^2\text{H}$ ,  $\delta^{18}\text{O}$ , and d-excess values for precipitation, stream, and  
 1343 groundwater over the two-year observation period including all sampling points.


Sample type	Mean $\pm$ SD		Min		Max		D-excess mean $\pm$ SD	N
	$\delta^2\text{H}$	$\delta^{18}\text{O}$	$\delta^2\text{H}$	$\delta^{18}\text{O}$	$\delta^2\text{H}$	$\delta^{18}\text{O}$		
	[‰]	[‰]	[‰]	[‰]	[‰]	[‰]		
Precipitation	-43.9 $\pm$ 23.4	-6.2 $\pm$ 3.1	-167.6	-22.4	-8.3	-1.2	5.9 $\pm$ 5.7	592
Vollnkirchener Bach	-58.0 $\pm$ 2.8	-8.4 $\pm$ 0.4	-66.3	-10.0	-26.9	-6.7	9.0 $\pm$ 2.3	332
Schwingbach	-58.2 $\pm$ 4.3	-8.4 $\pm$ 0.6	-139.7	-18.3	-47.2	-5.9	9.0 $\pm$ 2.2	463
Groundwater meadow	-57.6 $\pm$ 1.6	-8.2 $\pm$ 0.4	-64.9	-9.2	-50.8	-5.7	7.9 $\pm$ 5.5	375
Groundwater arable land	-56.2 $\pm$ 3.7	-8.0 $\pm$ 0.5	-91.6	-12.3	-49.5	-6.8	1.7 $\pm$ 5.0	338
Groundwater along stream	-59.9 $\pm$ 6.8	-8.5 $\pm$ 0.9	-94.5	-13.0	-49.5	-7.0	8.2 $\pm$ 1.5	108

1344

1345 Table 2. Mean and standard deviation (SD) for isotopic signatures and soil physical properties in 0.2 m and 0.5 m soil depth (N=52 per depth).

	$\delta^2\text{H}$ [‰]		$\delta^{18}\text{O}$ [‰]		water content [% w/w]		pH		bulk density [g cm <sup>-3</sup> ]	
	0.2 m	0.5 m	0.2 m	0.5 m	0.2 m	0.5 m	0.2 m	0.5 m	0.2 m	0.5 m
Mean±SD	-46.9±8.4	-58.5±8.3	-6.6±1.2	-8.2±1.2	16.8±7.2	16.1±8.3	5.0±1.0	5.3±1.0	1.3±0.2	1.3±0.2

1346




1347

1348

1349 Figure 1. Maps show (a) the location of the Schwingbach catchment in Germany, (b) the main  
 1350 main monitoring area, (c) the land use, elevation, and instrumentation, (d) the locations of the

1351 snapshot as well as the seasonal soil samplings, (e) soil types, and (f) geology of the  
1352 Schwingbach catchment including the Vollnkirchener Bach subcatchment boudaries.  
1353




1354

1355

1356 Figure 2. Temporal variation of precipitation amount, isotopic signatures ( $\delta^2\text{H}$  and  $\delta^{18}\text{O}$ )  
 1357 including snow samples (grey striped box), and d-excess values for the study area compared to  
 1358 monthly d-excess values (July 2011 to July 2013) of GNIP station Koblenz with reference d-  
 1359 excess of GMWL ( $d=10$ ; solid black line).


1360



1361

1362

1363 Figure 3. Local Meteoric Water Line for the Schwingbach catchment (LMWL) in comparison  
 1364 to GMWL, including comparisons between precipitation, stream water, groundwater, and soil  
 1365 water isotopic signatures and the respective EWLs.




1366

1367

1368 Figure 4. Boxplots of  $\delta^2\text{H}$  values comparing precipitation, stream, groundwater, and soil  
 1369 isotopic composition in 0.2 m and 0.5 m depth (N=52 per depth). Different letters indicate  
 1370 significant differences ( $p \leq 0.05$ ).

1371




1372

1373

1374 Figure 5. Dual isotope plot of soil water isotopic signatures in 0.2 m and 0.5 m depth compared  
 1375 by land use including precipitation isotope data from 19, 21, and 28 October 2011. Insets:  
 1376 Boxplots comparing  $\delta^2\text{H}$  isotopic signatures between different land use units and precipitation  
 1377 (small letters) in top and subsoil (capital letters). Different letters indicate significant  
 1378 differences ( $p \leq 0.05$ ).






1379

1380

1381 Figure 6. Seasonal  $\delta^2\text{H}$  profiles of soil water (upper panels) and water content (lower panels)  
 1382 for winter (28 March 2013), summer (28 August 2011), and spring (24 April 2013). Error bars  
 1383 represent the natural isotopic variation of the replicates taken during each sampling campaign.  
 1384 For reference, mean groundwater (grey shaded) and mean seasonal precipitation  $\delta^2\text{H}$  values are  
 1385 shown (coloured arrows at the top).



1386

1387

1388 Figure 7. Mean daily discharge at the Vollnkirchener Bach (13, 18) and Schwingbach (site 11,


1389 19, and 64) with automatically recorded data (solid lines) and manual discharge measurements

1390 (asterisks), temporal variation of  $\delta^2\text{H}$  of stream water in the Schwingbach (site 11, 19, and 64)

1391 and Vollnkirchener Bach (site 13, 18, and 94) including moving averages (MA) for streamflow

1392 isotopes.

1393




1394

1395

1396 Figure 8. Temporal variation of discharge at the Vollnkirchener Bach with automatically  
 1397 recorded data (solid line) and manual discharge measurements (asterisks) (site 18), groundwater  
 1398 head levels, and  $\delta^2\text{H}$  values (coloured dots) for selected piezometers under meadow (site 3 and

1399 21), arable land (site 26, 27, and 28), and beside the Vollnkirchener Bach (site 24 and 32)  
1400 including moving averages for groundwater isotopes.




1401

1402

1403 Figure 9. Network map of  $\delta^{18}\text{O}$  relationships between surface water (SW) and groundwater  
 1404 (GW) sampling points. Yellow circles represent groundwater sampling points on the arable  
 1405 field, light green circles are piezometers located on the grassland close to the conjunction of the  
 1406 Schwingbach with the Vollnkirchener Bach, and dark green circles represent piezometers along  
 1407 the Vollnkirchener Bach. Light blue circles stand for Schwingbach and darker blue circles for  
 1408 Vollnkirchener Bach surface water sampling points. See Figure 1 for an overview of all  
 1409 sampling points. Only statistically significant connections between  $\delta^{18}\text{O}$  time series ( $p < 0.05$ )  
 1410 are shown in the network diagram.

1411



1412

1413

1414 Figure 10. Maps of modelled groundwater ages (colour scheme) and flow directions (white  
 1415 arrows) of (a) the Vollnkirchner Bach subcatchment and (b) detail view of the northern part of  
 1416 the subcatchment. The intensity of flow is depicted by the length of the white arrows.

1417

1418

1419 **Appendix I**

1420 **Mean transit time estimation**

1421 We applied a set of five different models to estimate the MTT using the FlowPC software  
1422 (Maloszewski and Zuber, 2002): dispersion model (with different dispersion parameters  
1423  $D_p=0.05, 0.4, \text{ and } 0.8$ ), exponential model, exponential-piston-flow model, linear model, and  
1424 linear-piston-flow model. We evaluated these results using two goodness of fit criteria, i.e.  
1425 sigma ( $\sigma$ ) and model efficiency (ME) following Maloszewski and Zuber (2002):

$$\sigma = \frac{\sqrt{\sum(c_{mi} - c_{oi})^2}}{m} \quad (1)$$

$$ME = 1 - \frac{\sum(c_{mi} - c_{oi})^2}{\sum(c_{oi} - \bar{c}_o)^2} \quad (2)$$

1426 Where:

- 1427 •  $c_{mi}$ : The i-th model result
- 1428 •  $c_{oi}$ : The i-th observed result
- 1429 •  $\bar{c}_o$ : The arithmetic mean of all observations

1430

1431 A model efficiency  $ME=1$  indicates an ideal fit of the model to the concentrations observed,  
1432 while  $ME=0$  indicates that the model fits the data no better than a horizontal line through the  
1433 mean observed concentration (Maloszewski and Zuber, 2002). The same is true for sigma. For  
1434 calculations with FlowPC, weekly averages of precipitation and stream water isotopic  
1435 signatures are calculated. We firstly calculated the MTT from precipitation to the streams for  
1436 three sampling points in the Vollnkirchener Bach (sites 13, 18 and 94) and three points in the  
1437 Schwingbach (sites 11, 19 and 64). For the second set of simulations, the mean residence time  
1438 from precipitation to groundwater comprising thirteen groundwater sampling points was  
1439 determined. We also bias-corrected the precipitation input data with two different approaches:  
1440 the mean precipitation value is subtracted from every single precipitation value and then divided  
1441 by the standard deviation of precipitation isotopic signatures. Afterwards, this value is  
1442 subtracted from the weekly precipitation values (bias1). For the second approach, the difference  
1443 of the mean stream water isotopic value and the mean precipitation value is calculated and also  
1444 subtracted from the weekly precipitation values (bias2).

1445

## 1446 **Appendix II**

### 1447 **Model-based groundwater age dynamics**

1448 Objective:

1449 Stable water isotopes are only a tool to determine the residence time for a few years (McDonnell  
1450 et al., 2010). In cases of longer residence times and a strong mixing effect, seasonal variation  
1451 of isotopes vanishes and results in ~~stable flat lines of the~~barely varying isotopic ~~signals~~signals.  
1452 To get a rough estimate of residence times greater than the limit of stable water isotopes (>5  
1453 years), we split the water flow path in our catchment in two parts: the flow from precipitation  
1454 to groundwater, which was calculated via FlowPC and the longer groundwater transport. The  
1455 simplest method to estimate the residence time of groundwater transport is via the storage-to-  
1456 input-relation, with the storage as the aquifer size and the input as the groundwater recharge  
1457 time. However, this method ignores the topographic setting, and water input heterogeneity. In  
1458 our study we used a simplified groundwater flow model with tracer transport to calculate the  
1459 groundwater age dynamics. The numerical output of water ages cannot be validated with the  
1460 given isotope data, since the model is used to fill a residence time gap, where stable water  
1461 isotopes are not feasible to apply. The model is falsified however, if the residence time is short  
1462 enough (<5 years) to be calculable via FlowPC. Hence, the results of the groundwater age model  
1463 should be handled with care and only seen as the order of magnitude of flow time scales.

1464 Model setup:

1465 We set up a tailored hydrological model for the Vollnkirchener Bach subcatchment using the  
1466 *Catchment Modelling Framework* (CMF) by Kraft et al. (2011). CMF is a modular framework  
1467 for hydrological modelling based on the concept of finite volume method by Qu and Duffy  
1468 (2007). CMF is applicable for simulating one- to three-dimensional water fluxes but also  
1469 advective transport of stable water isotopes ( $^{18}\text{O}$  and  $^2\text{H}$ ). Thus, it is especially suitable for our  
1470 tracer study and can be used to study the origin (Windhorst et al., 2014) and age of water. To  
1471 avoid errors in transit time calculations from small differences between the isotopic signal in  
1472 groundwater and stream water, we are tracing the transit time of groundwater and not the real  
1473 isotopic values in this study. The generated model is a highly simplified representation of the  
1474 Vollnkirchener Bach subcatchment's groundwater body. The subcatchment is divided into 353  
1475 polygonal-shaped cells ranging from 100–40'000 m<sup>2</sup> in size based on land use, soil type, and  
1476 topography. The model is vertically divided in two compartments, the upper soft rock aquifer,  
1477 and the lower bed rock aquifer, referred to as upper and lower layer from now onwards.



1478 The layers of each cell are connected using a mass conservative Darcy approach with a finite  
 1479 volume discretization. The water storage dynamic of one layer in one cell  $i$  of the groundwater  
 1480 model is given as:

$$\frac{dV_{i,s}}{dt} = R_i - S_i - \sum_{j=1}^{N_i} \left( K_s \frac{\Psi_{i,s} - \Psi_{j,s}}{d_{ij}} A_{ij,s} \right) \quad (3)$$

$$\frac{dV_{i,b}}{dt} = S_i - \sum_{j=1}^{N_i} \left( K_b \frac{\Psi_{i,b} - \Psi_{j,b}}{d_{ij}} A_{ij,b} \right)$$

1481

1482 Where:

- 1483 •  $V_i$ : The water volume stored by the layer in  $\text{m}^3$  in cell  $I$  for soft rock ( $s$ ) and bedrock  
 1484 ( $b$ ), respectively
- 1485 •  $R_i$ : The groundwater recharge rate in  $\text{m}^3 \cdot \text{d}^{-1}$
- 1486 •  $S_i$ : the percolation from the soft rock to the bedrock aquifer, calculated by the  
 1487 gradient and geometric mean conductivity between the layers:  $S_i =$   
 1488  $\sqrt{K_s K_b} \frac{\Psi_{i,s} - \Psi_{i,b}}{d_{sb}} A_i$ , where  $d_{sb}$  is the distance between the layers and  $A_i$  is the cell area
- 1489 •  $N_i$ : Number of adjacent cells to cell  $i$
- 1490 •  $K$ : Saturated hydraulic conductivity in  $\text{m} \cdot \text{d}^{-1}$  for soft rock ( $s$ ) and bedrock ( $b$ ),  
 1491 respectively
- 1492 •  $\Psi$ : Water head in the current cell  $i$  and the neighbour cell  $j$  in m for soft rock ( $s$ ) and  
 1493 bedrock ( $b$ ), respectively
- 1494 •  $d_{ij}$ : The distance between the current cell  $i$  and the neighbour cell  $j$  in m
- 1495 •  $A_{i,j,x}$ : The wetted area of the joint layer boundary in  $\text{m}^2$  between cells  $i$  and  $j$  in layer  
 1496  $x$

1497 The volume head relation is linearized as  $\Psi = \phi \frac{V}{A}$ , with  $\phi$  being the fillable porosity and  $A$  the  
 1498 cell area. The resulting ordinary differential equation system is integrated using the CVODE  
 1499 solver by Hindmarsh et al. (2005), an error controlled Krylov-Newton multistep implicit solver  
 1500 with an adaptive order of 1–5 according to stability constraints.

1501 Boundary conditions:

1502 The upper boundary condition of the groundwater system – the mean groundwater recharge –  
1503 is modelled applying a Richard's equation based model using measured rainfall data (2011–  
1504 2013) and calculated evapotranspiration with the Shuttleworth-Wallace method (Shuttleworth  
1505 and Wallace, 1985) including land cover and climate data. To retrieve long-term steady state  
1506 conditions, the groundwater recharge is averaged and used as constant flow Neumann boundary  
1507 condition. The total outflow is calibrated against measured outflow data; hence, the unsaturated  
1508 model's role is mainly to account for spatial heterogeneity of groundwater recharge. As an  
1509 additional input, a combined sewer overflow (site 38, Fig. 1b) is considered based on findings  
1510 of Orłowski et al. (2014). Moreover, there are two water outlets in the two lowest cells for  
1511 efficient draining, reflecting measured groundwater flow directions throughout most of the year  
1512 at piezometers 1–6 (Fig. 1b). Both cells are located in the very north of the subcatchment and  
1513 their outlets are modelled as constant head Dirichlet boundary condition.

1514 Parameters:

1515 The saturated hydraulic conductivity of the groundwater body is set to  $0.1007 \text{ m d}^{-1}$ , as  
1516 measured in the study area. For the lower bedrock compartment there is no data available.  
1517 However, expecting a high rate of joints, preliminary testing revealed that a saturated hydraulic  
1518 conductivity of  $0.25 \text{ m d}^{-1}$  seemed to be a realistic estimation (based on field measurements).

1519 Water Age:

1520 To calculate the water age in each cell, a virtual tracer flows through the system using advective  
1521 transport. To calculate the water age from the tracer that enters the system with a unity  
1522 concentration by groundwater recharge, a linear decay is used to reduce the tracer concentration  
1523 with time:

$$\frac{dX_{i,s}}{dt} = 1 \frac{u}{m^3} R_i - S_i [X]_{i,s} - \sum_{j=1}^{N_i} \left( [X]_{i,s} K_s \frac{\Psi_{i,s} - \Psi_{j,s}}{d_{ij}} A_{ij,s} \right) - r X_{i,s} \quad (4)$$
$$\frac{dX_{i,b}}{dt} = S_i [X]_{i,s} - \sum_{j=1}^{N_i} \left( [X]_{i,b} K_b \frac{\Psi_{i,b} - \Psi_{j,b}}{d_{ij}} A_{ij,b} \right) - r X_{i,b}$$

$$t_{ix} = \frac{\ln[X]_{ix}}{r}$$

1524

1525 Where:

- 1526 •  $X_{i,x}$ : Amount of virtual tracer in layer  $x$  in cell  $i$  in virtual unit  $u$
- 1527 •  $1 \frac{u}{m^3} R_i$ : Tracer input with groundwater recharge  $R$  with unity concentration
- 1528 •  $[X]_{i,x}$ : Concentration of tracer in layer  $x$  of cell  $i$  in  $u m^{-3}$
- 1529 •  $r$ : Arbitrary chosen decay constant, for water age calculation in  $d^{-1}$ . Rounding errors
- 1530 occur due to low concentrations when  $r$  is set to a high value. We found a good
- 1531 numerical performance with values between  $10^{-6}$ – $10^{-9} d^{-1}$
- 1532 •  $t_{ix}$ : Water age in days in layer  $x$  in cell  $i$

1533

1534 To ensure long term steady state conditions, the model is run for 2000 years. However, after  
1535 300 years of model run time, steady state is reached.