

1 **Landscape heterogeneity drives contrasting concentration-**  
2 **discharge relationships in shale headwater catchments**

3  
4 **Elizabeth M. Herndon<sup>1,a</sup>, Ashlee L. Dere<sup>1,b</sup>, Pamela L. Sullivan<sup>1,c</sup>, David Norris<sup>2</sup>,**  
5 **Brian Reynolds<sup>2</sup>, and Susan L. Brantley<sup>1</sup>**

6  
7 **Author affiliations:**

8 [1] Department of Geosciences, The Pennsylvania State University, University Park, PA 16802

9 [2] Centre for Ecology and Hydrology, Bangor, Gwynedd LL57 2UW, United Kingdom

10  
11 **Current affiliations**

12 <sup>a</sup> Department of Geology, Kent State University, Kent OH 44242

13 <sup>b</sup> Department of Geography/Geology, University of Nebraska at Omaha, Omaha NE 68182

14 <sup>c</sup> Department of Geography, The University of Kansas, Lawrence KS 66045

15  
16  
17 **Corresponding author:**

18 Elizabeth Herndon (eherndo1@kent.edu)

19 Department of Geology, Kent State University, Kent OH 44242

25 **Abstract**

26 Solute concentrations in stream water vary with discharge in patterns that record complex  
27 feedbacks between hydrologic and biogeochemical processes. In a comparison of three shale-  
28 underlain headwater catchments located in Pennsylvania, USA (the forested Shale Hills Critical  
29 Zone Observatory) and Wales, U.K. (the peatland-dominated Upper Hafren and forest-  
30 dominated Upper Hore in the Plynlimon forest), dissimilar concentration-discharge behaviors are  
31 best explained by contrasting landscape distributions of soil solution chemistry – especially  
32 dissolved organic carbon (DOC) – that have been established by patterns of vegetation and soil  
33 organic matter (SOM). Specifically, elements that are concentrated in organic-rich soils due to  
34 biotic cycling (Mn, Ca, K) or that form strong complexes with DOC (Fe, Al) are spatially  
35 heterogeneous in pore waters because organic matter is heterogeneously distributed across the  
36 catchments. These solutes exhibit non-chemostatic behavior in the streams, and solute  
37 concentrations either decrease (Shale Hills) or increase (Plynlimon) with increasing discharge. In  
38 contrast, solutes that are concentrated in soil minerals and form only weak complexes with DOC  
39 (Na, Mg, Si) are spatially homogeneous in pore waters across each catchment. These solutes are  
40 chemostatic in that their stream concentrations vary little with stream discharge, likely because  
41 these solutes are released quickly from exchange sites in the soils during rainfall events.  
42 Furthermore, concentration-discharge relationships of non-chemostatic solutes changed  
43 following tree harvest in the Upper Hore catchment in Plynlimon, while no changes were  
44 observed for chemostatic solutes, underscoring the role of vegetation in regulating the  
45 concentrations of certain elements in the stream. These results indicate that differences in the  
46 hydrologic connectivity of organic-rich soils to the stream drive differences in concentration  
47 behavior between catchments. As such, in catchments where SOM is dominantly in lowlands  
48 (e.g. Shale Hills), we infer that non-chemostatic elements associated with organic matter are  
49 released to the stream early during rainfall events, whereas in catchments where SOM is  
50 dominantly in uplands (e.g. Plynlimon), these non-chemostatic elements are released later during  
51 rainfall events. The distribution of SOM across the landscape is thus a key component for  
52 predictive models of solute transport in headwater catchments.

53

54 **Keywords**

- 55 Critical Zone, Catchment Hydrology, Concentration-Discharge, Dissolved Organic Carbon, Soil
- 56 Organic Matter, Chemostasis

## 57 **1 Introduction**

58 Streams are regularly monitored to evaluate watershed geochemistry, ecosystem health, and  
59 suitability for human use. However, streams integrate hydrologic and biogeochemical processes  
60 over varied spatial and temporal scales, making it difficult to determine both the sources and  
61 flow paths of solutes. While many researchers examine short- to long-term element variability in  
62 stream water, it has remained difficult to derive generalized models quantifying solute  
63 concentration-discharge behavior (Fisher et al. 2004; Sivapalan 2005; Zimmer et al. 2012). Flow  
64 paths may dictate stream chemistry by controlling fluid residence times and chemical  
65 equilibration of flowing water with soil minerals within catchments (Maher, 2011). Therefore, it  
66 is necessary to understand how heterogeneous flow paths through distinct chemical sources  
67 within a catchment influence observed solute concentration patterns within streams.

68 When the discharge of a stream ( $Q$ ) increases, concentrations of solutes ( $C$ ) can either  
69 increase (enrichment behavior), decrease (dilution behavior), or, perhaps most paradoxically,  
70 change very little (chemostasis) (Kirchner, 2003; Godsey et al., 2009; Clow and Mast, 2010).  
71 Dilution can result during rainfall events as water stored in a catchment is diluted by less  
72 concentrated meteoric water. Enrichment can result if a more concentrated source (e.g.  
73 groundwater) mixes with stream water during large rainfall events (Johnson et al. 1969). In  
74 contrast, chemostasis cannot be explained by the simple mixing of multiple sources and therefore  
75 has been attributed to processes such as chemical reactions with the solid-phase along the  
76 pathway of water flow (Godsey et al. 2009). Although changing flow paths through soil horizons  
77 may explain differences in solute response to discharge along hillslope transects (e.g., Bishop et  
78 al., 2004), solutes often show different types of behavior in different streams due to landscape  
79 heterogeneity, and a unifying explanation for C-Q behavior has remained elusive.

80 Behavior differences amongst individual solutes in the stream have been linked to variability  
81 in solute concentrations within a catchment: in other words, discrete zones of element  
82 mobilization within soils and sediments can lead to pulses of solute transport into a stream  
83 (McClain et al., 2003; Andrews et al., 2011). This effect is furthermore affected by changes in  
84 hydrologic connectivity, defined as the water-mediated transfer of constituents between water  
85 sources (Pringle, 2001), within a catchment during rainfall events. Stream chemistry can vary  
86 during storm events as dominant water inputs to the stream shift from groundwater and riparian  
87 zones during base flow to hillslope runoff at high flow as pore waters stored in upland soils

88 become increasingly connected to the stream (McGlynn and McDonnell, 2003a). Throughout  
89 this paper, groundwater is defined as water that is stored in catchment soils and bedrock below  
90 the water table, and pore water is defined as water that is present in the pores of unsaturated soil  
91 in the vadose zone. Upland soils become hydrologically connected to the stream when soil layers  
92 become water-saturated, promoting downslope flow within the unsaturated zone. As a result,  
93 concentrations of solutes that are stored preferentially in the riparian zone, e.g. dissolved organic  
94 carbon (DOC) released from soil organic matter (SOM), peak in the stream prior to discharge or  
95 with rising discharge during storm events (McGlynn and McDonnell, 2003b; Hood et al. 2006).  
96 Variability in organic carbon dynamics across different landscape units can subsequently control  
97 metal export from headwater catchments and downstream hydrochemistry (Köhler et al., 2014).

98 Many previous studies examine single catchments and/or catchments that were developed on  
99 multiple lithologies (e.g., Johnson et al., 1969; Krám et al., 1997; Brown et al., 1999; Likens and  
100 Buso, 2006; Godsey et al., 2009), making the interpretation of solute behaviors difficult at best.  
101 When mono-lithologic catchments are compared, insights into other factors that influence the  
102 response of stream chemistry to discharge (e.g. biota, climatic) can be developed. To elucidate  
103 controls on stream chemistry not primarily driven by lithology, we examined C-Q relationships  
104 in three shale-underlain headwater streams with extensive hydrogeochemical datasets. Although  
105 these catchments are underlain by chemically similar shales, their soils have developed distinct  
106 and contrasting distributions of SOM across each landscape; i.e., organic-rich soils are  
107 predominantly in low-lands and swales in the Shale Hills Critical Zone Observatory but in  
108 upland peat regions of the Upper Hore and Upper Hafren catchments in the Plynlimon forest.  
109 Additionally, we investigate how C-Q patterns change following tree harvest in the forested  
110 Upper Hore. For these catchments, variations in stream chemistry with flow elucidate non-  
111 lithological factors that control solute transport to streams, yielding a paradigm that should help  
112 explain other catchments.

## 113 **2 Methods**

114 Water chemistry was compared for three sites: 1) the Susquehanna Shale Hills Critical Zone  
115 Observatory (Shale Hills) in central Pennsylvania, USA and 2) the Upper Hore and 3) Upper  
116 Hafren subcatchments in the Plynlimon experimental forest in Wales, UK (Figure 1). The Shale  
117 Hills and Plynlimon forests are underlain almost exclusively by Fe-rich, organic-poor, Silurian-

118 aged shale formations that are stratigraphically equivalent. Although these headwater catchments  
119 vary by size and location, their similar lithologies and extensive hydrogeochemical  
120 characterization (e.g. Kirby et al., 1991; Neal et al., 1997; Shand et al., 2005(a-b); Jin et al.,  
121 2010; Neal et al., 2011; Brantley et al., 2013(a-j); Dere et al., 2013; Neal et al., 2013(a-b)) allow  
122 development of a unifying theory on factors controlling concentration-discharge behavior.

## 123 **2.1 Susquehanna Shale Hills Critical Zone Observatory**

124 Shale Hills is an 8 ha forested headwater catchment nested within the larger Susquehanna  
125 River Basin in Pennsylvania, USA. Shale Hills contains primarily Inceptisol soils developed  
126 from shale residuum or colluvium of the Rose Hill Formation, which is dominantly comprised of  
127 clay minerals and quartz (Lin et al., 2006; Jin et al., 2010). Small areas of Ultisols are present  
128 near the stream (Lin et al., 2006). The regional mean annual temperature (MAT) is 10°C, and  
129 precipitation (MAP =  $105 \pm 17$  cm  $y^{-1}$ ) is acidic (e.g., pH averaged  $4.5 \pm 0.2$  for U.S. National  
130 Atmospheric Deposition Program (NADP) sites PA15 & PA42 during 2000-2011). Vegetation is  
131 dominated by deciduous oaks and hickories, and the elevation ranges from 256 m at the  
132 catchment outlet to 310 m on the ridge. Hillslopes are characterized as either “planar” (mildly  
133 convex-upward slopes with shallow soils that grade to concave-upward slopes toward valley  
134 floor) or “swale” (concave-upward depressions with deep soils and convergent water flow) (Lin  
135 et al., 2006) (Figure 1). Water flows vertically through pores when soils are unsaturated, or  
136 downslope along horizon interfaces when rainfall events create transiently perched saturated  
137 water zones (Lin, 2006; Jin et al., 2011). Soils in swales are generally wetter than soils on planar  
138 hillslopes and remain hydrologically connected to the stream during dry periods, whereas water  
139 flow through planar hillslopes is negligible under dry conditions and increases with increasing  
140 precipitation (Lin et al., 2006; Qu and Duffy, 2007; Takagi and Lin, 2012). Soils in the swales  
141 also store more organic carbon than soils on planar hillslopes and act as sources of DOC  
142 transport into streams (Andrews et al., 2011) (Table 1).

143 Water samples from Shale Hills were collected approximately daily from the stream outlet  
144 (2008 – 2010) and biweekly from soil lysimeters (2006 – 2011) from March through early  
145 December each year (Table 2). Detailed methods and results of chemical analyses, including  
146 isotopic variation and concentrations of major ions and DOC, have been reported elsewhere (Jin  
147 et al., 2011; Andrews et al., 2011; Brantley et al., 2013(a-j)). Aluminum concentrations in the

148 stream were consistently below detection limits; thus, Al data were not examined for Shale Hills.  
149 Daily discharge rates were estimated from continuous discharge measurements integrated over  
150 10 min intervals from the stream weir at the catchment's outlet (Duffy, 2012). Soil water was  
151 collected from suction lysimeters installed in the soil at 10 cm depth increments from 10 to 50  
152 cm depth in the south planar valley floor (SPVF) and from 10 to 90 cm depth in the south swale  
153 valley floor (SSVF). The groundwater was sampled from a 2.8 m deep well located 80 m  
154 upstream from the weir. Major cation (2000-2011; NADP, 2011) and trace element  
155 concentrations (Herndon, 2012) have been reported for precipitation samples collected from  
156 NADP sites PA-15 and PA-42. Vegetation chemistry was previously determined for green leaf  
157 and leaf litter samples collected throughout summer and fall seasons, respectively, in the Shale  
158 Hills catchment (Herndon et al., 2015).

## 159 **2.2 Plynlimon forest: Upper Hore and Upper Hafren catchments**

160 The Plynlimon forest is a 682 ha watershed located at the headwater of the River Severn, 20  
161 km from the west coast of Wales (Reynolds et al., 1997). MAT is 7.2 °C and MAP is 250 ± 78  
162 cm with an average pH of 4.98 ± 0.01. Vegetation is predominantly evergreen Sitka spruce  
163 (*Picea sitchensis*) with areas of heath, including *Sphagnum* and *Juncus* communities, dominating  
164 the uplands. Elevation in the Plynlimon forest ranges from 319 to 738 m.

165 We focus here on two adjacent headwater catchments within the Plynlimon watershed: the  
166 Upper Hore and the Upper Hafren (Figure 1). The Upper Hore (162 ha) is predominantly  
167 forested with periodically saturated, organic-rich Stagnopodzol soils and uplands that are  
168 dominated by grass and saturated Peat soils (Kirby et al., 1991). In contrast, the Upper Hafren  
169 (122 ha) is dominated by heath and Peat soils, with waterlogged and organic-rich Peaty gley soils  
170 located in riparian areas (Kirby et al., 1991). Generally, the main flow paths in both catchments  
171 are approximately orthogonal to the valley direction, with highly fractured shallow bedrock  
172 providing an important pathway and storage for water throughout the catchments, especially  
173 under base flow conditions (Haria and Shand, 2004; Shand et al., 2005(a-b); Shand et al., 2007).  
174 Shallow and deeper groundwater appear to be poorly connected but some mixing does occur  
175 (Haria and Shand, 2004; Shand et al., 2005b; Shand et al., 2007). Flow in organic horizons,  
176 however, tends to be largely lateral rather than vertical, providing minimal water-rock interaction

177 in peat dominated portions of the catchment and increasing contribution to streams during high  
178 flow conditions (Shand et al., 2009).

179 Stream chemistry data for the Upper Hore and Upper Hafren catchments were collected  
180 throughout the year for all years between 1983-2005 and 1990-2010, respectively (Neal et al.,  
181 2013a and 2013b). Due to extensive tree-cutting in the Upper Hore in 2005, data collected from  
182 2005-2010 were evaluated separately to examine the influence of tree removal on C-Q behavior.  
183 Stream discharge was measured every 15 min at weirs in both catchments and weekly stream  
184 grab samples were analyzed for major and trace ions. Likewise, bulk precipitation was collected  
185 weekly at the Carreg Wen meteorological station located between the Upper Hore and Upper  
186 Hafren catchments (Reynolds et al., 1997). Precipitation chemistry was influenced by seawater  
187 inputs, which varied with wind direction and season (Reynolds et al., 1987). Groundwater  
188 chemistry was estimated as average concentrations of solutes in seven shallow (< 3 m) wells  
189 located within the Plynlimon forest near the two catchments (Figure 1). Wells were sampled  
190 weekly from 1994 – 1999 (Neal et al., 1997) (Table A1). Average solute concentrations ( $\pm$   
191 standard error) were calculated for pore waters sampled from soils classified as Peat,  
192 Stagnopodzol, or Peaty Gley throughout Plynlimon (Reynolds et al., 1987; Reynolds et al., 1988;  
193 Stevens et al., 1997; Shand et al., 2005b) (Table 3; Table A2).

### 194 **2.3 Stream flow regimes and concentration-discharge (C-Q) behaviors**

195 To analyze stream chemistry under different flow regimes, stream water discharge ( $Q$ ,  $\text{m}^3 \text{d}^{-1}$ )  
196 was categorized as low-flow (lower quartile of  $Q$ ), moderate-flow (interquartile range), or  
197 high-flow (upper quartile) (Figure A1). Perennial stream flow with little seasonal variation in  
198 discharge was observed in the Upper Hafren and Upper Hore catchments, and the distributions of  
199  $Q$  were positively skewed by intermittent storms. Stream flow at Shale Hills was intermittent and  
200 highly seasonal, with extended periods of no- to low-flow during summer months (especially  
201 July and August). Solute concentrations for each flow regime at each site were averaged over all  
202 collection dates (Tables 2 and 3).

203 Linear regressions were fit to log-transformed C-Q data for each solute  $j$  ( $C_j$ ) for each  
204 catchment stream. We included all available data for all years for Shale Hills (2008 – 2010), the  
205 Upper Hafren (1990 – 2010), and the pre-harvest Upper Hore (1983 – 2004). The data collected  
206 for the Upper Hore following harvest (2005 – 2010) were evaluated separately. The slope of this



207 regression ( $m_j$ ) was used to identify solute behaviors as “chemostatic” or “non-chemostatic”.  
208 When  $m_j$  showed only minimal effects of dilution by meteoric water, i.e.,  $-0.1 < m_j < 0$ , the  
209 solutes were defined as chemostatic (Godsey et al., 2009). In contrast, non-chemostatic elements  
210 were defined to exhibit dilution behavior when concentrations decreased with increasing  $Q$  ( $m_j <$   
211  $-0.1$ ) or enrichment behavior when concentrations increased with increasing  $Q$  ( $m_j > 0$ ). Linear  
212 regressions and associated statistical parameters were calculated in Origin.

213 To investigate sources of solutes mobilized to the stream, element ratios in the stream under  
214 different flow regimes were compared to element ratios in pore waters, precipitation,  
215 groundwater, and leaves (where available). Element ratios have been used in other studies to link  
216 river chemistry to end member reservoirs (e.g. Gaillardet et al., 1999). Molar ratios of divalent  
217 cations (Ca:Mg) were compared to univalent cations (K:Na) in each reservoir to understand how  
218 elements exhibiting non-chemostasis (Ca, K) vary relative to chemostatic elements (Mg, Na). To  
219 further explore the association of certain non-chemostatic solutes with organic C, molar ratios of  
220 Mn (a non-chemostatic element) to Mg were compared to DOC concentrations. Average DOC  
221 concentrations were used to define soil waters as organic-rich or organic-poor, as discussed in  
222 section 3.2. In Shale Hills, green leaves were used to represent the most organic-rich end-  
223 member since pore waters could not be sampled from the thin O horizon.

## 224 **3 RESULTS**

### 225 **3.1 Solute concentration-discharge patterns**

226 Na and Mg behaved near-chemostatically in all catchments (Figure 2; Table 4) while Si and  
227 K were only chemostatic at Shale Hills. A subset of non-chemostatic solutes exhibited similar  
228 trends to DOC; however, trends were opposite between Shale Hills and Plynlimon. Specifically,  
229 when  $Q$  increased, concentrations of the non-chemostatic solutes Ca, Mn, Fe and DOC decreased  
230 at Shale Hills (i.e. dilution behavior;  $m_j < -0.1$ ) while the non-chemostatic solutes Mn, K, Al,  
231 Fe, and DOC increased at Plynlimon (i.e. enrichment behavior;  $m_j > 0$ ) (Figure 2b; Table 4).  
232 Note that Fe exhibited enrichment ( $m_{Fe} = 0.33 \pm 0.02$ ) similar to DOC in the Upper Hafren but  
233 was more consistent with chemostasis in the Upper Hore ( $m_{Fe} = -0.05 \pm 0.02$ ). Additionally, Si  
234 and Ca showed dilution patterns at Plynlimon that contrasted with DOC.

235 In the Shale Hills stream, higher concentrations of stream solutes were observed during the  
236 dry summer season (June through September) relative to the wetter spring and autumn (Figure  
237 A2). While concentrations of the chemostatic elements increased only slightly (~2X) during the  
238 summer, larger increases were observed for Ca (~4X), DOC (~7X), and Mn and Fe (> 100X).  
239 Increases in DOC, K, Fe, and Mn in the stream during summer were not consistent with  
240 increasing groundwater inputs because groundwater at Shale Hills is depleted in these elements  
241 relative to stream water at low flow (Table 2). In the Upper Hafren and Upper Hore streams,  
242 chemostatic elements Na and Mg, derived primarily from sea salts, showed no seasonality  
243 despite high seasonal variation in inputs from precipitation (Figures A3-A5; Reynolds et al.,  
244 1987), as if precipitation-derived solutes were buffered in the catchment soil pore waters before  
245 entering the stream (Neal and Kirchner, 2000). In contrast, solutes exhibiting enrichment (K, Al,  
246 Mn, Fe, DOC) varied by season (Figures A3 and A4).

247 In the Upper Hore where trees were harvested, solute concentrations and C-Q slopes  
248 increased following tree harvest for solutes showing enrichment behavior. Specifically, stream  
249 concentrations of DOC, K, Mn, and Fe increased after 2005 (Table 3). Post-harvest C-Q slopes  
250 for  $m_K$  ( $= 0.26 \pm 0.03$ ) and  $m_{Mn}$  ( $= 0.12 \pm 0.02$ ) increased relative to pre-harvest values ( $0.07 \pm$   
251  $0.01$  and  $0.05 \pm 0.01$ , respectively) (Figure 2; Table 4). No effects of tree harvest on  $C_j$  or  $m_j$   
252 were observed for chemostatic elements. Additionally, no changes in solute concentrations in the  
253 non-harvested Upper Hafren were observed over this time.

## 254 **3.2 Soil pore waters**

255 We examined the chemical composition of soil pore waters in order to investigate sources of  
256 solutes to the stream. Pore waters in each catchment were categorized into distinct chemical  
257 pools based on DOC concentrations (Table 2): “organic-rich” waters were defined by average  
258 DOC > 1 mM, while all other waters were “organic-poor”. At Shale Hills, pore waters collected  
259 from the A horizon (10 cm) of the swale (SSVF) were organic-rich ( $1.28 \pm 0.33$  mM DOC)  
260 while all other pore waters were organic-poor; i.e. the A horizon of SPVF ( $0.69 \pm 0.06$  mM  
261 DOC) and the B horizons of SSVF and SPVF (> 10 cm; averaged  $0.55 \pm 0.04$  mM DOC) were  
262 organic-poor. At Plynlimon, pore waters in organic horizons and Peat soils ( $1.2 \pm 0.2$  mM DOC)  
263 were organic-rich relative to mineral horizons of Stagnopodzol and Peaty gley soils ( $0.42 \pm 0.07$   
264 mM DOC) soils.

265 At Shale Hills, concentrations of the non-chemostatic solutes Mn, Fe, and Ca showed  
266 evidence of DOC-related behavior. For example, Mn and Fe were positively correlated with  
267 DOC across all pore waters ( $R^2 > 0.9$ ) and were highest in the organic-rich waters ( $6.8 \pm 1.9 \mu\text{M}$   
268 and  $1.7 \pm 0.3 \mu\text{M}$ , respectively) (Table 2). Calcium concentrations were enriched in the A  
269 horizon ( $72 \pm 11 \mu\text{M}$ ) relative to the B horizon ( $35 \pm 1 \mu\text{M}$ ) in SSVF. Furthermore, Fe and Mn  
270 concentrations were spatially variable across pore waters (% RSD = 100% and 140%,  
271 respectively). In contrast, chemostatic solutes Mg (33%), Na (19%), and Si (5%) were less  
272 variable. Thus, concentrations of non-chemostatic solutes were spatially heterogeneous in soil  
273 waters across the landscape while chemostatic solutes were distributed more homogeneously.

274 Like Shale Hills, concentrations of the chemostatic elements Na and Mg were spatially  
275 homogeneous in pore waters at Plynlimon amongst the different soils (RSD = 23% and 15%,  
276 respectively) (Table 3). Additional solutes chiefly derived from the atmosphere via precipitation  
277 (K, Ca) or through carbon fixation (DOC) were enriched in the organic horizons, while elements  
278 derived primarily from minerals (Si, Fe, Al, and Mn) were enriched in mineral horizons. In  
279 contrast to Shale Hills, many of the non-chemostatic elements at Plynlimon were not correlated  
280 with DOC in soil pore waters.

281 Element ratios in stream water under low, moderate, and high flow regimes were compared  
282 to element ratios in pore waters, precipitation, and groundwater (Figure 3). At Shale Hills, stream  
283 chemistry was most similar to pore waters from organic-rich soils and green leaves at low flow  
284 and approached values for pore waters from organic-poor soils at high flow. Ratios of  $C_{Ca}:C_{Mg}$   
285 and  $C_K:C_{Na}$  decreased slightly ( $< 2X$ ) with increasing discharge (Figure 3a), while  $C_{Mn}:C_{Mg}$   
286 decreased by 10X (Figure 3b). Stream water was more similar to soil pore waters than  
287 groundwater or precipitation under all flow regimes, documenting that flow through shallow  
288 soils and bedrock rather than deep groundwater sources dominated inputs to the stream. These  
289 trends further indicate a shift from inputs of organic-rich soil water to the stream at low flow to  
290 organic-poor soil water at high flow.

291 In contrast to this behavior at Shale Hills, stream chemistries in the Upper Hore and Upper  
292 Hafren catchments were most similar to organic-poor sources (precipitation, groundwater) at low  
293 flow and organic-rich sources (soil pore waters) at high flow (Figure 3c-f; Figure A6). Values of  
294  $C_{Mn}:C_{Mg}$ ,  $C_{DOC}$ , and  $C_K:C_{Na}$  increased while  $C_{Ca}:C_{Mg}$  decreased and converged towards the  
295 most organic-rich end-member in each system, either the peat (Upper Hafren) or peat and

296 organic horizon pore waters (Upper Hore), with increasing discharge. Stream  $C_{Ca}:C_{Mg}$  ratios  
297 were similar to groundwater at low flow in the Upper Hafren. The limited groundwater data that  
298 were available for Plynlimon indicate that groundwater was not chemically similar to stream  
299 water under any flow regime in the Upper Hore.

### 300 **3.3 Organic influence on concentration-discharge behavior**

301 Finally, we explored how chemical heterogeneity in soil pore waters influenced  
302 concentration-discharge relationships in the streams. Specifically, we evaluated solute  
303 heterogeneity due to redistribution by vegetation as the ratio of solute concentrations in “organic-  
304 rich” to “organic-poor” pore waters. As previously defined, these pore waters were collected  
305 from A versus B horizons at Shale Hills, and organic versus mineral soils in the Upper Hafren  
306 and Upper Hore. The slope of the concentration-discharge plot ( $m_j$ ) was used to define the  
307 magnitude of non-chemostatic behavior for each solute, i.e. the degree to which an element was  
308 diluted or enriched in the stream with increasing discharge.

309 At Shale Hills, elements concentrated in the organic-rich pore waters were diluted rapidly in  
310 the stream with increasing discharge, consistent with increasing inputs of water from mineral  
311 soils as the planar hillslope soils become saturated during storms (Qu and Duffy, 2007). This  
312 trend is documented in Figure 4a where the concentration ratios for organic-rich versus -poor soil  
313 waters were negatively correlated with respect to  $m_j$  ( $R^2 = 0.90$ ,  $p < 0.001$ ). According to these  
314 results, Fe and Mn were most concentrated in organic-rich pore waters and most rapidly diluted  
315 in the stream, followed by DOC, Ca, and K. Chemostatic elements Na, Mg, and Si were not  
316 concentrated in organic-rich pore waters.

317 No significant correlation ( $p > 0.05$ ) existed between organic to mineral pore water ratios and  
318  $m_j$  in the Upper Hore and Upper Hafren subcatchments, likely because organic horizons at  
319 Plynlimon have high concentrations of chemostatic solutes due large inputs of sea salts that  
320 dominate the chemical signature of near-surface pore waters. Provided this observation, organic-  
321 associations in each catchment were evaluated by inspecting the ratio of average solute  
322 concentrations in the pore water versus precipitation (Figure 4b,c), i.e., precipitation serves as the  
323 most organic-poor pool in the Plynlimon system. For both the Upper Hafren and the Upper Hore,  
324 the ratios of concentrations in soil water versus precipitation were positively correlated with  $m_j$   
325 ( $p < 0.05$ ). Elements exhibiting enrichment behavior, including DOC, Al, Mn, and K in both

326 catchments plus Fe in the Upper Hafren, were also enriched in pore water relative to  
327 precipitation. Chemostatic elements in pore waters were less enriched relative to precipitation. In  
328 contrast, the ratios for soil water versus precipitation were not significantly correlated with  $m_j$  at  
329 Shale Hills ( $p > 0.05$ ).

## 330 **4 Discussion**

331 Cross-site comparison of the Shale Hills and Plynlimon headwater catchments revealed that  
332 the behaviors of non-chemostatic solutes were controlled by the spatial variability of those  
333 elements in soil waters and the distribution of DOC. Conversely, chemostatic solutes were  
334 homogeneously distributed in pore waters across the catchments. In the following sections, we  
335 discuss how the landscape distribution of chemically distinct pools and the connectivity between  
336 organic-rich soils and the stream control how concentrations vary with discharge. We contend  
337 that the behavior of certain elements are non-chemostatic in these systems due to their  
338 association with organic matter. The distribution of soil organic matter across landscapes is in  
339 turn influenced by climate (e.g., SOM generally increases with increasing moisture and  
340 decreasing temperatures on large geographic scales) and geomorphology (e.g., organic matter  
341 accumulates in depressed areas such as swales on small geographic scales).

### 342 **4.1 Hydrologic connectivity of solute pools across landscapes**

343 At first glance, it may appear contradictory that concentrations of non-chemostatic elements  
344 in the streams at Shale Hills and Plynlimon trend in opposite directions with increasing  
345 discharge; however, the discrepancy can be explained by differences in the distributions of  
346 organic-rich source waters in each system. Similar to bioactive elements identified by Stallard  
347 and Murphy (2013), we attribute non-chemostatic concentration-discharge behavior to changing  
348 water flow through organic-rich soil matrices; however, we also observe that organic-rich  
349 sources and flow paths vary between the catchments (Figure 1).

350 At Shale Hills, meteoric water passes through the thin organic horizon and organic-rich A  
351 horizon (< 15 cm deep) and is transported along the horizon interfaces to the stream via  
352 preferential flow paths (Lin et al. 2006; Jin et al., 2011; Thomas et al., 2013). The stream  
353 receives water from organic-rich swales and surface soils during dry periods, and water inputs  
354 from organic-poor hillslope soils increase as the catchment saturates (Qu and Duffy, 2007;

355 Andrews et al. 2011). Consequently, we observed that stream water chemistry was similar to  
356 organic-rich soil waters at low flow and organic-poor soil waters at high flow (Figure 3). Solutes  
357 derived largely from organic-rich soils exhibited greater variability over different flow regimes  
358 due to their high spatial variability in soil pore water. Increasingly negative slopes for non-  
359 chemostatic elements at high discharge (Figure 2b) may reflect the transition in hydrologic  
360 connectivity and hillslope inputs to the stream. Stream chemistry did not reflect inputs from  
361 groundwater during dry periods, consistent with a previous finding that the water table drops to >  
362 2 m below the stream bed during late summer (Thomas et al., 2013).

363 In the grass-dominated Upper Hafren, which contains peat soils that experience minimal  
364 water-rock interaction (Kirby et al., 1991), concentrations of chemostatic elements in the soils  
365 never deviated far from an average precipitation signal (Figure 4). In contrast, concentrations of  
366 non-chemostatic elements were not driven by precipitation, and we propose that pore water  
367 concentrations of these elements are regulated by vegetation. During the drier growing season,  
368 certain non-chemostatic elements may be depleted from soil pore water and accumulated in  
369 vegetation, leading to lower concentrations in the stream. Indeed, it is well-established that  
370 seasonal uptake by vegetation regulates concentrations of nutrient elements in stream water (e.g.,  
371 Johnson et al., 1969; Vitousek, 1977; Mulholland, 1992). Warming and drying of the surface  
372 peat during this time increases microbial decomposition, thereby increasing mobility of elements  
373 that accumulate in vegetation by releasing them from storage in organic matter (Kirby et al.,  
374 1991). According to this conceptual model, once transpiration decreases and flow increases  
375 through the soil in autumn, concentrations of these elements increase in the stream because 1)  
376 transpiration is reduced and the soil water is no longer being depleted; 2) the surface peat is  
377 flushed of elements that have accumulated, providing elements in addition to precipitation. As  
378 observed at the Upper Hafren and Upper Hore, concentrations of non-chemostatic elements  
379 begin to increase in the stream as discharge increases following low-flow in the summer (Figures  
380 A3 and A4). This effect may be especially prominent in the peat regions since the grass  
381 vegetation decomposes annually with little aboveground storage (i.e. peat is leaky with respect to  
382 nutrients), and anoxic conditions limit complete conversion of SOM to CO<sub>2</sub>, enhancing release  
383 of DOC. Although groundwater can discharge to streams in Plynlimon during summer months  
384 (Shand et al., 2005b), the groundwater contains little DOC (Table 3) and therefore cannot  
385 explain changes in summer stream chemistry where DOC increases from spring to summer

386 (Figures A3 and A4). At high flow, decreases in Mn concentrations in the stream (Figures 2d and  
387 2g) may reflect the low abundance of Mn in the catchment relative to other elements; in other  
388 words, the supply of Mn is depleted at high flow and Mn is diluted in the stream.

389 In the spruce-forested Upper Hore, long-term storage of nutrient elements in above-ground  
390 biomass is expected to deplete soil pore waters of elements without the flushing effect due to  
391 rapid turnover observed in the Upper Hafren (Reynolds et al., 2000). Instead, the positive  
392 concentration-discharge slopes in the Upper Hore result from flushing of upland peat soils at  
393 high flow conditions (Neal et al., 1990). These effects can be observed by comparing pre- and  
394 post-harvest concentration-discharge slopes in the Upper Hore. Tree harvest impacted stream  
395 concentrations and C-Q slopes for K, Al, Fe, Mn, and DOC but had no observable effects on  
396 chemostatic elements. Specifically,  $m_K$  increased from 0.07 to 0.25 following harvest, while the  
397 peat-dominated Upper Hafren, which was not harvested, maintained  $m_K = 0.15$  during this time.  
398 Fe and DOC experienced increases in stream concentrations, but not  $m_j$ . Since neither Fe nor  
399 DOC is expected to be taken up into the trees in high quantities, increasing  $C_j$  may indicate that  
400 they were mobilized by increased decomposition of leaf litter debris in the forest region  
401 following harvest (Hughes et al., 1990).

402 Values of  $C_j$  and  $m_j$  for non-chemostatic, organic-associated elements likely increased  
403 following harvest because 1) these elements were no longer being taken up and stored in tree  
404 biomass (Stevens et al., 1997); 2) the organic debris left after harvest provided a labile pool of  
405 organic chelator molecules and organically-complexed elements, and 3) inputs from upland peats  
406 to the stream increased due to lack of interception by the forest (Neal et al., 1992; Neal et al.,  
407 2004). Such short-term increases in nutrient loss following ecosystem disturbance are well  
408 documented, especially for clear-cut systems (e.g., Likens et al., 1970; Vitousek, 1977; Neal et  
409 al., 1992). Although non-chemostatic in the Upper Hafren, Fe followed a chemostatic trend in  
410 the Upper Hore (Figure 2). This behavior could be due to Fe retention in the forested soils during  
411 downslope transport: Stagnopodzols on these slopes have Bs horizons that accumulate  
412 sesquioxides and immobilize leached Fe (Reynolds, 1982).

413 Previous studies have hypothesized that hydrologic connectivity within landscapes  
414 (McGlynn and McDonnell, 2003a; Hood et al. 2006; Clow and Mast 2010) and/or interactions  
415 between soil moisture and mineral reactive surface area (Godsey et al. 2009; Clow and Mast  
416 2010) can explain concentration-discharge relationships across multiple catchments. Our results

417 contribute to the understanding of solute behavior by highlighting the importance of hydrologic  
418 connectivity across landscapes and at mineral surfaces. At both Shale Hills and Plynlimon, the  
419 distribution of soil organic matter and its hydrologic connection to the stream governed non-  
420 chemostatic concentration discharge behavior of several solutes (Ca, K, Mn, Fe and Al), a  
421 process similarly invoked to explain stream DOC behavior in storm events (McGlynn and  
422 McDonnell, 2003b). Our results highlight the need to include or enhance reactive transport  
423 modules in spatially-distributed watershed-scale hydrologic models such as TOPMODEL  
424 (Beven and Kirkby 1979), the Penn State Integrated Hydrologic Model (PIHM; Qu and Duffy  
425 2007), and the Regional Hydro-Ecological Simulation System (RHESys; Brand et al. 1991).  
426 Specifically, combining RTM with the ability of spatially-distributed models to simulate soil  
427 moisture, temperature, and water fluxes at variable depths across geomorphic features (e.g.,  
428 swales vs. planar slopes) will allow researchers to elucidate specific flow water paths and transit  
429 times and better test drivers of chemostasis (cation exchange) and dynamics of mobile vs.  
430 immobile water. RT-Flux-PIHM is one model under development (Duffy et al. 2014) that will  
431 provide this platform, but it is imperative to cross-compare outputs from various models in order  
432 to reach consensus.

433

## 434 **4.2 Drivers of chemostasis**

435 Stream concentrations for most major weathering elements ( $j = \text{Na, K, Mg, Ca, Si}$ ) varied  
436 little over a wide range of discharge values (Figure 2): by this definition, most of the major  
437 elements were chemostatic, with the exception of Ca at Shale Hills and K at Plynlimon. Note that  
438 the standard descriptor “major elements” includes Na and Ca although both are low in abundance  
439 in the protolith at Shale Hills and Plynlimon ( $< 0.7 \text{ wt.}\%$  and  $< 0.2 \text{ wt.}\%$ , respectively), and K is  
440 lower at Plynlimon compared to Shale Hills ( $2.90 \text{ wt.}\%$  and  $3.76 \text{ wt.}\%$ , respectively) (Jin et al.,  
441 2010; Dere et al., 2013). Chemostasis could be expected for elements derived from minerals that  
442 are always equilibrated with long residence-time pore waters. However, the fast-dissolving  
443 minerals present at Shale Hills and Plynlimon, carbonates and pyrite, do not contain K, Na and  
444 Si and are also depleted in the upper protolith (Jin et al., 2010; Neal et al., 1997). Therefore,  
445 dissolution of these minerals cannot explain chemostasis of K, Na, Si, Mg and Ca. Godsey et al.  
446 (2009) suggested that changes in mineral-water interfacial area during periods of high and low  
447 discharge explain chemostasis; however, clay dissolution rates are too slow (Bandstra et al.,



448 2008) to provide a rapidly mobilized source of cations during the short-timescales of  
449 precipitation events.

450 The exchangeable cation pool is a likely source of chemostatic elements during rain events  
451 (e.g., Clow and Mast, 2010). The cation exchange capacity of soils along the planar hillslope at  
452 Shale Hills ranges from 35 to 71 meq kg<sup>-1</sup> (Jin et al., 2010). At Plynlimon, forest and peat soils  
453 have a cation exchange capacity of roughly 77.4 and 300 meq kg<sup>-1</sup>, respectively (Reynolds et al.,  
454 1988; Cuttle, 1983). Elements are displaced from cation exchange sites into solution by H<sup>+</sup> (i.e.  
455 protonation of the exchange sites), and we observe that the degree of chemostasis for  
456 chemostatic elements was inversely related to the elements' relative strengths of adsorption to  
457 cation exchange sites as reported by Evangelou et al. (2005). Furthermore, this explanation can  
458 even account for the highly chemostatic, neutrally charged solute Si(OH)<sub>4</sub><sup>0</sup>, which has also been  
459 documented in the exchangeable pool at Shale Hills (Jin et al., 2010). For these catchments  
460 where pH is low (pH < 7), Si should be weakly associated to exchange sites due to its neutral  
461 charge. The similar concentrations observed for major weathering elements in the planar and  
462 swale pore waters at Shale Hills as well as Plynlimon (Tables 2 and 3) are attributed to the quick  
463 exchange of protons in rain for cations in the exchange pool throughout the catchment. Thus,  
464 chemostasis is explained by increasing connectivity of the exchangeable pool, i.e., cations bound  
465 to surfaces of minerals and soil organic matter, to mobile water as soil saturation increases.

### 466 **4.3 Chemostatic nutrients become non-chemostatic under nutrient-limiting** 467 **conditions**

468 Although geochemically similar to Mg, K, and Na, the concentration-discharge pattern for  
469 Ca (Figure 2) is non-chemostatic at Shale Hills. The mixing model (Figure 3a) indicates leaves  
470 may be a primary source of Ca to the stream during low discharge. Indeed, these shallow soils  
471 are strongly leached of Ca (< 0.16% wt.; Jin et al. 2010), and organic matter may be a relatively  
472 large pool of Ca in this system. In contrast to Shale Hills where Ca trends are strongly influenced  
473 by organic matter, Ca at Plynlimon may be linked to groundwater, an effect most pronounced in  
474 the Upper Hore. Ratios of Ca:Mg trend towards organic-poor sources at low flow, likely due to  
475 inputs of Ca-rich groundwater during base flow that is diluted by increasing contribution from  
476 soil water at high flow. Although a lack of groundwater data from these two subcatchments

477 limits our ability to directly assess inputs to the stream, groundwater collected from lower  
478 elevations in the Plynlimon forest are rich in Ca and Si (Neal et al., 1997).

479       Similar to Ca at Shale Hills, K limitation may drive its increased association with organic  
480 matter at Plynlimon. Values of  $C_K:C_{Na}$  decrease at Shale Hills and increase at Plynlimon with  
481 increasing  $Q$  in a manner consistent with changing inputs from organic-rich sources (Figure A6).  
482 Although geochemically similar, Na is a nonessential element (Kirkby, 2012) that is readily  
483 weathered from soils whereas K is a plant macronutrient that accumulates in leaf tissue (Herndon  
484 et al., 2015). From the mixing diagrams and  $m_K$ , we infer that K has a stronger organic control at  
485 Plynlimon than at Shale Hills. One explanation for this is that net primary productivity (NPP) is  
486 higher ( $896 \text{ g C m}^{-2} \text{ y}^{-1}$ ; unpublished data) but bedrock K is lower ( $2.90 \pm 0.13\%$ ; Dere et al.,  
487 2013) at Plynlimon than at Shale Hills (NPP =  $550 \text{ g C m}^{-2} \text{ y}^{-1}$ ; Smith, 2013 and  $K = 3.76 \pm$   
488  $0.16\%$ ; Jin et al. 2010). These data suggest that K is limiting to vegetation at Plynlimon while Ca  
489 is limiting to vegetation at Shale Hills due to high demand and low supply.

## 490 **5 Conclusions**

491       A comparison of three shale-derived catchments, the Shale Hills CZO in central  
492 Pennsylvania, U.S.A. and the Upper Hafren and Upper Hore catchments in the Plynlimon forest,  
493 Wales, U.K., reveals that the concentration-discharge behaviors of elements are strongly  
494 impacted by the distribution of organic matter in soils and the hydrologic connectivity of these  
495 soils to the stream. At Shale Hills, stream water is derived from organic-rich swales at low flow  
496 and then from both swale and planar hillslopes with increasing flow. At Plynlimon, stream water  
497 is only dominated by water from organic-rich soils at high flow, and contributions from organic-  
498 rich upland soils increased following lower elevation tree harvest in the Upper Hore catchment.  
499 Solutes that are limiting nutrients or that are strongly retained by vegetation exhibit non-  
500 chemostatic behavior in the stream because they are released to the stream along with dissolved  
501 organic carbon. This non-chemostatic behavior is opposite between Plynlimon and Shale Hills  
502 due to the different landscape distribution of organic-rich soils. Due to minimal redistribution by  
503 vegetation, Na, Mg, and Si are equally concentrated in pore fluids for organic-rich and organic-  
504 poor soils, and concentrations of these elements in stream water remain relatively constant. From  
505 this, we conclude that the transport of elements associated with organic matter, termed  
506 previously as organomarker elements (Hausrath et al., 2009), is strongly controlled by the

507 movement of dissolved organic carbon, leading to a distinct non-chemostatic behavior in stream  
508 waters that contrasts with the chemostatic behavior of major weathering elements. Stream  
509 chemistry in headwater catchments are variable largely because of the chemical heterogeneities  
510 in distribution of organic-rich soils in catchments and how those soils connect to the stream.

511

### 512 **Author contributions**

513 EMH, PLS and ALD analyzed the data. EMH prepared the manuscript with contribution from all  
514 authors.

515

### 516 **Acknowledgements**

517 The authors thank Adam Wlostowski and two anonymous reviewers for helpful comments that  
518 improved the manuscript. This work was facilitated by NSF Critical Zone Observatory program  
519 grants to CJD (EAR 07-25019) and SLB (EAR 12-39285, EAR 13-31726). This research was  
520 conducted in Penn State's Stone Valley Forest, which is supported and managed by the Penn  
521 State Forestland Management Office in the College of Agricultural Sciences. Plynlimon data and  
522 support were provided by the Natural Environment Research Council Centre for Ecology and  
523 Hydrology (<https://gateway.ceh.ac.uk/home>). The authors acknowledge scientists who  
524 participated in sample collection and analysis, particularly L. Jin and D. Andrews. Special thanks  
525 to S. Grant for sample collection and D. Robinson, I. Robinson, S. Hughes and B. Emmett for  
526 logistical support at Plynlimon. Additional support was provided by NSF grant EAR-10-52614  
527 (to SLB), the Penn State Center for Global Studies, and the NSF GK-12 CarbonEARTH Grant  
528 (EHR-0947962).

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544 **References**

- 545  
546 Andrews, D. M., Lin, H., Zhu, Q., Jin, L., & Brantley, S. L. (2011). Hot spots and hot moments  
547 of dissolved organic carbon export and soil organic carbon storage in the Shale Hills  
548 catchment. *Vadose Zone Journal*, 10(3), 943-954.
- 549 Band, L.E., Peterson, D.L., Running, S.W., Coughlan, J., Lammers, R., Dungan, J., and Nemani,  
550 R., 1991. Forest ecosystem processes at the watershed scale: basis for distributed simulation.  
551 *Ecological Modelling*, 56, 171-196., DOI: 10.1016/0304-3800(91)90199-B
- 552 Bandstra JZ, Buss HL, Campen RK, Liermann LJ, Moore J, Hausrath EM, Navarre-Sitchler AK,  
553 Jang J, Brantley SL (2008) Compilation of mineral dissolution rates. In *Kinetics of Water-*  
554 *Rock Interaction* (eds. S. Brantley, J. Kubicki, A. White).
- 555 Beven, K J and M J. Kirkby. 1979. A physically based variable contributing area model of basin  
556 hydrology. *Hydrologic Science Bulletin*. 24(1):43-69.
- 557 Bishop K., Siebert J., Kohler S., Laudon H. (2004) Resolving the double paradox of rapidly  
558 mobilized old water with highly variable reponses in runoff chemistry. *Hydrol. Process.* **18**,  
559 185-189.
- 560 Brantley SL, Ketchum B, White T, Sullivan PL (2013) Susquehanna Shale Hills Critical Zone  
561 Observatory Porewater Chemistry (2006). EarthChem. DOI: 10.2136/vzj2010.0133
- 562 Brantley SL, Ravela R, Jin L, Nuester J, Sullivan PL (2013) Susquehanna Shale Hills Critical  
563 Zone Observatory Porewater Chemistry (2007). EarthChem. DOI: 10.2136/vzj2010.0133
- 564 Brantley SL, Jin L, Andrews D, Holmes G, Holleran M, Williams J, Herndon E, Sullivan PL  
565 (2013) Susquehanna Shale Hills Critical Zone Observatory Porewater Chemistry (2008).  
566 EarthChem. DOI: 10.1594/IEDA/100235
- 567 Brantley SL, Jin L, Andrews D, Holmes G, Bhatt M, Holleran M, Kaiser N, Williams J, Herndon  
568 E, Sullivan PL (2013) Susquehanna Shale Hills Critical Zone Observatory Porewater  
569 Chemistry (2009). EarthChem. DOI: 10.1594/IEDA/100236
- 570 Brantley SL, Bazilevskaya E, Andrews D, Williams J, Herndon E, Holmes G, Bhatt M, Holleran  
571 M, Yesavage T, Thomas E, Sullivan PL (2013) Susquehanna Shale Hills Critical Zone  
572 Observatory Porewater Chemistry (2010). EarthChem. DOI: 10.1594/IEDA/100237
- 573 Brantley SL, Ketchum B, White T, Sullivan PL (2013) Susquehanna Shale Hills Critical Zone  
574 Observatory Stream Water Chemistry (2006). EarthChem. DOI: 10.1594/IEDA/100239
- 575 Brantley SL, Ravela R, Jin L, Nuester J, Sullivan PL (2013) Susquehanna Shale Hills Critical  
576 Zone Observatory Stream Water Chemistry (2007). EarthChem. DOI: 10.1594/IEDA/100240
- 577 Brantley SL, Jin L, Andrews D, Holmes G, Holleran M, Williams J, Herndon E, Duffy CJ,  
578 Sullivan PL (2013) Susquehanna Shale Hills Critical Zone Observatory Stream Water  
579 Chemistry (2008). EarthChem. DOI: 10.1594/IEDA/100241
- 580 Brantley SL, Jin L, Andrews D, Holmes G, Bhatt M, Holleran M, Kaiser N, Williams J, Herndon  
581 E, Duffy CJ, Sullivan PL (2013) Susquehanna Shale Hills Critical Zone Observatory Stream  
582 Water Chemistry (2009). EarthChem. DOI: 10.1594/IEDA/100242
- 583 Brantley SL, Bazilevskaya E, Andrews D, Williams J, Herndon E, Holmes G, Bhatt M, Holleran  
584 M, Yesavage T, Thomas E, Duffy CJ, Sullivan PL (2013) Susquehanna Shale Hills Critical  
585 Zone Observatory Stream Water Chemistry (2010). EarthChem. DOI: 10.1594/IEDA/100243
- 586 Brantley, S. L., Holleran M. E., Jin L. and Bazilevskaya E. (2013) Probing deep weathering in  
587 the Shale Hills Critical Zone Observatory, Pennsylvania (USA): the hypothesis of nested  
588 chemical reaction fronts in the subsurface. *Earth Surface Processes and Landforms*. DOI:  
589 10.1002/esp.3415.

590 Brown, V. A., McDonnell, J. J., Burns, D. A., & Kendall, C. (1999). The role of event water, a  
591 rapid shallow flow component, and catchment size in summer stormflow. *Journal of*  
592 *Hydrology*, 217(3), 171-190.

593 Boyer E.W., Hornberger G.M., Bencala K.E., McKnight D.M. (1997) Response characteristics  
594 of DOC flushing in an alpine catchment. *Hydrological Processes* 11: 1635-1647.

595 Clow, D. W., & Mast, M. A. (2010). Mechanisms for chemostatic behavior in catchments:  
596 implications for CO<sub>2</sub> consumption by mineral weathering. *Chemical Geology*, 269(1), 40-  
597 51.

598 Cuttle, S. P. (1983) Chemical properties of upland peats influencing the retention of phosphate  
599 and potassium ions. *Journal of Soil Science*. 34:75-82.

600 Dere, A. L., White T. S., April R. H., Reynolds B., Miller T. E., Knapp E. P., McKay L. D. and  
601 Brantley S. L. (2013) Climate dependence of feldspar weathering along a latitudinal  
602 gradient. *Geochim. Cosmochim. Acta*. 122:101-126.

603 Duffy C. (2012) Stream flow data. Available at: <http://www.czo.psu.edu>

604 Duffy, C., Shi, Y., Davis, K., Slingerland, R., Li, L., Sullivan, P. L., & Brantley, S. L. (2014).  
605 Designing a Suite of Models to Explore Critical Zone Function. *Procedia Earth and*  
606 *Planetary Science*, 7-15. doi:10.1016/j.proeps.2014.08.003

607 Evangelou, V. P., Phillips, R. E., Tabatabai, M. A., Sparks, D. L., Al-Amoodi, L., & Dick, W. A.  
608 (2005). Cation exchange in soils. *Chemical processes in soils*, 343-410.

609 Fisher SG, Sponseller RA, Heffernan JB. 2004. Horizons in stream biogeochemistry: flowpaths  
610 to progress. *Ecology* 85: 2369-2379

611 Gaillardet, J., Dupré, B., Louvat, P., & Allegre, C. J. (1999). Global silicate weathering and CO  
612<sub>2</sub> consumption rates deduced from the chemistry of large rivers. *Chemical Geology*, 159(1),  
613 3-30.

614 Godsey SE, Kirchner JW, Clow DW. 2009. Concentration-discharge relationships reflect  
615 chemostatic characteristics of US catchments. *Hydrological Processes* 23: 1844-1864.  
616 DOI:10.1002/hyp.7315.

617 Herndon E.M. (2012) Biogeochemistry of manganese contamination in a temperate forested  
618 watershed. PhD Dissertation, The Pennsylvania State University.

619 Hood E, Gooseff MN, Johnson S. 2006. Changes in the character of stream water dissolved  
620 organic carbon during flushing in three small watersheds, Oregon. *Journal of Geophysical*  
621 *Research* 11: 1-8. DOI: 10.1029/2005JG000082

622 Hughes S., Reynolds B. and Roberts J.D. (1990) The influence of land management on  
623 concentrations of dissolved organic carbon and its effects on the mobilization of aluminium  
624 and iron in podzol soils in Mid-Wales. *Soil Use and Management*. 6:137-145.

625 Jin L, Andrews DM, Holmes GH, Lin H, Brantley SL. 2011. Opening the “Black Box”: water  
626 chemistry reveals hydrological controls on weathering in the Susquehanna Shale Hills  
627 Critical Zone Observatory. *Vadose Zone Journal* 10: 928-942. DOI:10/2136/vzj2010.0133

628 Jin L, Ravella R, Ketchum K, Bierman PR, Heaney P, White T, Brantley SL. 2010. Mineral  
629 weathering and elemental transport during hillslope evolution at the Susuehanna/Shale Hills  
630 Critical Zone Observatory. *Geochimica et Cosmochimica Acta* 74 : 3669-3691.

631 Johnson NM, Likens GE, Bormann FH, Fisher DW, Pierce RS. 1969. A working model for the  
632 variation in stream water chemistry at the Hubbard Brook Experimental Forest, New  
633 Hampshire. *Water Resources Research* 5: 1353-1363.

634 Haria A.H. and Shand P. 2004. Evidence for deep sub-surface flow routing in forested upland  
635 Wales: implications for contaminant transport and stream flow generation. *Hydrology and*  
636 *Earth System Sciences*. 8:334-344.

637 Kirby, C., Newson M.D. and Gilman K. (1991) Plynlimon Research: The first two decades.  
638 Institute of Hydrology Report No. 109, Wallingford, UK.

639 Kirkby E. (2012) Introduction, definition, and classification of nutrients. In *Marschner's Mineral*  
640 *Nutrition of Higher Plants*. DOI: 10.1016/B978-0-12-384905-2.00001-7.

641 Kirchner JW. 2003. A double paradox in catchment hydrology and geochemistry. *Hydrological*  
642 *Processes* 17: 872-874.

643 Köhler, S. J., Lidman, F., & Laudon, H. (2014). Landscape types and pH control organic matter  
644 mediated mobilization of Al, Fe, U and La in boreal catchments. *Geochimica et*  
645 *Cosmochimica Acta*, 135, 190-202.

646 Krám, P., Hruška, J., Wenner, B. S., Driscoll, C. T., & Johnson, C. E. (1997). The  
647 biogeochemistry of basic cations in two forest catchments with contrasting lithology in the  
648 Czech Republic. *Biogeochemistry*, 37(2), 173-202.

649 Likens, G. E., & Buso, D. C. (2006). Variation in streamwater chemistry throughout the Hubbard  
650 Brook Valley. *Biogeochemistry*, 78(1), 1-30.

651 Likens, G. E., Bormann, F. H., Johnson, N. M., Fisher, D. W., & Pierce, R. S. (1970). Effects of  
652 forest cutting and herbicide treatment on nutrient budgets in the Hubbard Brook watershed-  
653 ecosystem. *Ecological monographs*, 40(1), 23-47.

654 Lin H. 2006. Temporal stability of soil moisture spatial patterns and subsurface preferential flow  
655 pathways in the Shale Hills catchment. *Vadose Zone Journal* 5: 317-340. DOI:  
656 10.2136/vzj2005.0058

657 Lin HS, Kogelmann W, Walker C, Bruns MA. 2006. Soil moisture patterns in a forested  
658 catchment: A hydrogeological perspective. *Geoderma* 131: 345-368. DOI:  
659 10.1016/j.geoderma.2005.03.013

660 Maher, K. (2011). The role of fluid residence time and topographic scales in determining  
661 chemical fluxes from landscapes. *Earth and Planetary Science Letters*, 312(1), 48-58.

662 McClain, M. E., Boyer, E. W., Dent, C. L., Gergel, S. E., Grimm, N. B., Groffman, P. M., ... &  
663 Pinay, G. (2003). Biogeochemical hot spots and hot moments at the interface of terrestrial  
664 and aquatic ecosystems. *Ecosystems*, 6(4), 301-312.

665 McGlynn BL, McDonnell JJ. 2003a. Role of discrete landscape units in controlling catchment  
666 dissolved organic carbon dynamics. *Water Resources Research* 39: 1090-1108.  
667 DOI:10.1029/2002WR001525

668 McGlynn BL, McDonnell JJ. 2003b. Quantifying the relative contributions of riparian and  
669 hillslope zones to catchment runoff. *Water Resources Research* 39:1310-1330. DOI:  
670 10/1029/2003WE002091

671 Mulholland, P. J. (1992). Regulation of nutrient concentrations in a temperate forest stream:  
672 roles of upland, riparian, and instream processes. *Limnology and Oceanography*, 37(7),  
673 1512-152.

674 Neal, C., & Kirchner, J. W. (2000). Sodium and chloride levels in rainfall, mist, streamwater and  
675 groundwater at the Plynlimon catchments, mid-Wales: inferences on hydrological and  
676 chemical controls. *Hydrology and Earth System Sciences Discussions*, 4(2), 295-310.

677 Neal C., Smith C.J., Walls J., Billingham P., Hill S. and Neal M. (1990) Hydrogeochemical  
678 variations in Upper Hafren forest stream waters, mid-Wales. *Journal of Hydrology*. 116:185-  
679 200.

680 Neal, C., Fisher R., Smith C. J., Hill S., Neal M., Conway T., Ryland G. P. and Jeffrey H. A.  
681 (1992) The effects of tree harvesting on stream-water quality at an acidic and acid-sensitive  
682 spruce forested area: Plynlimon, mid-Wales. *Journal of Hydrology*, 135:305-319.

683 Neal, C., Robson A.J., Shand P., Edmunds W.M., Dixon A.J., Buckley D.K., Hill S., Harrow M.,  
684 Neal M., Wilkinson J. and Reynolds B. (1997) The occurrence of groundwater in the Lower  
685 Palaeozoic rocks of upland Central Wales. *Hydrology and Earth System Sciences*. 1:3-18.

686 Neal, C., Reynolds B., Neal M., Wickham H., Hill L. and Williams B. (2004) The impact of  
687 conifer harvesting on stream water quality: the Afon Upper Hafren, mid-Wales. *Hydrology  
688 and Earth System Sciences* 8:503-520.

689 Neal, C., Robinson M., Reynolds B., Neal M., Rowland P., Grant S., Norris D., Williams B.,  
690 Sleep Darren and Lawlor A. (2010) Hydrology and water quality of the headwaters of the  
691 River Severn: Stream acidity recovery and interactions with plantation forestry under an  
692 improving pollution climate. *Science of the Total Environment*. 408:5035-5051.

693 Neal, C., Reynolds B., Norris D., Kirchner J. W., Neal M., Rowland P., Wickham H., Harman S.,  
694 Armstrong L., Sleep D., Lawlor A., Woods C., Williams B., Fry M., Newton G. and Wright  
695 D. (2011) Three decades of water quality measurements from the Upper Severn experimental  
696 catchments at Plynlimon, Wales: and openly accessible data resource for research, modelling,  
697 environmental management and education. *Hydrological Processes*. DOI: 10.1002/hyp.8191.

698 Neal, C., Reynolds B., Rowland P., Norris D., Kirchner J. W., Neal M., Sleep D., Lawlor A.,  
699 Woods C., Thacker S., Guyatt H., Vincent C., Hockenhull K., Wickham H., Harman S. and  
700 Armstrong L. (2012) High-frequency water quality time series in precipitation and  
701 streamflow: From fragmentary signals to scientific challenge. *Science of the Total  
702 Environment*. 434:3-12.

703 Neal, C., Kirchner, J., Reynolds, B. (2013a) Plynlimon research catchment hydrochemistry.  
704 NERC-Environmental Information Data Centre. DOI: 10.5285/44095e17-43b0-45d4-a781-  
705 aab4f72da025

706 Neal, C., Kirchner, J., Reynolds, B. (2013b) Plynlimon research catchment high-frequency  
707 hydrochemistry data. NERC-Environmental Information Data Centre. DOI:  
708 10.5285/551A10ae-b8ed-4ebd-ab38-033dd597a374

709 Pringle, C. M. (2001). Hydrologic connectivity and the management of biological reserves: a  
710 global perspective. *Ecological Applications*, 11(4), 981-998.

711 Reynolds, B. (1982) Preliminary studies on the laboratory simulation of nutrient transfers within  
712 a podzol soil. Bangor Occasional Paper No. 12, Bangor, UK.

713 Reynolds, B., Hornung M. and Stevens P.A. (1987) Solute budgets and denudation rate estimates  
714 for a mid-Wales catchment. *Catena* 14, 13-23.

715 Reynolds, B., Neal C., Hornung M., Hughes S. and Stevens P.A. (1988) Impact of afforestation  
716 on the soil solution chemistry of stagnopodzols in mid-Wales. *Water, Air and Soil Pollution*.  
717 38:55-70.

718 Reynolds, B., Wood M. J., Truscott A. M., Brittain S. A. and Williams D. L. (2000) Cycling of  
719 nutrient base cations in a twelve year old Sitka spruce plantation in upland mid-Wales.  
720 *Hydrology and Earth System Sciences*. 4:311-321.

721 Rudeforth C.C., Hartnup R., Lea J.W., Thompson T.R.E. and Wright P.S. (1984) Soils and their  
722 use in Wales. *Soil Survey of England and Wales Bulletin* No. 11, Harpenden, UK. 336 pp.

723 Shand P, Abesser C., Farr G., Wilton N., Lapworth D.J., Gooddy D.C., Haria A. and Hargreaves  
724 R. 2005a. Baseline report series: 17. The Ordovician and Silurian meta-sedimentary aquifers  
725 of central and south-west Wales. British Geological Survey Commissioned Report



726 CR/04/03/034N.

727 Shand P., Haria A.H., Neal C., Griffiths K.J., Gooddy D.C., Dixon A.J., Hill T., Buckley D.K.  
728 and Cunningham J.E. 2005b. Hydrochemical heterogeneity in an upland catchment: further  
729 characterization of the spatial, temporal and depth variations in soils, streams and  
730 groundwaters of the Plynlimon forested catchment, Wales. *Hydrology and Earth System*  
731 *Sciences*. 9:621-644.

732 Shand P., Darbyshire D.P.F., Gooddy D. and Haria A. 2007.  $^{87}\text{Sr}/^{86}\text{Sr}$  as an indicator of  
733 flowpaths and weathering rates in the Plynlimon experimental catchments, Wales, U.K.  
734 *Chemical Geology* 236:247-265.

735 Shand P., Darbyshire D.P.F., Love A.J. and Edmunds W.M. 2009. Sr isotopes in natural waters:  
736 Applications to source characterization and water-rock interaction in contrasting landscapes.  
737 *Applied Geochemistry*. 24:574-586.

738 Smith L.A. (2013) Aboveground carbon distribution across a temperate watershed. MS Thesis,  
739 The Pennsylvania State University.

740 Sivapalan M. 2005. Pattern, process and function: elements of a unified theory of hydrology at  
741 the catchment scale. *In Encyclopedia of Hydrological Sciences*, Anderson MG (ed). John  
742 Wiley & Sons, Inc.: Chichester, UK; 193–220.

743 Stallard R.F. and Murphy S.F. (2013) A unified assessment of hydrologic and biogeochemical  
744 responses in research watersheds in eastern Puerto Rico using runoff-concentration relations.  
745 *Aquat. Geochem.* DOI: 10.1007/A10498-013-9216-5.

746 Stevens P.A., Reynolds B., Hughes S., Norris D.A. and Dickinson A.L. (1997) Relationships  
747 between spruce plantation age, solute and soil chemistry in the Upper Hafren forest.  
748 *Hydrology and Earth System Sciences*. 1:627-637.

749 Takagi, K., & Lin, H. S. (2012). Changing controls of soil moisture spatial organization in the  
750 Shale Hills Catchment. *Geoderma*, 173, 289-302.

751 Thomas, E. M., Lin, H., Duffy, C. J., Sullivan, P. L., Holmes, G. H., Brantley, S. L., & Jin, L.  
752 (2013). Spatiotemporal patterns of water stable isotope compositions at the Shale Hills  
753 Critical Zone Observatory: Linkages to subsurface hydrologic processes. *Vadose Zone*  
754 *Journal*, 12(4).

755 Qu, Y., & Duffy, C. J. (2007). A semidiscrete finite volume formulation for multiprocess  
756 watershed simulation. *Water Resources Research*, 43(8).

757 Vitousek, P. M. (1977). The regulation of element concentrations in mountain streams in the  
758 northeastern United States. *Ecological Monographs*, 65-87.

759 Zimmer MA, Bailey SW, McGuire KJ, Bullen TD. 2012. Fine scale variations of surface water  
760 chemistry in ephemeral and perennial drainage networks. *Hydrological Processes* 27: 3438-  
761 2451. DOI:10.1002/hyp.9449

762

Table 1. Soil profile descriptions and associated SOC (% wt.) and DOC (mM) averages

Site	Category	Horizon	Depth (cm)	DOC (mM)	SOC <sup>b</sup> (% wt.)
<b>Shale Hills<sup>c</sup></b>					
Planar, valley	Mineral	A	0-15	0.69	4.7
	Mineral	Bw	15-30	0.54	1.6
		Bt	30-53	--	--
		C	53-147	--	--
Swale, valley	Organic	A	0-11	1.28	2.0
	Mineral	Bw	11-38	0.55	1.2
		BC	38-60	--	--
		C	60-178	--	--
<b>Plynlimon<sup>d</sup></b>					
Peat	Organic	O	0-100+	1.10	40-50
Peaty gley	Organic	O	0-22	1.37	25
	Mineral	Eag	22-37	0.49	5-6
		Bs	37-86	--	--
		C	86-107	--	--
Stagnopodzol	Organic	O	0-19	1.12	46
	Mineral	Eag	19-24	0.35	5
		Bs	24-51	--	--
		C	51-89	--	--

<sup>a</sup> Category (organic or mineral) assigned to soil horizons in this paper; Note that reported SOC and DOC values are averaged over all mineral horizons

<sup>b</sup> Shale Hills SOC estimates from Jin et al. (2010) and Andrews (2011)

<sup>c</sup> Shale Hills soil descriptions from Lin (2006).

<sup>d</sup> Plynlimon soil descriptions from Ruderforth et al. (1984).

Table 2. Element concentrations ( $\pm$  standard error) in water (precipitation, soil pore water, stream water, groundwater) and vegetation (green leaves, leaf litter) averaged over all available data collected from the Susquehanna Shale Hills Critical Zone Observatory in Pennsylvania, USA between 2006 – 2011

	<b>pH</b>	<b>Na</b>	<b>K</b>	<b>Mg</b>	<b>Ca</b>	<b>Si</b>	<b>Fe</b>	<b>Mn</b>	<b>DOC</b>
		----- $\mu\text{mol L}^{-1}$ -----							<b>mmol L<sup>-1</sup></b>
<b>Precipitation</b>	4.5	2.1	0.67	0.73	2.65	--	0.24	0.05	0.08 <sup>a</sup>
$\pm$ std. err.	0.2	0.4	0.28	0.16	0.51	--	0.14	0.01	0.02
<b>Soil Pore Water</b>									
SPVF, A horizon	4.5	29	34	62	142	123	0.61	0.38	0.69
$\pm$ std. err.	0.2	3	3	10	26	5	0.15	0.06	0.06
SPVF, B horizon	4.7	34	29	89	146	125	0.25	0.44	0.54
$\pm$ std. err.	< 0.1	2	1	8	15	3	0.02	0.05	0.04
SSVF, A horizon	3.9	21	88	39	72	112	1.7	6.8	1.28
$\pm$ std. err.	0.1	3	14	5	11	11	0.26	1.9	0.33
SSVF, B horizon	4.4	28	29	79	35	126	0.21	1.3	0.55
$\pm$ std. err.	< 0.1	1	1	2	1	2	0.03	0.1	0.03
Average, all soils		28	45	68	99	121	0.69	2.2	0.77
RSD (%), all soils		19	64	33	55	5	100	140	46
<b>Stream Water</b>									
Low flow	6.3	39	50	162	450	108	14	5.2	0.81
$\pm$ std. err.	< 0.1	< 1	1	3	10	1	2	0.7	0.06
Moderate flow	5.7	32	31	122	240	101	1.9	1.1	0.45
$\pm$ std. err.	0.1	1	< 1	3	7	1	0.4	0.2	0.03
High flow	5.9	26	29	96	163	99	0.53	0.33	0.42
$\pm$ std. err.	0.1	< 1	< 1	2	6	1	0.02	0.03	0.03
<b>Groundwater</b>	--	145	24	404	758	124	0.14	2.9	0.20
$\pm$ std. err.	--	6	< 1	13	27	1	0.08	0.2	0.05
<b>Vegetation</b>									
		----- $\mu\text{mol g}^{-1}$ -----							
Leaf litter		n/a	69	59	263	n/a	n/a	49	
$\pm$ std. err.			6	3	15			3	
Green leaves		1.1	221	69	186		0.99	42	
$\pm$ std. err.		0.2	15	7	18		0.06	4	

<sup>a</sup>Andrews et al. (2011)

Table 3. Solute concentrations ( $\pm$  standard error) averaged over all available data collected from the Upper Hore (pre-harvest: 1983 – 2005; post-harvest: 2005 – 2010) and Upper Hafren (1990 – 2010) catchments in the Plynlimon forest in Wales, United Kingdom

	pH	Na	K	Mg	Ca	Si	Fe	Mn	Al	DOC
		----- $\mu\text{mol L}^{-1}$ -----								$\text{mmol L}^{-1}$
<b>Precipitation</b>	4.98	69.6	2.60	8.52	6.89	2.63	0.19	0.03	0.66	0.045
$\pm$ std. err.	0.01	3	0.10	0.32	0.59	0.35	0.02	< 0.01	0.05	0.001
<b>Soil Pore Water</b>										
Peat	3.24	143	5.55	27.1	12.6	9.40	2.97	0.16	2.22	1.10
$\pm$ std. err.	0.10	4	0.54	1.2	0.5	0.38	0.15	0.02	0.20	0.45
Peaty gley organic	3.56	239	29.7	30.0	8.32	47.6	9.29	---	26.6	1.37
$\pm$ std. err.	0.07	6	3.2	1.0	0.41	2.0	0.42	---	1.3	0.77
Peaty gley mineral	4.16	220	10.2	26.7	5.32	57.1	12.2	---	55.9	0.49
$\pm$ std. err.	0.01	1	0.2	0.1	0.02	0.3	0.5	---	0.2	0.09
Stagnopodzol O	3.82	165	25.7	29.8	22.7	18.9	4.25	0.27	8.00	1.12
$\pm$ std. err.	0.07	5	2.3	1.1	1.4	0.7	0.20	0.04	0.38	0.04
Stagnopodzol min	4.12	159	7.91	19.9	11.7	33.7	0.96	0.47	19.9	0.35
$\pm$ std. err.	0.01	< 1	0.04	< 0.1	0.04	0.2	0.02	0.01	0.2	< 0.01
<b>Upper Hore</b>										
Low flow	6.98	177	3.06	36.1	78.3	63.0	2.59	0.32	3.56	0.10
$\pm$ std. err.	0.02	1	0.06	0.3	1.0	0.9	0.11	0.01	0.40	< 0.01
Moderate flow	6.01	180	3.11	29.8	39.7	48.4	2.02	0.42	7.44	0.12
$\pm$ std. err.	0.02	1	0.05	0.1	0.5	0.3	0.04	< 0.01	0.12	< 0.01
High flow	4.75	184	3.78	25.6	17.5	30.4	2.08	0.40	16.2	0.21
$\pm$ std. err.	0.02	3	0.11	0.5	0.4	0.5	0.07	0.01	0.3	< 0.01
Post-harvest, all flows	5.96	169	8.9	30.2	41.3	45.0	3.33	0.51	7.9	0.31
$\pm$ std. err.	0.05	1	0.3	0.3	0.7	0.9	0.11	0.01	0.3	0.01
<b>Upper Hafren</b>										
Low flow	6.45	160	2.96	27.8	17.4	67.8	1.21	0.27	2.02	0.08
$\pm$ std. err.	0.02	1	0.06	0.2	0.1	1.0	0.06	0.01	0.18	< 0.01
Moderate flow	5.79	153	3.52	26.0	13.8	57.7	1.74	0.38	3.51	0.13
$\pm$ std. err.	0.02	< 1	0.06	0.1	0.08	0.5	0.04	< 0.01	0.09	< 0.01
High flow	4.85	142	5.04	22.8	10.1	34.5	2.68	0.34	7.44	0.24
$\pm$ std. err.	0.02	2	0.15	0.4	0.2	0.8	0.12	0.01	0.22	0.01
<b>Groundwater</b>	5.26	227	8.7	79	94	86	5.9	2.54	1.86	0.032
$\pm$ std. err.	0.07	8	1.7	11	20	4	1.5	0.33	0.11	0.002


Table 4. Slopes of regression lines fit to C-Q data ( $\log C = a + m * \log Q$ ) indicate chemostatic<sup>a</sup>, non-chemostatic<sup>b</sup>, or dilution<sup>c</sup> concentration-discharge patterns

	<b>Na</b>	<b>K</b>	<b>Ca</b>	<b>Mg</b>	<b>Si</b>	<b>Al</b>	<b>Fe</b>	<b>Mn</b>	<b>DOC</b>
<b>Shale Hills (2008 – 2010)</b>									
Slope ( <i>m</i> )	-0.04 <sup>a</sup>	-0.07 <sup>a</sup>	-0.12 <sup>b</sup>	-0.05 <sup>a</sup>	-0.01 <sup>a</sup>	-	-0.30 <sup>b</sup>	-0.19 <sup>b</sup>	-0.11 <sup>b</sup>
Std. Err.	< 0.01	< 0.01	0.01	< 0.01	< 0.01	-	0.01	0.02	0.02
R <sup>2</sup>	0.32	0.56	0.43	0.30	0.09	-	0.58	0.24	0.14
<b>Upper Hafren (1990 – 2010)</b>									
Slope ( <i>m</i> )	-0.05 <sup>a</sup>	0.18 <sup>b</sup>	-0.20 <sup>c</sup>	-0.08 <sup>a</sup>	-0.26 <sup>c</sup>	0.58 <sup>b</sup>	0.33 <sup>b</sup>	0.10 <sup>b</sup>	0.42 <sup>b</sup>
Std. Err.	0.00	0.01	0.00	0.00	0.01	0.02	0.02	0.01	0.02
R <sup>2</sup>	0.20	0.19	0.64	0.27	0.67	0.53	0.26	0.08	0.37
<b>Upper Hore (pre-harvest; 1983 – 2004)</b>									
Slope ( <i>m</i> )	0.00 <sup>a</sup>	0.07 <sup>b</sup>	-0.51 <sup>c</sup>	-0.12 <sup>a</sup>	-0.27 <sup>c</sup>	0.52 <sup>b</sup>	-0.05 <sup>a</sup>	0.05 <sup>b</sup>	0.26 <sup>b</sup>
Std. Err.	0.00	0.01	0.01	0.00	0.01	0.01	0.02	0.01	0.02
R <sup>2</sup>	0.00	0.04	0.89	0.43	0.75	0.59	0.01	0.04	0.23
<b>Upper Hore (post-harvest; 2005 - 2010)</b>									
Slope ( <i>m</i> )	-0.04 <sup>a</sup>	0.26 <sup>b</sup>	-0.49 <sup>c</sup>	-0.11 <sup>a</sup>	-0.23 <sup>c</sup>	0.42 <sup>b</sup>	-0.04 <sup>a</sup>	0.12 <sup>b</sup>	0.25 <sup>b</sup>
Std. Err.	0.00	0.03	0.01	0.01	0.01	0.02	0.03	0.02	0.03
R <sup>2</sup>	0.12	0.18	0.90	0.50	0.83	0.68	< 0.01	0.16	0.21


<sup>1</sup>Calculations exclude data where concentrations fell below the detection limit.

<sup>2</sup>All slopes are statistically different from zero ( $p < 0.001$ ) except Na (pre-harvest) and Fe (post-harvest) in the Upper Hore


**Figure 1.** Map views of the Susquehanna Shale Hill Critical Zone Observatory (Shale Hills, PA, USA; left) and Plynlimon (Wales, UK; right) catchments. Symbols mark locations of precipitation (black square), stream water (red circle), pore water (black triangle), and groundwater (green circle) samplers. Shading delineates major landscape features that are organic-rich or organic-poor in each catchment: swale (brown) versus planar (light green) slopes at Shale Hills or peat (tan) versus forested (dark green) regions at Plynlimon. Notably, the most organic-rich soils are in lowlands in Shale Hills but uplands in Plynlimon; consequently, inputs from organic-rich soils dominate stream flow under low-flow conditions in Shale Hills but high-flow conditions in Plynlimon.



**Figure 2.** Log  $C$  (solute concentration) versus log  $Q$  (discharge) in the Shale Hills catchment (a, b) and two Plynlimon subcatchments, the Upper Hafren (c-e) and Upper Hore (f-h). Data from the post-harvest period (2005 – 2010) in the Upper Hore are plotted in separate panels (i-k). For each catchment, the left panel shows elements that exhibit chemostatic behavior, the middle panel shows non-chemostatic elements that exhibit behavior similar to DOC (e.g. dilution at Shale Hills and enrichment at Plynlimon), and the right panel shows elements that exhibit dilution behavior.



**Figure 3.** Molar ratios of major divalent (Ca:Mg) versus univalent (K:Na) cations are plotted on the left and the molar ratios of Mn (mmol) to Mg (mol) versus dissolved organic carbon (mmol L<sup>-1</sup>) are plotted on the right for solute source pools in the forested Shale Hills (a, b), peatland Upper Hafren (c, d), and predominately forested Upper Hore (e, f) catchments. Arrows indicate direction of increasing discharge for average stream chemistries. Values for total C (~ 33 mmol kg<sup>-1</sup>) and K:Na (~60) in leaves are divided by 10 to fit on plots (a) and (b). For soil pore water, filled symbols indicate organic-rich waters and open symbols indicate organic-poor waters.





**Figure 4.** The degree of non-chemostatic behavior for a solute in stream water, denoted by  $m_j$ , was correlated with the relative concentration of that solute in organic-rich soil water in each catchment. Specifically,  $m_j$  was (a) negatively correlated with the ratio of solute concentrations in organic-rich versus organic-poor soil waters at Shale Hills ( $R^2 = 0.90$ ,  $p < 0.001$ ) and positively correlated with the ratio of solute concentrations in pore waters versus precipitation in (b) the Upper Hafren ( $R^2 = 0.48$ ;  $p < 0.05$ ) and (c) the Upper Hore ( $R^2 = 0.42$ ;  $p < 0.05$ ). Error bars represent the standard error of each value and are smaller than the symbol where not visible.

