

1 Dear editor, dr. Ehret,  
2 thank you for additional comments on our paper. Apart from improvements regarding your  
3 comments, we also slightly improved the method of detection the events (Sect. 2.1) so the  
4 values of the precipitation indices have changed a bit. The revised paper is provided in the  
5 required form; our responses to your comments are as follows:

6

7 *"I agree with the referees that the manuscript needs more focus, which you have indicated to  
8 put on the definition and application of the three indices. This is good, but if you do so, you  
9 should also, as proposed by the referees, provide an identification of extreme rainfall and  
10 flood events based on established 'benchmark' approaches, and compare the so-found rainfall  
11 and flood extremes in the manner you do for your indices. This should help to clarify the Pros  
12 and Cons of your approach and help readers to decide when to apply your approach (and  
13 when not)."*

14 We compared the WEI with two standard methods, the WAI with an index which was  
15 recently published, and the FEI with two indices mentioned in the review at the beginning of  
16 the paper. One person enlarged the team in this context. The results are presented in three new  
17 Figs. 9, 10, and 11 and broadly discussed in Sect. 4.1.

18

19 *"Page 292, line 13 pp: Like referee #1, I also had a hard time to understand the method from  
20 your explanation, even the revised one. Please rework this section again."*

21 The last paragraph in Sect. 2.4 was reworked; the original text was unclear because it  
22 included also the technic of the so called "mean day" which is not presented in the paper.

23

24 *"Page 293, line 14-15 (comment by referee #1): I think your answer has not sufficiently  
25 addressed the question of referee #1. I think he asked what conclusions i) about the  
26 agreement of rainfall and flood extremes in general and ii) about the applicability of your  
27 indices to identify them can be drawn from e.g. a 75% overlap of rainfall and flood extremes,  
28 and why."*


29 We have recognized that our statement (that the applicability of our indices can be proved  
30 from the overlap) is doubtful; therefore we had removed the sentence in the first improved

1 version of the text. Regarding the general agreement of extremes, we have enlarged the last  
2 paragraph in Sect. 3.1.

3

4 “*Referee #2 raises the issue of how the use of human-influenced flood hydrographs  
5 compromises the results and conclusions of your study. This is an important issue, which you  
6 have addressed in your reply. While I agree that you cannot avoid using such data (as there  
7 are not too many uninfluenced rivers left), you need to go into more detail here and give proof  
8 that the use of these data does not significantly alter the results of your study. E.g. estimate  
9 what the degree of uncertainty of the flood recurrence intervals induced by human inference  
10 is and how much this alters the FEI's and the comparison of WEI's and FEI's. As cited by you,  
11 Buchtele (1972) states that human influence becomes negligible for larger-than-20-year-  
12 floods. However, in your study you work with 50 events selected from 50 years (Sect. 3.1),  
13 which clearly includes less rare events.”*

14 In fact, approximately a half of stations used in the study are not influenced by reservoirs but  
15 their catchments are usually rather small. Anyway, we have computed another version of the  
16 FEI (FEIu) with encouraging results, see the figure below. A paragraph about it was added at  
17 the end of Sect. 2.3.



18

1

2 “When comparing the FEI and WEI values in Section 2.4, it is not clear to me whether the  
3 ranges of values that WEI, WAI and FEI can attain are identical or just comparable. If the  
4 latter is the case (which I think), I suggest not taking the ratio of the values, but compare the  
5 ranks of FEI and WEI from identical periods of time instead (e.g. take the difference of the  
6 ranks). A higher rank of FEI than WEI of a particular event then indicates an event that is  
7 more extreme wrt flood than rainfall.”

8 We agree that distributions of 50 maximum values of the three indices are not completely  
9 identical. Therefore, we compared the events with respect to their rankings as you have  
10 suggested (Fig. 12, originally Fig. 9). Nevertheless, values of the ratios FEI/WEI and  
11 FEI/WAI remained in Fig. 6 because it enables to determine the FEI also in case of  
12 precipitation events which did not resulted into an EFE.

13

14 “A final suggestion for further studies, which you already may have thought of: Quantify the  
15 extremity of antecedent conditions in the same manner as WEI and FEI. This gives you, in the  
16 framework you have built, additional possibilities to analyze the causes and propagation of  
17 ‘extremity’ of events.”

18 Thank you for your recommendation; we have already mentioned antecedent saturation in the  
19 present paper and we plan to employ it even more in the subsequent study.

20

21

22 Responses to reviewer’s comments are not included in this file because they are available in  
23 the online discussion. The marked-up text follows:

24

25

26

27

28

# Novel indices for the comparison of precipitation extremes and floods; an example from the Czech territory

**Odstraněno:** Evaluation  
**Odstraněno:** of  
**Odstraněno:** and comparison between their temporal distributions

M. Müller<sup>1,2</sup>, M. Kašpar<sup>1</sup>, A. Valeriánová<sup>3</sup>, L. Chrlová<sup>3</sup>, E. Holtanová<sup>3</sup>, and B. Gvoždíková<sup>2</sup>

**Odstraněno:** and

[1]{Institute of Atmospheric Physics AS CR, Prague, Czech Republic}

[2]{Charles University, Faculty of Science, Prague, Czech Republic}

[3]{Czech Hydrometeorological Institute, Prague, Czech Republic}

Correspondence to: M. Müller (muller@ufa.cas.cz)

**Odstraněno:** We proposed three analogous extremity indices based on the estimated return periods at individual sites and spatial averaging of the values; we optimized both the areal extent and the duration of individual events. The weather extremity index (WEI), the weather abnormality index (WAI), and the flood extremity index (FEI) were applied to the original precipitation data, the seasonally transformed precipitation data, and the runoff data to identify extreme precipitation events (EPEs), abnormal precipitation events (APEs), and extreme flood events (EFEs), respectively. We present 50 events of each type from the period of 1961–2010 in the Czech Republic and

**Odstraněno:** their

## Abstract

The paper presents three indices for evaluation of hydrometeorological extremes, considering them as areal precipitation events and trans-basin floods. In contrast to common precipitation indices, the weather extremity index (WEI) reflects not only the highest precipitation amounts at individual gauges but also the rarity of the amounts, the size of the affected area, and the duration of the event. Besides that, the aspect of precipitation seasonality was considered when defining the weather abnormality index (WAI) which enables the detection of precipitation extremes throughout the year. The precipitation indices were complemented with the flood extremity index (FEI) employing peak discharge data. A unified design of the three indices, based on return periods of station data, enables to compare easily inter-annual and seasonal distributions of precipitation extremes and large floods.

The indices were employed in evaluation of 50 extreme precipitation events, seasonally abnormal precipitation events, and large floods 1961–2010 in the Czech Republic. A preliminary study of discrepancies among historic values of the indices indicates that variations in the frequency and/or magnitude of floods can generally be due not only to variations in the magnitude of precipitation events but also to variations in their seasonal distribution and other factors, primarily the antecedent saturation. The events could be further studied with respect to circulation conditions, for example, by the circulation extremity index.

**Odstraněno:** Most of EFEs were produced by an EPE in warmer half-years, whereas fewer than half of the EFEs were produced by an APE in the remainders of the years because thawing can substantially enhance the discharge at those times. Most significant EPEs occurred in July and the first half of August, although their hydrological responses were also significantly influenced by the antecedent saturation and other factors. As a result, the accumulation of precipitation extremes during the 1977–1986 period produced less significant flooding than another accumulation after 1996. In general, the primary discrepancies between the magnitudes of EPEs and EFEs occurred in May and September, when consequent floods were usually much larger and smaller in relation to the WEI, respectively. The hydrological response to APEs was usually strong in December, whereas another accumulation of EFEs in March was usually not due to APEs. Neither precipitation nor flood extremes occurred from early April through early May. This

**Odstraněno:** confirms

**Odstraněno:** differences

## 1 1 Introduction

2 Precipitation is extensively studied due to its impacts on the hydrology, geomorphology, and  
3 economy of a given region. Precipitation extremes are of special interest because impacts  
4 rapidly increase with the precipitation extremity. There are three main concepts of extremity:  
5 severity, intensity, and rarity (Stephenson, 2008). Evaluation of the extremity of past events  
6 enables to determine return periods of heavy precipitation and to estimate the probable  
7 maximum precipitation (Řezáčová, 2005b). Currently, the main challenge in precipitation  
8 climatology is understanding past and possible future changes in the frequency and/or  
9 magnitude of precipitation extremes (e.g., Alexander et al., 2006). These changes could alter  
10 the frequency and/or magnitude of consequent floods. To properly assess this linkage, we  
11 need to define quantitative indices to investigate the coupling between precipitation extremes  
12 and flood events.

13 Floods are frequently evaluated by their severity, which can be defined as the amount of  
14 socioeconomic loss or number of casualties. For example, Barredo (2007) selected 23 flash  
15 floods and 21 river floods during the period of 1950–2005 in Europe based on two criteria:  
16 losses exceeding 0.005 % of EU GDP and/or more than 70 casualties. In contrast, severity can  
17 hardly be used for evaluation of precipitation events because the damage is usually produced  
18 not directly by the precipitation but by subsequent phenomena (floods, landslides, etc.).  
19 Moreover, factors of exposure and vulnerability can produce serious discrepancies between  
20 causes and consequences.

21 The concept of intensity better corresponds to physical causes; thus, it is more convenient for  
22 comparison between precipitation extremes and floods, Rodier and Roche (1984) and lately  
23 Herschy (2003) assessed the world's maximum floods with respect to their maximum  
24 instantaneous discharges. To compare the extremity of floods on various rivers, they used the  
25 Francou index, which normalizes the common logarithm of maximum discharge by the  
26 common logarithm of the catchment area. Not surprisingly, maximum floods were located in  
27 the雨iest regions.

28 The standard approach to the evaluation of precipitation intensity is to search data series from  
29 individual gauges using commonly accepted indices (Zhang et al., 2011), mainly those  
30 defined by the Expert Team on Climate Change Detection and Indices (ETCCDI). According  
31 to ETCCDI, precipitation extremes can be defined as days with maximum one-day ( $P_1$ ) or  
32 five-day ( $P_5$ ) precipitation total in a period. Nevertheless, the duration of events can vary

**Odstraněno:** we need to recognize various aspects of the relationship between extreme precipitation and flooding events

**Odstraněno:** There are many ways to evaluate precipitation and flooding depending primarily on the chosen concept of extremity. In general, three main concepts of weather extremes can be distinguished: severity, intensity, and rarity (Stephenson, 2008). ¶ Extreme flood events (EFEs)

**Odstraněno:** extreme

**Odstraněno:** (EPEs)

**Odstraněno:** seems to be the more promising for our purposes

**Odstraněno:** EPEs

**Odstraněno:** then

**Odstraněno:** most simply

**Odstraněno:** a

**Odstraněno:** 1

**Naformátováno:** Písmo: není Kurzíva

**Naformátováno:** Písmo: není Kurzíva

**Odstraněno:** ( $P_d$ ) exceeding a threshold at any gauge in the region. This approach was also used in classic hydro-meteorological works in search of EPEs in the Czech Republic. Štekla et al. (2001) studied a set of days with  $P_d \geq 150$  mm and their synoptic causes, seasonality, temporal distribution and other aspects. Based on this approach, the maximum Czech EPE occurred on 29 July 1897 when a  $P_d$  of 345.1 mm was measured at the Nová Louka gauge in the Jizerské Hory Mts. Indeed, this event was the maximum  $P_d$  recorded in Central Europe (Munzar et al., 2011). ¶

1 widely, and the precipitation intensity usually fluctuates during the event. Begueria et al.  
2 (2009) partly took account of this fact; they used declustering of daily precipitation totals to  
3 distinguish individual precipitation events and characterized them not only by magnitude and  
4 duration but also by peak intensity.

5 Even more important is the spatial aspect of precipitation events. They always affect a certain  
6 area; thus, precipitation extremes should be considered to be “regional events” (Ren et al.,  
7 2012). The latter approach is necessary if the intensity of precipitation and floods is to be  
8 compared, because the size of the affected area influences the hydrological response. In our  
9 previous paper (Kašpar and Müller, 2008), we used the concept of areal precipitation intensity  
10 and evaluated precipitation events based on the weighted average of daily areal precipitation  
11 totals on three consecutive days. Nevertheless, Konrad (2001) demonstrated that the extremity  
12 of an event depends on the size of the considered region. As a result, the areal average  
13 disadvantages events that were violent but affected only a part of the region over which the  
14 mean is taken (e.g., an administrative unit, a catchment).

**Odstraněno:** Moreover

**Odstraněno:**,

**Odstraněno:** precipitation

**Odstraněno:** s

15 Based on the concept of intensity, extreme events occur mainly in regions that are prone to  
16 heavy rains (in the Czech Republic, such regions are along the northern state border because  
17 of the orographic precipitation enhancement). In order to reflect regional climatic differences,  
18 the concept of rarity is applied; extreme precipitation and floods can thus be detected in the  
19 whole studied region. The concept is frequently used with regard to floods, and the intensity  
20 (magnitude of the peak flow) is usually compared with return levels. If a set of extreme floods  
21 is studied, they are defined as discharges with return periods exceeding a threshold.  
22 Nevertheless, Uhlemann et al. (2010) noted that flood events frequently affect several  
23 independent catchments and introduced the concept of trans-basin floods. We adopted and  
24 adapted this approach to our data because we compared flood extremity with precipitation,  
25 which also affects more than one catchment at a given time.

**Odstraněno:** In our previous paper  
(Kašpar and Müller, 2008), we used the  
concept of areal precipitation intensity and  
evaluated EPEs based on the weighted  
average of daily areal precipitation totals on  
three consecutive days.

**Odstraněno:** only

**Odstraněno:** them

**Odstraněno:** However, extreme  
precipitation and floods can occur  
anywhere on the Earth if a more-inclusive

26 To enable this comparison, we propose indices analogous to each other that are based on the  
27 point estimates of return periods of precipitation totals and peak discharges and on spatial  
28 averaging of the values (Sects. 2.1 and 2.3). The method is further enriched by the aspect of  
29 precipitation abnormality with respect to the season (Sect. 2.2). We demonstrate the method  
30 using data from the Czech Republic and present three sets of events: precipitation extremes  
31 regardless and regarding of the season and extreme floods (Sect. 3.1). These sets are further

**Odstraněno:** ion

**Odstraněno:** at individual sites

1 compared with regard to their inter-annual (Sect. 3.2) and seasonal (Sect. 3.3) distribution.  
2 The results obtained are discussed in Sect. 4.

3

## 4 **2 Proposed methods**

5 We proposed three ~~extremity~~ indices that enable to compare the temporal distribution of  
6 ~~precipitation extremes and floods. The indices are based on point return period estimates of~~  
7 ~~precipitation totals and peak flows, respectively, spatial~~ averaging of their values, and  
8 optimizing the areal extent and the duration of individual events.

**Odstraněno:** analogous

**Odstraněno:** various types of extreme events

**Odstraněno:** The indices are based on the estimated return periods at individual sites, spatial

### 9 **2.1 Evaluation of precipitation extremity**

10 The weather extremity index (WEI) ~~presented in detail by Müller and Kašpar (2014)~~ ~~was~~  
11 ~~employed in searching for extreme precipitation events (EPEs).~~ The WEI is based on return  
12 periods of ~~1-day to several-day~~ precipitation totals at individual sites. We used data from 711  
13 rain gauges with data series at least 20 years long between 1961 and 2010. The return periods  
14 were assessed using the generalized extreme value (GEV) distribution (Hosking and Wallis,  
15 1997) because it was confirmed to be a suitable model for precipitation extremes in most  
16 regions of the Czech Republic (Kyselý and Picek, 2007). The GEV distribution was applied  
17 as the parametric model for annual maxima of the totals. Parameters of the GEV distribution  
18 were estimated by means of the L-moment algorithm (Hosking and Wallis, 1997) and the  
19 region-of-influence (ROI) method (Burn, 1990). The ROI method is based on ‘homogenous  
20 regions’, in which all regional data, weighted by a dissimilarity measure, are used for  
21 estimating parameters of the distribution of extremes at the site of interest. Although the  
22 application of the ROI method makes the estimations more robust than local analyses (Kyselý  
23 et al., 2011), we did not accept return periods longer than 1000 years. Instead, we set the  
24 return period to 1000 years.

**Odstraněno:** was

**Odstraněno:** 1-day to 5

**Naformátováno:** Písmo: Kurzíva

25 The next step in the method was the interpolation of return periods from gauges in a regular  
26 grid with a horizontal resolution of 2 km. (Experiments have demonstrated that the results are  
27 not affected by the resolution if it is constant across all studied events.) Because of the  
28 exponential nature of the GEV distribution from which the return periods are derived, we  
29 interpolated their common logarithms instead of pure return period values. We chose linear  
30 kriging as the interpolation method. Using an inversion transformation, we obtained cell  
31 values of  $N_{ti}$ , which represents the return period of the precipitation total accumulated during  $t$

1 days in a cell  $i$ . The cells were then sorted in decreasing order with respect to  $N_{ti}$  and  
2 considered within a stepwise increasing area  $a$  of  $n$  pixels each representing  $4 \text{ km}^2$ .

3 The WEI was calculated by maximizing the variable  $E_{ta}$ . This variable is defined as the  
4 common logarithm of the spatial geometric mean  $G_{ta}$  of the return periods  $N_{ti}$  multiplied by  
5 the radius  $R$  of a circle of the same area as the one over which the geometric mean is taken.  
6 This relationship can be expressed as

$$7 E_{ta} = \log(G_{ta})R = \frac{\sum_{i=1}^n \log(N_{ti})}{n} \frac{\sqrt{a}}{\sqrt{\pi}}. \quad (1)$$

8 The optimization of  $a$  was performed using a step-by-step enlarging of the area under  
9 consideration. The variable  $E_{ta}$  initially increased as we accumulated cells with lengthy return  
10 periods; once the return periods were insufficiently long in the added cells, the value of  $E_{ta}$   
11 started to decrease. When we chose a time window for which  $E_{ta}$  reached its maximum during  
12 the entire event, the respective maximum  $E_{ta}$  equaled the WEI. Then, we were also able to  
13 determine the affected area  $a$ , the duration  $t$ , and the respective geometric mean of return  
14 periods  $G_{ta}$  complying with the relationship  $E_{ta} = \text{WEI}$ . The method is presented in Fig. 1,  
15 which shows the EPE from May/June 2013, which was subsequently added to the study  
16 because of catastrophic flooding observed during this time (Šercl et al., 2013). Although the  
17 maximum return period at a site belonged to the 1-day total on 1 June (Horní Maršov, 130.3  
18 mm,  $> 1000$  yr), maximum  $E_{ta}$  gradually increased with increasing  $t$ . The WEI corresponded  
19 to the five-day period from 30 May to 3 June 2013.

20 Nevertheless, the time distribution of maximum  $E_{ta}$  values can be more complex. Figure 2  
21 presents such a case from August 2002, when a subsequent EPE followed the previous one  
22 after a break of only three days. In this case, two distinct maxima of  $E_{ta}$  enabled us to  
23 recognize independent EPEs and determine the durations of both events (two and three days,  
24 respectively). Adding an extra day would cause the  $E_{ta}$  value to decrease. Therefore, we did  
25 not consider a longer time window (7 days) though the  $E_{ta}$  would be even higher as the two  
26 EPEs would be aggregated. Moreover, we also decided to consider precipitation events of the  
27 length from 1 to 5 days only because the thresholds correspond with two main indices of  
28 precipitation extremes by ETCCDI (Zhang et al., 2011).

29 Figure 2 also shows that the extremity of precipitation with respect to the maximum  $E_{ta}$  can  
30 substantially differ from the areal mean of daily precipitation totals throughout the Czech

<b>Odstraněno:</b> t
<b>Odstraněno:</b> , when a much larger area was affected by heavy rains
<b>Odstraněno:</b> display a more
<b>Odstraněno:</b> pattern
<b>Odstraněno:</b> one or two more
<b>Odstraněno:</b> d
<b>Odstraněno:</b> if
<b>Odstraněno:</b> had also
<b>Odstraněno:</b> ed
<b>Odstraněno:</b> an $E_{ta}$ spanning
<b>Odstraněno:</b> ,
<b>Odstraněno:</b> then
<b>Odstraněno:</b> have been
<b>Odstraněno:</b> However, if we had also considered an $E_{ta}$ spanning a longer time window (7 days), then the two EPEs would have been aggregated.
<b>Odstraněno:</b> This example induces a question about values of the parameter $t$ because it obviously cannot remain unlimited. W
<b>Odstraněno:</b> Therefore, we
<b>Odstraněno:</b> limit $t$ to a value of equal or less than 5 throughout the study. This threshold appears to be appropriate for the Czech Republic but could be slightly higher if the method were applied to a larger area.

1 Republic. Although the mean was nearly twice as high on 11 August than on 6 August, the  
2 respective  $E_{ta}$  maxima were the same. The advantage of the concept of  $E_{ta}$  is that the  
3 considered area and the time window are “event-adjusted”. Although comparably heavy rains  
4 were limited to the southwestern Czech Republic on both days, weaker rains occurred only on  
5 the latter day throughout the whole country. These non-extreme precipitation totals increased  
6 the mean, whereas they were not included in the  $E_{ta}$ .

## 7 **2.2 Precipitation abnormality with respect to the season**

8 Both precipitation long-term means and extremes are not equally distributed among the  
9 seasons in most places on the Earth. In the Czech Republic, higher precipitation totals  
10 generally occur in summer (Tolasz et al., 2007). As a result, the WEI maxima are also  
11 concentrated in the summer. However, even smaller precipitation totals can be considered to  
12 be “extreme” when they occur in a season when they are rare. If precipitation extremes are  
13 defined as events significantly different from seasonally normal conditions, then they can  
14 occur throughout the year. Precipitation extremes of this type will be referred to as abnormal  
15 precipitation events (APEs). They were evaluated using the weather abnormality index  
16 (WAI), which has the same design as the WEI, although it is calculated based on seasonally  
17 standardized precipitation totals.

**Odstraněno:** during warmer half-years

18 The standardization of daily precipitation totals reflects their annual distribution. The mean,  
19 variance and skewness fluctuate significantly during the year (Fig. 3), and thus none of these  
20 parameters can be avoided in the process of standardization. The same is true for the kurtosis  
21 which is very closely correlated with the skewness (not depicted). Furthermore, means and  
22 standard deviations are also closely correlated. Therefore, our standardization method consists  
23 of removing fluctuations of the mean  $\mu_d$  and the skewness  $\gamma_d$  from the daily totals on  
24 individual calendar days  $d$ .

25 There are two types of fluctuations in the data. First,  $\mu_d$  and  $\gamma_d$  change significantly from day  
26 to day depending on the presence or absence of heavy precipitation episodes on a given  
27 calendar day during the period of 1961–2010. These random fluctuations have to be smoothed  
28 using a proper time filter. Monthly means are sometimes used for these purposes, but we have  
29 excluded this method as it produces artificial edges in the data. Moving averages are only  
30 slightly better from this point of view. We use the Gaussian filter because it is considered the  
31 ideal time domain filter (Blinchikoff and Zverev, 2001). We tested several data series to

**Odstraněno:** are  
**Odstraněno:**, but we preferred the  
**Odstraněno:** of  
**Odstraněno:** in the manner of Kašpar et al. (2013)  
**Odstraněno:** of its still better performance

1 identify the most appropriate length of the filter and chose Gaussian smoothing with a  
2 standard deviation  $\sigma$  of 30 days and a time window of  $3\sigma$ . Time-smoothed values of the mean  
3 and skewness are hereinafter referred to as  $\mu_{dG}$  and  $\gamma_{dG}$ , respectively.

4 Even the values of  $\mu_{dG}$  and  $\gamma_{dG}$  fluctuate through the year because of seasonal changes in  
5 precipitation climatology. The actual daily totals  $P_d$  were standardized using the relationship

Odstraněno: d

6 
$$P_{ms} = \bar{P} \left( \frac{P_d}{\mu_{dG}} \right)^{\bar{\gamma}}. \quad (2)$$

7 where  $P_{ms}$  is the seasonally standardized daily total,  $\mu_{dG}$  is the time-smoothed mean,  $\gamma_{dG}$  is the  
8 time-smoothed skewness of the distribution of daily totals  $\geq 0.1$  mm for a calendar day  $d$ , and  
9  $\bar{P} = E(\mu_{dG})$  and  $\bar{\gamma} = E(\gamma_{dG})$ . The transformation (2) directly standardizes the mean and  
10 skewness and indirectly standardizes the standard deviation and kurtosis of the daily data. The  
11 correction using  $\bar{P}$  and  $\bar{\gamma}$  induces an important feature of seasonally standardized daily  
12 totals: their mean annual sum equals the actual mean annual total. This process only  
13 redistributes precipitation amounts within the data series: seasonally standardized daily totals  
14 become higher and lower in seasons that are less and more exposed to high precipitation,  
15 respectively.

16 An example from the mountain station Churáňov is presented in Figs. 3 and 4. The mean and  
17 skewness are at a maximum in the summer, whereas the winter is characterized by a  
18 minimum of smoothed means but a secondary maximum of the skewness. As a result,  
19 extreme totals are substantially reduced (approximately 30 %) by the dual standardization  
20 when they occur in the summer. The winter totals are increased (20 %) due to the  
21 standardization of means; the increase in the totals due to the standardization of the skewness  
22 is present primarily in the spring (15 %). Figure 4 confirms that both moments need to be  
23 standardized to obtain a rather even annual distribution of extremes throughout the year.

## 24 **2.3 Evaluation of flood extremity**

25 To compare the precipitation and flood extremity, we also designed the flood extremity index  
26 (FEI), which is analogous to the WEI and enables us to recognize extreme flood events  
27 (EFEs). The FEI is based on return periods of peak discharges at individual sites. The Czech  
28 Hydrometeorological Institute provided data from 198 gauges beginning in 1961. However,  
29 only the approximate return period values of  $N = 5, 10, 20, 50$  or  $100$  years were available.

1 Each site represents a catchment with an area exceeding 100 km<sup>2</sup>. If there are one or more  
2 considered sites upstream, the catchment area does not include the respective sub-catchments.  
3 By analogy to the WEI, we combine the extremity at each site  $j$  expressed by the return period  
4  $N_j$  with the area of the respective catchment  $a_j$ . The area of the basin indirectly represents the  
5 magnitude of the river. Return periods were considered without evaluating the possible  
6 human impact on peak discharge so the value of the FEI represents the actual course of the  
7 flood instead of the theoretical one. Nevertheless, the very high discharges that are crucial for  
8 the evaluation of an event are generally less affected by human activities (Langhammer,  
9 2008).

10 The FEI is defined as the maximum of the variable  $F_a$ , which is given by the equation

$$11 F_a = \frac{\sum_{j=1}^h (\log(N_{ti}) a_j)}{a_a} \frac{\sqrt{a_a}}{\sqrt{\pi}}. \quad (3)$$

12 The aggregated area  $a_a$  consists of  $h = 1, \dots, 198$  considered catchments, which are ordered  
13 according to  $N_j$  in descending order. Return periods shorter than 5 years were assigned a value  
14 of  $N_j = 1$  so that  $\log(N_j) = 0$  and the respective catchments do not contribute to the resultant  
15 FEI value.

16 The method is demonstrated in Fig. 5, which compares flooding due to the EPEs in August  
17 1977 and in May/June 2013. The  $F_a$  curve representing the 2013 EFE increases more rapidly  
18 and starts to decrease later than in 1977; maxima of the curves depict values of the FEI. The  
19 reason is that return periods of peak discharges did not reach 50 years during the first event,  
20 whereas during the latter event, the total area of catchments corresponding to gauges with  
21 peak flows of  $N_j \geq 100$  years exceeded 2000 km<sup>2</sup>, and the value of  $a_a$  corresponding to the FEI  
22 was larger. Unlike in Fig. 1, the curves are not fluent because only discrete values of  $N_j$  were  
23 used (see above). Because of similarities between Eqs. (1) and (3), the WEI and the FEI reach  
24 values of the same order. Nevertheless, values of the FEI are usually slightly smaller for  
25 several reasons, primarily because all return periods between two values (20 and 50 years, for  
26 example) are assigned the lower value. Moreover, the maximum values are 100 and 1000  
27 years when calculating the FEI and the WEI, respectively.

28 There is a question regarding the manner in which the FEI design can affect the results.  
29 Therefore, we also calculated a simpler index that only aggregates areas of affected

**Odstraněno:** , which can make the results slightly inaccurate. However,

1 catchments weighted by return period values of peak discharges. An approximate (non-linear)  
2 correlation between the two indices was noted, which indicates that the FEI concept is  
3 sufficiently robust.

4 A serious problem is the separation of individual EFEs if additional peaks occur during a  
5 short period in the same catchments. We decided to separate EFEs with respect to EPEs. For  
6 example, we distinguished two EFEs in August 2002 because they were produced by two  
7 independent EPEs (Fig. 2). Naturally, the extremity of the latter EFE was affected not only by  
8 the latter EPE but also by the previous one in such a case; this fact can partly explain the  
9 discrepancies between the extremity of precipitation and of subsequent flooding.

10 Finally, we tried to assess how much the FEI is influenced by human regulations. Therefore, a  
11 parallel calculation of the FEI was performed (hereinafter FEIu) using only data from 97  
12 stations where no reservoir with flood protection function is present upstream. A linear  
13 dependence was detected between the FEI and the FEIu with Pearson correlation coefficient  
14 as high as 0.86. For example, four maximum EFEs were evaluated as the extremes regardless  
15 either the FEI or the FEIu is used. Only four outliers were detected when the FEIu was  
16 substantially reduced; it happened because flooding occurred almost only in catchments not  
17 involved in the FEIu calculation. Nevertheless, since the FEIu represents only rather small  
18 catchments, we prefer to use the FEI which seems to be robust enough also the viewpoint of  
19 human regulations.

## 20 2.4 Comparison of precipitation extremes and floods

21 Extremity of precipitation and of subsequent flooding were compared using rankings of the  
22 events and, furthermore, by the normalized ratio of respective extremity indices:

$$23 C_e = 100 \frac{FEI}{WEI}. \quad (4)$$

24 and

$$25 C_a = 100 \frac{FEI}{WAI}. \quad (5)$$

26 Values significantly above and below average resulting from Eqs. (4) and (5) indicate that the  
27 hydrological response to the precipitation event was most likely affected by factors other than  
28 only the extremity of precipitation. One of these factors may be antecedent saturation. This

1 parameter was expressed using the antecedent precipitation index (Köhler and Linsley,  
2 1951) spanning 30 days ( $API_{30}$ ) before the first day of the EPE/APE, which is calculated  
3 using the relationship

$$4 API_n = \sum_{i=1}^n P_i k^{n-i+1}. \quad (6)$$

5 where  $P_i$  is the daily total during the  $i$ -th day of the period under consideration spanning  $n =$   
6 30 days, and 0.93 is the generally accepted value of the constant  $k$  for the Czech Republic  
7 (Brázdil et al., 2005).

8 One of important climatological aspects of EPEs, APEs, and EFEs is their seasonal  
9 distribution. To analyze this distribution, we adopted the directional characteristics method  
10 (e.g., Black and Werritty, 1997), which was applied also to floods on selected Czech rivers by  
11 Čekal and Hladný (2008). Individual extreme events were depicted in a radial diagram where  
12 directions of vectors, which originated in the diagram's center and ended in the signs  
13 representing individual events, account for calendar days. In contrast to the above mentioned  
14 papers, we modified the method so that signs representing individual events are not located on  
15 a unit circle but their distance from the diagram's center is proportional to the WEI, WAI, or  
16 FEI. The resultant diagram better represents the seasonality of extremes because strong events  
17 are assigned greater weighting.

18

### 19 3 Application to the Czech Republic

#### 20 3.1 Precipitation extremes and floods

21 Although the WEI, the WAI, and the FEI itself are independent of thresholds, it was  
22 necessary to limit their values to constrain the sets of events that would be classified as EPEs,  
23 APEs, and EFEs. This step was performed because there are no natural limits dividing  
24 extreme from non-extreme events. In fact, the extremity of events gradually decreases with  
25 even smaller differences among the events as less-extreme events are considered. We selected  
26 the 50 events of each type so that one extreme event occurs per year on average. Sets of EPEs,  
27 APEs, and EFEs during the period of 1961–2010 in the Czech Republic are listed in Fig. 6.

28 The sets of EPEs, APEs, and EFEs partly overlap. We identified 22 precipitation extremes  
29 that were classified both as EPEs and APEs. More than a half of the EPEs and nearly 50 % of

**Odstraněno:** by

**Odstraněno:** aiming from a point in various directions representing

**Odstraněno:** by vectors aiming from a point in various directions representing

**Odstraněno:** when they occurred

**Odstraněno:** The mean of the vectors indicates the mean day representing the seasonal centroid of the phenomenon under consideration. Moreover, the length of the resultant vector is proportional to the inequality of the distribution of events throughout the year.

**Odstraněno:** In studying the sets of extremes, we considered not only the frequency but also the extremity of the events. Therefore,

**Odstraněno:** the lengths

**Odstraněno:** y

**Odstraněno:** w

**Odstraněno:** ere

**Odstraněno:** of the vectors representing individual events reflect the magnitude of the events with respect

**Odstraněno:** vectors

the APEs produced EFEs. If only the warm half of the year is considered, the number of EFEs produced by EPEs increases to 75 %. This fact suggests that the magnitude of causal precipitation is the main factor conditioning most of floods in the Czech Republic. Nevertheless, we also identified cases in which the hydrological response to an EPE was too small or too big. These events confirm that flooding is also significantly influenced by other factors which are further discussed in Sect. 4.2.

**Odstraněno:** These findings indicate that the proposed indices reasonably reflect the extremity of the studied phenomena.

**Odstraněno:** These discrepancies will be

**Odstraněno:** These discrepancies will be

### 3.2 Inter-annual variability of extremes

The temporal distribution of extreme events during the period of 1961–2010 is shown in Fig. 7. Regardless of the type of extremes (EPEs, APEs, and EFEs), there are certain common features of their variations in time. One such feature is the below-normal frequency and magnitude of all types of extremes during the first 16 years of this time period. The situation dramatically changed in 1977, when three EPEs occurred in the span of less than one month; two of them produced EFEs (Fig. 6). Similar conditions occurred during the following two years. Moreover, the second and the fourth highest values of the WEI were recorded in July 1981 and August 1983, respectively. The wet years of 1985–6 ended a decade of all types of extremes with above-normal frequency and magnitude.

**Odstraněno:** The accumulated values of all three indices reached only approximately 15 % of their values for the entire period, whereas this 16-year time interval (1961–1976) spans 32 % of the studied period.¶

**Odstraněno:** these were so strong that they also qualified as APEs and

**Odstraněno:** During the five-year period of 1977–1981, 8 EPEs, 9 EFEs, and 10 APEs occurred. After three unremarkable years, t

In contrast, the following decade of 1987–1996 was lacking in hydrometeorological extremes. EFEs were completely absent from the warm half-years (MJJASO) and did not occur again until 1996. The rest of the 1990s would also be considered to be below normal if July 1997 were not included. The first of two EPEs in this month exhibited maximum values of both the WEI and WAI. Similar values were also observed five years later, in August 2002. Compared to similarly strong precipitation events in the early 1980s, these two events produced much greater flooding. Last five years of the study period were characterized by an abnormally high number of extremes, which were concentrated primarily in 2006 and 2010. If the 2013 flood were considered, four maximum EFEs occurred recently approximately every five years.

**Odstraněno:** EPEs were generally weaker than APEs during this dry period and produced no flooding.

**Odstraněno:** ¶

**Odstraněno:** During the period of 2003–5, again no EPEs and EFEs occurred during the warmer half-years. Finally, the 1

**Odstraněno:** (three EPEs and four EFEs in each of these years)

**Odstraněno:** through

**Odstraněno:** of any given year

**Odstraněno:** two

**Odstraněno:** both being

**Odstraněno:** and lasting five days

**Odstraněno:** In contrast, short events (1–2 days) occurred only from late May through early September. EPEs were as frequent in May as in September, but they were generally less extreme in May (WEI only up to 70). ¶

**Odstraněno:** four

### 3.3 Seasonal distribution of extremes

The seasonal distribution of EPEs was significantly unequal during the period of 1961–2010 (Fig. 8). Based on the selected threshold, these events occurred from May to December. Nevertheless, only three such events occurred since October, all rather weak. The months of greatest activity were clearly July and August, when the five highest values of WEI were

1 noted. The level of activity during the first half of August was particularly pronounced: this  
2 time period was the seasonal peak in EPEs during the period of 1961–2010.

**Odstraněno:** Nevertheless, there was a difference in the frequency of EPEs between July (9 events) and August, when they occurred twice as often. If another threshold were considered (e.g., WEI = 50), the dominance of August would be even more pronounced.

3 Naturally, APEs were distributed more equally from season to season during the 1961–2010  
4 period than were the EPEs (Fig. 8). We noted at least one event in every calendar month.  
5 From October to March, the distribution of APEs was very uniform in terms of both the  
6 number of events and the magnitude. The values of the WAI were less than 100 with only one  
7 exception, which occurred during the 5 days from 28 December 1986 to 1 January 1987. This  
8 event was so exceptional that it also qualified as an EPE (see above). In contrast, only one  
9 APE was noted in April. This event and two others in the first half of May lasted only one day  
10 each.

**Odstraněno:** As a result, the mean day of the seasonal distribution of EPEs is 27 July.

**Odstraněno:** (3–5 per month)

11 The seasonal distribution of EFEs partly correlates with the seasonality of precipitation  
12 extremes, but it is also significantly affected by snow accumulation during the winter and by  
13 changes in the saturation of basins. As a result, we identified three main periods when the  
14 frequency of EFEs is increased: (i) the period from May to August, when most of the EPEs  
15 occur; (ii) the second half of December and early January, when the values of the WAI are  
16 slightly increased in comparison with the months before and after; (iii) March and very early  
17 April, when the values of the WAI are not very high. Mainly the last period is affected by  
18 thawing. The rest of April is characterized by a distinct break both in precipitation and flood  
19 events. As in the winter apart from December, only weak EFEs occurred during the fall,  
20 when, in contrast, APEs were frequently noted. September appears to be the month with  
21 maximum differences between precipitation and flood extremity; although several high EPEs  
22 were noted then, the EFEs remained small.

**Odstraněno:** Another significant feature of the seasonal distribution of APEs during the 1961–2010 period was the lower number of such events in June and July. Apart from 3 EPEs with values of the WEI exceeding 100, which also qualified as the largest APEs, less significant EPEs dropped below the threshold of the WAI. However, August remained the most represented month, with 8 APEs. Even greater values of the WAI in excess of 100 were noted in September. This finding is most likely due to the significant decrease in mean precipitation in September, which results in such events being abnormal during this month. As a result, the mean day of the seasonal distribution of APEs is 5 October, although the length of the resulting vector is very small.¶

**Odstraněno:** because

## 23 24 4 Discussion of results

### 25 4.1 Comparison with standard indices

26 In this section, the presented evaluation of precipitation and flood events is compared with  
27 standard methods mentioned in Sect. 1.

**Odstraněno:** T

**Odstraněno:** extremity

**Odstraněno:** can be

**Odstraněno:** the

28 Regarding precipitation extremes, they are usually detected by maximum daily totals  $P_1$  at  
29 any individual rain gauge (Ustrnul et al., 2015) or by areal precipitation means  $A_d$  during  $d$   
30 days, for example for  $d=2$  days (Conrad, 2001). The first approach is commonly used in the

1 Czech Republic: Štekł et al. (2001) analyzed days during the period of 1876–2000 when a  
2 daily total reached at least 150 mm anywhere in the Czech Republic. The Czech maximum  
3 occurred on 29 July 1897 when a  $P_1$  of 345.1 mm was measured at the Nová Louka gauge in  
4 the Jizerské Hory Mts. Indeed, this event was the maximum  $P_1$  ever recorded in Central  
5 Europe (Munzar et al., 2011).

**Odstraněno:** The presented evaluation of precipitation extremity can be compared with the standard approach based on maximum daily totals recorded at individual gauges because

**Odstraněno:**

6 Figure 9 compares 50 maximum WEI values with maximum  $P_1$  and areal 1, 2, and 3-day  
7 means during the EPEs. It demonstrates that areal precipitation means generally decrease with  
8 decreasing WEI values but it is not fully true for point daily totals. As a result, the ranking of  
9 precipitation events significantly depends on the evaluation criterion. For example, daily point  
10 maxima were above 200 mm on 20 August 1972 and 22 July 1998 but the 1-day areal means  
11 were only 5.2 and 7.3 mm, respectively. The affected area was very small in both cases (see  
12 Fig. 6). In 1972, strong precipitation affected a large area but mainly outside the Czech state  
13 border (Müller et al., 2009) whereas the 1998 case was due to a strong convective storm  
14 within a limited area (Řezáčová and Sokol, 2003). Thus, the WEI seems to be a good  
15 compromise between the two standard approaches.

**Odstraněno:** ¶

**Odstraněno:** During the 40 years that overlap with our study, the authors identified 18 such days, 13 of which were days when EPEs occurred. Because of the high autocorrelation of the daily precipitation time series, on several occasions, this threshold was exceeded on two or even three consecutive days with an EPE.

**Odstraněno:** daily

**Odstraněno:** s

**Odstraněno:** These included

**Odstraněno:** ,

**Odstraněno:** , which

**Odstraněno:** a daily maximum

**Odstraněno:** the warmer half-years

**Odstraněno:** . However.. However,

**Odstraněno:** s

**Odstraněno:** ere

**Odstraněno:** in 1890

**Odstraněno:** T

**Odstraněno:** hese

**Odstraněno:** These two

**Odstraněno:** that it is necessary to take

**Odstraněno:** into account both

**Odstraněno:** that it is necessary to take

into account both the

**Odstraněno:** to make valid comparisons with consequent floods

16 In contrast, there were many significant EPEs with high areal totals but without any point  
17 daily total exceeding 150 mm. For example, the second largest EPE in July 1981 was  
18 characterized by  $P_1$  of only 122.0 mm but extra high areal totals (absolute maximum of  $A_1$ )  
19 produced the third largest EFE during MJJASO (Fig. 6). A still lower daily maximum (only  
20 97.6 mm) was recorded during the historical case on 2 September 1890 when the 3-day areal  
21 precipitation total (1-3 September) was nearly as high as in August 2002, and subsequent  
22 flooding was only slightly less catastrophic (Řezáčová et al., 2005a). The WEI reached very  
23 high value in July 1981 because the index reflects the size of the affected area and the fact  
24 that heavy rains occurred not only in mountains but also in lowlands where they are rare.

25 Moreover, some precipitation events lasted virtually only one day (e.g., 23 June 1992) but  
26 others lasted three or even more days (e.g., in July 1997). The first case is not properly  
27 evaluated by the areal 3-day mean total and the latter by the areal 1-day mean while the WEI  
28 accommodates to the real duration of the events. All above mentioned examples demonstrate  
29 the advantage of the WEI which combines the intensity of precipitation with its spatial extent  
30 and reflects also duration of precipitation events.

31 Precipitation extremes are usually evaluated regardless the season when they occurred.  
32 Recently, Ramos et al. (2014) suggested an index ranking daily precipitation extremes which

1 employs 1-day precipitation totals normalized by their climatology on the given calendar day.  
2 The index can be hardly compared with the WAI because Ramos et al. (2014) do not reflect  
3 duration of precipitation events. Nevertheless, we decided to accumulate daily values of their  
4 index on individual days of the APEs and compare the sums (hereinafter called RI) with the  
5 WAI values (Fig. 10).

6 Naturally, the WAI is more closely related to the WEI than the RI. Values of the WAI are  
7 higher than the WEI from September to the middle of May and lower in the rest of the year.  
8 Values of the RI seem to be a bit more equally distributed through the calendar year but they  
9 correspond less to the precipitation extremity. It can be demonstrated by the comparison of  
10 two July events: the maximum EPE from the beginning of July 1997 is represented by only  
11 the 16<sup>th</sup> highest RI value but significantly less extreme event from 1977 reached the 6<sup>th</sup>  
12 highest value of the RI though the turn of July and August is the top season of heavy rains in  
13 the Czech Republic. A possible reason can be the fact that daily precipitation totals are not  
14 normally distributed but the RI standardizes them only by the mean and the variance.

15 Finally, the FEI was compared with the Francou index (FI) and the index by Uhlemann et al.  
16 (2010) (hereinafter UI), see Sect. 1. Data from only eight large Czech catchments were at our  
17 disposal: Labe in Brandýs, Otava in Písek, Berounka in Beroun, Vltava in Prague, Labe in  
18 Děčín, Odra in Bohumín, Morava in Strážnice, and Dyje in Doní Věstonice / Nové Mlýny.  
19 We considered the maximum FI value from the eight catchments and the UI representing all  
20 these catchments together. To enable the comparison, the FEI was recalculated again using  
21 data from only the eight gauges (hereinafter FEI 8). Only 31 of EFEs reached the FEI 8  
22 values above zero (Fig. 11). Even such a rough comparison of the indices shows that the FEI  
23 is a compromise between the FI representing the most affected catchment and the UI which  
24 emphasizes the size of the affected area even more than the FEI. The difference between the  
25 FI and the UI was mainly evident in July 1997 when the floods were extreme in the eastern  
26 part of the Czech Republic but no flood occurred in the western part.

## 27 **4.2 Relationship between precipitation extremes and floods**

28 Though many precipitation extremes correspond with EFEs, there are still discrepancies  
29 between the WEI and the FEI and even greater discrepancies between the WAI and the FEI  
30 values. For example, the fourth largest EPE did not produce an EFE in August 1983; in fact,  
31 this EPE resulted in very limited flooding ( $C_e = 8\%$ , see Fig. 6). Several factors affected the

**Odstraněno:** Nevertheless

1 hydrological response of this event. These include unusually low antecedent saturation (mean  
2 API<sub>30</sub> only 9.3 mm!) and a moderately even distribution of rainfall over five days, whose  
3 maximum occurred on the second day of the event. Regulation processes by the dams could  
4 also play a role; nevertheless, Brázil et al. (2005) confirm that no even unaffected peak flow  
5 with the return period of two or more years occurred at Vltava River in Prague even though  
6 the catchment belonged to the most affected by heavy rains.

Odstraněno: T

7 All these factors will be studied in the future together with spatial patterns of precipitation to  
8 elucidate the discrepancies between individual precipitation and flood events. One of the  
9 factors to be considered should be the season when an EPE occurred, as discussed in Sect.  
10 3.3. The important role of this factor is confirmed by Fig. 12. It is clear that the hydrological  
11 effect of an EPE is typically strong in May, more ambiguous in summer, and considerably  
12 weaker in September. The very last event from the turn of May and June 2013 also supports  
13 this conclusion ( $C_e = 97\%$ , see Fig. 6). If an EPE occurs in the last month of the year, it can  
14 also produce rather big flooding, although such events are very rare.

Odstraněno: 9

15 The hydrological effect of APEs (Fig. 12) is substantially reduced in the winter, early spring  
16 (most likely due to precipitation in the form of snow) and autumn. In contrast, if precipitation  
17 events are sufficiently high to qualify as APEs in the late spring and summer, they are usually  
18 flood producing; surprisingly, this was also the case with all APEs in December. A  
19 subsequent detailed study of intra-annual variations in precipitation patterns is necessary to  
20 explain these findings.

Odstraněno: 6

Odstraněno: appears to

Odstraněno: 9

21 However, seasonality can hardly explain the difference in flood activity between two periods  
22 with unusually high EPEs (1977–1986 and 1997–2010). The FEI exceeded 50 in association  
23 with only two EFEs during the first period, whereas this occurred six times since 1997 (Fig.  
24 7). Several factors most likely explain the difference: (i) if two or more EPEs appeared during  
25 one year in the first period, they were separated by a much longer interval than in the latter  
26 period (the shortest interval between two EPEs was 8 days in 1977 but only 3 days in 2002);  
27 (ii) in cases of EPEs following one after another, the magnitude decreased in the first period  
28 (WEI decreasing from 113.5 to 49.6 in July/August 1977) but increased in the second period  
29 (WEI increasing from 104.3 to 174.3 in August 2002); (iii) just before the main EPEs, the  
30 mean antecedent saturation was much lower on 17 July 1981 (24.6 mm) and on 1 August  
31 1983 (9.3 mm) than on 4 July 1997 (34.5 mm) and on 11 August 2002 (50.1 mm because of  
32 the above-mentioned preceding EPE); (iv) the EPEs in 1981 and 1983 lasted one day longer

Odstraněno: is

Odstraněno: most

Odstraněno: 38.3

Odstraněno: 2

Odstraněno: 172

1 than those in 1997 and 2002. It demonstrates that changes in a flood regime can also occur  
2 without significant changes in only the magnitude of precipitation events.

3

## 4 **5 Conclusions**

5 The paper presents three indices for evaluation of hydrometeorological extremes. In contrast  
6 to common indices, the presented indices reflect not only maxima of precipitation amounts  
7 and peak discharges at individual gauges but also the rarity of values, the size of the affected  
8 area, and the duration of precipitation. Besides that, the aspect of precipitation seasonality was  
9 considered which enables the detection of precipitation extremes throughout the year. A  
10 unified design of the presented indices enables to compare easily inter-annual and seasonal  
11 distributions of precipitation extremes and large floods.

12 The application of the indices to the Czech territory demonstrates that this approach enables  
13 to compare the extremity of precipitation and consequent floods rather than if precipitation  
14 events are evaluated only by the maximum precipitation total at one station. Extreme floods  
15 correspond to precipitation extremes; nevertheless, not only the magnitude of precipitation  
16 extremes influences the hydrological response but also the season, the antecedent saturation  
17 and other factors. The study confirms that variations in the frequency and/or magnitude of  
18 floods can be due not only to variations in the magnitude of precipitation events but also to  
19 variations in these factors.

20 Additional detailed studies are necessary for elucidating the way in which seasonality  
21 influences the hydrological effect of precipitation extremes. This effect could be due to  
22 seasonal differences in evapotranspiration or to possible seasonal variations in the attributes  
23 of the precipitation itself. The events can differ, e.g., in the spatial distribution of precipitation  
24 within the affected area or in the temporal concentration of precipitation during the event  
25 (intensity can increase, remain the same or decrease). In addition, various circulation  
26 conditions could explain the differences among the extremes (Kašpar and Müller, 2010). In a  
27 next step, we plan to explore the dependences on the circulation extremity index (Kašpar and  
28 Müller, 2014), which completes the set of tools for studying the pathway of causation from  
29 circulation to precipitation and runoff.

30

## 31 **Acknowledgements**

**Odstraněno:** is study

**Odstraněno:** ,

**Odstraněno:** d to

**Odstraněno:** , precipitation extremes should be evaluated considering not only maxima at individual sites but also the spatial extent, duration and temporal concentration of the precipitation.

**Odstraněno:** . Although most of the EFEs were produced by an EPE in warmer half-years, fewer than half of all EFEs were produced by an APE because thawing can substantially enhance the discharge. Most significant EPEs occurred in July and the first half of August, but their hydrological responses were also significantly influenced by

**Odstraněno:** As a result, the accumulation of precipitation extremes during the 1977-1986 period produced less significant flooding than another accumulation after 1996. In general, the primary discrepancies between the magnitude of EPEs and EFEs were noted in May and September, when consequent floods were usually much larger and smaller in relation to the WEI, respectively. The hydrological response to APEs was usually strong in December, whereas another accumulation of EFEs in March was usually not due to APEs. Neither precipitation nor flood extremes occurred from early April through early May. ¶

**Odstraněno:** ir seasonal distribution and other factors, primarily the antecedent saturation

1 The research presented in this study is supported by the Czech Science Foundation under the  
2 GACR P209/11/1990 project. Precipitation data were provided by the Czech  
3 Hydrometeorological Institute. We would also like to thank J. Kyselý and L. Gaál from the  
4 Institute of Atmospheric Physics in Prague for implementing the ROI method.

5

1    **References**

- 2    Alexander, L. V., Zhang, X., Peterson, T. C., Caesar, J., Gleason, B., Klein Tank, A. M. G.,  
3    Haylock, M., Collins, D., Trewin, B., Rahimzadeh, F. Tagipour, A., Rupa Kumar, K.,  
4    Revadekar, J., Griffiths, G., Vincent, L., Stephenson, D. B., Burn, J., Aguilar, E., Brunet, M.  
5    Taylor, M., New, M., Zhai, P., Rusticucci, M., and Vazquez-Aguirre, J. L.: Global observed  
6    changes in daily climate extremes of temperature and precipitation, *J. Geophys. Res.*, 111,  
7    D05109, doi:10.1029/2005JD006290, 2006.
- 8    Barredo, J. I.: Major flood disasters in Europe: 1950–2005, *Nat Hazards*, 42, 125–148, 2007.
- 9    Begueria, S., Vicente-Serrano, S. M., Lopez-Moreno, J. I., and Garcia-Ruiz, J. M.: Annual  
10   and seasonal mapping of peak intensity, magnitude and duration of extreme precipitation  
11   events across a climatic gradient, northeast Spain, *Int. J. Climatol.*, 29, 1759–1779, 2009.
- 12   Black, A. R. and Werritty, A.: Seasonality of flooding: a case study of North Britain, *J.  
13   Hydrol.*, 195, 1–25, 1997.
- 14   | [Blinchikoff, H. J. and Zverev, A. I.: Filtering in the time and frequency domain, Scitech  
15   Publishing, Raleigh, United States of America, 2001.](#)
- 16   Brázdil, R., Dobrovolný, P., Elleder, L., Kakos, V., Kotyza, O., Květoň, V., Macková, J.,  
17   Müller, M., Štekl, J., Tolasz, R., and Valášek, H.: Historical and recent floods in the Czech  
18   Republic, (in Czech, including 83 pp. of an English abstract), Masaryk University, Brno,  
19   Czech Republic, 2005.
- 20   Burn, D. H.: Evaluation of regional flood frequency analysis with a region of influence  
21   approach, *Water Resour. Res.*, 26, 2257–2265, 1990.
- 22   Čekal, R. and Hladný, J.: Analysis of flood occurrence seasonality on the Czech Republic  
23   territory with directional characteristics method, *AUC Geographica*, 43, 3–14, 2008.
- 24   Herschy, R. W. (Ed.): World catalogue of maximum observed floods, IAHS-AISH Publ. 284,  
25   IAHS Press, Wallingford, 2003.
- 26   Hosking, J. R. M. and Wallis, J. R.: Regional frequency analysis: an approach based on L-  
27   moments, Cambridge University Press, New York, 1997.
- 28   Kašpar, M. and Müller, M.: Selection of historic heavy large-scale rainfall events in the Czech  
29   Republic, *Nat. Hazards Earth Syst. Sci.*, 8, 1359–1367, 2008.

- 1 Kašpar, M. and Müller, M.: Variants of synoptic patterns inducing heavy rains in the Czech  
2 Republic, *Phys. Chem. Earth*, 35, 477–483, 2010.
- 3 Kašpar, M. and Müller, M.: Combinations of large-scale circulation anomalies conducive to  
4 precipitation extremes in the Czech Republic, *Atmos. Res.*, 138, 205–212, 2014.
- 5 Köhler, M. A. and Linsley, R. K.: Predicting the runoff from storm rainfall. U.S. Weather  
6 Bureau Research Paper no. 34., Washington, 9 pp., 1951.
- 7 Konrad, C. E.: The most extreme precipitation events over the Eastern United States from  
8 1950 to 1996: Considerations of scale, *J. Hydrometeor.*, 2, 309–325, 2001.
- 9 Kyselý, J., Gaál, L., and Picek, J.: Comparison of regional and at-site approaches to modelling  
10 probabilities of heavy precipitation, *Int. J. Climatol.*, 31, 1457–1472, 2011.
- 11 Kyselý, J. and Picek, J.: Regional growth curves and improved design value estimates of  
12 extreme precipitation events in the Czech Republic, *Clim. Res.*, 33, 243–255, 2007.
- 13 Langhammer, J.: Modelling the impact of anthropogenic modifications to river channels on  
14 the course of extreme floods. Case study: August 2002 flood, Blanice River basin, Czechia,  
15 *Geografie*, 113, 237–252, 2008.
- 16 Müller, M. and Kašpar, M.: Event-adjusted evaluation of weather and climate extremes, *Nat.*  
17 *Hazards Earth Syst. Sci.*, 473–483, 2014.
- 18 Müller, M., Kašpar, M., and Matschullat, J.: Heavy rains and extreme rainfall-runoff events in  
19 Central Europe from 1951 to 2002, *Nat. Hazards Earth Syst. Sci.*, 9, 441–450, 2009.
- 20 Munzar, J., Ondráček, S., and Auer, I.: Central European one-day precipitation records,  
21 *Moravian Geographical Reports*, 19, 32–40, 2011.
- 22 Ramos, A. M., Trigo, R. M., and Liberato, M. L. R.: A ranking of high-resolution daily  
23 precipitation extreme events for the Iberian Peninsula, *Atmos. Sci. Lett.*, 15, 328–334, 2014.
- 24 Ren, F. M., Cui, D. L., Gong, Z. Q., Wang, Y. J., Zou, X. K., Li, Y. P., Wang, S. G., and  
25 Wang, X. L.: An objective identification technique for regional extreme events, *J. Clim.*, 25,  
26 7015–7027, 2012.
- 27 Řezáčová, D. and Sokol, Z.: A diagnostic study of a summer convective precipitation event in  
28 the Czech Republic using a nonhydrostatic NWP model, *Atmos. Res.*, 67–68, 559–572, 2003.

**Odstraněno:** Kašpar, M., Müller, M., and Pecho, J.: Comparison of meteorological conditions during May and August 2010 floods in Central Europe, *AUC Geographica*, 48, 27–34, 2013.¶

- 1 Řezáčová, D., Kašpar, M., Müller, M., Sokol, Z., Kakos, V., Hanslian, H., and Pešice, P.: A  
2 comparison of flood precipitation in August 2002 with historical extreme precipitation events  
3 from the Czech territory, *Atmos. Res.*, 77, 354–366, 2005a.
- 4 Řezáčová D., Pešice P., Sokol Z.: An estimation of the probable maximum precipitation for  
5 river basins in the Czech Republic, *Atmos. Res.*, 77, 407–421, 2005b.
- 6 Řezáčová D. and Sokol Z.: A diagnostic study of a summer convective precipitation event in  
7 the Czech Republic using a nonhydrostatic NWP model, *Atmos. Res.*, 67–68, 559–572,  
8 2003.
- 9 Rodier, J. A. and Roche M. (Eds.): World catalogue of maximum observed floods,  
IAHS Publ. 143, IAHS Press, Wallingford, 1984.
- 10 Šercl, P., Tyl, R., and Pecha, M.: The course and the extremity of the June 2013 floods,  
11 Meteorologické zprávy, 66, 197–202, 2013.
- 12 Štekł, J., Brázdil, R., Kakos, V., Jež, J., Tolasz, R., and Sokol, Z.: Extreme daily precipitation  
13 totals during 1879–2000 in the Czech territory a their synoptic causes, (in Czech), Národní  
14 klimatický program České republiky, 31, Praha, Czech Republic, 140 pp., 2001.
- 15 Stephenson, D. B.: Definition, diagnosis, and origin of extreme weather and climate events,  
16 in: Climate extremes and society, Diaz, H. F. and Murnane, R. J. (Eds.), Cambridge  
17 University Press, New York, 11–23, doi:10.1017/CBO9780511535840.004, 2008.
- 18 Tolasz, R., Míková, T., Valeriánová, A., and Voženílek, V. (Eds.): Climate atlas of Czechia,  
19 Czech Hydrometeorological Institute, Praha, Czech Republic, 256 pp., 2007.
- 20 Uhlemann, S., Thielen, A. H., and Merz, B.: A consistent set of trans-basin floods in  
21 Germany between 1952–2002, *Hydrol. Earth Syst. Sci.*, 14, 1277–1295, 2010.
- 22 Ustrnul, Z., Wypych, A., Henek, E., Maciejewski, M., and Bochenek, B.: Climatologically  
23 based warning system against meteorological hazards and weather extremes: the example for  
24 Poland. *Nat. Hazards*, 77, 1711–1729, 2015.
- 25 Zhang, X. B., Alexander, L., Hegerl, G. C., Jones, P., Klein Tank, A., Peterson, T. C., Trewin,  
26 B, and Zwiers, F. W.: Indices for monitoring changes in extremes based on daily temperature  
27 and precipitation data, *Wiley Interdiscip. Rev.-Clim. Chang.*, 2, 851–870, 2011.
- 28

## 1   Figure captions

2   Figure 1. Determination of the WEI of the EPE in May/June 2013 by maximizing  $E_{ta}$ . The  
3   inset maps present interpolated return periods of 1-day totals on 1 June (left) and of 5-day  
4   totals from 30 May to 3 June (right).

5   Figure 2. Mean daily precipitation totals  $P_{mean}$  in the Czech Republic (right axis) and  
6   respective maximum values of  $E_{ta}$  on 4–13 August 2002 (color bars, left axis). Selected  
7   maximum  $E_{ta}$  values for time windows with various lengths of  $t$  days are depicted including  
8   the hypothetical value of maximum  $E_{ta}$  for the 7-day period of 6–12 August.

9   Figure 3. Annual cycle of the mean, standard deviation, and skewness of daily precipitation  
10   totals at Churáňov station. Individual points represent the values of the mean, standard  
11   deviation, and skewness, calculated for each calendar day taking into account the period of  
12   1961–2010; curves depict data smoothed using the Gaussian filter.

13   Figure 4. Daily precipitation maxima on individual calendar days at Churáňov station during  
14   the 1961–2010 period: non-standardized totals ( $P_d$ ), totals standardized for the mean and  
15   variance only ( $P_m = \bar{P} P_d / \mu_{dG}$ ) and fully standardized totals ( $P_{ms}$ ). Dates of significant totals  
16   are indicated (day/month/yr).

17   Figure 5. Evaluation of extremity of floods in May/June 2013 and in August 1977 using the  
18   FEI. The step-by-step aggregated catchments are ordered in descending order with respect to  
19   return periods recorded there. The inset map presents return periods  $N_j$  of peak discharges at  
20   gauges in 2013; blue and green curves correspond to the main rivers and watersheds,  
21   respectively.

22   Figure 6. EPEs, APEs and EFEs in the Czech Republic, 1961–2010. Colors denote the  
23   assignment of events to one or more types of extremes; the ratios of the FEI to the WEI and to  
24   the WAI are designated the  $C_e$  and  $C_a$ , respectively. For comparison, an extra event from  
25   May/June 2013 is represented by values of the WEI and the FEI.

26   Figure 7. Temporal variability of extreme events, 1961–2010, in the Czech Republic. The red,  
27   blue, dark green, and light green symbols denote EPEs, APEs, EFEs during colder half-years  
28   (NDJFMA), and EFEs during warmer half-years (MJJASO), respectively. The symbol shapes  
29   denote the duration (# of days) of EPEs and APEs. The curves express relative cumulative

**Naformátováno:** Písmo: 12 b., není  
Tučné, Barva písma: Černá, Angličtina  
(USA)

**Odstraněno:** M

**Naformátováno:** Písmo: 12 b., není  
Tučné, Barva písma: Černá, Angličtina  
(USA)

**Odstraněno:** on individual calendar days

**Odstraněno:** during the period of 1961–  
2010;

**Naformátováno:** Angličtina (USA)

**Odstraněno:** d

1 values (right axis) of the WEI, the WAI, and the FEI (dark green denotes all seasons, light  
2 green denotes warmer half-years only).

3 Figure 8. Seasonal distribution of extreme events in the Czech Republic, 1961–2010. EPEs  
4 were evaluated using the WEI (red), APEs using the WAI (blue), and EFEs using the FEI  
5 (green). Values of the indices are depicted by the distance of symbols from the center of the  
6 diagram. The shape of symbols depicts how many days the precipitation event lasted.

7 [Figure 9. EPEs characterized by the WEI, maximum daily precipitation total \[mm\] at a site](#)  
8 [\( \$P\_1\$ \), and areal mean precipitation total \[mm\] in the Czech Republic recorded during 1 day](#)  
9 [\( \$A\_1\$ \), 2 days \( \$A\_2\$ \), and 3 days \( \$A\_3\$ \).](#)

10 [Figure 10. APEs characterized by the WAI, the WEI, and values of the index RI according to](#)  
11 [Ramos et al. \(2014\). The 50 APEs are ordered with respect to the calendar day in the](#)  
12 [increasing order.](#)

13 [Figure 11. EFEs characterized by the FEI, the index by Uhlemann et al. \(2010\) \(UI\), and by](#)  
14 [the maximum Francou index \(FI\) calculated within 8 selected large Czech catchments.](#)

15 [Figure 12. Relationship between rankings of 50 maximum EPEs \(left\) / APEs \(right\) and](#)  
16 [rankings of subsequent floods in various months \(distinguished by colors\). Precipitation](#)  
17 [events, whose hydrological response was too small \(no peak flow with return period at least 5](#)  
18 [years\), are depicted at the top of the diagram.](#)

**Odstraněno:** Figure 9. Relative magnitude of the hydrological response to individual EPEs (left) and APEs (right) with respect to their magnitude in different months (distinguished by colors).