

## RESPONSES TO THE REVIEWERS COMMENTS

We have considered all the suggestions and comments of the two reviewers, and we have made the modifications/corrections. We have detailed the replies when necessary in the letter (in black is the text of the reviewers and in blue our responses). The manuscript with the marked correction is up-loaded with this letter.

### REFEREE #1

The manuscript is based on a very detailed and extended data set, covering 10 years of monitoring at different stations in an estuary and tidal river. The data set is worthwhile to be published and the analysis of it reveals very interesting results. These results should convince authorities to extend monitoring of key ecological parameters, such as turbidity in other estuaries! The manuscript is (generally) well written and reasonably well structured. I recommend a publication of the manuscript in HESS after some revisions

#### a) GENERAL COMMENTS

**Referee Comment 1.** I was a little confused by the use of the terms 'low water' and 'high water' for discharge conditions. Usually they are used to indicate tidal elevation. I would therefore suggest replacing these words by 'low discharge' and 'high discharge' or 'low fresh water discharge' and 'high fresh water discharge'.

The terms were replaced in the text and figures, according the suggestion of RC#1, to avoid such confusion.

**Referee Comment 2.** The abstract starts with 'climate change and human activities'. I was immediately focused on these terms, but only very limited information was presented in the ms on these subjects. What do you mean by climate change? Is it global warming and its effect on e.g. precipitation or do you mean climate variability, such as the NAO?

Regarding the latter, the changes in e.g. duration of LW (low discharge) in Fig 15 correspond at a quick view (see [http://www.cpc.ncep.noaa.gov/data/teledoc/nao\\_ts.shtml](http://www.cpc.ncep.noaa.gov/data/teledoc/nao_ts.shtml)) with the variations in NAO index. The increase of LW duration in the 80's is correlated partly with a period (79/80-94/95) of positive NAO index. What is the effect of human impact (water usage for irrigation)? Do you have data that show the amount used for this purpose?

Yes, we begin the abstract with "climate change and human activities" to indicate the context of the article, however this is not the specific objective of the work. We know that the annual mean discharge of the Garonne-Gironde system has decreased over the last decades (see section 4.1 1<sup>st</sup> paragraph). There are already dedicated studies that confirm the influence of both human impact and climate variability on freshwater discharge variations, like those of Mazzega et al. (2014) and Hendrickx and Sauquet (2013), both cited in the article.


Such context is expected to affect natural distribution of estuarine SPM, as explained in the introduction. However we do not have historical data in the tidal Garonne River. The objective of the article is to detail turbidity dynamics in the upper reaches of the Gironde estuary in relation with the present-day hydrological conditions, based on a high-frequency turbidity records that covers only the last 10 years. We are convinced that figure 13 and 14 clearly demonstrate the decisive control of

fluvial discharge characteristics (duration of low discharge period; the river water volume passed during the previous high discharge water period) on TMZ occurrence in the tidal Garonne. The figure 15 is presented to show that these two indicators have already changed over the period 1959-2014.

We agree with the observation that the changes in duration of low discharge (Fig. 15) present some similarities with the NAO index. In fact, we have envisioned a first version of the figure 15 including the NAO index (see below). We had suggested Referee 1 the possibility of include this version of the Figure. However, in absence of response, entering in this discussion can be stodgy. Taking into account our main objective of presenting 10-yr high frequency turbidity records, we prefer do not present here the NAO index. However an additional discussion about the interest of this analysis for the future management of the tidal Garonne, considering the projections of river discharged in Europe for the next decades (Alfieri et al., 2015), is included (p16 lines 3-7),

Added reference:

Alfieri, L., Burek, P., Feyen, L., & Forzieri, G.: Global warming increases the frequency of river floods in Europe, *Hydrology and Earth System Sciences*, 19(5), 2247–2260. doi:10.5194/hess-19-2247-2015, 2015.



**Referee Comment 3.** How do you define 'mobile mud'? Is this the same as fluid mud, high concentrated mud suspension or are these low consolidated mud deposits? Are the data given any direct clue for the occurrence of these 'mobile muds', do you have other data that confirm the existence of these features or is their existence derived from the behavior of the turbidity variations?

The choice of the term "mobile mud" is better justified in the manuscript (p4 lines 19-23). We call "mobile mud" "low consolidated mud deposits", that are easily erodible. We discarded the term

“fluid mud” that refers precisely to a high-concentrated benthic suspension (several 10s of  $\text{g L}^{-1}$  to 100s of  $\text{g L}^{-1}$ ) often formed from settling of particles from the turbidity maximum. Field observations by Chanson et al. (2011) confirm the existence of mud deposits in the Garonne tidal, which are defined as “a cohesive mud mixture consisting of fine mud and silt materials” (the reference is included in the manuscript). For this reason we adopted this more neutral term, mobile mud, which is used in other estuaries (Uncles et al. 1996; 2006) to describe the seasonal occurrence of unconsolidated mud layers.

References:

Chanson, H., Reungoat, D., Simon, B., & Lubin, P. : High-frequency turbulence and suspended sediment concentration measurements in the Garonne River tidal bore, *Estuarine, Coastal and Shelf Science*, 95(2-3), 298–306. doi:10.1016/j.ecss.2011.09.012, 2011


Uncles, R. J., Barton, M.L., and Stephens, J. A. : Seasonal Variability of Mobile Mud Deposits in the Tamar Estuary, in *Mixing in Estuaries and Coastal Seas* (ed C. Pattiaratchi), American Geophysical Union, Washington, D. C., doi:10.1029/CE050p0374, 1996.

Uncles, R. J., Stephens, J. A., and Law, D. J.: Turbidity maximum in the macrotidal, highly turbid Humber Estuary, UK: flocs, fluid mud, stationary suspensions and tidal bores, *Estuarine, Coastal and Shelf Science*, 67, 30–52, doi:10.1016/j.ecss.2005.10.013, 2006


**Referee Comment 4.** I was not completely convinced by the definition of TMZ that you use, ie  $\text{NTU} > 1000$ . A TMZ in an estuary can be present even if the overall turbidity is lower than this value as it depends on the turbidity more upstream and downstream of it. The occurrence of TMZ in estuaries has been discussed a lot in literature, but I don't think that a certain turbidity value was proposed in order to have a TMZ. You refer to 'old' papers from Allen et al (1977) and Allen & Castaing (1981) where they have used a threshold for turbidity in a TMZ. What is the scientific basis of using such a threshold?

Indeed in any estuary, a TMZ is defined as a zone, where turbidity is higher than in upstream and downstream waters. However, for the Gironde estuary, there are pioneer works of Allen and Castaing that define the threshold of TMZ at  $1 \text{ g L}^{-1}$  based on numerous field determinations of suspended load distributions.

As our work is in the same fluvio-estuarine system, we use the same TMZ definition (in fact changing the threshold should have been justified), i.e. the TMZ is present when particle load values exceed  $1 \text{ g L}^{-1}$ . The critical point is in fact the turbidity – SPM relationship, as the automated stations measure turbidity and not particle load. Ongoing works on  $[\text{SSC}, \text{g L}^{-1}] = f(\text{Turbidity}, \text{NTU})$  curves based on particles collected at Bordeaux and Pauillac show that 1000 and 9999 NTU correspond to about 1 and 5-6  $\text{g L}^{-1}$  respectively (Schmidt, personal data, see figure). This justifies the use of 1000 NTU as the threshold of TMZ occurrence.



However this work could serve also to test the historical threshold. When the TMZ is installed at one of the four MAGEST station, we observe that the minimum value of tidally-averaged turbidity is always equal or higher to 1000 NTU, whatever the tidal range. This observation thus reinforces previous works.



#### b) SPECIFIC COMMENTS

**Referee Comment 1.** p2846 line 18-19: Mention here that the tides are semi-diurnal.

Done (p3 line 25)

**Referee Comment 2.** p2847 line 17-18: 'However, the suspended sediments dynamics in this estuarine region are largely unknown': this should not be in the a chapter that is intended to present facts.

Sentence suppressed

**Referee Comment 3.** p2848: I can live with NTU, but why are you not using mass concentration (mg/l). Is it because of a lack of calibration data? What is the correlation between NTU and mg/l? Is the correlation varying with season/discharge?

As already explained general comment 4, a work on  $[SSC, g L^{-1}] = f(\text{Turbidity, NTU})$  curves based on estuarine particles is ongoing. It is too preliminary to be presented here.

**Referee Comment 4.** p 2848 line 27-28 to p2489 line 1: Not clear what you mean with "to at least 70% of measured values for the considered period of time". Is it that 70% of the data during a tidal cycle have to be of good quality?

Automated stations measure turbidity each 10 min. unfortunately there are data missing due to different technical reasons. Normally the tidally-averaged turbidity relies on 75 successive measurements. Due to temporary stop in measurements or sensor failure, the number of effective measurements could be lower. We have tested that tidally-averaged turbidity are representative when it relies on, at least, 52 successive measurements.

As there is no question of the reviewer#2 on this point, we did not change the sentence.

**Referee Comment 5.** p2849 Fig 2 is not necessary.

As argued in the text, this is a key study for the management of the estuary. Since management policies use usually daily-averages, this figure is important for potentially users of this kind of data. We prefer to keep the figure because the result is not intuitive.

**Referee Comment 6.** p2849 line 5: "Previous works have defined the TMZ in the Gironde estuary by a SPM concentration  $> 1 \text{ gL}^{-1}$ . Is this near-bed, surface or vertical averaged SPM concentration?"

It is surface SPM concentration. The text is modified to clarify (p6 line 2).

**Referee Comment 7.** Last paragraph of §3.2: the results of the statistical analysis you describe here are not presented and discussed in the ms.

The results of the statistical analysis described in §3.2 were presented in the second paragraph of the section 4.3.1.

**Referee Comment 8.** 4.1 Hydrological trends: the paragraph is better suited in chapter 2 (study site) and chapter 3.2 (data treatment).

We prefer to keep the section §4.1 because it presents our own data treatment, calculated from data presented in chapter 3, and adapted for the time period of our study. The values presented in this section describe the hydrological regime of cases discussed in section §5, hence we consider essential to separate the description of the studied period from the general description of the estuary.

**Referee Comment 9.** Figure 3: quiet small and difficult to see. Possibly the figure will be more clear in the final version.

This is because of the orientation of the figure in the file. We suggest expanding this figure in landscape setup.

**Referee Comment 10.** Figure 4: Indicate the date of the example and the location in the caption of the figure. The figure is small and therefore not clear, e.g. sub-figure 4c (right) is hard to understand.

The location is included in the figure. The date is shown through the ticks. As suggested for Figure 3, the figure would benefit to be oversized.

**Referee Comment 11.** p2851 line 1: how do you know that 9999NTU is  $> 6 \text{ g/l}$ ? Add reference or explain.

The reference was already mentioned (p2848 line 5 of discussion version), however the correspondence NTU-SPM is suppressed here as the purpose was only to indicate that turbidity could be very high.

**Referee Comment 12. p2851 line 7:** Do you mean by mid-flood/mid-ebb mean water elevation? Is ebb and flood nicely correlated with tides, i.e. between LW→HW/HW→LW or is there a time shift? Maybe indicate in 'study site' to which part of the tidal cycle ebb and flood correspond.

We used mid-flood/mid-ebb for qualitatively denoting the moments of maximum current velocities. The terms are suppressed to avoid confusions.

**Referee Comment 13.** p2851 line 12: is the discharge peak for the Garonne?

Yes, Garonne is added in the text to clarify (p7 line 3).

**Referee Comment 14.** p2851 line 16: why do you call some floods 'peculiar' Is there not always a first flood after LW-periods? Better skip 'peculiar'. Can you show such a flood in a Figure?

The text is modified to clarify (p8 line 7). The different turbidity patterns associated to floods are detailed and graphically represented in Section §5.1.

**Referee Comment 15.** p2852 line 1-2: sentence is not needed.

The sentence is suppressed and included in a more appropriate position (now at p7 lines 15-16). It is indeed an, obvious but important, statement that needs to be repeated. There are still works in tidal environments that forget this point.

**Referee Comment 16.** p2852 line 15 and Fig 5: in the text you mention 'contrasting hydrological periods' in the figure you mention 'season'. I agree that hydrological regime is strongly seasonal, but are there not also wet periods in summer and dry ones in winter? Is the behaviour of the TMZ in the estuary similar in a wet/dry summer and a wet/dry winter period?

Section 4 is the "result section". We first detail and demonstrate the main marked seasonal trends in hydrological conditions in section 4.1. Based on these main "contrasting hydrological periods" we present only the associated main trends and statistics in turbidity (section §4.3). We precise in the figure caption the values of daily-discharges for these two periods. The paragraph was simplified for clarity (p9 lines 2-4).

But we agree that, apart from these main trends, there is inter-annual variability. It is the reason for which, after this overview (section 3 results), we discussed in deep the relationships between turbidity and river flow. We had showed and explained that, indeed, TMZ could be observed in the tidal Garonne during winter dry periods as in 2012 (see figure 6 by example).

**Referee Comment 17.** p 2854 line 7: showed → shown

Done

**Referee Comment 18.** p2855 line 7-8: 'probably remained TMZ-originated mud' → 'probably remains of a previous TMZ period'

Done

**Referee Comment 19.** p2855 line 27-28: What is different in 2010 as compared with dry and wet years? In the Table 1 several M(CC) pattern are indicated, however, in the text you write 'absence of CC pattern'.

M(CC) patterns are not CC pattern: see the table caption . However the text is modified to clarify (p12 lines 1-3).

**Referee Comment 20. p2856 line 9-10:** "The prediction of TMZ location is a challenge in the fluvial Gironde estuary". I would suggest to skip this part, especially the 'challenge'.

Done. "A challenge" is changed by "nowadays a need" (p12 line 12)

**Referee Comment 21.** p2856 line 12: 'presence of the TMZ': is it the presence or the position? See also my comment on the 1000 NTU as definition of TMZ. Possibly a weak TMZ is occurring?

Done. "Presence" is changed by "Position" (p12 line 15).

**Referee Comment 22.** Figure 3 caption: missing (e) Portets

Done

**Referee Comment 23.** Figure 4 caption: add date. What is a 'mean time step of river flow'? Do you mean measuring time interval? add 'one' before 'hour'.

Done

**Referee Comment 24.** Figure 9: Not clear, especially part B.

This is because of the size of the figure in the discussion version. We suggest enlarging this figure in the revised version.

**Referee Comment 25.** p2857 line 3-4: "On the opposite, the effect of tidal range is null during floods, when turbidity is associated to sediments transported from the watershed." Replace 'null' by 'small', not very clear from the figure 9.

"Null" is modified by "almost negligible" (p13 line2).

**Referee Comment 26.** p2858 line 12: "discharges between 200-300 m<sup>3</sup>/s". From the figure I would say discharges < 300 m<sup>3</sup>/s.

The purpose is to indicate the discharge threshold that corresponds of the observation of TMZ in Bordeaux, according to the figure 12. As explained in the previous paragraphs, there is also an influence of tidal range on turbidity that explained the range (min median max) in turbidity values at a given discharge (Figure 12). Therefore we prefer to keep "between 200-300 m<sup>3</sup>/s" that is more in agreement with the figure. Then below 200 m<sup>3</sup>/s, TMZ is always present. In a management perspective, this range is important.

**Referee Comment 27.** p2859 line 3: 'may lead' → 'may have lead'

Done

**Referee Comment 28.** p2860 line 7: 'especially since the 90': I see the duration increasing since the 80s.

Done

**Referee Comment 29.** p2860, line 9: see the general comment on climate change above. What is more important here climate change, climate variability (NAO) or human activity?

[See general comment 1](#)

**Referee Comment 30.** p2860 line 11: The volumes are 10 times smaller in Fig 15. What is the unit 'Hm<sup>3</sup>'?

[The value is the same in the text than in the figure. 30x10<sup>3</sup> Hm<sup>3</sup> is replaced by 3x10<sup>4</sup> to avoid confusions.](#)

**Referee Comment 31. p2860 line 15:** " As the TMZ is concentrated and persistent, the required water volume to expel it increases". Do you mean that the SPM concentration/turbidity is higher in the TMZ or that the length of the TMZ is smaller?

[The text is modified to clarify \(p16 line 1\).](#)

**Referee Comment 32.** p2860, line 20: has there been a deepening in the estuary and a change in tidal range and asymmetry? Not clear from the sentence. If yes, then an increase of SPM concentration in the estuary or the TMZ could also be due to a higher import from the sea. Is the origin of the mud terrestrial or also marine?

[The origin of mud is mainly terrestrial. It is written in the Section 2 with its correspondent reference \(p2847 line 6 of discursion version\). There is no works about the evolution of tidal range and asymmetry during the last decades, so we do not know if morphological changes have affected the sedimentary dynamics. The text is modified to clarify \(p16 lines 12-13\).](#)

## REFEREE #2

This article represents an important contribution to our understanding of the evolution of turbidity maxima in macrotidal estuaries under varying fresh water flow and tidal range. It is well laid out and clear, with a good structure. It is most useful to present a long term, high resolution data set in this way and to alert readers to the possibilities afforded by such a level of monitoring. Thus, I fully support publication.

My comments are generally minor and relate to the way in which the arguments are presented. I think readers will primarily be interested in the question of what the results mean for the future of the estuary and for management regimes. So, Section 5.3 on 'has the turbidity intensified' is important here and should be referred to in the abstract and aims.

[Done \(p1 lines 22-30 and p3 lines 16-17\)](#)

I am not in favour of the use of the terms 'installation' and 'expulsion' as, in my experience, these terms are not well known. Can I suggest 'Upstream migration' and 'downstream flushing'?

[Yes we admit that "installation" and "expulsion" terms are not usually. We replace these terms when we are referring the shift of the TMZ \(or mobile mud\) over the tidal rivers, considered in general. But our work is based on specific stations that are located at a specific location on the tidal rivers. It is the reason we prefer to use the terms "expulsion" and "installation", since it is denoting the TMZ](#)



presence or not in a specific location. We have introduced the definition of these terms (p6 lines 3-7) to clarify and avoid confusions.

Please be aware of the following paper on the Thames which I believe to be relevant: Mitchell S.B., Uncles R.J. and Akesson L. 2012. Observations of turbidity in the Thames estuary. Water and Environment Journal 26, 511-520

Yes indeed it is an important work that would have been cited. Done (p10 lines 13-14 and p13 lines 3-4)

I have tried to attach it to this comment, hopefully I have succeeded.

Please ensure to provide a clear key (legend) in Fig.15 to help interpretation of this figure.

Done

1 **Turbidity in the fluvial Gironde Estuary (S.-W. France)**  
2 **based on 10-year continuous monitoring: sensitivity to**  
3 **hydrological conditions**

4 **I. Jalón-Rojas, S. Schmidt and A. Sottolichio**

5 {UMR5805 EPOC, CNRS-Université de Bordeaux, Pessac, France}

6 Correspondence to: I. Jalón-Rojas (isabel.jalon-rojas@u-bordeaux.fr)

7

8 **Abstract**

9 Climate change and human activities impact the volume and timing of freshwater input to  
10 estuaries. These modifications in fluvial discharges are expected to influence estuarine  
11 suspended sediment dynamics, and in particular the turbidity maximum zone (TMZ). Located  
12 in the southwest France, the Gironde fluvial-estuarine systems has an ideal context to address  
13 this issue. It is characterized by a very pronounced TMZ, a decrease in mean annual runoff in  
14 the last decade, and it is quite unique in having a long-term and high-frequency monitoring of  
15 turbidity. The effect of tide and river flow on turbidity in the fluvial estuary is detailed,  
16 focusing on dynamics related to changes in hydrological conditions (river floods, periods of  
17 low-water, inter-annual changes). Turbidity shows hysteresis loops at different time scales:  
18 during river floods and over the transitional period between the installation and expulsion of  
19 the TMZ. These hysteresis patterns, that reveal the origin of sediment, locally resuspended or  
20 transported from the watershed, may be a tool to evaluate the presence of remained mud.  
21 Statistics on turbidity data bound the range of river flow that promotes the upstream migration  
22 of TMZ installation in the fluvial stations. Whereas the duration of the low discharge period  
23 mainly determines the TMZ persistence, the freshwater volume during high discharge periods  
24 explains the TMZ concentration at the following dry period. ~~Hydrological indicators of the~~  
25 ~~persistence and turbidity level of the TMZ are also defined.~~ The ~~long-term~~ evolution of these  
26 two hydrological indicators of TMZ persistence and turbidity level since 1960 confirms the  
27 ~~influenceeffect~~ of discharge decrease on the intensification of the TMZ in tidal rivers;  
28 ~~andboth~~ provides a tool to evaluate future scenarios.

29

# 1 1 Introduction

2 Macrotidal estuaries are highly variable systems as result of the strong influence of both tides  
3 and river discharge. In particular dynamics of suspended particulate matter (SPM) and the  
4 occurrence of a turbidity maximum zone (TMZ) are complex and difficult to predict (Fettweis  
5 et al., 1998; Mitchell and Uncles, 2013). Different processes can induce the formation of the  
6 TMZ (for details see Allen et al., 1980; Dyer, 1988; Jay and Musiak, 1994; Talke et al.,  
7 2009). This highly concentrated zone plays an important role on estuarine morphodynamics.  
8 | Sediment depositions from the TMZ, ~~termed fluid mud,~~ may generate gradual accretion of  
9 bed and banks (Pontee et al., 2004; Schrottke et al., 2006; Uncles et al., 2006). Therefore,  
10 many estuaries require regular dredging against ongoing siltation events to maintain the depth  
11 of navigation channels.

12 Quite recently, considerable attention has been paid to evaluate the effect of climate change  
13 (Fettweis et al., 2012) and human interventions (Schuttelaars et al., 2013; Winterwerp and  
14 Wang, 2013; Yang et al., 2013; De-Jonge et al., 2014) on natural distribution of SPM in  
15 estuaries. There are numerical evidences linking freshwater abstractions to an increased  
16 potential for up-estuary transport (Uncles et al., 2013). Nevertheless the effects of shifts in  
17 freshwater inflow on sediment regime are not yet totally understood (Mitchell and Uncles,  
18 2013). The longitudinally TMZ migration as result of seasonal variability of runoff was well  
19 described in many estuaries (Grabemann et al., 1997; Uncles et al., 1998; Guézennec et al.,  
20 1999). However, the effect of floods or long-term hydrological variability on sediment  
21 dynamics is scarcely documented. Only Grabemann and Krause (2001) showed differences in  
22 SPM concentrations of the TMZ in the Weser estuary between a dry and a wet year, although  
23 | the gaps in data hamper a detailed analysis. The transitional periods of upstream migration  
24 and downstream flushing ~~installation and expulsion~~ of the TMZ and of its associated mobile  
25 mud in fluvial sections have also not been detailed. These limitations are partly due to the  
26 absence of relevant long-term datasets, which are not so common in estuaries (Garel et al.,  
27 2009; Contreras and Polo, 2012).

28 The Gironde fluvio-estuarine system (SW France) is quite unique in having a long-term and  
29 high-frequency monitoring of water quality. This estuary presents a pronounced TMZ so far  
30 documented in the lower and central reaches (Allen and Castaing, 1973; Allen et al., 1980;  
31 Sottolichio and Castaing, 1999). The Gironde watershed has the largest water structural  
32 deficit in France (Mazzega et al., 2014). Warming climate over the basin induces a decrease

1 in mean annual runoff, a shift to earlier snow melting in mountainous areas and more severe  
2 low-flow conditions (Hendrickx and Sauquet, 2013). In addition, according to data of the  
3 agricultural census, irrigated areas have duplicated its surface in several regions of the  
4 watershed between 1988 and 2000, promoting strong water storage and abstractions. This  
5 context makes the Gironde estuary a good example to evaluate how changes in freshwater  
6 regime may affect the estuarine particle dynamic.

7 The goal of this work is to analyse the response of fine sediments to hydrological fluctuations,  
8 based on a 10-years, high-frequency database of turbidity in the fluvial Gironde estuary, in  
9 order to:

- 10 1. Document the trends of SPM at all representative time scales, from intratidal to  
11 interannual variability.
- 12 2. Analyse the role of floods on the sedimentary dynamic of the tidal rivers.
- 13 ~~3. Analyse~~Discuss—the influence of hydrological conditions on TMZ features  
14 (~~installation/expulsion~~upstream migration, downstream flushing, concentration,  
15 persistence).
- 16 ~~3.4.~~ Discuss the effect of the long-term decrease of runoff in the upstream  
17 intensification of the TMZ.

18

## 19 **2 The study site**

20 With a total surface area of 635 km<sup>2</sup>, the Gironde is a macrotidal fluvial-estuarine system  
21 located on the Atlantic coast (South-West of France, Fig. 1). The estuary shows a regular  
22 funnel shape of 75 km between the mouth and the junction of the Garonne and the Dordogne  
23 rivers. Tidal rivers present a single sinuous channel with weak slopes and narrow sections  
24 (about 300 m, 250 m and 200 m at Bordeaux, Portets and Libourne respectively, Fig. 1). At  
25 the Gironde mouth, tides are semidiurnal and the mean neap and spring tidal ranges are  
26 respectively 2.5 and 5 m (Bonneton et al., 2015). The tidal wave propagates up to 180 km  
27 from the estuary mouth. Thereby, the uppermost limits for the dynamic tidal zone are (Fig. 1):  
28 La Réole for the Garonne River (95 km from the river confluence); and Pessac for the  
29 Dordogne River (90 km from the river confluence). As the tide propagates upstream, tidal  
30 currents undergo an increasing ebb-flood asymmetry (longer and weaker ebb currents; shorter  
31 and stronger flood currents) and the wave is amplified (Allen et al., 1980). The tidal wave

1 reaches its maximum value at about 125 km from the mouth (Bonneton et al., 2015), before  
2 decaying in the fluvial narrow sections.

3 The tidal asymmetry toward upstream and the subsequent tidal pumping coupled to density  
4 residual circulation develop a turbidity maximum zone (TMZ). The high tidal ranges and the  
5 great length of the estuary promote a highly turbid TMZ (Uncles et al., 2002). In surface  
6 waters of the middle estuary, SPM concentrations range between 0.1 and 10 g L<sup>-1</sup> according  
7 to Sottolichio and Castaing (1999). Estuarine suspended sediments have a dominant terrestrial  
8 origin and are mainly composed of clays (45–65%) and silts (Fontugne and Jounneau, 1987).  
9 SPM residence time is comprised between 12 and 24 months, depending on river discharge  
10 (Saari et al., 2010). Freshwater inflow moves the TMZ along the estuary axis: during high  
11 river flow the TMZ moves down-estuary and vice versa (Castaing and Allen, 1981). There is  
12 also a secondary steady TMZ in the middle estuary possibly related to a high dynamic zone,  
13 so called the ‘erosion maximum zone’ (Allen et al., 1980; Sottolichio and Castaing, 1999). At  
14 slack water, sediment deposition occurs on the river bed and banks. In the channel fluid mud  
15 can form elongated patches, with concentrations up to 300 g L<sup>-1</sup> (Allen, 1971).

16 Contrary to the middle estuary, the tidal Garonne and Dordogne Rivers are still poorly  
17 documented. Measurements over a maximum of 3 days (Romaña 1983; Castaing et al. 2006)  
18 and satellite images (Doxaran et al., 2009) ~~showed~~revealed the seasonal presence of the TMZ  
19 ~~in the tidal rivers~~ during the summer-autumn period. Brief field observations in September  
20 2010 (Chanson et al, 2011) showed the presence of low consolidated mud deposits upstream  
21 Bordeaux. In the following, and according to Uncles et al. (2006), the term mobile mud is  
22 used for these low consolidated mud deposits that are easily erodible, and likely to shift  
23 seasonally with the TMZ. However, the suspended sediments dynamics in this estuarine  
24 region are largely unknown.

25

## 26 **3 Materials and Methods**

### 27 **3.1 The multiyear high-frequency monitoring system**

28 The Gironde estuary counts on an automated continuous monitoring network, called  
29 MAGEST (MAREL Gironde ESTuary), to address the current and future estuarine water  
30 quality. MAGEST network includes four sites (Fig. 1): Pauillac in the central estuary (52 km  
31 from the mouth); Libourne in the Dordogne tidal river (115 km from the mouth); and

1 Bordeaux and Portets in the Garonne tidal river (100 and 140 km from the mouth  
2 respectively). The automated stations record dissolved oxygen, temperature, turbidity and  
3 salinity every ten minutes at 1 m below the surface. In addition, an ultrasonic level controller  
4 measures the water depth in the stations of Bordeaux, Portets and Libourne. The turbidity  
5 sensor (Endress and Hauser, CUS31-W2A) measures values between 0 and 9999 NTU with a  
6 precision of 10%. The saturation value (9999 NTU) of turbidity sensor corresponds to about 6  
7  $\text{g L}^{-1}$  (Schmidt et al., 2014). One may refer to Etcheber et al. (2011) for a description of the  
8 MAGEST survey program, for the technical features of monitoring system and for examples  
9 of the trends in measured parameters; and to Lanoux et al. (2013) for a detailed analysis of  
10 oxygen records.

11 The first implemented station was Pauillac the 15 June 2004. Acquisition at Portets and  
12 Libourne stations began the 16 November 2004 and at Bordeaux stations the 1 March 2005.  
13 Portets station was stopped the 11 January 2012. The severe environmental conditions,  
14 electrical / mechanical failures and sensor malfunctions could cause missing or wrong data.  
15 Therefore the database needed a cleaning for erroneous values in turbidity. By example 9999  
16 NTU correspond to saturation values, but also to sensor errors in, these later need to be  
17 removed. A routine under Matlab was developed to retain only turbidity values corresponding  
18 to saturation. The validated database of turbidity corresponds to 1.223.486 data points  
19 recorded between 2005 and mid-2014. This corresponds to a rate of correct operating of 71%,  
20 70%, 70% and 57% for Bordeaux, Portets, Libourne and Pauillac stations respectively.

21 In addition, two tide gauges, managed by the port of Bordeaux (Grand Port Maritime de  
22 Bordeaux), record tide height at Pauillac and Bordeaux every 5 minutes. Hydrometric stations  
23 record each 1 to 24 hours discharges of the Dordogne River (Pessac ; Lamonzie Saint Martin)  
24 and of the Garonne River (La Réole ; Tonneins) (Fig. 1). Data are available on the national  
25 web site: <http://www.hydro.eaufrance.fr/>.

## 26 **3.2 Data treatment**

27 Turbidity was analysed as function of river flow and water height at different time scales. To  
28 better identify intertidal trends, we calculated tidal-averaged turbidity with its corresponding  
29 tidal range. In order to avoid biased averaged values, we only consider the tidal averages  
30 corresponding to at least 70% of measured values for the considered period of time. Since  
31 management directives are often based on daily values, tidally and daily averages were

1 compared. Figure 2 compares both turbidity averages and shows a very good agreement  
2 between the two calculations ( $R^2=0.993$ ). Previous works have defined the TMZ in the  
3 Gironde estuary by a SPM concentration  $> 1 \text{ g L}^{-1}$  in surface (Allen et al., 1977; Castaing and  
4 Allen, 1981), which corresponds to a turbidity of about 1000 NTU. We call TMZ installation  
5 and expulsion the transitional periods where turbidity oscillates around 1000 NTU in a given  
6 station, during the TMZ upstream and downstream migration (see Figure 3). River floods are  
7 defined by a daily increase of the Garonne discharge higher than  $480 \text{ m}^3 \text{ s}^{-1}$  (percentile 75 of  
8 river flow during the study period). A time shift was added to discharge time series for the  
9 study of floods, since hydrometric stations are located tens of kilometres upstream of the  
10 MAGEST ones. It has been estimated based on the velocity of the peak-floods peaks between  
11 two hydrometric stations.

12 We performed statistical analysis on tidal-averaged data. We compared turbidity values  
13 according to stations (Portets, Bordeaux, Libourne and Pauillac), period (months, and tidal  
14 range), and their interactions (e.g. station within period), by performing analysis of variance.  
15 We used parametric test (T-Test and ANOVA) when datasets or its their transforms (like log  
16 or cubic root) met the normality and homoscedasticity criteria. Otherwise we used non-  
17 parametric tests (Mann-Whitney U and Kruskal-Wallis tests). In the following, we refer to  
18 “significantly different” datasets when these tests on tidal-averaged data were significant at  $p$   
19  $< 0.5$ . These tests were carried out using STATA software (v. 12.1, StataCorp, 2011).

20

## 21 **4 Results**

### 22 **4.1 Hydrological trends**

23 The Gironde estuary drains a watershed of  $81000 \text{ km}^2$ , (Fig. 1.a) strongly regulated by dams  
24 and reservoirs. The Garonne and the Dordogne Rivers contributes respectively tofor 65% and  
25 35% of the freshwater input. Historical records reveal drastic changes in hydrological  
26 conditions: the annual mean discharge (Garonne + Dordogne) is decreasing, flood events are  
27 increasingly scarce and drought periods are becoming more durable. In the period between the  
28 60's and the 80's, the mean annual discharge was  $1000 \text{ m}^3 \text{ s}^{-1}$ . In contrast, during the studied  
29 period (January 2005 - July 2014), the mean annual discharge was  $680 \text{ m}^3 \text{ s}^{-1}$  (Fig. 3.A). For  
30 this period, the inter-annual variability in freshwater inflow was also remarkable: the driest  
31 year was 2011 with a mean discharge of  $433 \text{ m}^3 \text{ s}^{-1}$  and the wetter one was 2013 with a total

1 mean discharge of  $961 \text{ m}^3 \text{ s}^{-1}$ . River discharge varies also seasonally reaching the highest  
2 values in January to February and the ~~small~~lowest in August to September. For the studied  
3 period, mean discharges were  $720 \text{ m}^3 \text{ s}^{-1}$  in winter (December 21 to March 20) and  $190 \text{ m}^3 \text{ s}^{-1}$   
4 (June 21 to September 20) for the Garonne River (380 and  $105 \text{ m}^3 \text{ s}^{-1}$  for the Dordogne  
5 River).

6 Tides are semidiurnal (the main harmonic component is the  $M_2$ ) with a period of 12 h 25 min.  
7 Between January 2005 and July 2014, the ~~minimal~~, mean, minimal and maximal values of  
8 tidal ranges were respectively about 4.1, 1.9 and 6.1 m at Pauillac and about 4.9, 2.5 and 6.6  
9 m at Bordeaux (see the whole time series in Fig. 3.B). Spring and neap tides were defined as  
10 the tidal cycles which tidal range is respectively above the percentile 75 and below the  
11 percentile 25. These values were about 3.5 (p25) and 4.7 (p75) at Pauillac, and about 4.3  
12 (p25) and 5.4 (p75) at Bordeaux.

## 13 **4.2 Short-term variability in turbidity**

14 Figure 4 presents examples of high frequency (10 minutes) data recorded at Bordeaux under  
15 two contrasted conditions of fluvial discharge. Continuous measurements reveal turbidity  
16 patterns related to tidal cycles, and to changes in fluvial discharges. Only such a continuous  
17 record can capture turbidity signatures of a flood that often occurs for a few hours.

### 18 **4.2.1 Tidal cycles**

19 The first selected dataset (Fig. 4.I) corresponds to a low-water period: the Garonne discharge  
20 was below  $120 \text{ m}^3 \text{ s}^{-1}$ . Turbidity shows a large range of values between 740 and 9999 NTU,  
21 testifying the presence of the TMZ in the tidal river. It is noticeable that turbidity is higher  
22 than the saturation value, ~~i.e.  $> 6 \text{ g L}^{-1}$~~ ; during several hours per tidal cycle. These raw data  
23 illustrate the short-term changes in turbidity due to deposition-resuspension processes induced  
24 by changes in current velocities throughout the tidal cycles. This pattern was already reported  
25 in the central estuary (Allen et al., 1980; Castaing and Allen, 1981). Fig. 4.I.c relates turbidity  
26 and water level of the above raw data, showing more clearly the intratidal patterns: two  
27 turbidity peaks at mid-flood and mid-ebb, due to the resuspension by the maximum current  
28 velocities. In contrast minimum turbidity values are always recorded at high tide and low tide  
29 due to deposition processes.



## 1 4.2.2 Flood events

2 The second selected dataset (Fig. 4.II) represents the turbidity signal related to a spring flood  
3 with a discharge peak of the Garonne River at  $1730 \text{ m}^3 \text{ s}^{-1}$ . As shown in the middle and lower  
4 panels, throughout river floods turbidity is the lowest during rising tide when tidal currents  
5 are against river flow; from high tide, river sediments are transported downstream, turbidity  
6 values begin to increase and the SPM peak usually occurs between mid-ebbing and low tide.  
7 ~~Peculiar floods, specially the f~~First flood just after low-water periods can, present a turbidity  
8 peak also at mid-rising tide. These peaks are associated with local resuspension processes and  
9 their occurrences are likely to give an indication of the existence of remained mud trapped in  
10 the tidal rivers.

11 Table 1 collects maximum discharge value and its associated maximum turbidity (when  
12 recorded) of each flood event at Bordeaux and Portets stations. Flood events were identified  
13 in the time series of river discharge in figure 3.A. The associated turbidity peaks were  
14 calculated as the maximum of the turbidity values at low tide (fluvial signature) in order to  
15 consider only the sediments transported by river flow. As shown in Table 1, turbidity maxima  
16 during flood events are 5 to 30 times lower compared to TMZ maximum values (50% of the  
17 recorded floods present a maximum turbidity  $<1000 \text{ NTU}$ ). ~~Only such a continuous record~~  
18 ~~can capture turbidity signatures of a flood that often occurs for a few hours.~~

## 19 4.3 Long-term variability in turbidity

### 20 4.3.1 From fortnightly to seasonal variability

21 The 10-year time series of tidal averaged turbidity (Fig. 3) reveals short oscillations related to  
22 neap-spring tide cycles and seasonal trends induced by hydrology. Maximum turbidity values  
23 are recorded during spring tides, since higher current velocities favour the resuspension of  
24 sediments (Allen et al., 1980). The highest turbidities occur during low discharge  
25 periodswaters (usually between July and November) in the up estuary waters (Fig3.D, E and  
26 F) due to the upstream displacement of the TMZ. Turbidity is usually minimal in spring after  
27 the flood period. In the middle estuary (Fig 3.C) seasonal changes are more moderate and  
28 show an inverse trend. This is due to the existence of a permanent TMZ in this estuarine zone,  
29 which is possibly related to a mud-trapping zone (Sottolichio and Castaing, 1999).

1 Figure 5 summarizes the main characteristics (mean, percentiles) of turbidity to compare the  
2 four stations during high (February) and low (August) river discharges~~contrasted hydrological~~  
3 ~~periods~~ and tidal ranges. ~~Differences between periods of high and low river discharges are~~  
4 ~~obvious~~. In the fluvial stations (Bordeaux, Portets, Libourne), ~~In these stations~~, turbidity in  
5 August is significantly ( $p < 0.001$ ) higher than in February: mean values in August are 8, 27  
6 and 54 times higher than in February at Bordeaux, Portets et Libourne, respectively. By  
7 contrast, at Pauillac station, in the central estuary, turbidity remains relatively high throughout  
8 the year (see Fig. 3). However, turbidity in August is significantly lower ( $p < 0.0000001$ ) than  
9 in February, when TMZ moves upstream. Turbidity values are also significantly different  
10 between the three fluvial stations in both dry ( $p < 0.0018$ ) and wet ( $p < 0.0001$ ) months. Summer  
11 turbidity values at Bordeaux and Libourne are higher than at Portets and Pauillac, reaching  
12 values above 7500 NTU. In February, turbidity is lower in the most upstream stations, with  
13 mean tidally-averaged turbidity values of 1322, 401, 93 and 52 NTU at Pauillac, Bordeaux,  
14 Portets and Libourne, respectively. Turbidity at high tidal range is significantly (e.g.  
15  $p < 0.000025$  at Pauillac,  $p < 0.017$  at Libourne) higher than at low tide at all station in August:  
16 respectively for Pauillac, Bordeaux, Portets and Libourne, mean turbidity at high tide was 2.7,  
17 2.3, 1.7 and 1.6 times higher compared to low tide. However, in February tidal range does not  
18 induce significant differences in turbidity at the most upstream stations of Portets ( $p = 0.22$ )  
19 and Libourne ( $p = 0.37$ ). Only Pauillac and Bordeaux stations present turbidity values  
20 significantly higher ( $p < 0.000001$ ) at high tide than at low tide during this month.

### 21 **4.3.2 Interannual variability**

22 The observation of the entire dataset of tidally-averaged turbidity evidences a strong  
23 interannual variability in SPM in the fluvial Gironde estuary. Figure 3 allows to appreciate  
24 marked differences in the concentration and in the duration of the TMZ for the monitored  
25 years at Bordeaux, Portets and Libourne. The maximum turbidity values exceeded 7200 NTU  
26 in the years 2010, 2011 and 2012 at Bordeaux and in the years 2010 and 2012 at Libourne. By  
27 contrast, during the year 2008 tidal-averaged turbidity was always below 6700 and 4400 NTU  
28 at Bordeaux and Libourne respectively. Portets station is less documented: tidal-averaged  
29 turbidity maxima ranged between 4730 and 6880 NTU (years 2009 and 2006, respectively).  
30 The durations of the TMZ occurrence ( $\text{Duration}_{\text{TMZ}}$ ) were calculated per year as the number  
31 of days during which tidal-averaged turbidity overtakes 1000 NTU (Fig. 6). In general, the  
32 TMZ is less present for the more upstream reaches. Annual durations decrease from Bordeaux

1 (varying between 93 and 259 days; years 2013 and 2011, respectively), to Portets (varying  
2 between 91 and 171 days; years 2006 and 2008, respectively), and to Libourne (varying  
3 between 33 and 143 days; years 2007 and 2011, respectively). The TMZ appeared also during  
4 dry winters (striped bars in Fig. 6) like in 2012 (39 days at Bordeaux and 6 days at Libourne).

5

## 6 **5 Discussion**

7 The presence of TMZ (duration, turbidity level, hibernal occurrence) is more marked and  
8 better documented in Bordeaux waters. The following discussion is dedicated to the tidal  
9 Garonne.

### 10 **5.1 Mobile mud downstream flushing~~expulsion~~ rhythm based on sediment** 11 **dynamics during floods**

12 River floods expel the TMZ (and its associated mobile mud) from fluvial to middle estuary as  
13 shown in figure 3. Mitchell et al. (2012) related this downstream flushing to a lack of  
14 settling at high slack water during high river discharge. According to Castaing and Allen  
15 (1981), the repetition of strong flood events, along with spring tides, favours the flushing  
16 expulsion of a part of the TMZ toward the sea. Floods also transport eroded sediments from  
17 the watershed that contribute to the TMZ. Identifying both processes is important to discuss  
18 the role of floods on the sedimentary budget of tidal rivers. The literature proposes the  
19 hysteresis-based analysis to search specific patterns of sediment transport in rivers (Williams,  
20 1989; Klein, 1984; López-Tarazón et al., 2009). The relative position of sediment sources  
21 within the catchment is analysed through the flow sediment hysteresis shapes (clockwise or  
22 counterclockwise). In short, anti-clockwise loops correspond to a transport of sediments from  
23 upstream distant sources, while clockwise loops occur when the sediment source is the  
24 channel itself or adjacent areas. Based on the MAGEST turbidity database, flow sediment  
25 hysteresis shapes were systematically analysed for the 26 floods recorded at Bordeaux (13 at  
26 Portets; Table 1). Only the values at low tide were used to trace the loops in order to preserve  
27 the fluvial signal and to avoid the impact of local resuspension by tidal currents on the levels  
28 of turbidity. The succession of hysteresis shapes over several years follows a seasonal pattern  
29 in the Garonne tidal river (Table 1, illustrated for the year 2013 in Fig. 7). In the case of  
30 Bordeaux:

- 1 • The first floods that occur at the end of the low ~~discharge period~~water and  
2 expel the TMZ down estuary show clockwise (C) hysteresis loops (e.g. f3, f8,  
3 f11, f24, Table 1; f24 in Fig. 7). This indicates the advection of resuspended  
4 sediments from the close bed and banks. ~~During the occurrence of~~When  
5 TMZ is present in the fluvial section, there is an accretion of sediments that  
6 remain after the TMZ-~~downstream flushing~~expulsion. This mud is eroded by  
7 river flood.
- 8 • Winter and some early spring floods present mixed (M) hysteresis curves, i.e.,  
9 clockwise loops with a counterclockwise loop around the flood peak (f25 in  
10 Fig. 7). Some events show a predominance of the clockwise loop (M(C), e.g.  
11 floods f1, f4, f17, Table 1), or of the counterclockwise loop (M(CC), e.g.  
12 floods f15, f18 Table 1). This pattern suggests the presence of local sediments,  
13 probably remained of a previous TMZ period~~TMZ-originated mud~~, and also  
14 the transport of sediment from remote-~~areas~~places. The predominant loop  
15 could be interpreted in term of proportion of each sediment source.
- 16 • Spring floods follow counterclockwise (CC) hysteresis patterns (e.g. floods f2,  
17 f7, f10, f28, Table 1; f26, f27, f28 in Fig. 7). This means that sediments are  
18 mainly transported from upstream areas; the TMZ-derived mud is expected to  
19 be totally expelled.

20 A similar seasonal evolution of hysteresis also exists at Portets, but it is subtler probably in  
21 relation with its upstream position: the flow sediment curves of the first floods are mixed and  
22 counterclockwise loops already appear in winter (Table 1). For example, the flood f1 (31-01-  
23 2006) presented a mixed, but predominantly C, loop at Bordeaux indicating dominant local  
24 sediments, whereas the simultaneous CC loop at Portets traced a distant origin of sediments.  
25 The TMZ-originated mud is less present locally and more quickly expelled in the uppermost  
26 section.

27 Therefore, hysteresis curves are indicators of the presence of mobile mud in tidal rivers, as  
28 schematized in figure 8, and permit to discuss its rhythm of ~~downstream flushing~~expulsion for  
29 different hydrological conditions and positions along the tidal river axis. During the wet years  
30 2008 and 2009 the mud disappeared from Portets and Bordeaux in the beginning of winter  
31 with the first floods. In contrast, mud was only expelled in May during the dry years 2007 and  
32 2012. In the case of the period from January to Mayyear 2010, ~~we assume~~the observation of

1 mixed patterns shows that mobile mud ~~it~~ was not completely flushed (Table 1): this is  
2 explained by the absence of major floods until the following upstream migration of the  
3 TMZ~~present all the year in the absence of CC pattern and marked floods (Table 1).~~

4 This first detailed study of 10-year continuous turbidity records suggests that deposition of  
5 mobile mud also occurs in the tidal Gironde, as already reported in the central estuary (Allen,  
6 1971; Sottolichio and Castaing, 1999). Two-third of the floods from 2005 to mid-2014  
7 contributed to the progressive downstream flushing~~expulsion~~ of fluid~~mobile~~ mud from  
8 Bordeaux. As turbidity values associated to floods are significantly lower than those in the  
9 TMZ, this demonstrates that floods play a more important role in flushing sediment  
10 downstream than in increasing the TMZ concentration.

## 11 **5.2 Occurrence of the TMZ in the tidal river**

12 The prediction of TMZ location is nowadays a challengeneed in the fluvial Gironde estuary  
13 and of particular interest to improve regional sediment management. The present work, based  
14 on turbidity measurements over the last 10 years, reveals a seasonal occurrence of the TMZ at  
15 Portets, 40 km upstream Bordeaux. The position~~resenee~~ of the TMZ along the longitudinal  
16 axis depends mainly on the freshwater inflow in major macrotidal European estuaries (e.g.  
17 Weser, Seine, Scheldt, Humber, see Mitchell, 2013). To better understand the relationships  
18 between turbidity and river flow in the Garonne tidal river, figure 9 shows the tidally (A) and  
19 daily (B) averaged turbidity as a function of river flow (3-day average). In Pauillac (central  
20 estuary) the dependence on river flow is the weakest: turbidity is slightly lower when the  
21 TMZ elongates to the upper reaches, but also when floods push suspended sediments  
22 seaward. In the tidal Garonne River, turbidity increases with decreasing river flow for  
23 discharges lower than about 1000 and 600 m<sup>3</sup> s<sup>-1</sup> at Bordeaux and Portets respectively. At  
24 Bordeaux, the maximum turbidity values remain rather constant in the range 50-200 m<sup>3</sup> s<sup>-1</sup>  
25 because of the saturation of the turbidity sensor. For the highest discharges (>1500 m<sup>3</sup> s<sup>-1</sup>),  
26 turbidity increases up to about 2450 NTU with increasing river flow.

27 Determining a precise discharge threshold of the TMZ installation per station is tricky, due to  
28 the large variability in turbidity, more than one order of magnitude at 200 m<sup>3</sup> s<sup>-1</sup>, partly  
29 explained by the tidal range and the locally-available sediment stock. During spring tides,  
30 current velocities and thus bed shear stress are stronger, promoting sediment resuspension and  
31 hence higher turbidity. This process is visible and quantifiable in Fig. 9.A for different

1 discharges. The dependence is strong when the TMZ is installed in the fluvial estuary at low  
2 discharge period~~water~~. On the opposite, the effect of tidal range is almost negligible~~and~~  
3 during floods, when there are no sediments to resuspend from the river bed, as suggested  
4 Mitchell et al. (2012) for the Thames Estuary, and turbidity is then associated to sediments  
5 transported from the watershed. To detail the relationship between these variables, figure 10  
6 presents turbidity as a function of tidal range for raising-falling neap-spring cycles during the  
7 periods of installation, presence and expulsion of the TMZ at Bordeaux in 2009 (see periods  
8 in Fig. 3). During the TMZ installation (a) and when the TMZ is completely installed (b),  
9 turbidity was lower during neap-spring tide transition than during the spring-neap- tide one.  
10 This hysteresis pattern, already observed in other estuaries, is explained by the consolidation  
11 of deposited material during neap tides, when currents velocities and resuspension are lower  
12 (Grabemann et al., 1997; Guézennec et al., 1999; Grabemann and Krause, 2001). During the  
13 installation of the TMZ the maximum turbidity occurs 4-5 tidal cycles after the maximum  
14 tidal range (Fig. 10, curve a). This is explained by a gradual increase in sediment availability  
15 at the riverbed as river discharge decreases, promoting the upstream shift of the TMZ. During  
16 the TMZ expulsion period, following river flood, the hysteresis curve is reversed (Fig. 10,  
17 curve c), the sediments are progressively resuspended and expelled down estuary and the  
18 stock decreases. These behaviours were also found in Portets station.

19 Differences on turbidity between the periods of decreasing and increasing river flow are also  
20 notable in the fluvial estuary (Fig. 9.B). In the tidal Garonne, for same discharge intensity, the  
21 smallest turbidity values are always associated with the TMZ installation (decreasing  
22 discharge) and the highest values during the TMZ expulsion (increasing discharge). This  
23 indicates that the discharge turbidity curve follows a clockwise hysteresis over the transitional  
24 periods of installation and expulsion of the TMZ (Fig. 11). For example, for a river flow of  
25  $500 \text{ m}^3 \text{ s}^{-1}$ , daily-averaged turbidity at Bordeaux was 8 to 50 times higher during the falling  
26 discharge curve in the year 2009. Such hysteresis were also recorded in the Weser estuary  
27 (Grabemann et al., 1997; Grabemann and Krause, 1998), suggesting an association with  
28 delays in TMZ movements or with the local sediment inventory. We explain these hysteresis  
29 patterns by an accumulation of sediments during the presence of the TMZ that need large  
30 river flow to be expelled. This agrees with the existence a deposition flux of mud remained at  
31 upper reaches after the passage of the TMZ.

1 A distinction in turbidity values corresponding to the periods of TMZ installation or expulsion  
2 is then necessary to precise the discharge threshold of the TMZ installation in tidal rivers.  
3 Figure 12 summarizes the distribution of turbidity values as a function of river flow (intervals  
4 of  $30 \text{ m}^3 \text{ s}^{-1}$ ) during the transitional periods of installation and expulsion of the TMZ at  
5 Bordeaux station. It allows to associate a river discharge range to a probability of TMZ  
6 installation (as defined by tidal average turbidity  $>1000 \text{ NTU}$ , Fig. 12.A) or TMZ expulsion  
7 (tidal averaged turbidity  $<1000 \text{ NTU}$ , Fig. 12.B). The discharges between  $200\text{-}300 \text{ m}^3 \text{ s}^{-1}$   
8 present the highest probabilities to promote the TMZ installation. The expulsion threshold is  
9 less bounded since the intensity and the amount of first floods are variable. Discharges greater  
10 than  $350 \text{ m}^3 \text{ s}^{-1}$  promote the TMZ expulsion and discharges above over  $610 \text{ m}^3 \text{ s}^{-1}$  ensure the  
11 complete expulsion.

### 12 **5.3 Has the TMZ intensified in the tidal Garonne?**

13 In the absence of historical turbidity data in tidal rivers, it is difficult to judge the evolution of  
14 the TMZ. There are only few limited available dataset, issued from field campaigns. By  
15 example in September 1960, SPM concentrations of surface waters at Bordeaux range  
16 between  $1 \text{ g L}^{-1}$  (mean tide) and  $2.5 \text{ g L}^{-1}$  (spring tide) (Castaing et al., 2006). At Portets, SPM  
17 concentration reached  $2.5 \text{ g L}^{-1}$  just before high tide for spring tide, while at mean and neap  
18 tides, SPM concentrations was always bellow  $1 \text{ g L}^{-1}$ . Romaña (1983) presented also quasi-  
19 instantaneously turbidity measurements implemented by helicopter along the estuary for 3  
20 days of contrasted hydrological conditions in the years 1981-1982. At low-water TMZ  
21 appeared 10 km upstream Portets reaching a maximum value of  $1.7 \text{ g L}^{-1}$ . Although these  
22 values seem lower than current turbidity trends, the extremely limited measurement periods  
23 and the difference in sampling points prevent to draw conclusions about a possible TMZ  
24 intensification in the tidal river. However, the remarkable dependence of turbidity to river  
25 flow in the fluvial section (Fig. 9) suggests that the decreasing trend in river flow in the last  
26 decades (Section 4.1) may ~~have lead to promoted~~ an upstream intensification of the TMZ.

27 The 10-year dataset of the MAGEST stations of Bordeaux and Portets was used to evaluate  
28 the impact of hydrological conditions on TMZ (turbidity level and persistence) in the tidal  
29 Garonne. The annual maximum turbidity value ( $\text{Turbidity}_{\text{max}}$ , as an indicator of turbidity  
30 level) and the duration ( $\text{Duration}_{\text{TMZ}}$ ) of the TMZ were compared to three hydrological  
31 characteristics:



- 1 | 1. Duration<sub>LDW</sub> : the duration of low-~~discharge period~~water, calculated as the number of  
2 | days per year river flow is below 250 m<sup>3</sup> s<sup>-1</sup> at Bordeaux (Fig. 12) and 160 m<sup>3</sup> s<sup>-1</sup> at  
3 | Portets; these values were evaluated as the mean critical river flows above which the  
4 | TMZ is installed in two stations.
- 5 | 2. Vol<sub>HDW</sub> : the river water volume passed during the previous high ~~discharge~~ water  
6 | period, i.e., between the last expulsion and the reinstallation of the TMZ;
- 7 | 3. Vol<sub>TMZ</sub> : the river water volume passed during the presence of the TMZ at the  
8 | considered station.

9 | The Duration<sub>TMZ</sub> in both stations is well correlated to the Duration<sub>LDW</sub> (R<sup>2</sup>= 0.75) as shown in  
10 | Fig. 13. Years with a long low ~~discharge~~ water-period like 2011, 2006 or 2007 have a more  
11 | persistent TMZ than years like 2013 or 2010 characterized by shorter periods of low river  
12 | flow.

13 | There is also a good correlation between Turbidity<sub>max</sub> and Volume<sub>HDW</sub> (R<sup>2</sup>= 0.78, Fig. 14.A).  
14 | Years with numerous and large floods (like 2008, 2009 and 2013) present a less turbid TMZ.  
15 | This can be the result of the total ~~downstream flushing~~ expulsion of mobile sediment (as seen  
16 | in Section 5.2) and of the further flushing of the previous TMZ (Castaing & Allen, 1981). The  
17 | Volume<sub>LDW</sub> is not correlated to Turbidity<sub>max</sub>. However, the sum of Volume<sub>LDW</sub> and  
18 | Volume<sub>HDW</sub> improves the correlation (R<sup>2</sup>= 0.90). This is because the water volume during  
19 | very wet summers is enough to expel partly the TMZ.

20 | In summary, the duration of the low ~~water~~discharge period mainly determines the TMZ  
21 | duration, and the freshwater volume during high ~~discharge~~ water-periods the TMZ  
22 | concentration. High river flows are efficient in flushing the TMZ in the central estuary, even  
23 | to the coastal waters, and expel higher quantity of mobile mud, as seen in Section 5.2. In  
24 | order to discuss the potential evolution of the TMZ in the last decades, we calculated the  
25 | Duration<sub>LDW</sub> and the Volume<sub>HDW</sub> at Bordeaux since 1960 to 2013 (Fig. 15). There is a trend in  
26 | decreasing Volume<sub>HDW</sub> and increasing Duration<sub>LDW</sub>, especially since the 890', which has  
27 | changed the TMZ characteristics. The decrease in river discharge is attributed to climate  
28 | change and human activities (Mazzege et al., 2014). For example, in the years 1963 and 1976  
29 | the low ~~discharge~~ water-period lasted respectively only 20 and 9 day, and Volume<sub>HDW</sub> reached  
30 | 2.5 10<sup>34</sup> Hm<sup>3</sup> in 1969 and 1977 and 30 10<sup>34</sup> Hm<sup>3</sup> 1965 and 1976. Considering the relationship  
31 | between TMZ and hydrology (Fig. 13 and 14), we assume that the TMZ is at present more  
32 | persistent and turbid than 40-50 years ago. Furthermore, an accumulation effect can favour



1 this intensification. As the TMZ is concentrated in SPM and persistent, the required water  
2 volume to expel it increases, promoting the next TMZ to be more pronounced.

3 According to recent streamflow simulations from 1976 to 2100 based on 22 European river  
4 basins, including the Garonne watershed, average discharges are projected to decrease in  
5 southern Europe, and extreme events to increase (Alfieri et al., 2015). In this context, the  
6 finding of straightforward river discharge-based indicators of TMZ behaviour should be of  
7 great interest for future river basin management plans in the fluvial Garonne.

8 The effect of river discharge is assumed to be a major factor in the longitudinal shift of the  
9 TMZ. However morphological changes (natural or anthropogenic) may also contribute to the  
10 TMZ intensification (Winterwerp and Wang, 2013; De-Jonge et al., 2014), by amplifying  
11 tidal asymmetry and hence enhancing trapping of fine sediments, as suggested by Sottolichio  
12 et al. (2011). The existence and importance of these changes is not documented yet and will  
13 be the subject of future research. The combined effect of changes in topography and in river  
14 flow on the TMZ evolution needs be analysed by numerical modelling.

## 16 **6 Conclusions**

17 The high-frequency and long-term turbidity monitoring provides detailed informations on  
18 suspended sediment dynamics in the fluvial Gironde estuary over a wide range of time scales  
19 and hydrological conditions. Tide, river flow and sediment stock (mobile mud patches) induce  
20 large variability on turbidity levels. Suspended sediment dynamics related to tidal cycles  
21 (semidiurnal and fortnightly) follows the same cyclic processes in the tidal section, as  
22 previously described in the lower estuary (Allen et al., 1977). The TMZ occurrence in the  
23 tidal rivers is very sensitive to changes in hydrological conditions. River discharge is a key  
24 variable for to explain the upstream migration, downstream flushing installation, expulsion  
25 and concentration of the TMZ and its associated mobile mud. River discharge thresholds  
26 promoting the installation and expulsion of the TMZ at Bordeaux have been delimited, 250  
27 and at least  $350 \text{ m}^3 \text{ s}^{-1}$  respectively, showing the need to a higher “water effort” to expel the  
28 TMZ. Two hydrological indicators of the TMZ intensity have been defined: the duration of  
29 low discharge water periods as indicator of the persistence of the TMZ, and water volume  
30 passing before and during the presence of the TMZ as indicator of the TMZ turbidity level.  
31 Higher water volume contributes to move more efficiently the TMZ and to expel higher  
32 quantity of remained mobile mud, resulting in less concentrated TMZ. The existence of

1 mobile mud during and after the TMZ presence is confirmed through turbidity-discharge  
2 hysteresis patterns over different scales, which reveal the local or remote location of the  
3 sediment source. More particularly, these hysteresis patterns over river floods can serve as an  
4 indicator of the rhythm of downstream flushingexpulsion of mobile mud.

5 The extrapolation of hydrological conditions suggests an intensification of the TMZ  
6 occurrence in the fluvial Gironde during the last decades and could be used to evaluate future  
7 scenarios. This can be very useful to water management strategies in order to address the  
8 global change impacts as Garonne 2050 ([www.garonne2050.fr](http://www.garonne2050.fr)). The estimate of discharge  
9 thresholds of TMZ installation and expulsion is also of great interest to local public  
10 authorities. By example, a partner of the MAGEST network, the SMEAG, is in charge to  
11 maintain a minimum discharge level of the Garonne to ensure a water quality favourable to  
12 ecosystems (<http://www.smeag.fr/plan-de-gestion-detiage-garonne-ariege.html>). Their criteria  
13 to release water stocks from upstream dams are the levels of dissolved oxygen in Bordeaux  
14 waters. It appears from this work that the discharge threshold, below  $100\text{-}110\text{ m}^2\text{ s}^{-1}$ , actually  
15 retained by the SMEAG is far too low to prevent the installation of the TMZ, and the  
16 subsequent problems (dissolved oxygen consumption, pollutant accumulation ...).

17 Finally this work will be useful to improve the calibration of numerical models coupling  
18 hydrodynamics and suspended sediment transport. Numerical simulations will allow evaluate  
19 the turbidity in the upper estuary for different hydrological and climate scenarios (natural  
20 and anthropogenic), including the effect of morphological changes.

21

## 22 **Acknowledgements**

23 I. Jalón-Rojas thanks the Agence de l'Eau Adour-Garonne (AEAG) and the Aquitaine Region  
24 for the financial support of her PhD grant. The MAGEST network is financially supported by  
25 the following organisms: AEAG (Agence de l'Eau Adour-Garonne); SMIDDEST (Syndicat  
26 MIxte pour le Développement Durable de l'ESTuaire de la Gironde); SMEAG (Syndicat  
27 Mixte d'Etudes et d'Aménagement de la Garonne); EPIDOR (Etablissement Public  
28 Interdépartemental de la Dordogne); EDF; GPMB (Grand Port Maritime de Bordeaux); CUB  
29 (Communauté Urbaine de Bordeaux); Conseil Régional Aquitaine; CG-33 (Conseil Général  
30 de Gironde); Ifremer; CNRS; Université de Bordeaux. The authors thank also the support of  
31 the OASU (Observatoire Aquitain des Sciences de l'Univers) through the SOLAQUI (Service

- 1 d'Observation du Littoral AQUItain) program. MAGEST is a contribution to the CNRS
- 2 observation program DYNALIT.
- 3

## 1 **References**

- 2 [Alfieri, L., Burek, P., Feyen, L., & Forzieri, G.: Global warming increases the frequency of](#)  
3 [river floods in Europe, \*Hydrology and Earth System Sciences\*, 19\(5\), 2247–2260.](#)  
4 [doi:10.5194/hess-19-2247-2015, 2015.](#)
- 5 Allen, G. P.: Déplacements saisonniers de la lentille de "crème de vase" dans l'estuaire de la  
6 Gironde, in: C. R. Acad. Sc. Paris, 273, pp. 2429–2431, 1971.
- 7 Allen, G. P. and Castaing, P.: Suspended sediment transport from the Gironde estuary  
8 (France) onto the adjacent continental shelf, *Marine Geology*, 14, 47–53, 1973.
- 9 Allen, G. P., Sauzay, G., and Castaing, P.: Transport and deposition of suspended sediment in  
10 the Gironde Estuary, France, in: *Estuarine Processes*, edited by Wiley, M., pp. 63–81,  
11 Academic Press, New York, 2 edn., 1977.
- 12 Allen, G. P., Salomon, J. C., Bassoullet, P., Du Penhoat, Y., and De Grandpré, C.: Effects of  
13 tides on mixing and suspended sediment transport in macrotidal estuaries, *Sedimentary*  
14 *Geology*, 26, 69–90, 1980.
- 15 Bonneton, P., Bonneton, N., Parisot, J.-p., and Castelle, B.: Tidal bore dynamics in funnel-  
16 shaped estuaries, *Journal of Geophysical Research*, doi:10.1002/2014JC010267, 2015.
- 17 Castaing, P. and Allen, G. P.: Mechanisms controlling seaward escape of suspended sediment  
18 from the Gironde: A macrotidal estuary in France, *Marine Geology*, 40, 101–118, 1981.
- 19 Castaing, P., Etcheber, H., Sottolichio, A., and Cappe, R.: Evaluation de l'évolution  
20 hydrodologique et sédimentaire du système Garonne-Dordogne-Gironde, Tech. rep., Rapport  
21 Agence de l'eau Adour-Garonne - Université de Bordeaux, 2006.
- 22 [Chanson, H., Reungoat, D., Simon, B., and Lubin, P.: High-frequency turbulence and](#)  
23 [suspended sediment concentration measurements in the Garonne River tidal bore, \*Estuarine,\*](#)  
24 [\*Coastal and Shelf Science\*, 95\(2-3\), 298–306. doi:10.1016/j.ecss.2011.09.012, 2011.](#)
- 25 Contreras, E. and Polo, M. J.: Measurement frequency and sampling spatial domains required  
26 to characterize turbidity and salinity events in the Guadalquivir estuary (Spain), *Natural*  
27 *Hazards Earth System Sciences*, 12, 2581–2589, 2012.
- 28 De-Jonge, V. N., Schuttelaars, H. M., Van-Beusekom, J. E., Talke, S. A., and De-Swart, H.  
29 E.: The influence of channel deepening on estuarine turbidity levels and dynamics, as

1 exemplified by the Ems estuary, *Estuarine, Coastal and Shelf Science*, 139, 46–59,  
2 doi:10.1016/j.ecss.2013.12.030, 2014.

3 Doxaran, D., Froidefond, J. M., Castaing, P., and Babin, M.: Dynamics of the turbidity  
4 maximum zone in a macrotidal estuary (the Gironde, France): Observations from field and  
5 MODIS satellite data, *Estuarine, Coastal and Shelf Science*, 81, 321–332,  
6 doi:10.1016/j.ecss.2008.11.013, 2009.

7 Dyer, K. R.: Fine sediment particle transport in estuaries, in: *Physical Process in Estuaries*,  
8 edited by Dronkers, J. and van Leussen, W., pp. 427–445, Springer-Verlag, 1988.

9 Etcheber, H., Schmidt, S., Sottolichio, A., Maneux, E., Chabaux, G., Escalier, J. M.,  
10 Wennekes, H., Derriennic, H., Schmeltz, M., Quémener, L., Repecaud, M., Woerther, P., and  
11 Castaing, P.: Monitoring water quality in estuarine environments: lessons from the MAGEST  
12 monitoring programme in the Gironde fluvial-estuarine system, *Hydrology and Earth System  
13 Sciences*, 15, 831–840, doi:10.5194/hess-15-831-2011, 2011.

14 Fettweis, M., Sas, M., and Monbaliu, J.: Seasonal, Neap-spring and Tidal Variation of  
15 Cohesive Sediment Concentration in the Scheldt Estuary, Belgium, *Estuarine, Coastal and  
16 Shelf Science*, 47, 21–36, doi:10.1006/ecss.1998.0338, 1998.

17 Fettweis, M., Monbaliu, J., Baeye, M., Nechad, B., and Van den Eynde, D.: Weather and  
18 climate induced spatial variability of surface suspended particulate matter concentration in the  
19 North Sea and the English Channel, *Methods in Oceanography*, 3-4, 25–39,  
20 doi:10.1016/j.mio.2012.11.001, 2012.

21 Fontugne, M. R., and Jouanneau, J.-M.: Modulation of the particulate organic carbon flux to  
22 the ocean by a macrotidal estuary: Evidence from measurements of carbon isotopes in organic  
23 matter from the Gironde system. *Estuarine, Coastal and Shelf Science*, 24, 377–387, 1987.

24 Garel, E., Nunes, S., Neto, J. M., Fernandes, R., Neves, R., Marques, J. C., and Ferreira, O.:  
25 The autonomous Simpatico system for real-time continuous water-quality and current velocity  
26 monitoring: examples of application in three Portuguese estuaries, *Geo-Marine Letters*, 29,  
27 331–341, doi:10.1007/s00367-009-0147-5, 2009.

28 Grabemann, I. and Krause, G.: Response of the turbidity maximum in the Weser Estuary to  
29 pulses in freshwater runoff and to storms, in: *Physics of estuaries and coastal seas*, pp. 83–92,  
30 1998.

- 1 Grabemann, I. and Krause, G.: On Different Time Scales of Suspended Matter Dynamics in  
2 the Weser Estuary, *Estuaries*, 24, 688–698, 2001.
- 3 Grabemann, I., Uncles, R. J., Krause, G., and Stephens, J. A.: Behaviour of Turbidity Maxima  
4 in the Tamar (U.K.) and Weser (F.R.G.) Estuaries, *Estuarine, Coastal and Shelf Science*, 45,  
5 235–246, doi:10.1006/ecss.1996.0178, 1997.
- 6 Guézennec, L., Lafite, R., Dupont, J. P., Meyer, R., and Boust, D.: Hydrodynamics of  
7 Suspended Particulate Matter in the Tidal Freshwater Zone of a Macrotidal Estuary (The  
8 Seine Estuary, France), *Estuaries*, 22, 717–727, doi:10.2307/1353058, 1999.
- 9 Hendrickx, F. and Sauquet, E.: Impact of warming climate on water management for the  
10 Ariège River basin (France), *Hydrological Sciences Journal*, 58, 976–993,  
11 doi:10.1080/02626667.2013.788790, 2013.
- 12 Jay, D. A. and Musiak, J. D.: Particle trapping in estuarine tidal flows, *Journal of Geophysical*  
13 *Research*, 99(C10), 1994.
- 14 Klein, M.: Anti clockwise hysteresis in suspended sediment concentration during individual  
15 storms: Holbeck Catchment; Yorkshire, England, *Catena*, 11, 251–257, 1984.
- 16 Lanoux, A., Etcheber, H., Schmidt, S., Sottolichio, A., Chabaud, G., Richard, M., and Abril,  
17 G.: Factors contributing to hypoxia in a highly turbid, macrotidal estuary (the Gironde,  
18 France), *Environmental Science: Processes & Impacts*, 15, 585–595,  
19 doi:10.1039/c2em30874f, 2013.
- 20 López-Tarazón, J. A., Batalla, R. J., Vericat, D., and Francke, T.: Suspended sediment  
21 transport in a highly erodible catchment: The River Isábena (Southern Pyrenees),  
22 *Geomorphology*, 109, 210–221, doi:10.1016/j.geomorph.2009.03.003, 2009.
- 23 Mazzega, P., Therond, O., Debril, T., March, H., Sibertin-Blanc, C., Lardy, R., and Sant’ana,  
24 D.: Critical multi-level governance issues of integrated modelling: An example of low-water  
25 management in the Adour-Garonne basin (France), *Journal of Hydrology*, 519, 2515–2526,  
26 doi:10.1016/j.jhydrol.2014.09.043, 2014.
- 27 [Mitchell, S., Akesson, L., and Uncles, R.: Observations of turbidity in the Thames Estuary,](#)  
28 [United Kingdom, \*Water and Environment Journal\*, 26\(4\), 511–520. doi:10.1111/j.1747-](#)  
29 [6593.2012.00311.x, 2012.](#)

1 Mitchell, S.: Turbidity maxima in four macrotidal estuaries, *Ocean & Coastal Management*,  
2 79, 62–69, doi:10.1016/j.ocecoaman.2012.05.030, 2013.

3 Mitchell, S. B. and Uncles, R. J.: Estuarine sediments in macrotidal estuaries: Future research  
4 requirements and management challenges, *Ocean and Coastal Management*, 79, 97–100,  
5 doi:10.1016/j.ocecoaman.2012.05.007, 2013.

6 Pontee, N., Whitehead, P., and Hayes, C.: The effect of freshwater flow on siltation in the  
7 Humber Estuary, north east UK, *Estuarine, Coastal and Shelf Science*, 60, 241–249,  
8 doi:10.1016/j.ecss.2004.01.002, 2004.

9 Romaña, A.: Estuaire de la Gironde. Campagnes "Libellule". Distribution longitudinale des  
10 paramètres mesurés et calculés, Tech. rep., Rapport Ifremer, Dépt. Env. Litt. et Gest. du  
11 Milieu Marin, 1983.

12 Saari, H. K., Schmidt, S., Castaing, P., Blanc, G., Sautour, B., Masson, O., and Cochran, J.  
13 K.: The particulate  $^{7}\text{Be}/^{210}\text{Pb}_{\text{xs}}$  and  $^{234}\text{Th}/^{210}\text{Pb}_{\text{xs}}$  activity ratios as tracers for tidal-to  
14 seasonal particle dynamics in the Gironde estuary (France): Implications for the budget of  
15 particle-associated contaminants, *Science of the Total Environment*, 408, 4784–4794,  
16 doi:10.1016/j.scitotenv.2010.07.017, 2010.

17 Schmidt, S., Ouamar, L., Cosson, B., Lebleu, P., and Derriennic, H.: Monitoring turbidity as a  
18 surrogate of suspended particulate load in the Gironde Estuary: The impact of particle size on  
19 concentration estimates, in: ISOBAY, 2014.

20 Schrottke, K., Becker, M., Bartholomä, A., Flemming, B. W., and Hebbeln, D.: Fluid mud  
21 dynamics in the Weser estuary turbidity zone tracked by high-resolution side-scan sonar and  
22 parametric sub-bottom profiler, *Geo-Marine Letters*, 26, 185–198, doi:10.1007/s00367-006-  
23 0027-1, 2006.

24 Schuttelaars, H. M., de Jonge, V. N., and Chernetsky, A.: Improving the predictive power  
25 when modelling physical effects of human interventions in estuarine systems, *Ocean &*  
26 *Coastal Management*, 79, 70–82, doi:10.1016/j.ocecoaman.2012.05.009, 2013.

27 Sottolichio, A. and Castaing, P.: A synthesis on seasonal dynamics of highly-concentrated  
28 structures in the Gironde estuary, *Comptes Rendus de l'Academie de Sciences - Serie IIA:*  
29 *Sciences de la Terre et des Planetes*, 329, 795–800, doi:10.1016/S1251-8050(00)88634-6,  
30 1999.

1 Sottolichio, A., Castaing, P., Etcheber, H., Maneux, E., Schmeltz, M., and Schmidt, S.:  
2 Observations of suspended sediment dynamics in a highly turbid macrotidal estuary, derived  
3 from continuous monitoring, *Journal of Coastal Research*, pp. 1579–1583, 2011.

4 StataCorp: *Stata Statistical Software: Release 12*, 2011.

5 Talke, S. A., de Swart, H. E., and Schuttelaars, H. M.: Feedback between residual circulations  
6 and sediment distribution in highly turbid estuaries: An analytical model, *Continental Shelf*  
7 *Research*, 29, 119–135, doi:10.1016/j.csr.2007.09.002, 2009.

8 Uncles, R. J., Easton, A. E., Griffiths, M. L., Harris, C., Howland, R. J. M., King, R. S.,  
9 Morris, A. W., and Plummer, D. H.: Seasonality of the Turbidity Maximum in the Humber  
10 Ouse Estuary, UK, *Marine Pollution Bulletin*, 37, 206–215, 1998.

11 Uncles, R. J., Stephens, J. A., and Smith, R. E.: The dependence of estuarine turbidity on tidal  
12 intrusion length, tidal range and residence time, *Continental Shelf Research*, 22, 1835–1856,  
13 doi:10.1016/S0278-4343(02)00041-9, 2002.

14 Uncles, R. J., Stephens, J. A., and Law, D. J.: Turbidity maximum in the macrotidal, highly  
15 turbid Humber Estuary, UK: Flocs, fluid mud, stationary suspensions and tidal bores,  
16 *Estuarine, Coastal and Shelf Science*, 67, 30–52, doi:10.1016/j.ecss.2005.10.013, 2006.

17 Uncles, R. J., Stephens, J. A., and Harris, C.: Towards predicting the influence of freshwater  
18 abstractions on the hydrodynamics and sediment transport of a small, strongly tidal estuary:  
19 The Devonshire Avon, *Ocean and Coastal Management*, 79, 83–96,  
20 doi:10.1016/j.ocecoaman.2012.05.006, 2013.

21 Williams, G. P.: Sediment concentration versus water discharge during single hydrologic  
22 events in rivers, *Journal of Hydrology*, 111, 89–106, doi:10.1016/0022-1694(89)90254-0,  
23 1989.

24 Winterwerp, J. C. and Wang, Z. B.: Man-induced regime shifts in small estuaries—I: theory,  
25 *Ocean Dynamics*, 63, 1279–1292, doi:10.1007/s10236-013-0662-9, 2013.

26 Yang, Y., Li, Y., Sun, Z., and Fan, Y.: Suspended sediment load in the turbidity maximum  
27 zone at the Yangtze River Estuary: The trends and causes, *Journal of Geographical Sciences*,  
28 24, 129–142, doi:10.1007/s11442-014-1077-3, 2013.

29




1 Table 1. Discharge and turbidity characteristics of flood events for the period 2005 – mid  
 2 2014 in the tidal Garonne River (Bordeaux and Portets stations). Flood event were numbered  
 3 by f plus a number according to figure 3. Hysteresis loops were classified as: [C] clockwise;  
 4 [CC] counterclockwise; [M] mixed; [No] no trend. Mixed loops with a clear clockwise  
 5 [M(C)] or counterclockwise [M(CC)] predominance were specified. Flood without turbidity  
 6 record were included to facilitate the interpretation of the hysteresis succession.

7


	Date	Q <sub>max</sub> (m <sup>3</sup> s <sup>-1</sup> )	Bordeaux		Portets	
			T <sub>max</sub> (NTU)	Hysteresis	T <sub>max</sub> (NTU)	Hysteresis
2006						
	10.12.2005	865	-	-	-	-
	03.01.2006	938	-	-	-	-
f1	31.01.2006	1820	989	M(C)	908	CC
f2	12.03.2006	4160	1446	CC	1326	CC
2007						
f3	13.02.2007	2140	1460	C	975	M
f4	27.02.2007	1600	414	M(C)	292	M
f5	18.04.2007	1210	-	-	400	CC
f6	03.05.2007	953	349	C	-	-
f7	28.05.2007	1730	1794	CC	-	-
2008						
	11.12.2007	1270	-	-	-	-
f8	08.01.2008	1120	1008	C	313	M
f9	19.01.2008	2180	835	No	795	CC
	22.04.2008	3130	-	-	-	-
f10	28.05.2008	2640	495	CC	-	-
2009						

f11	03.11.2008	1450	2200	C	993	M(CC)
f12	06.12.2008	1830	476	CC	-	-
f13	25.01.2009	4750	1578	CC	-	-
f14	13.04.2009	1950	-	-	203	CC
	30.04.2009	2870	-	-	-	-
2010						
f15	16.01.2010	1880	747	M(CC)	-	-
f16	07.02.2010	1410	358	No	-	-
f17	02.04.2010	1070	471	M(C)	152	No
f18	06.05.2010	1770	971	M(CC)	474	CC
2011						
f19	24.12.2010	1480	425	M	-	-
f20	24.02.2011	1090	1598	M	(C)	164
f21	18.03.2011	2150	-	-	723	CC
2012						
	08.11.2012	1890	-	-	-	-
	07.01.2012	1390	-	-	-	-
f22	01.05.2012	1760	335	M	-	-
f23	23.05.2012	3110	963	CC	-	-
2013						
f24	07.12.2012	834	914	C	-	-
f25	21.01.2013	3460	1075	M	-	-
f26	09.03.2013	1150	175	CC	-	-
	31.03.2013	2510	-	-	-	-
f27	01.06.2013	4020	768	CC	-	-
f28	20.06.2013	1980	1304	CC	-	-



1  
2  
3  
4  
5  
6


Figure 1. The Gironde fluvial-estuarine system: a) Location map (SW France), the grey area shows the watershed of Garonne and Dordogne; b) the estuary with its main tributaries. Red circles locate the MAGEST stations; blue squares indicate the hydrometric stations.



1


2

3 Figure 2. Comparison of tidally-averaged turbidity and daily-averaged turbidity for Bordeaux station showing  
4 the correlation coefficient ( $R^2$ ).



1

2 Figure 3. (A) Daily mean flow of the Garonne River and the Dordogne River showing the river flood events of Table 1; (B) tidal range  
 3 recorded at Bordeaux tide gauge; and tidally-averaged turbidity at (C) Pauillac, (D) Bordeaux, and (E) Portets and (F) Libourne stations. Red  
 4 dotted lines represent the low-pass filtered data performed with running averages in order to highlight the turbidity trends. The labels f plus a  
 5 number refer to the flood events according to Table 1. (a), (b) and (c) indicate the neap-spring-neap cycles represented in Figure 10.




1

2

3

4 Figure 4. Examples of 48H raw data of (a) river flow, and (b) turbidity and water level (dotted  
 5 lines) at Bordeaux for two contrasted hydrological conditions. The mean time step of river  
 6 flow is one hour, while turbidity and water level were recorded everyeach-10 minutes. (c)  
 7 Relationships between turbidity and water level records of the middle panels.

8




1

2

3 Figure 5. Mean (red cross), median (red bars), percentiles 25-75 (blue bars) and minimum–  
 4 maximum (black bars) values of tidally-averaged turbidity depending on the season (months  
 5 of February and August) and the tidal range (TR) in each MAGEST station. The minimal,  
 6 mean and maximal values of river flow in February (2005-2014) are 176, 566 and 2994 m<sup>3</sup>s<sup>-1</sup>  
 7 respectively. These values in August are 56, 103 and 317 m<sup>3</sup>s<sup>-1</sup> respectively. High and low  
 8 tidal ranges correspond to values above the percentile 75 and below the percentile 25,  
 9 respectively, of the entire TR dataset of each station.

10

1




2

3

4 Figure 6. Duration of the TMZ presence per year at the three tidal rivers stations. Striped bars  
5 designate the duration of the TMZ when it appears in winter: 17, 18, 9 and 39 days  
6 respectively in the years 2005, 2008, 2011 and 2012 at Bordeaux; 6 days in the year 2012 at  
7 Libourne.


8






1  
2  
3  
4  
5  
6  
7  
8

Figure 7. Relationship between Garonne discharge and turbidity at Bordeaux, and corresponding hysteresis patterns for the successive floods occurring since the ~~departure~~ downstream flushing, in December 2012, and the following upstream migration ~~return of the~~ TMZ, in August 2013, of the TMZ. The labels f plus a number refer to the flood events according to Figure 3 and Table 1.




1  
2  
3  
4  
5  
6

Figure 8. Schematic representation of suspended sediment dynamics in tidal rivers associated to the different types of hysteresis (clockwise, ~~anti~~clockwise, mixed, and anticlockwise) during river floods.




1  
2  
3  
4  
5  
6  
7  
8

Figure 9. Tidally-averaged turbidity (A) and daily-averaged turbidity (B) as a function of 3-days averaged river flow for the MAGEST stations of Pauillac, Bordeaux and Portets (log-log representation). (A): values were classified in function of tidal range (TR). (B): values corresponding to the periods of installation (blue diamonds) and expulsion (green square) of the TMZ were differentiated.




1  
2  
3  
4  
5  
6

Figure 10. Turbidity as a function of tidal rage (2-days running averages) for three neap-spring-neap cycles (see the cycles in Fig.3.D) during a period of (a) installation, (b) presence and (c) expulsion of the TMZ at Bordeaux.




1  
2  
3  
4  
5

Figure 11. Examples of clockwise **D**ischarge/**T**urbidity hysteresis curves during the transition periods of installation and expulsion of the TMZ (see these periods in Fig. 3).




1  
2  
3  
4  
5  
6

Figure 12. Mean (red cross), median (red bars), percentiles 25-75 (blue bars) and percentiles 9-91 (black bars) values of tidally-averaged turbidity per  $30 m^3 s^{-1}$  intervals of river flow during the installation and expulsion of the TMZ at Bordeaux.



1




2

3

4 Figure 13. Duration of the TMZ presence as a function of the number of days per year where  
 5 the river flow was below  $250 \text{ m}^3 \text{ s}^{-1}$  at Bordeaux station and  $160 \text{ m}^3 \text{ s}^{-1}$  at Portets station.

6



1

2





3

4 Figure 14. Turbidity maxima of the TMZ as a function of the water volume passed: (A) during  
 5 the previous wet period; and (B) during the previous wet period + the presence of the TMZ.

6 PortetsOnly Bordeaux is ~~not~~ considered as it was not possible to estimate Turbidity<sub>max</sub> at  
 7 Portets due to the number of missing data.

8





1

2

3

4 Figure 15. Evolution of the duration of low discharge water-period (Duration<sub>LD</sub>) and the water  
 5 volume during high discharge water-periods (Volume<sub>HD</sub>) between 1960 and 2013 (calculated  
 6 from discharge data available on <http://www.hydro.eaufrance.fr/>); red lines represent the 5-  
 7 days running averages in order to highlight the trends.