

1 **Monitoring strategies of stream phosphorus under** 2 **contrasting climate-driven flow regimes**

3 **Goyenola, G.1; Meerhoff, M.1, 2; Teixeira-de Mello, F.1; González-Bergonzoni, I.1,**
4 **2, 3; Graeber, D.2; Fosalba, C.1: Vidal, N.1, 2, 3; Mazzeo, N.1; Ovesen, N. B.2;**
5 **Jeppesen, E.2, 3, & Kronvang, B.2**

6
7 [1]{Departamento de Ecología Teórica y Aplicada, CURE-Facultad de Ciencias, Universidad
8 de la República. Maldonado, Uruguay}

9 [2]{Department of Bioscience and Arctic Research Centre, Aarhus University, Silkeborg,
10 Denmark}

11 [3]{Sino-Danish Centre for Education and Research, Beijing, China}

12 Correspondence to: G. Goyenola (goyenola@gmail.com)

13 14 **Abstract**

15 Climate and hydrology are relevant control factors determining the timing and amount of
16 nutrient losses from land to downstream aquatic systems, in particular of P from agricultural
17 lands. The main objective of the study was to evaluate the differences in P export patterns and
18 the performance of alternative monitoring strategies in streams under contrasting climate-
19 driven flow regimes. We compared a set of paired streams draining lowland micro-catchments
20 under temperate climate and stable discharge conditions (Denmark) and under sub-tropical
21 climate and flashy conditions (Uruguay). We applied two alternative nutrient sampling
22 programs (high frequency composite sampling and low frequency instantaneous-grab
23 sampling) and estimated the contribution derived from point and diffuse sources fitting a source
24 apportionment model. We expected to detect a pattern of higher total and particulate phosphorus
25 export from diffuse sources in streams in Uruguay streams, mostly as a consequence of higher
26 variability in flow regime (higher flashiness). Contrarily, we found a higher contribution of

27 dissolved P in flashy streams. We did not find a notably poorer performance of the low
28 frequency sampling program to estimate P exports in flashy streams compared to the less
29 variable streams. We also found signs of interaction between climate/hydrology and land use
30 intensity, in particular in the presence of point sources of P, leading to a bias towards
31 underestimation of P in hydrologically stable streams and overestimation of P in flashy streams.
32 Based on our findings, we suggest that the evaluation and use of more accurate monitoring
33 methods, such as automatized flow-proportional water samplers and automatized bankside
34 analysers, should be prioritized whenever is logistically possible. However, it seems
35 particularly relevant in currently flashy systems and also in systems where climate change
36 predictions suggest an increase in stream flashiness.

37

38 **1 Introduction**

39 The global demand for phosphorus for fertilizers is projected to increase in the coming decades,
40 although the existing global reserves will likely be exhausted within this century (Cordell et al.,
41 2009). Nowadays, the global flow of phosphorus runs from mines through farms to aquatic
42 systems, contributing to eutrophication and triggering aquatic ecosystem degradation
43 (Carpenter et al., 1998; Elser and Bennett, 2011). Thus, humanity faces the challenge of
44 sustaining food production while reducing the associated environmental costs

45 The biogeochemical processes inside a catchment, which determine the loss of phosphorus from
46 the land to aquatic systems, are mainly dependent of climatic and hydrological regimes
47 (Bormann and Likens, 1967). Streams located in areas with short duration-high magnitude
48 rainfall events may respond with frequent and rapid changes in discharge (to be “flashy”
49 streams sensu Baker et al., 2004) and may potentially have a higher risk of diffuse transfer of
50 nutrients from land to water (Cassidy and Jordan, 2011; Haygarth et al., 1999). The importance
51 of understanding hydrology-driven variations in nutrient discharge increases in the current
52 climate change scenario where strong hydrological changes are expected in many different parts
53 of the world.

54

55 Irrespective of the hydrological nature of the catchments, nutrient monitoring programs are
56 frequently based on low frequency sampling at discrete intervals. This kind of sampling strategy
57 complemented by interpolation methods are prone to very high uncertainties due to typical
58 under-representation of high discharge, short-duration events (Defew et al., 2013; Jones et al.,
59 2012; Jordan and Cassidy, 2011; Stelzer and Likens, 2006). A priori, the risk of missing the key
60 moments when phosphorus is delivered to the streams is higher in flashy streams than in
61 hydrologically stable ones. An approach to advance the understanding of how different
62 monitoring schemes capture catchment phosphorus processes in contrasting climate-driven
63 flow regimes is to compare monitoring performances in catchments under different climatic
64 conditions and under different conditions of nutrient inputs.

65 The main objectives of our study were to evaluate the differences in stream P export patterns
66 and the performance of alternative monitoring strategies in contrasting climate-driven flow
67 regimes. We expected to detect higher total and particulate P exports in streams located in sites
68 with higher frequency of extreme rainfall events and higher stream hydrological variability
69 (flashiness). We also expected poorer performance of low frequency sampling programs under
70 such conditions.

71

72 **2 Material and Methods**

73 **2.1. Design rationale: Selection of case studies**

74 We conducted a comparative study of concentrations and export of different P forms in two
75 paired streams under two distinct climatic-hydrological conditions: temperate climate and
76 stable discharge conditions (Denmark) and subtropical climate and flashy conditions
77 (Uruguay). In both countries, the topography of the selected areas can be described as gently
78 rolling plains (mean slope < 5%). Two main causes explaining the differences in flashiness
79 between Denmark and Uruguay are precipitation patterns (annual average precipitation, 745
80 mm since 1990 in Denmark and 1300 mm in Uruguay, according to the Danish Meteorological
81 Institute (DMI, 2015) and the National Institute for Agriculture Research, INIA, Uruguay
82 (Castaño et al., 2011), respectively, and depth of soils and derived water storage capacity. The
83 long-term continuous monitoring data in Danish catchments and the existence of published

84 works evaluating alternative sampling strategies (e.g. Kronvang and Bruhn, 1996) decided us
85 to use the temperate streams as the reference systems in our comparison.

86 In each area, two lowland non-experimental micro-catchments (< 20 km²) were selected as
87 typical productive systems to represent extremes of land use intensity (hereafter LUI) in each
88 area.

89 As higher LUI catchments, we selected catchments where intensive farming comprising more
90 than 90% of the total land area, with arable cropping systems, intensive use of fertilizers, and
91 high density of livestock (Table 1). In the Uruguayan high-LUI catchment, the farming
92 production system was based on forage crops, no-till practices associated with intensive
93 application of mineral fertilizers over the soil surface (Derpsch et al., 2010), dairy cattle feeding
94 all year round in open fields, and no effluent treatment facilities on milking plants. In Denmark,
95 the high-LUI catchment included a rotation cropping system with winter cereals and confined
96 pig farms with slurry storage facilities. In Denmark, most loamy agricultural fields are drained
97 with subsurface tile drainage systems (Grant et al., 1996), and the manure originating from
98 farming activities is reutilized with a demand on a 75% reuse of N in slurry.

99 The lower-LUI catchments were chosen so as to represent local more preserved conditions. The
100 Uruguayan low-LUI catchment was dominated by the natural grasslands of the Pampa Biome
101 (Allaby, 2006) and sustained low density cattle production (70% of total area and below 1 head
102 by hectare; Table 1). In contrast, a mixture of deciduous and coniferous forests dominated the
103 Danish low-LUI catchment (Table 1).

104 The subtropical high-LUI catchment had 170 inhabitants and the low-LUI catchment only 20
105 inhabitants (National Institute of Statistics, 2015). In the former, the sewage from only 10
106 households is treated in a facultative pond. All other households in both subtropical catchments
107 had leaking septic tanks. The point sources in the temperate catchments were mainly scattered
108 dwellings without connection to sewage treatment plants. The temperate high-LUI stream
109 received stormwater outlets from a small village whose sewage water is pumped to a treatment
110 plant with tertiary treatment outside the catchment.

111 Maps including land use for each catchment were included as supplementary material.
112 Henceforth, we will refer to the temperate- Danish streams as “STABLE” and the subtropical-

113 Uruguayan streams as “FLASHY”, while LUI categorization of the intensive and extensive
114 production catchments will be referred to as “high-LUI” and “low-LUI”, respectively.

115

116 **2.2. Phosphorus monitoring**

117 Similar gauging stations were established in all four micro-catchments. We applied two
118 alternative nutrient sampling programs: low frequency instantaneous-grab sampling and high
119 frequency composite sampling. Instantaneous-grab sampling of water was conducted
120 fortnightly, and P exports were estimated by two daily step interpolation methods. High
121 frequency composite sampling was undertaken using Glacier refrigerated automatic samplers
122 (ISCO-Teledyne). The samplers collected an equal water volume every four hours, and the
123 composite samples were also collected fortnightly. The final phosphorus concentration in the
124 only sampler carboy thus represented a time-proportional average for the fortnightly sampling
125 period. As the high frequency composite samples integrated more information (i.e. shorter time
126 steps, with higher probability of capturing extreme events), we expected this method to provide
127 better estimates of the ‘true’ exported P from the catchments. Based on this assumption, we
128 evaluated the deviation relative to the estimation methods based on instantaneous-grab samples.

129 We analyzed the measurements and results from a two-year monitoring period (March 2010 to
130 March 2012 in STABLE; January 2011 to January 2013 in FLASHY).

131

132 **2.3. Meteorological and hydrometric monitoring**

133 In all catchments, CR10X data loggers (Campbell Scientific Ltd.) collected data every 10
134 minutes. In the FLASHY streams, we used CS450 Submersible Pressure Transducers
135 (Campbell Scientific Ltd.) for water stage monitoring as well as HMP45C temperature probes
136 (Campbell Scientific Ltd.) and Rain-O-Matic Professional rainfall automatized gauges
137 (Pronamic). In the STABLE catchments, water level was registered with PDCR 1830 pressure
138 sensors (Druck), while meteorological information was obtained from the Danish
139 Meteorological Institute monitoring network based on a 10 x10 km grid.

140 Periodic instantaneous flow measurements were taken using a C2-OTT Kleinflügel,
141 transferring data to software for the calculation of instantaneous discharge (VB-Vinge 3.0,
142 Mølgaard Hydrometri). Non-linear stable regressions between stage and discharge at each
143 monitoring station (rating curves) were fitted. The rating curves were used to generate a 10
144 minutes discharge data series utilizing the software HYMER (www.orbicon.com). For
145 comparisons, discharge data is reported as area-specific runoff.

146

147 **2.4. Phosphorus analysis**

148 All instantaneous-grab and composite water samples were analyzed for total phosphorus (TP),
149 total dissolved phosphorus (TDP), and particulate phosphorus (PP). In addition, also soluble
150 reactive phosphorus (SRP) and soluble non-reactive phosphorus (NSRP) were estimated from
151 instantaneous-grab samples.

152 Instantaneous-grab samples for TDP and SRP analysis were filtered through 0.45 μm
153 membranes. For TDP analysis, STABLE high frequency composite samples were filtered using
154 0.45 μm pore size membranes, while FLASHY-high frequency composite samples were filtered
155 using Whatman GF/C (pore size 1.2 μm). To detect possible bias derived from the type of filter
156 used, we performed a Kruskal-Wallis test on the proportional contribution of TDP to TP
157 between FLASHY instantaneous-grab and high frequency composite samples and found no
158 significant differences. Consequently, we consider grab and composite TDP samples to be
159 comparable. Particulate phosphorus (PP) was estimated as the difference between TP and TDP.
160 Soluble non-reactive phosphorus (SNRP) was also calculated as the difference between TDP
161 and SRP.

162 All the samples were determined as molybdate reactive P by equivalent spectrophotometric
163 methods, preceded by strong oxidation with a solution of potassium persulfate, boric acid, and
164 sodium hydroxide following Valderrama (1981).

165

166 **2.5. Data processing and analysis**

167 Climatic and runoff patterns were explored in order to investigate the main parameters relevant
168 for P temporal dynamics. As a proxy of catchment water balance, the runoff ratio (i.e. annual
169 percentage of rainfall water exported as runoff) was calculated (Wu et al., 2013). Additionally,
170 to quantify the variation in flow regime we calculated the Richards-Baker Index (hereafter R-
171 B Index; Baker et al., 2004). The R-B Index allows for evaluation of the “flashiness” or the
172 annual ratio of absolute day-to-day fluctuations of streamflow relative to total flow (Baker et
173 al., 2004). Increasing its value with increasing flashiness, the R-B Index varies between 0 and
174 infinity and assumes a value of 1 when the accumulated volume of daily oscillations has the
175 same magnitude as the annually accumulated discharge. The relative contribution of baseflow
176 to total stream flow was estimated from daily hydrographs using the automatic routine proposed
177 by Arnold et al. (1995) as an indicator of the deeper groundwater contribution. Percent
178 contribution of stormflow to total flow was estimated as complementary to the baseflow
179 contribution (Table 2).

180 The statistical relationship between all phosphorus compounds from instantaneous-grab
181 samples was analysed by Spearman rank order correlation. The temporal dynamics of P forms
182 were followed for total P (TP), particulate P (PP), total dissolved P (TDP) and soluble reactive
183 P (SRP), as minimum (min), median (med) and maximum (max) range and interquartile range
184 (IQR). The statistical comparisons of P temporal dynamics between the four streams were
185 conducted using Kruskal-Wallis tests (Zar, 2010), followed by a *post hoc* pairwise multiple
186 comparison procedure when appropriate (Dunn, 1964).

187 Three different methods were used for the calculation of stream P export. The first method was
188 based on multiplying the TP and TDP concentrations obtained from the high frequency
189 fortnight composite samples by the accumulated discharge for the same time period (Kronvang
190 and Bruhn, 1996). Missing data from the relatively short periods when the automatic samplers
191 were not in operation (e.g. frozen in Denmark) were re-generated through linear interpolation
192 of concentrations (Jones et al., 2012).

193

194 Secondly, we calculated exported TP and TDP from the low frequency instantaneous-grab data
195 by two alternative methods of concentration interpolation: linear (Kronvang and Bruhn, 1996)
196 and concentration-discharge relationships (Bowes et al., 2008). Daily real and interpolated

197 concentrations were subsequently multiplied by daily accumulated discharge to obtain daily
198 export estimates.

199 Concentration-discharge relationships (C-Q) were established based on instantaneous-grab
200 samples for all four streams by applying the load apportionment model developed by Bowes et
201 al. (2008). This simple modelling approach does not require GIS information on land use,
202 catchment size, population, or livestock density and may act as a valuable and versatile tool for
203 catchment managers to determine suitable catchment mitigation options (Bowes et al., 2008).
204 Several authors have found similar relationships and used them to characterize P dynamics (e.g.
205 Meyer and Likens, 1979), calculate P sources (e.g. Bowes et al., 2008), or to calculate P
206 transport (e.g. Kronvang and Bruhn, 1996).

207 The Bowes et al. (2008) method assumes that the load of phosphorus from point and diffuse
208 sources can be modelled as a power-law function of the river volumetric flow rate (Equation
209 1). The total load of P at the sampling point is then a linear combination of the loads from
210 diffuse and point source inputs, as shown in Eq. (1).

$$211 \quad PC = dso_PC + pso_PC = A \cdot Q^{B-1} + C \cdot Q^{D-1} \quad (1)$$

212 where PC is phosphorus concentration, dso_PC is diffuse source originated PC, and pso_PC
213 the point-source originated PC. Q is discharge (daily accumulated), while A, C, (proportionality
214 constants) and B, D (exponents) are empirically determined parameters. Parameter estimation
215 was conducted by using a nonlinear generalized reduced gradient method to select values that
216 minimize the residual sum of squares. Parameter B was constrained to values lower than 1
217 (dilution effect over point contributions) and D values greater than 1 (at no flow, the diffuse
218 inputs tend to be zero and to increase with increasing flow). Each established C-Q relationship
219 was used for the calculation of daily mean concentrations and then multiplied by the daily
220 discharge to achieve daily exports. The proportional annual contribution from point sources and
221 diffuse sources was also calculated with this method.

222 For FLASHY catchments, we estimated the maximum P contribution from human inhabitants
223 based on the composition of household wastewater (i.e. urine, faeces, and greywater) and
224 biodegradable solid waste per person and year based on Vinnerås (2002). For STABLE
225 catchments, we estimated the total annual load from scattered dwellings not connected to

226 sewage treatment plants and stormwater outlets from validated models (Wiberg-Larsen et al.,
227 2013).

228 The relative contribution of PP to total exported P was estimated based on data from low
229 frequency instantaneous-grab sampling and linear interpolation, high frequency composite
230 sampling, and flow-weighted concentrations (FWC) estimated from high frequency composite
231 samples on a monthly basis. FWC estimation allows calculation of a flow-normalized
232 comparison of P concentrations between catchments.

233

234 **3 Results**

235 **3.1. Climate and hydrology**

236 The climate characteristics of the study period can be considered typical years for both Denmark
237 and Uruguay. During the study period, minimum, mean, median, and maximum air
238 temperatures were between 8 and 12 °C lower in the temperate/STABLE catchments than in
239 the subtropical/FLASHY catchments (Fig. 1). The annual average temperature in the
240 temperate/STABLE catchments was around 8.8 °C and ranged between -7.0 to 20.4 °C. The
241 corresponding figures for the subtropical/FLASHY catchments were around 17.5 °C and ranged
242 between 3.7 to 32.2 °C.

243 In both climates, catchments showed similar intra-yearly distributed rain patterns, but with
244 marked differences in frequency and intensity (Fig. 2). In the STABLE catchments, it rained
245 almost 6 out of 10 days (58%), the rain frequency being nearly half in the FLASHY catchments
246 (31%). Although there were more rainy days in the STABLE catchments, the daily average
247 amount of rainfall was lower (3.4 mm d⁻¹) than in FLASHY catchment where it amounted to
248 10.7 mm d⁻¹. Only one event of 50 mm d⁻¹ was registered in the STABLE catchments during
249 the 2-year study period, while in the FLASHY catchments rainfall events > 50 mm d⁻¹ occurred
250 approximately 1.5% of the days, reaching extremes of > 100 mm d⁻¹. The annual rainfall was
251 1.44 times higher in the FLASHY than in the STABLE catchments (Table 2).

252 Most of the water flowing in the FLASHY streams was exported during stormflow conditions
253 (stormflow contribution > 60.8%), while in the STABLE streams water was exported during
254 baseflow conditions (stormflow contribution < 36.4%; Table 2). The STABLE/low-LUI stream

255 showed a very different hydrological behavior than the other three streams in that a very high
256 percentage of the rainfall was discharged ($> 62\%$), with high minimum flows and low temporal
257 variability (Fig. 2 and Table 2).

258 The Danish streams exhibited stable hydrological behaviour characterized by low inter-annual
259 variability of total discharge, and also low variability at daily scale (the R-B Index never
260 reached values higher than 0.3; Table 2). In contrast, the Uruguayan streams could be classified
261 as FLASHY systems, with an R-B index ranging around 1 (0.9-1.3; Table 2). The stream
262 draining the high-LUI catchments was the most flashy (Table 2).

263

264 **3.2. Phosphorus temporal dynamics**

265 Total P concentrations were positively correlated with all other P fractions (PP, TDP, SRP,
266 SNRP) in STABLE and FLASHY/high-LUI streams, whereas in the low-LUI streams TP only
267 showed a significant relationship with PP in the STABLE and with TDP and PP in the FLASHY
268 stream (Table 3). The relationships between TDP and SRP were weaker but significant ($p <$
269 0.05) in the low-LUI streams than in the high-LUI ones under both climatic conditions (Table
270 3). The contributions of PP to TP were relatively similar in the low and high-LUI catchments
271 in STABLE, but in FLASHY the proportion of PP decreased with declining intensity of land
272 use (Table 3). The strongest relationships between PP and TP were found in both STABLE
273 streams and the FLASHY/low-LUI stream (Table 3). In contrast, in FLASHY/high-LUI, TDP,
274 and particularly SNRP, showed the strongest relationship with TP (Table 3). Negative
275 relationships were found solely for low-LUI streams, between PP and TDP, PP and SNRP for
276 the STABLE stream and between SNRP and SRP for the FLASHY stream (Table 3).

277 Median TP concentrations calculated for the four streams differed significantly ($H = 107.8$; $p \leq$
278 0.001), being pronouncedly higher in the FLASHY/high-LUI stream than in any of the others
279 (min = 271; med = 1.024; max = 4436 $\mu\text{g P L}^{-1}$; Fig. 3). All other paired comparisons of TP
280 revealed no significant differences, except for the STABLE streams where the TP concentration
281 was significantly lower in the stream draining the high-LUI catchment (median = 76 $\mu\text{g P L}^{-1}$)
282 than in the low-LUI catchment (med = 108 $\mu\text{g P L}^{-1}$; Fig. 3). No differences were registered
283 between the STABLE and FLASHY/low-LUI catchments (med = 100 $\mu\text{g P L}^{-1}$; Fig. 3).

284 A significant difference ($H = 43.6$; $p \leq 0.001$) in median PP concentrations was found between
285 most streams (Fig. 3), with highest and lowest values being registered in the FLASHY high-
286 LUI and low-LUI streams (median = $146 \mu\text{g P L}^{-1}$ and $25 \mu\text{g P L}^{-1}$, respectively) and
287 intermediate values in the STABLE streams (median = $52 \mu\text{g P L}^{-1}$ in the high-LUI, and $80 \mu\text{g}$
288 P L^{-1} in the low-LUI stream). Particulate P concentrations were highest in the STABLE low-
289 LUI stream and *vice versa* in the FLASHY streams. As expected, the STABLE streams
290 exhibited lower temporal variation in PP than the FLASHY streams (IQR= 23-37 and 53-227
291 $\mu\text{g P L}^{-1}$, respectively).

292 Furthermore, a significant difference in median TDP concentrations occurred ($H = 133.3$; $p \leq$
293 0.001 ; Fig. 3). *Post hoc* analysis revealed statistical equivalence only for the STABLE streams
294 (median = $28 \mu\text{g P L}^{-1}$ and $23 \mu\text{g P L}^{-1}$ for low and high-LUI streams). Intermediate TDP
295 concentrations were found in FLASHY/low-LUI and the highest concentrations appeared in the
296 FLASHY/ high-LUI stream (median = $74 \mu\text{g P L}^{-1}$ and $756 \mu\text{g P L}^{-1}$ respectively; Fig. 3).

297 The median SRP concentrations also exhibited statistically significant differences between the
298 streams ($H = 141.2$; $p \leq 0.001$; Fig. 3). SRP levels resembled TDP, with the lowest
299 concentrations in the STABLE streams (median: $2 \mu\text{g P L}^{-1}$ in both), intermediate levels in the
300 FLASHY/low-LUI stream (median: $45 \mu\text{g P L}^{-1}$), and the highest levels in the FLASHY/ high-
301 LUI (median: $659 \mu\text{g P L}^{-1}$; Fig. 3). SRP in the STABLE streams never exceeded $23 \mu\text{g P L}^{-1}$,
302 and in the FLASHY/low-LUI stream it never exceeded $87 \mu\text{g P L}^{-1}$. In contrast, the
303 FLASHY/high-LUI stream never had SRP concentrations lower than $219 \mu\text{g P L}^{-1}$ and SRP
304 reached a maximum concentration of $1,920 \mu\text{g P L}^{-1}$ (Fig. 3).

305

306 **3.3. Modelling phosphorus inputs from diffuse and point sources**

307 Graphical exploration of C-Q relationships for the FLASHY streams showed the typical pattern
308 described by Bowes et al. (2008), with high TP concentrations at low discharges followed by
309 steeply declining TP concentrations with increasing discharge (dilution associated with point
310 source-originated P input), and a less pronounced increase in concentrations at higher
311 discharges (associated with diffuse source-originated P inputs; Fig. 4). The C-Q relationships
312 for the two STABLE streams did not show any dilution effect associated with point source
313 inputs; therefore, the best fitting was obtained when considering only a diffuse input signal (Fig.

314 4; Table 4). The performance of the models evaluated with the Nash-Sutcliffe model efficiency
315 coefficient was generally low (Moriassi et al., 2007), reaching a maximum value of 0.25 for the
316 STABLE/low-LUI stream (Table 4).

317 When considering the relationships established for point source-originated TP for the FLASHY
318 streams, we found a higher exponent (B) in the C-Q relationships for the high-LUI catchment
319 (Table 4). As a consequence, the decrease in TP with increasing flow (the dilution effect) was
320 less pronounced for the high than the low-LUI stream (at $1,000 \text{ L s}^{-1}$ the FLASHY/high-LUI
321 catchment reached $85 \mu\text{g P L}^{-1}$, while the FLASHY/low-LUI dropped to $5 \mu\text{g P L}^{-1}$, Fig. 4).

322 Considering only diffuse sources of TP, we found a higher coefficient (C) but slightly lower
323 exponents (D) in the C-Q relationships established for the high-LUI than for the low-LUI
324 catchments under both climate conditions (Fig. 4 and Table 4). However, the FLASHY/high-
325 LUI stream always had higher TP concentrations from diffuse sources than the other streams
326 (Fig. 4).

327

328 **3.4. Estimation of phosphorus export**

329 Comparing TP export estimates based on the high frequency composite sampling, we found an
330 underestimation pattern when applying the low frequency sampling/linear interpolation method
331 for both STABLE catchments and an overestimation for the FLASHY catchments (Table 6).
332 This bias was always higher for the low than for the high-LUI catchments (Table 6).

333 The TP and TDP export from the FLASHY/high-LUI catchment was higher than in the other
334 three catchments (Table 5). Moreover, for STABLE streams, the TP export was always higher
335 from the low-LUI than from the high-LUI catchment (Table 5).

336 Also comparing with the high frequency composite estimates, the C-Q relationships used to
337 calculate exported TP produced more accurate results than the linear interpolation for the two
338 high-LUI catchments, irrespective of climatic region (Table 6). The largest and disproportionate
339 deviations of exported TP were obtained when applying the C-Q model to the FLASHY/low-
340 LUI catchment compared to the high frequency composite sampling estimates (364-400%;
341 Table 6).

342 The contribution of PP to exported P was never lower than 65% of the annual exported TP in
343 the STABLE catchments (Table 5). A contrasting pattern was recorded for FLASHY streams
344 where the contribution of PP never exceeded 48% of TP, reaching values as low as 13.6%
345 (Table 5). This pattern of a major contribution of PP in the STABLE catchments repeats itself
346 in the estimations made with the low frequency instantaneous-grab samples, high frequency
347 composite samples, and flow-weighted concentrations (Fig. 5). We found a tendency to a
348 higher, though rarely significant, dissolved P contribution in streams draining high-LUI
349 catchments (Fig. 5).

350 The estimated contributions of TP from point sources and diffuse sources indicated that most
351 of the TP export from the STABLE catchments came from diffuse sources as point source
352 contribution from human sources only reached a maximum of 18% of the exported P (Table 5),
353 but it was still too low to be detectable in our established C-Q relationships as point sources
354 (Table 4). Contrarily, in the FLASHY catchments point sources dominated the P export, always
355 constituting more than 83% of the exported P, with human sources contributing < 8%, dairy
356 cattle being the most probable source of the remaining P (Table 5).

357

358 **4 Discussion**

359 Our results show that climatic and hydrological variability affects the temporal dynamics of P
360 in streams and that low frequency monitoring strategies may fail to adequately capture such
361 dynamics.

362 As expected, we found strong concordance between climatic characteristics and stream
363 hydrological regimes. Danish catchments have a lower rainfall that is more evenly distributed
364 during the year, with rare or no extreme rainfall events ($> 50 \text{ mm day}^{-1}$) and lower temperature
365 (and therefore low evapotranspiration). These conditions resulted in stream flows characterized
366 by higher stability (lower flashiness, R-B index = 0.1-0.3 in our study). In contrast, the
367 Uruguayan streams exhibited a much higher contribution of stormflow to total flow and higher
368 flashiness (R-B index = 0.9-1.3), reflecting the local climate and likely also soil conditions. As
369 reference, reported data in the literature includes daily maximum values of R-B Index reaching
370 0.43 in mountain streams in Slovakia and Austria (Holko et al., 2011), 0.65 in 30 agricultural
371 dominated catchments in seven Nordic and Baltic countries (Deelstra et al., 2014), 1.01 at 204

372 stations in Michigan temperate streams (USA) (Fongers, 2012), and 1.32 at 515 stations in
373 catchments located in six US Midwestern States (Baker et al., 2004).

374

375 The calculated annual TP export values in all streams fell within the range reported in the
376 literature for comparable micro-catchments, for instance streams with grassland-agriculture
377 production in Ireland (0.89 to 3.98 kg P ha⁻¹ y⁻¹; Campbell et al., 2015) Irish streams with
378 catchment farming activities (0.12 to 0.83 kg P ha⁻¹ y⁻¹; Melland et al., 2012), and Norwegian
379 streams (0.5 to 5.8 kg P ha⁻¹ y⁻¹; Kronvang et al., 2005a). The temporal variability of exported
380 PP followed the hydrological variability in the streams (Fig. 2), but the amount of exported P
381 and PP (the latter likely derived from diffuse sources) did not systematically increase with
382 increasing variability. This was valid for both Uruguayan catchments where the highest and the
383 lowest P loads were exported, and dissolved P forms always predominated over particulate
384 forms. The pattern of P loads exported in relation to conditions of high hydrological variability
385 was thus opposite to our *a priori* expectations (i.e. higher total and particulate P export from
386 diffuse sources in connection with higher flashiness) (Table 5), likely due to other factors
387 related to land use, less input of P with eroded stream bank material due to potential low content
388 of P and maybe also less erosion (Kronvang et al., 2012) and particularly the presence/absence
389 of point sources in the catchments.

390 Several factors may contribute to the predominance of dissolved P forms in the Uruguayan
391 streams. The direct access of cattle to the stream channels is one of those reasons, being a
392 practice that results in direct manure deposition in the water and trampling, and mobilization
393 from stream bed sediments (James et al., 2007; Jarvie et al., 2010; Kronvang et al., 2012; Laubel
394 et al., 2003; Sheffield et al., 1997; Trimble and Mendel, 1995). In the FLASHY/high-LUI
395 catchment this contribution was exacerbated and further aggravated by the additional effects of
396 the lack of slurry treatment in dairy facilities and the widespread no-till practices associated
397 with the application of fertilizers over the soil surface (Sharpley and Smith, 1994; Sharpley et
398 al., 1996). Some or all of these factors together may explain the high levels of TP, TDP, SRP,
399 and SNRP, exceeding the maximum range found in 35 comparable Nordic/Baltic micro-
400 catchments studied by Kronvang et al. (2007).

401 The contribution of TP from point sources was negligible in both STABLE catchment, but was
402 always higher than 83% in the two FLASHY catchments irrespective of land use intensity. The

403 magnitude of P point sources seems to have a much stronger influence on the hydrochemistry
404 of the Uruguayan stream waters than do hydrological variability and flashiness *per se*. In
405 contrast, the Danish pattern seems to be consistent with the well-documented reduced influence
406 of P from point sources in northern Europe (European Environment Agency, 2005; Kronvang
407 et al., 2005b). Apart from the climatic and hydrological differences between the analyzed
408 catchments, the extremely high TP concentrations (as high as 4,436 $\mu\text{g P L}^{-1}$), P exports (as
409 high as 5.20 $\text{kg P ha}^{-1} \text{ year}^{-1}$), and the extremely high proportion of dissolved P (as high as
410 86.4%) estimated for the FLASHY/high-LUI catchment may have implications for the
411 environmental regulations of farming production in Uruguay given the severe deterioration of
412 water quality in our case study. Our results also provide insight into the future behavior of P in
413 northern European temperate streams seen in the context of the predicted change in climate
414 towards more extreme, warmer, and wetter conditions, probably giving flashier streams
415 (Hanssen-Bauer et al., 2005).

416 The importance of understanding hydrology-driven variations in nutrient discharge will most
417 likely increase in the near future. In our study, the performance of the different monitoring
418 frequencies and P export estimation methods reflected the hydrological character or flashiness
419 of the investigated streams. The performance of the low frequency sampling programs in
420 estimating TP and TDP exports was low in comparison with the high frequency sampling
421 program. Although the performance of the models evaluated with the Nash-Sutcliffe efficiency
422 coefficient was generally low, the C-Q method did not have a comparatively poorer
423 performance.

424 Although our study was limited to four representative catchments (due to logistic reasons), our
425 results suggest that a clear interaction between climate/hydrology and land use intensity
426 occurred. This was shown by a lower deviation in the exported P estimations of the low
427 frequency sampling for the high-LUI than for the low-LUI catchments. However, the low
428 frequency sampling and linear interpolation method performed equally poor for the FLASHY
429 and the STABLE streams. Our results suggest that climate and hydrological conditions may
430 promote/may yield a bias in P load estimations at low sampling frequency, with a tendency
431 towards underestimation for the hydrologically STABLE streams and overestimation for the
432 FLASHY streams (Table 6). These results are in accordance with the findings of Richards and
433 Holloway (1987) that the loads of P from diffuse sources (whose water concentration increases
434 with runoff) tend to be underestimated, while the loads of P from point sources (whose

435 concentration decreases with runoff) tend to be overestimated. This underlines the potentially
436 high inadequacy of low frequency sampling programs to properly depict stream dynamics,
437 which is consistent with previous findings (e.g. Jones et al., 2012).

438 Research into P dynamics in subtropical FLASHY ~~flashy~~ streams is in its initial phase. The
439 expected intensification of agricultural production in many regions of the world, such as in
440 southern South America, highlights the need for appropriate stream monitoring programs.
441 Accurate estimation of P temporal dynamics and exports can help explain the linkages between
442 climate, hydrology, land use, and water quality. Based on our findings, we suggest that the
443 evaluation and use of more accurate monitoring methods, such as automatized flow-
444 proportional water samplers and automatized bankside analysers, should be prioritized
445 whenever is logistically possible. However, it seems particularly relevant in currently flashy
446 systems and also in systems where climate change predictions suggest an increase in stream
447 flashiness.

448 **Acknowledgements**

449 We thank the technicians of AU in Silkeborg for their work in the field and lab. We deeply
450 acknowledge ~~Carsten-~~ ~~Brian-~~ Nielsen for technical assistance in the set-up of monitoring
451 stations in Uruguay. We are grateful for the field assistance from colleagues and students, the
452 logistic support of ~~D-~~ aniella Agrati and family, and landowners (Mendiverri & ~~Θ-~~ Lature) in
453 Uruguay. We are grateful to ~~A-~~ nne ~~Mette-~~ Poulsen for manuscript editing.

454 GG, FTM, MM, IGB and NM received support from the SNI (Agencia Nacional de
455 Investigación e Innovación, ANII, Uruguay). GG was supported by a PhD scholarship from
456 PEDECIBA. The study was funded by the Danish Council for Independent Research, a grant
457 by ANII-FCE (2009-2749) Uruguay, and the National L'Oreal-UNESCO award for Women in
458 Science-Uruguay with support of DICYT granted to MM.

459 We also wish to thank the generous remarks by the Reviewers and Editor that helped us improve
460 this manuscript.

461 **References**

462 Allaby, M.: Grasslands, Biomes of the Earth, Chelsea House, 270 pp., 2006.

463 Arnold, J. G., Allen, P. M., Muttiah, R., and Bernhardt, G.: Automated base flow separation
464 and recession analysis techniques. *Ground Water*, 33, 1010-1018, 10.1111/j.1745-
465 6584.1995.tb00046.x, 1995.

466 Baker, D. B., Richards, R. P., Loftus, T. T., and Kramer, J. W.: A new flashiness index:
467 Characteristics and applications to midwestern rivers and streams. *J. Am. Water Resour. As.*,
468 40, 503-522, 10.1111/j.1752-1688.2004.tb01046.x, 2004.

469 Bormann, F. H., and Likens, G. E.: Nutrient cycling. *Science*, 155, 424-428, 1967.

470 Bowes, M. J., Smith, J. T., Jarvie, H. P., and Neal, C.: Modelling of phosphorus inputs to rivers
471 from diffuse and point sources. *Sci. Total Environ.*, 395, 125-138,
472 <http://dx.doi.org/10.1016/j.scitotenv.2008.01.054>, 2008.

473 Campbell, J. M., Jordan, P., and Arnscheidt, J.: Using high-resolution phosphorus data to
474 investigate mitigation measures in headwater river catchments. *Hydrol. Earth Syst. Sci.*, 19,
475 453-464, 10.5194/hessd-11-10965-2014, 2015.

476 Carpenter, S. R., Caraco, N. F., Correll, D. L., Howarth, R. W., Sharpley, A. N., and Smith, V.
477 H.: Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecol. Appl.*, 8, 559-
478 568, doi:10.1890/1051-0761(1998)008[0559:NPOSWW]2.0.CO;2, 1998.

479 Cassidy, R., and Jordan, P.: Limitations of instantaneous water quality sampling in surface-
480 water catchments: Comparison with near-continuous phosphorus time-series data. *J. Hydrol.*,
481 405, 182-193, <http://dx.doi.org/10.1016/j.jhydrol.2011.05.020>, 2011.

482 Castaño, J. P., Giménez, A., Ceroni, M., Furest, J., and Aunchayna, R.: Caracterización
483 agroclimática del Uruguay 1980-2009, INIA, Montevideo, 34, 2011.

484 Cordell, D., Drangert, J. O., and White, S.: The story of phosphorus: Global food security and
485 food for thought. *Global Environmental Change*, 19, 292-305, 2009.

486 Deelstra, J., Iital, A., Povilaitis, A., Kyllmar, K., Greipsland, I., Blicher-Mathiesen, G., Jansons,
487 V., Koskiaho, J., and Lagzdins, A.: Hydrological pathways and nitrogen runoff in agricultural
488 dominated catchments in Nordic and Baltic countries. *Agr. Ecosyst. Environ.*, 195, 211-219,
489 <http://dx.doi.org/10.1016/j.agee.2014.06.007>, 2014.

490 Defew, L. H., May, L., and Heal, K. V.: Uncertainties in estimated phosphorus loads as a
491 function of different sampling frequencies and common calculation methods. *Mar. Freshwater*
492 *Res.*, 64, 373-386, <http://dx.doi.org/10.1071/MF12097>, 2013.

493 Derpsch, R., Friedrich, T., Kassam, A., and Hongwen, L.: Current status of adoption of no-till
494 farming in the world and some of its main benefits. *IJABE*, 3, 1-25, 2010.

495 DMI: Precipitation and sun in Denmark, Danish Meteorological Institute, Copenhagen, 2015.

496 Dunn, O. J.: Multiple comparisons using rank sums. *Technometrics*, 6, 241-252, 1964.

497 Elser, J., and Bennett, E.: Phosphorus cycle: A broken biogeochemical cycle. *Nature*, 478, 29-
498 31, 2011.

499 European Environment Agency: Source apportionment of nitrogen and phosphorus inputs into
500 the aquatic environment, European Environment Agency, Copenhagen, Denmark, 1-48, 2005.

501 Fongers, D.: Application of the Richards-Baker Flashiness Index to gaged Michigan rivers and
502 streams, Water Resources Division, Michigan Department of Environmental Quality, 141,
503 2012.

504 Grant, R., Laubel, A., Kronvang, B., Andersen, H. E., Svendsen, L. M., and Fuglsang, A.: Loss
505 of dissolved and particulate phosphorus from arable catchments by subsurface drainage. *Water*
506 *Res.*, 30, 2633-2642, [http://dx.doi.org/10.1016/S0043-1354\(96\)00164-9](http://dx.doi.org/10.1016/S0043-1354(96)00164-9), 1996.

507 Hanssen-Bauer, I., Achberger, C., Benestad, R. E., Chen, D., and Førlund, E. J.: Statistical
508 downscaling of climate scenarios over Scandinavia. *Clim. Res.*, 29, 255-268, 2005.

509 Haygarth, P. M., Heathwaite, A. L., Jarvis, S. C., and Harrod, T. R.: Hydrological factors for
510 phosphorus transfer from agricultural soils, in: *Advances in Agronomy*, edited by: Donald, L.
511 S., Academic Press, San Diego, CA, 153-178, 1999.

512 Holko, L., Parajka, J., Kostka, Z., Škoda, P., and Blöschl, G.: Flashiness of mountain streams
513 in Slovakia and Austria. *J. Hydrol.*, 405, 392-401,
514 <http://dx.doi.org/10.1016/j.jhydrol.2011.05.038>, 2011.

515 James, E., Kleinman, P., Veith, T., Stedman, R., and Sharpley, A.: Phosphorus contributions
516 from pastured dairy cattle to streams of the Cannonsville Watershed, New York. *J. Soil Water*
517 *Conserv.*, 62, 40-47, 2007.

518 Jarvie, H. P., Withers, P. J. A., Bowes, M. J., Palmer-Felgate, E. J., Harper, D. M., Wasiak, K.,
519 Wasiak, P., Hodgkinson, R. A., Bates, A., Stoate, C., Neal, M., Wickham, H. D., Harman, S.
520 A., and Armstrong, L. K.: Streamwater phosphorus and nitrogen across a gradient in rural–
521 agricultural land use intensity. *Agr. Ecosyst. Environ.*, 135, 238-252,
522 <http://dx.doi.org/10.1016/j.agee.2009.10.002>, 2010.

523 Jones, A. S., Horsburgh, J. S., Mesner, N. O., Ryel, R. J., and Stevens, D. K.: Influence of
524 sampling frequency on estimation of annual total phosphorus and total suspended solids loads.
525 *J. Am. Water Resour. As.*, 48, 1258-1275, [10.1016/j.agwat.2009.01.011](http://dx.doi.org/10.1016/j.agwat.2009.01.011), 2012.

526 Jordan, P., and Cassidy, R.: Technical Note: Assessing a 24/7 solution for monitoring water
527 quality loads in small river catchments. *Hydrol. Earth Syst. Sci.*, 15, 3093-3100, [10.5194/hess-](http://dx.doi.org/10.5194/hess-15-3093-2011)
528 [15-3093-2011](http://dx.doi.org/10.5194/hess-15-3093-2011), 2011.

529 Kronvang, B., and Bruhn, A. J.: Choice of sampling strategy and estimation method for
530 calculating nitrogen and phosphorus transport in small lowland streams. *Hydrol. Process.*, 10,

531 1483-1501, 10.1002/(SICI)1099-1085(199611)10:11<1483::AID-HYP386>3.0.CO;2-Y,
532 1996.

533 Kronvang, B., Bechmann, M., Lundekvam, H., Behrendt, H., Rubaek, G. H., Schoumans, O.
534 F., Syversen, N., Andersen, H. E., and Hoffmann, C. C.: Phosphorus losses from agricultural
535 areas in river basins: Effects and uncertainties of targeted mitigation measures. *Journal of*
536 *Environmental Quality*, 34, 2129-2144, 10.2134/jeq2004.0439, 2005a.

537 Kronvang, B., Jeppesen, E., Conley, D. J., Søndergaard, M., Larsen, S. E., Ovesen, N. B., and
538 Carstensen, J.: Nutrient pressures and ecological responses to nutrient loading reductions in
539 Danish streams, lakes and coastal waters. *J. Hydrol.*, 304, 274-288, 2005b.

540 Kronvang, B., Vagstad, N., Behrendt, H., Bøgestrand, J., and Larsen, S. E.: Phosphorus losses
541 at the catchment scale within Europe: an overview. *Soil Use Manage.*, 23, 104-116,
542 10.1111/j.1475-2743.2007.00113.x, 2007.

543 Kronvang, B., Audet, J., Baattrup-Pedersen, A., Jensen, H. S., and Larsen, S. E.: Phosphorus
544 Load to Surface Water from Bank Erosion in a Danish Lowland River Basin. *J. Environ. Qual.*,
545 41, 304-313, 10.2134/jeq2010.0434, 2012.

546 Laubel, A., Kronvang, B., Hald, A. B., and Jensen, C.: Hydromorphological and biological
547 factors influencing sediment and phosphorus loss via bank erosion in small lowland rural
548 streams in Denmark. *Hydrol. Process.*, 17, 3443-3463, 10.1002/hyp.1302, 2003.

549 Melland, A. R., Mellander, P. E., Murphy, P. N. C., Wall, D. P., Mechan, S., Shine, O., Shortle,
550 G., and Jordan, P.: Stream water quality in intensive cereal cropping catchments with regulated
551 nutrient management. *Environ. Sci. Policy*, 24, 58-70,
552 <http://dx.doi.org/10.1016/j.envsci.2012.06.006>, 2012.

553 Meyer, J. L., and Likens, G. E.: Transport and transformation of phosphorus in a forest stream
554 ecosystem. *Ecology*, 60, 1255-1269, 1979.

555 Moriasi, D. N., Arnold, J. G., Van Liew, M. W., Bingner, R. L., Harmel, R. D., and Veith, T.
556 L.: Model evaluation guidelines for systematic quantification of accuracy in watershed
557 simulations. *T. ASABE*, 50, 885-900, 2007.

558 8° censo nacional de población 2011:
559 <http://www.ine.gub.uy/censos2011/resultadosfinales/florida.html>, access: 01/01/2015, 2015.

560 Richards, R. P., and Holloway, J.: Monte Carlo studies of sampling strategies for estimating
561 tributary loads. *Water Resour. Res.*, 23, 1939-1948, 10.1029/WR023i010p01939, 1987.

562 Sharpley, A. N., and Smith, S. J.: Wheat tillage and water quality in the Southern plains. *Soil*
563 *and Tillage Research*, 30, 33-48, [http://dx.doi.org/10.1016/0167-1987\(94\)90149-X](http://dx.doi.org/10.1016/0167-1987(94)90149-X), 1994.

564 Sharpley, A. N., Daniel, T. C., Sims, J. T., and Pote, D. H.: Determining environmentally sound
565 soil phosphorus levels. *J. Soil Water Conserv.*, 51, 160-166, 1996.

566 Sheffield, R. E., Mostaghimi, S., Vaughan, D., Collins Jr, E., and Allen, V.: Off-stream water
567 sources for grazing cattle as a stream bank stabilization and water quality BMP. *T. ASABE*, 40,
568 595-604, 1997.

569 Stelzer, R. S., and Likens, G. E.: Effects of sampling frequency on estimates of dissolved silica
570 export by streams: The role of hydrological variability and concentration-discharge
571 relationships. *Water Resour. Res.*, 42, 1-10, 2006.

572 Trimble, S. W., and Mendel, A. C.: The cow as a geomorphic agent — A critical review.
573 *Geomorphology*, 13, 233-253, [http://dx.doi.org/10.1016/0169-555X\(95\)00028-4](http://dx.doi.org/10.1016/0169-555X(95)00028-4), 1995.

574 Valderrama, J. C.: The simultaneous analysis of total nitrogen and total phosphorus in natural
575 waters. *Mar. Chem.*, 10, 109-122., 1981.

576 Vinnerås, B.: Possibilities for sustainable nutrient recycling by faecal separation combined with
577 urine diversion, PhD, Acta Universitatis Agriculturae Sueciae, Agraria 353, Swedish University
578 of Agricultural Sciences, Uppsala, 2002.

579 Wiberg-Larsen, P., Windolf, J., Bøgestrand, J., Baattrup-Pedersen, A., Kristensen, E. A.,
580 Larsen, S. E., Thodsen, H., Ovesen, N. B., Bjerring, R., Kronvang, B., and Kjeldgaard, A.:
581 Vandløb 2012. NOVANA, Aarhus Universitet, DCE - Nationalt Center for Miljø og Energi, 84
582 s. - Videnskabelig rapport fra DCE - Nationalt Center for Miljø og Energi nr. 75.
583 <http://dce2.au.dk/pub/SR75.pdf>, 2013.

584 Wu, J. Y., Thompson, J. R., Kolka, R. K., Franz, K. J., and Stewart, T. W.: Using the Storm
585 Water Management Model to predict urban headwater stream hydrological response to climate
586 and land cover change. *Hydrol. Earth Syst. Sci.*, 17, 4743-4758, 10.5194/hess-17-4743-2013,
587 2013.

588 Zar, J. H.: *Biostatistical Analysis*, 5th. edition ed., Prentice-Hall, Upper Saddle River, NJ, 2010.
589

590 Table 1. Site coordinates (datum WGS84), name and catchment size, dominant soils and land
 591 use. Danish catchments are part of Gudenå River basin, and the Uruguayan streams are part of
 592 the Santa Lucía Chico River basin. Source: (*) World Reference Soil Database classification,
 593 European Commission and European Soil Bureau Network (2004); (#) SOTERLAC database,
 594 ISRIC Foundation, (www.isric.org). STABLE: low flashiness Danish streams; FLASHY:
 595 Uruguayan flashy streams; low and high-LUI: low and high land use intensity.

596

Id	Name & size	Coordinates	Dominant soils	Land use (area %)
STABLE low-LUI	Granslev 740 ha	56°17'2"N 9°53'51"E	Haplic Luvisols (*)	Forest (59); arable farming (29); pasture/meadow (7);
STABLE high-LUI	Gelbæk 1180 ha	56°13'29"N 9°48'41"E	Gleyic Luvisols (*)	Arable farming (92); forest (2); urban (1); other (5)
FLASHY low-LUI	Chal-Chal 1880 ha	33°49'31"S 56°16'55"W	Luvic Phaeozem & Eutric	Extensive pasture (~70); arable farming (~30)
FLASHY high-LUI	Pantanoso 840 ha	33°54'13"S 56°00'23"W	Eutric Regosols (#)	Arable farming and dairy farms (90); extensive pasture (7); urban (3)

597

598

599 Table 2. Yearly accumulated rainfall and runoff in mm (direct measures), runoff ratio
600 (percentage of rainfall water exported as runoff), and R-B Index (Richards-Baker Index) of
601 flashiness for each monitored year. Stormflow contribution: Percent contribution of
602 stormflow to total flow was estimated for the complete data set (2 years). STABLE: low
603 flashiness Danish streams; FLASHY: Uruguayan flashy streams; low and high-LUI: low and
604 high land use intensity.

605

	STABLE	STABLE	FLASHY	FLASHY
	low-LUI	high-LUI	low-LUI	high-LUI
Accumulated rainfall 1st year	770 mm	778 mm	1030 mm	1196 mm
Accumulated rainfall 2nd year	756 mm	766 mm	1010 mm	1405 mm
Total accumulated runoff 1st year	515 mm	223 mm	170 mm	235 mm
Total accumulated runoff 2nd year	472 mm	198 mm	294 mm	255 mm
Runoff ratio 1st year	66.9%	28.6%	16.5%	19.6%
Runoff ratio 2nd year	62.4%	25.9%	29.1%	18.2%
R-B Index 1st year	0.1	0.3	1.0	1.3
R-B Index 2nd year	0.1	0.3	0.9	1.2
Stormflow contribution (%)	11.8%	36.4%	60.8%	70.6%

606

607

608 Table 3. Correlation matrices of total (TP), particulate (PP), total dissolved (TDP), soluble
609 reactive (SRP), and soluble non-reactive (SNRP) phosphorus from instantaneous-grab
610 samples. Upper and lower triangles of each matrix refer to high and low-LUI conditions,
611 respectively. Numeric values represent Spearman rank order correlation and were included
612 only when significant ($p \leq 0.05$). ns: non-significant. STABLE: low flashiness Danish
613 streams; FLASHY: Uruguayan flashy streams; low and high-LUI: low and high land use
614 intensity.

	TP	PP	TDP	SRP	SNRP		TP	PP	TDP	SRP	SNRP		
TP	\	0.78	0.75	0.68	0.57	STABLE/high-LUI	TP	\	0.63	0.90	0.67	0.84	FLASHY/high-LUI
PP	0.81	\	ns	ns	ns		PP	0.80	\	ns	ns	0.41	
TDP	ns	-0.31	\	0.93	0.70		TDP	0.58	ns	\	0.83	0.81	
SRP	ns	ns	0.86	\	0.44		SRP	ns	ns	0.56	\	0.40	
SNRP	ns	-0.34	0.56	ns	\		SNRP	ns	ns	0.41	-0.37	\	
STABLE/low-LUI							FLASHY/low-LUI						

615

616 Table 4. Estimated parameters for the load apportionment model fitted for each stream. RSS:
 617 residual sum of squares. NSC: Nash-Sutcliffe model efficiency coefficient. STABLE: low
 618 flashiness Danish streams; FLASHY: Uruguayan flashy streams; low and high-LUI: low and
 619 high land use intensity.

620

Source	Parameter	STABLE	STABLE	FLASHY	FLASHY
		low-LUI	high-LUI	low-LUI	high-LUI
Point	A	0	0	1915	2550
	B	-	-	0.140	0.501
Diffuse	C	7.145	20.677	3.658	399.000
	D	1.58	1.40	1.64	1.15
Global	RSS (10³)	42.5	460	253	36362
	NSC	0.25	0.12	0.12	0.10

621

622 Table 5. Alternative estimations of annual total phosphorus (TP) and total dissolved phosphorus (TDP) exported from the four catchments expressed
623 as kg P ha⁻¹ year⁻¹. Estimation strategies: “COMP”: high frequency composite sampling. “LFS-LI”: Low frequency sampling and linear
624 interpolation. “C-Q”: low frequency sampling and concentration-discharge relationships interpolation. % PP: percentage TP exported in particulate
625 form. “% hs” under COMP represents the percentage of annual exported P from human sources. “% ps” under C-Q represents the percentage of
626 the total annual exported load from point sources *sensu* the model. STABLE: low flashiness Danish streams; FLASHY: Uruguayan flashy streams;
627 low and high-LUI: low and high land use intensity.

		STABLE/low-LUI			STABLE/high-LUI			FLASHY/low-LUI			FLASHY/high-LUI		
		COMP	LFS-LI	C-Q	COMP	LFS-LI	C-Q	COMP	LFS-LI	C-Q	COMP	LFS-LI	C-Q
1 st year	TP	1.09		0.61	0.34		0.34	0.13		0.52	2.28		2.86
		(12.5% hs)	0.64	(0% ps)	(10.5% hs)	0.29	(0% ps)	(6.9% hs)	0.25	(89.7% ps)	(7.5% hs)	2.36	(83.1% ps)
	TDP	0.17	0.20	\	0.08	0.10	\	0.08	0.13	\	1.97	1.83	\
	% PP.	84.4	68.6	\	76.5	66.9	\	38.5	48	\	13.6	22.5	\
2 nd year	TP	0.74		0.54	0.35		0.33	0.25		0.91	5.20		5.19
		(17.7% hs)	0.47	(0% ps)	(10.4% hs)	0.25	(0% ps)	(3.6% hs)	0.27	(83,6% ps)	(4.2% hs)	5.76	(86.5% ps)
	TDP	0.10	0.11	\	0.07	0.06	\	0.14	0.21	\	4.07	4.7	\
	% PP.	86.5	76.4	\	80	76.9	\	44	22.2	\	21.7	18.4	\


628

629 Table 6. The percentages summarize the relative fit of alternative estimation methods in
630 relation to the reference annually exported load estimated by the composite sampling
631 programme (see Table 5 for references). 100% represents the same annual P exported in kg ha⁻¹
632 year⁻¹. Values below 100% (*italics*) represent underestimation (less than 90%) relative to the
633 estimation of composite sampling. Values over 110% (**bold**) represent overestimation relative
634 to the estimation of composite sampling. STABLE: low flashiness Danish streams; FLASHY:
635 Uruguayan flashy streams; low and high-LUI: low and high land use intensity.

	STABLE		STABLE		FLASHY		FLASHY		
	low-LUI		high-LUI		low-LUI		high-LUI		
	LFS-LI	C-Q	LFS-LI	C-Q	LFS-LI	C-Q	LFS-LI	C-Q	
1st year	TP	<i>58.7%</i>	<i>56.0%</i>	<i>85.3%</i>	100.0%	192.3%	400.0%	103.5%	125.4%
	TDP	117.6%	\	125.0%	\	162.5%	\	92.9%	\
2nd year	TP	<i>63.5%</i>	<i>73.0%</i>	<i>71.4%</i>	94.3%	108.0%	364.0%	110.8%	99.8%
	TDP	110.0%	\	<i>85.7%</i>	\	150.0%	\	115.5%	\

636

1 Fig. 1. Left: Mean daily air temperature variation for a temperate/STABLE (Danish) and a
2 subtropical/FLASHY (Uruguayan) catchment in 2011. Right: boxplots of the same data. The
3 boundary of the box closest to zero indicates the 25th percentile, a line within the box marks the
4 median, and the boundary of the box farthest from zero indicates the 75th percentile. Whiskers
5 above and below the box indicate the 90th and 10th percentiles. Black dots display outliers.





1 Fig. 2. Accumulated daily rainfall and specific runoff hydrographs for the four monitored
2 streams (temperate/STABLE: Danish; subtropical/FLASHY: Uruguayan). For each variable a
3 fixed scale was used to aid visual comparison. R-B Index (Richards-Baker Index) of
4 flashiness.



EXTENSIVE LAND USE

INTENSIVE LAND USE

STABLE



FLASHY



1 Fig. 3. From top to bottom: temporal variation of total (TP), particulate (PP), total dissolved
2 (TDP), and soluble reactive phosphorus (SRP) concentrations from grab samples for all the
3 monitored catchments. Log₁₀ scale was selected on the vertical axe to improve vizualisation.
4 The phosphorus concentration is always expressed as µg P L⁻¹. Right: boxplots are based on
5 the same data. Letters A, B, and C are used to display statistical groups according *post hoc*
6 paired comparison analysis. The boundary of the box closest to zero indicates the 25th
7 percentile, a line within the box marks the median, and the boundary of the box farthest from
8 zero indicates the 75th percentile. Whiskers above and below the box indicate the 90th and 10th
9 percentiles. Black dots display outliers. STABLE: low flashiness Danish streams; FLASHY:
10 Uruguayan flashy streams; low and high-LUI: low and high land use intensity.






Fig. 4. Scatterplots of total phosphorus (TP) concentrations of instantaneous-grab samples relative to discharge from the four streams. The dots connected by lines represent the predicted values according to the load apportionment model (see Table 4). For STABLE catchments, only the diffuse originated term of the model is included. Lowest graphs display the predicted TP concentration of all catchments from point (left) and diffuse sources (right) for a range of 0 to 1000 L s⁻¹. Note the Log₁₀ scale for TP. STABLE: low flashiness Danish streams; FLASHY: Uruguayan flashy streams; low and high-LUI: low and high land use intensity.

low Land Use Intesity





high Land Use Intesity




STABLE

FLASHY



MODELED POINT SOURCE TP CONTRIBUTION



MODELED DIFFUSE SOURCE TP CONTRIBUTION




Fig. 5. Contribution of particulate phosphorus (PP) to exported total phosphorus (TP). Left boxplots are based on instantaneous-grab sampling and linear interpolation estimation, centre boxplots are based on composite data, and the right boxplot on flow-weighted concentrations. Letters A, B, and C are used to display statistical groups according *post hoc* paired comparison analysis. The boundary of the box closest to zero indicates the 25th percentile, a line within the box marks the median, and the boundary of the box farthest from zero indicates the 75th percentile. Whiskers above and below the box indicate the 90th and 10th percentiles. STABLE: low flashiness Danish streams; FLASHY: Uruguayan flashy streams; low and high-LUI: low and high land use intensity.

