

Editor Decision: Publish subject to technical corrections (20 Dec 2015)

by Matjaz Mikos

line 589 & 592 - the two references should be twisted: Gamache and then Göke.

Answer: Thank you for your kindness comment. We changed the references list correctly.

line 656 - You et al., please check the journal title, volume and pages.

Answer: Thank you for your comment. According to Editor's recommend, we checked the structure of reference of this again and modified the form of reference is as follows;

[line 660-662]

You, C.-H., Lee, D.-I., and Kang, M.-Y.: Rainfall estimation using specific differential phase for the first operational polarimetric radar in Korea, *Advances in Meteorology*, 2014, 10 pages, 2014.

This reference form was modified from Endnote style form (Copernicus_Publications.ens) which provides HESS because of absence of page numbers.

Original form of reference shown by HESS's style is as follows

You, C.-H., Lee, D.-I., and Kang, M.-Y.: Rainfall estimation using specific differential phase for the first operational polarimetric radar in Korea, *Advances in Meteorology*, 2014, 2014.

Unfortunately, we could not find the policy of writing references form for online version. The journal of *Advances in Meteorology* operates by on-line version. Therefore, paper number was added in the form literally.

This paper number is started from 1 page.

If this form could cause some problem, please notify me.

Table 2 & Table 4 - please use dd month yyyy & hh:mm:ss style for dates - see http://www.hydrology-and-earth-system-sciences.net/for_authors/manuscript_preparation.html for details on Date and time

Answer: Thank you for your comment. The composition standard of HESS for date and time is as follows;

25 July 2007 (dd month yyyy), 15:17:02 (hh:mm:ss).

Often it is necessary to specify the time if referring to local time or universal time coordinated.

This can be done by adding "LT" or "UTC", respectively.

According to the composition standard of HESS, we modified the form of date correctly.

for Date style see also line 23 & 496, and for Time style lines 382-384, 388, 390-392, 395, 435-437, 439-440, 443, 446, 449, 455-459, 464.

Answer: Thank you for your comment. Same with aforementioned comment, we modified the form of time correctly.

Climatological Characteristics of Raindrop Size Distributions in Busan, Korea

Sung-Ho SUH¹

Cheol-Hwan YOU^{2*}

Email: youch@pknu.ac.kr

Dong-In LEE^{1,2}

Institutional addresses:

¹ Department of Environmental Atmospheric Sciences, Pukyong National University, Daeyeon campus 45, Yongso-ro, Namgu, Busan 608-737 Republic of Korea

² Atmospheric Environmental Research Institute, Daeyeon campus 45, Yongso-ro, Namgu, Busan 608-737 Republic of Korea

* Corresponding author

Abstract

Raindrop size distribution (DSD) characteristics within the complex area of Busan, Korea (35.12°N, 129.10°E) were studied using a Precipitation Occurrence Sensor System (POSS) disdrometer over a four-year period from ~~February–24th~~ February 2001 to ~~December–24~~ December~~th~~ 2004. Also, to find the dominant characteristics of polarized radar parameters which is differential radar reflectivity (Z_{dr}), specific differential phase (K_{dp}) and specific attenuation (A_h), T-matrix scattering simulation was applied in present study. To analyze the climatological DSD characteristics in more detail, the entire period of recorded rainfall was divided into 10 categories not only covering different temporal and spatial scales, but also different rainfall types. When only convective rainfall was considered, mean values of mass weighted mean diameter (D_m) and normalized number concentration (N_w) values for all these categories converged around a maritime cluster, except for rainfall associated with Typhoons. The convective rainfall of a Typhoon showed much smaller D_m and larger N_w compared with the other rainfall categories.

In terms of diurnal DSD variability, we analyzed maritime (continental) precipitation during the daytime (DT) (nighttime, NT), which likely results from sea (land) wind identified through wind direction analysis. These features also appeared in the seasonal diurnal distribution. The DT and NT Probability Density Function (PDF) during the Summer was similar to the PDF of the entire study period. However, the DT and NT PDF during the Winter season displayed an inverse distribution due to seasonal differences in wind direction.

Keyword: DSD, POSS disdrometer, Climatological characteristics, Land and sea wind.

1. Introduction

Raindrop Size Distribution (DSD) is controlled by the microphysical processes of rainfall and therefore it plays an important role in development of the Quantitative Precipitation Estimation (QPE) algorithms based on forward scattering simulations of radar measurements (Seliga and Bringi, 1976). DSD data accurately reflects local rainfall characteristics within an observation area (You et al., 2014). Many DSD models have been developed to characterize spatial-temporal differences in DSDs under various atmospheric conditions (Ulbrich, 1983). Marshall and Palmer (1948) developed an exponential DSD model using DSD data collected by a filter paper technique ($N(D) = 8 \times 10^3 \exp(-410R^{-0.21}D)$ in $m^{-3}mm^{-1}$, D in mm and R in $mm\ h^{-1}$). In subsequent studies, a lognormal distribution was assumed to overcome the problem of exponential DSD mismatching with real data (Mueller, 1966; Levin, 1971; Markowitz, 1976; Feingold and Levin, 1986).

To further investigate natural DSD variations, Ulbrich (1983) developed a gamma DSD that permitted changing the dimension of the intercept parameter (N_0 in $m^{-3}mm^{-1-\mu}$) with $N(D) = N_0 D^\mu \exp(-\Lambda D)$. In addition, to enable the quantitative analysis of different rainfall events, the development of a normalized gamma DSD model that accounted for the independent distribution of DSD from the disdrometer channel interval enabled a better representation of the actual DSD (Willis, 1984; Dou et al., 1999; Testud et al., 2001).

DSDs depend on the rainfall type, geographical and atmospheric conditions and observation time. Also these are closely linked to microphysical characteristics that control rainfall development mechanisms. In the case of stratiform rainfall, raindrops grow by the accretion mechanism because of the relatively long residence time in weak updraft condition, in which almost all water droplets are changed to ice particles. With time, the ice particles grow

sufficiently and fall to the ground. The raindrop size of stratiform rainfall observed at the ground level is larger than that of convective rainfall for a same rainfall intensity due to the resistance of the ice particles to break-up mechanisms. In contrast to stratiform rainfall, convective rainfall raindrops grow by the collision-coalescence mechanism associated with relatively strong vertical wind speeds and short residence time in the cloud. Fully-grown raindrops of maritime precipitation are smaller in diameter than those in stratiform rainfall due to the break-up mechanism in case of same rainfall rate (Mapes and Houze Jr, 1993; Tokay and Short, 1996). Convective rainfall can be classified into two types based on the origin and direction of movement. Rainfall systems occurring over ocean and land are referred to as maritime and continental rainfall, respectively (Göke et al., 2007). Continental rainfall is related to a cold-rain mechanism whereby raindrops grow in the form of ice particles. In contrast, maritime rainfall is related to a warm-rain mechanism whereby raindrops grow by the collision-coalescence mechanism. Therefore, the mass-weighted drop diameter (D_m) of continental rainfall observed on the ground is larger than that of maritime rainfall, and a smaller normalized intercept parameter (N_w) is observed in continental rainfall (Brangi et al., 2003).

Specific heat is a major climatological feature that creates differences between DSDs in maritime and continental regions. These two regions have different thermal capacities and thus different temperature variations are occurred with time. The surface temperature of the ocean changes slowly because of the higher thermal capacity compared with land. While the continental regions which have comparatively lower thermal capacity show greater diurnal temperature variability. Sea winds generally are occurred in from afternoon to early evening when the temperature gradient between the sea and land becomes negative, which is opposite gradient in the daytime (DT). In coastal regions, the land and sea wind effect causes a pronounced difference between the DT and nighttime (NT) DSD characteristics. Also, when

mountains are located near the coast, the difference is intensified by the effect of mountain and valley winds (Qian, 2008).

In the present study, we analyzed a four-year dataset spanning from 2001 to 2004, collected from Busan, Korea (35.12°N, 129.10°E) using a Precipitation Occurrence Sensor System (POSS) disdrometer, to investigate the characteristics of DSDs in Busan, Korea which consist complex mid-latitude region comprising both land and ocean. To quantify the effect of land and sea wind on these characteristics, we also analyzed diurnal variations in DSDs. The remainder of the manuscript is organized as follows. In Section 2 we review the normalized gamma model and explain the DSD quality control method and the classification of rainfall. In Section 3 we report the results of DSD analysis with respect to stratiform/convective and continental/maritime rainfall, and discuss diurnal variations. Finally, a summary of the results and the main conclusions are presented in Section 4.

2. Data and Methods

2.1. Normalized Gamma DSD

DSDs are defined by $N(D) = N_0 \exp(-\Lambda D)$ ($\text{m}^{-3}\text{mm}^{-1}$) and are reflect the microphysical characteristics of rainfall using the number concentration of raindrops ($N(D)$). Also, DSDs are able to calculate the many kind of parameters which show the dominant feature of raindrops. Normalization is used to define the DSD and to solve the non-independence of each DSD parameter (Willis, 1984; Dou et al., 1999; Testud et al., 2001). Furthermore, a normalized gamma DSD enables the quantitative comparison for rainfall cases regardless of time scale and rain rate. Here, we use the DSD model designed by Testud et al. (2001):

111

$$112 \quad N(D) = N_w f(\mu) \left(\frac{D}{D_m}\right)^\mu \exp \left[- (4 + \mu) \frac{D}{D_m} \right]. \quad (1)$$

113

114 where D is the volume equivalent spherical raindrop diameter (mm), and $f(\mu)$ is defined

115 using the DSD model shape parameter (μ) and gamma function (Γ) as follows:

116

$$117 \quad f(\mu) = \frac{6}{4^4} \frac{(\mu+4)^{4+\mu}}{\Gamma(4+\mu)}. \quad (2)$$

118

119 From the value of $N(D)$, the median volume diameter (D_0 in mm) can be obtained as follows:

120

$$121 \quad \int_0^{D_0} D^3 N(D) dD = \frac{1}{2} \int_0^{D_{\max}} D^3 N(D) dD. \quad (3)$$

122

123 Mass-weighted mean diameter (D_m in mm) is calculated as the ratio of the fourth to the third

124 moment of the DSD:

125

$$126 \quad D_m = \frac{\int_0^{D_{\max}} D^4 N(D) dD}{\int_0^{D_{\max}} D^3 N(D) dD}. \quad (4)$$

127

The normalized intercept parameter (N_w in $m^{-3}mm^{-1}$) is calculated as follows:

$$N_w = \frac{4^4}{\pi \rho_w} \left(\frac{LWC}{D_m^4} \right). \quad (5)$$

The shape of the DSD is calculated as the ratio of D_m to the standard deviation (SD) of D_m (σ_m in mm) (Ulbrich and Atlas, 1998; Bringi et al., 2003; Leinonen et al., 2012):

$$\sigma_m = \left[\frac{\int_0^{D_{max}} D^3 (D - D_m)^2 N(D) dD}{\int_0^{D_{max}} D^3 N(D) dD} \right]^{\frac{1}{2}}. \quad (6)$$

In addition, σ_m/D_m is related to μ as follows:

$$\frac{\sigma_m}{D_m} = \frac{1}{(4+\mu)^{1/2}}. \quad (7)$$

Liquid water content (LWC in $g m^{-3}$) can be defined from the estimated DSD:

$$LWC = \frac{\pi}{6} \rho_w \int_0^{D_{max}} D^3 N(D) dD. \quad (8)$$

where ρ_w is the water density (g m^{-3}) and it assumed as $1 \times 10^6 \text{ g m}^{-3}$ for a liquid. Similarly, the rainfall rate (R in mm h^{-1}) can be defined as follows:

$$R = \frac{3.6}{10^3} \frac{\pi}{6} \int_0^{D_{\max}} v(D) D^3 N(D) dD. \quad (9)$$

where the value of factor 3.6×10^3 is the unit conversion which converts the mass flux unit ($\text{mg m}^{-2} \text{s}^{-1}$) to the common unit (mm h^{-1}) for the convenience. $v(D)$ (m s^{-1}) is the terminal velocity for each raindrop size. The relationship between $v(D)$ and D (mm) is given by Atlas et al. (1973) who developed an empirical formula based on the data reported by Gunn and Kinzer (1949):

$$v(D) = 9.65 - 10.3 \exp[-0.6D]. \quad (10)$$

2.2 Quality Control of POSS Data

POSS is used to measure the number of raindrops within the diameter range of 0.34-5.34 mm, using bistatic, continuous wave X-band Doppler radar (10.525 GHz) across 34 channels (Fig. 1; Sheppard and Joe, 2008). To estimate DSDs, Doppler power density spectrum is calculated as follows;

$$S(f) = \int_{D_{\min}}^{D_{\max}} N(D_m) V(D_m, \rho, h, w) \bar{S}(f, D_m, \rho, h, w) dD_m. \quad (11)$$

Where $S(f)$ means Doppler spectrum power density, $V(D_m, \rho, h, w) \bar{S}(f, D_m, \rho, h, w)$ means weighting function of $S(f)$, \bar{S} is the mean of $S(f)$, ρ is density of precipitation distribution, h is the shape of precipitation distribution, w (m s^{-1}) is wind speed and $V(x)$ is sample volume and the symbol 'x' means arbitrary parameters which affect the sampling volume. The Doppler power density spectrum has a resolution of 16Hz and terminal velocity (v_t) has a resolution of 0.24 m s^{-1} . Transmitter and receiver skewed about 20° toward each other, and cross point of signal is located over 34 cm from transmitter-receiver. Transmitter-receiver toward upper side detects $N(D)$ in $V(x)$ (Sheppard, 1990). Also, Sheppard (1990) and Sheppard and Joe (1994) noted some shortcomings as the overestimation of small drops at horizontal wind larger than 6 m s^{-1} . However, in present study, the quality control of POSS for wind effect was not considered because it lies beyond this work. Detailed specifications and measurement ranges and raindrop sizes for each observation channel of the POSS disdrometer are summarized in Table 1.

A POSS disdrometer have been operating in Busan, Korea (35.12°N , 129.10°E), along with other atmospheric instruments, the locations of which are shown in Fig. 2. Estimating raindrop diameter correctly is challenging and care should be taken to ensure reliable data are collected. We performed the following quality controls to optimize the accuracy of the disdrometer estimates. i) Non-liquid type event data (e.g., snow, hail etc.) detected by POSS were excluded by routine observation and surface weather chart provided by Korea Meteorological Administration (KMA). ii) DSD spectra in which drops were not found in at least five

consecutive channels were removed as non-atmospheric. iii) Only data recorded in more than 10 complete channels were considered. iv) To compensate for the reduced capability to detect raindrops smaller than 1 mm when $R > 200 \text{ mm h}^{-1}$ (as recorded by the disdrometer), data for $R > 200 \text{ mm h}^{-1}$ were not included in the analyses, even though the number of samples was only 64 for the entire period. v) To eliminate wind and acoustic noise, data collected when $R < 0.1 \text{ mm h}^{-1}$ are removed (Tokay and Short, 1996).

After performing all quality control procedures, 99,388 spectra were left from original data (166,682) for 1-min temporal resolution. Accumulated rainfall amount from POSS during the entire period was 4261.49 mm. To verify the reliability of the POSS data, they were compared with data collected by a 0.5 mm tipping bucket rain gauge at an automatic weather system (AWS) located ~368 m from the POSS (Fig. 3).

2.3 Radar Parameters

First, the radar reflectivity factor (z in mm^6m^{-3}) and non-polarized radar reflectivity (Z in dBZ) were computed using the DSD data collected by POSS, as follows:

$$z = \int_0^{D_{\max}} D^6 N(D) dD. \quad (12)$$

$$Z = 10 \log_{10}(z). \quad (13)$$

The T-matrix method used in this study is initially proposed by Waterman (1965, 1971) to

calculate electromagnetic scattering by single non-spherical raindrops. The adaptable parameters for this calculation are frequency, temperature, hydrometeor types, raindrop's canting angle and axis ratio (γ) and explained the following sentences. Axis ratios of raindrops differ with atmospheric conditions and rainfall type. To derive the drop shape relation from the drop diameter, we applied the results of numerical simulations and wind tunnel tests employing a forth-polynomial equation, as in many previous studies (Beard and Chuang, 1987; Pruppacher and Beard, 1970; Andsager et al., 1999; Brandes et al., 2002). The axis ratio relation used in the present study is a combination of those from Andsager et al. (1999) and Beard and Chuang (1987) for three raindrop size ranges (Bringi et al., 2003). The raindrop axis ratio relation of Andsager et al. (1999) is applied in the range of $1 < D \text{ (mm)} < 4$, as follows:

$$r = 1.0048 + 0.0057D - 2.628D^2 + 3.682D^3 - 1.677D^4. \quad (14)$$

The drop-shape relation of Beard and Chuang (1987) is applied in the range of $D < 1 \text{ mm}$ and $D > 4 \text{ mm}$, as follows:

$$r = 1.012 + 0.01445D - 0.01028D^2. \quad (15)$$

We assumed SD and the mean canting angle of raindrops as 7° and 0° , respectively. The refractive indices of liquid water at 20°C were used (Ray, 1972). Also, the condition of

frequency for electromagnetic wave of radar is 2.85 GHz (S-band). We calculated dual polarized radar parameters based on these conditions. The parameters of differential reflectivity (Z_{dr} in dB), specific differential phase (K_{dp} in deg km^{-1}), and attenuation (A_h in dB km^{-1}), using DSD data were calculated and analyzed.

2.4. Classification of Rainfall Types and Rainfall Events

Rainfall systems can be classified as stratiform or convective in nature, via analysis of the following microphysical characteristics: i) DSD, using relationships between N_0 and R ($N_0 > 4 \times 10^9 R^{-4.3}$ in $\text{m}^{-3}\text{mm}^{-1}$ is considered as convective rainfall, Tokay and Short, 1996; Testud et al., 2001); ii) Z , where, according to Gamache and Houze (1982), a rainfall system that displays radar reflectivity larger than 38 dBZ is considered to be convective; and iii) R , where average value larger than 0.5 mm per 5 min is considered as convective rainfall (Johnson and Hamilton, 1988). Alternatively, rainfall that has 1-min $R > 5$ (0.5) mm h^{-1} and a SD of $R > (<) 1.5 \text{ mm h}^{-1}$ is considered as convective (stratiform) type (Bringi et al., 2003). The rainfall classification method proposed by Bringi et al. (2003) is applied in the present study. It is necessary to categorize different rainfall systems because their microphysical characteristics show great variation depending on the type of rainfall, as well as the type of rainfall event; e.g., Typhoon, Changma, heavy rainfall and seasonally discrete rainfall. To investigate the temporal variation in DSDs, we analyzed daily and seasonal DSDs. Likewise, to investigate diurnal variability in DSD, DT and NT data were considered by using the sunrise and sunset time in Busan (provided by the Korea Astronomy and Space Science Institute [KASI]). In the middle latitudes, and including Busan, the timings of sunrise and sunset vary due to solar culminating height. The earliest and latest sunrise (sunset) time of the entire period

is 0509 KST (1712 KST) and 0733 KST (1942 KST), respectively. DT (NT) is defined as the period from the latest sunrise (sunset) time to the earliest sunset (sunrise) time for the unity of classification of each time group (Table 2).

To analyze the predominant characteristics of DSDs for Typhoon rainfall, nine Typhoon events were selected from throughout the entire study period which is summarized in Table 2.

This study utilizes KMA rainfall warning regulations to identify heavy rainfall events. The KMA issues a warning if the accumulated rain amount is expected to be >70 mm within a 6-hour period, or >110 mm within a 12-hour period. Rainfall events classified as Changma and Typhoon were not included in the classification 'heavy rainfall'.

Changma is the localized rainfall system or rainy season that is usually present over the Korean Peninsula between mid-June and mid-July which is similar to the Meiyu (China) or Baiu (Japan). The selected dates and periods of each rainfall category are summarized in Table 2.

3. Results

3.1. DSD and Radar Parameters

Figure 4 shows the Probability Density Function (PDF) and Cumulative Distribution Function (CDF) of DSDs and radar parameters with respect to the entire, stratiform and convective rainfall. The PDFs of DSD and radar parameters were calculated using the non-parameterization kernel estimation to identify the dominant distribution of each parameter recorded in Busan. Non-parameterization kernel estimation was also used to identify continuous distributions of DSDs. The PDF of stratiform rainfall is more similar to that of the dataset for the entire analysis period due to the dominant contribution of stratiform rainfall

(about 62.93%) to the overall rainfall than that of convective rainfall. However, the PDF for convective rainfall is significantly different from that of the entire analysis period, and as the convective rainfall contributes only 6.11% of the overall rainfall (Table 3). When $\mu < 0$ the distribution of μ for convective rain has more value of PDF than that for stratiform rain (Fig. 4a). Alternatively, the frequency of μ for stratiform rainfall is higher than that of convective rainfall when $0 < \mu < 5$. The value of μ for convective rainfall is higher than that for stratiform rainfall because the break-up mechanism would be increase the number concentration of small raindrops. The number concentrations of mid-size raindrops increased due to the decrease in the number concentration of relatively large raindrops (Hu and Srivastava, 1995; Sauvageot and Lacaux, 1995). However, we observed a higher frequency of convective rainfall than stratiform rainfall in the negative μ range.

The PDF of D_m displays peak around 1.2 and 1.4 mm for stratiform rainfall and the entire rainfall dataset, respectively. We note that a gentle peak exists around 0.7 mm for both stratiform and convective rainfall datasets (Fig. 4b). These features are similar to the distribution of D_m observed in a high-latitude region at Järvenpää, Finland (Fig. 4 of Leinonen et al., 2012). For D_m values > 1.7 mm, the PDF for convective rainfall is higher than stratiform rainfall. Accordingly, the value of DSD for stratiform rainfall is higher than that of convective rainfall when $D_m < 1.7$ mm. Generally, stratiform rainfall that develops by the cold rain process displays weaker upward winds and less efficient break-up of raindrops. Therefore, in the same rainfall rate, stratiform rainfall tends to produce larger raindrops than convective rainfall that develops by the warm rain process. However, the average D_m values for convective and stratiform rain for the entire period are approximately 1.45 and 1.7 mm,

respectively. In short, D_m is proportional to R regardless of rainfall type. This finding is consistent with the results of Atlas et al. (1999) who found that the D_m of convective rainfall is larger than that of stratiform rainfall on Kapingamarangi Island, Micronesia.

The PDF of $\log_{10}(N_w)$ for the entire rainfall dataset was evenly distributed between 1.5 and 5.5, with a peak at $N_w = 3.3$ (Fig. 4c). The PDF of $\log_{10}(N_w)$ for stratiform rainfall is rarely > 5.5 , while for convective rainfall it is higher at > 5.5 than that of stratiform. There is a similar frequency in the stratiform and convective rainfall at 4.4.

The PDF distributions for $\log_{10}(R)$ and $\log_{10}(LWC)$ are similar each other (Fig. 4d and e). It is inferred that the similar results come from the using of alike moment of DSD as 3.67 and 3 for R and LWC , respectively. The PDF of $\log_{10}(R)$ for the entire rainfall dataset ranged between -0.5 and 2. A peak exists at 0.3 and the PDF rapidly decreases from the peak value as R increases. The PDF for stratiform rainfall has a higher frequency than that of the entire rainfall when $-0.3 < \log_{10}(R) < 0.7$, while the PDF for convective rainfall is denser between 0.4 and 2. Furthermore, the frequency of the PDF for convective rainfall was higher than that of stratiform rainfall in case of $\log_{10}(R) > 0.65$ and the peak value shown as 0.9.

The PDF and CDF for Z , Z_{dr} , K_{dp} and A_h are shown in Fig. 4f-i. The PDF of Z for stratiform rainfall (Fig. 4f) is widely distributed between 10 and 50 dBZ with the peak at approximately 27 dBZ. Conversely, for convective rainfall, the value of PDF lie between 27 and 55 dBZ and the peak frequency value at approximately 41 dBZ. The frequency value of reflectivity is higher for convective rainfall than for stratiform rainfall in the range of $\sim > 35$ dBZ. Furthermore, the shape of the PDF for convective rainfall is similar to that reported for Darwin, Australia (Steiner et al., 1995); however, for stratiform rainfall there are significant differences between Busan and Darwin in terms of the shape of the frequency distribution. The

PDF of Z_{dr} for the entire rainfall primarily exists between 0 and 2.5 dB, and the peaks are at 0.3 and 1.8 dB (Fig. 4g). The distribution of Z_{dr} for convective and stratiform rainfall is concentrated between 0.6 and 1.6 dB, and between 0.3 and 2 dB, respectively. The frequency of Z_{dr} for convective (stratiform) rainfall exists in ranges higher (lower) than stratiform (convective) at 0.9 dB.

The dominant distribution of K_{dp} for the entire dataset and for stratiform rainfall lies between 0 and 0.14 deg km⁻¹, with a peak value of 0.03 deg km⁻¹ and 0.08 deg km⁻¹. However, for convective rainfall the PDF of K_{dp} is evenly exist between 0.01 and 0.15 deg km⁻¹. Furthermore, when $K_{dp} > 0.056$ deg km⁻¹, the frequency of the PDF for convective rainfall is higher than that of stratiform rainfall (Fig. 4h).

The PDF of A_h is similar to that of K_{dp} and is exist between 0 and 0.01 dB km⁻¹. For the case of the entire rainfall dataset and for stratiform rainfall, the PDF of A_h is concentrated between 0 and 2.0×10^{-3} dB km⁻¹ and that of convective rainfall is strongly concentrated between 1.0×10^{-3} and 8.0×10^{-3} dB km⁻¹ (Fig. 4i). Unlike the PDF of A_h for convective rainfall, the PDF for stratiform rainfall shows a strong peak at about 7.0×10^{-4} dB km⁻¹.

3.2. Climatological Characteristics of DSD in Busan

The climatological characteristics of DSDs for 10 rainfall categories are analyzed in this study. Sample size and ratio rainfall for each category are summarized in Table 3. Figure 5a illustrates the distribution of all 1-min stratiform rainfall data, and Fig. 5b shows scatter plots of averaged D_m and $\log_{10}(N_w)$ for all 10 rainfall categories for stratiform rainfall data. Figure 5a displays a remarkable clear boundary in the bottom sector and shows that most of the data lie below the

reference line used by Bringi et al. (2003) to classify convective and stratiform rainfall. The average value of D_m and $\log_{10}(N_w)$ for all rainfall categories, except for heavy rainfall, exist between 1.4 mm and 1.6 mm and between 3.15 and 3.5, respectively (Fig. 5b). These values are relatively small compared with the reference line presented by Bringi et al. (2003). The distribution of 1-min convective rainfall data is displayed in Fig. 6a and the distribution of average values of D_m and N_w for the 10 rainfall categories in the case of convective rainfall in Fig. 6b. The blue and red plus symbols represent maritime and continental rainfall, respectively, as defined by Bringi et al. (2003). The scatter plot of 1-min convective rainfall data shows more in the continental cluster than the maritime cluster; however, the average values for the 10 rainfall categories are all located around the maritime cluster, except for the Typhoon category. By considering the entire average values including Typhoon event (Fig. 6b), we can induce the simple linear equation using D_m and $\log_{10}(N_w)$ as follows:

$$\log_{10}(N_w) = -1.8D_m + 6.9. \quad (16)$$

Even the coefficients in Eq. 16 might be changed slightly with the Typhoon values, this result does not represent in $D_m < 1.2$ mm and $D_m > 1.9$ mm. The D_m (N_w) value for the Typhoon category was considerably smaller (larger) than that of the other categories as well as that of stratiform type of Typhoon. This result does not agree with that reported by Chang et al. (2009), who noted that the D_m of convective rainfall Typhoon showed a large value compared with that associated with stratiform rainfall.

3.3 Diurnal Variation in Raindrop Size Distributions

3.3.1. Diurnal Variations in DSDs

Figure 7a shows a histogram of normalized frequency of 16 wind directions recorded by the AWS, which is the same instrument as that used to collect the data shown in Fig. 3. To establish the existence of a land and sea wind, the difference in wind direction frequencies between DT and NT were analyzed. Figure 7b shows the difference between DT and NT, difference frequency means normalized frequency of wind direction for DT subtract to that of NT for each direction, in terms of the normalized frequency of 16 wind directions. In other word, positive (negative) values indicate that the frequency of wind is more often observed during DT (NT). Also, land (sea) wind defined in present study from 225° (45°) to 45° (225°) according to the geographical condition in Busan. The predominant frequency of wind direction in the DT (NT), between 205° (22.5°) and 22.5° (205.5°), is higher than that in the NT (DT) (Fig. 7b). The observation site where the POSS was installed at western side from the closest coast line, distance is about 611 m, suggesting that the effect of the land and sea wind would have been recorded. To understand the effects of the land and sea wind on DSD characteristics, we analyzed the PDF and 2-hour averaged DSD parameters for DT and NT. Figure 8 illustrates the distributions of μ , D_m , $\log_{10}(N_w)$, $\log_{10}(LWC)$, $\log_{10}(R)$, and Z . There were large variations of μ with time. The μ values varied from 2.41 to 3.17 and the minimum and maximum μ values occurred at 08:00 KST and 12:00 KST, respectively (Fig. 8a). A D_m larger than 1.3 mm dominated from 00:00 KST to 12:00 KST, before decreasing remarkably between 12:00 and 14:00 KST. The minimum and maximum D_m appeared at 14:00 KST and 08:00 KST, respectively (Fig. 8b).

N_w distribution showed inversely to D_m ; however, no inverse relationship was identified between D_m and N_w in case of the time series (Fig. 8c). The maximum and minimum values of N_w were found at 06:00 KST and 22:00 KST.

Variability through time was similar for R , LWC , and Z_h as D_m . There was an increasing trend from 00:00 KST to 08:00 KST followed by a remarkably decreasing trend from 08:00 KST to 14:00 KST (Figs 8d, 11e and 11f). Note that the time of the sharp decline for R between 12:00 KST and 14:00 KST is simultaneous with a D_m decrease. Larger (smaller) drops would contribute to higher R in the morning (afternoon). These variations considerably matched with the diurnal sea wind time series (Fig. 8g). Sea wind is the sum value of normalized wind frequency between 45° and 225° . From 02:00 (14:00) KST to 12:00 (20:00) KST shows smaller (larger) value of sea wind frequency which is opposite to the relatively larger (smaller) parts of each parameter (D_m , R , LWC and Z_h).

The PDF distribution of μ between -2 and 0 is more concentrated for NT than for DT. Furthermore, when $\mu > 0$, DT and NT frequency distributions are similar (Fig. 9a). A larger $N(D)$ of small or large raindrops would be expected in NT than in DT.

The distribution of DT $D_m < 0.7 \text{ mm}$ is wider than that of the NT. However, between 0.7 and 1.5 mm the frequency for NT is higher than that for DT, whereas the distribution in the range $D_m > 1.5 \text{ mm}$ is similar for both DT and NT (Fig. 9b). We note that the smaller peak of D_m around 0.6 mm for the entire rainfall dataset (Fig. 4b) was observed only in DT.

The distribution of $\log_{10}(N_w)$ for DT has higher value of PDF at larger $\log_{10}(N_w)$ than that of NT at $\log_{10}(N_w) > 4$ (Fig. 9c).

Bringi et al. (2003) noted that the maritime climatology displayed larger N_w and smaller D_m values than the continental climatology, based on observed DSDs in the low and middle latitude.

Also, Göke et al. (2007) emphasized that rainfall type can be defined by the origin location and movement direction. In accordance with these previous results, we consider NT rainfall in the Busan region to be more likely caused by a continental convective system.

In the present study, the shape of the PDF of LWC and R for DT and NT are similar which is the same reason with the results of Fig. 4e-f. LWC and R distributions during the DT (NT) are higher (lower) than in the NT (DT) when $\log_{10}(LWC)$ and $\log_{10}(R)$ are larger (smaller) than -1.2 and 0, respectively (Fig. 9d and e). Z has similar pattern with LWC and R during the DT (NT) was higher (lower) than in the NT (DT) in the range below (above) about 27 dBZ (Fig. 9f).

3.3.2. Diurnal Variations of DSDs with respect to Season

Busan experiences distinct atmospheric conditions that are caused by the different frequencies and magnitudes of land and sea winds in response to variable sunrise and sunset times. To identify seasonal variations of DSDs with respect to the effect of the land and sea wind, we analyzed the DT and NT PDF of D_m and N_w in the Summer and Winter. The start and end times of DT (NT) were sorted using the latest sunrise (sunset) and the earliest sunset (sunrise) time for each season (Table 4) which is same method that of entire period classification.

Figure 10a shows a histogram of wind directions in Summer (light grey) and Winter (dark grey). The frequencies of Summer and Winter wind directions are similar to each other. However, in Fig. 10b, the DT and NT distributions of Winter wind direction display opposing frequencies. Note that Winter season shows remarkable frequency of land (sea) wind between 0° (157.5°) and 45° (202.5°) at DT (NT) compared with results of those for Summer season. The accumulated value of normalized wind frequencies at the sea and land wind show different

feature between Summer and Winter season (Table 5).

To identify the variability of DSDs caused by the land and sea wind in Summer and Winter, a 2-hour interval time series of D_m , N_w and R was analyzed. In the Summer, the time series of D_m displays considerably large values between 00:00 KST and 12:00 KST, compared with the period between 14:00 KST and 22:00 KST (Fig. 11a). The mean value of D_m decreases dramatically between 12:00 KST and 14:00 KST. $\log_{10}(N_w)$ has a negative relationship with D_m (Fig. 11b). However, the inverse relation between $\log_{10}(N_w)$ and D_m is not remarkable. $\log_{10}(R)$ tends to increase gradually from 00:00 KST to 08:00 KST and decrease from 08:00 KST to 14:00 KST, which is similar to the pattern that of entire period (Fig. 11c). Kozu et al. (2006) analyzed the diurnal variation in R at Gadanki (South India), Singapore, and Kototabang (West Sumatra) during the Summer monsoon season. All regions displayed maximum R at approximately 16:00 LST, except for Gadanki. Also, Qian (2008), who analyzed the diurnal variability of wind direction and R on Java Island during the Summer season using 30 years (from 1971 to 2000) of NCEP/NCAR reanalyzed data. They found that a land wind occurred from 01:00 LST to 10:00 LST and a sea wind from 13:00 LST to 22:00 LST (Fig. 7 of Qian (2008)). Normalized wind frequency for each direction is similar pattern to the results of Qian (2008) but pattern of R is different with that of Kozu et al. (2006). The diurnal variation of rain rate in the present study from 02:00 (12:00) KST to 10:00 (20:00) KST shows relatively smaller (larger) frequencies of sea wind. It is different pattern with the result of Kozu et al. (2006). However, these patterns matched with the time series of D_m and $\log_{10}(N_w)$. Larger frequency of sea wind direction shows counter-proportional (proportional) relationship to the smaller (larger) frequency of D_m ($\log_{10}(N_w)$).

Variability of D_m time series for Winter is the inverse of the Summer time series (Fig. 11a).

The mean value of D_m steadily increases from 00:00 KST to 16:00 KST and then decreases from 16:00 KST to 22:00 KST. The Winter $\log_{10}(N_w)$ time series displays a clear inverse pattern compared with the D_m variation with time and increases from 1600 KST to 0400 KST and then steadily decreases from 04:00 KST to 16:00 KST (Fig. 11b). The peak of $\log_{10}(N_w)$ occurs at 04:00 KST. However, the time series of $\log_{10}(R)$ for Winter season shows similar pattern with that of Summer unlike to another parameters (Fig. 11c). Based on the diurnal variation of R , the variations of D_m and N_w would be independent to R .

Alike to the D_m and $\log_{10}(N_w)$, normalized wind frequency of wind direction for Winter season shows inverse relationship to that of Summer season (Fig. 11d). The value of frequency generally decreases (increases) from 04:00 (14:00) KST to 14:00 (04:00) KST. Also, it shows symmetry pattern with that of Summer season.

The PDF distribution of Summer D_m displays a relatively large DT frequency compared with NT when $D_m < 1.65 \text{ mm}$, except for the range between 0.6 and 0.9 mm . However, in the range of $D_m > 1.65 \text{ mm}$, the NT PDF displays a larger frequency (Fig. 12a). The PDF of $\log_{10}(N_w)$ for DT (NT) has a larger frequency than the NT (DT) when $\log_{10}(N_w) > (<) 3.3$ but smaller frequency when $\log_{10}(N_w) < (>) 3.3$ (Fig. 12c).

The DT and NT PDFs of D_m and $\log_{10}(N_w)$ during Winter display an inverse distribution to that of Summer. For the PDF of D_m , there is a considerable frequency for NT (DT) when $D_m < (>) 1.6 \text{ mm}$ (Fig. 12b). The PDF of $\log_{10}(N_w)$ of Summer season for NT (DT) is larger than that of the DT when $\log_{10}(N_w) < (>) 3.5$ (Fig. 12d). In the PDF analysis, relatively large (small) D_m and small (large) $\log_{10}(N_w)$ are displayed during the NT (DT) when a land wind (sea wind) occurs.

Bringi et al. (2003) referred that the convective rainfall type is able to be classified as the

continental and maritime-like precipitation using D_m and N_w . As the previous study result, we analyzed the PDF of DSDs for Summer and Winter with respect to convective rainfall type. These feature would be shown more clearly in convective type. The convective rainfall type of PDFs of DT and NT for Summer show similar shape of distribution to that of all rainfall type (Fig. 3a). For the PDF of D_m , there is a more frequency for DT (NT) than NT (DT) when $D_m < (>) 2.0 \text{ mm}$ except for between 0.7 mm and 1.2 mm (Fig. 13a). The PDF of convective rainfall type's $\log_{10}(N_w)$ for DT (NT) has a larger frequency than the NT (DT) when $\log_{10}(N_w) > (<) 3.4$ except for between 4.3 and 5.5 (Fig. 12c). PDF distributions for Winter season show more clear pattern compared with those of the entire rainfall type. The value of PDF for D_m in DT (NT) have considerably larger than NT (DT) when $D_m > (<) 1.9 \text{ mm}$, especially between 2.15 mm and 2.3 mm (Fig. 13b). Also, those for $\log_{10}(N_w)$ in DT (NT) show dramatic values when $\log_{10}(N_w) < (>) 3.6$. Furthermore, PDF values significantly concentrated on between $3 < \log_{10}(N_w) < 3.2$ (Fig. 13d). In short, considering the DSD parameters with wind directions, the maritime (continental)-like precipitation would depend on the sea (land) wind.

4. Summary and Conclusion

Climatological characteristics of DSDs in Busan were analyzed using the DSD data observed by POSS over a four-year period from ~~February–24th~~ February 2001 to ~~December–24th~~ December 2004. Observed DSDs were filtered to remove errors by performing several quality control measures, and an AWS rain gauge installed nearby was used to verify the rainfall amount recorded by the POSS. We analyzed DSD characteristics of convective and stratiform rainfall types, as defined by Bringi et al. (2003). The rainfall dataset was thus divided into

stratiform and convective rainfall and their contributions to the total rainfall were 62.93% and 6.11%, respectively. Also, to find the climatological characteristics of DSD for rainfall case, the entire rainfall data was classified as 10 rainfall categories including the entire period case. According to the study by Bringi et al. (2003), the rainfall in Busan shows maritime climatological DSD characteristics. The mean values of D_m and N_w for stratiform rainfall are relatively small compared with the average line of stratiform rainfall produced by Bringi et al. (2003), except for heavy rainfall events. In case of convective type, mean values of D_m and N_w are converged around the maritime cluster, except for the Typhoon category. The convective rainfall associated with a Typhoon has considerably smaller D_m and larger N_w values compared with the other rainfall categories. This is likely caused by increased raindrop break-up mechanism as a result of strong wind effects. Furthermore, the distributions of mean D_m and N_w values for all rainfall categories associated with convective rainfall displays a linear relationship including the Typhoon category.

The analysis of diurnal variation in DSD yielded the following results: first, the frequency of μ is higher at NT than during the DT in the negative value. The PDF of R is higher at NT than during the DT when $\log_{10}(R) > 0.6$. The value of PDF for D_m during DT is larger than NT smaller than 0.65 mm. For N_w , which tends to be inversely related to D_m , its frequency is higher at NT than DT when $\log_{10}(N_w) > 3.8$. This feature is matched with the time series of normalized frequency of sea wind which shows inverse relationship to D_m . Smaller D_m corresponds to the larger sea wind frequency. In short, maritime (continental) –like precipitation are observed in the DT (NT) more often than in the NT (DT) according to the features of wind. The above-mentioned DSD characteristics are likely due to the land and sea wind caused by differences in specific heat between the land and ocean. These features are also apparent in the seasonal diurnal distribution. The PDF of DT and NT for convective rainfall type during the Summer is

similar to the PDF of the entire period; however, those of the Winter displays the significant inverse distribution compared to the Summer because of obvious seasonal differences in wind direction.

Author contributions

Cheol-Hwan You designed the study. Sung-Ho Suh modified the original study theme and performed the study. Cheol-Hwan You and Sung-Ho Suh performed research, obtained the results and prepared the manuscript along with contributions from all of the co-authors. Dong-In Lee examined the results and checked the manuscript.

Acknowledgement

This work was funded by the Korea Meteorological Industry Promotion Agency under Grant KMIPA 2015-1050. The authors acknowledge providing, weather chart, and AWS data for this work from the Korea Meteorological Administration.— and ~~The authors also acknowledge providing~~ codes for scattering simulation from Professor V. N. Bringi at Colorado State University.

564

565 **References**

566 Andsager, K., Beard, K. V., and Laird, N. F.: Laboratory measurements of axis ratios for large
567 raindrops, *Journal of the Atmospheric Sciences*, 56, 2673-2683, 1999.

568 Atlas, D., Srivastava, R., and Sekhon, R. S.: Doppler radar characteristics of precipitation at
569 vertical incidence, *Reviews of Geophysics*, 11, 1-35, 1973.

570 Atlas, D., Ulbrich, C. W., Marks, F. D., Amitai, E., and Williams, C. R.: Systematic variation
571 of drop size and radar-rainfall relations, *Journal of Geophysical Research: Atmospheres* (1984–
572 2012), 104, 6155-6169, 1999.

573 Beard, K. V., and Chuang, C.: A new model for the equilibrium shape of raindrops, *Journal of*
574 *the Atmospheric sciences*, 44, 1509-1524, 1987.

575 Brandes, E. A., Zhang, G., and Vivekanandan, J.: Experiments in rainfall estimation with a
576 polarimetric radar in a subtropical environment, *Journal of Applied Meteorology*, 41, 674-685,
577 2002.

578 Bringi, V., Chandrasekar, V., Hubbert, J., Gorgucci, E., Randeu, W., and Schoenhuber, M.:
579 Raindrop size distribution in different climatic regimes from disdrometer and dual-polarized
580 radar analysis, *Journal of the atmospheric sciences*, 60, 354-365, 2003.

581 Chang, W.-Y., Wang, T.-C. C., and Lin, P.-L.: Characteristics of the raindrop size distribution
582 and drop shape relation in Typhoon systems in the western Pacific from the 2D video
583 disdrometer and NCU C-band polarimetric radar, *Journal of Atmospheric and Oceanic*
584 *Technology*, 26, 1973-1993, 2009.

585 Dou, X., Testud, J., Amayenc, P., and Black, R.: The parameterization of rain for a weather
586 radar, *Comptes Rendus de l'Académie des Sciences-Series IIA-Earth and Planetary Science*,

328, 577-582, 1999.

Feingold, G., and Levin, Z.: The lognormal fit to raindrop spectra from frontal convective clouds in Israel, *Journal of Climate and Applied Meteorology*, 25, 1346-1363, 1986.

~~Göke, S., Ochs III, H. T., and Rauber, R. M.: Radar analysis of precipitation initiation in maritime versus continental clouds near the Florida coast: Inferences concerning the role of CCN and giant nuclei, *Journal of the Atmospheric Sciences*, 64, 3695-3707, 2007.~~

Gamache, J. F., and Houze Jr, R. A.: Mesoscale air motions associated with a tropical squall line, *Monthly Weather Review*, 110, 118-135, 1982.

Göke, S., Ochs III, H. T., and Rauber, R. M.: Radar analysis of precipitation initiation in maritime versus continental clouds near the Florida coast: Inferences concerning the role of CCN and giant nuclei, *Journal of the Atmospheric Sciences*, 64, 3695-3707, 2007.

Gunn, R., and Kinzer, G. D.: The terminal velocity of fall for water droplets in stagnant air, *Journal of Meteorology*, 6, 243-248, 1949.

Hu, Z., and Srivastava, R.: Evolution of raindrop size distribution by coalescence, breakup, and evaporation: Theory and observations, *Journal of the atmospheric sciences*, 52, 1761-1783, 1995.

Johnson, R. H., and Hamilton, P. J.: The relationship of surface pressure features to the precipitation and airflow structure of an intense midlatitude squall line, *Monthly Weather Review*, 116, 1444-1473, 1988.

Kozu, T., Reddy, K. K., Mori, S., Thurai, M., Ong, J. T., Rao, D. N., and Shimomai, T.: Seasonal and diurnal variations of raindrop size distribution in Asian monsoon region, *JOURNAL-METEOROLOGICAL SOCIETY OF JAPAN SERIES 2*, 84, 195, 2006.

Leinonen, J., Moisseev, D., Leskinen, M., and Petersen, W. A.: A climatology of disdrometer

611 measurements of rainfall in Finland over five years with implications for global radar
 612 observations, *Journal of Applied Meteorology and Climatology*, 51, 392-404, 2012.

613 Levin, Z.: Charge separation by splashing of naturally falling raindrops, *Journal of the*
 614 *Atmospheric Sciences*, 28, 543-548, 1971.

615 Mapes, B. E., and Houze Jr, R. A.: Cloud clusters and superclusters over the oceanic warm
 616 pool, *Monthly Weather Review*, 121, 1398-1416, 1993.

617 Markowitz, A. H.: Raindrop size distribution expressions, *Journal of Applied Meteorology*, 15,
 618 1029-1031, 1976.

619 Marshall, J. S., and Palmer, W. M. K.: The distribution of raindrops with size, *Journal of*
 620 *meteorology*, 5, 165-166, 1948.

621 Mueller, E. A.: Radar cross sections from drop size spectra, University of Illinois at Urbana-
 622 Champaign, 1966.

623 Pruppacher, H., and Beard, K.: A wind tunnel investigation of the internal circulation and shape
 624 of water drops falling at terminal velocity in air, *Quarterly Journal of the Royal Meteorological*
 625 *Society*, 96, 247-256, 1970.

626 Qian, J.-H.: Why precipitation is mostly concentrated over islands in the Maritime Continent,
 627 *Journal of the Atmospheric Sciences*, 65, 1428-1441, 2008.

628 Ray, P. S.: Broadband complex refractive indices of ice and water, *Applied Optics*, 11, 1836-
 629 1844, 1972.

630 Sauvageot, H., and Lacaux, J.-P.: The shape of averaged drop size distributions, *Journal of the*
 631 *Atmospheric Sciences*, 52, 1070-1083, 1995.

632 Seliga, T., and Bringi, V.: Potential use of radar differential reflectivity measurements at
 633 orthogonal polarizations for measuring precipitation, *Journal of Applied Meteorology*, 15, 69-
 634 76, 1976.

635 Sheppard, B. E.: Measurement of raindrop size distributions using a small Doppler radar. J.
636 Atmos. Oceanic Technol., 7, 255–268, 1990.

637 Sheppard, B. E. and P. I. Joe: Comparison of raindrop size distribution measurements by a
638 Joss–Waldvogel disdrometer, a PMS 2DG spectrometer and a POSS Doppler radar. J. Atmos.
639 Oceanic Technol., 11, 874–887, 1994.

640 Sheppard, B., and Joe, P.: Performance of the precipitation occurrence sensor system as a
641 precipitation gauge, Journal of atmospheric and Oceanic technology, 25, 196-212, 2008.

642 Steiner, M., Houze Jr, R. A., and Yuter, S. E.: Climatological characterization of three-
643 dimensional storm structure from operational radar and rain gauge data, Journal of Applied
644 Meteorology, 34, 1978-2007, 1995.

645 Testud, J., Oury, S., Black, R. A., Amayenc, P., and Dou, X.: The concept of “normalized”
646 distribution to describe raindrop spectra: A tool for cloud physics and cloud remote sensing,
647 Journal of Applied Meteorology, 40, 1118-1140, 2001.

648 Tokay, A., and Short, D. A.: Evidence from tropical raindrop spectra of the origin of rain from
649 stratiform versus convective clouds, Journal of applied meteorology, 35, 355-371, 1996.

650 Ulbrich, C. W.: Natural variations in the analytical form of the raindrop size distribution,
651 Journal of Climate and Applied Meteorology, 22, 1764-1775, 1983.

652 Ulbrich, C. W., and Atlas, D.: Rainfall microphysics and radar properties: Analysis methods
653 for drop size spectra, Journal of Applied Meteorology, 37, 912-923, 1998.

654 Waterman, P.C.: Matrix formulation of electromagnetic scattering. Proc. IEEE, 53, 805-812,
655 1965.

656 Waterman, P.C.: Symmetry, unitarity, and geometry in electromagnetic scattering, Physical
657 review D, 3, 825, 1971.

658 Willis, P. T.: Functional fits to some observed drop size distributions and parameterization of

rain, Journal of the Atmospheric Sciences, 41, 1648-1661, 1984.

You, C.-H., Lee, D.-I., and Kang, M.-Y.: Rainfall estimation using specific differential phase for the first operational polarimetric radar in Korea, Advances in Meteorology, 2014-, [10 pages](#), 2014.

Tables

Tables 1. Specification of POSS disdrometer.

Specifications	Detail
Manufacturer	ANDREW CANADA INC
Module	PROCESSOR
Model number	POSS-F01
Nominal power	100 mW
Bandwidth	Single frequency
Emission	43 mW
Pointing direction	20 ° (to the vertical side)
Antenna	Rectangular pyramidal horns
Range of sample area	< 2 m
Wavelength	10.525 GHz \pm 15 GHz
Physical dimension	277×200×200 cm ³
Net weights	Approximately 110 kg

672

673

674

675

676 **Table 2.** Designated date with respect to the source of rainfall.

<u>Rainfall Category</u>	<u>Period</u>			
	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>
<u>Typhoon</u>	-	<u>5-6 July, 31 August</u>	<u>29 May, 19 June, 7 August, 11-12 September</u>	<u>20 June, 19 August, 6 September</u>
<u>Changma</u>	<u>18-19 June, 23-26 June, 29-30 June, 1 July, 5-6 July, 11-14 July</u>	<u>23-25 June, 30 June, 1-2 July</u>	<u>12-14 June, 23 June, 27 June, 30 June, 1 July, 3-15 July</u>	<u>11-14 July</u>
<u>Heavy rainfall</u>	<u>15 April 2002, 20:13 KST to 16 April 2002, 06:29 KST</u>			
<u>Seasonal</u>	<u>Spring</u> <u>March to May</u>	<u>Summer</u> <u>June to August</u>	<u>Autumn</u> <u>September to November</u>	<u>Winter</u> <u>December to February</u>
<u>Diurnal</u>	<u>DT (KST)</u>		<u>NT (KST)</u>	
	<u>07:33-17:12</u>		<u>19:42-05:09</u>	

677 **Table 2.** Designated date with respect to the source of rainfall.

<u>Rainfall Category</u>	<u>Period</u>			
	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>
<u>Typhoon</u>	-	<u>7.5-7.6, 8.31</u>	<u>5.29, 6.19, 8.7, 9.11-12</u>	<u>6.20, 8.19, 9.6</u>

Changma	6.18-6.19, 6.23-6.26, 6.29-6.30, 7.1, 7.5-7.6 7.11-7.14	6.23-6.25, 6.30, 7.1-7.2	6.12-6.14, 6.23, 6.27 6.30, 7.1, 7.3- 7.15	7.11-7.13, 7.14
Heavy-rainfall	02.04.15. 20:13 to 02.04.16 06:29			
Seasonal	Spring	Summer	Autumn	Winter
	Mar. to May	Jun. to Aug.	Sep. to Nov.	Dec. to Feb.
Diurnal	DT (KST)		NT (KST)	
	0733-1712		1942-0509	

Table 3. Rainfall rate for each rainfall category and the number of sample size for 1-min data.

Rainfall Category	Total precipitation	Stratiform precipitation (%)	Convective precipitation (%)
Typhoon	5095	3118 (61.19)	652 (12.79)

Changma	18526	11099 (59.91)	1611 (8.69)
Heavy rainfall	359	153 (42.61)	150 (41.78)
Spring	30703	20370 (66.34)	1478 (4.81)
Summer	37187	22566 (60.68)	3409 (9.16)
Autumn	19809	12033 (60.74)	850 (4.29)
Winter	11689	7582 (64.86)	339 (2.90)
Daytime	41328	26373 (63.81)	2539 (6.14)
Nighttime	37455	23063 (84.00)	2242 (5.89)
Entire	99388	62551 (62.93)	6076 (6.11)

688

689

690

691

692 **Table 4.** DT and NT (KST) in Summer and Winter season.

Rainfall Category	Period	Beginning time (KST)	Finishing time (KST)
Summer	DT	05:33	19:27
	NT	19:42	05:09
Winter	DT	07:33	17:12

	NT	18:19	06:54
--	----	-------	-------

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707 **Table 5.** Sum of the normalized wind direction frequencies between Summer and Winter.

Sum of the normalized wind direction frequencies				
Season	Summer		Winter	
Type	Sea wind	Land wind	Sea wind	Land wind
Frequency	0.4139	0.5861	0.3137	0.6863
Difference of the normalized wind direction frequency between DT and NT (DT-NT)				

Season	Summer		Winter	
Type	Sea wind	Land wind	Sea wind	Land wind
Frequency	0.0731	-0.0731	-0.0697	0.0697

708

709

710

711

712

713

714

715

716

717

718

719

720

721 **Figures**

722



723

724 **Figure 1.**

725 Photograph of the POSS instrument used in this research.

726

727




Figure 2.

Locations of the POSS and the AWS rain gauge installed in Busan, Korea.




Figure 3.

Comparison of the recorded rainfall amounts between the POSS and AWS instrument.




Figure 4.

PDF and CDF for (a) μ , (b) D_m , (c) $\log_{10}(N_w)$, (d) $\log_{10}(R)$, (e) $\log_{10}(LWC)$, (f) Z , (g) Z_{dr} , (h) K_{dp} , and (i) A_h for the entire rainfall dataset (solid black line), stratiform rainfall (solid green line), and convective rainfall (solid blue line). The solid red line represents the CDF for entire rainfall dataset. The solid vertical line represents the mean value of each type.




Figure 5.

(a) Scatter plot of 1-min D_m and N_w for the 10 rainfall categories with respect to stratiform rainfall data. The broken grey line represents the average line as defined by Bringi et al. (2003).

(b) Scatter plot of mean D_m and $\log_{10}(N_w)$ values of the 10 rainfall categories with respect to stratiform rainfall and these mean values for each rainfall type are shown as circle symbols. The vertical line represents $\pm 1\sigma$ of $\log_{10}(N_w)$ for each category.




Figure 6.

(a) As in Figure 5(a), but for convective rainfall. The blue and red plus symbols represent maritime and continental rainfall, respectively, as defined by Bringi et al. (2003). (b) As in Figure 5(b), but for convective rainfall. The broken red line represents the mathematical expression described in Eq. (16).




Figure 7.

(a) Histogram of normalized frequency of 16 wind directions for the entire study period. (b) Difference values of wind direction frequencies between DT and NT (DT - NT).




Figure 8.

Two hour interval time series of (a) μ , (b) D_m , (c) $\log_{10}(N_w)$, (d) $\log_{10}(R)$, (e) $\log_{10}(LWC)$, (f) Z_h and (g) normalized frequency of wind direction for sea wind (45° to 225°) with quantiles for the total period. Solid lines are quantiles for each time.




Figure 9.

PDF and CDF curves for (a) μ , (b) D_m , (c) $\log_{10}(N_w)$, (d) $\log_{10}(R)$, (e) $\log_{10}(LWC)$, and (f) Z for DT and NT according to the entire period. The solid red and blue lines represent the PDF for DT and NT, respectively. The broken light red and blue lines represent the CDF for DT and NT, respectively.




Figure 10.

Histogram of normalized frequency for 16 wind directions in (a) the entire period and (b) differences of normalized frequency with respect to wind directions between DT and NT (DT – NT) for Summer (light grey) and Winter (dark grey) season.




Figure 11.

Two hour interval time series and quartiles of (a) D_m , (b) $\log_{10}(N_w)$ (c) $\log_{10}(R)$ and (d) normalized frequency of wind direction for sea wind (45° to 225°) for the Summer (red) and Winter (blue) season. Solid (broken) lines are quartiles of Summer (Winter) for each time, respectively.




Figure 12.

PDF and CDF of (a) D_m ((b) D_m) and (c) N_w ((d) N_w) for the entire rainfall type in the Summer (Winter) season. Red and blue solid lines represent the PDF of DT and NT, respectively. The light red and blue broken lines represent the CDF for each season




Figure 13.

As in Figure 12, but for convective rainfall type.