

# How to predict hydrological effects of local land use change: how the vegetation parameterisation for short rotation coppices influences model results

F. Richter<sup>1</sup>, C. Döring<sup>1</sup>, M. Jansen<sup>1</sup>, O. Panferov<sup>2,3</sup>, U. Spank<sup>4</sup> and C. Bernhofer<sup>4</sup>

[1]{Department of Soil Science of Temperate Ecosystems, Georg-August-Universität Göttingen, Büsgenweg 2, D-37077 Göttingen, Germany}

[2]{Department of Bioclimatology, Georg-August-Universität Göttingen, Büsgenweg 2, D-37077 Göttingen, Germany}

[3]{Institute of Climatology and Climate Protection, University of Applied Sciences, Bingen am Rhein, Berlinstr. 109, D-55411 Bingen am Rhein, Germany}

[4]{Institute of Hydrology and Meteorology, Technische Universität Dresden, Piennner Str. 23, D-01737 Tharandt, Germany}

Correspondence to: F. Richter (falk.richter@forst.uni-goettingen.de)

## Abstract

Among the different bioenergy sources short rotation coppices (SRC) with poplar and willow trees are one of the mostly promising options in Europe. SRC not only provide woody biomass, but often additional ecosystem services. However, one known shortcoming is the potentially lower groundwater recharge, caused by the potentially higher evapotranspiration compared to annual crops. The complex feedback between vegetation cover and water cycle can be only correctly assessed by application of well parameterized and calibrated numerical models. The present study implements the hydrological model system WaSim for the assessment of water balance. The special focus is the analysis of simulation uncertainties caused by the use of guidelines or transferred parameter sets from scientific literature compared to ‘actual’ parameterisations derived from local measurements of leaf area index (LAI), stomatal resistance (Rsc) and date of leaf unfolding (LU). It is clearly shown that uncertainties in parameterisation of vegetation lead to implausible model results. The present

study shows that LAI, Rsc as well as LU are the most sensitive plant physiological parameters when investigating the effects of an enhanced cultivation of SRC on water budget or on groundwater recharge. Mostly sensitive is the beginning of the growing season, i.e. LU. When this estimation is wrong, the accuracy of LAI and Rsc description plays a minor role. The analyses done here illustrate that the use of locally measured vegetation parameters like maximal LAI and meteorological variables like air temperature, to estimate LU, produce better results than literature data or data from remote network stations. However, the direct implementation of locally measured data is not always advisable or possible. In case of Rsc the adjustment of local measurements shows the best model evaluation. Especially if local investigations are in focus local measurements of model sensitive parameters like LAI, Rsc and LU are valuable information. The derivation of these model parameters based on local measurements show the best model fit. Additionally the adjusted seasonal course of LAI and Rsc is less sensitive to different estimates for LU. The different parametrisations, as they are all eligible either from local measurements or from scientific literature, can result in modelled ground water recharge to be present or completely absent in certain years under poplar SRC.

## **1 Introduction**

In the scope of climate change mitigation and reduction of greenhouse gases (GHGs) emissions bioenergy is one of the possible alternatives for fossil fuels. Among the different bioenergy sources short rotation coppices (SRC) with mainly poplar and willow trees are one of the mostly promising options in Europe (Djomo et al., 2011). SRC not only provide woody biomass, but also additional ecosystem services. Seepage water quality is enhanced, due to lower fertilizer requirements and higher nutrient use efficiency (Aronsson et al., 2000; Schmidt-Walter and Lamersdorf, 2012). Compared to conventional annual crops SRC sequester more carbon and emit less N<sub>2</sub>O (Don et al., 2012), which is one of the most important GHGs. As structural landscape elements in rural areas SRC might also contribute positively to biodiversity (Baum et al., 2012).

However, SRC are not without disadvantages. The most quantitatively assessable disadvantage is the potentially lower groundwater recharge (GWR) being caused by higher evapotranspiration of poplar and willow plantations in comparison to annual crops (Lasch et al., 2010; Schmidt-Walter and Lamersdorf, 2012). An assessment using hydrological models can help to minimize negative and maximize positive ecological effects due to the land use change from arable land to SRC at regional and local scales, e.g. to regional climate and/or to

1 adjacent ecosystems. To quantify effects of this land use change and to provide an adequate  
2 assessment, suitable hydrological models are required, correctly reproducing hydrological  
3 feedback effects of vegetation and land use management. However, in the same way as a  
4 proper model approaches are required, carefully executed parameterization of land use and  
5 vegetation is mandatorily needed to obtain reliable simulation results. The aim of present  
6 study is to assess the effect of parameter uncertainties of the land use type poplar SRC on  
7 modelling results.

8 The planting of SRC causes the occurrence of new factors and complex factor interactions  
9 influencing site water fluxes. One factor is the perennial vegetation cover with higher leaf  
10 area index (LAI,  $\text{m}^2 \text{m}^{-2}$ ) combined with a longer growing season compared to annual crops  
11 (Petzold et al., 2010).

12 The LAI directly affects canopy interception due to an almost linear correlation between LAI  
13 and canopy storage (Rutter et al., 1971). Outside the growing season LAI, more precisely plant  
14 area index, additionally provide canopy storage by woody biomass, i.e., stems and branches.  
15 Furthermore, the transpiration is positively correlated with the LAI, i.e., higher LAI causes  
16 higher transpiration rates. However, LAI is negatively correlated with soil evaporation as  
17 higher LAI means more shadowing and therefore less solar radiation input, and in  
18 consequence less evaporation below vegetation cover. Other important parameters controlling  
19 the water balance are the stomatal resistance ( $R_{sc}$ ), rooting depth and roots distribution. The  
20 structural and biophysical parameters do not remain stable during the year, but have a  
21 seasonal or even annual course. The most intensive changes occur during the growing period.  
22 Thus, the beginning and length of the growing season should be known for adequate  
23 description of seasonal dynamic of vegetation parameters.

24 The smaller the scale of interest the more detailed time-dependent parameterisation of land  
25 use and vegetation is necessary to capture the spatial and temporal variability of effects. There  
26 are two possibilities to obtain the required information: the first one, the labour- and time-  
27 consuming way, is to measure the parameters like LAI or  $R_{sc}$  directly at the investigated site.  
28 The other possibility is to apply parameters from scientific literature. This information is quite  
29 rare for SRC, due to the fact that this land use scarcely came into focus of investigation as a  
30 part of the renewable energy discussion during the last years (Surendran Nair et al., 2012).  
31 Very often not the annual course, but only one value is given in literature, e.g. the maximum  
32 value for LAI, or the minimum value for  $R_{sc}$ , describing together the maximum transpiration.

1 In such cases the annual course for these parameters has to be estimated for hydrological  
2 modelling. Hence, the question on transferability of published results obtained in a certain  
3 area and a specific year to other regions and years has to be solved in each study separately.  
4 However, it is well known that neither literature values nor direct measurements provide the  
5 true values of model parameters, but more or less representative approximations, because of  
6 spatial and temporal heterogeneity of vegetation stands including SRC.

7 The overarching aim of our research was the evaluation of land use change effects. However,  
8 this study does not focus on land use change effects in any way but rather on the evaluation of  
9 a suitable tool. In this study we used the results of our own measurements of LAI, Rsc and the  
10 estimation of leaf unfolding date (LU), determining the beginning of growing season, for a  
11 poplar SRC to parameterise the hydrological model system WaSim (Schulla and Jasper,  
12 2013). The aim of the present study is to assess the effect of parameterisation uncertainties of  
13 the land use type poplar SRC on modelling results. The hypothesis, that the parameterization  
14 of LAI, Rsc and LU based on measurements shows a better model fit than the use of values  
15 reported in literature in combination with approximation about the annual course should be  
16 proofed. In our study the following objectives should be met: 1) to quantify the WaSim  
17 response (sensitivity) to variations of following parameters: LAI, Rsc and LU, caused by  
18 different measurement methods and modelling approaches; 2) to estimate the most sensitive  
19 parameter and 3) to evaluate quantitatively whether it is advisable to implement locally point-  
20 measured values of sensitive parameters directly. We used GWR and plant available water as  
21 indicators.

## 22 **2 Material and methods**

### 23 **2.1 Study site**

24 The study site Reiffenhausen (51.67 °N, 10.65°E, 325 m a.s.l.) is located south of Göttingen,  
25 central Germany. According to the meteorological data provided by the German Weather  
26 Service (Deutscher Wetterdienst, DWD), for the station Göttingen (DWD Station-ID: 01691),  
27 nearest to the study site, the climate is characterized by an average temperature of 9.1 °C ( $\pm$   
28 0.7 °C) and a mean annual precipitation sum of 635 mm ( $\pm$  122 mm) for the period 1971-  
29 2010. The site Reiffenhausen was established as part of the interdisciplinary investigations of  
30 SRC by the joint integrated project BEST ("Bioenergie-Regionen stärken" - Boosting  
31 Bioenergy Regions), which ran from 2010 until 2014 and was funded by the German Ministry

of Education and Research (BMBF). The aim of BEST was to develop regionally appropriate concepts and innovative solutions for the production of biomass, with focus on SRC, and to evaluate ecological and economic impacts.

The soil in Reiffenhausen is characterized by a sedimentary deposits of Middle and Upper Buntsandstein, like sandstone, siltstone and clay stone. The main soil types present at the field level are stagnic cambisol and haplic stagnosol with a soil quality (Ackerzahl) of approximately 45 points. The maximum available points are 100 for very good agriculture fields (Blume et al., 2010). The soil texture is dominated by loamy sand or silty clay.

The plantation Reiffenhausen was established at a former arable field in March 2011 with the poplar clone *Populus nigra* x *Populus maximowiczii*, hereafter Max1. The poplar SRC were planted with 0.2 m long cuttings on 0.4 ha in a double row system with alternating inter-row distances of 0.75 m and 1.50 m, and a spacing of 1.0 m within the rows, yielding an overall planting density of 8800 cuttings per hectare. A detailed site description, including soil chemistry and biomass information is given by Hartmann et al. (2014). In 2011 the weather conditions at Reiffenhausen were unfavourably dry for the initial growth after planting. During the first months in 2011, from February to May, the precipitation sum was unusually low: only 42 % of the long term (1971-2010) precipitation (78 mm of 188 mm for the long term mean value of the same period of the year). This led to dry soil conditions, especially in the upper soil, and resulting in a survival rate of only 63 % for the mono-stem cycle of the poplar SRC. In 2013, when the most measurements and investigations took place the poplar SRC reached a height of 5-6 m, indicating that the unfavourable initial conditions were somewhat improved. All parameters influencing the hydrology in the model are in the order of a fully developed SRC, although it is in mono-stem cycle. The development of the rooting system was eventually enforced by the dry conditions during the initial phase (Broeckx et al., 2013). Rooting depth was more than 1 m in the second year after planting, exploiting the main part of soil layer above the bedrock (Kalberlah, 2013). The LAI reached 5 in 2012 and 6-7 in 2013, which is typical for a fully grown poplar SRC (Schmidt-Walter et al., 2014).

Observations for LU are also used from the site Großfahner, which is also part of the BEST-research project. Großfahner is located near Erfurt, Thuringia, Germany and was also established in 2011 with the poplar clone Max1 (Hartmann et al., 2014).

## 2.2 Measurements

### 2.2.1 Meteorological and local soil measurements

In Reiffenhausen, the micrometeorological measurements were carried out in the centre of the SRC stand; the instruments were installed above the vegetation on a 10 m mast. The air temperature and humidity were measured using Hmp45C (Campbell Sci.; Loughborough, UK); wind speed and wind direction (wind sensor compact and wind direction sensor compact, both ThiesClima; Göttingen, Germany), atmospheric pressure (pressure sensor, Theodor Friedrichs & Co.; Schenefeld, Germany) and solar radiation (CMP3, Kipp & Zonen, Delft, The Netherlands) were measured continuously with 1 Hz frequency and averaged over 15 minutes. 5 minutes precipitation sums were obtained using an ombrometer (Precipitation Transmitter, ThiesClima; Göttingen, Germany). The values were averaged and stored by a CR1000 data logger (Campbell Sci.; Loughborough, UK). Additionally to the meteorological measurements in the centre of the poplar SRC, a reference station was similarly equipped and installed approximately 500 m to the north from the stand in the open place (short-grass meadow) to measure the climate variables unaffected by the poplar SRC.

The soil moisture was measured continuously every 15 minutes using tensiometers and soil water content probes in 20, 60 and 120 cm depth, by six (tensiometers) and three (probes) sensors in every depth. Tensiometers were constructed in the Department of Soil Science for the study using the PCFA6D pressure sensor (Honeywell; Morristown, NJ, USA) and a P-80 ceramic (CeramTec AG; Marktreidwitz, Germany). Volumetric soil water content and soil temperature were measured using SM-300 probes (Delta-T Devices Ltd; Cambridge, UK). Additionally, descriptions of soil horizons and soil texture were assessed using a soil pit near the SRC.

Additionally to the data of Reiffenhausen, meteorological and phenological data are used from region of Tharandter Wald (Tharandt Forest) being located in the federal state Saxony (Germany), 15 km southwest of city of Dresden. As, climate characteristic of this region is comparable with Reiffenhausen, a proper set of comparison data are provided. Detailed information about measurement programs of Tharandter Wald can be found, i.a. in Bernhofer (2002) and Spank et al. (2013). In frame of this study, phenological observation data from the International phenological garden Tharandt-Hartha (IPG) and meteorological measurement data (air temperature, air humidity and precipitation) from climate stations Grillenburg and

Wildacker have special importance. Grillenburg and Wildacker are the nearest meteorological long-term measurements sites from IPG and are situated approx. 3 km away. Both stations provide meteorological and climatological information since 1958. The station Grillenburg represents a standard climate station fulfilling all guidelines and standards of World Meteorological Organisation (WMO) for large-scale representativeness of climatological observations. However, measurements on this site sometimes does not represent micro-scale climatic characteristic of the region, particularly related to daily minimum and maximum of air temperature. In contrary, climate station Wildacker, being not fulfil WMO standards of fetch and horizon heightening, better represent local climatic situation.

### 2.2.2 Leaf area index

For the present study we use the definition of leaf area index by Watson (1947) cited in Breda (2003) as the total one-sided area of leaf tissue per unit ground surface area with the dimension of  $\text{m}^2 \text{ m}^{-2}$  (or dimensionless). There are numerous ground-based as well as remote sensing-based techniques to estimate LAI. An extensive overview of ground-based methods is given by Breda (2003). Direct methods: allometric, litter collection and harvesting are based on statistically significant sampling of phytoelements and phytoelement dimensions. Among them only the harvesting can provide the information on the seasonal dynamic of LAI for the whole season or year. The obvious disadvantages of harvesting as destructive method, however, are that it is very time- and labour-consuming, that the canopy is irreversibly damaged and further statistically representative LAI-measurements for seasonal dynamics are affected.

Indirect ground-based methods are non-destructive and based on the inversion of the Beer-Lambert law, i.e. on measurements of the extinction of short-wave solar radiation by the canopy. The extinction is related to the vegetation structure parameters including LAI (Eq. 1).

$$\text{LAI} = \frac{\ln(I_0/I)}{k} \text{ with } k = \frac{G(\theta, \alpha)}{\cos \theta} \quad (1)$$

where LAI is the leaf area index for the vegetation layer,  $I_0$  is the radiation intensity incident to the vegetation layer,  $I$  – radiation intensity at the lower bound of vegetation layer and  $k$  is the extinction coefficient (Breda, 2003). The function  $G$  is the projection of unit foliage area on the plane normal to the direction  $\theta$ ,  $\theta$  – zenith angle and  $\alpha$  is the leaf angle distribution. It should be also noted that indirect methods estimate not LAI but Plant Area Index (PAI) as the light attenuation is caused not only by leaves but by branches and tree stems as well. To

1 derive LAI either the correction factors are applied which subtract the share of woody  
2 material from PAI, or the assumption is made (especially for dense canopies) that the  
3 attenuation is caused for the most part by leaves. The underlying assumptions, e.g. on stand  
4 homogeneity and small black opaque phytoelements that have to be considered to ensure the  
5 applicability of indirect methods, as well as advantages and disadvantages of various methods  
6 are presented, e.g. in LAI-2000 Manual (LI-COR INC, 1992), in Breda (2003) and  
7 (Jonckheere et al., 2004).

8 For the present study the data of one direct and two indirect methods for the estimation of  
9 LAI of the poplar SRC in Reiffenhausen were used. For the indirect method we used two  
10 different types of instruments. First - two LI-191 SA Line Quantum Sensors (LI-COR Inc.,  
11 USA) were used to measure incident ( $I_{0,PAR}$ ) and within-stand photosynthetic active radiation  
12 ( $I_{PAR}$ ) to calculate the LAI using Eq. (1). The  $k = 0.5$  for mixed broadleaved species was  
13 accepted in our study (Breda, 2003). Second - two plant canopy analysers LAI-2000 (LI-COR  
14 Inc., USA) were implemented in Two-Sensor mode (LI-COR INC, 1992) to obtain LAI and  $k$ .  
15 Measurements were performed weekly whenever possible from May till November 2013  
16 under homogenous illumination, i.e. at days with overcast conditions or during morning or  
17 evening hours. Sensor pairs were cross-validated at the beginning of each measurement day.

18 In the homogeneous poplar SRC ten evenly distributed plots were selected. To account for the  
19 double row planting of the SRC 3 m x 3 m square grids with 1 m distance between grid points  
20 were marked at every plot so that 16 grid points per plot were obtained for measurements. At  
21 each grid point two measurements were performed with instrument oriented along and  
22 perpendicularly to SRC rows. Thus 32 measurements were performed at each of ten plots  
23 during every measurement day. The LAI-2000 was used in two-instrument mode with 25 %  
24 view restriction caps to eliminate the influence of observer. The measurements with line  
25 quantum sensors LI-191SA were also carried out in two-instrument mode; the measurement  
26 design was absolutely identical to LAI-2000.

27 To obtain the reference values for leaf area the direct destructive sampling – harvesting were  
28 carried out. All phytoelements within the square column of 1 m<sup>2</sup> surface area were collected  
29 and measured with leaf area meter (LI-3100; LI-COR Inc., USA). The sampling was carried  
30 at 26 August 2013 at three plots within the investigated stand.



### 2.2.3 Stomatal resistance

The dominant factor controlling both the water loss from plant leaves and the uptake of CO<sub>2</sub> for photosynthesis is the resistance of stomata, regulated by the plant in response to environmental conditions. Stomatal resistance, or its reciprocal the stomatal conductance, is an important parameter in hydrological modelling, controlling the rate of transpiration for the different vegetation types. The version of WaSim applied in present study uses the Penman-Monteith approach for calculating evapotranspiration and requires a parameter of minimal surface resistance for a state when plants are fully supplied with water (Schulla, 1997; Schulla and Jasper, 2013). The real transpiration modelled is further influenced by meteorological boundary conditions and the available soil water.

For the Rsc measurements in poplar SRC we used the SC-1 leaf porometer (Decagon Devices Inc.; Pullman, WA, USA). The measurements took place in Reiffenhausen in 2013 and were carried out weekly or fortnightly from May till September only under favourable weather conditions promising minimal resistances: preferable sunny, but at least without rain and with dry leaves. The same 10 plots in the poplar SRC as for the LAI measurements were used, where 3 sun-leaves were marked to be measured at different times. All 10 locations were measured during one hour to minimize the effects of changing weather conditions. Measurements were started in the morning, when leaves are dry and continued till afternoon, or as long as weather conditions were appropriate.

### 2.2.4 Phenology – start of growing season

The phenological phases of plants, e.g., leaf unfolding, leaf colouring and fall of leaves, are controlled by environmental conditions and internal genetic characteristics of plants. Thus, the site and species specific phenological state is a result of complex interference between length of light period, meteorological drivers (mainly temperature and radiation), soil properties, plant provenance, age and height (Menzel, 2000).

Within WaSim a modified approach for estimating LU according to Cannell and Smith (1983) is implemented and used here. A detailed description of this model as presented in Eq. (2-5), as well as parameterisation examples (Tab. 1) is given by Menzel (1997).

The model has four parameters:  $T_0$ ,  $T_1$ ,  $a$  and  $b$  which are the threshold temperatures for chilling units and for forcing units and two tree specific regression parameters, respectively. The starting day for leaves unfolding is calculated according to Eq. (2), (3), (4) and (5).

$$T_S = \sum_{i=0}^n \begin{cases} T(t_1 + i\Delta t) - T_1 & T(t_1 + i\Delta t) \geq T_1 \\ 0 & T(t_1 + i\Delta t) < T_1 \end{cases} \quad (2)$$

Here  $T_S$  is the temperature sum,  $T$  is the daily mean temperature for a day  $t_1 + i\Delta t$ ,  $t_1$  is set as 1 February in present study and time step,  $\Delta t$ , as one day. The daily mean temperature is calculated according to Eq. (3).

$$T = \frac{T_{\min} + T_{\max}}{2} \quad (3)$$

Here  $T_{\min}$  is the daily minimum temperature and  $T_{\max}$  the daily maximum temperature. The LU occurs when  $T_S$  reaches the critical value  $T_{S,\text{crit}}$  (Eq. 4 and 5).

$$T_{S,\text{crit}} = a + b \ln(CD_n), \text{ with} \quad (4)$$

$$CD_n = \sum_{i=0}^n \begin{cases} 1 & T(t_0 + i\Delta t) \leq T_0 \\ 0 & T(t_0 + i\Delta t) > T_0. \end{cases} \quad (5)$$

$CD$  is the number of chilling days, i.e. when  $T < T_0$ , between days  $t_0$  and  $t_1$ . The date  $t_0$  was set as 1 November in present study. Values for  $T_0$ ,  $T_1$ ,  $a$  and  $b$  for *Populus tremula* (IPG235) are given by Menzel (1997) (Tab. 1).

Using these numbers as initial values we fitted the parameters  $T_0$ ,  $T_1$ ,  $a$  and  $b$  to observed LU for the poplar clone Max1 in Reiffenhausen for the years 2012 and 2013 using least squares method. Finally we evaluated the obtained model parameters against the independent observations in Reiffenhausen (for 2014), and observation in Großfahner for the years 2012 and 2013 (Lorenz and Müller, 2013) and 2014 (Lorenz, 2014). The observed LU in Reiffenhausen and Großfahner is comparable to the recommendations according to Volkert and Schnelle (1966). Because the estimation of LU, as used in this study is based on meteorological measures, the parameterisation should hold true for the same poplar clone in the same age, if other environmental factors are of minor importance. The results will confirm this.

For a long term comparison the data from the international phenological observation networks (IPG) are used (Chmielewski et al., 2013), namely LU of *Populus tremula* (IPG235) at the IPG station Tharandt-Hartha. For poplar clone Max1 we could not find parameter sets in literature, so we used for comparison the IPG235 parameters of Menzel (1997) for *Populus tremula*, which is better investigated. The IPG-data are used for long term comparison, because there were no long term investigations of LU available on the research plots of the BEST project or nearby. IPG235 is the acronym of the parameterisation for *Populus tremula*

used by (Menzel, 1997). We decided to retain this acronym to make it comparable to published results, and also because it is an acronym used in the data provided by the phenological garden network. The phenological phase of leaf unfolding is defined as the stage UL, according to the IPG webpage (International Phenological Gardens of Europe, 2014) and is obtained by daily observations of plant's development state. The IPG station Tharandt-Hartha is located at the eastern border of the Tharandter Wald. It is the nearest IPG station to the site Reiffenhausen, and comparable in climate and altitude. Modeling approach

For simulation, the deterministic spatially distributed hydrological catchment model system WaSim (version 9.05.04) was used. Complete and comprehensive descriptions of this model and its internal structure can be found inter alia in (Schulla, 1997; Schulla and Jasper, 2013). The setup of physically based parameters, such as LAI and Rsc as well as phenological state (date of leaf unfolding - LU here), are predicated on direct measurements and observations. Thus, physical nexus between model image and reality is reproduced as best as possible. The SRC described with these measurements and observations represents a poplar SRC in the 3rd growing season of its mono-stem cycle, which can be seen as a hydrological fully developed canopy, concerning LAI, Rsc and root development. The simulated local soil water contents were compared and evaluated with measurements.

Different model simulations are done to show the suitability of the direct use of specific plant physiological measurements, as well as the effects of an approximated parameter description in the model, i.e. the annual course and the quantity of LAI, Rsc and phenology.

All these model approaches were done on a plot model domain, which are 3x3 raster cells based on a digital elevation model with a spatial resolution of 12.5 m (LGLN - Landesbetrieb Landesvermessung und Geobasisinformation, 2013), provided by the project partner NW-FVA<sup>1</sup>. All topographic information, needed by the model is derived by the model itself. The research site, providing the measured soil water contents for model calibration is located in the centre of the domain. A retention curve required in hydrological modelling for the description of soil physical properties was taken from Van Genuchten (1980). The Van Genuchten retention parameters from Blume et al. (2010) were accepted based on a characterization of soil texture and soil horizons in Reiffenhausen.

---

<sup>1</sup> Nordwestdeutsche Forstliche Versuchsanstalt (NW-FVA), Northwest German Forest Research Station

The meteorological forcing data were taken from own measurements for the period 2011-2013, whereas the first two years were used for the model spin up. Analyses and the evaluation to measured local soil water contents were done for the year 2013, only. To show the effects of different parameterisations under various climate conditions the simulations were performed for the period from 1969 to 2013 using the forcing meteorological data from DWD station Göttingen. The period was chosen as the longest period without missing values. The parameterisation of land use is kept constant for the whole period. A WaSim control file including all information about parameterization and model setup is provided as supplementary material.

## **2.3 Data analysis**

All measured and applied meteorological, soil physical and eco physiological parameters have been checked for plausibility and measurement errors.

The data has been numerically analysed and graphically presented with the free software package GNU Octave, version 3.6.2 (Octave community, 2012). Parts of the statistical analysis were performed using the hydroGOF package (Mauricio Zambrano-Bigiarini, 2014) within the R software environment (R-Studio under Windows, version 0.98.501) for statistical computing and graphics (R Development Core Team, 2011).

The evaluation of model performance was done according to objective criteria of Moriasi et al. (2007). Important quality criterions of simulation runs are the Nash-Sutcliffe model efficiency criterion (NSC), the percent bias (PBIAS), and the ratio of the root mean square error to the standard deviation of measured data (RSR).

## **3 Results**

### **3.1 Measurements**

#### **3.1.1 Leaf area index**

Figure 1 shows the annual course of LAI, derived from two different indirect optical and one direct destructive method. LU started shortly before the first measurement at the 1 May 2013. Until the 1 August there is almost a linear increase of LAI up to 7.3 and 5.5 for the LI191SA and the LAI2000 measurements, respectively. After that LAI starts to decrease, with a more rapid decline toward the end of August 2013. Leaf fall was almost finished at the 25 October.

Differences between devices, i.e. LI191SA vs. LAI2000, are large. The LAI values obtained with the LI191SA are systematically higher ( $\approx 2 \text{ m}^2 \text{ m}^{-2}$ ). The values obtained by direct destructive sampling at the 26 August are rather on the level of the LAI2000 estimates.

### 3.1.2 Stomatal resistance

Figure 2 shows the stomatal resistance ( $R_{sc}$ ) as measured on well illuminated leaves in 2013. Values are ranging from  $100 \text{ s m}^{-1}$  to  $300 \text{ s m}^{-1}$  until August 2013. At the 18 June  $R_{sc}$  is higher with larger standard deviations as the previous measurement at the 14 June. Soil water supply is sufficient on both days. The two days significantly differ in temperature, although the 14 June is relatively colder with a daily maximum temperature of approximately  $17^\circ\text{C}$  and the 18 June is quite hot, reaching a maximum temperature of  $33^\circ\text{C}$ . This shows the effect of local environmental conditions to measurements, possibly influencing derived model parameters. Starting from August both mean  $R_{sc}$  and standard deviation are steadily increasing. This period is characterized by decreasing soil water availability leading to severe drought stress conditions. Due to higher  $R_{sc}$  the trees counteracting the drought stress to avoid water loss and xylem damage, e.g. embolism of xylem vessels. The increase of standard deviation is an expression of stand heterogeneity, single trees still have access to water, and other may already be limited or stressed. In September 2013 we stopped measurements because leaves were visibly affected by the drought stress event. The correlation of plant available soil water and plant regulation via stomata seems to be consistent, increasing confidence in the distinct measurements. The minimum observed stomatal resistance is  $80 \text{ s m}^{-1}$ .

### 3.1.3 Phenology – start of growing season

We used in situ phenological observations of the two years 2012 and 2013 in Reiffenhausen to calibrate the modified approach for estimating LU according to Cannell and Smith (1983), which is used in WaSim. In 2012 LU for Max1 in Reiffenhausen started at day of year (DOY) 88, i.e. the 28 March 2012 (Tab. 2). In 2013 LU was delayed by approximately 4 weeks due to low temperatures in spring and started at DOY 115 (25 April 2013). Our calibration resulted in values of  $10^\circ\text{C}$  and  $2^\circ\text{C}$  for  $T_0$  and  $T_1$ . The regression parameters  $a$  and  $b$  are 2200 and -403, respectively (Tab. 1). Estimates of LU using these values for  $T_0$ ,  $T_1$ ,  $a$  and  $b$  show deviations from the observed dates of +3 and -3 days for Reiffenhausen in 2012 and 2013, respectively (Tab. 2). Then the phenological model results with the obtained parameter set

1 and local temperatures were compared to phenological observations in Reiffenhausen in 2014  
2 and in Großfahner in 2012 and 2013. Observed LU in Großfahner are almost equal to that in  
3 Reiffenhausen, also showing the delay of approx. 4 weeks in 2013 compared to 2012. The  
4 phenological model using the Max1 parameters result in differences of -1 day for  
5 Reiffenhausen in 2014 and of +1 and -1 days for Großfahner in 2012 and 2013 compared to  
6 observations. Tab. 2 also shows the application of the IPG235 parameter set provided by  
7 Menzel (1997) for *Populus tremula*. Parameters of  $a$  and  $b$  for IPG235 are both smaller in  
8 magnitude and threshold temperatures for chilling and forcing units,  $T_0$  and  $T_l$  show smaller  
9 differences. Due to this wider spread between  $T_0$  and  $T_l$  the Max1-model is able to describe  
10 extreme values and therefore a higher variability of LU, which was observed in 2012 and  
11 2013. The model estimations of LU with Max1 and with the IPG235 parameters differ  
12 considerably. The IPG235 set produces systematically later dates. Differences to observations  
13 are +31 (2012), +10 (2013) and +28 (2014) days for Reiffenhausen and +23 (2012) and +7  
14 (2103) days for Großfahner using the local temperatures (Tab. 2, column: local).

15 To assess the effects of non-local micrometeorological data sources, the model was driven by  
16 temperature measurements from the nearest DWD stations, namely Göttingen for  
17 Reiffenhausen and Dachwig for Großfahner. Expectedly, the use of DWD data instead of the  
18 local measurements produces mostly larger estimation errors for both the Max1 and the  
19 IPG235 parameter set (Tab. 2, column: nearest DWD).

20 We will use the varying parameter sets, i.e. our Max1 model and the IPG235 parameter set  
21 (Tab. 1) to analyse the effect for hydrological modelling for the year 2013, where soil  
22 hydrological measurements are available to evaluate the hydrological model results.

23 To analyse the species-dependence of LU estimations the model with Max1 and IPG235  
24 parameters was also driven by temperature measurements during 2012-2014 at the  
25 phenological station Tharandt. Figure 3 (a-d) illustrates that, expectedly, the parameter sets  
26 correspond better to the observations at species for which they were calibrated: Max1  
27 parameters to Reiffenhausen and the IPG235 parameters to Tharandt observations, which  
28 were part of its calibration dataset. The differences between estimated and observed DOY of  
29 LU are smaller when local temperature measurements are used (Fig. 3a vs. 3b and Fig. 3c vs.  
30 3d).

31 Figure 3 (e-f) show the long term courses of estimated DOY of LU for Reiffenhausen and  
32 Tharandt using the temperatures of the nearest DWD stations (Göttingen and Wildacker,

respectively). The model with IPG235 is systematically later and shows less variability than with Max1 parameters. For Reiffenhausen no long term phenological observations are available. However, the average DOY of LU in Reiffenhausen is  $\text{DOY } 97 \pm 9$  using Max1 and  $\text{DOY } 124 \pm 5$  for IPG235. The long term phenological observations in Tharandt fit well to the IPG235 estimates, but showing less variability than observed. The average DOY of LU in Tharandt as observed is  $\text{DOY } 123 \pm 10$  days, estimated using Max1  $\text{DOY } 101 \pm 9$  days and with IPG235  $\text{DOY } 124 \pm 7$  days. In general estimates fit best to observations when the corresponding parameters are used, i.e. Max1 for Reiffenhausen and IPG235 for Tharandt. But variability is underestimated by IPG235 compared to observations.

## 3.2 Hydrological model simulations

Several model simulations were performed with different parameterisations of LAI, Rsc and LU. Table 3 summarizes the eight performed model simulations and introduces their abbreviations. The detailed descriptions of model simulations are given in text.

First the measured values of LAI, Rsc and LU are implemented for hydrological modelling (LAI2000 Rsc80 and LI191SA Rsc80). Starting from here we changed the parameter sets: i) to improve the model fit; ii) to adjust the suitability of applied parameterisations and iii) to show the effects of different parameterisations on hydrological model results.

### 3.2.1 Simulation using observed parameters and adaptation of stomatal resistance

First we used the measured annual courses of LAI for hydrological modelling. Rsc is set to the measured minimum of  $80 \text{ s m}^{-1}$ , when LAI is larger than 1. LU is not calculated for measured LAI from air temperature using the approach of Cannell and Smith (1983), because this information is already imprinted in LAI measurements and therefore fixed for the year 2013.

Figure 4 shows the applied model parameterisations for LAI and Rsc, as well as the plant available water (PAW), calculated until 1 m soil depth from measured and modelled soil water contents.

For all simulations using the measured value for Rsc (i.e.  $80 \text{ s m}^{-1}$ ) model results for soil water content were higher than measured values, resulting in larger PAW-values than observed. This is also reflected by the Nash-Sutcliffe criterion (NSC) calculated from PAW

(Tab. 4). The annual course of PAW is captured quite well by the model, but the drying up in summer is not sufficient, neither for the LI191SA, nor for the LAI2000 measurements. The NSC is better for the experiments with LAI measured by LI191SA (0.69) than with LAI2000 (0.44) because of the higher LAI values (Tab. 4).

As, especially the maximum of LAI measures using the LAI2000 showed better agreement with direct destructive measurements, we halved the value of  $R_{sc}$  from  $80 \text{ s m}^{-1}$  to  $40 \text{ s m}^{-1}$  to reach the low PAW values observed. This decision has two reasons. First it can be assumed that the measured  $R_{sc}$  is always higher than the minimum value needed for parameterisation, because the conditions by measuring  $R_{sc}$  are not satisfying the requirements for the parameter to be used in the model, i.e. optimal conditions for transpiration and no water stress. Another way to get lower PAW values would be to increase the LAI, as can be seen by comparing the results for LI191SA and LAI2000. However, in our experiment LAI has to be increased to unrealistically high values to minimize the differences to observed PAW. Additionally LAI is also affecting other processes in hydrological models, like interception evaporation and soil evaporation. Together the decrease of  $R_{sc}$  is a consistent way to minimize the deviations to observations and to improve the model fit.

The reduction of  $R_{sc}$ , from  $80 \text{ s m}^{-1}$  to  $40 \text{ s m}^{-1}$ , improved the NSC from 0.69 and 0.44 to 0.89 and 0.87 for the LI191SA and LAI2000, respectively (Tab. 4).

### 3.2.2 Approximation and adaptation of annual course of leaf area and stomatal resistance

In many cases when hydrological models should be applied for analyses involving vegetation there are no locally measured data on LAI and/or  $R_{sc}$ . Often only the literature data for the maximum and minimum values of LAI and  $R_{sc}$  are available. Then the annual course for these parameters has to be derived or approximated for modelling. The simplest approximation is a stepwise function, where the increase from minimum to maximum or decrease from maximum to minimum occurs within one time step. We applied this form to the LAI and  $R_{sc}$  as shown in Fig. 5. Here the maximum of LAI is set to  $6 \text{ m}^2 \text{ m}^{-2}$ , which is the observed maximum plus standard deviation of the LAI2000 measurements. The minimum of  $R_{sc}$  is set to  $40 \text{ s m}^{-1}$ . For this kind of approximation the start, and therefore the length of the growing season becomes important, because the maximum transpiration rate occurs



1 immediately after LU. In Fig. 5 we compare two different parameterisations for dynamical  
2 estimating LU, i.e. the Max1 and the IPG235 parameter set (Tab. 1).

3 The NSCs for both simulations are 0.89 (Tab. 4), which is even a bit better than for applying  
4 the direct LAI measurements. However, PBIAS values are negative for the step function  
5 simulations, where they are positive for the simulations using LAI measurements. Negative  
6 PBIAS values indicate a stronger drying signal.

7 Differences in PAW are only visible in the period when parameterisation is different (Fig. 5).  
8 The year 2013 shows no drought event in spring, where effects would be more obvious. There  
9 are small differences in May, where the step function simulation using the Max1 parameters  
10 are closer to observations.

11 However, the abrupt increase to the maximum transpirations rate immediately after LU is  
12 rather unrealistic as already shown by the LAI measurements. Unfolding of leaves in nature  
13 can happen very quickly, as everybody can observe when spring comes late in the year,  
14 followed by favourable growth conditions. When spring starts early the full leaf development  
15 can take much longer. To account for this and to further improve our model fit we changed  
16 the annual development of LAI and Rsc by using these parameters for manual model  
17 calibration, guided by the course of LAI measurements mainly (Fig. 6). Major changes are  
18 higher LAI and lower Rsc values at the date of leaf unfolding, i.e.  $2 \text{ m}^2 \text{ m}^{-2}$  and  $150 \text{ s m}^{-1}$ ,  
19 respectively. LU is estimated with the dynamic approach like in the step function simulations.  
20 This resulted in modelled higher transpiration rates in spring. Besides the annual course of  
21 LAI and Rsc is described more detailed and more similar to the observed LAI dynamics - the  
22 LAI increase and decline is smoother but also starts a bit earlier in the year and last a bit  
23 longer in autumn. Due to that smoother increase in spring the sensitivity to deviations in  
24 estimating LU is reduced.

25 Due to these changes the NSC increased to 0.90 for both Max1 and IPG235 parameter sets.  
26 This is the best fit obtained in manual calibration procedure (Tab. 4). PBIAS values are  
27 positive for the adjusted models, which is a slightly too small drying signal. However, the  
28 magnitudes of PBIAS and RSR values are smaller than for the step function simulations,  
29 indicating better agreement with observations and lower root mean square errors or residual  
30 variations (Moriassi et al., 2007).

### 3.2.3 Long term simulations

In all simulations shown for the year 2013, the effects on PAW caused by changes in estimated LU are quite low, due to the high soil water contents in spring 2013. Therefore, we applied all simulations for the years 1969-2013, which was the longest meteorological period without missing data. A focus is set to the year 2012, which was characterized by an early drought event in May. Because of missing data there is no complete set of soil water content information available for this year and there is no information about LAI and Rsc for 2012 that can be used to parameterise the hydrological model. So no evaluation of model fit is possible for 2012 or the other years of the period 1969-2013, like it is done for 2013.

To illustrate the effects for the different courses of LAI and Rsc development as well as the estimation of LU, Fig. 7 shows the precipitation, the plant available water and the GWR for the step function and for the adjusted simulations combined with the estimates of LU, i.e. the Max1 and IPG235 parameter set, respectively. Results in Fig. 7 show the last two years from the long term simulations 1969-2013, mean values for ETR and GWR for the whole period 1969-2013 are presented in Tab. 5. In 2012 and 2013 as well as for both estimations of LU, the adjusted simulation shows the highest GWR and the step function simulations results in the lowest GWR. The reason for this is the change in transpiration in spring, as described due to the different parameterisations of the step function and adjusted course for LAI and Rsc. However, the largest effects on GWR are caused by the different estimation of LU (Fig. 7). In the step function experiments GWR is zero in 2012 with both the Max1 and the IPG235 parameterisations of LU. Plant available water is reduced stronger for the Max1 parameterisation, due to the early start of the growing season. However, this early LU fits better to the observations in Reiffenhausen. For the adjusted simulations GWR in 2012 is only zero for the Max1 parameterisation. For the year 2012 data of the matrix potential (tensiometer measurements) in 20 cm, 60 cm and 120 cm soil depth are available. These data show a drought period in May 2012, where the tensiometers in 20 cm soil depth run out the measuring range, i.e. a matrix potential was lower than approx. -800 hPa. Starting from May 2012 the tensiometers in 60 cm and 120 cm soil depth indicated a consistent drying signal (not shown). Additionally the poplar SRC in 2012 is younger and therefore less water demanding than the poplar SRC parameterised in the model, applied for these analyses. This indicates that GWR after May 2012 is very unlikely in these simulations.

The parameterisation using the adjusted course for LAI and Rsc, based on the measured course of LAI, in combination with the Max1 parameterisation for LU, calibrated at local observations, seems to be the most realistic model simulation (LAIadjusted Rsc40 Max1). By comparing all four model parameter combinations shown in Fig. 7, one can switch completely from GWR present in 2012 to absent.

Table 5 summarizes the evapotranspiration (ETR) and GWR for all simulations, averaged over all years for the period 1969-2013, as well as for the 5 driest and 5 wettest years of this period. GWR averaged over all years varies from 80 mm year<sup>-1</sup> to 145 mm year<sup>-1</sup> depending on the approximation of the annual course of LAI and Rsc and the estimation of LU. The ratio of maximum and minimum of the all year averages of GWR for the different simulations is approx. 1.8. This factor is approx. 3 for the 5 driest years and approx. 1.7 for the 5 wettest years, showing that especially the model results for dry years are sensitive to the parameterisations used.

#### 4 Discussion

Not all necessary model parameters for WaSim could be measured in detail. One example is the implemented assumption on rooting depth which was measured in 2012, and was set to 1 m for modelling which is comparable to the commonly used values presented in Raissi et al. (2009).

Measuring Rsc in the field is rather challenging. For hydrological modelling we are interested in more theoretically minimum values, indicating optimal transpiration. These conditions are hardly found in reality. In addition the measurements are affected by soil water availability as well as rapidly changing atmospheric conditions. Breuer et al. (2003) summarises values for minimal stomatal resistance for various plants. Values for Populus clones (Populus grandidenata, P. tremula and P. tremuloides) are ranging from 102 s m<sup>-1</sup> to 400 s m<sup>-1</sup>. Our measured minimum of Rsc for poplar clone Max1 (Populus nigra x Populus maximowiczii) is lower: 80 s m<sup>-1</sup>. Yet we needed to further reduce Rsc to 40 s m<sup>-1</sup> for modelling in order to match the observed soil water contents. On the one hand the low observed minimum of 80 s m<sup>-1</sup> shows that specific measurements of Rsc are helpful. On the other hand the measurements of Rsc were still too high to produce plausible results with WaSim. One might interpret the reduction of Rsc from 80 s m<sup>-1</sup> to 40 s m<sup>-1</sup> as a shift from the often reported isohydric behaviour of poplar clones (Tardieu and Simonneau, 1998) to a more anisohydric behaviour. But the diurnal or seasonal variations of leaf water potential that are characteristic

for anisohydric plants are not expressed by the Rsc value in WaSim, which represents the minimal resistance for a state when plants are fully supplied with water. The reduction of transpiration in drought stress situations is done in a different way in WaSim. Furthermore, there are also more drought-tolerant, anisohydric water use strategies reported from greenhouse experiments for poplar clones (Ceulemans et al., 1988; Larchevêque et al., 2011). Schmidt-Walter et al. (2014) reported also a poor stomatal control of water loss estimated from field measurements of a poplar SRC.

The LAI measurements show a systematic difference between the two measurement devices, whereas the LI191SA seems to overestimate LAI taking the destructive method as a reference. In situ measurements of LAI are helpful to determine the maximum value, but differences due to the different estimation methods including underlying assumptions should be considered. The annual development of LAI is indispensable information to adjust and improve the model parameterisation of annual course. The measured LAI development represents local conditions and is therefore valid for the measurement site and time period only. Approximations of seasonal course are advisable to enable the transferability to other sites and years. A crucial factor here is LU, determining the start of LAI increase. For the determination of this date phenological stages have been defined. To describe LU various models are available, based on air temperature, soil temperature, photoperiod, day length or radiation. All models have to be calibrated for specific plants species. There is also evidence that local conditions like latitude or altitude of observations are influencing the calibration of the phenological model. Furthermore the derivation of parameters for the phenological model will depend on the observed data, e.g. the detection of extremely early or late LU as well as the climate data, which has to be appropriate for the observed site. For poplar clone Max1 we could not find parameter sets in literature, so we used for comparison the IPG235 parameters of Menzel (1997) for *Populus tremula*, which is better investigated. The period of parameter adjustment used by Menzel (1997) is 1959-1993, and is based on several phenological stations, whereas our derived parameters are based on 2 years at one site. However, these two years show a wide variability in LU. The parameters from Menzel (1997) should be generally more valid, because of the higher number of observations. Yet the use of IPG235 parameters resulted in an underestimation of observed variability, compared to the observations for *Populus tremula* in Tharandt. Differences between IPG235 and Max1 also show the importance of parameterisations for local site conditions and specific species. Comparing the parameter sets presented in Tab. 1 these effects become evident.

Especially threshold temperatures for chilling and forcing units,  $T_0$  and  $T_I$  vary more widely between IPG235 and Max1. Due to this wider range the model is able to describe extreme values and therefore a higher variability of LU, which was observed in our calibration years 2012 and 2013. We evaluated our parameter set on observations of the poplar SRC Reiffenhausen in 2014 and the poplar SRC Großfahner (2012-2014), which was planted with the same clone and in the same year like Reiffenhausen. The differences for LU between observations and the Max1 model setup are low and within the observed variability. The use of IPG235 parameters for the Max1 clone, which is a common procedure when specific values are missing, can result in large deviations as shown for ground water recharge (GWR), especially in the year 2012 with the drought period in spring.

The source of temperature data also influences the parameters derived for phenological models as well as the results obtained by applying these parameter sets. We compared the estimated DOY of LU derived with local temperature measurements and with temperatures of the nearest DWD stations. For Reiffenhausen, with the nearest DWD station Göttingen, we additionally tested an altitude correction using the vertical temperature gradient of  $-0.0065\text{ }^{\circ}\text{C m}^{-1}$  to account for 158 m altitude difference between Göttingen (167 m a.s.l.) and Reiffenhausen (325 m a.s.l.). Deviations in DOY of LU are small when using the DWD temperature instead of the local measurements. Interestingly, the altitude correction of temperature increases differences in DOY of LU comparing to observations. The reason could be the often occurring thermal inversion, when the air temperature in Reiffenhausen is higher than in Göttingen, so that implemented altitude reduction of temperature increases the differences even more, due to that also the differences of the estimated DOY of LU increase. The effects of the altitude correction are larger for the Max1 than for the IPG235 parameter set, because our model is more sensitive to extreme values due to higher  $T_0$  and lower  $T_I$  temperatures. This shows the importance of applying the local temperatures, associated to the phenological observations, to calibrate and use the temperature-dependent phenological models. The use of local temperatures improves the estimation of LU and better represents inter annual variability.

According to the criteria of Moriasi et al. (2007), the hydrological model results, using measured values of LAI and Rsc (start and development of LU is implemented), are satisfactory only for the simulation LI191SA with  $\text{Rsc} = 80\text{ s m}^{-1}$ . The simulation using the LAI values from the LAI2000 fails to satisfy the recommended criteria (Tab. 4). However the

model produces better agreement with observations when  $R_{sc}$  minima of  $40 \text{ s m}^{-1}$  are used with any LAI data. The reduction of  $R_{sc}$  is a consistent way to simulate the observed soil water conditions. An increase of LAI could lead to lower soil water contents as well, but is also effecting soil evaporation and interception evaporation. Additionally larger values for LAI, necessary to minimize the model deviations to measurements, have to be unrealistically high for the poplar SRC investigated here. When using a  $R_{sc}$  minimum of  $40 \text{ s m}^{-1}$  together with measured LAI the model evaluation is good for the year 2013, reaching NSC values of 0.87 and 0.89 for the LAI2000 and LI191SA simulations.

Data of such intense measurement campaigns are not available for all sites where hydrological modelling should be done. Therefore literature values, often providing maximum or minimum values for LAI and  $R_{sc}$  only, are used and the annual course has to be modelled. The question of transferability of these values to different sites, years or even species has to be solved. The applied step function is the simplest approximation of the annual course for LAI and  $R_{sc}$ . These simulations also pass the recommended criteria for a satisfactory model performance (Tab. 4).

For the year 2013 the best model fit could be obtained by the adjusted annual courses for LAI and  $R_{sc}$ . They are based on the observed course and maximum values of LAI measurements.

The weather regime and therefore the development of soil water conditions are not suitable in 2013 to show the effects of different estimates for the start of LU in spring. Drought conditions started not until July 2013. Therefore we performed scenario simulation by transferring the vegetation parameterisation for 2013 to the weather regime of 2012. This year was characterized by a drought period in spring. In consequence the effects of different estimates of DOY of LU are pronounced. The adjusted simulations using the IPG235 parameters to estimate LU, i.e. later LU by approx. 30 days in 2012 show GWR in this year. Due to the delayed start of the growing season the drought stress in spring is missed in the model mostly, leading to wetter soil conditions which benefits percolation and rewetting and finally enlarges GWR (Fig. 7). The tensiometer measurements available for 2012 suggest that GWR is rather unlikely for this year. The step function simulation using the IPG235 parameters for LU and both adjusted simulations (Max1 and IPG235) result in zero GWR for the year 2012. However, the strongest simplification of the course of LAI and  $R_{sc}$ , i.e. the step function, shows the lowest GWR for 2013 and for the long term simulations (Tab. 5).

In Fig. 7 the effects of the different simulations on GWR are presented for the years 2012 and 2013, which are characterized by rather different weather regimes. Whereas a realistic description of LAI and Rsc seems to be less important in 2013, it is even more essential in 2012, showing the importance of distinct spatial and temporal characteristics for local modelling.

We performed a long term simulation, by keeping the parameterisation for the vegetation constant for the period 1969-2013 to account for the effects of climate variability. This is a more theoretical scenario, because it accounts for changes in climate forcing only. In reality also the vegetation characteristics are changing over the years, as well as soil properties on a longer time scale, especially for SRC, whereas rotation cultivation is applied, e.g. harvesting and resprouting. Particularly the rotation cultivation can reduce extreme drought conditions, when dry years coincide with rotation stages that have a lower water demand. The vegetation parameterised here can be seen as fully developed in hydrological terms, characterized by a large water demand. The simulations here are rather artificial, especially by succeeding dry years when soil water storage is not refilled completely in winter and drought conditions are influencing the following growing season. Nevertheless the effects caused by different descriptions of vegetation parameterisations are quiet large (Tab. 5). Especially on a local scale such differences can be important by evaluating effects of land use change, particularly in dry years.

Taking into account, that the best model evaluation for 2013 is achieved with the adjusted course of LAI and Rsc, the adjusted simulation using the Max1 phenology parameters seems to be the most reasonable parameterisation. It fits best to the evaluation in 2013.

## **5 Conclusions**

For hydrological analysis of some area or sites with the focus on land use change or climate change the adequate parameterisation of the vegetation cover is important by determining processes like soil evaporation, interception evaporation and transpiration. Sources of model parameters for the vegetation cover are local measurements or scientific literature. The analysis shows simulation uncertainties evolving from the use of model parameters that are derived from i) non-local measurements or ii) some appropriate literature values.

Answering objective 1 our study shows that LAI, Rsc as well as the beginning and length of growing season are very sensitive parameters when effects of an enhanced cultivation of SRC on local water budget are investigated. Particularly, it reveals that correct information about

1 the beginning of the growing season is highly important to obtain correct and acceptable  
2 simulation results of evapotranspiration components and GWR. If the start of growing season  
3 is miscalled, such as shown for the different species as in the IPG235 and Max1  
4 parameterisation (Tab. 1), the accuracy of other parameters, i.e., LAI and Rsc; plays a minor  
5 role. Concerning GWR, LU is the most sensitive parameter. Its parameterisation is  
6 particularly important when inter annual variations and hydrological extreme conditions are  
7 on focus.

8 The implementation of locally measured vegetation parameters for hydrological modelling  
9 has both advantages and drawbacks. Measurements are expensive, time consuming and also  
10 not always feasible. In such cases the use of appropriate literature values and transposition of  
11 adjacent observations is necessary and common practice.

12 The present study displays for the parameter LAI, that the simulations using locally measured  
13 plant specific values show the suitability of data. The comparisons between locally measured  
14 and adjusted parameter sets reveal that simulation results are less affected by other model  
15 parameters, like Rsc or LU, when using adjusted parameters of LAI.

16 Contrary results appear for Rsc. Simulation results differ significantly when site specific  
17 values of Rsc are available. However, for Rsc the benefit of direct use of local measurements  
18 is arguable: minimum has to be reduced within WaSim to produce model results comparable  
19 with soil water measurements. In consequence the implementation of Rsc values from  
20 literature for hydrological modelling without accompanying measurement data for model  
21 evaluation can produce very uncertain results. The analyses illustrate that the locally adjusted  
22 vegetation parameterisation shows the best model fit. Additionally the adjusted course of LAI  
23 and Rsc is less sensitive to different estimates for LU, due to a slower increase in spring  
24 compared to a step functional annual course. But the adjusted courses are also approximations  
25 and not a distinct measure and are therefore more generally valid for different sites and years,  
26 than a direct use of measured parameters.

27 For the land use poplar SRC there are certain years where the modelled GWR is reduced to  
28 zero, like in the year 2012 (Fig. 7). Different parameterisations for vegetation characteristics  
29 are influencing modelled GWR for those years producing a wide range from GWR present or  
30 completely absent.

31 Hydrological models are often used to analyse effects of climate and land use changes on  
32 spatial and temporal scale becoming smaller and smaller. Approximations in the description



1 of vegetation, a lack of local information (also soil and climate description), the transfer of  
2 inappropriate parameters and lacks in model formulation can cause large differences in  
3 simulation results. To account for small-scale and local effects of land use change more  
4 detailed descriptions of sites and processes are necessary to capture the spatial and temporal  
5 variability of effects. Especially when analysing extremes, they are often underestimated  
6 when the description of site and processes are insufficient.

## 7 **Acknowledgements**

8 The work was funded by the German Federal Ministry of Education and Research (BMBF)  
9 and is part of the BEST-Research Framework (<http://www.best-forschung.de>), we gratefully  
10 acknowledge this support.

11 We give our special thanks to D. Fellert and H. Kreilein for valuable advices and assistance in  
12 handling the meteorological equipment; to D. Böttger for his assistance in field work, to J.  
13 Sauer, H. Wendler and G. Kalberlah for their assistance in measurements. Thanks to A. Knohl  
14 for the fruitful discussions and for the help in solution of research problems. Special thanks to  
15 J. Schulla for advising in all WaSim questions arose during the study, solving problems and  
16 providing new model versions.

## 1   **References**

- 2   Aronsson, P. G., Bergström, L. F. and Elowson, S. N. E.: Long-term influence of intensively  
3   cultured short-rotation Willow Coppice on nitrogen concentrations in groundwater, J.  
4   Environ. Manage., 58(2), 135–145, 2000.
- 5   Baum, S., Weih, M. and Bolte, A.: Stand age characteristics and soil properties affect species  
6   composition of vascular plants in short rotation coppice plantations, 2012.
- 7   Bernhofer, C., Institut für Hydrologie und Meteorologie, Professur Meteorologie: Exkursions-  
8   und Praktikumsführer Tharandter Wald Material zum “hydrologisch-meteorologischen  
9   Feldpraktikum,” Techn. Univ., Dresden., 2002.
- 10   Blume, H.-P., Brümmer, G. W., Horn, R., Kandeler, E., Kögel-Knabner, I., Kretzschmar, R.,  
11   Stahr, K., Thiele-Bruhn, S., Welp, G. and Wilke, B.-M.: Lehrbuch der Bodenkunde,  
12   Springer., 2010.
- 13   Breda, N. J.: Ground-based measurements of leaf area index: a review of methods,  
14   instruments and current controversies, J. Exp. Bot., 54(392), 2403–2417, 2003.
- 15   Breuer, L., Eckhardt, K. and Frede, H.-G.: Plant parameter values for models in temperate  
16   climates, Ecol. Model., 169(2), 237–293, 2003.
- 17   Broeckx, L. S., Verlinden, M. S., Berhongaray, G., Zona, D., Fichot, R. and Ceulemans, R.:  
18   The effect of a dry spring on seasonal carbon allocation and vegetation dynamics in a poplar  
19   bioenergy plantation, GCB Bioenergy, 2013.
- 20   Cannell, M. G. R. and Smith, R. I.: Thermal time, chill days and prediction of budburst in  
21   Picea sitchensis, J. Appl. Ecol., 951–963, 1983.
- 22   Ceulemans, R., Impens, I. and Imler, R.: Stomatal conductance and stomatal behavior in  
23   Populus clones and hybrids, Can. J. Bot., 66(7), 1404–1414, 1988.
- 24   Chmielewski, F.-M., Heider, S., Moryson, S. and Bruns, E.: International Phenological  
25   Observation Networks: Concept of IPG and GPM, in Phenology: An Integrative  
26   Environmental Science, pp. 137–153, Springer., 2013.
- 27   Djomo, S. N., Kasmioui, O. el and Ceulemans, R.: Energy and greenhouse gas balance of  
28   bioenergy production from poplar and willow: a review, GCB Bioenergy, 3(3), 181–197,  
29   2011.

- 1 Don, A., Osborne, B., Hastings, A., Skiba, U., Carter, M. S., Drewer, J., Flessa, H., Freibauer,  
2 A., Hyvönen, N., Jones, M. B., Lanigan, G. J., Mander, Ü., Monti, A., Djomo, S. N.,  
3 Valentine, J., Walter, K., Zegada-Lizarazu, W. and Zenone, T.: Land-use change to bioenergy  
4 production in Europe: implications for the greenhouse gas balance and soil carbon, *GCB*  
5 *Bioenergy*, 4(4), 372–391, 2012.
- 6 Van Genuchten, M. T.: A closed-form equation for predicting the hydraulic conductivity of  
7 unsaturated soils, *Soil Sci. Soc. Am. J.*, 44(5), 892–898, 1980.
- 8 Hartmann, L., Richter, F., Busch, G., Ehret, M., Jansen, M. and Lamersdorf, N.: Etablierung  
9 von Kurzumtriebsplantagen im Rahmen des Verbundprojektes BEST in Süd-Niedersachsen  
10 und Mittel-Thüringen – Standorteigenschaften und anfängliche Erträge, *Forstarchiv*, 134–150,  
11 2014.
- 12 International Phenological Gardens of Europe: IPG, Phasen [online] Available from:  
13 [http://ipg.hu-berlin.de/ipg/faces/list\\_phases.xhtml](http://ipg.hu-berlin.de/ipg/faces/list_phases.xhtml) (Accessed 9 September 2014), 2014.
- 14 Jonckheere, I., Fleck, S., Nackaerts, K., Muys, B., Coppin, P., Weiss, M. and Baret, F.:  
15 Review of methods for in situ leaf area index determination: Part I. Theories, sensors and  
16 hemispherical photography, *Agric. For. Meteorol.*, 121(1), 19–35, 2004.
- 17 Kalberlah, G.: Wurzelverteilungsmuster einer Pappel-Kurzumtriebsplantage in  
18 Südniedersachsen, Master Thesis, Georg-August-Universität Göttingen, Göttingen., 2013.
- 19 Larchevêque, M., Maurel, M., Desrochers, A. and Larocque, G. R.: How does drought  
20 tolerance compare between two improved hybrids of balsam poplar and an unimproved native  
21 species?, *Tree Physiol.*, 31(3), 240–249, 2011.
- 22 Lasch, P., Kollas, C., Rock, J. and Suckow, F.: Potentials and impacts of short-rotation  
23 coppice plantation with aspen in Eastern Germany under conditions of climate change, *Reg.*  
24 *Environ. Change*, 10(2), 83–94, 2010.
- 25 LGLN - Landesbetrieb Landesvermessung und Geobasisinformation: Digitale  
26 Geländemodelle – DGM - ATKIS, 2013.
- 27 LI-COR INC: LAI-2000 Plant Canopy Analyzer Instruction Manual, Lincoln, USA., 1992.
- 28 Lorenz, K.: Personal communication, 2014.

1 Lorenz, K. and Müller, J.: Ergebnisse zur Nettoprimärproduktion von mit biologisch  
2 geklärtem Abwasser bewässerten Pappeln und Weiden im Kurzumtrieb, Landbauforsch. -  
3 Appl. Agric. For. Res., (63), 307–320, 2013.

4 Mauricio Zambrano-Bigiarini: hydroGOF: Goodness-of-fit functions for comparison of  
5 simulated and observed hydrological time series. [online] Available from: [http://CRAN.R-](http://CRAN.R-project.org/package=hydroGOF)  
6 [project.org/package=hydroGOF](http://CRAN.R-project.org/package=hydroGOF), 2014.

7 Menzel, A.: Phänologie von Waldbäumen unter sich ändernden Klimabedingungen:  
8 Auswertung der Beobachtungen in den internationalen phänologischen Gärten und  
9 Möglichkeiten der Modellierung von Phänodaten, Frank., 1997.

10 Menzel, A.: Trends in phenological phases in Europe between 1951 and 1996, Int. J.  
11 Biometeorol., 44(2), 76–81, 2000.

12 Moriasi, D. N., Arnold, J. G., Van Liew, M. W., Bingner, R. L., Harmel, R. D. and Veith, T.  
13 L.: Model evaluation guidelines for systematic quantification of accuracy in watershed  
14 simulations, Trans ASABE, 50(3), 885–900, 2007.

15 Octave community: GNU Octave. [online] Available from: [www.gnu.org/software/octave/](http://www.gnu.org/software/octave/),  
16 2012.

17 Petzold, R., Wahren, A. and Feger, K.-H.: Steuerungsoptionen des Wasser- und  
18 Stoffhaushalts auf Landschaftsebene durch den Anbau von Kurzumtriebsplantagen – ein  
19 Forschungsansatz., in Bodenschutz in Europa – Ziele und Umsetzung, vol. Tagungsband 6,  
20 pp. 91–96, Marktredwitzer. [online] Available from: [http://boku.forst.tu-](http://boku.forst.tu-dresden.de/pdf/petzold_wahren_feger_Steuerungsoptionen.pdf)  
21 [dresden.de/pdf/petzold\\_wahren\\_feger\\_Steuerungsoptionen.pdf](http://boku.forst.tu-dresden.de/pdf/petzold_wahren_feger_Steuerungsoptionen.pdf), 2010.

22 Raissi, F., Müller, U. and Meesenburg, H.: Geofakten 9 Ermittlung der effektiven  
23 Durchwurzelungstiefe von Forststandorten, Red. Geofakten Landesamt Für Bergbau Energ.  
24 Geol. Hann., (4. Auflage) [online] Available from: <http://www.lbeg.niedersachsen.de>  
25 abgerufen werden, 2009.

26 R Development Core Team: RStudio for Windows, R Foundation for Statistical Computing,  
27 Vienna, Austria. [online] Available from: <http://www.R-project.org>, 2011.

28 Rutter, A. J., Kershaw, K. A., Robins, P. C. and Morton, A. J.: A predictive model of rainfall  
29 interception in forests, 1. Derivation of the model from observations in a plantation of  
30 Corsican pine, Agric. Meteorol., 9(0), 367–384, 1971.

1 Schmidt-Walter, P. and Lamersdorf, N.: Biomass Production with Willow and Poplar Short  
2 Rotation Coppices on Sensitive Areas—the Impact on Nitrate Leaching and Groundwater  
3 Recharge in a Drinking Water Catchment near Hanover, Germany, *BioEnergy Res.*, 5(3),  
4 546–562, 2012.

5 Schmidt-Walter, P., Richter, F., Herbst, M., Schuldt, B. and Lamersdorf, N. P.: Transpiration  
6 and water use strategies of a young and a full-grown short rotation coppice differing in  
7 canopy cover and leaf area, *Agric. For. Meteorol.*, 195–196, 165–178, 2014.

8 Schulla, J.: Hydrologische Modellierung von Flussgebieten zur Abschätzung der Folgen von  
9 Klimaänderungen, ETH Zürich. [online] Available from: [http://dx.doi.org/10.3929/ethz-a-](http://dx.doi.org/10.3929/ethz-a-001763261)  
10 001763261 (Accessed 20 August 2014), 1997.

11 Schulla, J. and Jasper, K.: Model Description WaSiM-ETH, Inst. Atmospheric Clim. Sci.  
12 Swiss Fed. Inst. Technol. Zür. [online] Available from:  
13 [www.wasim.ch/products/wasim\\_description.htm](http://www.wasim.ch/products/wasim_description.htm), 2013.

14 Spank, U.: Site water budget: influences of measurement uncertainties on measurement  
15 results and model results, Eigenverlag der Technischen Universität Dresden, Dresden., 2010.

16 Spank, U., Schwärzel, K., Renner, M., Moderow, U. and Bernhofer, C.: Effects of  
17 measurement uncertainties of meteorological data on estimates of site water balance  
18 components, *J. Hydrol.*, 492, 176–189, 2013.

19 Surendran Nair, S., Kang, S., Zhang, X., Miguez, F. E., Izaurralde, R. C., Post, W. M., Dietze,  
20 M. C., Lynd, L. R. and Wullschleger, S. D.: Bioenergy crop models: descriptions, data  
21 requirements, and future challenges, *GCB Bioenergy*, 4(6), 620–633, 2012.

22 Tardieu, F. and Simonneau, T.: Variability among species of stomatal control under  
23 fluctuating soil water status and evaporative demand: modelling isohydric and anisohydric  
24 behaviours, *J. Exp. Bot.*, 49(Special Issue), 419–432, 1998.

25 Volkert, E. and Schnelle, F.: Arboreta Phaenologica: Mitteilung der Arbeitsgemeinschaft  
26 Internationaler Phänologischer Gärten, Erläut. Zur Beob. Phänologischen Termins  
27 Blattentfaltung Bei Populus Tremula, 7, 1966.

28 Watson, D. J.: Comparative physiological studies on the growth of field crops: I. Variation in  
29 net assimilation rate and leaf area between species and varieties, and within and between  
30 years, *Ann. Bot.*, 11(1), 41–76, 1947.





Figure 1. Means and standard deviations of leaf area index of the poplar SRC Reiffenhausen in 2013. Measurements of two optical devices: LI191SA calculated with constant extinction coefficient  $k = 0.5$  and LAI2000 are shown. LAI values obtained by destructive harvesting at the 26 August on three plots are shown as green dots.



1  
2 Figure 2. Means and standard deviations of stomatal resistance of sun leaves derived from 10  
3 to 11 repetitions every day at 10 measurement plots and (a) the plant available water,  
4 calculated from soil water content measurements until 1 m soil depth and (b) daily maximum  
5 temperature at the poplar SRC Reiffenhausen in 2013. High temperatures effecting stomatal  
6 resistance (18 June); starting from August drought stress occurred, increasing the stomatal  
7 resistances.

1 Table 1. Parameters of the modified approach for estimating leaf unfolding (LU) according to  
2 Cannell and Smith (1983), which is used in WaSim. Estimated day of years (DOY) of LU for  
3 Reiffenhausen (2012 and 2013) are used to calibrate the Max1 parameters  $T_0$ ,  $T_1$ ,  $a$  and  $b$ , i.e.  
4 threshold temperature for chilling units and forcing units and two tree specific regression  
5 parameters, respectively. Additionally the parameter set for IPG235 according to (Menzel,  
6 1997) (appendix A7) is shown.

| Parameter | Max1 | IPG235 (Menzel, 1997) |
|-----------|------|-----------------------|
| T0 [°C]   | 10   | 8                     |
| T1 [°C]   | 2    | 5                     |
| a         | 2200 | 1693.4161             |
| b         | -403 | -301.9361             |

7



1 Table 2. Observed and estimated day of years (DOY) of leaf unfolding (LU) for  
2 Reiffenhausen and Großfahner. The Max1 parameters are calibrated at the observations in  
3 Reiffenhausen using local temperatures (2012 and 2013) and evaluated with Reiffenhausen  
4 (2014) and Großfahner (2012 – 2014). Additionally the DOY of LU is compared to estimates  
5 using the IPG235 parameter set as well as the temperatures of the nearest DWD climate  
6 station (Göttingen for Reiffenhausen, distance approx. 17 km; Dachwig for Großfahner,  
7 distance approx. 3.5 km) are presented.

|                        | observed | Max1  | Max1           | IPG235 | IPG235         |
|------------------------|----------|-------|----------------|--------|----------------|
| temperature data       |          | local | nearest<br>DWD | local  | nearest<br>DWD |
| Reiffenhausen DOY 2012 | 88       | 91    | 89             | 119    | 121            |
| Reiffenhausen DOY 2013 | 115      | 112   | 112            | 125    | 126            |
| Reiffenhausen DOY 2014 | 83       | 82    | 85             | 111    | 113            |
| Großfahner DOY 2012    | 88       | 89    | 88             | 111    | 112            |
| Großfahner DOY 2013    | 114      | 113   | 112            | 121    | 121            |
| Großfahner DOY 2014    | 89       | -     | 86             | -      | 103            |

8




Figure 3. Estimated day of year (DOY) of leaf unfolding (LU) using the Max1 and IPG235 parameters for the site Reiffenhausen with local temperature measurements (a) and temperatures from the nearest DWD station Göttingen (b) for the years 2012 to 2014; same for Tharandt, local temperatures (c) and nearest climate station Wildacker (d); with observations. The lower subplots show long term estimates for DOY of LU using the DWD temperatures for Reiffenhausen (e) and for Tharandt (f), where also long term observations are available.

1 Table 3. Description of the eight performed model simulations. All model parameters are  
2 constant, except leaf area index (LAI), stomatal resistance (Rsc) and the date of leaf unfolding  
3 (LU) for the two parameter sets Max1 and IPG235.

| version                                | LAI   | Rsc                                       | LU                         |
|--|---|---|----------------------------|
| LAI2000 Rsc80                          | LAI-2000<br>measurements                                | minimum 80 s m <sup>-1</sup><br>(LAI > 1) | defined by measured<br>LAI |
| LAI2000 Rsc40                          | LAI-2000<br>measurements                                | minimum 40 s m <sup>-1</sup><br>(LAI > 1) | defined by measured<br>LAI |
| LI191SA Rsc80                          | LI-191 SA<br>measurements                               | minimum 80 s m <sup>-1</sup><br>(LAI > 1) | defined by measured<br>LAI |
| LI191SA Rsc40                          | LI-191 SA<br>measurements                               | minimum 40 s m <sup>-1</sup><br>(LAI > 1) | defined by measured<br>LAI |
| LAIstep Rsc40<br>Max1                  | step function<br>(6 in growing<br>season; else 1)       | minimum 40 s m <sup>-1</sup><br>(LAI > 1) | Max1 model                 |
| LAIstep Rsc40<br>IPG235                | step function<br>(6 in growing<br>season; else 1)       | minimum 40 s m <sup>-1</sup><br>(LAI > 1) | IPG235 model               |
| LAIadjusted<br>Rsc40adjusted<br>Max1   | course calibrated to<br>improve model fit<br>(max. = 6) | minimum 40 s m <sup>-1</sup><br>(LAI > 1) | Max1 model                 |
| LAIadjusted<br>Rsc40adjusted<br>IPG235 | course calibrated to<br>improve model fit<br>(max. = 6) | minimum 40 s m <sup>-1</sup><br>(LAI > 1) | IPG235 model               |





Figure 4. Leaf area index and stomatal resistance as parameterised from measurements. Leaf area index is used as measured, LAI2000 (red) and LI191SA (blue). Stomatal resistance is set to the measured minimum, i.e. 80  $\text{s m}^{-1}$  (dashed line) and to 40  $\text{s m}^{-1}$  (solid line) as leaf area index is larger than 1. Length of growing season is determined by the leaf area observations. Simulation results using the four combinations of leaf area index and stomatal resistance are shown as plant available soil water, calculated until 1 m soil depth and compared to values based on soil water content measurements.

1 Table 4. Statistical parameters for model evaluation in terms of the accuracy of simulated data  
2 compared to measured values. Nash-Sutcliffe efficiency criterion (NSC), percent bias  
3 (PBIAS), and ratio of the root mean square error to the standard deviation of measured data  
4 (RSR) are calculated from plant available soil water till 1 m soil depth as derived from model  
5 simulations and soil water content measurements for the period from April till December  
6 2013, to cover the period of most variability.

|   | NSC     | RSR        | PBIAS [%] |
|---|---------|------------|-----------|
| Recommended as satisfactory by<br>(Moriassi et al., 2007) | $> 0.5$ | $\leq 0.7$ | $\pm 25$  |
| LAI2000 Rsc80   | 0.44    | 0.75       | 19.2      |
| LAI2000 Rsc40   | 0.87    | 0.37       | 5.5       |
| LI191SA Rsc80   | 0.69    | 0.56       | 13.9      |
| LI191SA Rsc40   | 0.89    | 0.33       | 0.0       |
| LAIstep Rsc40 Max1  | 0.89    | 0.33       | -2.0      |
| LAIstep Rsc40 IPG235                                      | 0.89    | 0.33       | -1.7      |
| LAIadjusted Rsc40adjusted Max1                            | 0.90    | 0.31       | 1.3       |
| LAIadjusted Rsc40adjusted IPG235                          | 0.90    | 0.31       | 1.6       |

7



1  
2 Figure 5. Leaf area index and stomatal resistance parameterised as step function, using  
3 maximum and minimum values in the growing season 2013, respectively. LAI is set to 6,  
4 stomatal resistance is set to 40 s m<sup>-1</sup> when LAI is larger than 1. Leaf unfolding (LU) is  
5 determined by the dynamic phenology approach implemented in WaSim, using the Max1  
6 parameterisation and IPG235. Simulations results using the two combinations of LAI and  
7 stomatal resistance are shown as plant available soil water, calculated till 1 m soil depth and  
8 compared to values based on soil water content measurements.




Figure 6. Leaf area index and stomatal resistance as parameterised from adjusted values for 2013. Maximum of leaf area index is set to 6, minimum of stomatal resistance is set to 40 s m<sup>-1</sup>. Leaf unfolding (LU) is determined by the dynamic phenology approach implemented in WaSim, using the Max1 and IPG235 parameterisation. The annual course of leaf area index and stomatal resistance is orientated on measurements for leaf area and used as calibration parameter for stomatal resistance. Simulations results using the two combinations of leaf area index and stomatal resistance are shown as plant available soil water, calculated till 1 m soil depth and compared to values based on soil water content measurements.




Figure 7. Measured daily precipitation (a). Plant available water and ground water recharge as simulated for the step-function and adjusted course for leaf area index and stomatal resistance, using the leaf unfolding (LU) parameters Max1 (b) and IPG235 (c) for simulating LU. The parameterisation for poplar is equal for 2012 and 2013, i.e. the same vegetation hydrological modelled driven by different weather conditions, i.e. a drier year 2012 with an earlier dry period in May.



1 Table 5. Precipitation ( $\text{mm year}^{-1}$ ), total evapotranspiration (ETR) and ground water recharge  
2 (GWR) for the period 1969-2013. Simulations are shown for the step-function simulation and  
3 the adjusted course for leaf area index and stomatal resistance, using the leaf unfolding (LU)  
4 according to the Max1 and IPG235 parameter set for simulating LU. The parameterisation for  
5 poplar is equal to that derived for 2013 for the whole period, i.e. the same vegetation  
6 hydrological modelled driven by different weather conditions. Values are summed up for all  
7 years (1969-2013) of the period and for the 5 driest and 5 wettest years, respectively.

|               | ETR         | ETR        | ETR         | GWR         | GWR        | GWR         |
|---------------|-------------|------------|-------------|-------------|------------|-------------|
|               | (all years) | (5 driest) | (5 wettest) | (all years) | (5 driest) | (5 wettest) |
| precipitation | 676.7       | 500.4      | 896.6       | 676.7       | 500.4      | 896.6       |
| LAIstep       |             |            |             |             |            |             |
| Rsc40         | 527.8       | 487.1      | 533.6       | 79.7        | 23.2       | 140.2       |
| Max1          |             |            |             |             |            |             |
| LAIstep       |             |            |             |             |            |             |
| Rsc40         | 488.5       | 463.0      | 482.6       | 105.0       | 37.9       | 170.4       |
| IPG235        |             |            |             |             |            |             |
| LAIadjusted   |             |            |             |             |            |             |
| Rsc40adjusted | 484.1       | 460.8      | 477.9       | 107.1       | 41.3       | 173.9       |
| Max1          |             |            |             |             |            |             |
| LAIadjusted   |             |            |             |             |            |             |
| Rsc40adjusted | 425.9       | 424.4      | 411.7       | 144.7       | 68.1       | 232.9       |
| IPG235        |             |            |             |             |            |             |