

Replies to the editor' comments

Authors' replies are in BLUE color.

Comments to the Author:

Dear Dr Ke,

Your manuscript “Variability in snow cover phenology of China from 1952 to 2010” has again been reviewed by two of the original reviewers. One of the reviewers argues that most of his/her points were not handled and points to serious flaws in the methodology of the study, like definitions of SCDs which varied over time, the special introduction of thin SCDs for a part of the country, and differences between calendar year and snow year. This reviewer suggests rejection. Given the fact that the other reviewer suggests acceptance of the manuscript, I give you a last chance to substantially improve the paper and address all comments of reviewer #2. Please motivate your responses to reviewer #2 better, both for the former comments and the new comments. As such, I think that again major revision of the manuscript is required.

Replies: Thank you very much for your detail comments and give us a last chance for major revision. We know the most important concern of reviewer #2 is about the data problem. In this revision, we reprocessed all our data to satisfy the reviewer #2 former comments and new comments, especially his comments you mentioned: “definitions of SCDs which varied over time, the special introduction of thin SCDs for a part of the country, and differences between calendar year and snow year”. What made us very happy is that the results after the reprocessing and recalculation are very similar to what we had before.

In your answer to the main points and detailed comments, please indicate how comments have been handled exactly, indicating also whether text has been deleted and what the position of newly included text blocks is. I am looking forward to the new version of the paper.

Best regards,

Harrie-Jan Hendricks Franssen - editor -

Replies to the comments of Anonymous Referee #2 (8 Sep.)

This manuscript, although revised somewhat, is basically the same as the previous version. It still suffers from the main problems raised by previous reviewer #2. Authors almost denied all comments and suggestions by reviewer #2. They only accepted to change the concepts of “heavy-snow year” and “light-snow year”. Even for this change, they made it only after reviewer #3 raised the same problem. In the initial version of the replies to reviewer #2 on the HESS web site, they claimed their definitions were right. As they did not give reasonable reasons for their defense, I would retain previous comments.

Replies: Thank you very much for your detail comments. As for the acceptance to change the concepts of “heavy-snow year” and “light-snow year” after reviewer #3 raised the same problem, it was just a good time with good suggestion to revise it. Reviewer #2 raised this problem, but the reviewer#2 did not give constructive suggestion on how to improve it. Moreover, we must respond all comments within a week, at that time we also did not have better idea to revise them, so we claimed our definitions were right. After Reviewer #3’s comments with suggested words to replace the original ones, we then accepted the reviewer #3’s suggestion and changed them. We did not intend to offend the reviewer#2 and we actually thank a lot for the reviewer’s valuable time to review our paper and all reviewers to review all papers in the academic community, including ourselves as authors and also as reviewers to other authors’ papers.

In addition, I have some other comments.

1. Thin SCDs should be separated from “non-thin SCDs” (snowcover existing for a long time, e.g. 10 days). As previously pointed out, the snow cover observations are commonly conducted at 8:00 (Beijing time) in the morning, and most of the thin snow covers correspond to the snowing events in which snow exists only several hours. Suppose two sites: one site with 30 thin SCDs and the other site with 30 “non-thin SCDs”, both of them belong to the same group of 30 SCDs and are of the same climatic significance according to their study.

However, in reality they could be quite different in meteorology and climatology. This makes their results of climatic analysis questionable. Actually, this issue is not a new one. Authors should refer to the existing statistical methods of snow cover, such as those in Japan.

Replies: Although many references suggested to think about thin SCDs, especially in western China (we gave many references in the last version), the reviewer #2 insists on removing the thin SCDs for consideration, we felt we were pushed to do so. But to satisfy the reviewer's requirement, we did it in this revision. The revision resulted in a large drop in number of stations with more than ten annual average SCDs from 352 to 296. We reprocessed and recalculated all other parameters and data, and updated tables 1-4, figures 3-9. However, the main conclusions are similar as before, moreover newly recalculated data enhanced the conclusions. We are happy about this.

2. Consistency of the data. This is also raised in the previous comments. For the Specifications for Surface Meteorological Observations of China, there have been several versions. There are some differences in the criteria between the versions (e.g. minimum snow depth of 0.5 cm in the 2003 version, whereas it is 1.0 cm in the previous versions). This is actually not a matter of version, but relates to the reliability of their results. Obviously, the criteria change of minimum snow depth would bring more SCDs for a same snowcover with no climate change. Again, they did not address this issue, which also makes their results questionable.

Replies: According to this comment, we checked all data and found that only data in the period of Jan. 1. 2003 to Dec. 31. 2010 has the issue due to improved precision in recent years. In the revision, we deleted snow days for each station with snow depth between 0.5 cm and 0.9 cm, and keep snow days for each station with snow depth equal to or more than 1.0 cm, so that all SCDs have the same minimum snow depth 1.0 cm to keep the data consistency. However, these changes have less impact on SCD reduction for each station, because 8 year time series is very short relative to more than 50 year time series for most stations in China.

Because both the previous comment "thin SCD" and this comment "data consistency" are data problem, we process them at the same time. We removed not only all thin SCDs, but

also deleted all snow days for each station with snow depth between 0.5 cm and 0.9 cm from Jan. 1, 2003 to Dec. 31, 2010, and to keep the data consistency.

Finally, we obtained a new SCD time series data for each station, and reprocessed and recalculated all data, and updated tables 1-4, figures 3-9. However, the main conclusions are similar to the former ones, moreover newly recalculated data enhanced the conclusions.

3. Presentation is confusing. They initially define the “snow year” (Lines 149-151, author reponse-version3, the same below). However, in the subsequent parts, they did not use it, but use the calendar year. For example, the 2008 SCDs (Lines 350-359) should belong to the 2009 snow year according to their definition. Readers don’t know which are snow years and which are calendar years. In addition, in the Results section, there are many statements which are not directly related to their results (e.g. Lines 329-340; Lines 343-347; Lines 356-360) and thus bring confusions to readers.

Replies: The original sentences are “We define a snow year as the period from 1 September of the previous year to 31 August of the current year. For instance, September, October, and November 2009 are treated as the autumn season of snow year 2010, December 2009 and January and February 2010 as the winter season of snow year 2010, and March, April, and May 2010 as the spring season of snow year 2010.”

In our definition, the previous year has 4 months, and the current year has 8 months, and the current year is the year we called. In calendar year, 2008 snow disaster happened in January 2008, the occurring time of this event is in accordance to our snow year definition, it is 2008, not 2009.

The results based on our data processing are described according to snow year definition throughout the paper and we made sure they are consistent now. In the previous version, there is an exception, snow disaster years cited from references, different from this definition, possibly calendar year. In the revised manuscript we revised all of them in order to be same as our snow year definition.

4. Key words or terms problems. Some key words or terms are very difficult to understand. For example, “Areas with SCDs of 10–60 are called unstable snow areas with annual periodicity (there is definitely snow in every winter)” (Lines 258-259). Are the stable/unstable snow areas determined according to the number of SCDs? If so, I don’t think it is a good classification. And annual periodicity means there is definitely snow in every winter??? Also, it is unclear with the concept of “year with a positive/negative SCD anomaly”. Why 70% and 30% (Lines 318-321)? Could “30%” be “the other 30%” (line 319)?

Replies: As for “Areas with SCDs of 10–60 are called unstable snow areas with annual periodicity (there is definitely snow in every winter)”, we cite some published references (Li, 1990) to support it, it is not our definition. “the stable/unstable snow areas determined according to the number of SCDs”, we also cited some references to support it (Li and Mi, 1983; Li, 1990). Moreover, they are statistical results in the climatology for a long time, in our opinion, the stable/unstable snow areas can be determined according to the number of SCDs. According to references and the descriptions in our paper, annual periodicity means there is definitely snow in every winter.

Originally, we called “heavy-snow year” and “light-snow year”. The concept of “year with a positive/negative SCD anomaly” is suggested by Reviewer #3, and we accepted his suggestion to revise the paper. Reviewer #2 also positively mentioned it in the first paragraph of this comment.

“70% and 30% (Lines 318-321)” are from our definition (for a given year, if both (1) 70% of the stations have a positive (negative) anomaly and (2) 30% of the stations have an SCD larger (smaller) than the mean \pm one standard deviation (1SD), it is regarded as a year with a positive (negative) SCD anomaly). It is a judge or speculation from a statistical significance to decide a year with a positive (negative) SCD anomaly. This definition is based on large volume data analysis (672 stations from 1952 to 2010), and the results of statistical analysis are reliable. “70%” refer to percentage of stations in all 672 stations, and “30%” also refer to percentage of stations in all 672 stations, it is not “the other 30%” mentioned by Reviewer #2.

There is “and” between “70% and 30%”, therefore a given year must meet the two requirements at the same time.

5. English should be improved. For example, “We define a snow year as the period from 1 September of the previous year to 31 August of the current year (Lines 150-151).” Is this sentence correct?

Replies: We invite an expert from the language editing company “American Journal Expert” whose native language is English to proofread this paper. The sentence is revised as “A snow year is defined at the time period from September 1 of the previous year to August 31 of the current year”

Replies to the comments of Anonymous Referee #2 (25 May)

Authors’ replies are in BLUE color.

Interactive comment on “Variability in snow cover phenology in China from 1952 to 2010” by C. Q. Ke et al.

Anonymous Referee #2

Received and published: 25 May 2015

This manuscript presents the spatio-temporal snow cover data of China on the timing (snow cover onset and end dates: SCOD and SCED) and duration (snow cover days: SCD) and analyses their relationships with air temperature and arctic oscillation. While substantial datasets were used, the data were not well interpreted and analysed, and no significant conclusions were drawn.

The results and conclusions are even suspicious considering the way they treated the data.

I suggest to reject and resubmit.

Given this recommendation, I would only give some major comments.

1. The data. “According to the Specifications for Surface Meteorological Observations (China

Meteorological Administration, 2003), an SCD is defined as a day when the snow cover in the area fulfils two requirements: at least half of the observation field is covered by snow, and the minimum snow depth is 1 cm. For any day with at least half of the observation field covered by snow but with snow depth of less than 1 cm, the snow depth is denoted as 0, i.e. a thin SCD.” (P4475: Lines 19-24). “: : in western China, station density is low, and the observation history is relatively short... If all stations with short time series are eliminated, and thin SCDs are not taken into account, the spatial representativeness of the dataset would be a problem. Therefore, a time series of at least 30 years is included in this study, including those thin SCDs.” (P4476: Lines 3-8).

In my opinion, however, including those thin SCDs is more problematic than excluding them. As far as I know, the snow cover observations are commonly conducted at 8:00 (Beijing time) in the morning, and most of the thin snow covers correspond to the snowing events in which snow exists only several hours. This is also the case for many SCDs with snow cover depths not less than 1 cm.

Replies: Although many references suggest to think about thin SCDs, especially in western China (we gave many references in the last version), Reviewer #2 insist on removing the thin SCDs, we accept this comments. Now we removed all thin SCDs, and SCDs of some stations in the west China decreased evidently. In particular, number of stations with more than ten annual average SCDs decreased from 352 to 296. We reprocessed and recalculated all data, and updated tables 1-4, figures 3-9. However, the main conclusions are similar to the former ones, moreover newly recalculated data enhanced the conclusions.

Except for several small regions, there have been not much snow in China during recent three decades. In this sense, there have been very few snow covers, but several snowfalls per year in a considerably large area of China (south, central and north China, and even a large area of western China) in recent _30 years. Therefore, for these areas, it may make more sense to conduct statistics of precipitation phase rather than the SCDs.

Replies: Unfortunately, analyzing precipitation is out of the scope of this paper.

2. Some basic information on the spatio-temporal distributions of snow cover water

equivalent or snow cover depth should be provided. Readers need these information for judgements.

Replies: In this paper, we only investigate several snow variables (snow cover days, onset and end date of snow cover), their spatiotemporal evolution, extreme years and trends, and also their relations to temperature variables and climate patterns (Arctic Oscillation, AO). It already has 36 pages, including 4 tables and 9 figures, longer enough, therefore we do not provide snow water equivalent or snow depth result. There are many studies aiming at only SCDs. The spatio-temporal distributions of snow cover water equivalent or snow cover depth will be provided in another paper in the future.

3. Analysis. Analysis is lacking on the climatic and physical interpretations/processes of the statistical results throughout the manuscript.

Replies: We analyzed the climatology of several snow variables (snow cover days, onset and end date of snow cover), their spatiotemporal evolution, extreme years and trends from 1952-2010 in China for a large number of stations. The relationships among SCDs and temperature and Arctic Oscillation are the climatic and physical interpretation in our view. All statistic results are conducted significant test, we only think about the results passed 90% or 95% significant test, and explanation is given in the manuscript.

4. Definition and analysis of heavy-snow and light-snow years (Sections 3.1.2 and 3.2.1). A heavy-snow year or a light-snow year was determined in terms of the relative time duration of SCDs of a region. This is logically problematic. Authors should know that, for a given station, a longer period of SCDs does not necessarily mean a year of more snowfall.

Replies: We think about comments from Reviewer #2 and Reviewer #3 at the same time, and accepted constructive suggestion from Reviewer #3, changed as “year with a positive/negative SCD anomaly”.

5. Consistency of data. As far as I know, for the Specifications for Surface Meteorological Observations of China, there have been several versions (1951?, 1980, 2003 and 2007?). There are some differences in the criteria between the versions (e.g. minimum snow depth of

0.5 cm in the 2007 version?). This should be addressed.

Replies: According to this comment, we checked all data from Jan. 1. 2003 to Dec. 31. 2010, and deleted snow days for each station with snow depth between 0.5 cm and 0.9 cm, and keep snow days for each station with snow depth equal to or more than 1.0 cm, so that all SCDs have the same minimum snow depth 1.0 cm and kept the data consistency. However, these changes have less impact on SCD reduction for each station, because 8 year time series is very short relative to more than 50 year time series for most stations in China.

Because both the first comment “thin SCD” and this comment “data consistency” are data problem, we process them at the same time. We removed not only all thin SCDs, but also deleted all snow days for each station with snow depth between 0.5 cm and 0.9 cm from Jan. 1. 2003 to Dec. 31. 2010, and kept the data consistency.

Finally, we obtained a new SCD time series data for each station, and reprocessed and recalculated all data, and updated tables 1-4, figures 3-9. However, the main conclusions are similar to the former ones, moreover newly recalculated data enhanced the conclusions.

6. The tilte. Authors used the word “phenology”. However, except the SCDs, SCODs and SCEDs, they did not analyze any of the important snow properties such as density. I would suggest not to use the word.

Replies: Yes, we did not analyze any of the important snow properties such as density, as well as snow depth, snow water equivalent. Because other experts used this word when they conducted the same research, we cited the following paper. In our opinion, the word “phenology” is correct.

Peng, S., Piao, S., Ciais, P., Friedlingstein, P., Zhou, L. and Wang, T.: Change in snow phenology and its potential feedback to temperature in the Northern Hemisphere over the last three decades, *Environ. Res. Lett.*, 8, 014008, 2013.

1 **Variability in snow cover phenology in China from 1952**
2 **to 2010**

3
4 Chang-Qing Ke^{1,2}, Xiu-Cang Li^{3,4}, Hongjie Xie⁵, Dong-Hui Ma^{1,6}, Xun Liu^{1,2} and
5 Cheng Kou^{1,2}

6 1. *Jiangsu Provincial Key Laboratory of Geographic Information Science and Technology,*
7 *Nanjing University, Nanjing, 210023, China.*

8 2. *Key Laboratory for Satellite Mapping Technology and Applications of State*
9 *Administration of Surveying, Mapping and Geoinformation of China, Nanjing University,*
10 *Nanjing, 210023, China.*

11 3. *National Climate Center, China Meteorological Administration, Beijing 100081, China.*

12 4. *Collaborative Innovation Center on Forecast and Evaluation of Meteorological*
13 *Disasters , Faculty of Geography and Remote Sensing, Nanjing University of*
14 *Information Science & Technology, Nanjing, 210044, China.*

15 5. *Department of Geological Sciences, University of Texas at San Antonio, Texas 78249,*
16 *USA.*

17 6. *Collaborative Innovation Center of South China Sea Studies, Nanjing, 210023, China*

18
19 Correspondence to: C. Q. Ke (kecq@nju.edu.cn)

20 Tel: 0086-25-89685860

21 Fax: 0086-25-83592686

22 | **Abstract** Daily snow observation data from 672 stations, particularly the [352–296](#)
23 | stations with over ten annual mean snow cover days (SCD), during 1952–2010 in China,
24 | are used in this study. We first examine spatiotemporal variations and trends of SCD,
25 | snow cover onset date (SCOD), and snow cover end date (SCED). We then investigate
26 | SCD relationships with number of days with temperature below 0°C (TBZD), mean air
27 | temperature (MAT), and Arctic Oscillation (AO) index, the latter two being constrained
28 | to the snow season of each snow year. The results indicate that years with positive SCD
29 | anomaly for the entire country include 1955, 1957, 1964, and 2010, and years with
30 | negative SCD anomaly include 1953, 1965, 1999, 2002, and 2009. The reduced TBZD
31 | and increased MAT are the main reasons for the overall late SCOD and early SCED
32 | since 1952, although it is not necessary for one station to experience both significantly
33 | late SCOD and early SCED. This explains why only [4512%](#) of the stations show
34 | significant shortening of SCD, while 75% of the stations show no significant change in
35 | the SCD trends. This differs with the overall shortening of the snow period in the
36 | Northern Hemisphere previously reported. Our analyses indicate that the SCD
37 | distribution pattern and trends in China are very complex and are not controlled by any
38 | single climate variable examined (i.e. TBZD, MAT, or AO), but a combination of
39 | multiple variables. It is found that the AO has the maximum impact on the SCD
40 | shortening trends in [the Shandong Peninsula, Changbai Mountains, Xiaoxingganling](#)
41 | and North Xinjiang, while the combined TBZD and MAT have the maximum impact
42 | on the SCD shortening trends in the Loess Plateau, [Xiaoxingganling, and Sanjiang](#)
43 | [Plain Tibetan Plateau, and Northeast Plain](#).

44 **Keywords:** snow cover day; snow cover onset date; snow cover end date;
45 spatiotemporal variation; trend; days with temperature below 0°C; Arctic Oscillation

46

47 **Abbreviations:**

48 Snow Cover Day (SCD)

49 Snow Cover Onset Date (SCOD)

50 Snow Cover End Date (SCED)

51 Days with Temperature Below 0°C (TBZD)

52 Mean Air Temperature (MAT)

53 Arctic Oscillation (AO)

54

55 **1 Introduction**

56 Snow has a profound impact on the surficial and atmospheric thermal conditions,
57 and is very sensitive to climatic and environmental changes, because of its high
58 reflectivity, low thermal conductivity, and hydrological effects via snowmelt ([Barnett et
59 al., 1989](#); [Groisman et al., 1994](#)). The extent of snow cover in the Northern Hemisphere
60 decreased significantly over the past decades because of global warming ([Robinson and
61 Dewey 1990](#); [Brown and Robinson 2011](#)). Snow cover showed the largest decrease in
62 the spring, and the decrease rate increased for higher latitudes in response to larger
63 albedo feedback ([Déry and Brown, 2007](#)). In North America, snow depth in central

64 Canada showed the greatest decrease (Dyer and Mote, 2006), and snowpack in the
65 Rocky Mountains in the U.S. declined (Pederson et al., 2013). However, *in situ* data
66 showed a significant increase in snow accumulation in winter but a shorter snowmelt
67 season over Eurasia (Bulygina et al., 2009). Decreases in snow pack have also been
68 found in the European Alps in the last 20 years of the 20th century (Scherrer et al.,
69 2004), but a very long time series of snow pack suggests large decadal variability and
70 overall weak long-term trends only (Scherrer et al., 2013). Meteorological data
71 indicated that the snow cover over northwest China exhibited a weak upward trend in
72 snow depth (Qin et al., 2006), with large but the spatiotemporal variations were large
73 (Ke et al., 2009; Ma and Qin 2012). Simulation experiments using climate models
74 indicated that, with continuing global warming, the snow cover variation in China
75 would show more variations differences and uncertainties in space and time than ever
76 before (Shi et al., 2011; Ji and Kang 2013). Spatiotemporal variations of snow cover are
77 also manifested as snowstorms or blizzards, particularly, excessive snowfall over a
78 short time duration (Bolsenga and Norton, 1992; Liang et al., 2008; Gao, 2009; Wang
79 et al., 2013; Llasat et al., 2014).

80 Snow cover day (SCD) is an important index that represents the environmental
81 features of climate (Ye and Ellison 2003; Scherrer et al., 2004), and is directly related
82 to the radiation and heat balance of the Earth-atmosphere system. The SCD varies in
83 space and time and contributes to climate change over short time scales (Zhang, 2005),
84 especially in the Northern Hemisphere. Bulygina et al. (2009) investigated the linear
85 trends of SCD observed at 820 stations from 1966 to 2007, and indicated that the

86 duration of snow cover decreased in the northern regions of European Russia and in the
87 mountainous regions of southern Siberia, while it increased in Yakutia and the Far East.
88 Peng et al. (2013) analysed trends in the snow cover onset date (SCOD) and snow
89 cover end date (SCED) in relation to temperature over the past 27 years (1980–2006)
90 from over 636 meteorological stations in the Northern Hemisphere. They found that the
91 SCED remained stable over North America, whereas there was an early SCED over
92 Eurasia. Satellite-derived snow data indicated that the average snow season duration
93 over the Northern Hemisphere decreased at a rate of 5.3 days per decade between
94 1972/73 and 2007/08 (Choi et al., 2010). Their results also showed that a major change
95 in the trend of snow duration occurred in the late 1980s, especially in the Western
96 Europe, central and East Asia, and mountainous regions in western United States.

97 There are large spatiotemporal differences in the SCD in China (Wang and Li,
98 2012). Analysis of 40 meteorological stations from 1971 to 2010 indicated that the
99 SCD had a significant decreasing trend in the western and south-eastern Tibetan
100 Plateau, with the largest decline observed in Nielamu, reaching 9.2 days per decade
101 (Tang et al., 2012). Data analysis also indicated that the SCD had a linear decreasing
102 trend at most stations in the Hetao region and its vicinity (Xi et al., 2009). However,
103 analysis of meteorological station data in Xinjiang showed that the SCD had a slight
104 increasing trend, occurring mainly in 1960–1980 (Wang et al., 2009b). Li et al. (2009)
105 analysed meteorological data from 80 stations in Heilongjiang Province, Northeast
106 China. Their results showed that the snow cover duration shortened, because of both the
107 late SCOD (by 1.9 days per decade) and early SCED (by 1.6 days per decade), which

108 took place mainly in the lower altitude plains.

109 The SCD is sensitive to local winter temperature and precipitation, latitude ([Hantel](#)
110 [et al., 2000](#); [Wang et al., 2009a](#); [Serquet et al., 2011](#); [Morán-Tejeda et al., 2013](#)), and
111 altitudinal gradient and terrain roughness ([Lehning et al., 2011](#); [Ke and Liu, 2014](#)).
112 Essentially, the SCD variation is mainly attributed to large-scale atmospheric
113 circulation or climatic forcing ([Beniston, 1997](#); [Scherrer and Appenzeller, 2006](#); [Ma](#)
114 [and Qin, 2012](#); [Birsan and Dumitrescu, 2014](#)), such as monsoons, El Niño/Southern
115 Oscillation (ENSO), North Atlantic Oscillation (NAO), and Arctic Oscillation (AO).
116 [Xu et al. \(2010\)](#) investigated the relationship between the SCD and monsoon index in
117 the Tibetan Plateau and indicated their great spatial differences. As an index of the
118 dominant pattern of non-seasonal sea-level pressure variations, the AO shows a large
119 impact on the winter weather patterns of the Northern Hemisphere ([Thompson and](#)
120 [Wallace, 1998](#); [Thompson et al., 2000](#); [Gong et al., 2001](#); [Wu and Wang, 2002](#); [Jeong](#)
121 [and Ho, 2005](#)). The inter-annual variation of winter extreme cold days in the northern
122 part of eastern China is closely linked to the AO ([Chen et al., 2013](#)). Certainly, the AO
123 plays an important role in the SCD variation. An increase in the SCD before 1990 and a
124 decrease after 1990 have been reported in the Tibetan Plateau, and snow duration has
125 positive correlations with the winter AO index ([You et al., 2011](#)), and a significant
126 correlation between the AO and snowfall over the Tibetan Plateau on inter-decadal
127 timescale was also reported by [Lü et al. \(2008\)](#).

128 The focus of this study is the variability in the snow cover phenology in China. A
129 longer time series of daily observations of snow cover is used for these spatial and

130 temporal analyses. We first characterize the spatial patterns of change in the SCD,
131 SCOD, and SCED in different regions of China; we then examine the sensitivity of
132 SCD to the number of days with temperature below 0°C (TBZD), the mean air
133 temperature (MAT), and the Arctic Oscillation (AO) index during the snow season
134 (between SCOD and SCED).

135 **2 Data and methods**

136 **2.1 Data**

137 We use daily snow cover and temperature data in China from [the](#) 1 September
138 1951 to [the](#) 31 August 2010, provided by the National Meteorological Information
139 Centre of China Meteorological Administration (CMA). According to the
140 Specifications for Surface Meteorological Observations ([China Meteorological
141 Administration, 2003](#)), an SCD is defined as a day when the snow cover in the area
142 meets the following requirement: at least half of the observation field is covered by
143 snow. For any day with at least half of the observation field covered by snow, snow
144 depth is recorded as a rounded-up integer. [For example, a normal SCD is recorded if
145 the snow depth is equal to or more than or equal to 1.00.5 cm \(measured with a ruler\),
146 or a thin SCD - i.e. a normal SCD, whereas the snow depth is denoted as 0 if the snow
147 depth is it is less than 1.0.5 cm, i.e. a thin SCD. A snow year is defined at the time
148 period from September 1 of the previous year to August 31 of the current year. We
149 define a snow year as the period from 1 September of the previous year to 31 August of
150 the current year.](#) For instance, September, October, and November 2009 are treated as
151 the autumn season of snow year 2010, December 2009 and January and February 2010

152 as the winter season of snow year 2010, and March, April, and May 2010 as the spring
153 season of snow year 2010.

154 Station density is high in eastern China, where the observational data for most
155 stations are complete, with relatively long histories (as long as 59 years). ~~Because of~~
156 ~~topography and climate conditions, the discontinuous nature of snowfall is obvious in~~
157 ~~western China, especially in the Tibetan Plateau, with patchy snow cover, and many~~
158 ~~thin SCDs in these station records (Ke and Li, 1998). At the same time, in western~~
159 ~~China, while~~ station density is low in western China, and the observation history is
160 relatively short, although two of the three major snow regions are located in western
161 China. If all stations with short time series are eliminated, ~~and thin SCDs are not taken~~
162 ~~into account~~, the spatial representativeness of the dataset would be a problem.
163 Therefore, a time series of at least 30 years is included in this study, ~~including those~~
164 ~~thin SCDs. Totally, there are 722 stations in the original dataset.~~

165 Because of topography and climate conditions, the discontinuous nature of
166 snowfall is obvious in western China, especially in the Tibetan Plateau, with patchy
167 snow cover, and there are many thin SCD records (Ke and Li, 1998). However, in order
168 to enhance data reliability, thin SCDs in the original dataset are not taken into account
169 in this paper according to the previous studies (An et al., 2009; Wang and Li, 2012).

170 Totally, there are 722 stations in the original dataset. Since station relocation and
171 changes in the ambient environment could cause inconsistencies in the recorded data,
172 we implement strict quality controls (such as inspection for logic, consistency, and
173 uniformity) on the observational datasets in order to reduce errors (Ren et al., 2005).

174 The standard normal homogeneity test ([Alexandersson and Moberg, 1997](#)) at the 95%
175 confidence level is applied to the daily SCD and temperature series data in order to
176 identify possible breakpoints. Time series gap filling is performed after all
177 inhomogeneities are eliminated, using nearest neighbour interpolation. After being
178 processed as mentioned above, the 672 stations with annual mean SCDs greater than
179 1.0 (day) are finally selected for subsequent investigation ([Fig. 1](#)).

180 The observation period for each station is different, varying between 59 years
181 (1951/1952–2009/2010) and 30 years (1980/1981–2009/2010). Overall, 588 stations
182 have observation records between 50 and 59 years, 47 stations between 40 and 49 years,
183 and 37 stations between 30 and 39 years ([Fig. 2](#)). Most of the stations with observation
184 records of less than 50 years are located in remote or high elevation areas. All 672
185 stations are used to analyse the spatiotemporal distribution of SCD in China, while only
186 [352–296](#) stations with more than ten annual mean SCDs are used to study the changes
187 of SCOD, SCED, and SCD relationships with TBZD, MAT, and the AO index.

188 The daily AO index constructed by projecting the daily (00Z) 1,000 mb height
189 anomalies poleward of 20°N from
190 http://www.cpc.ncep.noaa.gov/products/precip/CWlink/daily_ao_index/ao.shtml, is
191 used ~~in this paper~~. A positive (negative) AO index corresponds to low (high) pressure
192 anomalies throughout the polar region and high (low) pressure anomalies across the
193 subtropical and mid-latitudes ([Peings et al., 2013](#)). We average the daily AO indexes
194 during the snow season of each station as the AO index of the [snow](#) year. A time series
195 of AO indexes from 1952 to 2010, for each of the [352–296](#) stations, is then constructed.

196 A digital elevation model (DEM) ~~from~~[according to](http://srtm.csi.cgiar.org) the Shuttle Radar Topographic
197 Mission (SRTM, <http://srtm.csi.cgiar.org>) of the National Aeronautics and Space
198 Administration (NASA) with a resolution of 90 m and the administration map of China
199 are used as the base map.

200 **2.2 Methods**

201 We apply Mann–Kendall (MK) test to analyse the trends of SCD, SCOD, and
202 SCED. The MK test is an effective tool to extract the trends of time series, and is
203 widely applied to the analysis of climate series (Marty, 2008). The MK test is
204 characterized as being more objective, since it is a non-parametric test. A positive
205 standardized MK statistic value indicates an upward or increasing trend, while a
206 negative value demonstrates a downward or decreasing trend. Confidence levels of
207 90% and 95% are taken as thresholds to classify the significance of positive and
208 negative trends of SCD, SCOD, and SCED.

209 At the same time, if SCD, SCOD, or SCED at one climate station has significant
210 MK trend (above 90%), their linear regression analyses are performed against time,
211 respectively. The slopes of the regressions represent the changing trends and are
212 expressed in days per decade. The statistical significance of the slope for each of the
213 linear regressions is assessed by the Student's t test (two-tailed test of the Student t
214 distribution), and only confidence levels above 90% are considered.

215 Correlation analysis is used to examine the SCD relationships with the TBZD,
216 MAT, and the AO index, and the Pearson product-moment correlation coefficients
217 (PPMCC) have been calculated. The PPMCC is a widely used estimator for describing

218 the spatial dependence of rainfall processes, and it indicates the strength of the linear
219 covariance between two variables (Habib et al., 2001; Ciach and Krajewski, 2006). The
220 correlation coefficient can be defined as the covariance of the two variables (X , Y)
221 divided by the product of their standard deviations, giving a value between +1 and -1
222 inclusive, where 1 is total positive correlation, 0 is no correlation, and -1 is total
223 negative correlation. The statistical significance of the correlation coefficients is
224 calculated using the Student's t test, and only confidence levels above 90% are
225 considered in our analysis.

226 The spatial distribution of SCD, SCOD, and SCED, and their calculated results,
227 are spatially interpolated by applying the universal kriging method (assuming the data
228 is normally distributed). The universal kriging model is capable of simultaneously
229 treating multiple variables and their cross-covariance, and has been successfully applied
230 to spatial data interpolation (Kyriakidis and Goodchild, 2006). All mean errors are near
231 zero, all average standard errors are close to the corresponding root mean squared
232 errors, and all root mean squared standardized errors are close to 1 (Table 1). This fact
233 indicates that prediction errors are unbiased and valid, except for slightly overestimated
234 [coefficients of variation \(CV\)](#) and slightly underestimated SCD in 2002. Overall, the
235 interpolation results have fewer errors and are acceptable.

236 **3 Results**

237 **3.1 Spatiotemporal variations of SCD**

238 **3.1.1 Spatial distribution of SCD**

239 The analysis of observations from 672 stations indicates that there are three major

240 stable snow regions with more than 60 annual mean SCDs (Li, 1990): Northeast China,
241 North Xinjiang, and the Tibetan Plateau, with Northeast China being the largest of the
242 three (Fig. 3a). In the Daxingganling, Xiaoxingganling, and Changbai Mountains of
243 Northeast China, there are more than 90 annual mean SCDs, corresponding to a
244 relatively long snow season. The longest annual mean SCDs, ~~169-163~~ days, is at Arxan
245 Station (in the Daxinganling Mountains) in Inner Mongolia. In North Xinjiang, the
246 SCDs are relatively long in the Tianshan and Altun Mountains, followed by the Junggar
247 Basin. The annual mean SCDs in the Himalayas, Nyainqentanglha, Tanggula
248 Mountains, Bayan Har Mountains, Anemaqen Mountains, and Qilian Mountains of the
249 Tibetan Plateau are relatively long, although most of these ~~areas-regions~~ have less than
250 60 annual SCDs. The Tibetan Plateau has a high elevation, a cold climate, and many
251 glaciers, but its mean SCD is not as large as that of the other two stable snow regions.

252 Areas with SCDs of 10–60 ~~isare~~ called unstable snow ~~areas-regions~~ with annual
253 periodicity (~~there is~~ definitely ~~with snow cover in~~ every winter) (Li, 1990). It ~~;~~
254 ~~includ~~esing the peripheral parts of the three major stable snow regions, ~~and the~~ Loess
255 Plateau, Northeast Plain, North China Plain, Shandong Peninsula, and ~~areas-regions~~ in
256 north of the Qinling-Huaihe line (along the Qinling Mountains and Huaihe River to the
257 east). Areas with SCDs of 1–10 ~~isare~~ called unstable snow ~~areas-region~~ without annual
258 periodicity (the mountainous ~~areas-regions~~ are excluded, ~~not every winter there is snow,~~
259 ~~especially in a warm winter~~) (Li, 1990). It ~~;~~ includ~~esing~~ the ~~Tarim Basin,~~ Qaidam Basin,
260 Badain Jaran Desert, the peripheral parts of Sichuan Basin, the northeast part of the
261 Yungui Plateau, and the middle and lower Yangtze River Plain. Areas with occasional

262 snow and mean annual SCD of less than 1.0 (day) are distributed north of the Sichuan
263 Basin and in the belt along Kunming, ~~the~~ Nanling Mountains, and Fuzhou (approximate
264 latitude of 25°N). Because of the latitude or local climate and terrain, there is no snow
265 in the Taklimakan Desert, Turpan Basin, the Yangtze River Valley in the Sichuan Basin,
266 the southern parts of Yunnan, Guangxi, Guangdong and Fujian, and on the Hainan
267 Island.

268 The spatial distribution pattern of SCD based on climate data with longer time
269 series is similar to previous studies (Li and Mi, 1983; Li, 1990; Liu et al., 2012; Wang
270 et al., 2009a; Wang and Li, 2012). ~~The s~~ Snow distribution is closely linked to latitude
271 and elevation, and is generally consistent with the climate zones (Lehning et al., 2011;
272 Ke and Liu, 2014). ~~The higher the latitude, the lower the temperature and the more~~
273 ~~SCDs there are. Therefore, t~~ There are relatively more SCDs in Northeast China and
274 North Xinjiang, and fewer SCDs to the south (Fig. 3a). In the Tibetan Plateau, located
275 in south-western China, the elevation is higher than eastern areas at the same latitude,
276 and the SCDs are greater than in eastern China (Tang et al., 2012). The amount of
277 precipitation also plays a critical role in determining the SCD (Hantel et al., 2000). In
278 the north-eastern coastal areas of China, which are affected considerably by ~~the~~ ocean,
279 there is much precipitation. In North Xinjiang, which has a typical continental (inland)
280 climate, the precipitation is less than in Northeast China, and there are more SCDs in
281 the north of Northeast China than in North Xinjiang (Dong et al., 2004; Wang et al.,
282 2009b). Moreover, the local topography has a relatively large impact on the SCD
283 (Lehning et al., 2011). The Tarim Basin is located inland, with relatively little

284 precipitation, thus snowfall there is extremely rare except ~~for~~in the surrounding
285 mountains (Li, 1993). The Sichuan Basin is surrounded by high mountains, therefore
286 situated in the precipitation shadow in winter, resulting in fewer SCDs (Li and Mi,
287 1983; Li, 1990).

288 The three major stable snow regions, Northeast China, North Xinjiang, and the
289 eastern Tibetan Plateau, have smaller ~~coefficients of variation~~ (CV) in the SCD (Fig.
290 3b). Nevertheless, the SCDs in arid or semi-arid areasregions, such as South Xinjiang,
291 the northern and south-western Tibetan Plateau, and central and western Inner
292 Mongolia, have large fluctuation because there is little precipitation during the cold
293 seasons, and certainly little snowfall and large CVs of SCD. In particular, the
294 Taklimakan Desert in the Tarim Basin is an extremely arid region, with only occasional
295 snowfall. Therefore, it has a very large range of SCD fluctuations. Additionally, the
296 middle and lower Yangtze River Plain also has large SCD fluctuations because of
297 warm-temperate or sub-tropic climate with short winter and little snowfall. Generally,
298 the smaller the SCD, the larger the CV (Wang et al., 2009a). This is consistent with
299 other climate variables, such as precipitation (Yang et al., 2015).

300 **3.1.2 Temporal variations of SCD**

301 Seasonal variation of SCD is primarily controlled by temperature and precipitation
302 (Hantel et al., 2000; Scherrer et al., 2004; Liu et al., 2012). In North Xinjiang and
303 Northeast China, snow is primarily concentrated in the winter (Fig. 4). In these regions,
304 the SCD exhibits a 'single-peak' distribution. In the Tibetan Plateau, however, the
305 seasonal variation of SCD is slightly different, i.e. more snow in the spring and autumn

306 combined than in the winter. The mean temperature and precipitation at Dangxiong
307 station (30°29' N, 91°06'E, 4200.0 m) in winter are -7.73° C and 7.92 mm, respectively,
308 and those at Qingshuihe station (33°48' N, 97°08'E, 4415.4 m) are -15.8° C and 16.3
309 mm, respectively. It is too cold and dry to produce enough snow in the Tibetan Plateau
310 (Hu and Liang, 2014)

311 The temporal variation of SCD shows very large differences from one year to
312 another. We define a year with a positive (negative) SCD anomaly in the following way:
313 for a given year, if 70% of the stations have a positive (negative) anomaly and 30% of
314 the stations have an SCD larger (smaller) than the mean \pm one standard deviation
315 (1SD), it is regarded as a year with a positive (negative) SCD anomaly. The years with
316 a positive SCD anomaly in China are 1955, 1957, 1964, and 2010 (Table 2). Moreover,
317 the stations with SCDs larger than the mean + 2SD account for 29.25% and 26% of all
318 stations in 1955 and 1957, respectively, and these two years are considered as years
319 with an extremely positive SCD anomaly. In 1957, there was an almost nationwide
320 positive SCD anomaly except for North Xinjiang (Fig. 5a). This 1957 event had a great
321 impact on agriculture, natural ecology, and social-economic systems, and resulted in a
322 tremendous disaster (Hao et al., 2002). The year 2010 was also a year with a positive
323 SCD anomaly in China. At the same time, blizzards occurred in North America and
324 Europe (including Spain) (Llasat et al., 2014). Globally, an unusual cold weather
325 pattern caused by high pressure (the AO) brought cold, moist air from the north. Many
326 parts of the Northern Hemisphere experienced heavy snowfall and record-low
327 temperatures, leading to, among other things, a number of deaths, widespread transport

328 disruption, and power failures
329 (http://en.wikipedia.org/wiki/Winter_of_2009–10_in_Europe, [http://en.wikipedia.org](http://en.wikipedia.org/wiki/February_9–10,_2010_North_American_blizzard)
330 [/wiki/February_9–10,_2010_North_American_blizzard](http://en.wikipedia.org/wiki/February_9–10,_2010_North_American_blizzard)). The blizzards across the
331 Texas and Oklahoma panhandles in 1957 (Bolsenga and Norton, 1992; Changnon and
332 | Changnon, 2006) and ~~aeross~~ the east coast in 2010 were also recorded as the biggest
333 snowstorms of the United States from 1888 to the present
334 (<http://www.crh.noaa.gov/mkx/?n=biggestsnowstorms-us>).

335 Years with a negative SCD anomaly include 1953, 1965, 1999, 2002, and 2009
336 (Table 2). If there is too little snowfall in a specific year, a drought is possible. Drought
337 resulting from little snowfall in the cold season is a slow process and can sometimes
338 cause disasters. For example, East China displayed an apparent negative SCD anomaly
339 in 2002 (Fig. 5b), and had very little snowfall, leading to an extreme winter drought in
340 Northeast China, where snowfall is the primary form of winter precipitation (Fang et al.,
341 | 2014).

342 Because of different atmospheric circulation backgrounds, vapour sources, and
343 topographic conditions in different regions of China, there are great differences in the
344 SCD even in one year. For example, in 2008, there were more SCDs and longer snow
345 duration in the Yangtze River Basin, North China, and the Tianshan Mountains in
346 | Xinjiang (Fig. 5c), especially in the Yangtze River Basin, where large snowfall was
347 normally not observed. However, four episodes of severe and persistent snow, extreme
348 | low temperatures, and freezing weather occurred in ~~early~~ 2008, leading to a large-scale
349 catastrophe in this region where there were no mitigation measures for this type of a

350 disaster (Gao, 2009). As reported by the Ministry of Civil Affairs of China, the 2008
351 snow disaster killed 107 people and caused losses of US\$ 15.45 billion. Both the SCDs
352 and scale of economic damage broke records from the past five decades (Wang et al.,
353 2008). On the contrary, in the same year (2008), there was no snow disaster in North
354 Xinjiang, the Tibetan Plateau, and Pan-Bohai Bay region. Moreover, Northeast China
355 had an apparent negative SCD anomaly (Fig. 5c).

356 There are great differences in the temporal variations of SCD even in the three
357 major stable snow regions. If we redefine a year with a positive (negative) SCD
358 anomaly, using ~~at~~ much higher standard that 80% of stations should have a positive
359 (negative) anomaly and 40% of stations should have an SCD larger (smaller) than the
360 mean +/- 1SD. It is found that 1957, 1973, and 2010 are years with a positive SCD
361 anomaly in Northeast China, while 1959, 1963, 1967, 1998, 2002, and 2008 are years
362 with a negative SCD anomaly there (Table 3, Fig. 5a–c). Years with a positive SCD
363 anomaly in North Xinjiang include ~~1959~~, 1960, 1977, 1980, 1988, 1994, and 2010, and
364 years with a negative SCD anomaly include 1974, 1995, and 2008 (Table 3, Fig. 5c).
365 North Xinjiang is one of the regions prone to catastrophe, where frequent heavy
366 snowfall greatly affects the development of animal husbandry (Hao et al., 2002).

367 Years with a positive SCD anomaly in the Tibetan Plateau include 1983 and 1990,
368 whereas years with a negative SCD anomaly include 1965, 1969, and 2010 (Table 3).
369 The climate in the Tibetan Plateau is affected by the Indian monsoon from the south,
370 westerlies from the west, and the East Asian monsoon from the east (Yao et al., 2012).
371 Therefore, there is a regional difference in the SCD within the Tibetan Plateau, and

372 even a difference in the spatiotemporal distribution of snow disasters (Wang et al.,
373 2013). Our results differ from the conclusions drawn by Dong et al. (2001), as they
374 only used data from 26 stations, covering only a short period (1967–1996).

375 3.1.3 SCD trends

376 Changing trends of annual SCDs are examined, as shown in Figure 6a, and
377 summarized in Table 4. Among the ~~352-296~~ stations, there are ~~54-35~~ stations (~~1512~~%)
378 with a significant negative trend, and ~~35-37~~ stations (~~1013~~%) with a significant positive
379 trend (both at the 90% level), while 75% of stations show no significant trends. The
380 SCD exhibits a significant downward trend in ~~the Shandong Peninsula, and~~
381 ~~insignificant downward trends in the North China Plain, the Loess Plateau,~~ the
382 Xiaoxingganling, the Changbai Mountains, the Shandong Peninsula, the Qilian
383 Mountains, the North XinjiangTianshan Mountains, Northeast Qinghai, and the
384 peripheral zones in the south and eastern Tibetan Plateau (Fig. 6a). ~~Some station~~
385 ~~records indicate a decreasing rate of 1.3–7.2 days per decade.~~ For example, the SCD
386 decreased by ~~40-50~~ days from 1955 to 2010 at the Kuandian station in Northeast China,
387 ~~30-28~~ days from 1954 to 2010 at the Hongliuhe station in Xinjiang, and ~~15-10~~ days
388 from 1958 to 2010 at the Gangcha station on the Tibetan Plateau (Fig. 7a–c).

389 The SCDs in the Bayan Har Mountains, the Anemaqen Mountains, the Inner
390 Mongolia Plateau, and ~~Daxingganling~~the Northeast Plain, exhibit a significant upward
391 trend (Fig. 6a). For example, at the Shiqu station on the eastern border of the Tibetan
392 Plateau, the SCD increased 26 days from 1960 to 2010 (Fig. 7d). The coexistence of
393 negative and positive trends in the SCD change was also reported by Bulygina et al.

394 (2009) and Wang and Li (2012).

395 **3.2 Spatiotemporal variations of SCOD**

396 **3.2.1 SCOD variations**

397 The SCOD is closely related to both latitude and elevation (Fig. 8a). For example,
398 snowfall begins in September on the Tibetan Plateau, in early or middle October on the
399 Daxingganling, and in middle or late October on the Altai Mountains ~~of in~~ Xinjiang.
400 The SCOD also varies from one year to another (Table 2). Using the definition of a
401 year with a positive (negative) SCD anomaly, as introduced before (i.e. 70% stations
402 with positive (negative) SCOD anomaly and 30% stations with SCOD larger (smaller)
403 than the mean \pm 1SD), we consider a given year as a late (early) SCOD year. ~~Only~~
404 ~~Two~~ years, 1996 and 2006, can be considered as late SCOD years on a large scale
405 (Table 2), especially in 2006, in East China and the Tibetan Plateau (Fig. 6d5d), ~~while~~
406 ~~Only one year, (1982) not any single year,~~ can be considered as an early SCOD year.

407 **3.2.2 SCOD trends**

408 There are ~~136-196~~ stations (~~3966~~%) with a significant trend of late SCOD, and ~~23~~
409 ~~8~~ stations (~~73~~%) with a significant trend of early SCOD (both at the 90% level), while
410 ~~5431~~% of the stations show no significant trends (Table 4). The SCOD in ~~Northeast the~~
411 ~~major snow regions in~~ China, ~~the central and eastern Tibetan Plateau, the upper reach~~
412 ~~of the Yellow River, North Gansu, and North Xinjiang~~ exhibits a significant trend
413 towards late SCOD (Fig. 6b). These significantly late trends dominate the major snow
414 ~~areas regions of in~~ China. In particular, the late SCOD in Northeast China is consistent
415 with a previous study (Li et al., 2009). ~~Only~~ ~~the~~ SCOD in the ~~Pan-Bohai-East~~

416 ~~Liaoning Bay region and the Tianshan Mountains~~ exhibits a significant trend towards
417 early SCOD. ~~However, this trend is only significant in the Liaoxi corridor and the~~
418 ~~Tianshan Mountains.~~ For example, the SCOD at the Pingliang station in Gansu
419 Province shows a late rate of 5.2 days per decade from 1952 to 2010, but the SCOD at
420 the Weichang station in Hebei Province shows an early rate of 5.2 days per decade
421 from 1952 to 2010 (Fig. 7e–f).

422 **3.3 Spatiotemporal variations of SCED**

423 **3.3.1 SCED variations**

424 The pattern of SCED is similar to that of SCOD (Fig. 8b), i.e. places with early
425 snowfall normally show late snowmelt, while places with late snowfall normally show
426 early snowmelt. Like the SCOD, temporal variations of SCED are large (Table 2).
427 Using the same standard for defining the SCOD anomaly, we judge a given year as a
428 late (early) SCED year. Three years, 1957, 1976 and 1979, can be considered as late
429 SCED years on a large scale (Table 2). It is evident that 1957 was a typical year whose
430 SCED was late, which was also the reason for the great SCDs (Table 2, Fig. 5a and e).
431 The SCEDs in 1997 ~~and 2004 were very early. For example, in 1997, the SCED~~ was
432 early for almost all of China except for the Tibetan Plateau, western Tianshan
433 Mountains, and western Liaoning (Fig. 5f). ~~In general, the early SCED is dominant and~~
434 ~~more evident than the late SCED (Table 2).~~

435 **3.3.2 SCED trends**

436 For the SCED, there are 138-103 stations (3935%) with a significantly early trend
437 (at the 90% level), while 6064% of stations show no significant trends (Table 4). The

438 ~~Major-major~~ snow ~~areas-regions~~ in China all show early SCED, significant for
439 Northeast China, North Xinjiang and the Tibetan Plateau (Fig. 6c). The tendency of late
440 SCED is limited, with only ~~two-3~~ stations (1%) showing a significant trend. For
441 example, the SCED at the Jixi station in Northeast China shows an early rate of 4.43.5
442 days per decade from 1952 to 2010, while the SCED at the Maerkang station in
443 Sichuan Province shows a late rate of 4.2 days per decade from 1954 to 2010 (Fig.
444 7g-h).

445 **4 Discussion**

446 In the context of global warming, 136-196 stations (3966%) show significantly late
447 SCOD, and 138-103 stations (3935%) show significantly early SCED, all at the 90%
448 confidence level. It is not necessary for one station to show both significantly late
449 SCOD and early SCED. This explains why only 1512% of stations show a significantly
450 negative SCD trend, while 75% of stations show no significant change in the SCD
451 trends. The latter is inconsistent with the overall shortening of the snow period in the
452 Northern Hemisphere reported by Choi et al. (2010). One reason could be the different
453 time periods used in the two studies, 1972–2007 in Choi et al. (2010) as compared with
454 1952–2010 in this study. Below, we discuss the possible connections between the
455 spatiotemporal variations of snow cover and the warming climate and changing AO.

456 **4.1 Relationship with TBZD**

457 The number of days with temperature below 0°C (TBZD) plays an important role
458 in the SCD. There are 330-280 stations (9495% of all-296 stations) showing positive
459 correlations between TBZD and SCD, with 193-154 of them (5552%) having

460 significantly positive correlations (Table 4, Fig. 6d). For example, there is a
461 significantly positive correlation between SCD and TBZD at the Chengshantou station
462 (Fig. 9a). Therefore, generally speaking, the smaller the TBZD, the shorter the SCD.

463 For the SCOD, there are 287–245 stations with negative correlations with TBZD,
464 accounting for 8283% of 352–296 stations, whereas only 63–51 stations (1817%) show
465 positive correlations (Table 4). This means that for smaller TBZD, the SCOD is later.
466 For the SCED, there are 318–269 stations with positive correlations, accounting for
467 9091% of 352–296 stations, whereas only 34–27 stations (109%) have negative
468 correlations. This means that for smaller TBZD, the SCED is earlier.

469 Very similar results are found for the MAT (Table 4, Fig. 6e), and Fig. 9b shows
470 an example (the BaichengTieli station).

471 4.2 Relationship with AO

472 Although the AO index showed a strong positive trend in the past decades
473 (Thompson et al., 2000), its impact on the SCD in China is spatially distinctive.
474 Positive correlations (4746% of 352–296 stations) are found in ~~central China, i.e. the~~
475 ~~eastern Tibetan Plateau and the Loess Plateau, the upper reach of the Yangtze River,~~
476 ~~and the upper and middle reaches of the Yellow River~~ (Table 4, Fig. 6f), and Fig. 9c
477 shows an example (the Huajialing station). Negative correlations (5354% of 352–296
478 stations) exist in North Xinjiang, ~~the Changbaishan Mountain and the coasts of the~~
479 ~~Liaoning~~ Northeast China and the Shandong Peninsula, and Fig. 9d shows an example
480 (the Tonghua station).

481 5 Conclusion

482 This study examines the snow cover change based on 672 stations in 1952–2010 in
483 China. Specifically, the [352–296](#) stations with more than ten annual mean SCDs are
484 used to study the changing trends of SCD, SCOD, and SCED, and SCD relationships
485 with TBZD, MAT, and AO index during snow seasons. Some important results are
486 summarized below.

487 Northeast China, North Xinjiang, and the Tibetan Plateau are the three major snow
488 regions, with Northeast China being the largest. In North Xinjiang and in central and
489 north-eastern China, the SCDs are concentrated in the winter season. On the Tibetan
490 Plateau, however, snowfall is more frequent in the spring and fall. The overall
491 inter-annual variability of SCD is large in China. The years with a positive SCD
492 anomaly in China include 1955, 1957, 1964, and 2010, while the years with a negative
493 SCD anomaly are 1953, 1965, 1999, 2002, and 2009. Only [4512%](#) of stations show a
494 significantly negative SCD trend, while 75% of stations show no significant SCD
495 trends. This differs from the overall shortening of the snow period in the Northern
496 Hemisphere previously reported. One reason could be the different time periods used in
497 the two studies, 1972–2007 in the work of Choi et al. (2010) compared with 1952–2010
498 in this study. Our analyses indicate that the SCD distribution pattern and trends in
499 China are very complex and are not controlled by any single climate variable examined
500 (i.e. TBZD, MAT, or AO), but a combination of multiple variables. However, it seems
501 that the AO-[index](#) has the most impact on the SCD shortening trends in the Shandong
502 Peninsula, Changbai Mountains, [Xiaoxingganling](#), and North Xinjiang; the combination
503 of smaller TBZD and increasing MAT has the largest impact on the SCD shortening

504 trends on [the Tibetan Plateau](#), the Loess Plateau, [Xiaoxingganling](#), and the [Sanjiang](#)
505 [Northeast](#) Plain.

506 It is found that significantly late SCOD occurs in [nearly the entire](#)~~Northeast~~ China
507 [except for the east Liaoning Bay region, the central and eastern Tibetan Plateau, the](#)
508 [upper reach of the Yellow River, North Gansu, and North Xinjiang](#); significantly early
509 SCED occurs in [nearly all major snow arearegions in China](#)~~Northeast China and the~~
510 [Tibetan Plateau](#). Both the SCOD and SCED are closely related to the TBZD and MAT,
511 and are mostly controlled by local latitude and elevation. Owing to global warming
512 since 1950s, the reduced TBZD and increased MAT are the main reasons for overall
513 late SCOD and early SCED, although it is not necessary for one station to experience
514 both significantly late SCOD and early SCED. This explains why only [4512](#)% of
515 stations show significantly negative SCD trends, while 75% of stations show no
516 significant SCD trends.

517 Long-duration, consistent records of snow [cover and depth](#) are rare in China
518 because of many challenges associated with taking accurate and representative
519 measurements, especially in western China; ~~the~~ the station density and metric choice
520 also vary with time and locality. ~~T~~ therefore, more accurate and reliable observation
521 data are needed to further analyse the spatiotemporal distribution and features of snow
522 cover phenology. Atmospheric circulation causes variability in the snow cover
523 phenology, and [itsthis](#) effect ~~also~~ requires deeper investigations.

524

525

526 **Acknowledgments**

527 This work is financially supported by the Program for National Nature Science
528 Foundation of China (No. 41371391), and the Program for the Specialized Research
529 Fund for the Doctoral Program of Higher Education of China (No. 20120091110017).

530 This work is also partially supported by Collaborative Innovation Center of Novel
531 Software Technology and Industrialization. We would like to thank the National
532 Climate Center of China (NCC) in Beijing for providing valuable climate datasets. We
533 thank the three anonymous reviewers and the editor for valuable comments and
534 suggestions that greatly improved the quality of this paper.

535

536 **References**

537 An, D., Li, D., Yuan, Y. and Hui, Y.: Contrast between snow cover data of different
538 definitions, *J. Glaciol. Geocrol.*, 31(6), 1019-1027, 2009.

539 Alexandersson, H. and Moberg, A.: Homogenization of Swedish temperature data Part
540 1: homogeneity test for linear trends, *Int. J. Climatol.*, 17, 25-34, 1997.

541 Barnett, T. P., Dumenil, L. and Latif, M.: The effect of Eurasian snow cover on regional
542 and global climate variations, *J. Atmos. Sci.*, 46, 661-685, 1989.

543 Beniston, M: Variations of snow depth and duration in the Swiss Alps over the last 50
544 years: Links to changes in large-scale climatic forcings, *Clim. Change*, 36, 281-300,
545 1997.

546 Birsan, M. V. and Dumitrescu, A.: Snow variability in Romania in connection to
547 large-scale atmospheric circulation, *Int. J. Climatol.*, 34, 134-144, 2014.

548 Bolsenga, S. J., and Norton, D. C.: Maximum snowfall at long-term stations in the
549 U.S./Canadian Great Lakes, *Nat. Hazards*, 5, 221-232, 1992.

550 Brown, R. D. and Robinson, D. A.: Northern Hemisphere spring snow cover variability
551 and change over 1922-2010 including an assessment of uncertainty, *The Cryosphere*,
552 5, 219-229, 2011.

553 Bulygina, O. N., Razuvaev, V. N. and Korshunova, N. N.: Changes in snow cover over
554 Northern Eurasia in the last few decades, *Environ. Res. Lett.*, 4, 045026, 2009.

555 Changnon, S. A. and Changnon, D.: A spatial and temporal analysis of damaging
556 snowstorms in the United States, *Nat. Hazards*, 37, 373-389, 2006.

557 Chen, S., Chen, W. and Wei, K.: Recent trends in winter temperature extremes in
558 eastern China and their relationship with the Arctic Oscillation and ENSO, *Adv.*
559 *Atmos. Sci.*, 30, 1712-1724, 2013.

560 China Meteorological Administration: Specifications for Surface Meteorological
561 Observations, Beijing, China Meteorological Press, 1-62, 2003.

562 Choi, G., Robinson, D. A. and Kang, S.: Changing Northern Hemisphere snow seasons,
563 *J. Climate*, 23, 5305-5310, 2010.

564 Ciach, G. J. and Krajewski, W. F.: Analysis and modeling of spatial correlation
565 structure in small-scale rainfall in Central Oklahoma, *Adv. Water Resour.*, 29(10),
566 1450–1463, 2006.

567 Déry, S. J. and Brown, R. D.: Recent Northern Hemisphere snow cover extent trends
568 and implications for the snow-albedo feedback, *Geophys. Res. Lett.*, 34, L22504,
569 2007.

570 Dong, A., Guo, H., Wang, L. and Liang, T.: A CEOF analysis on variation about yearly
571 snow days in Northern Xinjiang in recent 40 years, *Plateau Meteorol.*, 23, 936-940,
572 2004.

573 Dong, W., Wei, Z. and Fan, J.: Climatic character analysis of snow disasters in east
574 Qinghai-Xizang Plateau livestock farm, *Plateau Meteorol.*, 20, 402-406, 2001.

575 Dyer, J. L. and Mote, T. L.: Spatial variability and trends in observed snow depth over
576 North America, *Geophys. Res. Lett.*, 33, L16503, 2006

577 Fang, S., Qi, Y., Han, G., Zhou, G. and Cammarano, D.: Meteorological drought trend
578 in winter and spring from 1961 to 2010 and its possible impacts on wheat in wheat
579 planting area of China, *Sci. Agricul. Sin.*, 47, 1754-1763, 2014

580 Gao, H.: China's snow disaster in 2008, who is the principal player? *Int. J. Climatol.*, 29,
581 2191-2196, 2009.

582 Gong, D. Y., Wang, S. W. and Zhu, J. H.: East Asian winter monsoon and Arctic
583 oscillation, *Geophys. Res. Lett.*, 28, 2073-2076, 2001.

584 Groisman, P. Y., Karl, T. R. and Knight, R. W.: Observed impact of snow cover on the
585 heat-balance and the rise of continental spring temperatures, *Science*, 263, 198-200,
586 1994.

587 Habib, E., Krajewski, W. F. and Ciach, G. J.: Estimation of rainfall interstation
588 correlation, *J. Hydrometeorol.*, 2(6), 621–629, 2001.

589 Hantel, M., Ehrendorfer, M. and Haslinger, A.: Climate sensitivity of snow cover
590 duration in Austria, *Int. J. Climatol.*, 20, 615-640, 2000.

591 Hao, L., Wang, J., Man, S. and Yang, C.: Spatio-temporal change of snow disaster and
592 analysis of vulnerability of animal husbandry in China, *J. Nat. Disaster*, 11, 42-48,
593 2002.

594 He, L. and Li, D.: Classification of snow cover days and comparing with satellite
595 remote sensing data in west China, *J. Glaciol. Geocrol.*, 33(2), 237-245, 2011.

596 Hu, H. and Liang, L.: Temporal and spatial variations of snowfall in the east of
597 Qinghai-Tibet Plateau in the last 50 years, *Acta Geogr. Sin.*, 69, 1002-1012, 2014.

598 Jeong, J. H. and Ho, C. H.: Changes in occurrence of cold surges over East Asia in
599 association with Arctic oscillation, *Geophys. Res. Lett.*, 32, L14704, 2005.

600 Ji, Z. and Kang, S.: Projection of snow cover changes over China under RCP scenarios
601 *Clim. Dyn.*, 41, 589-600, 2013.

602 Ke, C. Q. and Li, P. J.: Spatial and temporal characteristics of snow cover over the
603 Tibetan plateau, *Acta Geogr. Sin.*, 53, 209-215, 1998.

604 Ke, C. Q. and Liu, X.: MODIS-observed spatial and temporal variation in snow cover in
605 Xinjiang, China, *Clim. Res.*, 59, 15-26, 2014.

606 Ke, C. Q., Yu, T., Yu, K., Tang, G. D. and King, L.: Snowfall trends and variability in
607 Qinghai, China, *Theor. Appl. Climatol.*, 98, 251-258, 2009.

608 Kyriakidis, P. C. and Goodchild, M. F.: On the prediction error variance of three
609 common spatial interpolation schemes, *Int. J. Geogr. Info. Science*, 20(8), 823-855,
610 2006.

611 Lehning, M., Grünewald, T. and Schirmer, M.: Mountain snow distribution governed by
612 an altitudinal gradient and terrain roughness, *Geophys. Res. Lett.*, 38, L19504, 2011.

613 Li, D., Liu, Y., Yu, H. and Li, Y.: Spatial-temporal variation of the snow cover in
614 Heilongjiang Province in 1951-2006, *J. Glaciol. Geocrol.*, 31, 1011-1018, 2009.

615 Li, J. and Wang, J.: A modified zonal index and its physical sense, *Geophys. Res. Lett.*,
616 30, 1632, 2003.

617 Li, L. Y. and Ke, C. Q.: Analysis of spatiotemporal snow cover variations in Northeast
618 China based on moderate-resolution-imaging spectroradiometer data, *J. Appl.*
619 *Remote Sens.*, 8, 084695, doi: 10.1117/1.JRS.8.084695. 2014.

620 Li, P. J.: Dynamic characteristic of snow cover in western China, *Acta Meteorol. Sin.*,
621 48, 505-515, 1993.

622 Li, P. J.: A preliminary study of snow mass variations over past 30 years in China, *Acta*
623 *Geogr. Sin.*, 48, 433-437, 1990.

624 Li, P. J. and Mi, D.: Distribution of snow cover in China, *J. Glaciol. Geocrol.*, 5, 9-18,
625 1983.

626 Liang, T. G., Huang, X. D., Wu, C. X., Liu, X. Y., Li, W. L., Guo, Z. G. and Ren, J. Z.:
627 An application of MODIS data to snow cover monitoring in a pastoral area: A case
628 study in Northern Xinjiang, China, *Remote Sens. Environ.*, 112, 1514-1526, 2008.

629 Liu, Y., Ren, G. and Yu, H.: Climatology of Snow in China, *Sci. Geogr. Sin.*, 32,
630 1176-1185, 2012.

631 Llasat, M. C., Turco, M., Quintana-Seguí, P. and Llasat-Botija, M.: The snow storm of
632 8 March 2010 in Catalonia (Spain): a paradigmatic wet-snow event with a high
633 societal impact, *Nat. Hazards Earth Syst. Sci.*, 14, 427-441, 2014.

634 Lü, J. M., Ju, J. H., Kim, S. J., Ren, J. Z. and Zhu, Y. X.: Arctic Oscillation and the
635 autumn/winter snow depth over the Tibetan Plateau, *J. Geophys. Res.*, 113, D14117,
636 2008.

637 Ma, L. and Qin, D.: Temporal-spatial characteristics of observed key parameters of
638 snow cover in China during 1957-2009, *Sci. Cold Arid Reg.*, 4, 384-393, 2012.

639 Marty, C.: Regime shift of snow days in Switzerland, *Geophys. Res. Lett.*, 35, L12501,
640 2008.

641 Morán-Tejeda, E., López-Moreno, J. I. and Beniston, M.: The changing roles of
642 temperature and precipitation on snowpack variability in Switzerland as a function of
643 altitude, *Geophys. Res. Lett.*, 40, 2131-2136, 2013.

644 Pederson, G. T., Betancourt, J. L. and Gregory, J. M.: Regional patterns and proximal
645 causes of the recent snowpack decline in the Rocky Mountains, U.S., *Geophys. Res.*
646 *Lett.*, 40, 1811-1816, 2013.

647 Peings, Y., Brun, B., Mauvais, V. and Douville, H.: How stationary is the relationship
648 between Siberian snow and Arctic Oscillation over the 20th century, *Geophys. Res.*
649 *Lett.*, 40, 183-188, 2013.

650 Peng, S., Piao, S., Ciais, P., Fang, J. and Wang, X.: Change in winter snow depth and its
651 impacts on vegetation in China, *Glob. Change Biol.*, 16, 3004-3013, 2010.

652 Peng, S., Piao, S., Ciais, P., Friedlingstein, P., Zhou, L. and Wang, T.: Change in snow
653 phenology and its potential feedback to temperature in the Northern Hemisphere over
654 the last three decades, *Environ. Res. Lett.*, 8, 014008, 2013.

655 Qin, D., Liu, S. and Li, P.: Snow cover distribution, variability, and response to climate
656 change in western China, *J. Climate*, 19, 1820-1833, 2006.

657 Ren, G. Y., Guo, J., Xu, M. Z., Chu, Z. Y., Zhang, L., Zou, X. K., Li, Q. X. and Liu, X.
658 N.: Climate changes of China's mainland over the past half century, *Acta. Meteorol.*
659 *Sin.*, 63, 942-956, 2005.

660 Robinson, D. A. and Dewey, K. F.: Recent secular variations in the extent of northern
661 hemisphere snow cover, *Geophys. Res. Lett.*, 17, 1557-1560, 1990.

662 Scherrer, S. C., Appenzeller, C. and Laternser, M.: Trends in Swiss Alpine snow days:
663 The role of local- and large-scale climate variability, *Geophys. Res. Lett.*, 31, L13215,
664 2004.

665 Scherrer, S. C. and Appenzeller, C.: Swiss Alpine snow pack variability: major patterns
666 and links to local climate and large-scale flow, *Clim. Res.*, 32(3), 187-199, 2006.

667 Scherrer, S. C., Wüthrich, C., Croci-Maspoli, M., Weingartner, R. and Appenzeller, C.:
668 Snow variability in the Swiss Alps 1864-2009, *Int. J. Clim.*, 33(15), 3162 – 3173,
669 2013, doi: 10.1002/joc.3653.

670 Serquet, G., Marty, C., Dulex, J-P. and Rebetez, M.: Seasonal trends and temperature
671 dependence of the snowfall/precipitation-day ratio in Switzerland, *Geophys. Res.*
672 *Lett.*, 38, L07703, 2011.

673 Shi, Y., Gao, X., Wu, J. and Giorgi, F.: Changes in snow cover over China in the 21st
674 century as simulated by a high resolution regional climate model, *Environ. Res. Lett.*,
675 6, 045401, 2011.

676 Tang, X., Yan, X., Ni, M. and Lu, Y.: Changes of the snow cover days on Tibet Plateau
677 in last 40 years, *Acta. Geogr. Sin.*, 67, 951-959, 2012.

678 Thompson, D. W. J. and Wallace, J. M.: The Arctic oscillation signature in the
679 wintertime geopotential height and temperature fields, *Geophys. Res. Lett.*, 25,
680 1297-1300, 1998.

681 Thompson, D. W. J., Wallace, J. M. and Hegerl, G. C.: Annular modes in the
682 extratropical circulation, part II: Trends, *J. Climate*, 13, 1018-1036, 2000.

683 Wang, C. and Li, D.: Spatial-temporal variations of the snow cover days and the
684 maximum depth of snow cover in China during recent 50 years, *J. Glaciol. Geocrol.*,
685 34, 247-256, 2012.

686 Wang, C., Wang, Z. and Cui, Y.: Snow cover of China during the last 40 years: Spatial
687 distribution and interannual variation, *J. Glaciol. Geocrol.*, 31, 301-310, 2009a.

688 Wang, J. and Hao, X.: Responses of snowmelt runoff to climatic change in an inland
689 river basin, Northwestern China, over the past 50 years, *Hydrol. Earth Syst. Sci.*, 14,
690 1979-1987, 2010.

691 Wang, L. et al.: Characteristics of the extreme low-temperature, heavy snowstorm and
692 freezing disasters in January 2008 in China, *Meteorol. Mon.*, 34, 95-100, 2008.

693 Wang, Q., Zhang, C., Liu, J. and Liu, W.: The changing tendency on the depth and days
694 of snow cover in Northern Xinjiang, *Adv. Clim. Change Res.*, 5, 39-43, 2009b.

695 Wang, W., Liang, T., Huang, X., Feng, Q., Xie, H., Liu, X., Chen, M. and Wang, X.:
696 Early warning of snow-caused disasters in pastoral areas on the Tibetan Plateau, *Nat.*
697 *Hazards Earth Syst. Sci.*, 13, 1411-1425, 2013.

698 Wu, B. Y. and Wang, J.: Winter Arctic oscillation, Siberian high and East Asian winter
699 monsoon, *Geophys. Res. Lett.*, 29, 1897, 2002.

700 Xi, Y., Li, D. and Wang, W.: Study of the temporal-spatial characteristics of snow
701 covers days in Hetao and its vicinity, *J. Glaciol. Geocrol.*, 31, 446-456, 2009.

702 Xu, L., Li, D. and Hu, Z.: Relationship between the snow cover day and monsoon index
703 in Tibetan Plateau, *Plateau Meteorol.*, 29, 1093-1101, 2010.

704 Yang, H., Yang, D., Hu, Q. and Lv, H.: Spatial variability of the trends in climatic
705 variables across China during 1961-2010, *Theor. Appl. Climatol.*, 2015 (in press).

706 Yao, T. et al.: Different glacier status with atmospheric circulations in Tibetan Plateau
707 and surroundings, *Nature Clim. Change*, 2, 663-667, 2012.

708 Ye, H. and Ellison, M.: Changes in transitional snowfall season length in northern
709 Eurasia, *Geophys. Res. Lett.*, 30, 1252, 2003.

710 You, Q., Kang, S., Ren, G., Fraedrich, K., Pepin, N., Yan, Y. and Ma, L.: Observed
711 changes in snow depth and number of snow days in the eastern and central Tibetan
712 Plateau, *Clim. Res.*, 46, 171-183, 2011.

713 Zhang, T.: Influence of the seasonal snow cover on the ground thermal regime: An
714 overview, *Rev. Geophys.*, 43, 1-23, 2005.

715

716 **Table Captions**

717 **Table 1.** Prediction errors of cross validation for the spatial interpolation with the
 718 universal kriging method.

<u>Item (Figures)</u>	<u>Mean error</u>	<u>Average standard error</u>	<u>Root mean squared error</u>	<u>Root mean squared standardized error</u>
<u>SCD (Fig.3a)</u>	<u>-0.0230</u>	<u>11.0558</u>	<u>13.7311</u>	<u>1.1097</u>
<u>CV (Fig.3b)</u>	<u>0.0017</u>	<u>0.7364</u>	<u>0.5510</u>	<u>0.7579</u>
<u>SCD in 1957 (Fig.5a)</u>	<u>-0.0015</u>	<u>11.1561</u>	<u>13.4662</u>	<u>1.1898</u>
<u>SCD in 2002 (Fig.5b)</u>	<u>0.0306</u>	<u>6.6185</u>	<u>8.5887</u>	<u>1.2522</u>
<u>SCD in 2008 (Fig.5c)</u>	<u>0.0477</u>	<u>7.3167</u>	<u>8.1968</u>	<u>1.0969</u>
<u>SCED in 1957 (Fig.5d)</u>	<u>-0.0449</u>	<u>15.0528</u>	<u>18.9860</u>	<u>1.1921</u>
<u>SCED in 1997 (Fig.5e)</u>	<u>0.0696</u>	<u>15.5722</u>	<u>17.7793</u>	<u>1.1040</u>
<u>SCOD in 2006 (Fig.5f)</u>	<u>0.0482</u>	<u>15.4503</u>	<u>16.1757</u>	<u>1.0449</u>
<u>SCOD (Fig.8a)</u>	<u>0.0293</u>	<u>11.2458</u>	<u>13.9078</u>	<u>1.1712</u>
<u>SCED (Fig.8b)</u>	<u>-0.0222</u>	<u>15.2265</u>	<u>18.3095</u>	<u>1.1308</u>

719

720

721

722

723

724

725

726

727 **Table 2.** Percentage (%) of stations with anomalies (P for positive and N for negative)
728 of snow cover day (SCD), snow cover onset date (SCOD), and snow cover end date
729 (SCED). Percentage (%) of stations with anomalies of SCD, SCOD, and SCED larger
730 (smaller) than the mean +/- one or two standard deviations (1SD or 2SD), with the bold
731 number denoting years with a positive (negative) SCD anomaly, and late (early) years
732 for SCOD or SCED in China. All the percentages are calculated based on 672 stations.

733

Year	P	SCD					SCOD					SCED						
		1SD	2SD	-2SD	-1SD	N	P	1SD	2SD	-2SD	-1SD	N	P	1SD	2SD	-2SD	-1SD	N
1952	31	<u>2</u>	<u>0</u>	<u>13</u>	<u>33</u>	<u>69</u>	<u>69</u>	<u>40</u>	<u>21</u>	<u>2</u>	<u>9</u>	<u>31</u>	<u>55</u>	<u>17</u>	<u>2</u>	<u>12</u>	<u>17</u>	<u>45</u>
1953	28	<u>7</u>	<u>0</u>	<u>3</u>	<u>36</u>	<u>72</u>	<u>40</u>	<u>8</u>	<u>2</u>	<u>2</u>	<u>18</u>	<u>60</u>	<u>37</u>	<u>8</u>	<u>1</u>	<u>10</u>	<u>18</u>	<u>63</u>
1954	57	<u>31</u>	<u>12</u>	<u>0</u>	<u>8</u>	<u>43</u>	<u>35</u>	<u>8</u>	<u>4</u>	<u>1</u>	<u>18</u>	<u>65</u>	<u>56</u>	<u>11</u>	<u>0</u>	<u>0</u>	<u>10</u>	<u>44</u>
1955	79	<u>45</u>	<u>25</u>	<u>1</u>	<u>5</u>	<u>21</u>	<u>37</u>	<u>9</u>	<u>4</u>	<u>1</u>	<u>22</u>	<u>63</u>	<u>77</u>	<u>21</u>	<u>2</u>	<u>1</u>	<u>6</u>	<u>23</u>
1956	46	<u>10</u>	<u>0</u>	<u>0</u>	<u>4</u>	<u>54</u>	<u>69</u>	<u>20</u>	<u>2</u>	<u>0</u>	<u>9</u>	<u>31</u>	<u>61</u>	<u>24</u>	<u>1</u>	<u>2</u>	<u>12</u>	<u>39</u>
1957	85	<u>62</u>	<u>26</u>	<u>0</u>	<u>3</u>	<u>15</u>	<u>26</u>	<u>6</u>	<u>1</u>	<u>0</u>	<u>15</u>	<u>74</u>	<u>84</u>	<u>35</u>	<u>5</u>	<u>1</u>	<u>4</u>	<u>16</u>
1958	48	<u>15</u>	<u>4</u>	<u>0</u>	<u>14</u>	<u>52</u>	<u>46</u>	<u>17</u>	<u>0</u>	<u>0</u>	<u>18</u>	<u>54</u>	<u>52</u>	<u>17</u>	<u>3</u>	<u>4</u>	<u>18</u>	<u>48</u>
1959	28	<u>7</u>	<u>1</u>	<u>4</u>	<u>23</u>	<u>72</u>	<u>53</u>	<u>26</u>	<u>8</u>	<u>1</u>	<u>18</u>	<u>47</u>	<u>59</u>	<u>23</u>	<u>3</u>	<u>1</u>	<u>5</u>	<u>41</u>
1960	37	<u>13</u>	<u>3</u>	<u>0</u>	<u>16</u>	<u>63</u>	<u>49</u>	<u>11</u>	<u>2</u>	<u>0</u>	<u>10</u>	<u>51</u>	<u>59</u>	<u>24</u>	<u>6</u>	<u>4</u>	<u>18</u>	<u>41</u>
1961	36	<u>7</u>	<u>1</u>	<u>1</u>	<u>18</u>	<u>64</u>	<u>25</u>	<u>9</u>	<u>2</u>	<u>1</u>	<u>27</u>	<u>75</u>	<u>30</u>	<u>6</u>	<u>1</u>	<u>7</u>	<u>26</u>	<u>70</u>
1962	41	<u>11</u>	<u>3</u>	<u>0</u>	<u>10</u>	<u>59</u>	<u>44</u>	<u>13</u>	<u>4</u>	<u>2</u>	<u>10</u>	<u>56</u>	<u>58</u>	<u>18</u>	<u>3</u>	<u>0</u>	<u>11</u>	<u>42</u>
1963	25	<u>5</u>	<u>2</u>	<u>2</u>	<u>27</u>	<u>75</u>	<u>34</u>	<u>14</u>	<u>5</u>	<u>1</u>	<u>23</u>	<u>66</u>	<u>51</u>	<u>14</u>	<u>0</u>	<u>8</u>	<u>17</u>	<u>49</u>
1964	76	<u>36</u>	<u>11</u>	<u>0</u>	<u>1</u>	<u>24</u>	<u>31</u>	<u>3</u>	<u>1</u>	<u>4</u>	<u>24</u>	<u>69</u>	<u>64</u>	<u>18</u>	<u>1</u>	<u>0</u>	<u>5</u>	<u>36</u>
1965	26	<u>8</u>	<u>0</u>	<u>1</u>	<u>32</u>	<u>74</u>	<u>59</u>	<u>18</u>	<u>5</u>	<u>1</u>	<u>8</u>	<u>41</u>	<u>55</u>	<u>14</u>	<u>2</u>	<u>3</u>	<u>17</u>	<u>45</u>
1966	28	<u>6</u>	<u>1</u>	<u>0</u>	<u>13</u>	<u>72</u>	<u>46</u>	<u>21</u>	<u>6</u>	<u>0</u>	<u>13</u>	<u>54</u>	<u>67</u>	<u>12</u>	<u>1</u>	<u>2</u>	<u>5</u>	<u>33</u>
1967	31	<u>5</u>	<u>0</u>	<u>3</u>	<u>23</u>	<u>69</u>	<u>40</u>	<u>11</u>	<u>3</u>	<u>2</u>	<u>15</u>	<u>60</u>	<u>43</u>	<u>5</u>	<u>0</u>	<u>3</u>	<u>12</u>	<u>57</u>
1968	61	<u>29</u>	<u>12</u>	<u>3</u>	<u>8</u>	<u>39</u>	<u>35</u>	<u>8</u>	<u>1</u>	<u>0</u>	<u>13</u>	<u>65</u>	<u>34</u>	<u>13</u>	<u>0</u>	<u>4</u>	<u>26</u>	<u>66</u>
1969	42	<u>18</u>	<u>5</u>	<u>4</u>	<u>21</u>	<u>58</u>	<u>45</u>	<u>13</u>	<u>1</u>	<u>3</u>	<u>20</u>	<u>55</u>	<u>67</u>	<u>20</u>	<u>1</u>	<u>1</u>	<u>7</u>	<u>33</u>
1970	46	<u>15</u>	<u>1</u>	<u>2</u>	<u>11</u>	<u>54</u>	<u>38</u>	<u>10</u>	<u>3</u>	<u>2</u>	<u>24</u>	<u>62</u>	<u>62</u>	<u>19</u>	<u>3</u>	<u>0</u>	<u>7</u>	<u>38</u>
1971	53	<u>12</u>	<u>1</u>	<u>1</u>	<u>9</u>	<u>47</u>	<u>38</u>	<u>15</u>	<u>4</u>	<u>1</u>	<u>17</u>	<u>62</u>	<u>53</u>	<u>9</u>	<u>1</u>	<u>1</u>	<u>8</u>	<u>47</u>
1972	55	<u>23</u>	<u>11</u>	<u>0</u>	<u>8</u>	<u>45</u>	<u>37</u>	<u>9</u>	<u>2</u>	<u>1</u>	<u>21</u>	<u>63</u>	<u>46</u>	<u>16</u>	<u>4</u>	<u>1</u>	<u>9</u>	<u>54</u>
1973	50	<u>19</u>	<u>2</u>	<u>1</u>	<u>7</u>	<u>50</u>	<u>35</u>	<u>10</u>	<u>1</u>	<u>2</u>	<u>23</u>	<u>65</u>	<u>43</u>	<u>9</u>	<u>1</u>	<u>1</u>	<u>8</u>	<u>57</u>
1974	33	<u>8</u>	<u>0</u>	<u>3</u>	<u>23</u>	<u>67</u>	<u>53</u>	<u>29</u>	<u>6</u>	<u>1</u>	<u>11</u>	<u>47</u>	<u>52</u>	<u>12</u>	<u>1</u>	<u>1</u>	<u>10</u>	<u>48</u>
1975	41	<u>10</u>	<u>4</u>	<u>1</u>	<u>15</u>	<u>59</u>	<u>26</u>	<u>7</u>	<u>2</u>	<u>1</u>	<u>21</u>	<u>74</u>	<u>43</u>	<u>15</u>	<u>3</u>	<u>2</u>	<u>16</u>	<u>57</u>
1976	35	<u>11</u>	<u>3</u>	<u>1</u>	<u>23</u>	<u>65</u>	<u>60</u>	<u>25</u>	<u>12</u>	<u>0</u>	<u>5</u>	<u>40</u>	<u>77</u>	<u>31</u>	<u>5</u>	<u>1</u>	<u>3</u>	<u>23</u>
1977	45	<u>20</u>	<u>3</u>	<u>0</u>	<u>9</u>	<u>55</u>	<u>28</u>	<u>5</u>	<u>1</u>	<u>0</u>	<u>25</u>	<u>72</u>	<u>57</u>	<u>14</u>	<u>3</u>	<u>2</u>	<u>12</u>	<u>43</u>
1978	60	<u>22</u>	<u>8</u>	<u>0</u>	<u>2</u>	<u>40</u>	<u>43</u>	<u>13</u>	<u>2</u>	<u>2</u>	<u>13</u>	<u>57</u>	<u>55</u>	<u>10</u>	<u>1</u>	<u>0</u>	<u>8</u>	<u>45</u>
1979	41	<u>8</u>	<u>1</u>	<u>0</u>	<u>7</u>	<u>59</u>	<u>43</u>	<u>11</u>	<u>1</u>	<u>0</u>	<u>20</u>	<u>57</u>	<u>79</u>	<u>32</u>	<u>2</u>	<u>0</u>	<u>4</u>	<u>21</u>
1980	39	<u>12</u>	<u>1</u>	<u>0</u>	<u>5</u>	<u>61</u>	<u>41</u>	<u>9</u>	<u>1</u>	<u>1</u>	<u>16</u>	<u>59</u>	<u>82</u>	<u>27</u>	<u>2</u>	<u>0</u>	<u>4</u>	<u>18</u>

<u>1981</u>	<u>42</u>	<u>13</u>	<u>2</u>	<u>0</u>	<u>13</u>	<u>58</u>	<u>45</u>	<u>20</u>	<u>4</u>	<u>2</u>	<u>18</u>	<u>55</u>	<u>44</u>	<u>13</u>	<u>1</u>	<u>2</u>	<u>15</u>	<u>56</u>
<u>1982</u>	<u>40</u>	<u>12</u>	<u>1</u>	<u>1</u>	<u>15</u>	<u>60</u>	<u>23</u>	<u>9</u>	<u>2</u>	<u>0</u>	<u>30</u>	<u>77</u>	<u>58</u>	<u>23</u>	<u>6</u>	<u>6</u>	<u>16</u>	<u>42</u>
<u>1983</u>	<u>50</u>	<u>19</u>	<u>6</u>	<u>0</u>	<u>12</u>	<u>50</u>	<u>44</u>	<u>14</u>	<u>1</u>	<u>1</u>	<u>11</u>	<u>56</u>	<u>67</u>	<u>26</u>	<u>2</u>	<u>1</u>	<u>9</u>	<u>33</u>
<u>1984</u>	<u>26</u>	<u>9</u>	<u>1</u>	<u>1</u>	<u>28</u>	<u>74</u>	<u>68</u>	<u>32</u>	<u>16</u>	<u>0</u>	<u>5</u>	<u>32</u>	<u>48</u>	<u>8</u>	<u>1</u>	<u>2</u>	<u>13</u>	<u>52</u>
<u>1985</u>	<u>66</u>	<u>24</u>	<u>3</u>	<u>0</u>	<u>3</u>	<u>34</u>	<u>32</u>	<u>8</u>	<u>1</u>	<u>1</u>	<u>24</u>	<u>68</u>	<u>46</u>	<u>8</u>	<u>2</u>	<u>1</u>	<u>8</u>	<u>54</u>
<u>1986</u>	<u>50</u>	<u>14</u>	<u>2</u>	<u>0</u>	<u>12</u>	<u>50</u>	<u>32</u>	<u>5</u>	<u>1</u>	<u>1</u>	<u>19</u>	<u>68</u>	<u>63</u>	<u>18</u>	<u>4</u>	<u>3</u>	<u>10</u>	<u>38</u>
<u>1987</u>	<u>67</u>	<u>23</u>	<u>4</u>	<u>0</u>	<u>4</u>	<u>33</u>	<u>40</u>	<u>7</u>	<u>1</u>	<u>2</u>	<u>15</u>	<u>60</u>	<u>60</u>	<u>23</u>	<u>3</u>	<u>1</u>	<u>8</u>	<u>40</u>
<u>1988</u>	<u>56</u>	<u>17</u>	<u>1</u>	<u>0</u>	<u>2</u>	<u>44</u>	<u>24</u>	<u>6</u>	<u>1</u>	<u>3</u>	<u>26</u>	<u>76</u>	<u>69</u>	<u>23</u>	<u>0</u>	<u>1</u>	<u>7</u>	<u>31</u>
<u>1989</u>	<u>47</u>	<u>18</u>	<u>4</u>	<u>0</u>	<u>11</u>	<u>53</u>	<u>71</u>	<u>29</u>	<u>7</u>	<u>1</u>	<u>6</u>	<u>29</u>	<u>41</u>	<u>6</u>	<u>1</u>	<u>3</u>	<u>18</u>	<u>59</u>
<u>1990</u>	<u>56</u>	<u>19</u>	<u>2</u>	<u>0</u>	<u>7</u>	<u>44</u>	<u>52</u>	<u>9</u>	<u>1</u>	<u>0</u>	<u>9</u>	<u>48</u>	<u>49</u>	<u>12</u>	<u>1</u>	<u>2</u>	<u>10</u>	<u>51</u>
<u>1991</u>	<u>34</u>	<u>4</u>	<u>0</u>	<u>2</u>	<u>9</u>	<u>66</u>	<u>60</u>	<u>21</u>	<u>3</u>	<u>0</u>	<u>4</u>	<u>40</u>	<u>72</u>	<u>26</u>	<u>3</u>	<u>1</u>	<u>4</u>	<u>28</u>
<u>1992</u>	<u>50</u>	<u>13</u>	<u>4</u>	<u>1</u>	<u>7</u>	<u>50</u>	<u>54</u>	<u>18</u>	<u>5</u>	<u>0</u>	<u>4</u>	<u>46</u>	<u>50</u>	<u>13</u>	<u>1</u>	<u>5</u>	<u>19</u>	<u>50</u>
<u>1993</u>	<u>58</u>	<u>19</u>	<u>2</u>	<u>1</u>	<u>4</u>	<u>42</u>	<u>43</u>	<u>9</u>	<u>1</u>	<u>0</u>	<u>17</u>	<u>57</u>	<u>49</u>	<u>18</u>	<u>2</u>	<u>2</u>	<u>21</u>	<u>51</u>
<u>1994</u>	<u>58</u>	<u>19</u>	<u>2</u>	<u>0</u>	<u>4</u>	<u>42</u>	<u>28</u>	<u>6</u>	<u>2</u>	<u>1</u>	<u>22</u>	<u>72</u>	<u>39</u>	<u>11</u>	<u>0</u>	<u>3</u>	<u>18</u>	<u>61</u>
<u>1995</u>	<u>36</u>	<u>10</u>	<u>3</u>	<u>3</u>	<u>15</u>	<u>64</u>	<u>57</u>	<u>24</u>	<u>3</u>	<u>1</u>	<u>15</u>	<u>43</u>	<u>49</u>	<u>8</u>	<u>1</u>	<u>7</u>	<u>18</u>	<u>51</u>
<u>1996</u>	<u>26</u>	<u>8</u>	<u>2</u>	<u>2</u>	<u>22</u>	<u>74</u>	<u>71</u>	<u>30</u>	<u>4</u>	<u>0</u>	<u>5</u>	<u>29</u>	<u>55</u>	<u>11</u>	<u>1</u>	<u>2</u>	<u>15</u>	<u>45</u>
<u>1997</u>	<u>37</u>	<u>3</u>	<u>0</u>	<u>1</u>	<u>18</u>	<u>63</u>	<u>44</u>	<u>13</u>	<u>3</u>	<u>2</u>	<u>12</u>	<u>56</u>	<u>18</u>	<u>4</u>	<u>2</u>	<u>9</u>	<u>49</u>	<u>82</u>
<u>1998</u>	<u>34</u>	<u>8</u>	<u>2</u>	<u>4</u>	<u>18</u>	<u>66</u>	<u>37</u>	<u>11</u>	<u>3</u>	<u>1</u>	<u>20</u>	<u>63</u>	<u>30</u>	<u>9</u>	<u>1</u>	<u>7</u>	<u>25</u>	<u>70</u>
<u>1999</u>	<u>25</u>	<u>4</u>	<u>1</u>	<u>1</u>	<u>35</u>	<u>75</u>	<u>61</u>	<u>23</u>	<u>12</u>	<u>1</u>	<u>7</u>	<u>39</u>	<u>51</u>	<u>11</u>	<u>2</u>	<u>5</u>	<u>15</u>	<u>49</u>
<u>2000</u>	<u>64</u>	<u>17</u>	<u>4</u>	<u>0</u>	<u>5</u>	<u>36</u>	<u>59</u>	<u>18</u>	<u>2</u>	<u>0</u>	<u>9</u>	<u>41</u>	<u>39</u>	<u>7</u>	<u>0</u>	<u>5</u>	<u>22</u>	<u>61</u>
<u>2001</u>	<u>67</u>	<u>29</u>	<u>8</u>	<u>0</u>	<u>5</u>	<u>33</u>	<u>39</u>	<u>16</u>	<u>2</u>	<u>1</u>	<u>22</u>	<u>61</u>	<u>42</u>	<u>17</u>	<u>1</u>	<u>3</u>	<u>15</u>	<u>58</u>
<u>2002</u>	<u>17</u>	<u>2</u>	<u>0</u>	<u>5</u>	<u>32</u>	<u>83</u>	<u>59</u>	<u>22</u>	<u>4</u>	<u>1</u>	<u>4</u>	<u>41</u>	<u>31</u>	<u>6</u>	<u>0</u>	<u>12</u>	<u>30</u>	<u>69</u>
<u>2003</u>	<u>57</u>	<u>29</u>	<u>4</u>	<u>1</u>	<u>8</u>	<u>43</u>	<u>36</u>	<u>6</u>	<u>1</u>	<u>0</u>	<u>21</u>	<u>64</u>	<u>50</u>	<u>9</u>	<u>2</u>	<u>6</u>	<u>18</u>	<u>50</u>
<u>2004</u>	<u>35</u>	<u>3</u>	<u>1</u>	<u>0</u>	<u>16</u>	<u>65</u>	<u>42</u>	<u>11</u>	<u>2</u>	<u>1</u>	<u>26</u>	<u>58</u>	<u>32</u>	<u>7</u>	<u>1</u>	<u>13</u>	<u>33</u>	<u>68</u>
<u>2005</u>	<u>60</u>	<u>18</u>	<u>1</u>	<u>0</u>	<u>4</u>	<u>40</u>	<u>48</u>	<u>15</u>	<u>2</u>	<u>0</u>	<u>11</u>	<u>52</u>	<u>33</u>	<u>4</u>	<u>0</u>	<u>2</u>	<u>19</u>	<u>67</u>
<u>2006</u>	<u>48</u>	<u>11</u>	<u>3</u>	<u>0</u>	<u>8</u>	<u>52</u>	<u>70</u>	<u>33</u>	<u>7</u>	<u>0</u>	<u>5</u>	<u>30</u>	<u>57</u>	<u>16</u>	<u>0</u>	<u>1</u>	<u>10</u>	<u>43</u>
<u>2007</u>	<u>30</u>	<u>6</u>	<u>1</u>	<u>0</u>	<u>22</u>	<u>70</u>	<u>69</u>	<u>25</u>	<u>5</u>	<u>1</u>	<u>6</u>	<u>31</u>	<u>29</u>	<u>3</u>	<u>1</u>	<u>7</u>	<u>26</u>	<u>71</u>
<u>2008</u>	<u>43</u>	<u>19</u>	<u>5</u>	<u>3</u>	<u>20</u>	<u>57</u>	<u>68</u>	<u>27</u>	<u>7</u>	<u>0</u>	<u>8</u>	<u>32</u>	<u>41</u>	<u>10</u>	<u>1</u>	<u>4</u>	<u>24</u>	<u>59</u>
<u>2009</u>	<u>24</u>	<u>6</u>	<u>0</u>	<u>1</u>	<u>31</u>	<u>76</u>	<u>73</u>	<u>23</u>	<u>9</u>	<u>0</u>	<u>5</u>	<u>27</u>	<u>27</u>	<u>4</u>	<u>0</u>	<u>3</u>	<u>25</u>	<u>73</u>
<u>2010</u>	<u>75</u>	<u>42</u>	<u>11</u>	<u>0</u>	<u>10</u>	<u>25</u>	<u>42</u>	<u>11</u>	<u>2</u>	<u>1</u>	<u>18</u>	<u>58</u>	<u>72</u>	<u>20</u>	<u>1</u>	<u>1</u>	<u>7</u>	<u>28</u>

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748 **Table 3.** The same as Table 2, but only for the years with a positive (negative) SCD
749 anomaly and only for the three major stable snow regions: Northeast China (78
750 stations), North Xinjiang (21 stations) and the Tibetan Plateau (63 stations).

751

Year	Northeast China						North Xinjiang						Tibetan Plateau					
	P	1SD	2SD	-2SD	-1SD	N	P	1SD	2SD	-2SD	-1SD	N	P	1SD	2SD	-2SD	-1SD	N
1957	<u>98</u>	<u>72</u>	<u>16</u>	0	0	2	<u>22</u>	0	0	2	<u>33</u>	<u>78</u>	<u>74</u>	<u>52</u>	<u>13</u>	0	4	<u>26</u>
1959	<u>2</u>	0	0	<u>15</u>	<u>73</u>	<u>98</u>	<u>88</u>	<u>38</u>	0	0	0	<u>12</u>	<u>37</u>	<u>11</u>	3	0	6	<u>63</u>
1960	<u>39</u>	<u>14</u>	1	0	<u>26</u>	<u>61</u>	<u>100</u>	<u>88</u>	<u>29</u>	0	0	0	<u>23</u>	0	0	3	<u>30</u>	<u>77</u>
1963	<u>11</u>	0	0	<u>6</u>	<u>41</u>	<u>89</u>	26	0	0	5	<u>26</u>	<u>74</u>	<u>20</u>	0	0	0	<u>28</u>	<u>80</u>
1965	<u>66</u>	<u>24</u>	0	1	<u>16</u>	<u>34</u>	<u>21</u>	0	0	0	<u>37</u>	<u>79</u>	<u>12</u>	4	0	4	<u>50</u>	<u>88</u>
1967	<u>16</u>	0	0	<u>14</u>	<u>59</u>	<u>84</u>	<u>78</u>	<u>22</u>	0	0	6	<u>22</u>	<u>23</u>	6	0	0	<u>15</u>	<u>77</u>
1969	<u>21</u>	1	0	<u>15</u>	<u>43</u>	<u>79</u>	<u>78</u>	<u>28</u>	0	0	6	<u>22</u>	4	0	0	6	<u>53</u>	<u>96</u>
1973	<u>89</u>	<u>60</u>	4	0	0	11	<u>42</u>	0	0	5	<u>11</u>	<u>58</u>	<u>36</u>	<u>11</u>	2	0	<u>21</u>	<u>64</u>
1974	<u>55</u>	<u>18</u>	0	3	<u>21</u>	<u>45</u>	5	0	0	<u>21</u>	<u>58</u>	<u>95</u>	<u>38</u>	3	0	2	<u>14</u>	<u>62</u>
1977	<u>73</u>	<u>32</u>	4	0	5	<u>27</u>	<u>95</u>	<u>74</u>	0	0	5	5	<u>36</u>	<u>19</u>	7	0	7	<u>64</u>
1980	<u>65</u>	<u>18</u>	1	0	8	<u>35</u>	<u>95</u>	<u>63</u>	5	0	0	5	<u>45</u>	<u>10</u>	2	0	3	<u>55</u>
1983	<u>62</u>	<u>23</u>	3	0	3	<u>38</u>	<u>26</u>	0	0	0	<u>21</u>	<u>74</u>	<u>95</u>	<u>60</u>	<u>19</u>	0	0	5
1988	<u>70</u>	<u>23</u>	0	0	3	<u>30</u>	<u>100</u>	<u>68</u>	<u>11</u>	0	0	0	<u>52</u>	<u>22</u>	5	0	2	<u>48</u>
1990	<u>40</u>	0	0	0	<u>11</u>	<u>60</u>	<u>32</u>	5	0	0	<u>21</u>	<u>68</u>	<u>81</u>	<u>41</u>	3	0	0	<u>19</u>
1994	<u>94</u>	<u>29</u>	1	0	0	6	<u>95</u>	<u>53</u>	0	0	0	5	<u>46</u>	<u>14</u>	2	0	<u>11</u>	<u>54</u>
1995	<u>33</u>	1	0	3	<u>15</u>	<u>67</u>	5	0	0	<u>21</u>	<u>74</u>	<u>95</u>	<u>75</u>	<u>42</u>	<u>11</u>	0	0	<u>25</u>
1998	4	0	0	<u>14</u>	<u>64</u>	<u>96</u>	<u>63</u>	5	0	5	<u>11</u>	<u>37</u>	<u>82</u>	<u>39</u>	<u>12</u>	0	0	<u>18</u>
2002	4	0	0	<u>19</u>	<u>63</u>	<u>96</u>	<u>26</u>	0	0	5	<u>21</u>	<u>74</u>	<u>22</u>	2	0	0	<u>15</u>	<u>78</u>
2008	7	0	0	<u>11</u>	<u>48</u>	<u>93</u>	5	0	0	5	<u>47</u>	<u>95</u>	<u>59</u>	6	0	2	<u>14</u>	<u>41</u>
2010	<u>92</u>	<u>69</u>	<u>17</u>	0	3	8	<u>100</u>	<u>67</u>	<u>11</u>	0	0	0	<u>15</u>	6	0	2	<u>50</u>	<u>85</u>

752
753
754
755
756
757
758
759
760
761
762
763
764
765

766 **Table 4.** Number of stations with SCD, SCOD, and SCED trends, number of stations
767 with relationships of SCD, SCOD, and SCED, respectively, with TBZD, number of
768 stations with relationship between SCD and MAT, and number of stations with
769 relationship between SCD and AO (~~352~~296 stations in total). All of them have two
770 significance levels, the 90% and 95%.

		<u>SCD</u>			<u>SCOD</u>			<u>SCED</u>		
		<u>95%</u>	<u>90%</u>	<u>I*</u>	<u>95%</u>	<u>90%</u>	<u>I*</u>	<u>95%</u>	<u>90%</u>	<u>I*</u>
<u>Trend</u>	<u>Positive</u>	<u>19</u>	<u>37</u>	<u>125</u>	<u>178</u>	<u>196</u>	<u>74</u>	<u>1</u>	<u>3</u>	<u>37</u>
	<u>Negative</u>	<u>26</u>	<u>35</u>	<u>99</u>	<u>5</u>	<u>8</u>	<u>18</u>	<u>72</u>	<u>103</u>	<u>153</u>
<u>TBZD</u>	<u>Positive</u>	<u>124</u>	<u>154</u>	<u>126</u>	<u>0</u>	<u>1</u>	<u>50</u>	<u>72</u>	<u>99</u>	<u>170</u>
	<u>Negative</u>	<u>1</u>	<u>1</u>	<u>15</u>	<u>61</u>	<u>87</u>	<u>158</u>	<u>0</u>	<u>2</u>	<u>25</u>
<u>MAT</u>	<u>Positive</u>	<u>0</u>	<u>2</u>	<u>22</u>						
	<u>Negative</u>	<u>114</u>	<u>148</u>	<u>124</u>						
<u>AO</u>	<u>Positive</u>	<u>31</u>	<u>45</u>	<u>90</u>						
	<u>Negative</u>	<u>33</u>	<u>48</u>	<u>113</u>						

771 (Note: I* for insignificant trends or relations)

772

773

774

775

776

777

778

779

780

781

782

783 **Figure Captions**

784 **Figure 1.** Locations of weather stations and major basins, mountains and plains
785 mentioned in the paper, overlying the digital elevation model for China.

786 **Figure 2.** Percentage of weather stations with different measurement lengths.

787 **Figure 3.** Annual mean snow cover days (SCDs) from 1980/81 to 2009/10 (a), and their
788 coefficients of variation (CV) (b).

789 **Figure 4.** Seasonal variation of SCDs; the number in the centre denotes annual mean
790 SCDs, the blue colour in the circle the SCDs for winter season, the ~~yellow-green~~
791 colour for spring, and the red colour for autumn.

792 **Figure 5.** SCD anomalies in 1957 (a), 2002 (b) and 2008 (c), anomaly of snow cover
793 onset date (SCOD) in 2006 (d), and anomalies of snow cover end date (SCED) in
794 1957 (e) and 1997 (f).

795 **Figure 6.** Trends of annual mean SCDs (a), SCOD (b), and SCED (c) from the ~~352-296~~
796 stations of more than ten annual mean SCDs with Mann–Kendall test, and
797 relationships among the SCD and day with temperature below 0°C (TBZD) (d), mean
798 air temperature (MAT) (e), and Arctic Oscillation (AO) index (f).

799 **Figure 7.** SCD variations at Kuandian (40°43' N, 124°47'E, 260.1 m) (a), Hongliuhe
800 (41°32' N, 94°40'E, 1573.8 m) (b), Gangcha (37°20' N, 100°08'E, 3301.5 m) (c) and
801 Shiqu (32°59' N, 98°06'E, 4533.0 m) (d), SCOD at Pingliang (35°33' N, 106°40'E,
802 1412.0 m) (e) and Weichang (41°56' N, 117°45'E, 842.8 m) (f), and SCED at Jixi
803 (45°18' N, 130°56'E, 280.8 m) (g) and Maerkang (31°54' N, 102°54'E, 2664.4 m) (h).

804 (The unit on the Y-axis in the figures e, f, g, h denotes the Julian day using 1st
805 September as reference).

806 **Figure 8.** Spatial distribution of SCOD (a) and SCED (b) based on the stations with an
807 average of more than ten SCDs.

808 **Figure 9.** SCD relationships with TBZD at Chengshantou (37°24' N, 122°41'E, 47.7 m)
809 (a), MAT at ~~Baicheng-Tieli~~ (~~4146°4759'~~ N, ~~81128°5401'~~E, ~~1229210.25~~ m) (b), and
810 AO index at Huajialing (35°23' N, 105°00'E, 2450.6 m) (c) and Tonghua (41°41' N,
811 125°54'E, 402.9 m) (d).

812

813

814

815

816

817

818


819

820

821

822

823




824

825

Figure 1


826



827

828

Figure 2




829

830

Figure 3


831



832

833

Figure 4




835

836

Figure 5

837

838




840

841

Figure 6

842

843



845


846

Figure 7

847

848

849




850

851

Figure 8

852



853

854

Figure 9

855