

Dear Editor and Reviewers,

Again, the manuscript faced considerable changes and we think that the manuscript has improved even more. We re-arranged the manuscript thoroughly as suggested by reviewer #3 and put all of the describing Figures of the Methods to the Results section. We also changed and unified the layout of the Figures slightly and also deleted Figure 3, since most information was also appearing in (old) Figure 5.

We added more evaluation of the final gridded product by comparing it to the INCA nowcasting system of the Austrian weather service. We think it was really worth the effort, since it clearly shows the benefit of our method and the added value.

We hope you agree and wait for your feedback.

The authors

Klaus Haslinger

Annett Bartsch

Reviewer #1

This revision addresses all my earlier comments well. However, I don't quite agree with the specific numerical approach taken in the new Figure 13. See below.

New Figure 13 and accompanying discussion: I think you have the right physical idea, but the math you give isn't quite right. Since DTR is *less* related to global radiation at high elevations, your reasoning would say that a *lower* value of Cadj should be found by the analysis, not a higher value (right?) Since we think of DTR*Cadj as an approximation of radiation in the HM method. Yet you find a very high value of Cadj at these times and places.

So, instead, I think the key mechanism is that the *absolute values* of DTR are so much lower in winter and/or high elevations, compared to "normal" evaporative environments in which the HM was developed. That is, for any given value of cloudiness (or sunshine hours or similar), DTR is much smaller in winter/high elevation than in normal environments (due to the free tropospheric ventilation / mixing mechanism you describe.) So this small DTR needs to be multiplied by a much higher value of Cadj to compensate for this and achieve the right value of radiation relative to Ra.

One way to see this would be to plot the *regression* coefficient of DTR against global radiation, instead of the correlation coefficient. If my idea above is right, this regression coefficient should be much smaller in high elevations (especially in winter) than in low elevations. Even better would be a zero-intercept ($y=mx$) regression of $\sqrt{\text{DTR}}$ against the ratio of global to extraterrestrial radiation, since this is the best mathematical match to the HM. (In the HM, $\sqrt{\text{DTR}}$ seems to be parameterizing this ratio.)

1 **Thank you for your valuable idea of displaying the regression coefficients instead of the**
2 **correlations. We deleted the Figure with the correlations and created a new one showing**
3 **the linear regression coefficient of the sqrt(DTR) against the ratio between**
4 **extraterrestrial and global radiation. The paragraph describing the Figure and**
5 **discussing the implications was also changed of course. We think that the new approach**
6 **is better explaining the physics behind, thanks again for this recommendation.**

7

Reviewer #3

This manuscript introduces a new approach to construct daily 1 km fields of reference evaporation. The Hargreaves method is improved and calibrated over space and time against Penman-Monteith. The methods are applied and analyzed over Austria from 1961 to 2013. While I find the method and description logic and appealing and the contribution thus relevant, I am not fully convinced by its presentation in the manuscript. Regarding this I have two major concerns/suggestions:

1) For the content I would recommend to have a separate part that is reserved for the evaluation of the product. This starts with including existing similar products (for instance the maps mentioned by reviewer #2) in the introduction over methods for comparison of the new products with the old ones and general plausibility (as done with the correlation between diurnal temperature range and global radiation in comparison to the Cadj values already) in the methods to their presentation in the results and discussion. This would lead to a clearer message regarding the novelty and the ability of the introduced method.

Thanks your comment. We now added a further evaluation of the final gridded product against another ET0 dataset. As we also replied to Reviewer #1 that the intention of this dataset was not to provide new long term climatological means of ET0, but rather to create ET0 fields on a daily time step. So we consider a comparison with climatological mean maps not useful and therefore aimed to compare our ET0 estimates with another high resolution gridded dataset, the ET0 of the INCA (Integrated Nowcasting through Comprehensive Analysis, Haiden et al., 2011) system. This dataset is limited going back in time, but provides a good basis for evaluating our ET0 estimates.

A whole new section was introduced in the results with a comparison of INCA and our dataset compared to station data.

2) The structure of the paper was not always clear. Often there is a mix between Methods and Results but also between Discussion and Results (see more detailed comments in the

1 specific comments below). I would hence recommend revising the structure of the document
2 substantially.

3

4 **We changed the whole structure as recommended. We think that the manuscript clearly**
5 **improved through these measures, thank you for this recommendation.**

6

7 Specific comments

8 P5L28- P6L14 These are Results not Method

9 **Shifted to Results**

10 P6L24- P7L2 Results not Method

11 **Shifted to Results**

12 P7L7 - L23 Results not Method

13 **Shifted to Results**

14 P8L3 from “As was shown...” to P8L23

15 For all the above indicated lines, I see that these are meant to be preliminary results that lead
16 to the further development of the study. However, it disrupted from reading and I had to
17 check several times which section I was reading actually. I would recommend moving all
18 these to Results. The Method section then could be structured to 1) which parts of the method
19 were needed for the development of the method, 2) the application of the method and then
20 finally 3) the evaluation of the method. 1) specifically, could then be structures like this: this
21 and that was tested to...and then mention that based on these results... further steps were
22 taken

23

24 **As mentioned before, the whole manuscript has been restructured. The methods section**
25 **was re-ordered, not referring to Figures which are in the Results section now. We also**
26 **added a paragraph on the evaluation with the INCA dataset**

27

28 P9L31-P10L10 should be reformulated to stress the benefit more clearly

1

2 **This passage was reformulated and text was added:**

3 **“Considering Austrian topography it comes clear that using a method like HM without**
4 **calibration has major impacts on the result. Using non-calibrated HM ET0 data for**
5 **rainfall-runoff modelling for example would introduce large errors and uncertainties.**
6 **Given the fact that gridded ET0 based on PM are only available for a rather short time**
7 **period from the INCA system, the AET dataset provides a sound alternative for ET0**
8 **estimates on a high spatial resolution covering the last 53 years.”**

9

10 P10L11 From here it would be great to have this evaluation part including the comparison to
11 existing maps as pointed out by reviewer #2, which could impact the discussion on Cadj, and
12 add credibility as well as general advantage of the method proposed to existing methods

13

14 **As mentioned above we extended the manuscript with an evaluation with INCA ET0**
15 **fields where we compared different error characteristics against station data (Bias,**
16 **RMSE, RE). We added a new Figure which shows exemplarily two fields of INCA and**
17 **AET on the same day and the associated station values. We also added a table showing**
18 **the monthly and all year round error characteristics of both datasets.**

19

20 P10L26-27 This sentence is not well connected to this paragraph, it would be linked better
21 in a paragraph dealing with this relationship, the consequent Caj values per day and grip point
22 and a comparison to other products and/or the pronounced events that were anyhow
23 exemplary used in the study

24

25 **Right. This sentence made not much sense at that passage, so we decided to delete it,**
26 **since it is not essential.**

27

28 P11L7-16 this is not Discussion but Results

We did not move this part of the text to the Results, since we think it is better placed in the Discussion. We think, that this part dealing with the reasons behind this altitude dependence is very well suited in the Discussions, since it is not a Result per se, but rather a *discussion* of the result, the implications and the characteristics behind the Hargreaves method. We kindly hope you accept our suggestion.

P12L10-16 Results not dicussion

Shifted from Discussion to evaluation part of Results section

Technical comments

There are many formulations that could be expressed easier and more concise, it would be advisable to have a native English person correct for the language in the manuscript. Some of these formulations I picked up below. Generally, the own results are presented in past tense.

P1L19 in -> over

Done

P1L20 the sole predictor -> the only predictor

Done

P1L22 the statistical -> a statistical

Done

P1L24 Having -> with these

Done

P1L26-29 Reformulate to be clearer

We reformulated in the following manner: “. This approach is opening opportunities to create high resolution reference evapotranspiration fields based only temperature observations, but being closest as possible to the estimates of the Penman-Monteith approach.”

1 P2L2 by the fluxed -> by fluxes

2 **Done**

3 P2L3 the latter -> evapotranspiration

4 **Done**

5 P2L5 “agriculture” does this imply science or the farmer?

6 **Agriculture was changed to agricultural sciences**

7 P2L24 “have been developed...data.” -> “as alternatives for regions where the input data is

8 not sufficient to used PM.”

9 **Done**

10 P2L25 add “one of these simpler methods” before “, the methods of Hargreaves...”; move “in

11 this paper” to the end of the sentence

12 **Done**

13 P2L27 by -> from

14 **Done**

15 P2L27-30 rewrite to make this clear: sth like: “Hence, HM is much broader applicable for

16 many regions, because temperature observations are dense and easily accessible.

17 Nevertheless, like most temperature based methods, HM has been developed for distinct

18 studies and regions representing also distinct climate conditions.”

19 **Done**

20 P2L31 add “temperature-based” before methods

21 **Done**

22 P3L1 to -> in; add “in those” before they

23 **Done**

24 P3L3 add comma after “In this paper” and “is presented” after ET0

25 **Done**

26 P3L5 on -> in?

27 **Done**

1 P3L8 similar results compared to PM -> results comparable to PM

2 **Done**

3 P3L12 and the interpolating -> followed by an interpolation of

4 **Done**

5 P3L13 in this paper -> here

6 **Done**

7 P3L22 to use the better -> at using the best

8 **Done**

9 P3L28 The foundation of the ET0 calculations is -> The ET0 calculations are based on

10 **Done**

11 P3L32 remove “and reaching down to the present day”; remove “the conduction of”

12 **Done**

13 P4L2 for -> to

14 **Done**

15 P4L3 to ensure -> at ensuring

16 **Done**

17 P4L4 whole -> full

18 **Done**

19 P4L6 (DEM). The SRTM DEM was also applied in this study.

20 **Done**

21 P4L9 remove the before HM

22 **Done**

23 P4L11 is -> are; change to “to calibrate the HM to PM on a monthly basis” (PM was already

24 introduced before)

25 **Done**

26 P4L12 among the Austrian domain and also comprise rather -> over Austria and cover

- 1 **Done**
- 2 P4L15 covering -> for
- 3 **Done**
- 4 P4L20 Remove “As explained above”
- 5 **Done**
- 6 P4L22 they -> these methods
- 7 **Done**
- 8 P5L25 the original publication of
- 9 **Done**
- 10 P6L28 remove underscore
- 11 **Done**
- 12 P6L30 remove “time of”, change “rather high” to “the highest”
- 13 **Done**
- 14 P9L1 remove “the” before higher; it -> This higher ET0
- 15 **Done**
- 16 P9L2 higher -> longer?; termed -> known?
- 17 **Done**
- 18 P9L4 Northwest no capital (also for the other cardinal directions in the manuscript); specify
19 the “it” not clear like this
- 20 **Done**
- 21 P9L6 remove “the” before ET0
- 22 **Done**
- 23 P9L7 again -> also; the higher proportion of -> longer
- 24 **Done**
- 25 P9L8 enhances indirectly ET0 -> enhance ET0 indirectly
- 26 **Done**

1 P9L9 remove exemplary

2 **Done**

3 P9L12 bringing -> which brought

4 **Done**

5 P9L13 This is -> These conditions were

6 **Done**

7 P9L14 nearly -> almost

8 **Done**

9 P9L24 Remove “on the other hand” or replace it with instead after “cool,”

10 **Done**

11 P9L28 “which is“ before equivalent

12 **Done**

13 P9L30 remove “the” before some

14 **Done**

15 P910 is generally above the original -> has generally a higher value than Hargreaves

16 original value; in -> during

17 **Done**

18 P10L29 add “was done” after analysis

19 **Done**

20 P10L31 will be -> is

21 **Done**

22 P10L32 are dropping -> drop

23 **Done**

24 P11L1-4 Change to “Numerous studies investigating the relationship between DTR and

25 radiation (Pan...) showed considerable correlations. For example,”

26 **Done**

- 1 P11L31 add “the coefficient” before C
- 2 **Done**
- 3 P12L24 -25 Split sentence in two
- 4 **Done**
- 5 P12L29 may be -> is
- 6 **Done**
- 7 P12L30 aims to provide -> provides; tries to combine -> combined
- 8 **Done**
- 9 Table 1, caption Figure 1: remove underscores
- 10 **Done**
- 11 Caption Figure 2 station-> stations remove for before the spread
- 12 **Done**
- 13 Figure 15 add July and January at a more pronounced place of the figure not just in the
- 14 caption similar to Fig 13
- 15 **Done**
- 16 |

Creating long term gridded fields of reference evapotranspiration in Alpine terrain based on a re-calibrated Hargreaves method

K. Haslinger¹, A. Bartsch¹

[1]{Central Institute for Meteorology and Geodynamics (ZAMG), Climate Research Department, Vienna, Austria}

Correspondence to: K. Haslinger (klaus.haslinger@zamg.ac.at)

Abstract

A new approach for the construction of high resolution gridded fields of reference evapotranspiration for the Austrian domain on a daily time step is presented. Gridded data of minimum and maximum temperatures are used to estimate reference evapotranspiration based on the formulation of Hargreaves. The calibration constant in the Hargreaves equation is recalibrated to the Penman-Monteith equation in a monthly and station-wise assessment. This ensures on one hand eliminated biases of the Hargreaves approach compared to the formulation of Penman-Monteith and on the other hand also reduced root mean square errors and relative errors on a daily time scale. The resulting new calibration parameters are interpolated ~~in~~-over time to a daily temporal resolution for a standard year of 365 days. The overall novelty of the approach is the use of surface elevation as the ~~sole~~-only predictor to estimate the re-calibrated Hargreaves parameter in space. A third order polynomial is fitted to the re-calibrated parameters against elevation at every station which yields ~~the~~-a statistical model for assessing these new parameters in space by using the underlying digital elevation model of the temperature fields. ~~Having~~-With these newly calibrated parameters for every day of year and every grid point, the Hargreaves method is applied to the temperature fields, yielding reference evapotranspiration for the entire grid and time period from 1961-2013. ~~With this approach it is possible to generate high resolution reference evapotranspiration fields starting when only temperature observations are available but re-calibrated to meet the requirements of the recommendations defined by the Food and Agricultural Organisation (FAO).~~ This approach is opening opportunities to create high resolution reference

- 1 | [evapotranspiration fields based only temperature observations, but being closest as possible to](#)
- 2 | [the estimates of the Penman-Monteith approach.](#)

1 Introduction

The water balance in its most general form is determined by ~~the~~ fluxes of precipitation, change in storage and evapotranspiration (Shelton 2009). Particularly for ~~the latter~~ evapotranspiration, measurement is rather costly, since it requires sophisticated techniques like eddy correlation methods or lysimeters. In hydrology as well as agricultural sciences the actual evapotranspiration as part of the water balance equation is mostly assessed from the potential evapotranspiration (PET). PET refers to the maximum moisture loss from the surface, determined by meteorological conditions and the surface type, assuming unlimited moisture supply (Lhomme 1997). Since surface conditions determine the amount of PET, the concept of reference evapotranspiration (ET₀) was introduced (Doorenbos and Pruitt, 1977). ET₀ refers to the evapotranspiration from a standardized vegetated surface (grass) under unrestricted water supply, making ET₀ independent of soil properties. Numerous methods exist for estimating ET₀; differences arise in the complexity and the amount of necessary input data for calculation.

A standard method, recommended by the Food and Agricultural Organisation (FAO; Allen et al. 1998), is the Penman-Monteith (PM) formulation of ET₀. There are of course countless other methods as thoroughly described in McMahon et al. (2013), but the PM equation is considered the most reliable estimate and serves as a standard for comparisons with other methods (Allen et al. 1998). PM is fully physically based and requires four meteorological parameters (air temperature, wind speed, relative humidity and net radiation). It utilizes energy balance calculations at the surface to derive ET₀ and is therefore considered a radiation based method (Xu and Singh 2000).

On the contrary, much simpler methods which use air temperature as a proxy for radiation (Xu and Singh 2001) ~~have been developed to overcome the shortcoming of PM of not having sufficient input data~~ are applied as alternatives for regions where the input data is not sufficient to use PM. In this paper, One of these simpler methods; the method of Hargreaves (HM, Hargreaves et al. 1985), is used in this paper. It requires minimum and maximum air temperature and extra-terrestrial radiation, which can be derived ~~by~~ from the geographical location and the day of year. ~~Though much easier to calculate, as temperature observations are dense and easily accessible, one has to be aware that the HM, among most temperature based estimates, are developed for distinct studies and/or regions, representing a rather distinct climatic setting~~ Hence, HM is much broader applicable for many regions, because temperature

1 [observations are dense and easily accessible. Nevertheless, like most temperature based](#)
2 [methods, HM has been developed for distinct studies and regions representing also distinct](#)
3 [climate conditions](#) (Xu and Singh, 2001). To avoid large errors, these [temperature-based](#)
4 methods need to undergo a recalibration procedure to make them applicable ~~to~~ [in](#) different
5 climatic regions than [in those](#) they were originally designed for (Chattopadhyay and Hulme
6 1997, Xu and Chen 2005).

7 In this paper, the method for constructing a dataset of ET0 [is presented](#) on a daily time
8 resolution and a 1 km spatial resolution based on the method of Hargreaves ~~is presented~~. The
9 HM is calibrated to the PM ~~on~~ [in](#) a station-wise assessment. Many studies describe re-
10 calibration procedures for ET0 estimations in general (Tegos et al., 2015; Oudin et al. 2005)
11 and for the HM in particular (Pandey et al. 2014; Tabari and Talaei, 2011; Bautista et al.,
12 2009; Gavilán et al. 2006) in order to achieve ~~similar results compared to PM results~~
13 [comparable to PM](#). There are also some studies describing methods for creating interpolated
14 ET0 estimates (e. g. Aguila and Polo, 2011; Todorovic et al, 2013). However, two main
15 methodological frameworks emerged for the interpolation of ET0 (McVicar et al., 2007): (i)
16 interpolation of the forcing data and then calculating ET0, or (ii) calculating ET0 at every
17 weather station ~~and the interpolating followed by an interpolation of~~ ET0 onto the grid. ~~In this~~
18 ~~paper~~ [Here](#) we follow the first approach and combine it with methods proposed by Tegos et al.
19 (2015) and Mancosu et al. (2014) which use spatially interpolated ET0 model parameters.
20 Gridded data of minimum and maximum temperatures are used as forcing fields for the
21 application of the Hargreaves formulation of ET0. The novelty of this study is the application
22 of elevation as a predictor for the interpolation of the re-calibrated HM calibration parameter.
23 Furthermore, these new calibration parameters are also variable in time, by changing day-by-
24 day for all days of the year. This approach goes a step further than the method of Aguilar and
25 Polo (2011) which derived one new calibration parameter for the dry and one for the wet
26 season of the year. [An evaluation of the final gridded product is carried out by assessing](#)
27 [different error metrics at grid points next to weather stations where PM ET0 is available, and](#)
28 [also by comparing the ET0 fields with those of the operational ET0 estimates based on INCA](#)
29 [\(Integrated Nowcasting through Comprehensive Analysis, Haiden et al. 2011\), the nowcasting](#)
30 [system of the Austrian weather service.](#)

31 The presented dataset aims ~~to use the better~~ [at using the best](#) of two worlds by (i) using a
32 method for estimating ET0 that is calibrated to the standard algorithm as defined by the FAO

and (ii) being applicable to a comprehensive, long-term forcing dataset~~and~~, on a high temporal and spatial resolution.

2 Forcing Data

~~The foundation of the ET0 calculations is~~The ET0 calculations are based on a high resolution gridded dataset of daily minimum and maximum temperatures calculated for the Austrian domain (SPARTACUS, see Hiebl and Frei 2015), whereas the actual data stretches beyond Austria to entirely cover catchments close to the border. SPARTACUS is an operationally, daily updated dataset starting in 1961~~and reaching down to the present day~~. For ~~the conduction of~~ the ET0 fields, the SPARTACUS temperature forcing is used for the period 1961-2013. The interpolation algorithm is tailored ~~for to~~ complex, mountainous terrain with spatially complex temperature distributions. SPARTACUS also aims ~~to ensure~~at ensuring temporal consistency through a fixed station network over the ~~whole full~~ time period, providing robust trend estimations in space. SPARTACUS uses the SRTM (Shuttle Radar Topography Mission, Farr and Kobrick 2000) version 2 Digital Elevation Model (DEM)~~, so the~~. The SRTM DEM is also applied in the present study.

SPARTACUS provides the input data for calculating ET0 following the Hargreaves method (HM, Hargreaves and Samani 1982, Hargreaves and Allen 2003). However, a recalibration of ~~the~~ HM is necessary to avoid considerable estimation errors. This is carried out in a station wise assessment. Data of 42 meteorological stations (provided by the Austrian ~~Weather~~ weather Service-service ZAMG) ~~is are~~ used to ~~monthly calibrate the HM to the Penman-Monteith Method (PM)~~calibrate the HM to PM on a monthly basis. Figure 1 shows the location of these stations, which are spread homogeneously ~~among the over~~ Austrian ~~domain~~ ~~and also comprise~~and cover rather different elevations and environmental settings (Table 1). Data of daily global radiation, wind speed, humidity, maximum and minimum temperatures ~~covering for~~ the period 2004-2013 are used to calculate ET0 simultaneously with HM and PM.

3 Methods

3.1 Estimating reference evapotranspiration

~~As explained above,~~ Numerous methods exist for the estimation of ET₀, which is defined as the maximum moisture loss from a standardized, vegetated surface, determined by the meteorological forcing (Shelton, 2009). ~~They~~ These methods can roughly be classified as temperature based and radiation based estimates (Xu and Singh, 2000, Xu and Singh, 2001, Bormann, 2011). Following the recommendations of the FAO (Allen et al. 1998) the radiation-based Penman-Monteith Method (PM) provides most realistic results and generally outperforms temperature based methods. The overall shortcoming of the PM is the data intense calculation algorithm which requires daily values of net radiation, wind speed, humidity, maximum and minimum temperatures. Data coverage for these variables is usually rather sparse, particularly if gridded data is required. ET₀ following the PM is calculated as displayed in Equation 1:

$$ET_{0-P} = \frac{0.408\Delta(R_N - G) + \gamma \frac{900}{T + 273} u_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34u_2)} \quad (1)$$

Feldfunktion geändert

where E is the reference evapotranspiration [mm day⁻¹], R_N is the net radiation at the crop surface [MJ m⁻² day⁻¹], G is the soil heat flux density [MJ m⁻² day⁻¹], T is the mean air temperature at 2 m height [°C], u₂ is the wind speed at 2 m height [m s⁻¹], e_s is the saturation vapour pressure [kPa], e_a is the actual vapour pressure [kPa]; giving the vapour pressure deficit by subtracting e_a from e_s; Δ is the slope of the vapour pressure curve [kPa °C⁻¹] and γ is the psychrometric constant [kPa °C⁻¹]. Given the time resolution of one day the soil heat flux term is set to zero. The calculation of the other individual terms of Equation 1 is described in Allen et al. (1998). It should be mentioned, that the original Penman-Monteith equation contains a “surface resistance” term, expressing the response of different vegetation types, which is set constant for FAO PM, since it uses a standardized vegetated surface.

In contrast to the radiation based PM, the HM is based on daily minimum and maximum temperatures (T_{min}, T_{max}). Hargreaves (1975) stated from regression analysis between meteorological variables and measured ET₀ that temperature multiplied by surface global radiation is able to explain 94 % of the variance of ET₀ for a five day period (see Hargreaves and Allen 2003). Furthermore, wind and relative humidity explained only 10 and 9 %

respectively. Additional investigations by Hargreaves led to an assessment of surface radiation which can be explained by extra-terrestrial radiation at the top of the atmosphere and the diurnal temperature range as an indicator for the percentage of possible sunshine hours. The final form of the Hargreaves equation is given by:

$$ET0_h = C(T_{mean} + 17.78)(T_{max} - T_{min})^{0.5} R_a \quad (2)$$

Feldfunktion geändert

where $ET0_h$ is the reference evapotranspiration [mm day^{-1}], T_{mean} , T_{max} and T_{min} are the daily mean, maximum and minimum air temperatures [$^{\circ}\text{C}$] respectively and R_a is the water equivalent of the extra-terrestrial radiation at the top of the atmosphere [mm day^{-1}]. C is the calibration parameter of the HM and was set to 0.0023 in the original [publication of Hargreaves et al. \(1985\)](#)~~publication.~~

Following these formulations the $ET0$ for all stations ~~was~~is calculated for the period 2004-2013. ~~Figure 2a shows, as an example, the daily time series of $ET0$ as derived by PM ($ET0_p$) and HM ($ET0_h$) in the year 2004 at the station Grossenzersdorf. The differences between those two are obvious as $ET0_p$ shows clearly higher variability, with $ET0_h$ underestimating the upward peaks in the cold season and downward peaks in the warm season. This feature is more noticeable in Figure 2b, which shows the monthly averages over all stations, indicating the spread among all 42 stations. Here, an underestimation of the $ET0_h$ compared to $ET0_p$ from October to April is counteracted by an overestimation between May and September. On the other hand, $ET0_h$ shows higher spread among stations compared to $ET0_p$ except for November to January.~~

~~These features are also reflected in the bias of $ET0_h$ compared to $ET0_p$ as can be seen in Figure 3a. The average monthly bias over all stations is negative in the cold season with largest deviations in February of 0.3 mm day^{-1} , compared to the peak average positive bias in June of 0.4 mm day^{-1} . The annual cycle of the Root Mean Squared Error (RMSE) of $ET0_h$ as displayed in Figure 3b shows peak values in summer mainly due to the higher absolute values in the warm season compared to wintertime. The RMSE in December is around 0.5 mm day^{-1} compared to 1.1 mm day^{-1} in July, showing some more spread in wintertime compared to summer.~~

3.2 Calibration

In order to achieve a meaningful representation of ET₀ by HM, an adjustment of the calibration parameter (C_{adj}) of HM is necessary, with respect to ET₀ derived from PM. This is carried out on an average monthly basis for every station by the following equation, as also proposed by Bautista et al. (2009):

$$C_{adj} = 0.0023 / (E_H / E_P) \quad (3)$$

Feldfunktion geändert

where C_{adj} represents the new calibration parameter of the HM, E_H is the original ET_{0_h} from HM, using a C of 0.0023 and E_P is the ET_{0_p} from PM. As a result, a new set of C values for every month and every station is available. [An analysis on the behaviour of \$C_{adj}\$ in space revealed rather strong altitude dependence, particularly in the cold season. This feature enables to estimate \$C_{adj}\$ in space for every grid point by using the underlying DEM of the temperature fields as a predictor.](#)

~~Figure 4 shows the adjusted C values for three exemplary stations. C_{adj} is generally higher in winter and autumn compared to the original value indicated by the dashed line at 0.0023. It is also obvious that at station Grossenzersdorf the original value is matching rather well to the C_{adj} from April to October, in the other months the adjusted values are clearly higher. On the contrary, at station Weissensee_Gatschach C_{adj} is lower than 0.0023 except for the months from November to February. At station Rudolfshuette Alpinzentrum the adjusted values are above the original ones all time of the year, reaching rather high values in wintertime of about 0.007. These results clearly underpin the necessity for a re-calibration of C in order to receive sound ET₀ from temperature.~~

~~After determining the values for C_{adj} the ET₀ was re-calculated with these new calibration parameter values (ET_{0_h.e}). For simplicity for this first assessment the monthly values of C_{adj} were used for all days of the month, no temporal interpolation was conducted. As a result, the monthly mean bias, as was shown in Figure 4a, is reduced to zero at every station. Furthermore, the RMSE has also slightly decreased by 0.1 to 0.2 mm day⁻¹, as can be seen in Figure 5a. The Relative Error (RE) has also decreased, from around 50 % to fewer than 40 % in January for example (cf. Figure 5b). The improvements regarding RE in summer are lower due to the higher absolute values of ET₀ in the warm season.~~

~~The complete monthly mean time series from 2004 to 2013 of ET_{0_p}, ET_{0_h} and ET_{0_h.e} for three stations are shown in Figure 6. At station Grossenzersdorf the underestimation of~~

~~ET0_h in winter is reduced as well as the overall underestimation at station Rudolfshütte Alpinzentrum. On the other hand, the overestimation in summer at station Weissensee Gatschaach is considerably reduced with ET0_h.e. These features in combination with the information on the altitude of the given stations provide some information on more general characteristics of C_{adj} and the effects of the calibration. It seems that there is an altitude-dependence of C_{adj} , which is displayed in more detail in Figure 7. It shows the monthly average C_{adj} for stations which were binned to distinct classes of altitude ranging from 100 to 2300 m in steps of 100 m. As already seen in Figure 4 as an example for three stations, C_{adj} is clearly higher in winter than the unadjusted value. From April to September C_{adj} is lower than 0.0023 up to altitudes of 1500 m.a.s.l., lowest values are visible in May to August between altitudes of 400 to 1000 m.a.s.l.~~

3.3 Temporal and spatial interpolation of the Hargreaves calibration parameter C_{adj}

~~The monthly adjusted calibration parameters are now interpolated in space and time in order to receive a congruent overlay of C_{adj} over the SPARTACUS grid for every day of year.~~ As a first step, the monthly C_{adj} values at every station are linearly interpolated to daily values to avoid stepwise changes and therefore abrupt shifts of C_{adj} between months. This is carried out for a standard year with length of 365 days. The result is a time series of daily changing values of C_{adj} over the course of the year, available for every station, stretching over different altitudes and therefore yielding 42 different annual time series of C_{adj} .

Subsequently the daily, station-wise values of C_{adj} are interpolated in space. The analysis of the C_{adj} -altitude relationship indicated non-linear characteristics, so a third order polynomial fit was chosen. Using the underlying DEM of the SPARTACUS dataset it is possible to determine adjusted calibration parameters for every grid point in space by this relationship. The polynomial fit is applied for every day of the daily interpolated station-wise C_{adj} values, since these are changing day by day as well. The result is a gridded dataset of C_{adj} for the SPARTACUS domain for 365 time steps from January 1st to December 31st. ~~As was shown in the previous section, C_{adj} changes with altitude. Figure 8 shows the adjusted calibration parameters plotted against altitude for the monthly means of C_{adj} . From this Figure it comes clear that this relationship is not linear. C_{adj} is decreasing from the very low situated stations until altitudes between 500 and 1000 m.a.s.l. Going further up C_{adj} increases and one could say it might be a linear increase, particularly in winter. On the other hand, looking at the summer months the station with the highest elevation (Sonnblick, 3106 m.a.s.l.) shows~~

Formatiert: Standard

1 somewhat lower or at least equal values of C_{adj} compared to the cluster of stations between
2 2000 and 2400 m.a.s.l. This feature indicates that the relationship above 1000 m.a.s.l. might
3 not be linear. Taking all this characteristics into account, a higher order polynomial fit was
4 chosen to describe the C_{adj} altitude relation. As shown in Figure 8 a third order polynomial fit,
5 indicated by the red line, is applied. Using the underlying DEM of the SPARTACUS dataset
6 it is possible to determine adjusted calibration parameters for every grid point in space by this
7 relationship. The polynomial fit is applied for every day of the daily interpolated station wise
8 C_{adj} values, since these are changing day by day as well. The result is a gridded dataset of C_{adj}
9 for the SPARTACUS domain for 365 time steps from January 1st to December 31st. Figure 9
10 shows two examples of C_{adj} distribution in space on January 1st (a) and July 1st (b).
11 Particularly in January the altitude dependence of the calibration parameter is clearly standing
12 out, showing rather high values of C_{adj} in the mountainous areas. In contrast to winter the
13 spatial variations in summer are smaller, only some central Alpine areas between 1000 and
14 3000 m.a.s.l. are appearing in somewhat different shading than the surrounding low lands.

15 Having these gridded C_{adj} values the $ET0_{h.c}$ is calculated for every grid point and day since
16 1961 to 2013. In the case of leap years the C_{adj} grid of February 28th is also used for February
17 29th. The final gridded product is termed AET (Austrian reference EvapoTranspiration
18 dataset) throughout the rest of the paper.

19 The AET fields are finally evaluated against station data and another $ET0$ product.
20 Unfortunately there is no long-term gridded dataset of $ET0$ for the Austrian domain, so we
21 used the $ET0$ of the nowcasting system INCA (Integrated Nowcasting through
22 Comprehensive Analysis, Haiden et al., 2011) which yields daily fields of $ET0$ based on PM
23 on 1 km grid resolution. INCA uses weather stations, remote sensing data, rainfall radar data
24 as well as DEM information to derive nowcasting fields of several meteorological variables.
25 INCA is operational for several years, but due to constant changes in data input quality and
26 other improvements we chose to use only the 5-year period from 2009-2013.

27 For the skill assessment of the AET dataset we calculate mean monthly values of mean bias,
28 Root Mean Squared Error (RMSE) and Relative Error (RE) of those grid points in AET as
29 well as INCA closest to a station with PM $ET0$.

4 Results

Figure 2a shows, as an example, the daily time series of ET₀ as derived by PM (ET_{0_p}) and HM (ET_{0_h}) in the year 2004 at the station Grossenzersdorf. The differences between those two are obvious as ET_{0_p} shows clearly higher variability, with ET_{0_h} underestimating the upward peaks in the cold season and downward peaks in the warm season. This feature is more noticeable in Figure 2b, which shows the monthly averages over all stations, indicating the spread among all 42 stations. Here, an underestimation of the ET_{0_h} compared to ET_{0_p} from October to April is counteracted by an overestimation between May and September. On the other hand, ET_{0_ph} shows higher spread among stations compared to ET_{0_hp} except for November to January.

These features are also reflected in the bias of ET_{0_h} compared to ET_{0_p} as can be seen in Figure 3a. The average monthly bias over all stations is negative in the cold season with largest deviations in February of 0.3 mm day^{-1} , compared to the peak average positive bias in June of 0.4 mm day^{-1} . The annual cycle of the Root Mean Squared Error (RMSE) of ET_{0_h} as displayed in Figure 3b shows peak values in summer mainly due to the higher absolute values in the warm season compared to wintertime. The RMSE in December is around 0.5 mm day^{-1} compared to 1.1 mm day^{-1} in July, showing some more spread in wintertime compared to summer. Figure 4 shows the adjusted C values for three exemplary stations. C_{adj} is generally higher in winter and autumn compared to the original value indicated by the dashed line at 0.0023. It is also obvious that at station Grossenzersdorf the original value is matching rather well to the C_{adj} from April to October, in the other months the adjusted values are clearly higher. On the contrary, at station Weissensee Gatschach C_{adj} is lower than 0.0023 except for the months from November to February. At station Rudolfshuette-Alpinzentrum the adjusted values are above the original ones all year round, reaching the highest values in wintertime of about 0.007. These results clearly underpin the necessity for a re-calibration of C in order to receive sound ET₀ from temperature observations.

For simplicity for this first assessment the monthly values of C_{adj} were used for all days of the month, no temporal interpolation was conducted. As a result, the monthly mean bias, as was shown in Figure 4a, is reduced to zero at every station. Furthermore, the RMSE has also slightly decreased by 0.1 to 0.2 mm day^{-1} , as can be seen in Figure 4b. The Relative Error (RE) has also decreased, from around 50% to fewer than 40% in January for example

(cf. Figure 45b). The improvements regarding RE in summer are lower due to the higher absolute values of ET0 in the warm season.

The complete monthly mean time series from 2004 to 2013 of ET0_p, ET0_h and ET0_{h.c} for three stations are shown in Figure 56. At station Grossenzersdorf the underestimation of ET0_h in winter is reduced as well as the overall underestimation at station Rudolfshuette-Alpinzentrum. On the other hand, the overestimation in summer at station Weissensee-Gatschach is considerably reduced with ET0_{h.c}. These features in combination with the information on the altitude of the given stations provide some information on more general characteristics of C_{adj} and the effects of the calibration. It seems that there is, which underpins an altitude-dependence of C_{adj} , which is displayed in more detail in Figure 67. It shows the monthly average C_{adj} for stations which were binned to distinct classes of altitude ranging from 100 to 2300 m in steps of 100 m. As already seen in Figure 34 as an example for three stations, C_{adj} is clearly higher in winter than the unadjusted value. From April to September C_{adj} is lower than 0.0023 up to altitudes of 1500 m.a.s.l., lowest values are visible in May to August between altitudes of 400 to 1000 m.a.s.l. As was shown in the previous section, C_{adj} changes with altitude. Figure 78 shows displays the adjusted calibration parameters plotted against altitude for the monthly means of C_{adj} . From this Figure it comes clear that this relationship is not linear. C_{adj} is decreasing from the very low situated stations until altitudes between 500 and 1000 m.a.s.l. Going further up C_{adj} increases and one could say it might be a linear increase, particularly in winter. On the other hand, looking at the summer months the station with the highest elevation (Sonnblick, 3106 m.a.s.l.) shows somewhat lower or at least equal values of C_{adj} compared to the cluster of stations between 2000 and 2400 m.a.s.l. This feature indicates that the relationship above 1000 m.a.s.l. might not be linear. Taking all this characteristics into account, a higher order polynomial fit was chosen to describe the C_{adj} -altitude relation. As shown in Figure 8 a third order polynomial fit, indicated by the red line, is applied. Using the underlying DEM of the SPARTACUS dataset it is possible to determine adjusted calibration parameters for every grid point in space by this relationship. The polynomial fit is applied for every day of the daily interpolated station-wise C_{adj} values, since these are changing day by day as well. The result is a gridded dataset of C_{adj} for the SPARTACUS domain for 365 time steps from January 1st to December 31st.

The results of the spatial interpolation of C_{adj} are displayed in Figure 89, where shows two examples of C_{adj} distribution in space are displayed; on January 1st (a) and July 1st (b).

Particularly in January the altitude dependence of the calibration parameter is clearly standing out, showing rather high values of C_{adj} in the mountainous areas. In contrast to winter the spatial variations in summer are smaller, only some central Alpine areas between 1000 and 3000 m.a.s.l. are appearing in somewhat different shading than the surrounding low lands.

The climatological mean (1961-2013) of the final AET fields is displayed in Figure 9.10a shows the climatological mean (1961-2013) of the daily ET0 fields over the whole domain.

Lowest daily mean values of below 1.5 mm day^{-1} are apparent on the highest mountain ridges of the main Alpine crest. Highest values of 2.4 mm day^{-1} and above are found in the eastern and southern low lands. Other spatial features are visible as well, for example the higher ET0 in the valleys in the far western part of Austria. ~~It-This higher ET0~~ is driven by the ~~higher~~ longer sunshine hours in these areas, which are also ~~termed-known~~ as “inner alpine dry valleys”, because rainfall approaching from the west is often screened by the mountain chains in the ~~Northwest~~northwest. In the ET0 estimate ~~it-this feature of less cloud cover and therefore longer sunshine durations~~ is reflected in the higher Diurnal Temperature Range (DTR), yielding larger values in that particular area. A similar characteristic is apparent in the very south of Austria. Here ~~the~~ET0 is higher as well, compared to topographically similar regions on the northern rim of the Alps. This is ~~again-also~~ connected to the ~~higher-proportion of longer~~ sunshine hours which enhances indirectly ET0 through higher DTR values.

Figure 9.10b shows ~~exemplary~~the ET0 field of August 8th 2013. For the first time on that particular day, temperatures reached above 40°C in Austria at some stations in the ~~e~~East and ~~South~~south. Values of ET0 are particularly high, reaching up to 7 mm day^{-1} in some areas in the ~~Southeast~~southeast. That day was also characterized by an approaching cold front, ~~bringing-which brought~~ rain, dropping temperatures and overcast conditions from the ~~West~~west. ~~This-is~~These conditions were featured as well in the ET0 field, showing a considerable gradient from ~~West-west~~ to ~~East~~east, with ~~nearly-almost~~ zero ET0 at the headwaters of the Inn River in the far ~~Southwest~~southwest of the domain. Furthermore, the implications of overcast conditions in the ~~West-west~~ with lower altitudinal gradients of ET0 compared to the ~~East-east~~ with sunny conditions and distinct gradients along elevation are visible.

July, the month with the highest absolute values of ET0 shows considerable variations in the last 53 years. As an example, the mean anomaly of ET0 in July of 1983 with respect to the July mean of 1961-2013 is displayed in Figure 10.11a. This month was characterized by a

considerable heat wave and mean temperature anomalies of +3.5 °C which also affected ET₀. The absolute anomaly of ET₀ reaches above 1 mm day⁻¹ with respect to the climatological mean in some areas. The relative anomaly is in a range between 10 to 30 % (Figure 10+c). ~~On the other hand,~~ July of 1979 was rather cool instead with temperatures 1.5 °C below the climatological mean and accompanied by a strong negative anomaly in sunshine duration, particularly in the areas north of the main Alpine crest. These characteristics implicated a distinctly negative anomaly of ET₀ in this particular month (Figure 10+b). The absolute anomaly stretches between 0 and more than -1 mm day⁻¹, which is equivalent to a relative anomaly of 0 to -30 % (Figure 10+d). The negative signal is stronger in the areas north of the Alpine crest, zero anomalies are found in ~~the some areas south of the main Alpine crest~~ in the south.

In Figure 11 the overall benefits of the re-calibration of the HM are revealed. It shows the mean ET₀ in July 2012, a month accompanied by a considerable heat wave at the beginning and an overall temperature anomaly of around +2 °C. In Figure 11b the ET₀ field of the original HM formulation without calibration is shown, and Figure 11a displays the results with re-calibration as described in this study. Overall, the gradient along elevation of ET₀ is larger in the non-calibrated field. Particularly in this time of the year with large absolute values, the re-calibration has a considerable impact, although C_{adj} in July is relatively small compared to winter. As shown before (cf. Figure 34), the ET₀ estimation using the original C is good for July in the very lowlands, since biases tend to be rather small. However, going to higher elevations, the overestimation of the original HM is rather pronounced. Mean biases reach +1 mm day⁻¹ or +30 % over large parts of the domain. This signal switches to negative biases of -0.5 mm day⁻¹ (-25 %) above 1500 m.a.s.l. Considering Austrian topography it comes clear that using a method like HM without calibration has major impacts on the result. Using non-calibrated HM ET₀ data for rainfall-runoff modelling for example would introduce large errors and uncertainties. Given the fact that gridded ET₀ based on PM are only available for a rather short time period from the INCA system, the AET dataset provides a sound alternative for ET₀ estimates on a high spatial resolution covering the last 53 years.

The overall performance of the final gridded dataset AET compared to the station wise PM estimates is displayed in Figure 12. 12a shows the monthly bias of the original HM ET₀ and the calibrated ET₀ of the nearest grid point. The bias is clearly reduced in nearly all months. However, in April, as the only exception, the bias of the calibrated grid point values

is larger than the bias of the original estimation. The biases concerning different levels of altitude are reduced as well, as can be seen in Figure 125b which shows the biases in July and Figure 125c displaying the biases in January.

A comparison between AET and INCA ET0 and station based PM ET0 is given in Figure 13, showing ET0 on two different days in summer 2013. The first example (Figures 13a and 13b) is June the 4th 2013, a day with mostly overcast conditions, lower than average temperatures of between 7 to 12 °C and high relative humidity, it was the time after a big flood event in northern Austria. AET is clearly overestimating ET0 by a median difference of +1 mm day⁻¹ across all stations as shown by the boxplot in Figure 13c. INCA has a median difference of nearly zero, although the spread is larger than in AET. Under the given circumstances AET cannot compete with INCA, which considers, through using PM, information on relative humidity, which might has a strong forcing on ET0 on that particular day, information that is not available in the AET estimate. Another example is July 23rd 2013 (Figure 13d and 13e) which characterized by temperatures ranging between 20 °C in the West and 29 °C in the east, accompanied by some rainfall in the West and South. ET0 in both AET and INCA range between 3 and 6 mm day⁻¹, although INCA shows a general overestimation with a median difference around +0.5 mm day⁻¹ (Figure 13f). On the other hand median differences of AET compared to stations are around zero. There might be some biases in the global radiation in INCA, which is derived based on sunshine duration estimates (blended remote sensing and station data) and a simple radiation model.

However, comparing error characteristics in AET and INCA against station data (Table 2) for the period 2009-2013 reveals only minor differences. The mean bias all year round is lower in INCA (0.03 mm day⁻¹) compared to AET (0.12 mm day⁻¹). Considering monthly mean values the spread is rather similar spanning -0.30 to 0.66 mm day⁻¹ in INCA and -0.17 to 0.80 mm day⁻¹ in AET. The highest monthly mean values are in both dataset found in April (AET: 0.80 mm day⁻¹, INCA: 0.66 mm day⁻¹) and May (AET: 0.79 mm day⁻¹, INCA: 0.51 mm day⁻¹). The RMSE is slightly lower in AET reaching maximum values in June of 1.42 mm day⁻¹ compared to INCA with 1.80 mm day⁻¹. The overall mean RMSE is 0.89 mm day⁻¹ in AET and 1.05 mm day⁻¹ in INCA. Concerning the RE the characteristics are similar to the bias and the RMSE, with only minor differences between AET and INCA. The RE in AET ranges between +35 % (April) and -15 % (November) and in INCA these are rather similar spanning +25 % (February) and -18 % (November).

Formatiert: Hochgestellt

Formatiert: Hochgestellt

5 Discussion

By comparing the characteristics of ET₀ based on HM and PM on a daily time step it came clear that a re-calibration of C within the formulation of Hargreaves follows distinct patterns. The values of C_{adj} show markedly variations in space and time (over the course of the year). It turned out, that a monthly re-calibration of C reveals an annual cycle of C_{adj}, with C_{adj} being close to the original value of 0.0023 in the warm season (April-October) and low elevations. Going to higher elevations, C_{adj} decreases until roughly 1000 m.a.s.l. Reaching altitudes above 1700 m.a.s.l., C_{adj} ~~is generally above the original 0.0023~~ has generally a higher value than Hargreaves' original value, particularly ~~in~~ during the cold season (November-March). This altitude dependency of the calibration parameter in HM is mentioned in Samani (2000), but the authors also claimed that this relationship may be affected by different latitudes. Aguila and Polo (2011) also found that the original HM using a C of 0.0023 underestimates ET₀ at higher elevations and defined a value of 0.0038 at an elevation of 2500 m.a.s.l. However, this altitude dependency of C turned out to be more complex, as we are able to display, showing a distinct variation throughout the year along with elevation. ~~So this relationship is used to derive C_{adj} values for every day of year and every grid point of the forcing fields.~~

To reveal the sources of this altitude dependence of C ~~we accomplished~~ some additional analysis was done. In general, the HM utilizes the Diurnal Temperature Range (DTR, T_{max} minus T_{min}) to mimic the amount of global radiation at the land surface. Clear sky conditions are usually associated with higher DTR. There ~~will be~~ is more heating during daytime due to large proportions of direct solar radiation, whereas at night time temperatures ~~are dropping~~ drop further down since the outgoing long-wave radiation is not reflected by clouds. ~~The connection between DTR and radiation is shown in numerous studies~~ Numerous studies investigating the relationship between DTR and radiation (Pan et al., 2013; Makowski et al., 2009; Bindi and Miglietta, 1991; Bristow and Campbell, 1984). ~~All these investigations showed, which show~~ considerable correlations. ~~For example~~ For example Makowski et al. 2009 reported a correlation coefficient of 0.87 of the annual means of DTR and solar radiation averaged over 31 stations across Europe.

Figure 14 ~~3~~ shows ~~the correlation of DTR and global radiation~~ the linear regression coefficients of the square root of DTR and Global Top-Of-Atmosphere (TOA) radiation ratio on a daily

time scale at the 42 stations used in this study. The idea is to get a better understanding of the parameterization embedded in HM, which tries to assess the amount of global radiation via the DTR and the TOA radiation. The coefficients show a distinct altitudinal dependency, particularly in winter. In January ~~the correlations are above 0.90 at some stations and the coefficients are~~ generally high at altitudes between ~~400-300~~ and ~~1000-1100~~ m.a.s.l. At higher elevations ~~the correlations~~ they are dropping considerably, getting ~~negative~~ slightly negative above 3000 m.a.s.l. at station Sonnblick. This altitude dependency is also apparent in the transitional season (c.f. Figure 14; April and October) although not as pronounced as in winter. ~~between 1500 and 2000 m.a.s.l.~~ In July the ~~correlations~~ coefficients are generally higher, roughly ranging between 0.15 and 0.30, with no change along altitude. ~~Apart from two stations the correlations lie between 0.45 and 0.98, but again accompanied by a decline with altitude, which is also seen in the year round correlations. Interestingly, the patterns of the correlations along altitude are rather similar to the C_{adj} patterns as can be seen in Figure 8. Therefore we think that the DTR global radiation nexus is the crucial point in the altitude dependence of C_{adj} .~~

The reasons for the ~~correlation~~ patterns in Figure ~~13-14~~ seem to be rooted in the lower atmospheric mixing ratios at the lowest stations, some of them located in, or nearby cities, which might dampen the DTR, although clear sky conditions are apparent. At moderate altitudes between 400 and 1500 m.a.s.l. the daily temperature amplitude is more dominantly driven by surface energy balance processes which reflects ~~the higher correlations~~ higher regression coefficients. Going further up, the proportion of the DTR which is determined by large scale air mass changes rises, as the station locations reach up above the planetary boundary layer into the free atmosphere, ~~causing considerably low correlations at higher elevations, particularly in winter.~~ So for any given value of cloudiness, DTR is much smaller in winter and high elevations than in low elevation environments where boundary layer processes are dominant. This means for yielding realistic values of global radiation relative to TOA radiation, a much higher C_{adj} value is needed to compensate for this.

Although these circumstances seem to be a drawback of the methodology, the overall effect is only minor. Figure ~~14-15~~ shows the HM ET0 in dependence of the DTR and the daily mean temperature. At low daily mean temperatures, between -10 and +10 °C, the contour lines determining the value of ET0 are rather steep. This implies that a change in DTR has only

1 minor effects on the ET₀ outcome, whereas a change in daily mean temperature is more
2 important.

3 However, the procedure of altering [the coefficient](#) C has also implications on the variability of
4 ET₀ on a daily time scale. As was visible in Figure 2a the variability of ET₀ based on HM is
5 lower than using PM. The presented re-calibration has only little effect on the enhancement of
6 variability. By scaling C, variability is slightly enhanced in those areas and time of the year
7 where C_{adj} is higher than 0.0023. This is the case for most of the time and widespread areas,
8 but there are regions or altitudinal levels where the opposite is taking place. As is visible in
9 Figure 8-6 areas up to 1500 m.a.s.l. show lower than original values of C_{adj} in the summer
10 months. There are particular areas in June between altitudes of 500 to 1000 m.a.s.l. that show
11 the largest deviation from the original value. In these areas variability is lower in the re-
12 calibrated version. On the other hand the benefit of an ET₀ formulation being unbiased
13 compared to the reference of PM may overcome these shortcomings.

~~14 The overall performance of the final gridded dataset compared to the PM estimates is
15 displayed in Figure 15. 15a shows the monthly bias of the original HM ET₀ and the calibrated
16 ET₀ of the nearest grid point. The bias is clearly reduced in nearly all months. However, in
17 April, as the only exception, the bias of the calibrated grid point values is larger than the bias
18 of the original estimation. The biases concerning different levels of altitude are reduced as
19 well, as can be seen in Figure 15b which shows the biases in July and Figure 15c displaying
20 the biases in January.~~

22 6 Conclusion

23 In this paper a gridded dataset of ET₀ for the Austrian domain from 1961-2013 on daily time
24 step is presented. The forcing fields for estimating ET₀ are daily minimum and maximum
25 temperatures from the SPARTACUS dataset (Hiebl and Frei 2015). These fields are used to
26 calculate ET₀ by the formulation of Hargreaves et al. (1985). The HM is calibrated to the
27 Penman-Monteith equation, which is the recommended method by the FAO (Allen et al.
28 1998). ~~¶~~ [This is done using](#) a set of 42 meteorological stations from 2004-2013, which have
29 full data availability for calculating ET₀ by PM. The adjusted monthly calibration parameters
30 C_{adj} are interpolated in time (resulting in daily C_{adj} for a standard year) and space (resulting in
31 C_{adj} for every grid point of SPARTACUS and day of year). With these gridded C_{adj} the daily
32 fields of reference evapotranspiration are calculated for the time period from 1961-2013.

1 This dataset ~~may be~~^{is} highly valuable for users in the field of hydrology, agriculture, ecology
2 etc. as it ~~aims to provide~~^{provides} ET0 in a high spatial resolution and a long time period. Data
3 for calculating ET0 by recommended PM is usually not available for such long time spans
4 and/or with this spatial and temporal resolution. However, the method presented in this study
5 ~~tries to combine~~^{combined} both strengths of long time series, high spatial and temporal
6 resolution provided by the temperature based HM and the physical more realistic radiation
7 based PM by adjusting HM.

10 **Acknowledgements**

11 The authors want to thank the Federal Ministry of Science, Research and Economy (Grant
12 1410K214014B) for financial support. We also like to thank Johann Hiebl for providing the
13 SPARTACUS data and for fruitful discussions on the manuscript. The Austrian Weather
14 Service (ZAMG) is acknowledged for providing the data of 42 meteorological stations. We
15 would also like to thank two anonymous reviewers for the valuable comments which
16 improved the manuscript substantially.

References

- Aguila, C., and Polo, M. J.: Generating reference evapotranspiration surfaces from the Hargreaves equation at watershed scale, *Hydrol. Earth Syst. Sci.*, 15, 2495–2508, 2011.
- Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration – Guidelines for computing crop water requirements, *FAO Irrigation and drainage paper* 56, Rome, 15 pp, 1998.
- Bautista, F., Bautista, D., and Delgado-Carranza, C.: Calibrating the equations of Hargreaves and Thornthwaite to estimate the potential evapotranspiration in semi-arid and subhumid tropical climates for regional applications, *Atmósfera* 22(4), 331-348, 2009.
- Bindi, M., and Miglietta, F.: Estimating daily global radiation from air temperature and rainfall measurements, *Clim. Change*, 1, 117-124, 1991.
- Bormann, H.: Sensitivity analysis of 18 different potential evapotranspiration models to observed climatic change at German climate stations, *Clim. Ch.*, 104, 729-753, 2011.
- Bristow, K. L., and Campbell, G. S.: On the relationship between incoming solar radiation and daily maximum and minimum temperature, *Agric. Forest. Meteorol.*, 31(2), 159-166, 1984.
- Chattopadhyay N. and Hulme M.: Evaporation and potential evapotranspiration in India under conditions of recent and future climate changes, *Agric. Forest. Meteorol.* 87, 55-74, 1997.
- Doorenbros, J., and Pruitt, O. W.: Crop water requirements. *FAO Irrigation and Drainage Paper* 24, Rome, 144 pp, 1977.
- Farr, T.G., Kobrick, M.: Shuttle Radar Topography Mission produces a wealth of data, *Amer. Geophys. Union Eos*, 81, 583-585, 2000.
- Gavilán, P., Lorite, I. J., Tornero, S., and Berengena, J.: Regional calibration of Hargreaves equation for estimating reference ET in a semiarid environment, *Agr. Water Manage.*, 81, 257-281, 2006.
- [Haiden, T., Kann, A., Wittmann, C., Pistotnik, G., Bica, B. and Gruber C.: The Integrated Nowcasting through Comprehensive Analysis \(INCA\) System and Its Validation over the Eastern Alpine Region, *Weather Forecast.*, 26, 166-183, 2011.](#)

1 Hargreaves, G. H., and Allen, R.: History and Evaluation of Hargreaves Evapotranspiration
2 Equation, *J. Irrig. Drain Eng.*, 129(1), 53-63, 2003.

3 Hargreaves, G. H., and Samani, Z. A.: Estimating potential evapotranspiration, *J. Irrig. Drain*
4 *Eng.*, 108(3), 225-230, 1982.

5 Hargreaves, G. H., and Samani, Z. A.: Reference crop evapotranspiration from temperature,
6 *Appl. Eng. Agric.*, 1, 96-99, 1985.

7 Hargreaves, G. H.: Moisture Availability and Crop Production, *Trans. ASABE*, 18 (5): 980-
8 984, 1975.

9 Hargreaves, G. L., Hargreaves, G. H., and Riley, J. P.: Irrigation water requirements for
10 Senegal River Basin, *J. Irrig. Drain. Eng.*, 111(3), 265-275, 1985.

11 Hiebl, J., and Frei, C.: Daily temperature grids for Austria since 1961 – concept, creation and
12 applicability, submitted to *Theor. Appl. Climatol.*

13 Lhomme, J.-P.: Towards a rational definition of potential evapotranspiration, *Hydrol. Earth*
14 *Sys. Sci.*, 1(2), 257-264, 1997.

15 Mancosu, N., Snyder, R. L., and Spano, D.: Procedures to Develop a Standardized Reference
16 Evapotranspiration Zone Map, *J. Irrig. Drain Eng.*, 140, A4014004, 2014.

17 Makowski, K., Jaeger E. B., Chiacchio, M., Wild, M., Ewen, T., and Ohmura, A.: On the
18 relationship between diurnal temperature range and surface solar radiation in Europe, *J.*
19 *Geophys. Res.*, 114, D00D07, 2009.

20 McMahon, T. A., Peel, M. C., Lowe, L., Srikanthan, R., and McVicar, T. R.: Estimating
21 actual, potential, reference crop and pan evaporation using standard meteorological data: a
22 pragmatic synthesis, *Hydrol. Earth Sys. Sci.*, 17, 1331–1363, 2013.

23 McVicar, T. R., Van Niel, T. G., Li, L., Hutchinson, M. F., Mu, X.-M., and Liu, Z.-H.:
24 Spatially distributing monthly reference evapotranspiration and pan evaporation considering
25 topographic influences, *J. Hyd.*, 338, 196-220, 2007.

26 Pan, T., Wu, S., Dai, E., and Liu, Y.: Estimating the daily global solar radiation spatial
27 distribution from diurnal temperature ranges over the Tibetan Plateau in China, *Ap. Energy*,
28 107, 384-393, 2013.

1 Pandey, V., Pandey, P. K., and Mahanta, A. P.: Calibration and performance verification of
2 Hargreaves Samani equation in a humid region, *Irrig. and Drain.*, 63, 659–667, 2014.

3 Samani, Z.: Estimating Solar Radiation and Evapotranspiration Using Minimum
4 Climatological Data (Hargreaves-Samani equation), *J. Irr. Drain. Eng.*, 126 (4), 265-267,
5 2000.

6 Shelton, M. L.: *Hydroclimatology*, Cambridge University Press, Cambridge, United
7 Kingdom, 2009.

8 Tabari, H. and Talaei, P.: Local Calibration of the Hargreaves and Priestley-Taylor Equations
9 for Estimating Reference Evapotranspiration in Arid and Cold Climates of Iran Based on the
10 Penman-Monteith Model, *J. Hydrol. Eng.*, 16(10), 837-845, 2011.

11 Tegos, A., Malamos, M., and Koutsoyiannis, D.: A parsimonious regional parametric
12 evapotranspiration model based on a simplification of the Penman–Monteith formula, *J. Hyd.*,
13 524, 708-714, 2015.

14 Todorovic, M., Karic, B., and Pereira, L. S.: Reference evapotranspiration estimate with
15 limited weather data across a range of Mediterranean climates, *J. Hyd.*, 481, 166-176, 2011.

16 Xu, C.-Y., and Chen D.: Comparison of seven models for estimation of evapotranspiration
17 and groundwater recharge using lysimeter measurement data in Germany, *Hydrol. Processes*,
18 19, 3717-3734, 2005.

19 Xu, C.-Y., and Singh, V. P.: Evaluation and generalization of radiation-based equations for
20 calculating evaporation, *Hydrol. Processes*, 14, 339-349, 2000.

21 Xu, C.-Y., and Singh, V. P.: Evaluation and generalization of temperature-based equations for
22 calculating evaporation, *Hydrol. Processes*, 14, 339-349, 2001.

23

1 Table 1. Location, altitude and setting of the 42 meteorological stations used for calibration.

	Station	Lon (°)	Lat (°)	Alt (m)	Setting
1	Aflenz	15.24	47.55	783	Mountainous
2	Alberschwende	9.85	47.46	715	Mountainous
3	Arriach	13.85	46.73	870	Mountainous
4	Bregenz	9.75	47.50	424	Lakeside
5	Dornbirn	9.73	47.43	407	Valley
6	Feldkirchen	14.10	46.72	546	Valley
7	Feuerkogel	13.72	47.82	1618	Summit
8	Fischbach	15.64	47.44	1034	Mountainous
9	Galzig	10.23	47.13	2084	Alpine
10	Graz <small>—</small> Universitaet	15.45	47.08	366	City
11	Grossenzersdorf	16.56	48.20	154	Lowland
12	Gumpoldskirchen	16.28	48.04	219	Lowland
13	Irdning <small>—</small> Gumpenstein	14.10	47.50	702	Valley
14	Ischgl <small>—</small> Idalpe	10.32	46.98	2323	Alpine
15	Jenbach	11.76	47.39	530	Valley
16	Kanzelhoehe	13.90	46.68	1520	Summit
17	Krems	15.62	48.42	203	Lowland
18	Kremsmünster	14.13	48.06	382	Lowland
19	Langenlois	15.70	48.47	207	Lowland
20	Lilienfeld <small>—</small> Tarschberg	15.59	48.03	696	Mountainous
21	Lofereralm	12.65	47.60	1624	Alpine
22	Lunz <small>—</small> am <small>—</small> See	15.07	47.85	612	Valley
23	Lutzmannsburg	16.65	47.47	201	Lowland

24	Mariapfar	13.75	47.15	1153	Mountainous
25	Mariazell	15.30	47.79	864	Mountainous
26	Neumarkt	14.42	47.07	869	Mountainous
27	Patscherkofel	11.46	47.21	2247	Summit
28	Poertschach	14.17	46.63	450	Lakeside
29	Retz	15.94	48.76	320	Lowland
30	Reutte	10.72	47.49	842	Valley
31	Rudolfshuette-Alpinzentrum	12.63	47.13	2304	Alpine
32	Schaerding	13.43	48.46	307	Lowland
33	Schmittenhoehe	12.74	47.33	1973	Alpine
34	Sonnblick	15.96	47.05	3109	Summit
35	Spittal-Drau	13.49	46.79	542	Valley
36	Villacheralpe	13.68	46.60	2156	Summit
37	Virgen	12.46	47.00	1212	Valley
38	Weissensee-Gatschach	13.29	46.72	945	Lakeside
39	Wien-Donaufeld	16.43	48.26	161	City
40	Wien-Hohewarte	16.36	48.25	198	City
41	Wien-Unterlaa	16.42	48.12	201	City
42	Wolfsegg	13.67	48.11	638	Lowland

1

2

1

Table 2. Error Characteristics of AET and INCA against station data

	Bias [mm/d]		RMSE [mm/d]		RE [%]	
	AET	INCA	AET	INCA	AET	INCA
<u>January</u>	<u>-0.01</u>	<u>-0.05</u>	<u>0.29</u>	<u>0.34</u>	<u>1</u>	<u>-7</u>
<u>February</u>	<u>-0.17</u>	<u>-0.30</u>	<u>0.60</u>	<u>0.65</u>	<u>-12</u>	<u>-25</u>
<u>March</u>	<u>0.04</u>	<u>-0.23</u>	<u>0.84</u>	<u>0.89</u>	<u>4</u>	<u>-14</u>
<u>April</u>	<u>0.80</u>	<u>0.66</u>	<u>1.34</u>	<u>1.59</u>	<u>35</u>	<u>28</u>
<u>May</u>	<u>0.79</u>	<u>0.51</u>	<u>1.38</u>	<u>1.58</u>	<u>29</u>	<u>19</u>
<u>June</u>	<u>0.19</u>	<u>-0.24</u>	<u>1.42</u>	<u>1.80</u>	<u>6</u>	<u>-8</u>
<u>July</u>	<u>0.39</u>	<u>0.31</u>	<u>1.29</u>	<u>1.58</u>	<u>12</u>	<u>9</u>
<u>August</u>	<u>-0.09</u>	<u>-0.01</u>	<u>1.16</u>	<u>1.42</u>	<u>-1</u>	<u>1</u>
<u>September</u>	<u>-0.14</u>	<u>-0.10</u>	<u>0.96</u>	<u>1.11</u>	<u>-6</u>	<u>-4</u>
<u>October</u>	<u>-0.15</u>	<u>-0.06</u>	<u>0.57</u>	<u>0.69</u>	<u>-8</u>	<u>-3</u>
<u>November</u>	<u>-0.03</u>	<u>0.01</u>	<u>0.43</u>	<u>0.54</u>	<u>2</u>	<u>5</u>
<u>December</u>	<u>-0.16</u>	<u>-0.18</u>	<u>0.39</u>	<u>0.43</u>	<u>-15</u>	<u>-18</u>
<u>Year</u>	<u>0.12</u>	<u>0.03</u>	<u>0.89</u>	<u>1.05</u>	<u>4</u>	<u>-1</u>

2

Formatierte Tabelle

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Zentriert

Formatiert: Englisch (Großbritannien)

1

2

3

4

5

6

7




Figure 1. Location of the meteorological stations used for calibration; coloured circles around points indicate stations that are exemplary displayed in other plots: Grossenzersdorf (blue), Weissensee-Gatschach (green) and Rudolfshuette-Alpinzentrum (red).




Figure 2. Daily time series of ET0 in 2004 for ET0 based on PM (ET0_p) and HM (ET0_h) at the station Grossenzersdorf (a); Monthly mean ET0 from 2004 to 2013 averaged over all stations, error bars denote ~~for~~ the spread among all stations (b).




Figure 3. Monthly Bias (a) and monthly Root Mean Square Error (b) between daily ET0_p and ET0_h for all stations; the grey shading indicates the spread among the different stations.





Figure 34. Monthly values of C_{adj} at three different stations, the dashed black lines indicates the original C value of 0.0023 from Hargreaves et al. (1985).




Figure 45. Monthly Root Mean Square Error (a) and monthly Relative Error (b) between daily ET0_p and ET0_h (black) and ET0_p and ET0_h.c (red).









Figure 56. Monthly ET0 sums derived from ET0_p, ET0_h and ET0_h.c for three stations located at different altitudes.





Figure 67. Monthly variations of C_{adj} with respect to altitude; the black contour line defines the original Hargreaves Calibration Parameter C value of 0.0023; stations are binned to classes of altitude from 100 to 2300 m every 100 m; white areas denote classes of altitude with no station available.






Figure 78. Station-wise monthly third-order polynomial fit of the Hargreaves Calibration Parameter C_{adj} against altitude; the blue dotted line indicates the original C value of 0.0023.




Figure 89. Spatially interpolated C_{adj} values for January 1st (a) and July 1st (b).




Figure 9+0. Climatological daily mean ET0 from 1961-2013 (a); example of a daily field of ET0 on August 8th 2013 (b).




Figure 10.4. Upper panel: absolute anomalies of ET0 sum in July 1983 (a) and July 1979 (b) with respect to the climatological mean in July from 1961-2013; lower panel: corresponding relative anomaly (c, d).






Figure 11.2. July 2012 monthly mean ET0 based on C_{adj} values – ET0_h.c (a), using the original C of 0.0023 for the whole grid ET0_h (b) and the corresponding absolute (c) and relative bias (d); the dots in (a) and (b) denote for the PM ET0 at the stations.



Formatiert: Block, Abstand Vor: 6 Pt.,
Zeilenabstand: 1.5 Zeilen

Figure 12. Boxplots of monthly mean bias of the station-wise original Hargreaves ET0 (grey) and the AET, re-calibrated ET0 (red) against Penman-Monteith ET0 (a); stratified by different classes of altitude in July (b) and January (c).



Formatiert: Block, Abstand Vor: 6 Pt.,
Zeilenabstand: 1.5 Zeilen

Figure 13. ET0 fields of AET (a, d) and INCA (b, e) and station wise PM ET0 on June 4th 2013 and July 23rd 2013 and corresponding differences at grid points closest to a station with PM ET0 of both datasets displayed as boxplots (c, f).

Formatiert: Hochgestellt

Formatiert: Hochgestellt




Figure 143. Station-wise ~~Correlation-linear regression coefficient~~ of ~~the TOA radiation -~~ Global Radiation ~~ratio and~~ ~~against the square root of the~~ Diurnal Temperature Range (T_{\max} - T_{\min}) against altitude represented by black dots in January, ~~April, July and October.~~ ~~(left), July~~ ~~(middle) and all year (right); the red line represents a third order polynomial fit.~~




Figure 154. ET0 response to varying Daily Mean Temperature and Diurnal Temperature Range; ET0 values are calculated with 1st of April Top of the Atmosphere Radiation and the original C value of 0.0023.




Figure 15. Boxplots of monthly mean bias of the station-wise original Hargreaves ET0 (grey) and the final gridded, re-calibrated ET0 (red) against Penman-Monteith ET0 (a); stratified by different classes of altitude in July (b) and January (c);