

## Revision guide:

This document contains two parts: part I addresses the comments from the referees and our comments to them including our changes on the manuscript. Part II starting at page 9 contains the marked-up manuscript that compares the current revision to the re-submitted manuscript of the previous iteration.

### 1. Point-by-point reply to the comments

We thank Daphné Freudiger and the two further referees for their comments and remarks and would like to answer the general remarks in detail. The specific remarks have been applied (see marked-up manuscript). First we reply to referee #1. The comments of referee #2 are written in *blue italic*, the comments of Daphné Freudiger are formatted in *orange italic*. Our replies are formatted in black, while changes on the manuscript are formatted in *black italic* and covered in quotation marks.

#### Response to referee #1

We thank the referee #1 for his comment on deleting the sentence about the errors of INCA in 2003 and 2004. We deleted this sentence.

#### Response to the referee #2

We thank referee #2 for the response and suggestions. In the following, we will respond to every point separately.

*General comments: The authors have put in a lot of effort in improving the manuscript. I am especially pleased to the analysis on parameter uncertainty, although I found it difficult to understand the conclusions. I think you need to explain more clearly how you can state “narrower boundaries of the snowmelt factors” (p.14 l28). To me the analysis shows clearly a huge problem of equifinality in the model. 6 calibration parameters just in the snow module is a lot.*

Most modelers tend to use snowmelt factors up to about 8 or 10 mm °C<sup>-1</sup> d<sup>-1</sup> (sometimes even higher) in high elevations to control snowmelt in these altitudes. The physical basis of the snowmelt factor however does not allow for that as shown by Kling et al. (2006). In their paper, Kling et al. (2006) derived meaningful values for the snowmelt factor on a 1x1 km grid in Austria by accounting for solar radiation, aspect of the slope and elevation of the cell of interest at both the summer and winter solstice. These two values than can be interpreted as the upper and lower boundaries of the snowmelt factor per cell. They found values ranging from 1.2 to 2.2 for the lower and 2.0 to 3.0 mm °C<sup>-1</sup> d<sup>-1</sup> for the upper boundary. Our model (model A) allows the modeller to use snowmelt factors in this range and achieve a good model performance regarding both snow accumulation and runoff. Model B however produces these snow towers mentioned in our paper and would need much higher values to account for this problem (not to mention that high snowmelt factors alone are not capable of solving the problem due to less days with positive air temperature values during the year). Consequently, one can conclude that model A allows the modeller to use “narrower boundaries of the snowmelt factors” than model B does while giving good results.

As every model using parameters that cannot be derived physically (and even such), the model suffers from equifinality issues. Obviously, the more parameters need to be estimated during calibration, the greater the problem. However, Gharari et al. (2014) showed that this might not necessarily be true if the parameters can be estimated using experts knowledge. In our opinion, this is the case in model A.

We edited the manuscript and added the paragraph 5.5 in the discussion concerning equifinality:

*“Like most hydrological models COSERO requires calibration of some parameters. This necessarily causes equifinality issues (Beven and Freer, 2001). The more adjustable parameters a model provides, the more important this problem may become (e. g. Gupta et al., 2008). On the other hand, some authors pointed out that more complex models may produce more feasible results if the parameters can be estimated within realistic boundaries (Gharari et al., 2012, 2014; Hrachowitz et al., 2014). Applying COSERO with the presented snow redistribution routine requires two additional parameters: the vegetation threshold HV (estimated a priori) and the calibration parameter C (see Eq. 12). Yet, accounting for snow redistribution allows the modeller to use DU values within or close to the range proposed by Kling et al., (2006), while the standard version of the model leads to the best results if higher and therefore unrealistic DU values are used (see supplements of this article).”*

*However, I do not think this is the main problem of the paper, namely that the authors model a process at a spatial scale for which it has not been observed.*

In this point we disagree with referee #2. As pointed out by Liston (2004) and in our introduction, snow transport occurs at several spatial scales ranging from micro, e.g. several meters, to macro scales, e.g. several kilometres. The process forcing snow to be redistributed clearly is different at different scales. But at the scale of 1x1 km lateral snow transport does occur driven by a. wind and b. gravitation (e.g. avalanches, snow drift etc). But even if snow transport occurs only at smaller scales as referee #2 suggests, the 1x1 km scale is affected by that, too. If snow is transported only several 10s to 100s of meters, some of snow passes the border between two 1x1 grid cell sizes and consequently ends up in the neighbour cell of its origin.

However, as stated in the introduction of our paper, our model is a conceptual model not trying to account for the physical process of snow mobilisation. This, we agree with referee #2, would be very hard to model on such that spatial scale. In fact, also the temporal scale would be inefficient for such a model approach.

*In the previous review I suggested a few explanations to why such a problem is present. The authors only considered that of meteorological input, and not that of the spatial frequency distribution of SWE. In several rainfall- runoff models (including this one) snowfall is distributed lognormally, often with a fixed coefficient of variation (i.e which gives it a fixed skewness). The problem with this is that the new snowfall is distributed such that (a unrealistic) perfect spatial correlation is implicitly assumed, i.e. the fractional area receiving the most snow from the first event, also receives the most snow in future events. In this way, the spatial distribution of accumulated snow has the same skewness as the individual events (often skewed too much to the left), and the quantiles (i.e. spatial fractions, or here, snow classes) that continuously receive maximum snow will eventually build these “snowtowers”. The paper is not detailed enough so that I can claim this to be the problem the COSERO model, but I suggest the problem of snow distribution to be investigated as an alternative to*

*the presented model which suggest snow redistribution processes at a spatial scale for which they have not really been observed.*

The lognormal distribution of snowfall leads to high accumulations in the most left quantile (in this case the most right snow class (see fig. 2b) of the distribution function. Editing the distribution function can be done by adjusting parameter  $N_{VAR}$  which we did in calibrating the model and in the uncertainty analysis shown in the appendix of the paper. However, even if snowfall is evenly distributed throughout the five quantiles of a single grid cell, i.e. is equally distributed, snow will accumulated in elevations, where the air temperature scarcely rises above 0°C.

*The introduction unfortunately appears as a rather patchy review and not really leading to why the proposed model is a good idea. Some discussion of scale is present, but the proposed model with its 1X1 km scale comes as a surprising conclusion.*

We disagree in this point with referee #2. The introduction, in our opinion, gives an insight of what has been done in solving the problem of snow accumulations and shows that, on small scales of 10s to 100s of square meters there are several approaches leading to good results. When applying models using coarser cell sizes, this problem exists as well.

Nevertheless, we edited the introduction and added some paragraphs:

*“Scipi3n et al. (2013) however identified significant discrepancies between precipitation patterns obtained by a Doppler X-band radar and the final seasonal snow accumulation at the end of the winter period which gives clear indications that snowfall is redistributed on different driving forces. Consequently, the variability of the meteorological input (alone) cannot explain the variability of snow cover patterns.”*

...

*“Due to some available databases for vegetation and meteorology (Haiden et al., 2011; Masson et al., 2003; Oubeidillah et al., 2014), many models operate on cell sizes of 1 km<sup>2</sup> or more (e. g. Andersen et al., 2001; Henriksen et al., 2003; Mauser and Bach, 2009; Safeeq et al., 2014). To our knowledge, no model for redistributing snow on a 1x1 km grid size exists. In this paper we present a simple approach to deal with snow in high mountainous regions and its application in the catchment of 3tztaler Ache in Tyrol, Austria. Since the model uses meteorological input from INCA (Haiden et al., 2011) that already account for meteorological corrections, we focus on snow redistribution rather than to edit the input data. The paper has two foci: a. to achieve a better model efficiency regarding runoff and b. to avoid the existence of snow towers at high altitudes.”*

*Can you please be more specific in your description of “snow classes”. I understand now (I think) it is quantiles of the lognormal distribution, but it really took me quite some time.*

We edited the paragraph concerning the log-normal distribution within paragraph 2.1:

*“Numerous studies have shown that sub-grid variability of snow depths can be described by a two parameter log-normal distribution (e. g. Donald et al., 1995; Pomeroy et al., 1998). COSERO uses five snow classes per cell (i.e. the log-normal distribution is subdivided into five quantiles) to approximate this sub-grid log-normal distribution under accumulation conditions (see Fig. 2 b)), i. e. snowfall is*

*distributed log-normally into snow classes., where the sum of the snow water equivalent (SWE) of each classes represent the mean conditions in the grid cell. [...]"*

*The notations need a make over, for example,  $S\_SWEA$  (Eq. 12) does not do. You can also use  $S\_SWE(t)$  for example in eq. 1. Or why not  $SWE(t)$  ?*

We would have liked to keep the notations more simple, but unfortunately, the editor suggested this way of notations. Changed  $S_{SWEA}$ , to  $S_{SWE(A)}$ .

Response to Daphné Freudiger (referee #3):

We thank Daphné Freudiger for her comments. In the following, we will respond to every point separately.

*(1) Aims of the study: Throughout the paper, the authors focus on saying that the main goal of the study is to better model the discharge. In fact, the improvement of model efficiency achieved by the simulation with snow distribution compared to the usual model is very little for discharge (KGE 0.92 vs 0.9). In my opinion, the main improvement achieved by the model is the elimination of “snow towers” and the better representation of the snow cover according to the efficiency coefficient  $r^2$  (0.66 vs 0.74), which can indicate, as suggested in the conclusion, that the model works right for the right reasons. Therefore I suggest that the authors focus their discussion more on the improved modelling of the snow cover, snow melt, and glacier melt and less on the improvement of the simulated discharge. This would also allow being more optimistic regarding the potential of such a conceptual model, since it is stated in the discussion that snow redistribution is not important for discharge modelling at the large scale (p.13, l.20-21).*

We agree that the main achievement of the model is in eliminating the snow towers in the summit regions. In comparison with MODIS data, the improved model leads to a coefficient  $r^2$  of 0.74 instead of 0.66. We agree on emphasize the improvements of the model regarding of snow cover, snowmelt and glacier melt in the discussion. However, we think that the aim of the study still remains on modelling discharge. If the focus would be on the improvement of snowcover, the methodology would differ (e.g. a more detailed description of processes in and on the snow layer).

We added to the results (4.3):

*“[...] It can be shown that net loss is evident in the zones of ridges and high elevations, where the maximum net gain is along the valley bottoms.”*

And to the discussion:

Paragraph 5.3:

*“[...] The resulting net loss and gain areas shown in Fig. 11 give some indication that redistribution algorithm is plausible.”*

Paragraph 5.4:

*“The model provides results that have been found by other models, too. For instance the elevation where the highest snow accumulations occurs (2800 to 3000 m a.s.) as was found by LiDAR measurements in the same catchment (Helfricht et al., 2012) as well as by modelling (Frey, 2015). [...]”*

*(2) Transferability of the method: The presented method showed very interesting results in the Ötztaler Ache catchment in Austria, but was it also tested in other catchments? Can you find similar results in other alpine catchments with different topographical properties, land cover, etc.? How transferable is this model? Since the model is conceptual and therefore should not need long calculation time, I suggest that the authors support their findings and test the model in 2-3 other catchments or at least discuss the transferability of their model in the discussion. This would add validity to the presented study.*

In this point we disagree with Daphné Freuding. As you state in your comment, the model is conceptual. This means, as long as a catchment provides high elevations and rather steep slopes, the model will work and eliminate snow towers. Of course, it would need, as all conceptual models do, some calibration. We think that modelling and presentation of any additional catchment does not improve the manuscript. In fact, we think it would make the paper rather difficult to read since it would a. make the article longer and b. provide additional data with the same conclusions.

We agree to discuss transferability of the method to other catchments. We added the following paragraph:

#### *“5.4 Transferability to other catchments*

*The model provides results that have been found by other models, too. For instance the elevation where the highest snow accumulations occurs (2800 to 3000 m a.s.) as was found by LiDAR measurements in the same catchment (Helfricht et al., 2012) as well as by modelling (Frey, 2015). Given that and the needs of the model (slope angles, snow density) for transporting snow, it produces valid results as long as a catchment features relatively steep slopes in the summit regions (which is the case in most catchments in the Alps). Obviously, the model needs calibration if it is transferred to another catchment.”*

*(3) Reliability of the model: The authors should discuss how their results are situated compared to other studies. Are their results reliable? How did other studies model snow redistribution? Do they come to similar results (physically-based as well as conceptual model)? If not, why not? The model results for example show enhanced glacier melt (100 mm/yr) because of the glacier area being snow free earlier during the year (p. 12, lines 16-17; p.14, line 10-12, fig.8). Are 100 mm/year in agreement with mass balances studies made on glaciers in the catchment or in catchment with similar properties? A comparison of the results with other studies related on snow redistribution would improve the reliability of the results and therefore also of the model.*

We think that reliability of the model is given by the fact, that the model is able to reproduce both the discharge and the snow cover extent (MODIS) more accurate than the model without the snow redistribution routine does. In addition fig. 11 shows reliable zones of loss and gain of snow due to the redistribution routine. Of course, small scale (10s to 100s of square meters)

variations as sinks or shelters cannot be modelled. This, however, is not the purpose of the model, but may lead to other results regarding the mass balance of glaciers. Bernhardt et al. (2010), for instance, found a surplus snow blown on the “Blau eisferner” when they used a spatial resolution of 30 m but didn’t find this when they used the coarser resolution of 300 m.

In the added paragraph 5.4 (concerning transferability) we describe some results that match our findings. These results were found by both measurements and models.

*(4) Structure of the paper: The structure of the paper presents some weaknesses that need to be improved:*

*a. Overall, the introduction makes a good review of the snow transport processes and the existing models. However this section needs to be restructured more logically. For example in section 1.1 processes responsible for the variability of the snow cover are described, that are not all snow transport processes. I suggest that the authors rename this section accordingly. As other processes are also involved in the variability of the snow cover, they should better explain why their model focuses on the snow redistribution. The objectives of the study need to be reformulated (see above, point 1).*

We agree on that point with the referee #3 and applied some changes to the introduction. See the marked-up manuscript.

*b. The authors should pay attention not to put results or part of the discussion or the methods in the figure captions. The figure captions should describe the figure but not analyze it. For example fig. 7: “Specific runoff at the outlet at Huben modelled with (model A) and without (model B) using the snow redistribution routine.” This is enough for the caption of the figure, the rest “In the early snow melt period, more runoff is generated by model A because snow accumulates rather in lower than in higher levels. In summer, enhanced glacier melt leads to more runoff by model A.” is part of the results and should be moved to the results section. The caption of fig. 2 to fig. 11 should be revised.*

We agree on that point with the referee #3 and applied some changes to the introduction. See the marked-up manuscript.

*c. The conclusion should be reformulated. It should summarize the main findings of the study and relate to the importance of the work. Part of the conclusion should be moved to the discussion. The second paragraph for example (p. 14, lines 13-20) is rather a discussion of the results than a conclusion. This is also the case for the last paragraph (p.14, lines 26-29). In comparison, paragraph 3 belongs perfectly to the conclusion since it points out the contribution of the paper and the further work that needs to be done to have better hydrological models.*

It always is a matter of personal preference what exactly belongs to the discussion and what is a conclusion. While we agree on the second paragraph, we do not agree on the last paragraph. This is a short link to the supplements of the article and the demand for further work on the problem of equifinality. We moved the second paragraph to the discussion.

*d. The language of the paper can be improved. Especially, long sentences should be avoided, since it makes them difficult to read and to understand.*

We agree and edited sentences throughout the whole paper. See marked-up manuscript.

## References

Bernhardt, M., Liston, G. E., Strasser, U., Zängl, G. and Schulz, K.: High resolution modelling of snow transport in complex terrain using downscaled MM5 wind fields, *Cryosph.*, 4(1), 99–113, doi:10.5194/tc-4-99-2010, 2010.

Gharari, S., Hrachowitz, M., Fenicia, F., Gao, H. and Savenije, H. H. G.: Using expert knowledge to increase realism in environmental system models can dramatically reduce the need for calibration, *Hydrol. Earth Syst. Sci.*, 18(12), 4839–4859, doi:10.5194/hess-18-4839-2014, 2014.

Kling, H., Fürst, J. and Nachtnebel, H. P.: Seasonal, spatially distributed modelling of accumulation and melting of snow for computing runoff in a long-term, large-basin water balance model, *Hydrol. Process.*, 20(10), 2141–2156, doi:10.1002/hyp.6203, 2006.

Liston, G. E.: Representing Subgrid Snow Cover Heterogeneities in Regional and Global Models, *J. Clim.*, 17(6), 1381–1397, doi:10.1175/1520-0442(2004)017<1381:RSSCHI>2.0.CO;2, 2004.

Figure changed (see next page):






Figure 6. Elevation levels of the Ötztal using a 1x1 km grid. (b). Frequency distribution of slope angles derived from 1x1 km grid are shown (a). Slopes in general are steeper in the summit regions than in the valleys. Note that instead of the average slope of a grid cell only steepest vertical gradients are plotted.

## 2. Marked-up manuscript showing the changes made with respect to the last iteration of review

(See following pages)



# 1 **A conceptual, distributed snow redistribution model**

2 **S. Frey<sup>1</sup> and H. Holzmann<sup>1</sup>**

3 [1]{Institute of Water Management, Hydrology and Hydraulic Engineering, University of  
4 Natural Resources and Life Sciences, Vienna, Austria}

5 Correspondence to: S. Frey (simon.frey@boku.ac.at)

6

## 7 **Abstract**

8 When applying conceptual hydrological models using a temperature index approach for  
9 snowmelt to high alpine areas often accumulation of snow during several years can be  
10 observed. Some of the reasons why these “snow towers” do not exist in nature are vertical and  
11 lateral transport processes. While snow transport models have been developed using grid cell  
12 sizes of tens to hundreds of square meters and have been applied in several catchments, no  
13 model exists using coarser cell sizes of one km<sup>2</sup>, which is a common resolution for mean and  
14 large scale hydrologic modelling. In this paper we present an approach that uses only gravity  
15 and snow density as a proxy for the age of the snow cover and land-use information to  
16 redistribute snow in Alpine basins. The results are based on the hydrological modelling of the  
17 Austrian Inn basin in Tyrol, the detailed description of the current paper refer to the  
18 catchment of Ötztaler Ache, Austria, but the findings hold for other tributaries of the river  
19 Inn. This transport model is implemented in the distributed rainfall-runoff model COSERO. A  
20 comparison for model validation between the standard model without parameterization for  
21 lateral snow redistribution and the updated version is done using observed discharge and  
22 MODIS derived snow covered areas. While the signal of snow redistribution can hardly be  
23 seen in the binary classification compared with MODIS, snow accumulation over several  
24 years can be prevented. In a seven year period the standard model would lead to snow  
25 accumulation of approximately 2900 mm SWE in high elevated regions whereas the updated  
26 version of the model does not show accumulation and does also predict discharge more  
27 precisely leading to a Kling-Gupta-Efficiency of 0.93 instead of 0.9. A further improvement  
28 can be shown in the comparison of MODIS snow cover data and the calculated depletion  
29 curve, where the redistribution model increased the efficiency ( $R^2$ ) from 0,70 to 0,78  
30 (calibration) and from 0,66 to 0,74 (validation).

1

## 2 **1 Introduction**

3 Conceptual models are widely used in hydrology. Examples are the HBV model (Bergström,  
4 1976), PDM (Moore, 2007), GSM-SOCONT (Schaepli et al., 2005) or VIC (Wood et al.,  
5 1992) just to name a few. Many of these conceptual models use a temperature index approach  
6 to model snow melt and snow accumulation and even in some physically based models as  
7 e. g. versions of the SHE model (Bøggild et al., 1999) this method can be found. This  
8 approach has the advantage of being quite simple since it uses only temperature as input to  
9 determine whether precipitation occurs in the form of snow or rain and whether snow can be  
10 melted or not. A typical example of a temperature index method for snow modelling is the  
11 degree-day approach (see for example Hock 2003). A disadvantage is that snow accumulates  
12 as long as the air temperature does not rise above a certain threshold (often 0 °C) regardless of  
13 any other processes that may lead to snow melt like radiation or turbulent fluxes of latent  
14 energy. In high mountainous areas this may be the case for most days in the year leading to an  
15 intensive computational accumulation of snow in these areas. In the modellers terminology  
16 these artefacts are often called “snow towers”. In nature, however, these accumulations are  
17 barely existent.

18 The reasons for that are either wind or gravitationally induced lateral snow distribution  
19 processes (Elder et al., 1991; Winstral et al., 2002). Resulting snow depths are not uniformly  
20 distributed in space but vary within large ranges (Helfricht et al., 2014). When changing the  
21 focus from micro (e. g. several square meters) to macro scales (e. g. one to several square  
22 kilometres), variations become less (Melvold and Skaugen, 2013). The intention of the  
23 applied snow redistribution concept was (a) to prevent the artefacts of “snow towers” and (2)  
24 to develop a concept which considers gravity driven lateral snow transport with reasonable  
25 and plausible process depiction.

### 26 **1.1 Theoretical background of snow cover variations**

27 During the accumulation period, according to Liston (2004), primarily three mechanisms are  
28 responsible for these variations: (i) snow-canopy interactions in forest covered regions, (ii)  
29 wind induced snow redistribution and (iii) orographic influences on snow fall. These  
30 mechanisms influence snow cover patterns on scales ranging from the micro to the macro  
31 scale. Spatial snow cover variability beneath canopies is mainly affected by different tree

1 species (deciduous vs coniferous trees) influencing LAI, height and density of the canopy and  
2 gap sizes (Garvelmann et al., 2013; Liston, 2004; Pomeroy et al., 2002).

3 Besides the impact of vegetation, wind is the most dominant factor influencing snow patterns  
4 in alpine terrain. Snow is transported from exposed ridges to the lee side of these ridges,  
5 valleys and vegetation covered areas (Essery et al., 1999; Liston and Sturm, 1998; Rutter et  
6 al., 2009; Winstral et al., 2002). One has to be aware that besides of the physical transport of  
7 solid snow wind also stimulates sublimation processes (Liston and Sturm, 1998; Strasser et  
8 al., 2008). Wind influences snow depth distributions on scales of some 100s to 1000 square  
9 metres (Dadic et al., 2010a).

10 The third mechanism (orographic effect) influences snow patterns on a larger scale of one to  
11 several kilometres (e. g. Barros and Lettenmaier, 1994). Non-uniform snow distributions are  
12 caused by interactions of the atmosphere (air pressure, humidity, atmospheric stability) with  
13 topography (Liston, 2004).

14 In addition to these processes, avalanches play a role in snow redistribution (Lehning and  
15 Fierz, 2008; Lehning et al., 2002; Sovilla et al., 2010). In steep terrain, avalanches depend  
16 mainly on the slope angle and are capable of transporting large snow masses over distances of  
17 tens to hundreds of metres (Dadic et al., 2010b; Sovilla et al., 2010).

18 During the ablation period, spatial snow distributions are mainly influenced by differences in  
19 snow melt behaviour. On the northern hemisphere, on south-facing slopes, rates of snow melt  
20 are generally enhanced compared to north-facing slopes due to the inclination of radiation.  
21 Also vegetation influences melting behaviour. Shading reduces snowmelt compared to direct  
22 sunlight. Enhanced emitted long wave radiation due to warm bare rocks or trees increases the  
23 melt rate (Garvelmann et al., 2013; Pohl et al., 2014).

## 24 **1.2 Modelling approaches**

25 A common approach avoiding intensive accumulation of snow is editing the meteorological  
26 input (Dettinger et al., 2004). For instance, many models use a constant yet adjustable lapse  
27 rate for interpolating temperature with elevation (Holzmann et al., 2010; Koboltschnig et al.,  
28 2008). Besides temperature, precipitation gradients are often adjusted to fit observed and  
29 modelled target variables (e. g. snow patterns or runoff) (Huss et al., 2009b; Schöber et al.,  
30 2014). Justification for doing so is the general lack of gauging stations in the summit regions  
31 (Daly et al., 1994, 2008) along with the high error of precipitation gauges (Rasmussen et al.,

1 2011; Williams et al., 1998). An approach presented by Jackson (1994) defining a  
2 precipitation correction matrix was successfully applied in several studies (Farinotti et al.,  
3 2010; Huss et al., 2009a). Scipión et al. (2013) however identified significant discrepancies  
4 between precipitation patterns obtained by a Doppler X-band radar and the snow  
5 accumulation at the end of the winter period which gives clear indications that snowfall is  
6 redistributed based on different driving forces. Consequently, the variability of the  
7 meteorological input cannot explain the variability of snow cover patterns.

8 Models trying to deal with snow accumulation and redistribution apart from input corrections  
9 may be classified into two major approaches. One is the consideration of process based snow  
10 distribution patterns the other approach is empirical. Examples for process oriented model are  
11 SNOWPACK (Bartelt and Lehning, 2002) used in avalanche research or SnowTran3D  
12 (Liston et al., 2007; Liston and Sturm, 1998). Empirical models use the fact, that snow  
13 patterns resemble each other every year (Helfricht et al., 2012, 2014). The presented paper  
14 concentrates on the empirical approach.

15 Helfricht et al. (2012) used airborne LiDAR measurements to determine snow accumulation  
16 gradients for elevation bands in the Ötztaler Alps. These could be used to improve  
17 hydrological models regarding snow cover distributions and subsequently to achieve better  
18 runoff predictions. LiDAR data, however, are relatively expensive. Often wind speed  
19 and -direction are used to model snow drift (e.g. Bernhardt et al., 2009; 2010; Shulski and  
20 Seeley, 2004; Winstral et al., 2002; Liston and Sturm, 1998). Kirchner et al. (2014) concluded  
21 from LiDAR measurements in combination with meteorological stations in a catchment in  
22 California, USA that wind measurements from only one meteorological station are of too poor  
23 quality for a useful description of wind fields for snow transport. The computed wind fields  
24 generated by regional circulation models (RCM) have also shown to be erroneous (Nikulin et  
25 al., 2011) and therefore are not useful for direct implementation in redistribution models.  
26 Additionally models using wind have in common that they are computationally intensive as  
27 they require data in high spatial resolution (e. g. 100 to 1000s of square metres). Schöber et al.  
28 (2014) combined gravitational and wind induced snow transport using a distributed energy  
29 balance model with a resolution of 50x50 m.

30 However, the difficulties of snow accumulation also occur when models with coarser cell  
31 sizes are applied. Due to some available databases for vegetation and meteorology (Haiden et  
32 al., 2011; Masson et al., 2003; Oubeidillah et al., 2014), many models operate on cell sizes of

1 1 km<sup>2</sup> or more (e. g. Andersen et al., 2001; Henriksen et al., 2003; Mauser and Bach, 2009;  
2 Safeeq et al., 2014). To our knowledge, no model for redistributing snow on a 1x1 km grid  
3 size exists. In this paper we present a simple approach to deal with snow in high mountainous  
4 regions and its application in the catchment of Ötztaler Ache in Tyrol, Austria. Since the  
5 model uses meteorological input from INCA (Haiden et al., 2011) that already account for  
6 meteorological corrections, we focus on snow redistribution rather than to edit the input data.  
7 As already mentioned the two main objectives in this respect are to achieve a better model  
8 efficiency regarding runoff and to avoid the existence of snow towers at high altitudes.

9

## 10 **2 Model description**

### 11 **2.1 Hydrological Model COSERO**

12 COSERO is a spatially distributed conceptual hydrological model which is similar to the  
13 HBV model (Bergström, 1976). In the presented paper it uses 1x1 km grid cells. Originally  
14 developed for modelling discharge of the Austrian rivers Enns and Steyer (Nachtnebel et al.,  
15 1993), it has recently been used for different purposes like climate change studies (e. g. Kling  
16 et al., 2012, 2014b; Stanzel and Nachtnebel, 2010), investigating the role of  
17 evapotranspiration in high alpine regions (Herrnegger et al., 2012) and operational runoff  
18 forecasting (Stanzel et al., 2008). Potential evapotranspiration is calculated using the  
19 Thornthwaite method (Thornthwaite, 1948). Discharge due to rainfall and snow-/ice melt is  
20 estimated using the same non-linear function of soil moisture as the original HBV. In this  
21 study, the model is run using daily time steps. It is, however, capable of using hourly or  
22 monthly time steps. In the latter case, intra-monthly variations are considered for snow and  
23 interception processes as well as for soil moisture (Kling et al., 2014a). A schematic overview  
24 of the model is given by Fig. 1 and a detailed description of the model can be found in Kling  
25 et al. (2014a), where the model was applied to several catchments across Europe, Africa and  
26 Australia. However, in Kling et al. (2014a) snow parameters were not calibrated and therefore  
27 the snow module is not fully explained in detail in their paper. This will be done in the  
28 following. Equations (1) to (7) and (10) were taken from the original model by Stanzel and  
29 Nachtnebel (2010), all other methods were developed in the present study.

30 Numerous studies have shown that sub-grid variability of snow depths can be described by a  
31 two parameter log-normal distribution (e. g. Donald et al., 1995; Pomeroy et al., 1998).

1 COSERO uses five snow classes per cell (i.e. the log-normal distribution is subdivided into  
2 five quantiles) to approximate this sub-grid log-normal distribution under accumulation  
3 conditions (see Fig. 2 b)), i. e. snowfall is distributed log-normally into snow classes, where  
4 the sum of the snow water equivalent (SWE) of each classes represent the mean conditions in  
5 the grid cell. This distribution can be interpreted as a statistical description of snow  
6 distribution processes taking place at the subgrid scale (Pomeroy et al., 1998). This method  
7 has the potential to indirectly consider the influence of curvature, shelter, vegetation or  
8 elevation (Hiemstra et al., 2006). The properties of each class are treated unique as equations  
9 (1) to (13) apply to every snow class separately. Consequently the log-normal distribution  
10 within a grid cell may be disturbed by the processes of melting, sublimation, refreezing and  
11 redistribution to other grid cells. Once fallen, snow redistribution between the snow classes  
12 within a single grid cell is not considered. A scheme of the composition of a snow class is  
13 illustrated in Fig. 2 a). The snow water equivalent ( $S_{SWEt}$ ) of a given day  $t$  per class is  
14 calculated by Eq. (1) where  $P_{Rt}$  and  $P_{St}$  are liquid and solid precipitation in mm, respectively,  
15  $M_t$  is snow melt and  $E_{St}$  is sublimation of snow. All variables are given in mm SWE.

$$16 \quad S_{SWEt} = S_{SWE_{t-1}} + P_{Rt} + P_{St} - M_t - E_{St} \quad (1)$$

17 Snow melt is calculated by a temperature index approach (see for example Hock 2003). Eq.  
18 (2) is used:

$$19 \quad M_t = \min(S_{SWEt}; P_{Rt} \cdot \varepsilon \cdot T_{AIRt} + D_{ft} \cdot T_{AIRt}) \quad (2)$$

20 where  $M_t$  is snowmelt [mm],  $\varepsilon$  is the ratio of specific heat of water and melting energy,  $T_{AIRt}$   
21 is the (mean) daily air temperature [ $^{\circ}\text{C}$ ] and  $D_{ft}$  [ $\text{mm } ^{\circ}\text{C}^{-1}$ ] is the snow melt factor of a given  
22 day  $t$  estimated by Eq. (3):

$$23 \quad D_{ft} = \left( -\cos\left(J \cdot \frac{2\pi}{365}\right) \cdot \frac{D_U - D_L}{2} + \frac{D_U - D_L}{2} \right) \cdot M_{REDt} \quad (3)$$

24 with

$$25 \quad M_{REDt} = \begin{cases} D_{RED}, & S_{fresh} \geq S_{CRIT} \\ M_{RED_{t-1}} + \frac{(1 - M_{RED_{t-1}})}{5}, & S_{fresh} < S_{CRIT} \end{cases} \quad (4)$$

26 where  $J$  is the Julian day of the year [-],  $D_U$  and  $D_L$  are the upper and lower boundaries of  $D_f$   
27 [ $\text{mm } ^{\circ}\text{C}^{-1}$ ], respectively, and  $M_{RED}$  [-] is a reduction factor to account for the higher albedo  
28 caused by freshly fallen snow calculated by Eq. (4).  $S_{CRIT}$  [mm] is the critical snow depth of  
29 fresh snow necessary to increase the albedo, whereas  $S_{fresh}$  is the actual depth of fresh snow



1 [mm] fallen within one time step. For fresh snow depth larger than  $S_{CRIT}$ ,  $M_{RED}$  is set to a  
2 reduced melting factor  $D_{RED}$  [-].

3 Whether precipitation occurs in form of snow or rain is controlled by two parameters  $T_{PS}$  and  
4  $T_{PR}$ , defining the temperature range where snow and rain occur simultaneously. At and above  
5 temperature  $T_{RP}$  precipitation is pure liquid, at and below  $T_{PS}$  precipitation is pure solid. In  
6 between those two boundaries, the proportion of solid to liquid precipitation is estimated  
7 linearly.

8 For the estimation of snow sublimation, Eq. (5) is used, where  $E_{SP}$  [mm] refers to potential  
9 sublimation of snow,  $E_P$  [mm] is the potential evapotranspiration and  $E_R$  is a correction factor  
10 to reduce  $E_P$ . Sublimation is considered only for snow classes actually covered by snow.  
11 Hence, if a grid cell is partly snow free (this can be the case if one subgrid class has no snow  
12 cover due to melting) sublimation is estimated for the snow covered part only. For the  
13 uncovered classes evapotranspiration according to the Thornthwaite method is applied.

$$14 \quad E_{SP_t} = E_{P_t} \cdot E_R \quad (5)$$

15 The snow cover in COSERO is treated as porous medium and therefore is able to store a  
16 certain amount of liquid water ( $S_l$ ) in dependency of the snow pack density ( $\rho$ ) calculated  
17 using Eq. (6).

$$18 \quad S_{l_t} = (S_{SWE_t} - S_{l_{t-1}}) \cdot (S_{l_{MAX}} - (\rho - \rho_{MAX}) \cdot S_{l_\rho}) \quad (6)$$

19 Where  $S_{l_{MAX}}$  [ $m^3 \text{ kg}^{-1}$ ] is the maximum water holding capacity at the maximum snow density  
20 of the snow pack  $\rho_{MAX}$  [ $\text{kg m}^{-3}$ ] and  $S_{l_\rho}$  describes the decrease of water holding capacity with  
21 increasing snow density  $\rho$ .

22 At negative air temperatures, retained melt water has the ability to refreeze in the snow pack.  
23 The potential amount of refrozen water ( $S_R$ ) is estimated by Eq. (7), where  $R_f$  is the refreezing  
24 factor [ $\text{mm } ^\circ\text{C}^{-1}$ ]. As long as there is enough liquid water in the snow pack, actual refreezing  
25 will be equal to potential refreezing.

$$26 \quad S_R = \begin{cases} 0, & T_{AIR_t} > 0 \\ R_f \cdot (T_{AIR_t} \cdot (-1)), & T_{AIR_t} \leq 0 \end{cases} \quad (7)$$

27 Refrozen water is treated in the same way as snow. The amount of water leaving the snow  
28 cover then equals snowmelt minus retained water.

1 Snow density ( $\rho_t$ ) of each class is calculated using a sigmoid function shown in Eqs. (8) and  
 2 (9) where  $\rho_{MAX}$  and  $\rho_{MIN}$  are the respective maximum and minimum values of  $\rho$ ,  $T_{AIR}$  is the  
 3 temperature of the air mass above the snow layer and  $\rho_{scale}$  and  $T_{scale}$  are scaling coefficients  
 4 to calculate a transition temperature ( $T_{tr}$ ) for the estimation of the snow density. Herby,  $\rho_{scale}$   
 5 adjusts the slope of the function, whereas  $T_{scale}$  is responsible for a shift on the x-axis. These  
 6 two parameters are set to fixed values of 1.2 and 1, respectively. The solution of Eqs. (8)  
 7 and (9) is illustrated in Fig. 3 for a range of typical air temperatures, where snowfall occurs.  
 8 Already fallen snow can reach a higher density ( $\rho_{OLD}$ ) than fresh snow. Its density is  
 9 calculated using a time settling constant ( $\rho_{SET}$ , derived from Riley et al., 1973) until the  
 10 maximum density is reached (Eq. 10).

$$11 \quad \rho_t = (\rho_{MAX} - \rho_{MIN}) \cdot \left( \frac{T_{tr}}{\sqrt{1+(T_{tr})^2}} + 1 \right) \cdot 0.5 + \rho_{MIN} \quad (8)$$

12 with

$$13 \quad T_{tr} = \frac{T_{AIR_t}}{\rho_{scale}} + T_{scale} \quad (9)$$

$$14 \quad \rho_{OLD} = \frac{\rho_{SET} \cdot \left( \frac{S_{SWE_t} + S_t}{2} \right)}{1 + \frac{\rho_{SET}}{2}} \quad (10)$$

15 The COSERO model considers both snow and glacier ice melt processes. Ice melt ( $M_{ICE}$ ) is  
 16 computed by means of a degree-day method (see Eq. 11) and uses separate parameter sets.  
 17 Here,  $D_{ICE}$  refers to the ice melt factor [ $\text{mm } ^\circ\text{C}^{-1}$ ]. A prerequisite of ice melt is the full  
 18 depletion of the overlying snow cover. Spatial information of glaciers are taken from the  
 19 Randolph Glacier Inventory version 3.2 (Arendt et al., 2012).

$$20 \quad M_{ICE} = D_{ICE} \cdot T_{AIR} \quad (11)$$

## 21 **2.2 Snow transport model**

22 Several authors reported that the slope angle has an important influence on snow depths  
 23 (Bernhardt and Schulz, 2010; Kirchner et al., 2014; Schöber et al., 2014). The model  
 24 redistributes snow only to grid cells providing the steepest slope (acceptor cell) in the direct  
 25 neighbourhood of the raster cell it searches from (donor cell). Only downward transportation  
 26 is considered. If more than one cell show the same (largest) difference in elevation, the  
 27 amount of donated snow is distributed equally to the number of acceptor cells. The actual  
 28 amount of snow being redistributed depends on the steepness of the slope, the age of the snow

1 cover, considered by the density of snow, the type of land cover of the donor cell and the  
 2 snow depth of the donor cell. The drier (less dense) the snow pack the higher the snow rate  
 3 available for the redistribution routine ( $f_\rho$ , Eq. 13). Thus the defined maximum density of  
 4 snow ( $450 \text{ kg m}^{-3}$ ) determines the threshold for snow redistribution. The availability of snow  
 5 for transport is determined by a vegetation-based threshold value ( $H_v$ ) for each class of land  
 6 cover. This value can also be interpreted as a roughness coefficient for areas where no or  
 7 hardly any vegetation is present like in alpine and nival elevations. If the snow depth ( $S$   
 8 [mm]) of a snow class of a raster cell exceeds  $H_v$  [mm], snow transport from that cell is  
 9 activated and redistribution is calculated by solving Eqs. (12) and (13).

$$10 \quad = \max(S_D - H_v; 0) \cdot f_\rho \cdot \frac{1}{\Sigma A} \cdot C \quad (12)$$

11 With

$$12 \quad f_\rho = \left( \frac{(\rho_{MAX} - \rho_D)}{\rho_{MAX}} \cdot e^{\left( -\frac{\rho_D}{\rho_{MAX}} \right)} \right) \cdot \frac{\alpha}{90} \quad (13)$$

13 Where  $S_{SWE(A)}$  is the amount of snow water equivalent that is redistributed from the donor cell  
 14 (D) to the available acceptor cell(s) (A),  $\rho_D$  is the density of snow in the donor cell,  $\rho_{MAX}$  is  
 15 the possible maximum density of snow,  $\alpha$  is the angle of the slope between the donor and  
 16 acceptor cells in degree and  $C$  is a correction coefficient that can be calibrated.

17 Notwithstanding, that other geomorphological properties than slope angle influencing snow  
 18 patterns are important on scales smaller than the grid size of COSERO (see section 1.1), slope  
 19 was selected as driving force for the model. One has to be aware that this is a simplification  
 20 and under realistic conditions snow might not necessarily be transported only on the steepest  
 21 route (Bernhardt and Schulz, 2010; Winstral et al., 2002).

22 Fig. 4 illustrates the shape of the distribution coefficient  $f_\rho$  as a function of different elevation  
 23 gradients between the acceptor and donor cells and of the snow density. In acceptor cells  
 24 redistributed snow is treated as fresh snow in the sense that it is distributed to the snow  
 25 classes according to the log-normal distribution.

26 The model is organized in form of a loop starting at the highest grid cell (summit region) and  
 27 ending at the lowest cell (outlet of the catchment). This ensures that snow cannot be  
 28 redistributed into already processed grid cells. Snow will be transported downslope as long as  
 29 the slope is big enough to allow for transportation given that the density of snow is low  
 30 enough. Consequently, snow accumulates rather in flat regions of the catchment. The concept

1 of the redistribution model is sketched in Fig. 5. Note that although snow depths in the highest  
2 cell are prevented by the model, the number of snow covered cells remains the same.

3

### 4 **3 Case study in the catchment the Ötztaler Ache, Tyrol, Austria**

#### 5 **3.1 Catchment description**

6 The catchment of Ötztaler Ache at gauge Huben, situated in western Austria close to the  
7 Italian border, covers an area of 511 km<sup>2</sup> and has an altitudinal range between 1185 m a.s.l at  
8 the gauge at Huben and 3770 m a.s.l at its highest peaks. Due to the use of a 1x1 km gridded  
9 DEM, the highest grid cell has a mean elevation of 3450 m a.s.l, whereas the lowest cell has  
10 an elevation of 1250 m a.s.l. (Fig. 6). About 30 % of its area is covered by vegetation, mainly  
11 pastures and meadows. Glaciers cover about 19 % leading to an annual ice melt contribution  
12 of about 25 % of the total runoff at Huben, while 41 % of the discharge has its origin in  
13 snowmelt (Weber et al., 2010). Table 1 gives an overview of the land cover.

14 In Fig. 6 the elevations of the Ötztal basin are described. Frequency distribution of slope  
15 angles derived from 1x1 km grid are shown (6 a). This frequency distribution exhibits the  
16 highest frequencies in the slope classes between 20 and 25 degrees for higher elevations. In  
17 lower elevated regions slope classes between 0 and 15 degrees dominate. However, also  
18 glacier covered areas at the summits can have flat slopes. Note that the listed slopes are based  
19 on the steepest vertical gradients of the neighbour elements.

#### 20 **3.2 Input data**

21 Gridded meteorological data of precipitation and air temperature are required to run the  
22 model. These data are provided by the INCA dataset (Haiden et al., 2011) with the same grid  
23 spacing like the hydrological model, allowing a direct use in the model without the need for  
24 pre-processing. INCA data are available since 2003. The years 2003 and 2004 have been used  
25 as a warm-up period for the model. In the subsequent years no correction of meteorological  
26 data was done since INCA already accounts for elevation gradients regarding air temperature  
27 and precipitation. Six land use classes were derived from the most recent CORINE data set  
28 (CLC2006 version 17, see EEA, 1995). These classes and their areal fractions in the  
29 catchment of Ötztaler Ache are given in Table 1. It should be pointed out, that neither  
30 radiation nor wind speed or wind direction data are necessary to run the model.

### 1 **3.3 Model calibration**

2 The hydrological model was calibrated for the period from 2005 to 2008 using a  
3 Rosenbrock's automated optimization routine (Rosenbrock, 1960). Although the model is rich  
4 of parameters, the vast majority of them have been estimated *a priori* according to literature  
5 (Liston and Sturm, 1998; Prasad et al., 2001) and previous work on the model (Fuchs, 2005;  
6 Kling, 2006; Nachtnebel et al., 2009). In the snow model including snow redistribution only  
7 six parameters have been calibrated: upper and lower boundaries of snow melt factors  $D_U$  and  
8  $D_L$ , respectively, the threshold values that control the range where liquid and solid  
9 precipitation occur simultaneously ( $T_{PR}$ ,  $T_{PS}$ ), the standard deviation of the log-normal  
10 distribution of snow depth in one grid cell ( $N_{VAR}$ ) and the calibration parameter for snow  
11 redistribution  $C$  (see Eq. 12). The limited number of optimization parameters reduces  
12 equifinality problems. For a more detailed description of equifinality issues see the  
13 supplements of this article. The target of the calibration was a good fit of runoff using the  
14 Kling-Gupta-Model-Efficiency (Gupta et al., 2009; Kling et al., 2012) as objective function.  
15 The model was validated for the years 2009 and 2010. Both calibration and validation have  
16 been done with and without using the snow drift module. In the following model A refers to  
17 the model using snow transport, whereas model B stands for the standard model. Vegetation  
18 threshold values for snow detention were taken from previous studies (Liston and Sturm,  
19 1998; Prasad et al., 2001). These are given in Table 1. Maximum snow density was assumed  
20  $450 \text{ kg m}^{-3}$  which matches long term snow measurements (Jonas et al., 2009; Schöber et al.,  
21 2014). Besides discharge in the validation period also snow cover data from MODIS (8 day  
22 maximum snow cover, version 5) satellite images (Hall et al., 2002) were used to compare the  
23 performance of both models.

24

## 25 **4 Results**

### 26 **4.1 Discharge**

27 Fig. 7 shows a comparison of total discharge using model A and B at the gauge Huben for the  
28 year 2006. Both models result in similar quality criteria in the calibration as well as in the  
29 validation period (see Table 2). Nevertheless, the model efficiency could be improved by 0.05  
30 in the calibration period and 0.02 in the validation period by accounting for lateral snow  
31 transport. Maximum differences in the mean daily discharges between the two models reach

1 up to 2 mm per day (which equals to  $12.1 \text{ m}^3 \text{ s}^{-1}$ ) leading to a relative difference of minus 9 up  
2 to 44 % of model A in respect to model B. In total, model A generates a surplus of about  
3 300 mm discharge in five years compared to model B (Fig. 8). About 2/3 of the additional  
4 discharge originate in enhanced snowmelt the rest occurs due to enhanced glacier melt.

## 5 **4.2 Spatially distributed snow cover data**

6 Fig. 9 compares model A and B with MODIS snow depletion data. Both the accumulation  
7 period in winter and the ablation period in spring and summer are represented well by both  
8 models. Cold snowfall periods in summer generate sharp peaks in the depletion curve, which  
9 could be calculated by both model versions, where Model A computed slightly smaller peaks  
10 during the snowmelt period (May to July). This leads to a moderate increase of the  
11 determination factor  $R^2$  from 0.70 to 0.78 (calibration) and from 0.66 to 0.74 (validation).

## 12 **4.3 Inter annual snow accumulation**

13 The main reason for developing a snow transport model was the prevention of “snow towers”  
14 – accumulation of snow over several years in high mountainous regions. Fig. 10 presents  
15 model behaviour of model A and B with respect to the accumulation of snow in elevations  
16 above 2800 m a.s.l. This elevation was chosen because here none of the models indicates  
17 snow accumulation for more than one year and therefore snow accumulation in lower  
18 altitudes is no problem. By the end of seven years of modelling, model B shows snow depths  
19 of approx. 2900 mm SWE in elevations above 3400 m a.s.l. whereas model A does hardly  
20 show any accumulation behaviour in these altitudes. Spatially distributed net loss and gain of  
21 snow for all raster cells within the period of one year in the watershed are presented in  
22 Fig. 11. It can be shown that net loss is evident in the zones of ridges and high elevations,  
23 where the maximum net gain is along the valley bottoms.

## 24 **4.4 Parameter equifinality**

25 Since the model uses several parameters that need calibration it suffers from equifinality  
26 issues. To investigate those issues, Monte Carlo simulations have been carried out varying the  
27 snow relevant parameters that cannot be estimated a priori. Since the aim of this paper is  
28 snow transport, the results of the Monte Carlo simulations can be found in the supplements of  
29 this article.



1

## 2 **5 Discussion**

### 3 **5.1 Discharge**

4 In spring, at the beginning of the melting season, higher runoff is generated by model A due  
5 to a larger amount of snow in lower altitudes (see Fig. 7). Later in the year enhanced glacier  
6 melt is mainly responsible for higher discharge rates. About 200 mm have their origin in  
7 enhanced snowmelt, while the remaining 100 mm originate in amplified melt of glaciers.  
8 Since glacier cover about 19.4 % of the catchment's area 100 mm of additional mean basin  
9 runoff corresponds to an enhanced negative glacier mass balance of -500 mm. The reason for  
10 this is transport of snow in warmer altitudes and therefore earlier and more snow free glacier  
11 surfaces producing higher discharge due to glacier melt (see Fig. 8) and explains the peak in  
12 July and August in runoff difference (see Fig 7).

### 13 **5.2 Spatially distributed snow cover data**

14 Fig. 9 shows the snow depletion curve of the year 2009 based on MODIS data and the  
15 comparison of model runs A and B. Only little differences between model A and B can be  
16 identified. The reason for this is the vegetation threshold. Even if snow is being transported, a  
17 residual of snow remains in the donor cell resulting in the cell marked as snow covered. Grid  
18 cells covering the summits only donate snow to their respective acceptor cells. However, a  
19 certain amount of snow is held back according to the threshold due to vegetation and  
20 roughness of the surface. As indicated in Fig. 5 grid cells nested in the intermediate slope  
21 regions receive and donate snow at the same time. Thus their snow depth changes little if  
22 comparing model A and model B. In flat valley regions, grid cells only receive snow, where  
23 relatively high air temperature values often allow for melting.

24 Satellite based snow cover information by MODIS are binary and so is the model output for  
25 comparing these results. In a binary system, no difference can be distinguished between cells  
26 covered by much or little snow.

### 27 **5.3 Snow accumulation**

28 While using model B, the higher the elevation the more snow is accumulated. Contrary,  
29 model A shows less pronounced and in some high altitudes even contrary behavior(see Fig.

1 10). This is a result of the slope dependency of the distribution model that the amount of snow  
2 distributed to other grid cells is higher with increasing vertical distance to the downward grid  
3 cell. In general and in the Ötztal as well mountains are steeper in the summit regions than at  
4 the bottom (see Fig. 6). Consequently in the summit regions snow will be preferentially  
5 eroded while it accumulates at the rather flat valleys where the vertical distances between the  
6 grid cells are less than at the peaks. This does reflect snow accumulations that can be  
7 observed in nature where summits might be nearly snow free in spring while flatter parts are  
8 still covered with snow. While the raster cells covering peak regions act as donators only  
9 those cells located on slopes may receive and distribute snow at the same time (Fig. 11).  
10 Valley regions only receive snow. The resulting net loss and gain areas shown in Fig. 11 give  
11 some indication that the redistribution algorithm is plausible.

12 Although snow accumulation behaviour of model A is more realistic than model B snow  
13 accumulation can still be observed in the highest elevations zone (see Fig. 10). This is based  
14 on the parameterization of the snow holding capacity  $H_v$ , where even bare ground assigns a  
15 value of 200 mm (see Table 1). The influence of the highest elevation class ( $> 3400$  m a.s.l.)  
16 on both the hydrograph and snow covered area however is very small, since this elevation  
17 level is represented by only four grid cells. Consequently the objective function during  
18 calibration using an automated optimization routine like Rosenbrock's routine does not differ  
19 much when underestimating the correction coefficient in these grid cells.

20 The smaller the portion of high altitude areas in a catchment compared to the total catchment  
21 area the less important is snow redistribution for modelling runoff. This ratio of summit  
22 regions to total catchment size is normally smaller for bigger catchments. The catchment of  
23 river Inn, for instance, covers an area of about 10000 km<sup>2</sup> yet only 733 km<sup>2</sup> are located at  
24 elevations where intensive snow accumulations and mobilizations occur (above  
25 2800 m a.s.l.). In the Ötztal basin 204 out of 511 km<sup>2</sup> are located higher than 2800 m a.s.l. If  
26 model A is applied to the catchment of river Inn in five years of modelling about 15 mm SWE  
27 (with respect to the entire river basin) remain in the catchment due to snow accumulation  
28 processes instead of 300 mm in the Ötztal.

#### 29 **5.4 Transferability to other catchments**

30 The model provides results that have been found by other models, too. For instance the  
31 elevation where the highest snow accumulations occurs (2800 to 3000 m a.s.) as was found by

1 LiDAR measurements in the same catchment (Helfricht et al., 2012) as well as by modelling  
2 (Frey, 2015). Given that and the needs of the model (slope angles, snow density) for  
3 transporting snow, it produces valid results as long as a catchment features relatively steep  
4 slopes in the summit regions (which is the case in most catchments in the Alps). Obviously,  
5 the model needs calibration if it is transferred to another catchment.

#### 6 **5.4 Parameter equifinality**

7 Like most hydrological models COSERO requires calibration of some parameters. This  
8 necessarily causes equifinality issues (Beven and Freer, 2001). The more adjustable  
9 parameters a model provides, the more important this problem may become (e. g. Gupta et al.,  
10 2008). On the other hand, some authors pointed out that more complex models may produce  
11 more feasible results if the parameters can be estimated within realistic boundaries (Gharari et  
12 al., 2012, 2014; Hrachowitz et al., 2014). Applying COSERO with the presented snow  
13 redistribution routine requires two additional parameters: the vegetation threshold  $H_V$   
14 (estimated *a priori*) and the calibration parameter  $C$  (see Eq. 12). Yet, accounting for snow  
15 redistribution allows the modeller to use  $D_U$  values within or close to the range proposed by  
16 Kling et al., (2006), while the standard version of the model leads to the best results if higher  
17 and therefore unrealistic  $D_U$  values are used (see supplements of this article).

18

### 19 **6 Conclusions**

20 A model for redistribution of snow on a coarse 1x1 km raster has been developed and tested  
21 in the catchment of Ötztaler Ache, Austria. While only little improvement of snow cover  
22 compared to MODIS data could be achieved, appearance of “snow towers” in high altitudes  
23 could be prevented. In terms of discharge at the outlet of the basin, both models show good  
24 results. However, the Kling-Gupta-efficiency of model A could be improved by 0.05 in the  
25 calibration and by 0.02 in the validation period. With respect to the entire watershed area the  
26 model using snow redistribution generates about 200 mm more runoff originated from  
27 snowmelt in five years than without considering this process. This does not only affect the  
28 water balance of the catchment but also amplifies glacier melt about 500 mm in five years,  
29 with respect to glaciated areas, due to longer time periods where glacier surfaces are fully  
30 snow free.

1 The integration of a snow transport module promotes the demand, that models work “right for  
2 the right reasons” and is an attempt to integrate more real process understanding into the  
3 model approach. Further work needs to be carried out with respect to validation of spatially  
4 distributed snow patterns. For this purpose, satellite images from Landsat might be of use  
5 providing a higher spatial resolution than MODIS.

6 Even though the vast majority of parameters were estimated *a priori* in this work, equifinality  
7 remains an issue. However, redistribution of snow requires only two additional parameters but  
8 allows for more realistic boundaries (see Kling et al., 2006) of the snow melt factors (see  
9 supplements of this article). However, more work needs to be carried out to account for that  
10 issue.

## 11 **Acknowledgements**

12 The authors thank their colleagues for continuing support and discussion around the coffee  
13 breaks, especially to Matthias Bernhardt for friendly reviewing and commenting on the  
14 manuscript. Special thanks to Herbert Formayer and David Leidinger of the institute of  
15 meteorology, BOKU, for supplying the INCA data. Many thanks to Daphné Freudiger and  
16 two further anonymous referees for their comments and valuable suggestions. This study was  
17 part of a research project in cooperation with Verbund AG.

18

## 19 **References**

20 Andersen, J., Refsgaard, J. C. and Jensen, K. H.: Distributed hydrological modelling of the  
21 Senegal River Basin — model construction and validation, *J. Hydrol.*, 247(3-4), 200–214,  
22 doi:10.1016/S0022-1694(01)00384-5, 2001.

23 Arendt, A., Bolch, T., Cogley, J. G., Gardner, A., Hagen, J.-O., Hock, R., Kaser, G., Pfeffer,  
24 W. T., Moholdt, G., Paul, F., Radić, V., Andreassen, L., Bajracharya, S., Barrand, N., Beedle,  
25 M., Berthier, E., Bhambri, R., Bliss, A., Brown, I., Burgess, D., Burgess, E., Cawkwell, F.,  
26 Chinn, T., Copland, L., Davies, B., De Angelis, H., Dolgova, E., Filbert, K., Forester, R. R.,  
27 Fountain, A., Frey, H., Giffen, B., Glasser, N., Gurney, S., Hagg, W., D., H., Haritashya, U.  
28 K., Hartmann, G., Helm, C., Herreid, S., Howat, I., Kapusti, G. G., Khromova, T., Kienholz,  
29 C., Köönig, M., Kohler, J., Kriegel, D., Kutuzov, S., Lavrentiev, I., Le Bris, R., Lund, J.,  
30 Manley, W., Mayer, C., Miles, E., Li, X., Menounos, B., Mercer, A., Mölg, N., Mool, P.,  
31 Nosenko, G., Negrete, A., Nuth, C., Pettersson, R., Racoviteanu, A., Ranzi, R., Rastner, P.,  
32 Rau, F., Raup, B., Rich, J., Rott, H., Schneider, C., Seliverstov, Y., Sharp, M., Siguonson, O.,  
33 Stokes, C., Wheate, R., Winsvold, S., Wolken, G., Wyatt, F. and Zheltyhina, N.: Randolph  
34 Glacier Inventory - A Dataset of Global Outliners: Version 3.2, Boulder Colorado, USA.,  
35 2012.

- 1 Barros, A. P. and Lettenmaier, D. P.: Dynamic modeling of orographically induced  
2 precipitation, *Rev. Geophys.*, 32(3), 265, doi:10.1029/94RG00625, 1994.
- 3 Bartelt, P. and Lehning, M.: A physical SNOATACK model for the Swiss avalanche warning  
4 Part I: numerical model, *Cold Reg. Sci. Technol.*, 35(3), 123–145, doi:10.1016/S0165-  
5 232X(02)00074-5, 2002.
- 6 Bergström, S.: Development and application of a conceptual runoff model for Scandinavian  
7 catchments, *SMHI Reports RHO*, No. 7, Norrköping, 1976.
- 8 Bernhardt, M., Liston, G. E., Strasser, U., Zängl, G. and Schulz, K.: High resolution  
9 modelling of snow transport in complex terrain using downscaled MM5 wind fields,  
10 *Cryosph.*, 4(1), 99–113, doi:10.5194/tc-4-99-2010, 2010.
- 11 Bernhardt, M. and Schulz, K.: SnowSlide: A simple routine for calculating gravitational snow  
12 transport, *Geophys. Res. Lett.*, 37(11), L11502, doi:10.1029/2010GL043086, 2010.
- 13 Bernhardt, M., Zängl, G., Liston, G. E., Strasser, U. and Mauser, W.: Using wind fields from  
14 a high-resolution atmospheric model for simulating snow dynamics in mountainous terrain,  
15 *Hydrol. Process.*, 23(7), 1064–1075, doi:10.1002/hyp.7208, 2009.
- 16 Beven, K. and Freer, J.: Equifinality, data assimilation, and uncertainty estimation in  
17 mechanistic modelling of complex environmental systems using the GLUE methodology, *J.*  
18 *Hydrol.*, 249, 11–29, doi:10.1016/S0022-1694(01)00421-8, 2001.
- 19 Bøggild, C. E., Knudby, C. J., Knudsen, M. B. and Starzer, W.: Snowmelt and runoff  
20 modelling of an Arctic hydrological basin in west Greenland, *Hydrol. Process.*, 13(12-13),  
21 1989–2002, doi:10.1002/(SICI)1099-1085(199909)13:12/13<1989::AID-HYP848>3.0.CO;2-  
22 Y, 1999.
- 23 Dadic, R., Mott, R., Lehning, M. and Burlando, P.: Parameterization for wind-induced  
24 preferential deposition of snow, *Hydrol. Process.*, 24(June), 1994–2006,  
25 doi:10.1002/hyp.7776, 2010a.
- 26 Dadic, R., Mott, R., Lehning, M. and Burlando, P.: Wind influence on snow depth distribution  
27 and accumulation over glaciers, *J. Geophys. Res. Earth Surf.*, 115(1), F01012,  
28 doi:10.1029/2009JF001261, 2010b.
- 29 Daly, C., Halbleib, M., Smith, J. I., Gibson, W. P., Doggett, M. K., Taylor, G. H., Curtis, J.  
30 and Pasteris, P. P.: Physiographically sensitive mapping of climatological temperature and  
31 precipitation across the conterminous United States, *Int. J. Climatol.*, 28, 2031–2064,  
32 doi:10.1002/joc.1688, 2008.
- 33 Daly, C., Neilson, R. P. and Phillips, D. L.: A Statistical-Topographic Model for Mapping  
34 Climatological Precipitation over Mountainous Terrain, *J. Appl. Meteorol.*, 33, 140–158,  
35 doi:10.1175/1520-0450(1994)033<0140:ASTMFM>2.0.CO;2, 1994.

- 1 Dettinger, M., Redmond, K. and Cayan, D.: Winter Orographic Precipitation Ratios in the  
2 Sierra Nevada—Large-Scale Atmospheric Circulations and Hydrologic Consequences, *J.*  
3 *Hydrometeorol.*, 5(1992), 1102–1116, doi:10.1175/JHM-390.1, 2004.
- 4 Donald, J. R., Soulis, E. D., Kouwen, N. and Pietroniro, A.: A Land Cover-Based Snow  
5 Cover Representation for Distributed Hydrologic Models, *Water Resour. Res.*, 31(4), 995–  
6 1009, doi:10.1029/94WR02973, 1995.
- 7 EEA: CORINE Land Cover Project, [online] Available from:  
8 <http://www.eea.europa.eu/publications/COR0-landcover>, 1995.
- 9 Elder, K., Dozier, J. and Michaelsen, J.: Snow accumulation and distribution in an Alpine  
10 Watershed, *Water Resour. Res.*, 27(7), 1541–1552, doi:10.1029/91WR00506, 1991.
- 11 Essery, R., Li, L. and Pomeroy, J.: A distributed model of blowing snow over complex  
12 terrain, *Hydrol. Process.*, 13(14-15), 2423–2438, doi:10.1002/(SICI)1099-  
13 1085(199910)13:14/15<2423::AID-HYP853>3.0.CO;2-U, 1999.
- 14 Farinotti, D., Magnusson, J., Huss, M. and Bauder, A.: Snow accumulation distribution  
15 inferred from time-lapse photography and simple modelling, *Hydrol. Process.*, 24(15), 2087–  
16 2097, doi:10.1002/hyp.7629, 2010.
- 17 Frey, S.: Possible Impacts of Climate Change on the Water Balance with Special Emphasis on  
18 Runoff and Hydropower Potential, University of Natural Resources and Life Sciences,  
19 Vienna. PhD thesis. [online] Available from: <http://permalink.obvsg.at/AC10777542>, 2015.
- 20 Fuchs, M.: Auswirkungen von möglichen Klimaänderungen auf die Hydrologie verschiedener  
21 Regionen in Österreich. (PhD thesis; in German), University of Natural Resources and Life  
22 Sciences, Vienna., 2005.
- 23 Garvelmann, J., Pohl, S. and Weiler, M.: From observation to the quantification of snow  
24 processes with a time-lapse camera network, *Hydrol. Earth Syst. Sci.*, 17(4), 1415–1429,  
25 doi:10.5194/hess-17-1415-2013, 2013.
- 26 Gharari, S., Hrachowitz, M., Fenicia, F., Gao, H. and Savenije, H. H. G.: Using expert  
27 knowledge to increase realism in environmental system models can dramatically reduce the  
28 need for calibration, *Hydrol. Earth Syst. Sci.*, 18(12), 4839–4859, doi:10.5194/hess-18-4839-  
29 2014, 2014.
- 30 Gharari, S., Hrachowitz, M., Fenicia, F. and Savenije, H. H. G.: Moving beyond traditional  
31 model calibration or how to better identify realistic model parameters: sub-period calibration,  
32 *Hydrol. Earth Syst. Sci. Discuss.*, 9(2), 1885–1918, doi:10.5194/hessd-9-1885-2012, 2012.
- 33 Gupta, H. V., Kling, H., Yilmaz, K. K. and Martinez, G. F.: Decomposition of the mean  
34 squared error and NSE performance criteria: Implications for improving hydrological  
35 modelling, *J. Hydrol.*, 377(1-2), 80–91, doi:10.1016/j.jhydrol.2009.08.003, 2009.



- 1 Gupta, H. V., Wagener, T. and Liu, Y.: Reconciling theory with observations: elements of a  
2 diagnostic approach to model evaluation, *Hydrol. Process.*, 22(18), 3802–3813,  
3 doi:10.1002/hyp.6989, 2008.
- 4 Haiden, T., Kann, A., Wittmann, C., Pistotnik, G., Bica, B. and Gruber, C.: The Integrated  
5 Nowcasting through Comprehensive Analysis (INCA) System and Its Validation over the  
6 Eastern Alpine Region, *Weather Forecast.*, 26(2), 166–183,  
7 doi:10.1175/2010WAF2222451.1, 2011.
- 8 Hall, D. K., Riggs, G. A., Salomonson, V. V., DiGirolamo, N. E. and Bayr, K. J.: MODIS  
9 snow-cover products, *Remote Sens. Environ.*, 83(1-2), 181–194, doi:10.1016/S0034-  
10 4257(02)00095-0, 2002.
- 11 Helfricht, K., Schöber, J., Schneider, K., Sailer, R. and Kuhn, M.: Interannual persistence of  
12 the seasonal snow cover in a glacierized catchment, *J. Glaciol.*, 60(223), 889–904,  
13 doi:10.3189/2014JoG13J197, 2014.
- 14 Helfricht, K., Schöber, J., Seiser, B., Fischer, A., Stötter, J. and Kuhn, M.: Snow  
15 accumulation of a high alpine catchment derived from LiDAR measurements, *Adv. Geosci.*,  
16 32, 31–39, doi:10.5194/adgeo-32-31-2012, 2012.
- 17 Henriksen, H. J., Troldborg, L., Nyegaard, P., Sonnenborg, T. O., Refsgaard, J. C. and  
18 Madsen, B.: Methodology for construction, calibration and validation of a national  
19 hydrological model for Denmark, *J. Hydrol.*, 280(1-4), 52–71, doi:10.1016/S0022-  
20 1694(03)00186-0, 2003.
- 21 Herrnegger, M., Nachtnebel, H.-P. and Haiden, T.: Evapotranspiration in high alpine  
22 catchments – an important part of the water balance!, *Hydrol. Res.*, 43(4), 460,  
23 doi:10.2166/nh.2012.132, 2012.
- 24 Hiemstra, C. A., Liston, G. E. and Reiners, W. A.: Observing, modelling, and validating snow  
25 redistribution by wind in a Wyoming upper treeline landscape, *Ecol. Modell.*, 197(1-2), 35–  
26 51, doi:10.1016/j.ecolmodel.2006.03.005, 2006.
- 27 Hock, R.: Temperature index melt modelling in mountain areas, *J. Hydrol.*, 282(1-4), 104–  
28 115, doi:10.1016/S0022-1694(03)00257-9, 2003.
- 29 Holzmann, H., Lehmann, T., Formayer, H. and Haas, P.: Auswirkungen möglicher  
30 Klimaänderungen auf Hochwasser und Wasserhaushaltskomponenten ausgewählter  
31 Einzugsgebiete in Österreich (in german), *Österreichische Wasser- und Abfallwirtschaft*,  
32 62(1-2), 7–14, doi:10.1007/s00506-009-0154-9, 2010.
- 33 Hrachowitz, M., Fovet, O., Ruiz, L., Euser, T., Gharari, S., Nijzink, R., Freer, J., Savenije, H.  
34 H. G. and Gascuel-Oudou, C.: Process consistency in models: The importance of system  
35 signatures, expert knowledge, and process complexity, *Water Resour. Res.*, 50(9), 7445–  
36 7469, doi:10.1002/2014WR015484, 2014.
- 37 Huss, M., Bauder, A. and Funk, M.: Homogenization of long-term mass-balance time series,  
38 *Ann. Glaciol.*, 50(50), 198–206, doi:10.3189/172756409787769627, 2009a.

- 1 Huss, M., Farinotti, D., Bauder, A. and Funk, M.: Modelling runoff from highly glacierized  
2 drainage basins in a changing climate, *Mitteilungen der Versuchsanstalt für Wasserbau,*  
3 *Hydrol. und Glaziologie an der Eidgenoss. Tech. Hochschule Zurich*, 22(213), 123–146,  
4 doi:10.1002/hyp.7055, 2009b.
- 5 Jackson, T. H. R.: A Spatially Distributed Snowmelt-Driven Hydrologic Model applied to the  
6 Upper Sheep Creek Watershed. PhD thesis., Utah State University, Logan, Utah, USA., 1994.
- 7 Jonas, T., Marty, C. and Magnusson, J.: Estimating the snow water equivalent from snow  
8 depth measurements in the Swiss Alps, *J. Hydrol.*, 378(1-2), 161–167,  
9 doi:10.1016/j.jhydrol.2009.09.021, 2009.
- 10 Kirchner, P. B., Bales, R. C., Molotch, N. P., Flanagan, J. and Guo, Q.: LiDAR measurement  
11 of seasonal snow accumulation along an elevation gradient in the southern Sierra Nevada,  
12 California, *Hydrol. Earth Syst. Sci. Discuss.*, 11, 5327–5365, doi:10.5194/hessd-11-5327-  
13 2014, 2014.
- 14 Kling, H.: Spatio-Temporal Modelling of the Water Balance of Austria (PhD-thesis).,  
15 University of Natural Resources and Life Sciences, Vienna., 2006.
- 16 Kling, H., Fuchs, M. and Paulin, M.: Runoff conditions in the upper Danube basin under an  
17 ensemble of climate change scenarios, *J. Hydrol.*, 424-425(0), 264–277,  
18 doi:10.1016/j.jhydrol.2012.01.011, 2012.
- 19 Kling, H., Fürst, J. and Nachtnebel, H. P.: Seasonal, spatially distributed modelling of  
20 accumulation and melting of snow for computing runoff in a long-term, large-basin water  
21 balance model, *Hydrol. Process.*, 20(10), 2141–2156, doi:10.1002/hyp.6203, 2006.
- 22 Kling, H., Stanzel, P., Fuchs, M. and Nachtnebel, H.-P.: Performance of the COSERO  
23 precipitation-runoff model under non-stationary conditions in basins with different climates,  
24 *Hydrol. Sci. J.*, 141217125340005, doi:10.1080/02626667.2014.959956, 2014a.
- 25 Kling, H., Stanzel, P. and Preishuber, M.: Impact modelling of water resources development  
26 and climate scenarios on Zambezi River discharge, *J. Hydrol. Reg. Stud.*, 1, 17–43,  
27 doi:10.1016/j.ejrh.2014.05.002, 2014b.
- 28 Koboltschnig, G. R., Schöner, W., Zappa, M., Kroisleitner, C. and Holzmann, H.: Runoff  
29 modelling of the glacierized Alpine Upper Salzach basin (Austria): Multi-criteria result  
30 validation, in *Hydrological Processes*, vol. 22, pp. 3950–3964, John Wiley & Sons, Ltd.,  
31 2008.
- 32 Lehning, M., Bartelt, P., Brown, B., Fierz, C. and Satyawali, P.: A physical SNOWPACK  
33 model for the Swiss avalanche warning: part II. Snow microstructure, *Cold Reg. Sci.*  
34 *Technol.*, 35(3), 147–167, doi:10.1016/S0165-232X(02)00073-3, 2002.
- 35 Lehning, M. and Fierz, C.: Assessment of snow transport in avalanche terrain, *Cold Reg. Sci.*  
36 *Technol.*, 51(2-3), 240–252, doi:10.1016/j.coldregions.2007.05.012, 2008.

- 1 Liston, G. E.: Representing Subgrid Snow Cover Heterogeneities in Regional and Global  
2 Models, *J. Clim.*, 17(6), 1381–1397, doi:10.1175/1520-  
3 0442(2004)017<1381:RSSCHI>2.0.CO;2, 2004.
- 4 Liston, G. E., Haehnel, R. B., Sturm, M., Hiemstra, C. a., Berezovskaya, S. and Tabler, R. D.:  
5 Simulating complex snow distributions in windy environments using SnowTran-3D, *J.*  
6 *Glaciol.*, 53(181), 241–256, doi:10.3189/172756507782202865, 2007.
- 7 Liston, G. and Sturm, M.: A snow-transport model for complex terrain, *J. Glaciol.*, 44(148),  
8 1998.
- 9 Masson, V., Champeaux, J.-L., Chauvin, F., Meriguet, C. and Lacaze, R.: A Global Database  
10 of Land Surface Parameters at 1-km Resolution in Meteorological and Climate Models, *J.*  
11 *Clim.*, 16(9), 1261–1282, doi:10.1175/1520-0442-16.9.1261, 2003.
- 12 Mauser, W. and Bach, H.: PROMET - Large scale distributed hydrological modelling to study  
13 the impact of climate change on the water flows of mountain watersheds, *J. Hydrol.*, 376(3-4),  
14 362–377, doi:10.1016/j.jhydrol.2009.07.046, 2009.
- 15 Melvold, K. and Skaugen, T.: Multiscale spatial variability of lidar-derived and modeled  
16 snow depth on Hardangervidda, Norway, *Ann. Glaciol.*, 54(62), 273–281,  
17 doi:10.3189/2013AoG62A161, 2013.
- 18 Moore, R. J.: The PDM rainfall-runoff model, *Hydrol. Earth Syst. Sci.*, 11(1), 483–499,  
19 doi:10.5194/hess-11-483-2007, 2007.
- 20 Nachtnebel, H. P., Baumung, S. and Lettl, W.: Abflussprognosemodell für das Einzugsgebiet  
21 der Enns und Steyer (in German), Vienna., 1993.
- 22 Nachtnebel, H. P., Senoner, T., Stanzel, P., Kahl, B., Hernegger, M., Haberl, U. and  
23 Pfaffenwimmer, T.: Inflow prediction system for the Hydropower Plant Gabčíkovo, Part 3 -  
24 Hydrologic Modelling, Bratislava., 2009.
- 25 Nikulin, G., Kjellström, E., Hansson, U., Strandberg, G. and Ullerstig, A.: Evaluation and  
26 future projections of temperature, precipitation and wind extremes over Europe in an  
27 ensemble of regional climate simulations, *Tellus A*, 63(1), 41–55, doi:10.1111/j.1600-  
28 0870.2010.00466.x, 2011.
- 29 Oubeidillah, A. A., Kao, S.-C., Ashfaq, M., Naz, B. S. and Tootle, G.: A large-scale, high-  
30 resolution hydrological model parameter data set for climate change impact assessment for  
31 the conterminous US, *Hydrol. Earth Syst. Sci.*, 18(1), 67–84, doi:10.5194/hess-18-67-2014,  
32 2014.
- 33 Pohl, S., Garvelmann, J., Wawerla, J. and Weiler, M.: Potential of a low-cost sensor network  
34 to understand the spatial and temporal dynamics of a mountain snow cover, *Water Resour.*  
35 *Res.*, 50(3), 2533–2550, doi:10.1002/2013WR014594, 2014.

- 1 Pomeroy, J. W., Gray, D. M., Hedstrom, N. R. and Janowicz, J. R.: Prediction of seasonal  
2 snow accumulation in cold climate forests, *Hydrol. Process.*, 16(18), 3543–3558,  
3 doi:10.1002/hyp.1228, 2002.
- 4 Pomeroy, J. W., Gray, D. M., Shook, K. R., Toth, B., Essery, R. L. H., Pietroniro, A. and  
5 Hedstrom, N.: An evaluation of snow accumulation and ablation processes for land surface  
6 modelling, *Hydrol. Process.*, 12(15), 2339–2367, doi:10.1002/(SICI)1099-  
7 1085(199812)12:15<2339::AID-HYP800>3.0.CO;2-L, 1998.
- 8 Prasad, R., Tarboton, D. G., Liston, G. E., Luce, C. H. and Seyfried, M. S.: Testing a blowing  
9 snow model against distributed snow measurements at Upper Sheep Creek, Idaho, United  
10 States of America, *Water Resour. Res.*, 37(5), 1341–1356, doi:10.1029/2000WR900317,  
11 2001.
- 12 Rasmussen, R. M., Hallett, J., Purcell, R., Landolt, S. D. and Cole, J.: The hotplate  
13 precipitation gauge, *J. Atmos. Ocean. Technol.*, 28, 148–164,  
14 doi:10.1175/2010JTECHA1375.1, 2011.
- 15 Riley, J., Israelsen, E. and Eggleston, K.: Some approaches to snowmelt prediction, *Role*  
16 *Snowmelt Ice Hydrol. IAHS Publ.* 107, 956–971, 1973.
- 17 Rosenbrock, H.: An automatic method for finding the greatest or least value of a function,  
18 *Comput. J.*, 3(3), 175–184, doi:10.1093/comjnl/3.3.175, 1960.
- 19 Rutter, N., Essery, R., Pomeroy, J., Altimir, N., Andreadis, K., Baker, I., Barr, A., Bartlett, P.,  
20 Boone, A., Deng, H., Douville, H., Dutra, E., Elder, K., Ellis, C., Feng, X., Gelfan, A.,  
21 Goodbody, A., Gusev, Y., Gustafsson, D., Hellström, R., Hirabayashi, Y., Hirota, T., Jonas,  
22 T., Koren, V., Kuragina, A., Lettenmaier, D., Li, W. P., Luce, C., Martin, E., Nasonova, O.,  
23 Pumpanen, J., Pyles, R. D., Samuelsson, P., Sandells, M., Schädler, G., Shmakin, A.,  
24 Smirnova, T. G., Stähli, M., Stöckli, R., Strasser, U., Su, H., Suzuki, K., Takata, K., Tanaka,  
25 K., Thompson, E., Vesala, T., Viterbo, P., Wiltshire, A., Xia, K., Xue, Y. and Yamazaki, T.:  
26 Evaluation of forest snow processes models (SnowMIP2), *J. Geophys. Res. Atmos.*, 114,  
27 doi:10.1029/2008JD011063, 2009.
- 28 Safeeq, M., Mauger, G. S., Grant, G. E., Arismendi, I., Hamlet, A. F. and Lee, S.-Y.:  
29 Comparing Large-Scale Hydrological Model Predictions with Observed Streamflow in the  
30 Pacific Northwest: Effects of Climate and Groundwater\*, *J. Hydrometeorol.*, 15(6), 2501–  
31 2521, doi:10.1175/JHM-D-13-0198.1, 2014.
- 32 Schaefli, B., Hingray, B., Niggli, M. and Musy, A.: A conceptual glacio-hydrological model  
33 for high mountainous catchments, *Hydrol. Earth Syst. Sci.*, 9(1/2), 95–109, doi:10.5194/hess-  
34 9-95-2005, 2005.
- 35 Schöber, J., Schneider, K., Helfricht, K., Schattan, P., Achleitner, S., Schöberl, F. and  
36 Kirnbauer, R.: Snow cover characteristics in a glacierized catchment in the Tyrolean Alps -  
37 Improved spatially distributed modelling by usage of Lidar data, *J. Hydrol.*, 519, 3492–3510,  
38 doi:10.1016/j.jhydrol.2013.12.054, 2014.

- 1 Scipi3n, D. E., Mott, R., Lehning, M., Schneebeli, M. and Berne, A.: Seasonal small-scale  
2 spatial variability in alpine snowfall and snow accumulation, *Water Resour. Res.*, 49, 1446–  
3 1457, doi:10.1002/wrcr.20135, 2013.
- 4 Shulski, M. D. and Seeley, M. W.: Application of Snowfall and Wind Statistics to Snow  
5 Transport Modeling for Snowdrift Control in Minnesota, *J. Appl. Meteorol.*, 43(11), 1711–  
6 1721, doi:10.1175/JAM2140.1, 2004.
- 7 Sovilla, B., Mcelwaine, J. N., Schaer, M. and Vallet, J.: Variation of deposition depth with  
8 slope angle in snow avalanches: Measurements from Vall3e de la Sionne., 2010.
- 9 Stanzel, P., Kahl, B., Haberl, U., Herrnegger, M. and Nachtnebel, H. P.: Continuous  
10 hydrological modelling in the context of real time flood forecasting in alpine Danube tributary  
11 catchments, *IOP Conf. Ser. Earth Environ. Sci.*, 4, 012005, doi:10.1088/1755-  
12 1307/4/1/012005, 2008.
- 13 Stanzel, P. and Nachtnebel, H. P.: M3gliche Auswirkungen des Klimawandels auf den  
14 Wasserhaushalt und die Wasserkraftnutzung in 3sterreich (in German), *3sterreichische  
15 Wasser- und Abfallwirtschaft*, 62(9-10), 180–187, doi:10.1007/s00506-010-0234-x, 2010.
- 16 Strasser, U., Bernhardt, M., Weber, M., Liston, G. E. and Mauser, W.: Is snow sublimation  
17 important in the alpine water balance?, *Cryosph.*, 2, 53–66, doi:10.5194/tc-2-53-2008, 2008.
- 18 Thornthwaite, C. W.: An Approach toward a Rational Classification of Climate, *Geogr. Rev.*,  
19 38(1), 55–94, 1948.
- 20 Weber, M., Braun, L., Mauser, W. and Prash, W.: Contribution of rain, snow- and icemelt in  
21 the upper Danube discharge today and in the future, *Geogr. Fis. e Din. Quat.*, 33(2), 221–230,  
22 2010.
- 23 Williams, M. W., Bardsley, T. and Ridders, M.: Overestimation of snow depth and inorganic  
24 nitrogen wetfall using NADP data, Niwot Ridge, Colorado, *Atmos. Environ.*, 32, 3827–3833,  
25 doi:10.1016/S1352-2310(98)00009-0, 1998.
- 26 Winstral, A., Elder, K. and Davis, R. E.: Spatial Snow Modeling of Wind-Redistributed Snow  
27 Using Terrain-Based Parameters, *J. Hydrometeorol.*, 3(5), 524–538, doi:10.1175/1525-  
28 7541(2002)003<0524:SSMOWR>2.0.CO;2, 2002.
- 29 Wood, E. F., Lettenmaier, D. P. and Zartarian, V. G.: A land-surface hydrology  
30 parameterization with subgrid variability for general circulation models, *J. Geophys. Res.*,  
31 97(D3), 2717, doi:10.1029/91JD01786, 1992.

32

1 Table 1. Land use classes used in COSERO (derived from CORINE land cover data, EEA,  
 2 1995) and their proportion in the Ötztal. Snow holding capacities  $H_v$  for each type of land use  
 3 are taken from (Liston and Sturm, 1998; Prasad et al., 2001).


Land use class	proportion [%]	Snow holding capacity $H_v$ [mm]
Build-up areas	1.2	100
Pastures and meadows	20.9	500
Coniferous forests	8.1	2500
Sparsely vegetated areas	20.9	300
Bare rocks	29.5	200
Glaciers	19.4	200

4

1 Table 2. Comparison of performances of model A and B with respect to snow cover and  
 2 runoff. For snow cover coefficient of determination ( $R^2$ ) was used, whereas Kling-Gupta-  
 3 Efficiency (Gupta et al., 2009) was used for runoff.

	Calibration		Validation	
	Snow cover	Runoff	Snow cover	Runoff
	( $R^2$ )	(KGE)	( $R^2$ )	(KGE)
MODEL A	0.78	0.93	0.74	0.92
MODEL B	0.70	0.88	0.66	0.90


4



1  
2


3 Figure 1. Flow chart of the conceptual model COSERO. Potential evapotranspiration is  
 4 estimated using the Thornthwaite method (Thornthwaite, 1948). White parts represent  
 5 distributed processes, greyish parts are calculated on a subbasin scale. Snow transport is  
 6 implemented in the snow cover module.






1  
2  
3  
4  
5  
6  
7  
8

Figure 2. Schematic view of the snow cover in COSERO. a) Composition of one snow class. Vegetation or surface roughness defines the threshold value ( $H_v$ ) to hold back an amount of snow. b) View of one grid cell including five snow classes each of which is composed in the way shown in a). Snowfall is distributed log-normally throughout the classes (dashed lines in b)). Note that snow depth  $S$  is given in mm while all other parameters regarding snow are given in mm SWE.



1  
2  
3  
4  
5


Figure 3. Estimation of the density of snow using Eqs. (8) and (9). Minimum and maximum densities of fresh snow are 100 and 300 kg m<sup>-3</sup>, respectively.



1

2


3 Figure 4. Shapes of the distribution coefficient in dependency of different slope angles and  
 4 snow densities. If cold snow with a density of  $100 \text{ kg m}^{-3}$  is located on a slope of  $35^\circ$ , a  
 5 portion of 25% of the available snow is transported to the neighbour cell. If the snow density  
 6 reaches its maximum value, no transport occurs regardless of the slope.



1

2


3 Figure 5. Conceptual snow accumulations in mountainous regions without (a) and with (b)  
 4 considering lateral snow transport processes. Dotted blocks represent exaggerated snow  
 5 accumulations. Applying the redistribution model snow is transported from the highest grid  
 6 cell to its neighbour where it is treated like solid precipitation. From this grid cell a portion of  
 7 snow gets transported to the downward neighbour again and so forth until either the terrain is  
 8 too flat or snow depths do not exceed the threshold for vegetation (see Fig. 4). Consequently  
 9 less snow remains in the summit region whereas lower grid cells show enhanced  
 10 accumulation. Although snow depths in the summits are lower, the amount of snow covered  
 11 cells stay similar as some residual snow remains in all cells due to  $H_V$  parameterization.



1


2

3 Figure 6. Elevation levels of the Ötztal using a 1x1 km grid (b). Frequency distribution of  
 4 slope angles derived from 1x1 km grid are shown (a). Slopes in general are steeper in the  
 5 summit regions than in the valleys. Note that instead of the average slope of a grid cell only  
 6 steepest vertical gradients are plotted.




1  
2

3 Figure 7. Specific runoff at the outlet at Huben is modelled with (model A) and without  
 4 (model B) using the snow redistribution routine. In the early snow melt period, more runoff is  
 5 generated by model A because snow accumulates rather in lower than in higher levels. In  
 6 summer, enhanced glacier melt leads to more runoff by model A.



1  
2  
3  
4  
5


Figure 8. Accumulated differences (model A minus model B) in discharge at gauge Huben. Using model B, about 300 mm SWE in five years are remaining in the catchment due to snow accumulation processes and less glacier melt.



1  
2  
3  
4  
5


Figure 9. Snow cover in 2009 modelled by both model A and B compared with MODIS data. Error bars refer to uncertainties due to cloud coverage.





1  
2  
3  
4  
5  
6  
7  
8  
9

Figure 10. Behaviour of snow accumulation and melt of model A (a) and B (b) in the upper elevations. Model B leads to “snow towers” of approx. 2900 mm SWE in regions above 3400 m a.s.l. in seven years of modelling, whereas model A does not show such behaviour. In elevations lower 2800 m a.s.l. neither model A nor B show accumulation behaviour. Note that model results are shown from 2005 to 2010 without the warm-up period for clarity reasons. Therefore snow depth does not start at zero in the figure while it does at the beginning of the modelling.



1

2

3 Figure 11. Net snow deposition in the catchment during the time period of one year. Negative  
 4 values refer to a net loss, positive to a net gain of snow. Raster cells in the peak regions act as  
 5 donor cells and do not receive any snow whereas mean elevated cells may act as donor and  
 6 acceptor in the same time. Note that, since only the net deposition of snow based on lateral  
 7 transport is shown, values cannot be linked to snow depths at the end of the time period.