

Revision guide:

This document contains two parts: part I addresses the comments from the referees and our comments to them including our changes on the manuscript. We start with the general comments from both referees and continue with the specific ones. First we answer the ones from referee #2 and later from Daphné Freudiger. Part II starting at page 7 contains the marked-up manuscript that compares the current revision to the re-submitted manuscript of the previous iteration.

1. Point-by-point reply to the comments

We thank Daphné Freudiger and one further referee for their comments and remarks and would like to answer the general remarks in detail. The specific remarks have been applied (see marked-up manuscript). The comments of referee #2 are written in *blue italic*, the comments of Daphné Freudiger are formatted in *orange italic*. Our replies are formatted in black, while changes on the manuscript are formatted in *black italic* and covered in quotation marks. Response to the referee #2

Since both referees criticize the introduction and why a model on the 1x1 km scale only taking into account gravitation was developed, we answer these comments at once.

General comments.

This paper keeps improving, but I still think the problem of mismatch between the spatial scale of the model and those of the physical processes of redistribution (wind and avalanches) remains.

The authors address the maladies of snow distribution and -melt models and I agree that that “snow towers” in conceptual hydrological models is a commonly observed problem. I also agree on that temperature- index models for melting represents a problem for mountainous catchments which contains areas in elevations which never experience temperature above zero. I agree on that, most likely, the reason for the “snow tower” problems are due to how the spatial distribution of snow is modelled. What I find difficult to agree on is that we experience such problems because we have not included redistribution due to wind and avalanches in our 1x1km² scale model. The authors try to justify this approach in the introduction by listing redistribution mechanisms and scale. In section 1.1 we find that “wind redistribution influences snow depth distribution on scales of 100- 1000 square meters” (1000 m² is 0.001 km²). “Avalanches are capable of transporting large snow masses over distance of tens to hundreds of meters.” These scales are far from the scale of the proposed model (1x1km²). It comes hence as a surprise that wind and avalanches are responsible for the erroneous spatial distribution of snow in our conceptual 1x1km² model. I might be wrong in this, but I think that the author have to justify this aspect better so that the reader think that redistribution of snow is the logical solution to snow distribution problems at the 1x1km² scale.

Introduction: As explained in my precedent review, the introduction gives a good review of the snow transport processes and the existing models, but I agree with Referee #1 on the fact, that the introduction, as it is structured now, “appears as a patchy review” and that it is difficult to follow how this literature review leads to the development of the proposed snow transport model. The introduction describes all processes for the variability of snow cover but the model focuses only on snow transport. Since all processes are explained in detail, it is not clear enough why the authors “only” account for the snow transport and not for other processes. Are the processes really taken into account in the model or is only the problem of snow towers solved? I would suggest the authors to completely restructure the introduction in order to lead more logically to the aim of the study.

We agree that the introduction was kind of patchy in a way, that it did not clearly lead to the conclusions, that a snow distribution model on the 1x1 km scale is necessary. We also agree on the better explanation why only gravity was used. We therefore restructured essential parts of the introduction and added some paragraphs to the section 1.2 Modelling approaches:

- *“There exist a plenty of model concepts to simulate the snow redistribution in mountainous areas, ranging from simple conceptual models to complex, physically based ones. The latter attempt to consider all energy fluxes and therefore show a huge data demand with respect to meteorological input. Usually this type requires high spatial resolution of the model domain. Alternatively the conceptual models can also be easily applied for meso and macro scale basins, where the spatial resolution generally shows coarser grid spacing. This is the case for the introduced study where the entire basin of the river Inn was modelled (see Frey et al., 2014) and selective results of the Ötztal are presented. Generally, there are several ways of coping with intensive snow accumulations in hydrological models, in particular (i) adapting the meteorological input data, (ii) the application of physically based models to solve the full energy balance and considering the wind induced snow drift and (iii) conceptual models using topographic information for lateral snow transport. In the following some descriptions and references of the respective concepts are given.”*
- *“Due to the mentioned lack of meteorological stations in high elevations, the use of computer generated wind fields seems appealing. These information have been successfully used to model snow redistributions in small scales of 30 metres (Bernhardt et al., 2009, 2010, 2012). Those models using wind information have in common that they are computationally intensive as they require data in high spatial resolution (e. g. 100 to 1000s of square metres).”*
- *“While the main driving physical processes on the scale of these datasets might differ from the scale of the modelling approaches described above the difficulties of snow accumulations also occur when models with grid cell sizes of 1x1 km are applied to mountainous regions.”*

For further changes, regarding both smaller changes and structural changes see the marked up manuscript.

Discussion:

- What I am still missing in the discussion is a specific discussion on the spatial scale of the model (1x1 km²) (see my precedent comment: p.13, discussion: A discussion on the meaning and the influence of the chosen scale is missing). Since this is an unusual scale for the modeling of snow transport, it would be interesting to read in the discussion what can be reached with a model of this scale and what are the limitations.

We agree on that comment and added a new paragraph to the discussion:

5.5 Scaling issues

“Notwithstanding, that other geomorphological properties than slope angle influencing snow patterns are important on scales smaller than the grid size of COSERO (see section 1.1), slope was selected as driving force for the model. One has to be aware that this is a simplification and under realistic conditions snow might not necessarily be transported only on the steepest route (Bernhardt and Schulz, 2010; Winstral et al., 2002). Also the response of glaciers might change when finer spatial resolutions are applied. In a study at the Blaueisferner, Germany, (Bernhardt et al., 2010) found additional snow getting blown on glacier surfaces when they used a 30 m resolution. At the coarser resolution of 300 m this result could not be found, though. However, as indicated in the introduction,

the 1x1 km scale is often used when hydrological models are applied to medium or large catchments of 100s or 1000s of square kilometres. Not only because of a variety of existing input data sets on that resolution but also for performance issues. The results of this study show that the model operating on that scale is able to reproduce the spatial snow distribution patterns in the catchment and prevent the model from accumulating snow over several years."

- Regarding the transferability of the model, I disagree with the author's response: " We think that modeling and presentation of any additional catchment does not improve the manuscript". I personally think that modeling different catchments would in fact give more reliability to the model results, and especially if the results in other catchments lead to similar conclusions. If for example the comparison of the modeled snow cover with the LiDAR snow cover data is also better in other catchments, it would support the fact that the snow paths chosen by the model for snow transport are more realistic. However, including a paragraph on model transferability can be enough but the argumentation must be developed a little bit more. Especially the comparison with other models is in my opinion in this form not detailed enough. On page 13, line 30, the authors talk about several other models that agree with their results and only the study of Frey (2015) is cited there. What are the other models? And how does this study differ from the present paper? What model was used?

We agree on this comment about developing the argumentation a little bit more, too. We added a study from Switzerland, where several catchments were analysed with different methods (LiDAR, ALS,...) and the elevation where the highest snow accumulations were measured was determined. We briefly described the model used in Frey et al. (2015), too. We think, the combination of both in combination with the better modelling results gives clear indication, that the model is transferable to other catchments.

- Reliability of the model: I disagree with the authors response: " We think that reliability of the model is given by the fact, that the model is able to reproduce both, the discharge and the snow cover extent (MODIS) more accurate." Of course, the fact that the new model gives better efficiency is a good sign that the model is better. But the model could still work "right for the wrong reasons". Therefore, I think that it is important to compare the obtained results with the literature. Of course, glacier mass balance is not the purpose of the paper, but if more glacier melt is produced by Model A compared to Model B, it has to be discussed. As I explained in my earlier comment, I would expect more snow on glacier area with a snow transport model, why is that not the case? The model transports the snow from the summit to the valley, this can be compared to other studies with snow transport models that were successfully conducted in other basins and that could support the results found in this paper. Overall, I find the discussion very interesting but I think it would improve the quality if the results and the model would be compared to existing studies (e.g. the points mentioned about the "Blaueiferner" in the response to the comments). Please include this in your revisions.

We agree on discussing the response of glaciers in more detail. This, however, overlaps with scaling issues. We therefore point to the added paragraph 5.5. In addition, we added a paragraph to section 5.1:

"One has to be aware that the glacier model is very simple, since it treats glaciers as surfaces with infinite depths and static properties. However, besides an advanced algorithm that increases model complexity, a dynamic glacier model would require high spatial resolution information of both the glaciers surfaces as well as the terrain (Farinotti et al., 2009). Since the intention of this paper was to develop a model operating at the 1x1 km scale, this is not feasible."

Figure captions: Most figure captions still contain too much information. Most of the text in the figure captions belongs to the methods, results, or discussion. Please shorten.

We drastically shortened the most of the figure's captions. See marked up manuscript.

It would be good to remind sometimes throughout the paper what model A and B is. It makes it easier to read.

At the beginning of the discussion, we added: “[...] the model accounting for lateral snow transport (model A) [...]”

Detailed comments of referee # 2:

P1,I 13:..resolution for meso- and large scale ?

Right. Changed to “meso”

P1,I 16: ..of the Austrian Inn basin in Tyrol, more specifically the öztaler Ache catchment, but...

Applied the suggested change

P1,I 19-22: reformulate sentence

Reformulated into: *“The results of both model concepts with and without consideration of lateral snow redistribution are compared against observed discharge and snow covered areas derived from MODIS satellite images.”*

P1,I 26:.. predict discharge with more accuracy leading...

Applied the suggested change

P2,I 23: determine whether you will have numbers or letters

We use roman letters

P4,I 13-7: I think you will have to include that the precipitation pattern from the radar was measured a few hundred meters above ground and for a tiny area, 1.5 km. Scale is important here as well.

We changed the passage into: *“Using a Doppler X-band radar capable of a spatial resolution of 75 m Scipi3n et al. (2013) identified significant discrepancies between precipitation patterns in 300 to 600 metres above ground and the snow accumulation at the end of the winter period in a small area of 1.5 km² in the vicinity of Davos, Switzerland. They conclude that snowfall variability at the height of some hundreds of meters above ground is not the driving factor of snow accumulation variabilities at the scale of the radar’s resolution. Consequently, the variability of the meteorological input cannot explain the variability of snow cover patterns.”*

P4,I 26: In addition, ...

Applied the suggested change

P11,I 16-17: drift or transport?, decide on terminology

We deleted “drift” and only use transport.

P12,I 1-2:reformulate sentence

We split the one into two sentences: *“Maximum differences in the mean daily discharges between the two models reach up to 2 mm per day (which equals to 12.1 m³ s⁻¹). This equals a leading to a relative difference of minus -9 up to 44 % of model A in respect to model B.”*

P14-15, l 30: reformulate sentences starting with "For instance.." and the next one starting with "Given that..".

We reconstructed large parts of that paragraph. See marked up manuscript.

P15, l 29:..five years due to longer..

We reconstructed large parts of that paragraph. See marked up manuscript.

Detailed comments of referee # 3 (Daphné Freudiger):

p. 4, l.27: "Schöber et al. [...] with a resolution of 50x50m." I don't understand the relation between this sentence and the previous ones. Please reformulate.

Reformulated into: "A combined approach using gravity and wind induced snow transport was presented by Schöber et al. (2014) who used a distributed energy balance model with a resolution of 50x50 m."

p.7, l.20: the unit of S_{lp} is missing

Units added

p.9, l.17 – 21: move this paragraph to the discussion. This would be a good point to discuss the effects of the spatial scale of the model (see above)

We moved it to the discussion where we added paragraph 5.5. See marked-up manuscript.

p.11, l.21: insert a space before " Besides".

Applied the suggested change

p.12, l.1: replace "minus 9" by "-9".

Applied the suggested change

p.12, l.14: Snow tower was already defined in the introduction, delete "-accumulation of snow cover over several years in high mountain regions".

Applied the suggested change

p.12, l.16-18: "This elevation was chosen..." I don't understand the meaning of this sentence. How did you come to the elevation 2800?

Reformulated the sentence to: "Below that elevation none of the models indicates snow accumulation for more than one year [...]"

p.13, l. 29: insert a space after behavior

Applied the suggested change

p.14, l.1-4: "This is a result..." this sentence is very difficult to understand, please rephrase.

We split the sentence into two ones: "This is a result of the slope dependency of the distribution model. The amount of snow distributed to other grid cells is higher with increasing vertical distance to the downward grid cell (steeper slope)."

p.14, l.17: Please give the percentage of area instead of the amount of grid cells “four grid cells”.

Applied the suggested change

p.14, l.20-21: Was this observed somewhere else? References?

We cannot find a reference for that. Deleted the sentence “*This ratio of summit regions to total catchment size is normally smaller for bigger catchments.*” since it is not necessary for the further discussions.

p.14, l.25: Are there other studies that found 2800m asl? (The whole discussion is lacking references!)

References added.

p.14, l.27: What do you mean with “remain”? I thought that the snow melted more thanks to the snow transport, therefore it should not remain in the basin. Unclear.

Confused model A with B. Should be clear now.

p.14, l.26: Where do the results for the Inn basin come from? Own modeling? Other studies?

Added references and descriptions of where the results come from.

p.15, l.28: -500 mm is the mass balance and 100 mm is the glacier runoff. It is not clear what is meant here.

Rephrased some parts of the conclusions. See marked-up manuscript.

2. Marked-up manuscript showing the changes made with respect to the last iteration of review

(See following pages)

1 A conceptual, distributed snow redistribution model

2 S. Frey¹ and H. Holzmann¹

3 [1]{Institute of Water Management, Hydrology and Hydraulic Engineering, University of
4 Natural Resources and Life Sciences, Vienna, Austria}

5 Correspondence to: S. Frey (simon.frey@boku.ac.at)

7 Abstract

8 When applying conceptual hydrological models using a temperature index approach for
9 snowmelt to high alpine areas often accumulation of snow during several years can be
10 observed. Some of the reasons why these “snow towers” do not exist in nature are vertical and
11 lateral transport processes. While snow transport models have been developed using grid cell
12 sizes of tens to hundreds of square meters and have been applied in several catchments, no
13 model exists using coarser cell sizes of one km², which is a common resolution for meso-~~an~~
14 and large scale hydrologic modelling (hundreds to thousands of square kilometres). In this
15 paper we present an approach that uses only gravity and snow density as a proxy for the age
16 of the snow cover and land-use information to redistribute snow in Alpine basins. The results
17 are based on the hydrological modelling of the Austrian Inn basin in Tyrol, ~~the detailed~~
18 ~~description of the current paper refer to the catchment of Ötztaler Ache,~~ Austria, more
19 specifically the Ötztaler Ache catchment but the findings hold for other tributaries of the river
20 Inn. This transport model is implemented in the distributed rainfall-runoff model COSERO.
21 The results of both model concepts with and without consideration of lateral snow
22 redistribution are compared against observed discharge and snow covered areas derived from
23 MODIS satellite images. A comparison for model validation between the standard model
24 without parameterization for lateral snow redistribution and the updated version is done using
25 observed discharge and MODIS derived snow covered areas. While the signal of snow
26 redistribution can hardly be seen in the binary classification compared with MODIS, By
27 means of the snow redistribution concept snow accumulation over several years can be
28 prevented and the snow depletion curve compared with MODIS data could be improved, too.
29 In a seven year period the standard model would lead to snow accumulation of approximately
30 2900 mm SWE in high elevated regions whereas the updated version of the model does not
31 show accumulation and does also predict discharge more precisely with more accuracy leading

1 to a Kling-Gupta-Efficiency of 0.93 instead of 0.9. A further improvement can be shown in
2 the comparison of MODIS snow cover data and the calculated depletion curve, where the
3 redistribution model increased the efficiency (R^2) from 0,70 to 0,78 (calibration) and from
4 0,66 to 0,74 (validation).

5

6 **1 Introduction**

7 Conceptual models are widely used in hydrology. Examples are the HBV model (Bergström,
8 1976), PDM (Moore, 2007), GSM-SOCONT (Schaepli et al., 2005) or VIC (Wood et al.,
9 1992) just to name a few. Many of these conceptual models use a temperature index approach
10 to model snow melt and snow accumulation and even in some physically based models as
11 e. g. versions of the SHE model (Bøggild et al., 1999) this method can be found. This
12 approach has the advantage of being quite simple since it uses only temperature as input to
13 determine whether precipitation occurs in the form of snow or rain and whether snow can be
14 melted or not. A typical example of a temperature index method for snow modelling is the
15 degree-day approach (see for example Hock 2003). A disadvantage is that snow accumulates
16 as long as the air temperature does not rise above a certain threshold (often 0 °C) regardless of
17 any other processes that may lead to snow melt like radiation or turbulent fluxes of latent
18 energy. In high mountainous areas this may be the case for most days in the year leading to an
19 intensive computational accumulation of snow in these areas. In the modellers terminology
20 these artefacts are often called “snow towers”. In nature, however, these accumulations are
21 barely existent.

22 The reasons for that are either wind or gravitationally induced lateral snow distribution
23 processes (Elder et al., 1991; Winstral et al., 2002). Resulting snow depths are not uniformly
24 distributed in space but vary within large ranges (Helfricht et al., 2014). When changing the
25 focus from micro (e. g. several square meters) to ~~meso~~ scales (e. g. one to several square
26 kilometres), variations become less (Melvold and Skaugen, 2013). The intention of the
27 applied snow redistribution concept was (i) to prevent the artefacts of “snow towers” and
28 (ii) to develop a concept which considers gravity driven lateral snow transport with
29 reasonable and plausible process depiction.

1 **1.1 Theoretical background of snow cover variations**

2 During the accumulation period, according to Liston (2004), primarily three mechanisms are
3 responsible for these variations: (i) snow-canopy interactions in forest covered regions, (ii)
4 wind induced snow redistribution and (iii) orographic influences on snow fall. These
5 mechanisms influence snow cover patterns on scales ranging from the micro to the macro
6 scale. Spatial snow cover variability beneath canopies is mainly affected by different tree
7 species (deciduous vs coniferous trees) influencing LAI, height and density of the canopy and
8 gap sizes (Garvelmann et al., 2013; Liston, 2004; Pomeroy et al., 2002).

9 Besides the impact of vegetation, wind is the most dominant factor influencing snow patterns
10 in alpine terrain. Snow is transported from exposed ridges to the lee side of these ridges,
11 valleys and vegetation covered areas (Essery et al., 1999; Liston and Sturm, 1998; Rutter et
12 al., 2009; Winstral et al., 2002). One has to be aware that besides of the physical transport of
13 solid snow wind also stimulates sublimation processes (Liston and Sturm, 1998; Strasser et
14 al., 2008). Wind influences snow depth distributions on scales of some 100s to 1000 square
15 metres (Dadic et al., 2010a).

16 The third mechanism (orographic effect) influences snow patterns on a larger scale of one to
17 several kilometres (e. g. Barros and Lettenmaier, 1994). Non-uniform snow distributions are
18 caused by interactions of the atmosphere (air pressure, humidity, atmospheric stability) with
19 topography (Liston, 2004).

20 In addition to these processes, avalanches play a role in snow redistribution (Lehning and
21 Fierz, 2008; Lehning et al., 2002; Sovilla et al., 2010). In steep terrain, avalanches depend
22 mainly on the slope angle and are capable of transporting large snow masses over distances of
23 tens to hundreds of metres (Dadic et al., 2010b; Sovilla et al., 2010).

24 During the ablation period, spatial snow distributions are mainly influenced by differences in
25 snow melt behaviour. On the northern hemisphere, on south-facing slopes, rates of snow melt
26 are generally enhanced compared to north-facing slopes due to the inclination of radiation.
27 Also vegetation influences melting behaviour. Shading reduces snowmelt compared to direct
28 sunlight. Enhanced emitted long wave radiation due to warm bare rocks or trees increases the
29 melt rate (Garvelmann et al., 2013; Pohl et al., 2014).

1 1.2 Modelling approaches

2 There exist a plenty of model concepts to simulate the snow redistribution in mountainous
3 areas, ranging from simple conceptual models to complex, physically based ones. The latter
4 attempt to consider all energy fluxes and therefore show a huge data demand with respect to
5 meteorological input. Usually this type requires high spatial resolution of the model domain.
6 Alternatively the conceptual models can also be easily applied for meso and macro scale
7 basins, where the spatial resolution generally shows coarser grid spacing. This is the case for
8 the introduced study where the entire basin of the river Inn was modelled (see Frey et al.,
9 2014) and selective results of the Ötztal are presented.

10 Generally, there are several ways of coping with intensive snow accumulations in
11 hydrological models, in particular (i) adapting the meteorological input data, (ii) the
12 application of physically based models to solve the full energy balance and considering the
13 wind induced snow drift and (iii) conceptual models using topographic information for lateral
14 snow transport. In the following some descriptions and references of the respective concepts
15 are given.

16 A common approach ~~avoiding intensive accumulation of snow~~ is editing the meteorological
17 input (Dettinger et al., 2004). For instance, many models use a constant yet adjustable lapse
18 rate for interpolating temperature with elevation (Holzmann et al., 2010; Koboltschnig et al.,
19 2008). Besides temperature, precipitation gradients are often adjusted to fit observed and
20 modelled target variables (e. g. snow patterns or runoff) (Huss et al., 2009b; Schöber et al.,
21 2014). Justification for doing so is the general lack of gauging stations in the summit regions
22 (Daly et al., 1994, 2008) along with the high error of precipitation gauges (Rasmussen et al.,
23 2011; Williams et al., 1998). An approach presented by Jackson (1994) defining a
24 precipitation correction matrix was successfully applied in several studies (Farinotti et al.,
25 2010; Huss et al., 2009a). Using a Doppler X-band radar capable of a spatial resolution of
26 75 m Scipión et al. (2013) ~~however~~ identified significant discrepancies between precipitation
27 patterns in 300 to 600 metres above ground obtained by a Doppler X-band radar and the snow
28 accumulation at the end of the winter period in a small area of 1.5 km² in the vicinity of
29 Davos, Switzerland. They conclude which gives clear indications that snowfall is
30 redistributed based on different driving forces snowfall variability at the height of some
31 hundreds of meters above ground is not the driving factor of snow accumulation variabilities

1 at the scale of the radar's resolution. ~~Consequently~~ Consequently, the variability of the
2 meteorological input cannot explain the variability of snow cover patterns.

3 Models trying to deal with snow accumulation and redistribution apart from input corrections
4 may be classified into two major approaches. One is the consideration of process based snow
5 distribution patterns. ~~the other approach is empirical.~~ ~~Examples for process-oriented model~~ are
6 SNOWPACK (Bartelt and Lehning, 2002) used in avalanche research or SnowTran3D
7 (Liston et al., 2007; Liston and Sturm, 1998). The other approach is empirical. Empirical
8 ~~models~~ Models following the second approach use the fact, that snow patterns resemble each
9 other every year (Helfricht et al., 2012, 2014). Since our model is following the empirical
10 approach, too, ~~F~~ the presented paper concentrates on ~~the empirical~~ that approach.

11 Snow accumulation gradients determined by airborne LiDAR measurements (Helfricht et al.
12 2012) were used by (Schöber et al., (2014) to improve hydrological modelling using the
13 distributed energy balance model SES (Asztalos, 2004). ~~Helfricht et al. (2012) used airborne~~
14 ~~LiDAR measurements to determine snow accumulation gradients for elevation bands in the~~
15 ~~Ötztaler Alps. These could be used to improve hydrological models regarding snow cover~~
16 ~~distributions and subsequently to achieve better runoff predictions.~~ LiDAR data, however, are
17 relatively expensive to obtain. A common way of dealing with snow accumulations without
18 the need for intensive field campaigns is using ~~Often~~ wind speed and -direction ~~are used~~ to
19 model lateral snow transport drift (e.g. Bernhardt et al., 2009; 2010; Shulski and Seeley,
20 2004; Winstral et al., 2002; Liston and Sturm, 1998). Wind information may be obtained by
21 meteorological stations or by computed wind fields. Kirchner et al. (2014) concluded from
22 LiDAR measurements in combination with meteorological stations in a catchment in
23 California, USA that wind measurements from only one meteorological station are of too poor
24 quality for a useful description of wind fields for snow transport. Due to the mentioned lack
25 of meteorological stations in high elevations, the use of computer generated wind fields seems
26 appealing. These information have been successfully used to model snow redistributions in
27 small scales of 30 metres (Bernhardt et al., 2009, 2010, 2012). Those models using wind
28 information have in common that they are computationally intensive as they require data in
29 high spatial resolution (e. g. 100 to 1000s of square metres). Wind fields may also be
30 generated by regional circulation models (RCM). ~~The~~ ~~However~~ ~~computed wind fields~~
31 ~~generated by regional circulation models (RCM)~~ these wind fields have ~~also~~ shown to be
32 erroneous (Nikulin et al., 2011) and therefore are not useful for direct implementation in

1 redistribution models. ~~Additionally models using wind have in common that they are~~
2 ~~computationally intensive as they require data in high spatial resolution (e. g. 100 to 1000s of~~
3 ~~square metres). A combined approach using gravity and wind induced snow transport was~~
4 ~~presented by~~ Schöber et al. (2014) ~~combined gravitational and wind induced snow transport~~
5 ~~who used~~ing a distributed energy balance model with a resolution of 50x50 m. ~~The~~
6 ~~SnowSlide model (Bernhardt and Schulz, 2010) applied to the Watzmann massif, Germany,~~
7 ~~even only accounts for gravitational induced snow transport.~~

8 ~~The mentioned model approaches have in common that they operate on spatial scales of 10s~~
9 ~~to 100s of metres. However, the difficulties of snow accumulation also occur when models~~
10 ~~with coarser cell sizes are applied. Due~~ However, due to some available databases for
11 vegetation and meteorology (Haiden et al., 2011; Hiebl and Frei, 2015; Masson et al., 2003;
12 Oubeidillah et al., 2014), many models operate on cell sizes of 1 km² or more (e. g. Andersen
13 et al., 2001; Henriksen et al., 2003; Mauser and Bach, 2009; Safeeq et al., 2014). ~~While the~~
14 ~~main driving physical processes on the scale of these datasets might differ from the scale of~~
15 ~~the modelling approaches described above the difficulties of snow accumulations also occur~~
16 ~~when models with grid cell sizes of 1x1 km are applied to mountainous regions. Yet,~~ To our
17 knowledge, no model for redistributing snow on a 1x1 km grid size exists. In this paper we
18 present a simple approach to deal with snow in high mountainous regions and its application
19 in the catchment of Ötztaler Ache in Tyrol, Austria. Since the model uses meteorological
20 input from INCA (Haiden et al., 2011) that already account for meteorological corrections, we
21 focus on snow redistribution rather than to edit the input data. As already mentioned the two
22 main objectives in this respect are to achieve a better model efficiency regarding runoff and to
23 avoid the existence of snow towers at high altitudes.

24

25 **2 Model description**

26 **2.1 Hydrological Model COSERO**

27 COSERO is a spatially distributed conceptual hydrological model which is similar to the
28 HBV model (Bergström, 1976). In the presented paper it uses 1x1 km grid cells. Originally
29 developed for modelling discharge of the Austrian rivers Enns and Steyer (Nachtnebel et al.,
30 1993), it has recently been used for different purposes like climate change studies (e. g. Kling
31 et al., 2012, 2014b; Stanzel and Nachtnebel, 2010), investigating the role of

1 evapotranspiration in high alpine regions (Herrnegger et al., 2012) and operational runoff
2 forecasting (Stanzel et al., 2008). Potential evapotranspiration is calculated using the
3 Thornthwaite method (Thornthwaite, 1948). Discharge due to rainfall and snow-/ice melt is
4 estimated using the same non-linear function of soil moisture as the original HBV. In this
5 study, the model is run using daily time steps. It is, however, capable of using hourly or
6 monthly time steps. In the latter case, intra-monthly variations are considered for snow and
7 interception processes as well as for soil moisture (Kling et al., 2014a). A schematic overview
8 of the model is given by Fig. 1 and a detailed description of the model can be found in Kling
9 et al. (2014a), where the model was applied to several catchments across Europe, Africa and
10 Australia. However, in Kling et al. (2014a) snow parameters were not calibrated and therefore
11 the snow module is not fully explained in detail in their paper. This will be done in the
12 following. Equations (1) to (7) and (10) were taken from the original model by Stanzel and
13 Nachtnebel (2010), all other methods were developed in the present study.

14 Numerous studies have shown that sub-grid variability of snow depths can be described by a
15 two parameter log-normal distribution (e. g. Donald et al., 1995; Pomeroy et al., 1998).
16 COSERO uses five snow classes per cell (i.e. the log-normal distribution is subdivided into
17 five quantiles) to approximate this sub-grid log-normal distribution under accumulation
18 conditions (see Fig. 2 b)), i. e. snowfall is distributed log-normally into snow classes, where
19 the sum of the snow water equivalent (SWE) of each classes represent the mean conditions in
20 the grid cell. This distribution can be interpreted as a statistical description of snow
21 distribution processes taking place at the subgrid scale (Pomeroy et al., 1998). This method
22 has the potential to indirectly consider the influence of curvature, shelter, vegetation or
23 elevation (Hiemstra et al., 2006). The properties of each class are treated unique as equations
24 (1) to (13) apply to every snow class separately. Consequently the log-normal distribution
25 within a grid cell may be disturbed by the processes of melting, sublimation, refreezing and
26 redistribution to other grid cells. Once fallen, snow redistribution between the snow classes
27 within a single grid cell is not considered. A scheme of the composition of a snow class is
28 illustrated in Fig. 2 a). The snow water equivalent (S_{SWEt}) of a given day t per class is
29 calculated by Eq. (1) where P_{Rt} and P_{St} are liquid and solid precipitation in mm, respectively,
30 M_t is snow melt and E_{St} is sublimation of snow. All variables are given in mm SWE.

$$1 \quad S_{SWE_t} = S_{SWE_{t-1}} + P_{R_t} + P_{S_t} - M_t - E_{S_t} \quad (1)$$

2 Snow melt is calculated by a temperature index approach (see for example Hock 2003). Eq.
3 (2) is used:

$$4 \quad M_t = \min(S_{SWE_t}; P_{R_t} \cdot \varepsilon \cdot T_{AIR_t} + D_{f_t} \cdot T_{AIR_t}) \quad (2)$$

5 where M_t is snowmelt [mm], ε is the ratio of specific heat of water and melting energy, T_{AIR_t}
6 is the (mean) daily air temperature [$^{\circ}\text{C}$] and D_{f_t} [$\text{mm } ^{\circ}\text{C}^{-1}$] is the snow melt factor of a given
7 day t estimated by Eq. (3):

$$8 \quad D_{f_t} = \left(-\cos \left(J \cdot \frac{2\pi}{365} \right) \cdot \frac{D_U - D_L}{2} + \frac{D_U - D_L}{2} \right) \cdot M_{RED_t} \quad (3)$$

9 with

$$10 \quad M_{RED_t} = \begin{cases} D_{RED}, & S_{fresh} \geq S_{CRIT} \\ M_{RED_{t-1}} + \frac{(1 - M_{RED_{t-1}})}{5}, & S_{fresh} < S_{CRIT} \end{cases} \quad (4)$$

11 where J is the Julian day of the year [-], D_U and D_L are the upper and lower boundaries of D_f
12 [$\text{mm } ^{\circ}\text{C}^{-1}$], respectively, and M_{RED} [-] is a reduction factor to account for the higher albedo
13 caused by freshly fallen snow calculated by Eq. (4). S_{CRIT} [mm] is the critical snow depth of
14 fresh snow necessary to increase the albedo, whereas S_{fresh} is the actual depth of fresh snow
15 [mm] fallen within one time step. For fresh snow depth larger than S_{CRIT} , M_{RED} is set to a
16 reduced melting factor D_{RED} [-].

17 Whether precipitation occurs in form of snow or rain is controlled by two parameters T_{PS} and
18 T_{PR} , defining the temperature range where snow and rain occur simultaneously. At and above
19 temperature T_{RP} precipitation is pure liquid, at and below T_{PS} precipitation is pure solid. In
20 between those two boundaries, the proportion of solid to liquid precipitation is estimated
21 linearly.

22 For the estimation of snow sublimation, Eq. (5) is used, where E_{SP} [mm] refers to potential
23 sublimation of snow, E_P [mm] is the potential evapotranspiration and E_R is a correction factor
24 to reduce E_P . Sublimation is considered only for snow classes actually covered by snow.
25 Hence, if a grid cell is partly snow free (this can be the case if one subgrid class has no snow
26 cover due to melting) sublimation is estimated for the snow covered part only. For the
27 uncovered classes evapotranspiration according to the Thornthwaite method is applied.

$$28 \quad E_{SP_t} = E_{P_t} \cdot E_R \quad (5)$$

1 The snow cover in COSERO is treated as porous medium and therefore is able to store a
 2 certain amount of liquid water (S_l [$\text{m}^3 \text{kg}^{-1}$]) in dependency of the snow pack density (ρ)
 3 calculated using Eq. (6).

$$4 \quad S_{l_t} = (S_{SWE_t} - S_{l_{t-1}}) \cdot (S_{lMAX} - (\rho - \rho_{MAX}) \cdot S_{l\rho}) \quad (6)$$

5 Where S_{lMAX} [$\text{m}^3 \text{kg}^{-1}$] is the maximum water holding capacity at the maximum snow density
 6 of the snow pack ρ_{MAX} [kg m^{-3}] and $S_{l\rho}$ [-] describes the decrease of water holding capacity
 7 with increasing snow density ρ .

8 At negative air temperatures, retained melt water has the ability to refreeze in the snow pack.
 9 The potential amount of refrozen water (S_R) is estimated by Eq. (7), where R_f is the refreezing
 10 factor [$\text{mm } ^\circ\text{C}^{-1}$]. As long as there is enough liquid water in the snow pack, actual refreezing
 11 will be equal to potential refreezing.

$$12 \quad S_R = \begin{cases} 0, & T_{AIR_t} > 0 \\ R_f \cdot (T_{AIR_t} \cdot (-1)), & T_{AIR_t} \leq 0 \end{cases} \quad (7)$$

13 Refrozen water is treated in the same way as snow. The amount of water leaving the snow
 14 cover then equals snowmelt minus retained water.

15 Snow density (ρ_t) of each class is calculated using a sigmoid function shown in Eqs. (8) and
 16 (9) where ρ_{MAX} and ρ_{MIN} are the respective maximum and minimum values of ρ , T_{AIR} is the
 17 temperature of the air mass above the snow layer and ρ_{scale} and T_{scale} are scaling coefficients
 18 to calculate a transition temperature (T_{tr}) for the estimation of the snow density. Herby, ρ_{scale}
 19 adjusts the slope of the function, whereas T_{scale} is responsible for a shift on the x-axis. These
 20 two parameters are set to fixed values of 1.2 and 1, respectively. The solution of Eqs. ~~(8)(8)~~
 21 and ~~(9)(9)~~ is illustrated in Fig. 3 for a range of typical air temperatures, where snowfall
 22 occurs. Already fallen snow can reach a higher density (ρ_{OLD}) than fresh snow. Its density is
 23 calculated using a time settling constant (ρ_{SET} , derived from Riley et al., 1973) until the
 24 maximum density is reached (Eq. 10).

$$25 \quad \rho_t = (\rho_{MAX} - \rho_{MIN}) \cdot \left(\frac{T_{tr}}{\sqrt{1+(T_{tr})^2}} + 1 \right) \cdot 0.5 + \rho_{MIN} \quad (8)$$

26 with

$$27 \quad T_{tr} = \frac{T_{AIR_t}}{\rho_{scale}} + T_{scale} \quad (9)$$

$$1 \quad \rho_{OLD} = \frac{\rho_{SET} \cdot \left(\frac{S_{SWE_t} + S_t}{2} \right)}{1 + \frac{\rho_{SET}}{2}} \quad (10)$$

2 The COSERO model considers both snow and glacier ice melt processes. Ice melt (M_{ICE}) is
3 computed by means of a degree-day method (see Eq. 11) and uses separate parameter sets.
4 Here, D_{ICE} refers to the ice melt factor [$\text{mm } ^\circ\text{C}^{-1}$]. A prerequisite of ice melt is the full
5 depletion of the overlying snow cover. Spatial information of glaciers are taken from the
6 Randolph Glacier Inventory version 3.2 (Arendt et al., 2012).

$$7 \quad M_{ICE} = D_{ICE} \cdot T_{AIR} \quad (11)$$

8 **2.2 Snow transport model**

9 Several authors reported that the slope angle has an important influence on snow depths
10 (Bernhardt and Schulz, 2010; Kirchner et al., 2014; Schöber et al., 2014). The model
11 redistributes snow only to grid cells providing the steepest slope (acceptor cell) in the direct
12 neighbourhood of the raster cell it searches from (donor cell). Only downward transportation
13 is considered. If more than one cell show the same (largest) difference in elevation, the
14 amount of donated snow is distributed equally to the number of acceptor cells. The actual
15 amount of snow being redistributed depends on the steepness of the slope, the age of the snow
16 cover, considered by the density of snow, the type of land cover of the donor cell and the
17 snow depth of the donor cell. The drier (less dense) the snow pack the higher the snow rate
18 available for the redistribution routine (f_ρ , Eq. 13). Thus the defined maximum density of
19 snow (450 kg m^{-3}) determines the threshold for snow redistribution. The availability of snow
20 for transport is determined by a vegetation-based threshold value (H_v) for each class of land
21 cover. This value can also be interpreted as a roughness coefficient for areas where no or
22 hardly any vegetation is present like in alpine and nival elevations. If the snow depth (S
23 [mm]) of a snow class of a raster cell exceeds H_v [mm], snow transport from that cell is
24 activated and redistribution is calculated by solving Eqs. ~~(12)(12)~~ and ~~(13)(13)~~.

$$25 \quad = \max(S_D - H_v; 0) \cdot f_\rho \cdot \frac{1}{\Sigma A} \cdot C \quad (12)$$

26 With

$$27 \quad f_\rho = \left(\frac{(\rho_{MAX} - \rho_D)}{\rho_{MAX}} \cdot e^{\left(-\frac{\rho_D}{\rho_{MAX}} \right)} \right) \cdot \frac{\alpha}{90} \quad (13)$$

1 Where $S_{SWE(A)}$ is the amount of snow water equivalent that is redistributed from the donor cell
2 (D) to the available acceptor cell(s) (A), ρ_D is the density of snow in the donor cell, ρ_{MAX} is
3 the possible maximum density of snow, α is the angle of the slope between the donor and
4 acceptor cells in degree and C is a correction coefficient that can be calibrated.

5 ~~Notwithstanding, that other geomorphological properties than slope angle influencing snow~~
6 ~~patterns are important on scales smaller than the grid size of COSERO (see section 1.1), slope~~
7 ~~was selected as driving force for the model. One has to be aware that this is a simplification~~
8 ~~and under realistic conditions snow might not necessarily be transported only on the steepest~~
9 ~~route (Bernhardt and Schulz, 2010; Winstral et al., 2002).~~

10 Fig. 4 illustrates the shape of the distribution coefficient f_p as a function of different elevation
11 gradients between the acceptor and donor cells and of the snow density. In acceptor cells
12 redistributed snow is treated as fresh snow in the sense that it is distributed to the snow
13 classes according to the log-normal distribution.

14 The model is organized in form of a loop starting at the highest grid cell (summit region) and
15 ending at the lowest cell (outlet of the catchment). This ensures that snow cannot be
16 redistributed into already processed grid cells. Snow will be transported downslope as long as
17 the slope is big steep enough to allow for transportation given that the density of snow is low
18 enough (see Fig. 4). ~~Consequently, snow accumulates rather in flat regions of the catchment.~~
19 Consequently less snow remains in the summit region whereas lower, rather flat grid cells
20 show enhanced accumulation. Although snow depths in the summits are lower, the amount of
21 snow covered cells stay similar as some residual snow remains in all cells due to HV
22 parameterization.

23 The concept of the redistribution model is sketched in Fig. 5. Note that although snow depths
24 in the highest cell are prevented by the model, the number of snow covered cells remains the
25 same.

26

27 **3 Case study in the catchment the Ötztaler Ache, Tyrol, Austria**

28 **3.1 Catchment description**

29 The catchment of Ötztaler Ache at gauge Huben, situated in western Austria close to the
30 Italian border, covers an area of 511 km² and has an altitudinal range between 1185 m a.s.l at

1 the gauge at Huben and 3770 m a.s.l at its highest peaks. Due to the use of a 1x1 km gridded
2 DEM, the highest grid cell has a mean elevation of 3450 m a.s.l, whereas the lowest cell has
3 an elevation of 1250 m a.s.l. (Fig. 6). About 30 % of its area is covered by vegetation, mainly
4 pastures and meadows. Glaciers cover about 19 % leading to an annual ice melt contribution
5 of about 25 % of the total runoff at Huben, while 41 % of the discharge has its origin in
6 snowmelt (Weber et al., 2010). [Table 1](#) gives an overview of the land cover.

7 In Fig. 6 the elevations of the Ötztal basin are described. Frequency distribution of slope
8 angles derived from 1x1 km grid are shown (6 a). This frequency distribution exhibits the
9 highest frequencies in the slope classes between 20 and 25 degrees for higher elevations. In
10 lower elevated regions slope classes between 0 and 15 degrees dominate. However, also
11 glacier covered areas at the summits can have flat slopes. Note that the listed slopes are based
12 on the steepest vertical gradients of the neighbour elements.

13 **3.2 Input data**

14 Gridded meteorological data of precipitation and air temperature are required to run the
15 model. These data are provided by the INCA dataset (Haiden et al., 2011) with the same grid
16 spacing like the hydrological model, allowing a direct use in the model without the need for
17 pre-processing. INCA data are available since 2003. The years 2003 and 2004 have been used
18 as a warm-up period for the model. In the subsequent years no correction of meteorological
19 data was done since INCA already accounts for elevation gradients regarding air temperature
20 and precipitation. Six land use classes were derived from the most recent CORINE data set
21 (CLC2006 version 17, see EEA, 1995). These classes and their areal fractions in the
22 catchment of Ötztaler Ache are given in [Table 1](#). It should be pointed out, that neither
23 radiation nor wind speed or wind direction data are necessary to run the model.

24 **3.3 Model calibration**

25 The hydrological model was calibrated for the period from 2005 to 2008 using a
26 Rosenbrock's automated optimization routine (Rosenbrock, 1960). Although the model is rich
27 of parameters, the vast majority of them have been estimated *a priori* according to literature
28 (Liston and Sturm, 1998; Prasad et al., 2001) and previous work on the model (Fuchs, 2005;
29 Kling, 2006; Nachtnebel et al., 2009). In the snow model including snow redistribution only
30 six parameters have been calibrated: upper and lower boundaries of snow melt factors D_U and

1 D_L , respectively, the threshold values that control the range where liquid and solid
2 precipitation occur simultaneously (T_{PR} , T_{PS}), the standard deviation of the log-normal
3 distribution of snow depth in one grid cell (N_{VAR}) and the calibration parameter for snow
4 redistribution C (see Eq. 12). The limited number of optimization parameters reduces
5 equifinality problems. For a more detailed description of equifinality issues see the
6 supplements of this article. The target of the calibration was a good fit of runoff using the
7 Kling-Gupta-Model-Efficiency (Gupta et al., 2009; Kling et al., 2012) as objective function.
8 The model was validated for the years 2009 and 2010. Both calibration and validation have
9 been done with and without using the snow ~~transport~~~~drift~~ module. In the following model A
10 refers to the model using snow transport, whereas model B stands for the standard model.
11 Vegetation threshold values for snow detention were taken from previous studies (Liston and
12 Sturm, 1998; Prasad et al., 2001). These are given in ~~Table 1~~~~Table 1~~. Maximum snow density
13 was assumed 450 kg m^{-3} which matches long term snow measurements (Jonas et al., 2009;
14 Schöber et al., 2014). Besides discharge in the validation period also snow cover data from
15 MODIS (8 day maximum snow cover, version 5) satellite images (Hall et al., 2002) were used
16 to compare the performance of both models.

17

18 **4 Results**

19 **4.1 Discharge**

20 Fig. 7 shows a comparison of total discharge using model A and B at the gauge Huben for the
21 year 2006. Both models result in similar quality criteria in the calibration as well as in the
22 validation period (see ~~Table 2~~~~Table 2~~). Nevertheless, the model efficiency could be improved
23 by 0.05 in the calibration period and 0.02 in the validation period by accounting for lateral
24 snow transport. Maximum differences in the mean daily discharges between the two models
25 reach up to 2 mm per day (~~which equals to~~ $12.1 \text{ m}^3 \text{ s}^{-1}$). ~~This equals a leading to a~~ relative
26 difference of ~~minus~~~~9~~ up to 44 % of model A in respect to model B. In total, model A
27 generates a surplus of about 300 mm discharge in five years compared to model B (Fig. 8).
28 About 2/3 of the additional discharge originate in enhanced snowmelt the rest occurs due to
29 enhanced glacier melt.

1 **4.2 Spatially distributed snow cover data**

2 Fig. 9 compares model A and B with MODIS snow depletion data. Both the accumulation
3 period in winter and the ablation period in spring and summer are represented well by both
4 models. Cold snowfall periods in summer generate sharp peaks in the depletion curve, which
5 could be calculated by both model versions, where Model A computed slightly smaller peaks
6 during the snowmelt period (May to July). This leads to a moderate increase of the
7 determination factor R^2 from 0.70 to 0.78 (calibration) and from 0.66 to 0.74 (validation).

8 **4.3 Inter annual snow accumulation**

9 The main reason for developing a snow transport model was the prevention of “snow towers”
10 ~~—accumulation of snow over several years in high mountainous regions.~~ Fig. 10 presents
11 model behaviour of model A and B with respect to the accumulation of snow in elevations
12 above 2800 m a.s.l. ~~This elevation was chosen because here none of the models~~ Below that
13 ~~elevation none of the models~~ indicates snow accumulation for more than one year and
14 therefore snow accumulation in lower altitudes is no problem. By the end of seven years of
15 modelling, model B shows snow depths of approx. 2900 mm SWE in elevations above
16 3400 m a.s.l. whereas model A does hardly show any accumulation behaviour in these
17 altitudes. Spatially distributed net loss and gain of snow for all raster cells within the period of
18 one year in the watershed are presented in Fig. 11. It can be shown that net loss is evident in
19 the zones of ridges and high elevations, where the maximum net gain is along the valley
20 bottoms.

21 **4.4 Parameter equifinality**

22 Since the model uses several parameters that need calibration it suffers from equifinality
23 issues. To investigate those issues, Monte Carlo simulations have been carried out varying the
24 snow relevant parameters that cannot be estimated a priori. Since the aim of this paper is
25 snow transport, the results of the Monte Carlo simulations can be found in the supplements of
26 this article.

27

1 **5 Discussion**

2 **5.1 Discharge**

3 In spring, at the beginning of the melting season, higher runoff is generated by the model
4 accounting for lateral snow transport (model A) due to a larger amount of snow in lower
5 altitudes (see Fig. 7). Later in the year enhanced glacier melt is mainly responsible for higher
6 discharge rates. About 200 mm have their origin in enhanced snowmelt, while the remaining
7 100 mm originate in amplified melt of glaciers. Since glacier cover about 19.4 % of the
8 catchment's area 100 mm of additional mean basin runoff corresponds to an enhanced
9 negative glacier mass balance of -500 mm. The reason for this is transport of snow in warmer
10 altitudes and therefore earlier and more snow free glacier surfaces producing higher discharge
11 due to glacier melt (see Fig. 8) and explains the peak in July and August in runoff difference
12 (see Fig 7).

13 One has to be aware that the glacier model is very simple, since it treats glaciers as surfaces
14 with infinite depths and static properties. However, besides an advanced algorithm that
15 increases model complexity, a dynamic glacier model would require high spatial resolution
16 information of both the glaciers surfaces as well as the terrain (Farinotti et al., 2009). Since
17 the intention of this paper was to develop a model operating at the 1x1 km scale, this is not
18 feasible.

19 **5.2 Spatially distributed snow cover data**

20 Fig. 9 shows the snow depletion curve of the year 2009 based on MODIS data and the
21 comparison of model runs A and B. Only little differences between model A and B can be
22 identified. The reason for this is the vegetation threshold. Even if snow is being transported, a
23 residual of snow remains in the donor cell resulting in the cell marked as snow covered. Grid
24 cells covering the summits only donate snow to their respective acceptor cells. However, a
25 certain amount of snow is held back according to the threshold due to vegetation and
26 roughness of the surface. As indicated in Fig. 5 grid cells nested in the intermediate slope
27 regions receive and donate snow at the same time. Thus their snow depth changes little if
28 comparing model A and model B. In flat valley regions, grid cells only receive snow, where
29 relatively high air temperature values often allow for melting.

1 Satellite based snow cover information by MODIS are binary and so is the model output for
2 comparing these results. In a binary system, no difference can be distinguished between cells
3 covered by much or little snow.

4 **5.3 Snow accumulation**

5 While using model B, the higher the elevation the more snow is accumulated. Contrary,
6 model A shows less pronounced and in some high altitudes even contrary ~~behavior~~behaviour
7 (see Fig. 10). This is a result of the slope dependency of the distribution model. ~~that~~ The
8 amount of snow distributed to other grid cells is higher with increasing vertical distance to the
9 downward grid cell (steeper slope). In general and in the Ötztal as well mountains are steeper
10 in the summit regions than at the bottom (see Fig. 6). Consequently in the summit regions
11 snow will be preferentially eroded while it accumulates at the rather flat valleys where the
12 vertical distances between the grid cells are less than at the peaks. This does reflect snow
13 accumulations that can be observed in nature where summits might be nearly snow free in
14 spring while flatter parts are still covered with snow. While the raster cells covering peak
15 regions act as donators only those cells located on slopes may receive and distribute snow at
16 the same time (Fig. 11). Valley regions only receive snow. The resulting net loss and gain
17 areas shown in Fig. 11 give some indication that the redistribution algorithm is plausible.

18 Although snow accumulation behaviour of model A is more realistic than model B snow
19 accumulation can still be observed in the highest elevations zone (see Fig. 10). This is based
20 on the parameterization of the snow holding capacity H_v , where even bare ground assigns a
21 value of 200 mm (see Table 1). The influence of the highest elevation class (> 3400 m a.s.l.)
22 on both the hydrograph and snow covered area however is very small, since this elevation
23 level is represented by only ~~four grid cells~~0.78 % of the catchment's area. Consequently the
24 objective function during calibration using an automated optimization routine like
25 Rosenbook's routine does not differ much when underestimating the correction coefficient in
26 these grid cells.

27 The smaller the portion of high altitude areas in a catchment compared to the total catchment
28 area the less important is snow redistribution for modelling runoff. ~~This ratio of summit~~
29 ~~regions to total catchment size is normally smaller for bigger catchments.~~ The catchment of
30 river Inn, for instance, covers an area of about 10000 km² yet only 733 km² are located at
31 elevations where intensive snow accumulations and mobilizations occur (above

1 2800 m a.s.l.). In the Ötztal basin 204 out of 511 km² are located higher than 2800 m a.s.l. If
2 model **BA** is applied to the catchment of river Inn in five years of modelling about 15 mm
3 SWE (with respect to the entire river basin) remain in the catchment due to snow
4 accumulation processes instead of 300 mm in the Ötztal. These findings are based on an
5 applied research project for the Austrian Verbund AG, where a hydrological model was
6 applied for the assessment of the hydropower potential of the river Inn (see Frey and
7 Holzmann, 2014; Frey et al., 2014)

8 **5.4 Transferability to other catchments**

9 The model provides results that have been found by other models and field observations, too.
10 The largest snow accumulations occur at the elevation range between 2800 and 3000 m a.s.l.
11 (Fig. 10). This was also found by LiDAR measurements carried out in the same catchment
12 (Helfricht et al., 2012) as well as in several catchments in Switzerland (Grünwald et al.,
13 2014). By applying a simple hydrological model that uses elevation bands instead of raster
14 cells to a variety of other catchments in the Alps, Frey (2015) could identify this elevation
15 range, too. For instance the elevation where the highest snow accumulations occurs (2800 to
16 3000 m a.s.) as was found by LiDAR measurements in the same catchment (Helfricht et al.,
17 2012) as well as by modelling (Frey, 2015). Given that and the needs of the model (slope
18 angles, snow density) for transporting snow, it produces valid results as long as a catchment
19 features relatively steep slopes in the summit regions (which is the case in most catchments in
20 the Alps). Obviously, the model needs calibration if it is transferred to another catchment.

21 **5.4 Parameter equifinality**

22 Like most hydrological models COSERO requires calibration of some parameters. This
23 necessarily causes equifinality issues (Beven and Freer, 2001). The more adjustable
24 parameters a model provides, the more important this problem may become (e. g. Gupta et al.,
25 2008). On the other hand, some authors pointed out that more complex models may produce
26 more feasible results if the parameters can be estimated within realistic boundaries (Gharari et
27 al., 2012, 2014; Hrachowitz et al., 2014). Applying COSERO with the presented snow
28 redistribution routine requires two additional parameters: the vegetation threshold H_V
29 (estimated *a priori*) and the calibration parameter C (see Eq. 12). Yet, accounting for snow
30 redistribution allows the modeller to use D_U values within or close to the range proposed by

1 Kling et al., (2006), while the standard version of the model leads to the best results if higher
2 and therefore unrealistic D_U values are used (see supplements of this article).

3 **5.5 Scaling issues**

4 Notwithstanding, that other geomorphological properties than slope angle influencing snow
5 patterns are important on scales smaller than the grid size of COSERO (see section 1.1), slope
6 was selected as driving force for the model. One has to be aware that this is a simplification
7 and under realistic conditions snow might not necessarily be transported only on the steepest
8 route (Bernhardt and Schulz, 2010; Winstral et al., 2002). Also the response of glaciers might
9 change when finer spatial resolutions are applied. In a study at the Blaueisferner, Germany,
10 found additional snow getting blown on glacier surfaces when they used a 30 m resolution. At
11 the coarser resolution of 300 m this result could not be found, though. However, as indicated
12 in the introduction, the 1x1 km scale is often used when hydrological models are applied to
13 medium or large catchments of 100s or 1000s of square kilometres. Not only because of a
14 variety of existing input data sets on that resolution but also for performance issues. The
15 results of this study show that the model operating on that scale is able to reproduce the
16 spatial snow distribution patterns in the catchment and prevent the model from accumulating
17 snow over several years.

19 **6 Conclusions**

20 A model for redistribution of snow on a coarse 1x1 km raster has been developed and tested
21 in the catchment of Ötztaler Ache, Austria. While only little improvement of snow cover
22 compared to MODIS data could be achieved, appearance of “snow towers” in high altitudes
23 could be prevented. In terms of discharge at the outlet of the basin, both models show good
24 results. However, the Kling-Gupta-efficiency of model A could be improved by 0.05 in the
25 calibration and by 0.02 in the validation period. With respect to the entire watershed area the
26 model using snow redistribution generates a surplus of about 200 mm ~~more~~-runoff originated
27 from snowmelt in five years than without considering this process. This does not only affect
28 the water balance of the catchment, ~~but~~Due to longer time periods where the overlaying snow
29 cover on glacier surfaces is fully depleted, glacier melt is amplified by about 100 mm in five
30 years. With respect to the glaciated area this means that glaciers lose an additional 500 mm of
31 their ice during that period. Since glaciers are represented in a very simple way in this model,

1 these results need to be treated with caution. Nevertheless, glaciers play an important role in
2 the water balance of Alpine catchments and respond on snow redistribution also on the 1x1 km
3 scale. also amplifies glacier melt about 500 mm in five years, with respect to glaciated areas,
4 due to longer time periods where glacier surfaces are fully snow free.

5 The integration of a snow transport module promotes the demand, that models work “right for
6 the right reasons” and is an attempt to integrate more real process understanding into the
7 model approach. Further work needs to be carried out with respect to validation of spatially
8 distributed snow patterns. For this purpose, satellite images from Landsat might be of use
9 providing a higher spatial resolution than MODIS.

10 Even though the vast majority of parameters were estimated *a priori* in this work, equifinality
11 remains an issue. However, redistribution of snow requires only two additional parameters but
12 allows for more realistic boundaries (see Kling et al., 2006) of the snow melt factors (see
13 supplements of this article). However, more work needs to be carried out to account for that
14 issue.

15 **Acknowledgements**

16 The authors thank their colleagues for continuing support and discussion around the coffee
17 breaks, especially to Matthias Bernhardt for friendly reviewing and commenting on the
18 manuscript. Special thanks to Herbert Formayer and David Leidinger of the institute of
19 meteorology, BOKU, for supplying the INCA data. Many thanks to Daphné Freudiger and
20 two further anonymous referees for their comments and valuable suggestions. This study was
21 part of a research project in cooperation with Verbund AG.

23 **References**

24 Andersen, J., Refsgaard, J. C. and Jensen, K. H.: Distributed hydrological modelling of the
25 Senegal River Basin — model construction and validation, *J. Hydrol.*, 247(3-4), 200–214,
26 doi:10.1016/S0022-1694(01)00384-5, 2001.

27 Arendt, A., Bolch, T., Cogley, J. G., Gardner, A., Hagen, J.-O., Hock, R., Kaser, G., Pfeffer,
28 W. T., Moholdt, G., Paul, F., Radić, V., Andreassen, L., Bajracharya, S., Barrand, N., Beedle,
29 M., Berthier, E., Bhambri, R., Bliss, A., Brown, I., Burgess, D., Burgess, E., Cawkwell, F.,
30 Chinn, T., Copland, L., Davies, B., De Angelis, H., Dolgova, E., Filbert, K., Forester, R. R.,
31 Fountain, A., Frey, H., Giffen, B., Glasser, N., Gurney, S., Hagg, W., D., H., Haritashya, U.
32 K., Hartmann, G., Helm, C., Herreid, S., Howat, I., Kapusti, G. G., Khromova, T., Kienholz,

- 1 C., Köönig, M., Kohler, J., Kriegel, D., Kutuzov, S., Lavrentiev, I., Le Bris, R., Lund, J.,
2 Manley, W., Mayer, C., Miles, E., Li, X., Menounos, B., Mercer, A., Mölg, N., Mool, P.,
3 Nosenko, G., Negrete, A., Nuth, C., Pettersson, R., Racoviteanu, A., Ranzi, R., Rastner, P.,
4 Rau, F., Raup, B., Rich, J., Rott, H., Schneider, C., Seliverstov, Y., Sharp, M., Siguosson, O.,
5 Stokes, C., Wheate, R., Winsvold, S., Wolken, G., Wyatt, F. and Zheltyhina, N.: Randolph
6 Glacier Inventory - A Dataset of Global Outliners: Version 3.2, Boulder Colorado, USA.,
7 2012.
- 8 Asztalos, J.: Ein Schnee- und Eisschmelzmodell für vergletscherte Einzugsgebiete, Vienna
9 University of Technology, Vienna, Austria., 2004.
- 10 Barros, A. P. and Lettenmaier, D. P.: Dynamic modeling of orographically induced
11 precipitation, *Rev. Geophys.*, 32(3), 265, doi:10.1029/94RG00625, 1994.
- 12 Bartelt, P. and Lehning, M.: A physical SNOATACK model for the Swiss avalanche warning
13 Part I: numerical model, *Cold Reg. Sci. Technol.*, 35(3), 123–145, doi:10.1016/S0165-
14 232X(02)00074-5, 2002.
- 15 Bergström, S.: Development and application of a conceptual runoff model for Scandinavian
16 catchments, SMHI Reports RHO, No. 7, Norrköping, 1976.
- 17 Bernhardt, M., Liston, G. E., Strasser, U., Zängl, G. and Schulz, K.: High resolution
18 modelling of snow transport in complex terrain using downscaled MM5 wind fields,
19 *Cryosph.*, 4(1), 99–113, doi:10.5194/tc-4-99-2010, 2010.
- 20 Bernhardt, M. and Schulz, K.: SnowSlide: A simple routine for calculating gravitational snow
21 transport, *Geophys. Res. Lett.*, 37(11), L11502, doi:10.1029/2010GL043086, 2010.
- 22 Bernhardt, M., Schulz, K., Liston, G. E. and Zängl, G.: The influence of lateral snow
23 redistribution processes on snow melt and sublimation in alpine regions, *J. Hydrol.*, 424-425,
24 196–206, doi:10.1016/j.jhydrol.2012.01.001, 2012.
- 25 Bernhardt, M., Zängl, G., Liston, G. E., Strasser, U. and Mauser, W.: Using wind fields from
26 a high-resolution atmospheric model for simulating snow dynamics in mountainous terrain,
27 *Hydrol. Process.*, 23(7), 1064–1075, doi:10.1002/hyp.7208, 2009.
- 28 Beven, K. and Freer, J.: Equifinality, data assimilation, and uncertainty estimation in
29 mechanistic modelling of complex environmental systems using the GLUE methodology, *J.*
30 *Hydrol.*, 249, 11–29, doi:10.1016/S0022-1694(01)00421-8, 2001.
- 31 Bøggild, C. E., Knudby, C. J., Knudsen, M. B. and Starzer, W.: Snowmelt and runoff
32 modelling of an Arctic hydrological basin in west Greenland, *Hydrol. Process.*, 13(12-13),
33 1989–2002, doi:10.1002/(SICI)1099-1085(199909)13:12/13<1989::AID-HYP848>3.0.CO;2-
34 Y, 1999.
- 35 Dadic, R., Mott, R., Lehning, M. and Burlando, P.: Parameterization for wind-induced
36 preferential deposition of snow, *Hydrol. Process.*, 24(June), 1994–2006,

- 1 doi:10.1002/hyp.7776, 2010a.
- 2 Dadic, R., Mott, R., Lehning, M. and Burlando, P.: Wind influence on snow depth distribution
3 and accumulation over glaciers, *J. Geophys. Res. Earth Surf.*, 115(1), F01012,
4 doi:10.1029/2009JF001261, 2010b.
- 5 Daly, C., Halbleib, M., Smith, J. I., Gibson, W. P., Doggett, M. K., Taylor, G. H., Curtis, J.
6 and Pasteris, P. P.: Physiographically sensitive mapping of climatological temperature and
7 precipitation across the conterminous United States, *Int. J. Climatol.*, 28, 2031–2064,
8 doi:10.1002/joc.1688, 2008.
- 9 Daly, C., Neilson, R. P. and Phillips, D. L.: A Statistical-Topographic Model for Mapping
10 Climatological Precipitation over Mountainous Terrain, *J. Appl. Meteorol.*, 33, 140–158,
11 doi:10.1175/1520-0450(1994)033<0140:ASTMFM>2.0.CO;2, 1994.
- 12 Dettinger, M., Redmond, K. and Cayan, D.: Winter Orographic Precipitation Ratios in the
13 Sierra Nevada—Large-Scale Atmospheric Circulations and Hydrologic Consequences, *J.*
14 *Hydrometeorol.*, 5(1992), 1102–1116, doi:10.1175/JHM-390.1, 2004.
- 15 Donald, J. R., Soulis, E. D., Kouwen, N. and Pietroniro, A.: A Land Cover-Based Snow
16 Cover Representation for Distributed Hydrologic Models, *Water Resour. Res.*, 31(4), 995–
17 1009, doi:10.1029/94WR02973, 1995.
- 18 EEA: CORINE Land Cover Project, [online] Available from:
19 <http://www.eea.europa.eu/publications/COR0-landcover>, 1995.
- 20 Elder, K., Dozier, J. and Michaelsen, J.: Snow accumulation and distribution in an Alpine
21 Watershed, *Water Resour. Res.*, 27(7), 1541–1552, doi:10.1029/91WR00506, 1991.
- 22 Essery, R., Li, L. and Pomeroy, J.: A distributed model of blowing snow over complex
23 terrain, *Hydrol. Process.*, 13(14-15), 2423–2438, doi:10.1002/(SICI)1099-
24 1085(199910)13:14/15<2423::AID-HYP853>3.0.CO;2-U, 1999.
- 25 Farinotti, D., Huss, M., Bauder, A. and Funk, M.: An estimate of the glacier ice volume in the
26 Swiss Alps, *Glob. Planet. Change*, 68(3), 225–231, doi:10.1016/j.gloplacha.2009.05.004,
27 2009.
- 28 Farinotti, D., Magnusson, J., Huss, M. and Bauder, A.: Snow accumulation distribution
29 inferred from time-lapse photography and simple modelling, *Hydrol. Process.*, 24(15), 2087–
30 2097, doi:10.1002/hyp.7629, 2010.
- 31 Frey, S.: Possible Impacts of Climate Change on the Water Balance with Special Emphasis on
32 Runoff and Hydropower Potential, University of Natural Resources and Life Sciences,
33 Vienna. PhD thesis. [online] Available from: <http://permalink.obvsg.at/AC10777542>, 2015.
- 34 Frey, S., Goler, R., Leidinger, D., Formayer, H. and Holzmann, H.: POWERCLIM - Final
35 Report (in German), Vienna., 2014.
- 36 Frey, S. and Holzmann, H.: Berücksichtigung von Schneeverlagerungsprozessen bei der

1 hydrologischen Modellierung alpiner Einzugsgebiete (in German)., in Tri-Nationaler
2 Workshop – Hydrologische Prozesse im Hochgebirge, Obergurgl, Austria., 2014.

3 Fuchs, M.: Auswirkungen von möglichen Klimaänderungen auf die Hydrologie verschiedener
4 Regionen in Österreich. (PhD thesis; in German)., University of Natural Resources and Life
5 Sciences, Vienna., 2005.

6 Garvelmann, J., Pohl, S. and Weiler, M.: From observation to the quantification of snow
7 processes with a time-lapse camera network, *Hydrol. Earth Syst. Sci.*, 17(4), 1415–1429,
8 doi:10.5194/hess-17-1415-2013, 2013.

9 Gharari, S., Hrachowitz, M., Fenicia, F., Gao, H. and Savenije, H. H. G.: Using expert
10 knowledge to increase realism in environmental system models can dramatically reduce the
11 need for calibration, *Hydrol. Earth Syst. Sci.*, 18(12), 4839–4859, doi:10.5194/hess-18-4839-
12 2014, 2014.

13 Gharari, S., Hrachowitz, M., Fenicia, F. and Savenije, H. H. G.: Moving beyond traditional
14 model calibration or how to better identify realistic model parameters: sub-period calibration,
15 *Hydrol. Earth Syst. Sci. Discuss.*, 9(2), 1885–1918, doi:10.5194/hessd-9-1885-2012, 2012.

16 Grünewald, T., Bühler, Y. and Lehning, M.: Elevation dependency of mountain snow depth,
17 *Cryosph.*, 8(2013), 2381–2394, doi:10.5194/tc-8-2381-2014, 2014.

18 Gupta, H. V., Kling, H., Yilmaz, K. K. and Martinez, G. F.: Decomposition of the mean
19 squared error and NSE performance criteria: Implications for improving hydrological
20 modelling, *J. Hydrol.*, 377(1-2), 80–91, doi:10.1016/j.jhydrol.2009.08.003, 2009.

21 Gupta, H. V., Wagener, T. and Liu, Y.: Reconciling theory with observations: elements of a
22 diagnostic approach to model evaluation, *Hydrol. Process.*, 22(18), 3802–3813,
23 doi:10.1002/hyp.6989, 2008.

24 Haiden, T., Kann, A., Wittmann, C., Pistotnik, G., Bica, B. and Gruber, C.: The Integrated
25 Nowcasting through Comprehensive Analysis (INCA) System and Its Validation over the
26 Eastern Alpine Region, *Weather Forecast.*, 26(2), 166–183,
27 doi:10.1175/2010WAF2222451.1, 2011.

28 Hall, D. K., Riggs, G. A., Salomonson, V. V, DiGirolamo, N. E. and Bayr, K. J.: MODIS
29 snow-cover products, *Remote Sens. Environ.*, 83(1-2), 181–194, doi:10.1016/S0034-
30 4257(02)00095-0, 2002.

31 Helfricht, K., Schöber, J., Schneider, K., Sailer, R. and Kuhn, M.: Interannual persistence of
32 the seasonal snow cover in a glacierized catchment, *J. Glaciol.*, 60(223), 889–904,
33 doi:10.3189/2014JoG13J197, 2014.

34 Helfricht, K., Schöber, J., Seiser, B., Fischer, A., Stötter, J. and Kuhn, M.: Snow
35 accumulation of a high alpine catchment derived from LiDAR measurements, *Adv. Geosci.*,
36 32, 31–39, doi:10.5194/adgeo-32-31-2012, 2012.

- 1 Henriksen, H. J., Troldborg, L., Nyegaard, P., Sonnenborg, T. O., Refsgaard, J. C. and
2 Madsen, B.: Methodology for construction, calibration and validation of a national
3 hydrological model for Denmark, *J. Hydrol.*, 280(1-4), 52–71, doi:10.1016/S0022-
4 1694(03)00186-0, 2003.
- 5 Herrnegger, M., Nachtebel, H.-P. and Haiden, T.: Evapotranspiration in high alpine
6 catchments – an important part of the water balance!, *Hydrol. Res.*, 43(4), 460,
7 doi:10.2166/nh.2012.132, 2012.
- 8 Hiebl, J. and Frei, C.: Daily temperature grids for Austria since 1961—concept, creation and
9 applicability, *Theor. Appl. Climatol.*, doi:10.1007/s00704-015-1411-4, 2015.
- 10 Hiemstra, C. A., Liston, G. E. and Reiners, W. A.: Observing, modelling, and validating snow
11 redistribution by wind in a Wyoming upper treeline landscape, *Ecol. Modell.*, 197(1-2), 35–
12 51, doi:10.1016/j.ecolmodel.2006.03.005, 2006.
- 13 Hock, R.: Temperature index melt modelling in mountain areas, *J. Hydrol.*, 282(1-4), 104–
14 115, doi:10.1016/S0022-1694(03)00257-9, 2003.
- 15 Holzmann, H., Lehmann, T., Formayer, H. and Haas, P.: Auswirkungen möglicher
16 Klimaänderungen auf Hochwasser und Wasserhaushaltskomponenten ausgewählter
17 Einzugsgebiete in Österreich (in german), *Österreichische Wasser- und Abfallwirtschaft*,
18 62(1-2), 7–14, doi:10.1007/s00506-009-0154-9, 2010.
- 19 Hrachowitz, M., Fovet, O., Ruiz, L., Euser, T., Gharari, S., Nijzink, R., Freer, J., Savenije, H.
20 H. G. and Gascuel-Oudou, C.: Process consistency in models: The importance of system
21 signatures, expert knowledge, and process complexity, *Water Resour. Res.*, 50(9), 7445–
22 7469, doi:10.1002/2014WR015484, 2014.
- 23 Huss, M., Bauder, A. and Funk, M.: Homogenization of long-term mass-balance time series,
24 *Ann. Glaciol.*, 50(50), 198–206, doi:10.3189/172756409787769627, 2009a.
- 25 Huss, M., Farinotti, D., Bauder, A. and Funk, M.: Modelling runoff from highly glacierized
26 drainage basins in a changing climate, *Mitteilungen der Versuchsanstalt für Wasserbau,*
27 *Hydrol. und Glaziologie an der Eidgenoss. Tech. Hochschule Zurich*, 22(213), 123–146,
28 doi:10.1002/hyp.7055, 2009b.
- 29 Jackson, T. H. R.: A Spatially Distributed Snowmelt-Driven Hydrologic Model applied to the
30 Upper Sheep Creek Watershed. PhD thesis., Utah State University, Logan, Utah, USA., 1994.
- 31 Jonas, T., Marty, C. and Magnusson, J.: Estimating the snow water equivalent from snow
32 depth measurements in the Swiss Alps, *J. Hydrol.*, 378(1-2), 161–167,
33 doi:10.1016/j.jhydrol.2009.09.021, 2009.
- 34 Kirchner, P. B., Bales, R. C., Molotch, N. P., Flanagan, J. and Guo, Q.: LiDAR measurement
35 of seasonal snow accumulation along an elevation gradient in the southern Sierra Nevada,
36 California, *Hydrol. Earth Syst. Sci. Discuss.*, 11, 5327–5365, doi:10.5194/hessd-11-5327-

- 1 2014, 2014.
- 2 Kling, H.: Spatio-Temporal Modelling of the Water Balance of Austria (PhD-thesis),
3 University of Natural Resources and Life Sciences, Vienna., 2006.
- 4 Kling, H., Fuchs, M. and Paulin, M.: Runoff conditions in the upper Danube basin under an
5 ensemble of climate change scenarios, *J. Hydrol.*, 424-425(0), 264–277,
6 doi:10.1016/j.jhydrol.2012.01.011, 2012.
- 7 Kling, H., Fürst, J. and Nachtnebel, H. P.: Seasonal, spatially distributed modelling of
8 accumulation and melting of snow for computing runoff in a long-term, large-basin water
9 balance model, *Hydrol. Process.*, 20(10), 2141–2156, doi:10.1002/hyp.6203, 2006.
- 10 Kling, H., Stanzel, P., Fuchs, M. and Nachtnebel, H.-P.: Performance of the COSERO
11 precipitation-runoff model under non-stationary conditions in basins with different climates,
12 *Hydrol. Sci. J.*, 141217125340005, doi:10.1080/02626667.2014.959956, 2014a.
- 13 Kling, H., Stanzel, P. and Preishuber, M.: Impact modelling of water resources development
14 and climate scenarios on Zambezi River discharge, *J. Hydrol. Reg. Stud.*, 1, 17–43,
15 doi:10.1016/j.ejrh.2014.05.002, 2014b.
- 16 Koboltschnig, G. R., Schöner, W., Zappa, M., Kroisleitner, C. and Holzmann, H.: Runoff
17 modelling of the glacierized Alpine Upper Salzach basin (Austria): multi-criteria result
18 validation, in *Hydrological Processes*, vol. 22, pp. 3950–3964, John Wiley & Sons, Ltd.,
19 2008.
- 20 Lehning, M., Bartelt, P., Brown, B., Fierz, C. and Satyawali, P.: A physical SNOWPACK
21 model for the Swiss avalanche warning: part II. Snow microstructure, *Cold Reg. Sci.*
22 *Technol.*, 35(3), 147–167, doi:10.1016/S0165-232X(02)00073-3, 2002.
- 23 Lehning, M. and Fierz, C.: Assessment of snow transport in avalanche terrain, *Cold Reg. Sci.*
24 *Technol.*, 51(2-3), 240–252, doi:10.1016/j.coldregions.2007.05.012, 2008.
- 25 Liston, G. E.: Representing Subgrid Snow Cover Heterogeneities in Regional and Global
26 Models, *J. Clim.*, 17(6), 1381–1397, doi:10.1175/1520-
27 0442(2004)017<1381:RSSCHI>2.0.CO;2, 2004.
- 28 Liston, G. E., Haehnel, R. B., Sturm, M., Hiemstra, C. a., Berezovskaya, S. and Tabler, R. D.:
29 Simulating complex snow distributions in windy environments using SnowTran-3D, *J.*
30 *Glaciol.*, 53(181), 241–256, doi:10.3189/172756507782202865, 2007.
- 31 Liston, G. and Sturm, M.: A snow-transport model for complex terrain, *J. Glaciol.*, 44(148),
32 1998.
- 33 Masson, V., Champeaux, J.-L., Chauvin, F., Meriguet, C. and Lacaze, R.: A Global Database
34 of Land Surface Parameters at 1-km Resolution in Meteorological and Climate Models, *J.*
35 *Clim.*, 16(9), 1261–1282, doi:10.1175/1520-0442-16.9.1261, 2003.
- 36 Mauser, W. and Bach, H.: PROMET - Large scale distributed hydrological modelling to study

1 the impact of climate change on the water flows of mountain watersheds, *J. Hydrol.*, 376(3-4),
2 362–377, doi:10.1016/j.jhydrol.2009.07.046, 2009.

3 Melvold, K. and Skaugen, T.: Multiscale spatial variability of lidar-derived and modeled
4 snow depth on Hardangervidda, Norway, *Ann. Glaciol.*, 54(62), 273–281,
5 doi:10.3189/2013AoG62A161, 2013.

6 Moore, R. J.: The PDM rainfall-runoff model, *Hydrol. Earth Syst. Sci.*, 11(1), 483–499,
7 doi:10.5194/hess-11-483-2007, 2007.

8 Nachtnebel, H. P., Baumung, S. and Lettl, W.: Abflussprognosemodell für das Einzugsgebiet
9 der Enns und Steyer (in German), Vienna., 1993.

10 Nachtnebel, H. P., Senoner, T., Stanzel, P., Kahl, B., Hernegger, M., Haberl, U. and
11 Pfaffenwimmer, T.: Inflow prediction system for the Hydropower Plant Gabčíkovo, Part 3 -
12 Hydrologic Modelling, Bratislava., 2009.

13 Nikulin, G., Kjellström, E., Hansson, U., Strandberg, G. and Ullerstig, A.: Evaluation and
14 future projections of temperature, precipitation and wind extremes over Europe in an
15 ensemble of regional climate simulations, *Tellus A*, 63(1), 41–55, doi:10.1111/j.1600-
16 0870.2010.00466.x, 2011.

17 Oubeidillah, A. A., Kao, S.-C., Ashfaq, M., Naz, B. S. and Tootle, G.: A large-scale, high-
18 resolution hydrological model parameter data set for climate change impact assessment for
19 the conterminous US, *Hydrol. Earth Syst. Sci.*, 18(1), 67–84, doi:10.5194/hess-18-67-2014,
20 2014.

21 Pohl, S., Garvelmann, J., Wawerla, J. and Weiler, M.: Potential of a low-cost sensor network
22 to understand the spatial and temporal dynamics of a mountain snow cover, *Water Resour.*
23 *Res.*, 50(3), 2533–2550, doi:10.1002/2013WR014594, 2014.

24 Pomeroy, J. W., Gray, D. M., Hedstrom, N. R. and Janowicz, J. R.: Prediction of seasonal
25 snow accumulation in cold climate forests, *Hydrol. Process.*, 16(18), 3543–3558,
26 doi:10.1002/hyp.1228, 2002.

27 Pomeroy, J. W., Gray, D. M., Shook, K. R., Toth, B., Essery, R. L. H., Pietroniro, A. and
28 Hedstrom, N.: An evaluation of snow accumulation and ablation processes for land surface
29 modelling, *Hydrol. Process.*, 12(15), 2339–2367, doi:10.1002/(SICI)1099-
30 1085(199812)12:15<2339::AID-HYP800>3.0.CO;2-L, 1998.

31 Prasad, R., Tarboton, D. G., Liston, G. E., Luce, C. H. and Seyfried, M. S.: Testing a blowing
32 snow model against distributed snow measurements at Upper Sheep Creek, Idaho, United
33 States of America, *Water Resour. Res.*, 37(5), 1341–1356, doi:10.1029/2000WR900317,
34 2001.

35 Rasmussen, R. M., Hallett, J., Purcell, R., Landolt, S. D. and Cole, J.: The hotplate
36 precipitation gauge, *J. Atmos. Ocean. Technol.*, 28, 148–164,

1 doi:10.1175/2010JTECHA1375.1, 2011.

2 Riley, J., Israelsen, E. and Eggleston, K.: Some approaches to snowmelt prediction, Role
3 Snowmelt Ice Hydrol. IAHS Publ. 107, 956–971, 1973.

4 Rosenbrock, H.: An automatic method for finding the greatest or least value of a function,
5 Comput. J., 3(3), 175–184, doi:10.1093/comjnl/3.3.175, 1960.

6 Rutter, N., Essery, R., Pomeroy, J., Altimir, N., Andreadis, K., Baker, I., Barr, A., Bartlett, P.,
7 Boone, A., Deng, H., Douville, H., Dutra, E., Elder, K., Ellis, C., Feng, X., Gelfan, A.,
8 Goodbody, A., Gusev, Y., Gustafsson, D., Hellström, R., Hirabayashi, Y., Hirota, T., Jonas,
9 T., Koren, V., Kuragina, A., Lettenmaier, D., Li, W. P., Luce, C., Martin, E., Nasonova, O.,
10 Pumpanen, J., Pyles, R. D., Samuelsson, P., Sandells, M., Schädler, G., Shmakin, A.,
11 Smirnova, T. G., Stähli, M., Stöckli, R., Strasser, U., Su, H., Suzuki, K., Takata, K., Tanaka,
12 K., Thompson, E., Vesala, T., Viterbo, P., Wiltshire, A., Xia, K., Xue, Y. and Yamazaki, T.:
13 Evaluation of forest snow processes models (SnowMIP2), J. Geophys. Res. Atmos., 114,
14 doi:10.1029/2008JD011063, 2009.

15 Safeeq, M., Mauger, G. S., Grant, G. E., Arismendi, I., Hamlet, A. F. and Lee, S.-Y.:
16 Comparing Large-Scale Hydrological Model Predictions with Observed Streamflow in the
17 Pacific Northwest: Effects of Climate and Groundwater*, J. Hydrometeorol., 15(6), 2501–
18 2521, doi:10.1175/JHM-D-13-0198.1, 2014.

19 Schaefli, B., Hingray, B., Niggli, M. and Musy, A.: A conceptual glacio-hydrological model
20 for high mountainous catchments, Hydrol. Earth Syst. Sci., 9(1/2), 95–109, doi:10.5194/hess-
21 9-95-2005, 2005.

22 Schöber, J., Schneider, K., Helfricht, K., Schattan, P., Achleitner, S., Schöberl, F. and
23 Kirnbauer, R.: Snow cover characteristics in a glacierized catchment in the Tyrolean Alps -
24 Improved spatially distributed modelling by usage of Lidar data, J. Hydrol., 519, 3492–3510,
25 doi:10.1016/j.jhydrol.2013.12.054, 2014.

26 Scipión, D. E., Mott, R., Lehning, M., Schneebeli, M. and Berne, A.: Seasonal small-scale
27 spatial variability in alpine snowfall and snow accumulation, Water Resour. Res., 49, 1446–
28 1457, doi:10.1002/wrcr.20135, 2013.

29 Shulski, M. D. and Seeley, M. W.: Application of Snowfall and Wind Statistics to Snow
30 Transport Modeling for Snowdrift Control in Minnesota, J. Appl. Meteorol., 43(11), 1711–
31 1721, doi:10.1175/JAM2140.1, 2004.

32 Sovilla, B., Mcelwaine, J. N., Schaer, M. and Vallet, J.: Variation of deposition depth with
33 slope angle in snow avalanches: Measurements from Vallée de la Sionne., 2010.

34 Stanzel, P., Kahl, B., Haberl, U., Herrnegger, M. and Nachtnebel, H. P.: Continuous
35 hydrological modelling in the context of real time flood forecasting in alpine Danube tributary
36 catchments, IOP Conf. Ser. Earth Environ. Sci., 4, 012005, doi:10.1088/1755-
37 1307/4/1/012005, 2008.

- 1 Stanzel, P. and Nachtnebel, H. P.: Mögliche Auswirkungen des Klimawandels auf den
2 Wasserhaushalt und die Wasserkraftnutzung in Österreich (in German), Österreichische
3 Wasser- und Abfallwirtschaft, 62(9-10), 180–187, doi:10.1007/s00506-010-0234-x, 2010.
- 4 Strasser, U., Bernhardt, M., Weber, M., Liston, G. E. and Mauser, W.: Is snow sublimation
5 important in the alpine water balance?, *Cryosph.*, 2, 53–66, doi:10.5194/tc-2-53-2008, 2008.
- 6 Thornthwaite, C. W.: An Approach toward a Rational Classification of Climate, *Geogr. Rev.*,
7 38(1), 55–94, 1948.
- 8 Weber, M., Braun, L., Mauser, W. and Prasch, W.: Contribution of rain, snow- and icemelt in
9 the upper Danube discharge today and in the future, *Geogr. Fis. e Din. Quat.*, 33(2), 221–230,
10 2010.
- 11 Williams, M. W., Bardsley, T. and Ridders, M.: Overestimation of snow depth and inorganic
12 nitrogen wetfall using NADP data, Niwot Ridge, Colorado, *Atmos. Environ.*, 32, 3827–3833,
13 doi:10.1016/S1352-2310(98)00009-0, 1998.
- 14 Winstral, A., Elder, K. and Davis, R. E.: Spatial Snow Modeling of Wind-Redistributed Snow
15 Using Terrain-Based Parameters, *J. Hydrometeorol.*, 3(5), 524–538, doi:10.1175/1525-
16 7541(2002)003<0524:SSMOWR>2.0.CO;2, 2002.
- 17 Wood, E. F., Lettenmaier, D. P. and Zartarian, V. G.: A land-surface hydrology
18 parameterization with subgrid variability for general circulation models, *J. Geophys. Res.*,
19 97(D3), 2717, doi:10.1029/91JD01786, 1992.
- 20

1 Table 1. Land use classes used in COSERO (derived from CORINE land cover data, EEA,
 2 1995) and their proportion in the Ötztal. Snow holding capacities H_v for each type of land use
 3 are taken from (Liston and Sturm, 1998; Prasad et al., 2001).


Land use class	proportion [%]	Snow holding capacity H_v [mm]
Build-up areas	1.2	100
Pastures and meadows	20.9	500
Coniferous forests	8.1	2500
Sparsely vegetated areas	20.9	300
Bare rocks	29.5	200
Glaciers	19.4	200

4

1 Table 2. Comparison of performances of model A and B with respect to snow cover and
 2 runoff. For snow cover coefficient of determination (R^2) was used, whereas Kling-Gupta-
 3 Efficiency (Gupta et al., 2009) was used for runoff.

	Calibration		Validation	
	Snow cover	Runoff	Snow cover	Runoff
	(R^2)	(KGE)	(R^2)	(KGE)
MODEL A	0.78	0.93	0.74	0.92
MODEL B	0.70	0.88	0.66	0.90

4



1


2

3 Figure 1. Flow chart of the conceptual model COSERO. ~~Potential evapotranspiration is~~

4 ~~estimated using the Thornthwaite method (Thornthwaite, 1948).~~ White parts represent

5 distributed processes, greyish parts are calculated on a subbasin scale. Snow transport is


6 implemented in the snow cover module.



1


2

3 Figure 2. Schematic view of the snow cover in COSERO. a) Composition of one snow class.
 4 Vegetation or surface roughness defines the threshold value (H_v) to hold back an amount of
 5 snow. b) View of one grid cell including five snow classes each of which is composed in the
 6 way shown in a). Snowfall is distributed log-normally throughout the classes (dashed lines in
 7 b)). Note that snow depth S is given in mm while all other parameters regarding snow are
 8 given in mm SWE.



1
2
3
4
5


Figure 3. Estimation of the density of snow using Eqs. (8) and (9). Minimum and maximum densities of fresh snow are 100 and 300 kg m⁻³, respectively.



1

2


3 Figure 4. Shapes of the distribution coefficient in dependency of different slope angles and
 4 snow densities. ~~If cold snow with a density of 100 kg m^{-3} is located on a slope of 35° , a~~
 5 ~~portion of 25% of the available snow is transported to the neighbour cell. If the snow density~~
 6 ~~reaches its maximum value, no transport occurs regardless of the slope.~~



1

2


3 Figure 5. Conceptual snow accumulations in mountainous regions without (a) and with (b)
 4 considering lateral snow transport processes. Dotted blocks represent exaggerated snow
 5 accumulations. ~~Applying the redistribution model snow is transported from the highest grid
 6 cell to its neighbour where it is treated like solid precipitation. From this grid cell a portion of
 7 snow gets transported to the downward neighbour and so forth until either the terrain is
 8 too flat or snow depths do not exceed the threshold for vegetation (see Fig. 4). Consequently
 9 less snow remains in the summit region whereas lower grid cells show enhanced
 10 accumulation. Although snow depths in the summits are lower, the amount of snow covered
 11 cells stay similar as some residual snow remains in all cells due to H_v parameterization.~~



1


2

3 Figure 6. Elevation levels of the Ötztal using a 1x1 km grid (b). Frequency distribution of
 4 slope angles derived from 1x1 km grid are shown (a). Slopes in general are steeper in the
 5 summit regions than in the valleys. Note that instead of the average slope of a grid cell only
 6 steepest vertical gradients are plotted.




1
2

3 Figure 7. Specific runoff at the outlet at Huben is modelled with (model A) and without
 4 (model B) using the snow redistribution routine. ~~In the early snow melt period, more runoff is~~
 5 ~~generated by model A because snow accumulates rather in lower than in higher levels. In~~
 6 ~~summer, enhanced glacier melt leads to more runoff by model A.~~




1
2
3
4
5

Figure 8. Accumulated differences (model A minus model B) in discharge at gauge Huben. ~~Using model B, about 300 mm SWE in five years are remaining in the catchment due to snow accumulation processes and less glacier melt.~~




1
2
3
4
5

Figure 9. Snow cover in 2009 modelled by both model A and B compared with MODIS data. Error bars refer to uncertainties due to cloud coverage.



1
2
3
4
5
6
7
8
9

Figure 10. Behaviour of snow accumulation and melt of model A (a) and B (b) in the upper elevations. ~~Model B leads to “snow towers” of approx. 2900 mm SWE in regions above 3400 m a.s.l. in seven years of modelling, whereas model A does not show such behaviour. In elevations lower 2800 m a.s.l. neither model A nor B show accumulation behaviour.~~ Note that model results are shown from 2005 to 2010 without the warm-up period for clarity reasons. Therefore snow depth does not start at zero in the figure while it does at the beginning of the modelling.



1

2

3 Figure 11. Net snow deposition in the catchment during the time period of one year. Negative
 4 values refer to a net loss, positive to a net gain of snow. ~~Raster cells in the peak regions act as~~
 5 ~~donor cells and do not receive any snow whereas mean elevated cells may act as donor and~~
 6 ~~acceptor in the same time.~~ Note that, since only the net deposition of snow based on lateral
 7 transport is shown, values cannot be linked to snow depths at the end of the time period.