

Date: 12-07-2015

To,
The Editor
Hydrology and Earth System Sciences Discussions.

Sub: Submission of revised manuscript hess-2015-173-discussions

Dear Sir,

I would like to submit the revised version of the manuscript entitled 'Investigating the impact of land-use land-cover change on Indian summer monsoon daily rainfall and temperature during 1951–2005 using a regional climate model' by authors S. Halder, S. K. Saha, P. A. Dirmeyer, T. N. Chase and B.N. Goswami. Along with revised version I would also like to submit other files that show point-by-point response to the reviewers' and Editor's comments, abstract, tracked changes version of the article file that shows the edits, supplementary file and figures.

We thank the Editor as well as the reviewers for their insightful comments and suggestions that helped in improvement of the manuscript in several ways. We have incorporated the suggestions in the revised manuscript and mentioned about the changes made while responding to them. I would also like to mention that the line numbers increase continuously from the start of the revised manuscript till the end. Hence, such numbers have been used while referring to the changes made in the text.

I certify the manuscript has not been published in any language in any other journal and that it is not being submitted to any other journal. Also all co-authors have agreed to its submission. I look forward to your final decision.

Thanking you,
Yours sincerely,

Subhadeep Halder.

Subhadeep Halder, Ph.D.
Post-doctoral Research Fellow
Center for Ocean-Land-Atmosphere Studies
George Mason University
261 Research Hall, Mail Stop 6C5
4400 University Drive
Fairfax-22030, VA

Off.: (703)-993-5728
Mob.: (703)-901-9257

I) Reply to comments by the Editor

1) Comments to the Author: Dear Dr. Halder, We have now at last received to reviewer reports on your manuscript - my apologies for the delay. It is my pleasure to report that the reviews report positively on the topic of study and the relevance of the obtained outcomes. However, both reviewers raise relevant points about the experimental setup, and I agree with the reviewers that these will have to be addressed before the manuscript can be considered for publication in HESS, if not by adjusting the experiment then at least with a detailed justification of the current setup.

Ans. We would like to thank the editor for his comments and suggestions. We have provided justification for the current experiments with a RCM in the sections Introduction (P6 and P7) and Design of Experiments and Methodology (P11). Our justifications for using a RCM are as follows:

‘Krishnan et al. (2015) made several experiments with a high resolution global atmospheric model and concluded that a multitude of factors such as aerosols, land-use change, Indian Ocean warming as well as GHGs have together contributed to the observed weakening of the south-Asian monsoon and changes in frequency distribution of daily rainfall events during the later half of the 20th century. However, the impact of LULCC as a lone forcing component on the Indian summer monsoon has not been quantified. It is also plausible that feedbacks due to variations in remote SSTs and snow cover may have modulated the local impacts due to LULCC. In this study, we hypothesize and demonstrate that LULCC has partly contributed to the observed decrease in moderate rainfall events over CI during the monsoon season from 1951 through 2005, apart from the increasing trend in daily mean and maximum temperatures. We have conducted experiments with a high-resolution regional climate model (RCM) RegCM4.0 and accurate LULC data over the Indian region to prove our hypothesis. No added external forcing in terms of aerosols or GHG concentration is used in our experiments. Furthermore, additional experiments by removing the positive trend in Indian Ocean SSTs have also been made to isolate the impact of LULCC.

RCMs have shown much improvement over global climate models (GCMs) in terms of representation of spatio-temporal details of climate (Giorgi and Mearns, (1999); Laprise et al., 2008; Leduc and Laprise, 2009) and dynamical downscaling ability (Xue et al., 2014) and add value in simulation of topography induced phenomena and extremes of short spatio-temporal character (Feser et al., 2011; Feser and Barcikowska, 2012; Shkol’nik et al., 2012). Saha et al. (2011, 2012) and Halder et al. (2015) have made experiments with the RCM RegCM3 and RegCM4, respectively to better resolve regional land-atmosphere feedback processes and demonstrate their role in the mean and variability of the Indian summer monsoon. When time-dependent lateral boundary conditions are used as forcing for a RCM in one-way mode, feedback from the model-simulated climate to the global climate is not allowed. Such interactions between large-scale forcing such as El Niño–Southern Oscillation (ENSO) that is external to the Indian monsoon region and internal monsoon dynamics may lead to more variability than due to

local feedback processes. Therefore, our methodology helps in segregation of the impact of regional LULCC on the Indian summer monsoon and its changes. However, one of the drawbacks of regional climate modeling is that lateral boundary conditions are not perfect.'

'Our objective is to analyze changes in the climatological mean of the number of moderate rainfall events over CI and intensity of rainfall in that category, between PLC and HLC experiments. The lateral boundary forcing and prescribed SST in our experiments are transient in nature and impose the global warming signal on the model climate. As each year of forcing is different from the other, long term mean is expected to be closer to the reality. However, use of climatological boundary conditions is not an option, as in that case the model will have problem in properly capturing the synoptic and intraseasonal rainfall variability that contribute to the seasonal mean rainfall significantly. Similarly, a single year (ENSO/non-ENSO year) of boundary condition cannot be repeated as that may lead to biased response of the model climate to LULCC. As time varying lateral boundary conditions also impose the effect of variations in remote SSTs, such as that of the Pacific Ocean on the model, partial remote influence on the nature of response due to regional LULCC is possible. Although our RCM simulations are not time-slice experiments in the true sense, the statistics of their difference are expected to reveal the effect of LULCC and associated regional land-atmosphere feedbacks on daily temperature and rainfall variability in a climatological sense.'

Furthermore, we have also addressed the comments made by the respective reviewers in detail.

Additionally, reading carefully through the manuscript, I identify a couple of further issues with the presentation, which I think need addressing:

2) I believe that the manuscript needs further editing for language, but can also benefit from a more condensed, less repetitive, and more precise writing style. For instance, try to avoid qualitative statements such as "reasonably well" (p6593, 6594, 6595), use the more formal "until" instead of "till". On p.6588, l.1-4: remove this section; it is sufficient to outline the structure of the manuscript once (p6583) and then simply follow the outlined structure. These are just a couple of examples, but I encourage you to go carefully through the whole manuscript.

Ans. We thank the editor for his suggestions and have incorporated the suggested changes in the revised manuscript.

P16, l386: 'It is evident that the RCM reproduces the regions of rainfall maxima and the spatial pattern very well, particularly over the Western Ghats section over peninsular India, CI, north-east India and the Himalayan foothills.'

P16, l402: 'Therefore, the model performs well in capturing the observed interannual variability of seasonal rainfall over CI (which is about 10% of the seasonal mean), although it underestimates both quantities.'

P17, 1420: ‘The RCM captures the location of these large-scale low-level features very well in both PLC and PLCS (Fig. S2b and c).’

The line in page P6595 that was the older version of the manuscript has now been deleted.

The section on p.6588, 1.1-4 in earlier version of the manuscript has been removed in order to avoid repetition.

3) The literature review reads rather like a listing of past findings with references. I would encourage you to make this more strongly into a coherent argumentation about the state of the art in the research field, and how your experimental setup will push the boundaries of current knowledge.

Ans. We appreciate the editor’s suggestion about this aspect. Keeping that in mind, we have modified the Introduction (P3-P7) to make it a more coherent argument about the state of the art in the research field, and how our experimental setup will help in investigating the impact of regional LULCC on the Indian summer monsoon and its changes during 1951 until 2005.

II) Reply to Interactive comments by Reviewer 1 (Dr. O. Tuinenburg) on **“Investigating the impact of land-use land-cover change on Indian summer monsoon daily rainfall and temperature during 1951–2005 using a regional climate model”** by S. Halder et al.

General comments

The authors study realistic land-use and land-cover changes from 1951-2005 over India in a regional climate model and show that moderate rainfall events are impacted significantly by these changes. Moreover, they have repeated the simulations without the observed ocean warming trend to confirm the effects of the land surface on the Indian climate.

In my opinion, this is an important subject to study as India is a part of the world where land-use has changed significantly and the effects of changes in rainfall distribution are important, for example for agriculture.

I think the study is well executed and the conclusions drawn are supported by the experiments. My main concern is the fact that only one model is used. As regional climate models respond different to these kinds of land surface changes, this uncertainty should be discussed more thorough. Moreover, some textual improvements should be made.

Ans. We appreciate the views of the reviewer and the encouragement. Our reply to his queries may be found below.

Specific comments

1) P6593, l20: Here, you state that RCMs have trouble to capture the observed frequency distribution of rainfall. I think this is true. However, in the discussion (section 5), it is only briefly mentioned. I think that is a place to convince the reader that the results found are due to the changes in the model and not due to other factors (parameterization schemes, resolution differences, boundary forcing, etc.).

Ans. We appreciate the reviewer's comments in this aspect. We have now modified that portion in section 5 (P28, l719-l725).

‘It is important to note that the deficiency in the RCM in terms of capturing the frequency distribution of daily very heavy rainfall events over CI realistically could have a bearing on our inferences. Hence, our results are partly dependent on the choice of model parameterization schemes. However, this is a well-known problem related to climate models (Frei et al., 2003; Kang et al., 2014) and similar studies when repeated with other RCMs is expected to give us further evidence on the role of LULCC in affecting the frequency distribution of daily rainfall events over India.’

We have further added few lines at the end of the Abstract (P2, l25-l27 and P3 l28-l29).

‘Although, the regional climate model helps in better resolving land-atmosphere

feedbacks over the Indian region, the inferences do depend on the fidelity of the model in capturing the features of Indian monsoon realistically. It is proposed that similar studies using a suite of climate models will further enrich our understanding about the role of LULCC in Indian monsoon climate.'

2) In many places, the text is vague or grammatically incorrect. Please have a look at it. (See textual comments below for some examples.)

Ans. We thank the reviewer for identifying these deficiencies in the text. Due care has been taken in rectifying them in the revised manuscript.

Minor and textual comments:

3) Use of brackets to state the opposite is confusing to me, maybe separate sentences can be made. One example, but many occurrences in the text: '... region of decreased (increased) forest (crop) ...' (P6578, 110).

Ans. We thank the reviewer for identifying this aspect. We have modified that sentence (P2, 19-111) and other such sentences wherever it was suitable, in the revised manuscript.

'Model simulations show that the increase in seasonal mean and extreme temperature over central India coincides with the region of decrease in forest and increase in crop cover.'

4) P6578, 115: argues → argue.

Ans. We have corrected that word (P3, 141)

'On the contrary, Seneviratne et al. (2012) opine that ...'

5) P6580, 120: the effects → the climate effects

Ans. We thank the reviewer for this suggestion. The correction has been made accordingly (P4, 155).

6) P6580, 126: fix accent in Ducroudre.

Ans. We thank the reviewer suggesting the correction for the accent. That has now been fixed (P4, 165).

7) P6584, 113: I visited the website, but found that the data are only available for purchase. This makes it hard to repeat the detection of precipitation changes. Would it not be better to use the APHRODITE dataset, which is publicly available?

Ans. We have repeated the analysis with APHRODITE data (P8, 1165-1169) and replaced the earlier figure. The new Figure 2 is based on APHRODITE data.

‘For analysis of trends in daily rainfall events and their intensities in different categories daily gridded data for 55 years (1951–2005) from the Asian Precipitation Highly-Resolved Observational Data Integration Towards Evaluation of Water Resources (APHRODITE Water Resources, Yatagai et al., 2012) at 0.5° x 0.5° resolution is used.’

8) P6586, l21: remove degree signs.

Ans. We have removed the degree signs (P10, l220-l225).

‘CLM3.5 also uses global datasets on canopy top and bottom height (resolution 0.5x0.5), percentage of glacier (resolution 0.5x0.5), lake and wetland (resolution 1x1) with corresponding spatial resolution included in brackets (Elguindi et al., 2010).’

9) Paragraph 2.3: How is the orography treated? This is a very important aspect, so maybe include it here.

Ans. We thank the reviewer for this suggestion. We have included information on the orography in P10, l215-l217.

‘The GTOPO30 topography data at 30 arc seconds resolution courtesy of the U.S.G.S. (United States Geological Survey) EROS Data Center has been used in this study.’

10) P6588, l4-5: "Extreme rainfall ... return periods.", this is a strange sentence, seems to be a truism, maybe it can be changed into something more informative.

Ans. We thank the reviewer for this suggestion. The sentence has now been modified (P11, l265).

‘Extreme rainfall events are short lived, less frequent but intense.’

11) P6589, l21: till → to.

Ans. We have corrected the word from ‘till’ to ‘to’ (P13, l301).

12) Section 3.2: Some of this section could go to the introduction.

Ans. We appreciate the suggestion. We have moved some of the discussion to the Introduction (P5, l84-l91).

‘Over central India (CI, 74.5–86.5° E; 16.5–26.5° N), daily heavy and very heavy rainfall events during the monsoon season (June–September, JJAS) have shown significant increasing trend during 1951 to 2000, whereas moderate rainfall events have shown significant decreasing trend (Goswami et al., 2006; Rajeevan et al. 2008; Pai and Sridhar, 2015). Above studies proposed that significant warming of sea surface temperatures (SSTs) over the equatorial Indian Ocean in recent decades could be the plausible reason

for increase in precipitation extremes, however the mechanism for changes in moderate rainfall events remained unexplored.'

Furthermore, we have made modifications to the text in section 3.2 (P13, 1305-1311) based on the additional analysis using APHRODITE data.

'After counting daily rainfall at each grid point over CI as an event during JJAS from 1951 until 2005 and fitting a linear trend, we find that the number of moderate rainfall events in these 55 years have significantly decreased by about 640, (which is about 3% of the initial value in 1951), (Fig. 2a). Associated with that, total rainfall in the moderate category has also decreased during JJAS (Fig. 2b). The number of extreme rainfall events over CI has significantly increased by 8 (almost double the value in 1951) between 1951 and 2005 (figure not shown).'

13) P6590, 17-8: "On the contrary ...", this is a strange sentence.

Ans. Earlier studies by Goswami et al. (2006) and Rajeevan et al. (2008) have analyzed changes in both moderate and extreme rainfall events over CI. Hence, for information, we have included that sentence and modified it in the revised manuscript (P13, 1309-1311).

'The number of extreme rainfall events over CI has significantly increased by 8 (almost double the value in 1951) between 1951 and 2005 (figure not shown).'

However, a figure has not been included as investigating the reason for changes in extreme rainfall events is out of the scope of this study.

14) P6590, 126: "Daily maximum... day.", Remove this sentence or replace with something more informative.

Ans. We thank the reviewer for this suggestion. However, we would like to state that the sentence is meant to give an estimate about the change in observed mean and extreme surface temperature that has taken place during the mentioned period. We have now modified that sentence in the revised manuscript (P14, 1326-1327).

'We have also analyzed the maximum temperature attained during the day, that represents the higher extreme.'

15) P6591, 17: Can the statement about the coincidence of decreasing trend in daily temperature and forest cover be substantiated with a spatial correlation?

Ans. This is a very good question. Unfortunately, we do not have data on forest cover or crops at high temporal resolution (yearly) that can be used to analyze the spatial correlation.

16) P6592, 125: This rain-shadow area relates strongly to the orography in the model, so please include details on the orography.

Ans. We appreciate this suggestion. The GTOPO30 topography data at 30 arc seconds (approximately 1 km) resolution courtesy of the U.S.G.S. (United States Geological Survey) EROS Data Center and obtained from the following source <http://users.ictp.it/~pubregcm/RegCM4/globedat.htm>, has been used in this study. We have now included details about orography (P10, 1215-1217). A figure based on the orography used in the model is also included as panel (f) of Figure 4.

17) P6593, 19: "On closer examination...", this wording could be a bit more formal.

Ans. We have corrected that wording (P16, 1386-1388).

‘It is evident that the RCM reproduces the regions ...’

18) P6599, 19: warmer → higher.

Ans. We thank the reviewer for suggesting this correction (P21, 1544).

19) P6599, 110: "relatively less than the observed increased", this is an unclear statement, please be more specific.

Ans. We thank the reviewer for this correction. That statement has been deleted in the revised manuscript (P21, 1544-545). The revised statement is:

‘Daily 2m mean air temperature during JJAS in PLC is higher than HLC by a maximum of 0.3 °C over CI and parts of south (Fig. 8a).’

20) P6604, 14: Difference → The difference.

Ans. The correction has been incorporated (P26, 1676).

21) P6621, caption: switch crop and forest to have them in the same order as the figures are.

Ans. We appreciate this correction. Please see the corrected caption in P43.

‘Figure 1. PFT distribution of forest and crop cover (in %) used as fixed lower boundary condition in the model experiments.’

22) P6624, fig 4: Could the shading be made with the same scale, that would be clearer. Moreover, the APHRODITE and IMD data do not seem to differ very much, so why not use only APHRODITE as that one is freely available.

Ans. We appreciate the reviewer’s suggestion. The intent behind keeping the scale in the lower panel different is to bring out the high resolution spatial features of daily seasonal mean rainfall over land, which is lesser than that simulated over the ocean. However, we

have now used slightly different shading for the upper and lower panels to show the contrast.

We admit that only the APHRODITE rainfall data could be used and have agreed to the suggestion. Furthermore, we have included a plot based on the orography used in the model simulations to depict the rain-shadow region (P46).

23) P6599, l24: 925 the hPa level → the 925 hPa level (or at 925 hPa?).

Ans. We have corrected this error (P22, l558).

‘However, we also noted an increase in temperature at the 925 hPa level (figure not shown), implying that the surface warming extends further up to the depth of the boundary layer.’

24) In some places in the text, words are capitalized that should not be, some examples:

Ans. We appreciate the reviewer’s efforts in identifying these mistakes in detail.

- P6601, l22: Bowen Ratio.

Ans. Correction has been incorporated (P24, l110-611).

- P6601, l24: Tropics (not sure whether this should be capitalized)

Ans. The said portion has been deleted in the revised manuscript.

III) Reply to comments by anonymous Reviewer #2 on the paper “**Investigating the impact of land-use land-cover change on Indian summer monsoon daily rainfall and temperature during 1951–2005 using a regional climate model**” by S. Halder et al.

In this study, the authors investigate the impact of land use land cover change over Indian region on the change in the moderate rainfall events and surface temperature using a regional climate model. Four sets of experiments are performed, with pre and present LULC, and with/without the impact of SST trend. In questions posed in the manuscript is important in terms of understanding the variability in Monsoon and is adequate to address through a small study like this. The results are also positive.

However, there are some limitations in the experimental designs those need to be addressed before this can be accepted for publication.

We are thankful to the referee for his comments and encouragement, and also for addressing certain limitations. Our answers to the referee’s comments may be found below.

1. This study uses a regional model, where the lateral boundary condition is provided from reanalysis data, from 1982-2008. The surface boundary condition (over ocean) in the PLC and HLC experiments are also from Reynolds SST during the same period. This makes the model simulations transient in time and any change such as LULC will need time to get into equilibrium. Ideally, such a study need to be done in a global modelling framework with fully coupled components. Or, even to avoid a transient simulation, the boundary conditions can be repeated (e.g., climatology) for several years to get ensemble simulations at equilibrium.

Ans. We thank the reviewer for the queries. Our replies to the comments are as follows.

1) Our objective is to analyze changes in the climatological mean of the number of moderate rainfall events over CI and intensity of rainfall in that category, between PLC and HLC experiments. The lateral boundary forcing and prescribed SSTs in our experiments are transient in nature. As each year of forcing is different from the other, long-term mean (in our case, it is 27 years) is expected to be closer to the reality.

2) Use of climatological boundary conditions is a possibility. However, in that case the model will have problem in properly capturing the synoptic and intraseasonal rainfall variability that contribute to the seasonal mean rainfall of Indian summer monsoon significantly.

3) Repetition of a single year (such as ENSO/non-ENSO year) of boundary conditions is not a suitable option as that may lead to biased response of the model climate to LULCC. These aspects are now mentioned in the revised manuscript (P11, 1249-1264).

These justifications may also be found in the text as:

‘Our objective is to analyze changes in the climatological mean of the number of moderate rainfall events over CI and intensity of rainfall in that category, between PLC and HLC experiments. The lateral boundary forcing and prescribed SST in our experiments are transient in nature and impose the global warming signal on the model climate. As each year of forcing is different from the other, long-term mean is expected to be closer to the reality. However, use of climatological boundary conditions is not an option, as in that case the model will have problem in properly capturing the synoptic and intraseasonal rainfall variability that contribute to the seasonal mean rainfall significantly. Similarly, a single year (ENSO/non-ENSO year) of boundary condition cannot be repeated as that may lead to biased response of the model climate to LULCC. As time varying lateral boundary conditions also impose the effect of variations in remote SSTs, such as that of the Pacific Ocean on the model, partial remote influence on the nature of response due to regional LULCC is possible. Although our RCM simulations are not time-slice experiments in the true sense, the statistics of their difference are expected to reveal the effect of LULCC and associated regional land–atmosphere feedbacks on daily temperature and rainfall variability in a climatological sense.’

Use of simulations with a coupled global model is another option. However, that would also require large computational power. We have proposed that method as an option for possible future experiments in the Conclusions section (P30, 1792-794).

‘Investigation of the impact of LULCC in a high-resolution coupled global climate model where the land cover changes with time or dynamic vegetation is used, would make another interesting study.’

2. The prescription of the boundary conditions virtually imposes the remote impact over this region to the present climate scenario (1982-2008). Therefore, the impact seen between HLC and PLC simulations are really only local impact, which might get enhanced or even eliminated when remote feedback is present.

Ans. The climatology of long-term simulations made with a RCM is determined by a dynamic equilibrium between the time-varying lateral boundary forcing as well as internal model dynamics (Giorgi and Mearns, 1999). The respective model climates simulated in the PLC and HLC experiments are a result of interactions between the prescribed LULC, time varying lateral boundary conditions that impose the remote impact over this region and also internal variability of the model. Hence, partial remote influence on the response of the regional LULCC is a possibility. However, this study tries to point out that for a given large scale condition, what would be the effects of the LULC change on the rainfall events.

3. In the detrended SST experiments, SST trend is removed only over Indian Ocean (the domain of the model). Therefore, these experiments do not include the impact of change in SST over other regions like the Pacific or Atlantic Oceans, where multidecadal oscillations in SST are observed.

Ans. The Indian Ocean has shown significant warming trend in the past few decades and studies suggest that SSTs over that ocean can affect rainfall variability over the Indian region significantly (Roxy et al., 2015). Therefore, the Indian Ocean SST warming trend is removed in our additional set of experiments to further support our hypothesis about the impact of LULCC on changes rainfall and temperature variability. However, the impact of the changes in SST over other regions like the Pacific or Atlantic Oceans is imposed over the Indian monsoon domain through the imposed lateral boundary conditions (NCEP/NCAR reanalysis). This aspect is now mentioned in the revised manuscript (P11, 1258-1261).

‘As time varying lateral boundary conditions also impose the effect of variations in remote SSTs, such as that of the Pacific Ocean on the model, partial remote influence on the nature of response due to regional LULCC is possible.’

It is only ensured that the two-way feedback between model generated climate and climate in the remote region is not allowed. Therefore, our set of experiments does include the impact of changes in remote SST in the monsoon domain.

References

Giorgi, F. and Mearns, L. O.: Introduction to special section: Regional climate modeling revisited, *J. Geophys. Res.*, 104, D6, 6335–6352, 1999.

Roxy, K. M., Kapoor, R., Terray, P., Murtugudde, R., Ashok, K., and Goswami, B. N.: Drying of Indian subcontinent by rapid Indian Ocean warming and a weakening land-sea thermal gradient, *Nat. Commun.*, 6, doi:10.1038/ncomms8423, 2015.

IV) Reply to interactive comment by Dr. R. Pielke Sr. on “**Investigating the impact of land-use land-cover change on Indian summer monsoon daily rainfall and temperature during 1951–2005 using a regional climate model**” by S. Halder et al.

I found this an informative and important paper that further documents the role of land use change as a first order climate forcing. I only have recommendations for papers the authors might like to add to their citations:

We would like to thank Dr. Roger Pielke Sr. for the encouraging comments. The suggested papers have been included in the revised manuscript.

1) Roy, S.S., R. Mahmood, D. Niyogi, M. Lei, S.A. Foster, K.G. Hubbard, E. Douglas, and R.A. Pielke Sr., 2007: Impacts of the agricultural Green Revolution - induced land use changes on air temperatures in India. J. Geophys. Res. - Special Issue, 112, D21108, doi:10.1029/2007JD008834.

Ans. We have included the reference in P5, line 106.

2) Douglas, E., A. Beltrán-Przekurat, D. Niyogi, R.A. Pielke, Sr., and C. J. Vörösmarty, 2009: The impact of agricultural intensification and irrigation on land-atmosphere interactions and Indian monsoon precipitation – A mesoscale modeling perspective. Global Planetary Change, 67, 117–128, doi:10.1016/j.gloplacha.2008.12.007.

Ans. We have included the reference in P6, line 113.

3) Niyogi, D., H.-I. Chang, F. Chen, L. Gu, A. Kumar, S. Menon, and R.A. Pielke Sr., 2007: Potential impacts of aerosol-land-atmosphere interactions on the Indian monsoonal rainfall characteristics. Natural Hazards, Special Issue on Monsoons, Invited Contribution, DOI 10.1007/s11069-006-9085-y.

Ans. We have included the reference in P6, line 113.

4) Lei, M., D. Niyogi, C. Kishtawal, R. Pielke Sr., A. Beltrán-Przekurat, T. Nobis, and S. Vaidya, 2008: Effect of explicit urban land surface representation on the simulation of the 26 July 2005 heavy rain event over Mumbai, India. Atmos. Chem. Phys. Discussions, 8, 8773–8816.

Ans. We have included the reference in P6, line 114-115.

5) Kishtawal, C.M., D. Niyogi, M. Tewari, R.A. Pielke Sr., and J. Marshall Shepherd, 2009: Urbanization signature in the observed heavy rainfall climatology over India. Int. J. Climatol., doi:10.1002/joc.2044.

Ans. We have included the reference in P6, line 115.

V) The marked-up manuscript version has been created separately using LATEX, whereas the responses to the reviewer's comments have been prepared in MS Word. Hence, the marked-up manuscript 'markup_changes_hess-2015-173-discussions.pdf' file is attached separately along with the e-mail to the Editor.

Investigating the impact of land-use land-cover change on Indian summer monsoon daily rainfall and temperature during 1951–2005 using a regional climate model

S. Halder¹, S. K. Saha², P. A. Dirmeyer¹, T. N. Chase³, and B. N. Goswami⁴

¹Center for Ocean–Land–Atmosphere Studies, George Mason University, Fairfax 22030, VA, USA

²Indian Institute of Tropical Meteorology, Dr. Homi Bhabha Road, Pashan, Pune-411008, India

³Cooperative Institute for Research in Environmental Sciences (CIRES), Boulder, CO 80309, USA

⁴Indian Institute of Science Education and Research, Dr. Homi Bhabha Road, Pashan,
Pune-411008, India

Correspondence to: S. Halder (shalder3@gmu.edu)

Abstract

Daily moderate rainfall events, ~~that~~ which constitute a major portion of seasonal summer monsoon rainfall over central India, have decreased significantly during the period 1951 ~~till~~ through 2005. ~~Mean~~ On the other hand, mean and extreme near surface daily temperature during the monsoon season have ~~also~~ increased by a maximum of 1–1.5 °C. Using simulations made with a high-resolution regional climate model (RegCM4) ~~with~~ and prescribed vegetation cover of years 1950 and 2005, it is demonstrated that part of the ~~above observed~~ changes in moderate rainfall events and temperature have been caused by ~~land-use land-cover change~~ Land-Use/Land-Cover Change (LULCC) which is mostly anthropogenic. Model simulations show that the increase in seasonal mean and extreme temperature over central India coincides with the region of ~~decreased (increased) forest (crop) cover~~ The decrease in forest and increase in crop cover. Our results also show that ~~land-use land-cover~~ LULCC alone causes warming in the extremes of daily mean and maximum temperatures by maximum of 1–1.2 °C, that is comparable with the observed increasing trend in the extremes. Decrease ~~(increase) in forest (crop) cover~~ in forest cover and simultaneous increase in crops not only reduces the evapotranspiration over land and large-scale convective instability, ~~apart from decreasing the moisture convergence but also contributes toward decrease in moisture convergence through reduced surface roughness~~. These factors act together ~~not only in reducing~~ in reducing significantly the moderate rainfall events ~~over central India but also~~ and the amount of rainfall in that category ~~, significantly. This is the most interesting result of this study.~~ over central India. Additionally, the model simulations are repeated by removing the warming trend in sea surface temperatures over the Indian Ocean. As a result, ~~there is~~ enhanced warming at the surface and greater decrease in moderate rainfall events over central India compared to the earlier set of simulations are noticed. Results from ~~the~~ these additional experiments corroborate our initial findings and confirm the contribution of ~~land-use land-cover change on~~ LULCC in the decrease in moderate rainfall events and increase in daily mean and extreme temperature ~~and decrease in moderate rainfall events. This study not only~~ over

India. Therefore, this study demonstrates the important implications of LULCC over India ;
but also shows the necessity for inclusion of projected anthropogenic changes in land-use
land-cover in future climate change scenarios for developing better adaptation and mitigation
strategies during the monsoon season. Although, the regional climate model helps in better
resolving land-atmosphere feedbacks over the Indian region, the inferences do depend
on the fidelity of the model in capturing the features of Indian monsoon realistically. It
is proposed that similar studies using a suite of climate models will further enrich our
understanding about the role of LULCC in Indian monsoon climate.

1 Introduction

Observational evidences show that globally averaged annual mean surface temperature
has increased by about 0.85 °C between 1880 and 2012, with rapid warming in the recent
past decades (about 0.72 °C after 1951) (IPCC AR5, Stocker et al., 2014). The IPCC AR5
also summarizes that the number of cold (warm) days and nights have also decreased
(increased) globally, with increase in the frequency of heat waves over large parts of Europe,
Asia and Australia. Apart from surface temperature, There has also been an increase in
extreme (heavy) precipitation events have also increased over most of the global land areas
(Alexander et al., 2006; Stocker et al., 2014). The According to Allen and Ingram (2002),
the increase in mean precipitation is expected to be much less than the extremes as it is
constrained by the net rate of cooling of the troposphere, which, in turn also depends on its
temperature and presence of Greenhouse Gases (GHGs) and aerosols (Allen and Ingram;
2002). The IPCC AR5 summarizes that there is medium confidence on the contribution
of human influence on large-scale changes in land precipitation. Furthermore, On the
contrary, Seneviratne et al. (2012) argues opine that there is no general relationship
between changes in total and extreme precipitation. However, It is intriguing to note that
seasonal and regional or local changes in these extremes extreme weather events can be
of different magnitude and sign than global changes due to complex regional feedbacks
associated with the GHGs, clouds, aerosols and other anthropogenic activities - such as

Land-Use/Land-Cover Change (LULCC). For example, Haerter and Berg (2009) argue that changes in humidity, atmospheric stability, wind direction etc. can strongly influence the local temperature variability. Observational However, due to observational uncertainty, challenges in modeling and natural variability affect proper detection and attribution of the changes regional climate changes often becomes difficult. Therefore, quantification of the changes in regional climate and as well as proper attribution are both very important for the policy makers for devising better adaptation and mitigation strategies.

~~Studies on trends in extreme temperature and precipitation events over the Indian sub-continent have focused on different regions and periods. Kothawale et al. (2010; 2005) found an overall increase of about 0.5C (0.71C) in the annual mean (maximum) temperature over seven homogeneous regions of India in the last century. They also reported an accelerated warming trend of 0.22 in the mean annual temperature in the recent decades (1971–2003). Jaswal et al. (2015) using station observations reported a significant increase in high temperature in summer over India during the period 1969–2013. Pai et al. (2013) also reported a significant increase in occurrence of heat waves in summer during 1961–2010. Sheikh et al. (2014), using daily station data found a general increase in warm temperature (precipitation) extremes during 1971–2000 (1961–2000), with enhanced warming over the desert of Thar. They associated those changes with the effect of increasing aerosols. Over central India (CI, 74.5–86.5E; 16.5–26.5N), heavy and very heavy (moderate) rainfall events during the monsoon season (June–September, JJAS) have been reported to show a significant increasing (decreasing) trend from 1951 to 2000 (Goswami et al., 2006). Rajeevan et al. (2008) and Pai and Sridhar (2015) supported that result and reported a similar but relatively weaker trend in heavy and moderate rainfall events over that region from 1901 to 2000. Singh et al. (2014) reported a significant decrease in monsoon precipitation over that core (CI) region of India from 1951 till 2011, along with a significant increase in frequency of dry spells and intensity of wet spells and decrease in intensity of dry spells. Niranjana Kumar et al. (2013) found an increase in the intensity of droughts during 1901 till 2010 over India. Panda et al. (2014) reported a significant decrease in wet days and moderate and total rainfall during the summer~~

monsoon (1971–2005) over the northeast, central and southwest of India. Both studies attributed their observed trends with evolving SST anomalies over the Pacific and Indian Oceans. Although Goswami et al. (2006) and Rajeevan et al. (2008) pointed at warming of the Indian Ocean sea surface temperature (SST) to be the governing factor for increase in precipitation extremes, the probable reason for decrease in moderate rainfall events was left unaddressed. Observed changes in daily high temperature and moderate/extreme rainfall events over the Indian region may be attributed to both natural variability and anthropogenic activity. Extreme high or low temperature phases are not only harmful for crop growth and yield but heat/cold waves and heavy rainfall events also cause fatality of living beings. Light to moderate rainfall events constitute a major portion of the seasonal rainfall over India and are an important source for replenishment of ground water. The aim of this study is to understand the processes responsible for observed changes in moderate rainfall events and daily mean and maximum temperature over India during the monsoon season.

1.1 Role of LULCC in climate

1.2 ~~Role of land use/land cover change in climate~~

~~Land use/land cover change (LULCC)~~ LULCC is an important driver of climate change at local, regional and possibly, ~~also on~~ global scale (Snyder, 2010) and ~~on~~ time scales inter-decadal and beyond (Pitman et al., 2012; Mahmood et al., 2014; Dirmeyer et al., 2010; Solomon et al., 2007). ~~LULCC affects the surface albedo, fluxes of radiation and that of momentum, heat, water vapor, carbon dioxide (CO₂) and other trace gases, aerosols and dust, and turbulence in the boundary layer (Pielke et al., 2011; Mahmood et al., 2014).~~ However, the climate effects of deforestation and agricultural intensification vary regionally and also depend on the seasons, making resulting land-atmosphere interactions complex. In the last 300 years (1700–2000), about 42–68 % of the global land surface has been affected due to land use practices (Hurtt et al., 2006). ~~There has been an increase in global~~ ; Pielke et al. 2011), resulting in an increase in cropland (Ramankutty et al., 1999, 2008) and pastures (Goldewijk et al., 2001). ~~For a concise~~

summary of the evolution of LULCC across centuries, refer Fig. 1 of the paper by Pielke et al. (2011) that is based on the Land Use Harmonization data (). Cropland areas and pastures in general, decrease the surface temperature on account of their higher albedo than forests. Likewise, irrigated agricultural landscapes also contribute to surface cooling and rainfall (Pielke et al., 2011; Mahmood et al., 2014). However, the effects of deforestation and agricultural intensification vary regionally and also depend on the seasons, making resulting land-atmosphere interactions complex. Robust results show Robust results have shown that albedo changes due to increase in croplands and pastures leading to decrease in surface temperature tend to dominate over the mid-latitudes, whereas decrease in evapotranspiration (ET) and roughness length, roughness length and cloudiness play a primary role in increasing surface warming in the Tropics tropics (Garratt, 1993; Bounoua et al. 2002; Fedora et al. 2005; Sampaio et al., 2007; Davin and De Noblet-Ducoudré, 2010; Pitman et al., 2012). The above conclusions are supported in studies by Lawrence and Chase (2010), Bounoua et al. (2002) and Feddema; Pitman et al. (2005). Deforestation and increases in crop cover reduces the surface roughness and also decreases the moisture convergence and (2012). Furthermore, deforestation can affect moisture convergence, atmospheric stability and changes in rainfall (Sud et al., 1998). Spatial heterogeneity in vegetation has been shown to affect convective rainfall through changes in surface fluxes of heat and moisture and atmospheric stability (Pielke et al., 2001). Lawrence and Chase (2010) further demonstrated that alteration of potential vegetation to present day land cover state can lead to regional warming as well as reduction in precipitation. In an interesting study, Avila et al. (2012) demonstrated that Studies also suggest that changes in the temperature extremes due to LULCC could be of comparable magnitude but of similar or opposite sign as due to increase in CO₂, depending on the region. Pitman (Avila et al. (2012) supported those conclusions using a multi-model ensemble study, and also extended their analysis for rainfall extremes. Since; Pitman et al. 2012). As it is difficult to segregate the impact of LULCC on temperature and precipitation extremes when analyzed in a globally averaged sense (Pielke et al., 2011; Pitman et al., 2012), one needs to carefully design carefully designed sensitivity studies

with climate models ~~and focus focussing~~ on specific regions ~~to better understand its role~~ are required.

1.2 LULCC Changes in the context of temperature, rainfall and LULC over India

Over India, industrialization and urbanization has grown immensely from the middle of the 20th century, apart from the Green revolution. This has lead to widespread deforestation and changes in land-use practices towards agriculture. A recent study over India by Tian Observed changes in daily temperature and rainfall extreme events over the Indian region may be attributed to both natural variability and anthropogenic activity. There has been an increase of about 0.5°C in the annual mean and 0.71°C in the maximum temperature over India in the last century, but increased warming in the recent decades (1971–2003) (Kothawale et al. 2010, 2005). Pai et al. 2013 have noted a significant increase in the occurrence of heat waves in summer (1961–2010), whereas Jaswal et al. (2014) using high-resolution remote sensing datasets from the Resourcesat-1 satellite and historical archives at district and state level show loss (gain) of about 26 (48.1) million ha of forests (crops) over the period 1880–2010. Greater cropland expansion and urbanization occurred during 1950–1980 but mostly over central, eastern and northwest India and eastern and western parts of peninsular India (refer Fig. 4 in their paper). Such regional changes in LULC could have caused changes in the Indian climate through their impact on the surface moisture and heat budget, and may have amplified or compensated other potential changes due to increased GHGs, aerosols, large-scale circulation changes or natural variability. Therefore, it is important to quantify whether the decrease in (2015) have shown an increase in temperature extremes (1969–2013). Observed changes in temperature in recent decades have been associated with the effect of increasing aerosols (Pai et al. 2013; Sheikh et al. 2014), as reported earlier by Krishnan and Ramanathan, (2002). Over central India ($74.5\text{--}86.5^{\circ}\text{E}$; $16.5\text{--}26.5^{\circ}\text{N}$), daily heavy and very heavy rainfall events during the monsoon season (June–September, JJAS) have shown significant increasing trend during 1951 to 2000, whereas moderate rainfall events over CI and how much of the observed

changes in daily mean and extreme surface temperature over India in the later half of the 20th century is attributed to LULCC?

As mentioned earlier, the increasing trend in heavy rainfall events over CI has been potentially associated with a significant increasing trend in have shown significant decreasing trend (Goswami et al., 2006; Rajeevan et al. 2008; Pai and Sridhar, 2015). Above studies proposed that significant warming of sea surface temperatures (SSTs) over the equatorial Indian Ocean in recent decades could be the plausible reason for increase in precipitation extremes, however the mechanism for changes in moderate rainfall events remained unexplored. There has also been an increase in the intensity of droughts over India during 1901 to 2010 (Niranjan Kumar et al. 2013) and a significant decrease in wet days and moderate and total rainfall during the summer monsoon (1971–2005) (Panda et al. 2014). Both studies have associated the observed changes to variations in SST over the tropical Indian Ocean (Goswami Indian and Pacific Ocean. Furthermore, rapid warming of the Indian Ocean compared to land has been shown to have significantly affected the land-sea thermal contrast and decreased summer rainfall during 1901 through 2012 (Roxy et al., 2006; Rajeevan 2015). Apart from regional changes in the concentration of anthropogenic aerosols and Greenhouse Gases (GHGs) or Indian Ocean SSTs, industrialization and urbanization over India have lead to widespread deforestation and changes in land-use practices in recent decades. According to Tian et al., 2008) that has been also reported in Roxy et al. (2014). 2014), there has been loss of about 26 million ha of forests and gain of about 48.1 million ha of crops in India during 1880 through 2010 (cf. Fig. 4 in their paper). Therefore, the impact of LULCC alone on changes in the distribution of moderate rainfall events or surface temperature extremes during 1951 through 2000 needs to be investigated.

Studies have attempted to understand the mechanisms through which LULCC affects the regional climate over India. For example, Sen Roy et al. (2007, 2011) found demonstrated that irrigation can lead to a significant decrease (increase) in pre-monsoon (March–May) surface temperature (precipitation) over India due to irrigation activity. Lee et al. (2009) showed that pre-monsoon irrigation activity could affect the early part of the. Irrigation

activity has also been shown to affect the Indian Summer Monsoon Rainfall (ISMR) through changes in land-ocean temperature contrast ~~–(Lee et al. 2009) or land-atmosphere~~ feedbacks (Niyogi et al. (2010) ~~found an overall decrease in rainfall during monsoon season over the northwest of India due to irrigation and land-use change. Similarly, ; Tuinenburg et al. (2011) used model simulations to conclude that large-scale irrigation might increase local precipitation over eastern and southern India through land-atmosphere feedbacks. Nayak and Mandal (2012) used model studies to attribute increased warming over western India (1973–2009) to LULCC.~~ There have been ~~similar other studies as well like several~~ other studies addressing the effects of LULCC over the Indian region (Lohar and Pal (1995), Douglas et al. (2006), (2009), Niyogi et al. (2007), Saeed et al. (2009), Dutta et al. (2009): ~~Kishtawal, Nayak and Mandal (2012)). Apart from them, Lei et al. (2010) reported an influence of growing urbanization on the increasing trend of extreme rainfall events over India (2008), Kishtawal et al. (2009) and Ali et al. (2014) ,on the contrary, found have~~ explored the impact of growing urbanization in India and large-scale climate variability and not urbanization in India responsible for observed in the changes in extreme rainfall events. ~~LULCC not only involves~~ Interesting time slice experiments made with a global model have shown that an increase in ~~irrigated lands but also deforestation, afforestation, conversion to bare, pasture or cropland and urbanization as well. Global model simulations by crop and pasture land lead to a decrease in seasonal rainfall over India during the pre-industrial period (years 1700 through 1850) when the impact of anthropogenic activity or natural climate variations were minimal (Takata et al. (2009) using historical reconstructed LULC (years 1700 and 1850 during the pre-industrial period) showed a . Krishnan et al. (2015) made several experiments with a high resolution global atmospheric model and concluded that a multitude of factors such as aerosols, land-use change, Indian Ocean warming as well as GHGs have together contributed to the observed~~ weakening of the Indian summer monsoon circulation and a decrease in seasonal rainfall due to an increase in crop and pasture land. Earlier studies have not focused on the changes in the frequency and intensity south-Asian monsoon and changes in frequency distribution of daily rainfall or temperature extremes. Likewise, the mechanism for decrease in light and moderate rainfall events is

~~not understood and has also not been investigated.~~ events during the later half of the 20th century. However, the impact of LULCC as a lone forcing component on the Indian summer monsoon has not been quantified. It is also plausible that feedbacks due to variations in remote SSTs and snow cover may have modulated the local impacts due to LULCC. In this study, we hypothesize and demonstrate that LULCC ~~over India has contributed partly~~ has partly contributed to the observed decrease in moderate rainfall events over CI during the monsoon season from 1951 ~~till 2005. Apart from that, we also show that LULCC has contributed to the significant through 2005, apart from the~~ increasing trend in surface daily mean and maximum temperatures ~~during summer season. For that purpose, we have used high resolution improved and up-to-date.~~ We have conducted experiments with a high-resolution regional climate model (RCM) RegCM4.0 and accurate LULC data over the Indian region -

~~High-resolution regional climate model (RCM) simulations with RegCM4 have to prove~~ our hypothesis. No added external forcing in terms of aerosols or GHG concentration is ~~used in our experiments. Furthermore, additional experiments by removing the positive trend in Indian Ocean SSTs have also been made to support our hypothesis.~~ isolate the impact of LULCC.

RCMs have shown ~~improvements over global models~~ much improvement over global climate models (GCMs) in terms of representation of spatio-temporal details of climate (Giorgi and Mearns, (1999); Laprise et al., 2008; Leduc and Laprise, 2009) and ~~in~~ dynamical downscaling ability (Xue et al., 2014) ~~. RCMs and~~ add value in simulation of topography induced phenomena and extremes of short spatio-temporal character (Feser et al., 2011; Feser and Barcikowska, 2012; Shkol'nik et al., 2012). ~~They have also shown improvement in representation of large-scale monsoon features over East Asia (Gao Saha et al. (2011, 2012) . The main advantage of a RCM in the context and Halder et al. (2015) have made experiments with the RCM RegCM3 and RegCM4, respectively to better resolve regional land-atmosphere feedback processes and demonstrate their role in the mean and variability of the Indian monsoon is that it can isolate external forcing generated in remote areas by the local feedback processes within the monsoon region or else, they may~~

255 ~~interact with the~~ summer monsoon. When time-dependent lateral boundary conditions are
used as forcing for a RCM in one-way mode, feedback from the model-simulated climate
to the global climate is not allowed. Such interactions between large-scale forcing such
as El Niño–Southern Oscillation (ENSO) that is external to the Indian monsoon region
260 ~~and~~ internal monsoon dynamics ~~and produce more variability~~ may lead to more variability
than due to local feedback processes. Therefore, our methodology helps in segregation
of the impact of regional LULCC on the Indian summer monsoon and its changes. ~~In that~~
~~way, a RCM better resolves regional land–atmosphere feedback processes than a GCM.~~
However, one of the ~~disadvantages is that the~~ drawbacks of regional climate modeling is
265 ~~that~~ lateral boundary conditions ~~may not be perfect. Studies by Saha et al. (2011, 2012)~~
~~have demonstrated the capability of the RCM RegCM3 (previous version of RegCM4) in~~
~~simulating the mean and interannual variability of the Indian summer monsoon.~~

~~This are not perfect. Our~~ paper is organized ~~as follows~~in the following way. Observed
data, the RCM and the design of experiments are described in detail Sect. 2. Method of
preparation of the LULC data used for model experiments is described in the Supplement.
270 ~~Herein, a brief description about preparation of the LULC data that is used in model~~
~~simulations is also presented.~~ The observed changes in near surface temperature and
rainfall and LULC over the Indian subcontinent in the last 55 years are discussed in Sect. 3.
Results from model experiments pertaining to changes in rainfall and surface temperature
are discussed in detail in Sect. 4. Discussions are presented in Sect. 5 and the conclusions
275 drawn are summarized in Sect. 6.

2 Data and methods

2.1 Observed data

Daily 2 m mean and maximum temperature data (1969–2005, at $1^\circ \times 1^\circ$ resolution) from
the India Meteorological Department (IMD) (Srivastava et al., 2009) are used for analysis
280 of trends and validation of the model simulations. In addition to that, we have also

used daily 2 m mean temperature data (at $2.0^{\circ} \times 2.0^{\circ}$ resolution) of the twentieth century reanalysis (20CR) project (Compo et al., 2011) that is available for a longer period (1951–2005). For analysis of trends in daily rainfall events and their intensities in different categories daily gridded data ~~(at $1^{\circ} \times 1^{\circ}$ resolution) for 57 for 55 years (1951–2007) from IMD (Rajeevan 1951–2005) from the Asian Precipitation Highly—Resolved Observational Data Integration Towards Evaluation of Water Resources (APHRODITE Water Resources, Yatagai et al., 2006)–2012) at $0.5^{\circ} \times 0.5^{\circ}$ resolution~~ is used. ~~These observed data available from IMD are one of the most reliable data sets at high resolution (–).~~ Monthly rainfall from GPCP version 2.2 (Adler et al., 2003) for the period ~~1979–1982~~ to 2008 (at $2.5^{\circ} \times 2.5^{\circ}$ resolution) is also used for validation of model simulated rainfall. Apart from that, the National Centers for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR) monthly reanalysis winds, temperature and specific humidity (Kalnay et al., 1996) for the period ~~1982–2008~~ 1982–2007, at $2.5^{\circ} \times 2.5^{\circ}$ horizontal resolution and multiple pressure levels are used for validation of the model simulated large-scale features during monsoon.

2.2 LULC data

Annual harmonized LULCC data (LUHa.v1) from the University of New Hampshire (UNH, <http://luh.unh.edu>) at $0.5^{\circ} \times 0.5^{\circ}$ horizontal resolution (Hurt et al., 2006) that comprises of crop, pasture and primary and secondary vegetation types has been used. This data has been transformed in the form of 17 Plant Functional Type (PFT) mosaics for use as a time invariant lower boundary condition for ~~RCM simulations coupled with the Community Land Model (CLM) land surface model in this study~~ simulations with the RCM. The four UNH vegetation categories are converted into different ~~GLM~~ PFT distributions based on present day and potential vegetation GLM for Community Land Model (CLM) land surface parameters. We have used the resulting PFT distributions and associated vegetation dependent parameters such as ~~LAI, Stem Area Index~~ leaf area index (LAI), stem srea index (SAI), roughness length etc. for the present day conditions (year 2005) and historical period

(year 1950) for our model simulations. Detailed methodology of preparation of the LULC data is given in the Supplement.

2.3 RegCM4.0 and the CLM3.5 land surface model

The RCM ~~RegCM-version4~~RegCM4.0 (Elguindi et al., 2010; Giorgi et al., 2012) is used for this study. The dynamical core of RegCM4 is from the NCAR-Pennsylvania State University (PSU) Mesoscale Model version 4 (MM4), which is a compressible, finite difference model with hydrostatic balance and vertical σ -coordinates. The NCAR CCM3 radiation scheme (Kiehl et al., 1996) and a planetary boundary layer scheme based on a non-local diffusion concept (Holtslag et al., 1999) are used for our simulations. We have also used the ~~new~~ parameterization scheme of Zeng et al. (2005) that allows for a realistic representation of the diurnal variation of sea surface skin temperature. Apart from that, the Grell convective parameterization scheme (Grell, 1993) with the Fritsch and Chappell closure (Fritsch and Chappell, 1980) is used. The model configuration comprises of 23 vertical sigma coordinate levels in the atmosphere and a horizontal resolution of $60^\circ \times 60^\circ$ km with Normal Mercator map projection. The model domain extends from $40.2\text{--}116.3^\circ$ E and 10.8° S– 47.7° N with the Indian subcontinent at the center. ~~Recent studies with a similar model configuration as this (Saha et al., 2011, 2012) has demonstrated fidelity of the RCM in simulating the climatological features of Indian summer monsoon.~~ The NCEP/NCAR reanalysis data (Kalnay-Kalyan et al., 1996) at $2.5^\circ \times 2.5^\circ$ horizontal resolution and 6 hourly frequency for the period 1982 to 2008 is used as lateral boundary conditions for the model. Reynolds weekly SST at $1^\circ \times 1^\circ$ horizontal resolution (Reynolds et al., 2002) interpolated to daily values is prescribed over the ocean.

RegCM4.0 is coupled to the Community Land Model (~~GLMv3~~CLM3.5) (Oleson et al., 2008; Stockli et al., 2008) land surface model. There are 10 soil layers of variable depth and up to 5 layers of snow. CLM3.5 uses a nested sub-grid hierarchy of mosaics in the form of glaciers, lakes, wetlands, urban and vegetated land to better represent surface heterogeneity in a grid box. The vegetated land portion of a grid cell may be composed of multiple columns. Furthermore, in each column ~~4~~the four most abundant PFTs out of

possible 17 that include forests, grasses, crops and bare ground co-exist. The fractional areas of the 4 PFTs do not vary with time but their ~~monthly LAI~~ leaf and stem area ~~index (SAI) values~~ indices vary seasonally, which are all interpolated from global datasets at $0.5^\circ \times 0.5^\circ$ horizontal resolution to the model grid. Fluxes are computed at the PFT level and their weighted averages constitute the net upward flux from a column. Several PFT based parameters are also prescribed in the model. The GTOPO30 topography data at 30 arc seconds resolution courtesy of the U.S.G.S. (United States Geological Survey) EROS Data Center has been used in this study. A global soil texture dataset at 5 min spatial resolution from the International Geosphere Biosphere Program (IGBP) (Bonan et al., 2002) is used with varying sand and clay content in each of the 10 layers. Soil color dataset (8 classes) at ~~2.8~~ 2.8×2.8 spatial resolution is from Zeng et al. (2002). CLM3.5 also uses global datasets on canopy top and bottom height (resolution ~~$0.5^\circ \times 0.5^\circ$~~ 0.5×0.5), percentage of glacier (resolution ~~$0.5^\circ \times 0.5^\circ$~~ 0.5×0.5), lake and wetland (resolution ~~$1^\circ \times 1^\circ$~~ 1×1) with corresponding spatial resolution included in brackets (Elguindi et al., 2010). Details about the ~~land surface~~ parameterization schemes in CLM3.5 are also presented in Oleson et al. (2010) and ~~Tawfik and Steiner (2011)~~ Halder et al. (2015).

2.4 Design of experiments and methodology

Two sets of model ~~simulations~~ simulations, each for 27 years are carried out with similar Lateral Boundary Conditions (LBCs) from NCEP/NCAR reanalysis and Reynolds weekly SST prescribed at the lower boundary, but different LULC ~~of for~~ the years 1950 and 2005 as fixed lower boundary condition. The LULC of 1950 and 2005 correspond to different PFT distributions. The RCM is initialized at 00:00 GMT on 1 November 1981 and the simulation is continued up to 18:00 GMT on 31 December 2008. In CLM3.5 coupled to RegCM4, soil moisture is initialized based on ~~climatology~~ climatological values (as in ~~Tawfik and Steiner, 2011~~; Giorgi and Bates, 1989; Halder et al., 2015), in order to reduce model spin-up time for ~~deeper layers~~ the deeper soil layers. Therefore, we have discarded the initial seven months for model spin-up. Soil points are initialized with temperatures of 283 K (Oleson et al., 2010). Hereafter, these simulations with LULC of 2005 and 1950

will be referred as Present Land Cover (PLC) and Historical Land Cover (HLC) experiment, respectively. ~~No added external forcing in terms of aerosols or GHG concentration is used in our RCM experiments. Although, our RCM simulations are not time-slice experiments in the true sense, the statistics of their difference are expected to reveal the effect of LULGC and associated regional land-atmosphere interactions on daily temperature and rainfall variability in a climatological sense. Studies have suggested that changes in surface~~ temperature (Kothawale et al. (2010) and ; Chowdary et al. (2013) ~~have shown surface temperatures over India and the Indian Ocean SSTs to be strongly linked on low-frequency interannual time scales. Apart from that, Goswami et al. (2006) and Rajeevan and extreme rainfall events~~ (Krishnan et al. (2008) ~~speculated that the increasing trend of extreme rainfall events might be associated with the warming trend in SST over 2015) over India are related with variations in~~ the Indian Ocean SSTs. Therefore, in order to isolate the effect of Indian Ocean SSTs on the temperature and rainfall variability over the ~~monsoon Indian~~ region, another ~~two sets pair~~ of model simulation for the same 27 years are carried out using the fixed LULC of years 1950 and 2005, but with de-trended ~~weekly~~ Reynolds SSTs over the Indian Ocean. The ~~de-trended weekly SSTs from November 1981 till December 2008 are interpolated to daily values for model simulation. The~~ LBCs from NCEP/NCAR reanalysis and the initial conditions ~~for land~~ remained exactly same as in the earlier experiments. Henceforth, these RCM simulations will be referred as Present Land Cover de-trended SST experiment (PLCS) and Historical Land Cover de-trended SST experiment (HLCS), respectively. The four experiments are briefly summarized in Table 1.

~~Observed changes in LULC used as fixed boundary condition in the RCM are described in Sect. 3, which also includes discussion on observed changes in surface temperature and daily rainfall distribution during 1951 to 2005. Simulation of the large-scale climatological features of Indian summer monsoon by the RCM in the PLC and PLCS experiments are discussed in~~ Our objective is to analyze changes in the following Sect. 4.1. Differences in simulated rainfall statistics between experiments with past (1950) and present land cover (2005) are discussed in Sect. 4.2. Of particular importance is our discussion on changes in moderate rainfall events (Sect. 4.2.1). climatological mean of the number of

moderate rainfall events over CI and intensity of rainfall in that category, between PLC and HLC experiments. The lateral boundary forcing and prescribed SST in our experiments are transient in nature and impose the global warming signal on the model climate. As each year of forcing is different from the other, long term mean is expected to be closer to the reality. However, use of climatological boundary conditions is not an option, as in that case the model will have problem in properly capturing the synoptic and intraseasonal rainfall variability that contribute to the seasonal mean rainfall significantly. Similarly, a single year (ENSO/non-ENSO year) of boundary condition cannot be repeated as that may lead to biased response of the model climate to LULCC. As time varying lateral boundary conditions also impose the effect of variations in remote SSTs, such as that of the Pacific Ocean on the model, partial remote influence on the nature of response due to regional LULCC is possible. Although our RCM simulations are not time-slice experiments in the true sense, the statistics of their difference are expected to reveal the effect of LULCC and associated regional land–atmosphere feedbacks on daily temperature and rainfall variability in a climatological sense.

Extreme rainfall events are short lived, less frequent but intense, ~~with long return periods~~. They are associated with deep convective activity that is triggered by local instabilities or large-scale moisture convergence and drain out the atmospheric moisture content very fast, thus increasing the atmospheric stability. On the other hand, light and moderate rainfall events are relatively less intense and long-lived and require time for large-scale moisture and instability to build up ~~for them to sustain~~ and be sustained. Thus, due to the smaller spatial scale and random frequency of occurrence of extreme rainfall events, analysis of their trends over stations sparsely spaced or individual grid points is not expected to give a robust or consistent result about their temporal variability. ~~Therefore~~ However, more meaningful information on the statistics of extreme rainfall events can be obtained when analyzed in a spatially aggregated sense (Goswami et al., 2006; Rajeevan et al., 2008). ~~This approach has also been followed by~~ Singh et al. (2014). For our study, the CI domain that is considered homogeneous in terms of the mean and variability of the Indian summer monsoon rainfall (Goswami et al., 2006) is used for the analysis of moderate

and extreme rainfall events. Significance of the results have been tested on the basis of Student's t test. ~~The effect of LULCC on surface air temperature extremes over India in the experiments with realistic prescribed SSTs and the experiments with de-trended SSTs are discussed in Sect. 4.3.~~ For the analysis on temperature extremes in the model we have used data for the period JJAS (instead of JAS used for observation) that will be further discussed in Sect. 4.1. ~~Based on the experiments and further analysis, we propose a physical mechanism responsible for the observed changes in Sect. 4.4.~~

3 Observed changes

3.1 LULCC

Figure 1 shows the distribution of PFTs ~~of all forests and crops (including grasses)~~ in the year 1950 and 2005 used as lower boundary condition in the RCM and also gives an overview of past changes in land cover. The northwest of India, the hilly terrain over CI, western states of Gujarat and Maharashtra, foothills of the Himalayas and northeastern states are mostly dominated by forest cover (Fig. 1a and c). Agriculture or crop cover is mostly concentrated along the northern states of India such as Punjab, Haryana, Delhi, the Gangetic plains, the plains of east and west CI and peninsular India (Fig. 1b and d). Difference between PFT distribution under present climatic condition (year 2005) and historical period (year 1950) show that forest cover is reduced and crop cover is increased in the recent period by about 5–30 % (Fig. 1e and f). Maximum increase in crop fraction is seen largely over CI, peninsular India, north and northwest India and extreme northern part around the plains of river Indus. This increase in crop fraction also matches very well with the changes shown in Fig. 5 in the paper study by Tian et al. (2014) over the period from 1950 to 2010. ~~Observed~~ It is interesting to note that observed surface evaporation has significantly decreased over continental India during the monsoon season from 1971 to 2000 (Jaswal et al., 2008) ~~and which~~

3.2 Rainfall over central India

There is no clear trend in the all India mean summer monsoon rainfall during JJAS from 1951 ~~till to~~ 2000, but extreme and moderate rainfall events have changed over CI significantly. Following, Goswami et al., ~~(2006)~~; Rajeevan et al., ~~(2008)~~. ~~While moderate rainfall events (are defined in this study as $5 > \text{rainfall} \leq 100 \text{ mm day}^{-1}$) show a significant decreasing trend, the heavy (rainfall $\geq 100 \text{ mm day}^{-1}$) and very heavy rainfall (rainfall $> 150 \text{ mm day}^{-1}$) events show a significant increasing trend. As mentioned earlier, increase in heavy and very heavy rainfall events over CI have been potentially associated with a significant increasing trend in SST over the Indian Ocean. However, the physical mechanism for decrease in moderate rainfall events over CI is unknown and is investigated in this study., respectively.~~ After counting daily rainfall at each grid point ($1^\circ \times 1^\circ$) over CI as an event during JJAS from 1951 until 2005 and fitting a linear trend, we find that the number of moderate rainfall events between 1951 and 2005 in these 55 years have significantly decreased by about ~~520~~640, (which is about ~~103~~ % of the initial value in 1951), (Fig. 2a). Associated with that, ~~the~~ total rainfall in the moderate category has also decreased during JJAS (Fig. 2b). ~~On the contrary,~~ The number of extreme rainfall events over CI has significantly increased by ~~about 10~~8 (almost double the value in 1951) between 1951 and 2005 (figure not shown). ~~On the basis of earlier modeling studies that addressed the impact of vegetation cover on rainfall and observed changes in vegetation over India we propose that, LULCC~~ We propose that LULCC during these 55 years might have contributed to the observed decrease in moderate rainfall over CI. ~~We and~~ substantiate our hypothesis using multiple simulations with the RCM RegCM4.

3.3 Surface air temperature

The pre-monsoon season in India (March–April–May) is characterized by days that are hot and dry. The climatological onset date of the southwest monsoon over Kerala (southern tip of India) is 1 June. There is large year-to-year variability in the date of onset and in

many years, onset takes place during the middle of June (Wang et al., 2009). Therefore, to investigate the changes in observed daily mean temperature and its extreme during the monsoon season, trends are calculated using temperature of the months July–September (JAS) only. A warming trend in the JAS mean temperature by $0.2\text{--}0.4\text{ }^{\circ}\text{C decade}^{-1}$ is observed over the northwest, northeast and southern parts of India (Fig. 3a). Similar to the mean, extreme of daily mean temperature in JAS (its 90th percentile) also shows a warming trend, but over a larger region (Fig. 3b). Based on this trend from 1969 ~~till~~ until 2005, it is estimated that the daily mean surface temperature and its extreme in JAS have increased by a maximum amount of about $1.11\text{--}1.48\text{ }^{\circ}\text{C}$. ~~Daily maximum temperature represents~~ We have also analyzed the maximum temperature attained during the day, that represents the higher extreme. Figure 3c shows the trend in JAS averaged daily maximum temperature. It is evident that warming in the daily maximum temperature is of the same order, but is more widespread as compared to the daily mean and includes areas north of CI. Furthermore, the 90th percentile of daily maximum temperature has increased by more than $1.48\text{ }^{\circ}\text{C}$ over north-central India, which is greater than the increase in the mean (Fig. 3d). It may be noted that the spatial pattern of increase in daily temperature over CI is consistent with the area of increase in crop PFTs over CI and northwest (Fig. 1f). Similarly, the decreasing trend in daily temperature over areas south of CI also coincides well with the small increase in forest cover over that region (Fig. 1e). Increased observed temperature over the western coast of peninsular India may have happened due to its region specific mean climate ~~that is~~ predominantly determined by the adjoining Arabian Sea and Indian Ocean.

~~We also analyzed the trend~~ Trends in daily 20CR 2 m mean temperature data and its extreme (90th percentile) during JAS ~~for the period 1950–2005~~. are further analyzed for the extended period 1951–2005. A significantly increasing trend is evident both in the mean and its extreme, over northern India (Fig. 3e), north of CI (Fig. 3f) and southern parts of peninsular India. The maximum increase in daily mean temperature in JAS is about $1.11\text{ }^{\circ}\text{C}$. The pattern of increase in daily maximum temperature is not only more widespread (possibly due to coarser resolution of the ~~model data~~) but its magnitude is also comparable to that seen over the 37-year period (1969–2005). Apart from that, a decrease south of

CI and an increase towards the extreme south is also evident. However, while comparing the trends shown by the above two datasets we note that the model used to generate the 20CR data did not assimilate surface temperature observations. Therefore the resulting trend is also partially model dependent. Observed increasing trend in daily mean surface temperature and its higher extreme may be attributed to forcing of natural (solar, volcanic) or anthropogenic origin (GHGs, aerosols, LULCC etc.) or both. We aim to quantify the contribution to such increase due to LULCC over India.

4 Results from RCM experiments

In the PLC and HLC experiments, we keep the atmospheric and oceanic boundary conditions during 1982 to 2008 ~~similar~~ same but the distribution of PFTs are altered corresponding to years 2005 and 1950 respectively. This experimental set-up is meant to help us understand the statistics of changes in rainfall and temperature due to LULCC.

4.1 Indian summer monsoon features in PLC and PLCS experiment

~~The Indian summer monsoon is a large-scale phenomenon and an important driver of boreal climate in the tropics as well as extra-tropics. The~~ skill of the RCM in capturing the mean spatial distribution of seasonal (JJAS) rainfall and its interannual variability are assessed here. The observed seasonal mean monsoon rainfall in GPCP (Fig. 4a) shows a region of maxima over the Western Ghats, head Bay of Bengal (BoB), hilly terrain of Central India and north-east India. There is also a region of maximum over east equatorial Indian Ocean. In comparison, rainfall in the PLC experiment is overestimated near the Arabian Sea coast and over BoB. Apart from that, a secondary rainfall maximum ~~exists~~ which-that is shifted to the west equatorial Indian Ocean region is also noted in PLC. Although rainfall is also captured over CI and the northeast region, the magnitude ~~seems~~ appears to be underestimated, particularly over western India. Earlier studies have shown that the rainfall bias in ~~RCMs over this~~ this RCM over the ocean is attributed to the lack of coupling with the atmosphere and also to the choice of convective parameterization

schemes (Chow et al., 2006; Ratnam et al., 2009; Saha et al., 2011; [Halder et al., 2015](#)). However, it is interesting to note that compared to an earlier version of the RCM (RegCM3) used for simulation of the Indian summer monsoon with a similar model set-up (Saha et al., 2011, 2012), this positive bias over the west-equatorial Indian Ocean region and western part of BoB is relatively reduced. The ~~rain-shadow area over peninsular India is captured by the RCM. The~~ dashed (solid) lines in Fig. 4b represent the CI (big-India, BI) domain used for our analysis related ~~distribution to the statistics~~ of daily rainfall. Seasonal mean rainfall in [the](#) PLCS experiment follows a similar spatial pattern as in [the](#) PLC and captures the locations of rainfall maxima [very well](#) (Fig. 4c). However, the magnitude is relatively less everywhere compared to ~~PLC~~[the PLC experiment](#). Maximum decrease in seasonal total rainfall over CI between [the](#) PLCS and PLC experiments is about 4 cm (figure not shown). This decrease is possibly ~~due to associated with~~ relatively colder SSTs over the Indian Ocean that leads [to](#) lesser evaporation over ocean and hence moisture ~~convergence over land in the atmosphere~~. These aspects will be discussed further in Sect. 4.2.

Seasonal rainfall over the land part in PLC (Fig. 4e) is ~~compared further compared in detail~~ with that from IMD and APHRODITE data (Fig. ~~4c and d~~). ~~On closer examination it is revealed that the~~ [4d](#)). The representation of orography in the model is depicted in [Fig. 4f](#) which suggests that the surface topography is very well captured by the model. It is evident that the RCM reproduces the regions of rainfall maxima and the spatial pattern ~~reasonably well. very well, particularly over the Western Ghats section over peninsular India, CI, north-east India and the Himalayan foothills. The rain-shadow area east of the Western Ghats is also captured by very well the RCM.~~ However, it ~~seems to slightly underestimate slightly underestimates~~ the magnitude of rainfall over [the](#) peninsular and western part of India ~~—(also reported in Halder et al., 2015).~~ The pattern correlation between rainfall in [the](#) PLC experiment and APHRODITE is 0.71. The Mean Bias calculated over the presented domain with respect to APHRODITE rainfall for the period 1982–2007 is $-0.48 \text{ mm day}^{-1}$ and the RMSE is 3.53 mm day^{-1} . Although ~~there are more station observations aggregated into the IMD data, the APHRODITE dataset has also been chosen for its greater spatial extent over land part compared to IMD. Although~~ daily CI averaged

rainfall during JJAS in both observation (~~IMD~~) and the PLC experiment (CI domain for the RCM is 75.30–86.63° E, 16.92–26.43° N) follows the Poisson distribution, the number of very heavy rainfall events simulated in the RCM is relatively less (figure not shown). This deficiency in climate models in terms of capturing the observed frequency distribution of daily rainfall realistically is a well-known problem (Frei et al., 2003; Kang et al., 2014) and may be attributed to the model dynamics, choice of convective parameterization schemes and their interplay (Frei et al., 2003). The mean and interannual standard deviation of CI averaged rainfall (1982–2007) in PLC (~~IMDAPHRODITE~~) are 77.59 cm (87.28 cm) and 7.57 cm (8.8 cm), respectively. ~~The model performs reasonably~~ Therefore, the model performs well in capturing the observed interannual variability of seasonal rainfall over CI (which is about 10 % of the seasonal mean), although it underestimates both quantities.

~~Air temperature at 2 has been used in modeling studies to validate simulated features of the Indian summer monsoon by the RCM (Pal et al., 2007; Saha et al., 2011).~~ We further evaluate JJAS averaged 2 m near-surface air temperature simulated by the model ~~is compared here~~ with the IMD data for the period 1982 ~~till~~ until 2005. Mean surface temperature in observation is highest over the north, ~~east~~, north-west, east and the rain-shadow region ~~to the east of~~ over the peninsular India (Fig. S1a in the Supplement). In ~~the model contrast~~, surface temperature simulated by the model is high particularly over the north-west (Fig. S1b). ~~As a result, a~~ A cold bias of about 3–4 °C in temperature is found over ~~northern, western and eastern India that is probably linked with rest of the Indian region that is linked with biases in~~ the land-surface ~~and as well as~~ other parameterization schemes in the model such as radiation, convection etc. (Fig. S1c). ~~Bias is much higher over extreme north and north-east India.~~ Such biases have also been noted and discussed in Halder et al., 2015. The pattern correlation between IMD and RegCM4 simulated JJAS 2 m air temperature is 0.76. A similar pattern of cold bias in 2 m near surface air temperature is also seen in PLCS experiment (figure not shown). As our objective is to analyze the mean differences between model simulations, these biases are not expected to have significant effect on the results.

Observed large-scale circulation features from [the](#) NCEP/NCAR reanalysis in the lower troposphere (850 hPa) shows the location of the low-level Somali Jet over the Arabian Sea, cross-equatorial flow and the easterlies south of the Equator (Supplement Fig. S2a). The RCM captures the location of these large-scale low-level features [reasonably-very](#) well in both PLC and PLCS (Fig. S2b and c). However, the wind speed is slightly overestimated in [the](#) PLC experiment, particularly along the core of the Somali Jet and the BoB. [This As mentioned earlier, this](#) overestimation conforms to the positive rainfall bias over the Ocean, the Arabian Sea and the BoB in the RCM. ~~Increased precipitation in the model leads to greater release of latent heat in the atmosphere. As circulation and precipitation are convectively coupled, this heating invigorates convection and vertical motion in the atmosphere, which further increases large-scale low-level convergence and hence the strength of circulation.~~ On the contrary, low-level wind speed is reduced around the core of the Jet, over the Indian Ocean, BoB and also over land in the PLCS experiment, which is associated with the reduction in precipitation.

~~At the upper level (200 hPa) large-scale~~ circulation in observations show the sub-Tropical Westerly Jet stream about north of 30° N, the Tropical Easterly Jet over the equatorial Indian Ocean and the Tibetan Anticyclone south of 30° N (Fig. S2d). ~~Location~~ [The location](#) of these characteristic circulation features is also well captured in the simulations PLC and PLCS (Fig. S2e and f). The model simulated wind speed is stronger than observations in the PLC experiment at upper level like at 850 hPa. The pattern correlation between NCEP/NCAR reanalysis and PLC simulated wind at 850 hPa (200 hPa) is 0.81 (0.95). [We infer that the model RegCM4 performs well in simulating the climatological mean features of Indian summer monsoon. This gives us confidence to conduct sensitivity experiments with the model.](#)

The climatological onset date of ~~Indian summer monsoon rainfall (ISMR)~~ [ISMR](#) based on the Tropospheric Temperature Gradient (TTG) index (Xavier et al., 2007) in the PLC experiment is around 20 May, with interannual standard deviation of about 8 days. Hence, it is advanced by about 10 days from the observed onset. Unlike in the observations, ISMR onset in the model happens to be in the month of May for most years. Therefore,

for our analysis of temperature extremes in the model we have used data for the period JJAS (instead of JAS) in order to have a longer time series and more confidence in the model results. ~~We infer that the model RegCM4 performs reasonably well in simulating the climatological mean features of Indian summer monsoon. This gives us confidence to conduct sensitivity experiments with the model.~~

4.2 Changes in circulation and seasonal rainfall due to LULCC

Mean surface winds (at 10 m) during JJAS blow from west to east over peninsular India and the Indian Ocean, carrying moisture from the Arabian Sea. They turn anti-clockwise over the BOB to move north-west over the Gangetic plains, thus forming the monsoon trough all along CI where the mean wind speed is very low (Fig. 5a). As forest cover in the HLC experiment is replaced by crop PFTs over most of the land part in the PLC experiment, surface roughness length is decreased due to reduction in vegetation height and LAI. We note that surface wind has become more westerly (easterly) over southern and western (northern) India and shows increased anti-cyclonic circulation (Fig. 5b). It has intensified significantly by about 1 m s^{-1} over peninsular India and 0.5 m s^{-1} over the northern India (Fig. 5c). This implies less convergence of moisture and also a reduction in rainfall in the PLC experiment (see Sud et al., 1998; Takata et al., 2009) ~~to be that is~~ discussed in the following paragraph. This intensification of surface wind speed further extends up to the depth of the boundary layer that interacts more directly with the large-scale circulation (figure not shown). Surface and boundary layer winds also intensify in a similar fashion in the PLCS experiment when compared to HLCS and depict the effect of reduced roughness length due to LULCC. It is interesting to note that these significant changes take place ~~only mainly~~ over the land portion of the domain and partly over water bodies close to its boundaries.

The climatological seasonal (JJAS) distribution of rainfall over the Indian subcontinent has been discussed in detail in relation to Fig. 4. ~~Difference~~ Differences in seasonal rainfall between PLC and HLC ~~shows show~~ a significant reduction over a large part of CI, peninsular and northwest India (Fig. 6a). It is interesting to note that the pattern of

decrease matches very well with the regions that show an increase in crop PFTs from 1950 to 2005, with maximum changes over the northwest of India. The magnitude of decrease
645 in seasonal rainfall is quite high (by 5–7 cm) over certain regions, however it is difficult to find out exact reason for such changes at every grid. It may be due to changes in local instability brought about by land–atmosphere feedback processes or changes in large-scale moisture convergence or both. A part of these changes also depend on the choice of parameterization schemes in the RCM. Observational evidence suggests that despite
650 increase in water holding capacity of the atmosphere on a large scale, changes in rainfall are very localized. It is plausible that large-scale conditions and moisture convergence in the PLC experiment might be relatively unfavorable for formation of rainfall compared to the HLC experiment. In order to ~~find out~~ analyze changes in the large-scale moisture convergence, we ~~first~~ calculated vertically integrated moisture flux (qV) from surface to
655 300 hPa. Following Helmholtz's theorem ~~and the methodology of Behera et al. (1999)~~, velocity potential is further calculated that represents the divergent component of that moisture flux ~~. It turns out, from the difference of that divergent component, (cf. Behera et al. 1999). From the difference, it turns out~~ that large-scale moisture convergence is reduced in the PLC experiment and contributes to the reduction in rainfall over CI (Fig. 6a).
660 However, it also remains to be explored how much do the changes in LULC contributes to reduction in surface evaporation and hence moisture convergence over land. This will be discussed in Sects. 4.3 and 4.4. ~~It is known from earlier studies~~ Studies have shown that precipitation variance is amplified by land–atmosphere feedback over those regions that are least affected by SST (~~see cf.~~ Koster et al., 2000). Therefore, it is possible that higher
665 decrease in precipitation over the northwest semi-arid northwest region of India, ~~which is far that is farther~~ away from the influence of SST is dominated by changes in local land surface processes.

As the monsoon circulation in the PLCS experiment is relatively weaker than in the PLC experiment and SSTs are cooler, large-scale moisture flux into the monsoon domain is also
670 expected to be less. Therefore, changes in rainfall over land would better reflect the impact of local land–atmosphere feedbacks due to LULCC. It is evident from Fig. 6b that there is

indeed a significant reduction in seasonal rainfall in PLCS, and over a much wider area of CI and the Western Ghats region than in PLC. Enhancement of rainfall is also evident over some parts of the north and west of India that depict an increase in forests. ~~Maximum decrease~~ (Fig. 1e). ~~Decrease~~ in seasonal rainfall ~~(, by a maximum of about 3–4 cm)-occurs~~ is evident over most parts of CI. Decrease in large-scale moisture convergence in the PLCS experiment is also much widespread extending up to the Arabian Sea, and stronger than in PLC experiment (as evident from denser dashed contours).

4.2.1 Changes in frequency of daily rainfall and intensity

We ~~shall~~ study next how changes in seasonal rainfall over CI are also associated with the changes in daily rainfall in the moderate and extreme category. We ~~adopted~~ adopt the criteria for determining thresholds for categorizing moderate and extreme daily rainfall events over CI in the model, that is not exactly the same but is consistent with the method of Goswami et al., 2006. Any daily rainfall total averaged in a grid box is considered as an event. Percentiles of observed (~~IMD~~ APHRODITE) daily rainfall over CI during JJAS are calculated for the period 1982 ~~till~~ until 2007 to identify the value that corresponds to the range of moderate rainfall and lower threshold of extreme rainfall events (see Sect. 3.3). The observed percentiles are then compared with those calculated for the model to categorize daily moderate and extreme rainfall events in the model. In this way, moderate ~~(extreme)~~ events are identified in the model when $5.34 < \text{daily rainfall} \leq 41.72 \text{ mm day}^{-1}$ ~~(and extreme events are identified when~~ daily rainfall $> 59.94 \text{ mm day}^{-1}$). Figure 7a (Fig. 7b) depicts the difference in total number of moderate rainfall events (intensity of rainfall in moderate category) between PLC and HLC experiments during JJAS from 1982 ~~till~~ until 2008. Note that for PLC, there is a significant and widespread decrease over CI and the spatial pattern coincides with the increase in crop PFTs in PLC experiment. It ~~may~~ can also be noted, that the pattern of decrease matches very well with that of the changes in seasonal rainfall. Following the above methodology, moderate ~~(extreme)~~ rainfall events are identified in the PLCS and HLCS experiments when $4.97 < \text{daily rainfall} \leq 41.62 \text{ mm day}^{-1}$. Likewise, extreme rainfall events are identified when daily rainfall $> 59.80 \text{ mm day}^{-1}$. The spatial

700 pattern of changes depict that this decrease takes place over a larger part of CI as well as the BI domain (shown in Fig. 4b). On the contrary, changes in extreme rainfall events or the intensity of rainfall in that category are not found to be significant between PLC and HLC or between PLCS and HLCS, whether ~~looked-at~~ analyzed spatially (figure not shown) or in an aggregated sense over CI.

705 The mean decrease in the number of moderate rainfall events between PLC and HLC ~~is 388, and that between~~ PLCS and HLCS ~~) is 388 (is 450)~~, which is significant at the 90 % level of significance. Over the larger BI domain, decrease in moderate rainfall events between PLC and HLC is even greater and is about 642 (significant at 95 %). We note that the order of decrease is comparable to the observed decrease in the number of moderate
710 rainfall events over CI (about ~~500~~ 640) in the last 55 years. Along with the number of events, intensity of rainfall amount in a season in the moderate category also decreases significantly at the 95 % level of significance. The decrease in number of moderate events and corresponding rainfall intensity between PLCS and HLCS is even greater, aided by further reduction in large-scale convergence of moisture apart from local land–atmosphere
715 feedbacks. ~~The additional set~~ Therefore, the additional pair of sensitivity experiments with de-trended SSTs further help in establishing our hypothesis. As moderate rainfall events constitute a major portion of the seasonal (JJAS) rainfall ~~;~~ (85 % in observations) we conclude that decrease in seasonal mean rainfall over CI is mainly attributed to differences in the moderate rainfall category due to increase in crop PFTs. Inclusion of light rainfall
720 events ($1 < \text{daily rainfall} < 5.34 \text{ mm day}^{-1}$) in the ~~experiments~~ analysis along with the moderate category does not change our result. We further investigate changes in surface temperature over land and other associated fluxes in order to better understand the above large-scale changes.

4.3 Changes in surface air temperature

4.3.1 PLC and HLC experiments

Daily 2 m mean air temperature during JJAS in PLC is ~~warmer~~^{higher} than HLC by a maximum of 0.3 °C over CI and parts of south (Fig. 8a); ~~which is relatively less than the observed increase~~. A significant increase in daily maximum temperature (0.4 °C) over the same region as in the mean is also evident (Fig. 8b). The pattern of increase does coincide with increase in crop fraction in PLC (Fig. 1f). Widespread warming is also seen beyond the dry northwestern region of India where the increase in fraction of crop PFTs is more than over CI. Significant cooling is found along a thin belt ~~of~~^{around} the Himalayan foothills in the north that may be attributed to an increase in precipitation (see Fig. 4) as well as changes in albedo and net radiation. A decrease in mean and extreme temperature over small ~~regions~~^{parts} of western India is attributed to an increase in forest PFTs (Fig. 1e). At night, the land surface gets de-coupled from the overlying atmosphere on account of cooling, is capped by a layer of inversion and the effect of land surface processes or vegetation on 2 m temperature is minimized. Therefore, and as discussed in Kothawale et al. (2010), the increase in daily mean temperature is mostly dominated by the increase in daily maximum temperature. However, we also noted an increase in temperature at ~~925 the~~^{the 925} hPa level (figure not shown), implying that the surface warming extends further up to the depth of the boundary layer.

Apart from changes in the mean temperature, there are also changes in the variability of daily mean and maximum temperature as evident from Fig. 9a and b. There is significant increase in temperature variability over the central and eastern part of India that is attributed to LULCC as well as changes in surface net radiation and advection of moisture and heat. Increase in the variance of extreme is more widespread than in the mean. As the mean and variance of daily surface temperature are altered over CI, it is expected that daily extremes will also change. In order to find out the differences in the extreme temperatures, percentiles are calculated using a time series of 122 days for 27 years (June–September 1982–2008). Difference in the 90th percentiles of daily mean and maximum temperature

(in JJAS) between and PLC and HLC is shown in Fig. 9c and d. The 90th percentiles represent the higher temperature extremes attained within the season in the PLC and HLC experiment. The higher extreme values of both daily mean and maximum temperature are about 1 °C more in the PLC experiment over CI. We note that the area of increase coincides very well with the region of maximum increase in the fraction of crop PFTs from 1950 to 2005 (Fig. 1f). It is also interesting to note that the higher extremes ~~get hotter~~ warm by the same order as depicted in observations. Apart from that, the model does not capture the observed warming over the northwest and peninsular India despite changes in the LULC. Over the northwest of India, the mean as well as extreme temperatures decrease on account of an increase in forest cover over a small region (Fig. 1e).

~~We further analyzed changes~~ Changes in other surface variables and cloud cover during JJAS are further analyzed to better understand the causes for surface temperature ~~changes due to LULC change~~. The black contours in different panels in Fig. 10 represent the JJAS mean value from the PLC experiment, while the values in shaded color show the difference. Areas enclosed within the green contours depict ~~significant changes~~ changes that are significant. One would expect the surface pressure over land to decrease and an increase in the land-ocean temperature gradient in the PLC experiment on account of an increase in the surface temperature. However, from Fig. 10a it is evident that surface pressure has increased over most of north, northwest and the Gangetic plains of India in PLC compared to HLC. Although a part of CI and its west, ~~the region that captures the monsoon trough shows a decrease in surface pressure~~ (Fig. 6b) ~~and has the lowest surface pressure during JJAS, shows a decrease,~~ the changes are not significant. Therefore, ~~such~~ changes in surface ~~pressure are not responsible for~~ roughness length mainly dominate the increase in surface wind speed, ~~but changes in roughness length are compared to changes in surface pressure~~. There is also a significant decrease in ~~surface~~ soil moisture (Fig. 10b) associated with the decrease in precipitation, and specific humidity at 2 m (Fig. 10c) over those regions where the fraction of crop PFTs has increased in PLC.

We note a significant increase in surface albedo over the land part (Fig. 10d) that is attributed to the increase in crop PFTs and reduction in precipitation, that leads to drier

soils. ~~However the changes are much less compared to the mean albedo in the PLC.~~ An increase in albedo would tend to reduce the surface temperature. However, we also ~~note that the percentage of~~ find that the cloud cover has decreased significantly over a large part of CI, the west, north and peninsular India in PLC (Fig. 10e). This conforms to the reduction in seasonal precipitation in PLC compared to HLC. Due to reduction in cloud cover, there is also an increase in surface Net Radiation (NRAD) over those regions, although changes are not found to be significant over CI (Fig. 10f). The increase ~~(decrease)~~ in NRAD over central and southern India ~~(the Himalayan foothills)~~ in PLC is contributed by a significant increase ~~(decrease)~~ in net shortwave (SW) radiation ~~(in~~. Whereas the decrease in NRAD over the Himalayan foothills is associated with a significant decrease in both net SW and long-wave longwave (LW) radiation). Over CI, decrease in net LW radiation partly compensates for the increase in net SW radiation in PLC (figure not shown), hence changes in NRAD are small. Decrease in net LW radiation in PLC dominates over the northwest of India. ~~We note~~ The increased NRAD in PLC contributes to a significant enhancement (reduction) in the mean surface sensible heat flux (SHF) in the PLC experiment over those areas that show an increase in crop (forest) cover (Fig. 10g). On the contrary, ~~the~~ latent heat flux (LHF) that is directly associated with the ~~total evapotranspiration shows a significant change~~ ET shows significant changes in the opposite sense (Fig. 10h), leading to an overall enhancement in the ~~Bowen Ratio~~ bowen ratio in PLC (figure not shown). Therefore, we infer that an increase in NRAD and SHF in PLC dominates over changes in surface albedo over India south of 30° N ~~(towards the Tropics)~~ that also conforms to conclusions and contributes to the increase in surface temperature. Our results also conform to the inferences reported in earlier studies (Lawrence and Chase, 2010; Sampaio et al., 2007; Davin and Noblet-Ducoudre, 2010).

It is interesting to note that about 30 % of the ~~decrease in ET (or LHF)~~ changes in LHF over CI, west and southern India in PLC (Fig. 10h) ~~is~~ are mainly contributed by a reduction in transpiration from vegetation and evaporation of canopy-intercepted water due to LULCC (Fig. 10i). Although this decrease is relatively ~~more~~ higher over eastern India than towards ~~the center~~ CI, enhanced ground evaporation arising from increased

precipitation in PLC compared to HLC partly compensates for that. As a result changes in total ET are not significant towards the east of India. Therefore, due to a reduction in surface ET, the increased NRAD absorbed at the surface over central and southern India is ~~used up~~ primarily used in enhancing the SHF and that further contributes to the increase in mean and higher extreme surface temperatures in PLC during JJAS. As mentioned ~~earlier in the introduction~~, the daily spatio-temporal variability of surface temperature may be attributed to local thermodynamic effects due to changes in low-level moisture and surface fluxes as well as large-scale dynamics. In this regard, we note that our ~~inferences~~ results differ from earlier studies that ~~showed a~~ have shown a decrease in growing season surface temperatures over India due to irrigated crops (e.g. Sen Roy et al., 2007; Lee et al., 2009) because we did not use any parameterization scheme for irrigation. Irrigation provides an enhanced source of soil moisture ~~;~~ and hence cools the surface and lowers the temperature due to evaporation.

4.3.2 PLCS and HLCS experiment

We find similar changes when simulated surface temperatures in the ~~remaining two experiments are analyzed~~ PLCS and HLCS experiments are compared. Daily 2 m mean as well as maximum temperature ~~is~~ are significantly enhanced in the PLCS experiment by maximum of 0.5°C , but over a much larger area covering central and southern ~~part~~ parts of India compared to HLCS (Fig. 11a and b). We note that the increase in temperature over CI ~~is greater and due to similar LULCC is higher and more~~ is greater and due to similar LULCC is higher and more widespread than in ~~earlier experiments~~ PLC. Likewise, over the northwest of India, the spatial pattern of increase extends further to the north and shows ~~greater~~ higher increase (0.5°C) in the maximum. Significant cooling is also evident over western and northern India in PLCS over those areas that show increase in forest cover. The higher extremes (*i.e.* 90th ~~percentile~~ percentiles) of the daily mean (maximum) temperature have also increased in the PLCS experiment by 1.2°C (1.0°C), which is ~~greater than in~~ more than in the earlier set of experiments PLC and HLC (Fig. 11c and d). Increase in higher extreme temperature in the PLCS experiment extends further to the west and hence covers a much larger part of CI than in the PLC

experiment. We ~~also~~ further note that the order of increase in temperature as evident from these two experiments is comparable to that inferred from observations.

There ~~is a~~ are significant and widespread ~~decrease in surface decreases in~~ soil moisture, ~~specific humidity and LHF and increase LHF and specific humidity but increases~~ in NRAD and sensible heat flux ~~that contributes toward the increase in surface temperature in PLGS experiment (figure in the PLCS experiment compared to HLCS (figures not shown) .It may be noted that contribute to the increase in surface temperature. It is interesting~~ that in both set of experiments, the increase in surface temperature is slightly towards south of the area that depicts ~~increase in observation~~ an increase in observations. Apart from that, mean monsoonal features ~~in PLGS experiment simulated in the PLCS experiment also~~ convey that there is a decrease in large-scale moisture flow as well as precipitation over the land. As a result alterations in ~~surface fluxes and radiation net radiation and surface fluxes between PLCS and HLCS experiments~~ have a greater impact on changes in ~~air temperature in these additional experiments~~ surface air temperature. Therefore, our experiments with de-trended SST further confirm the proposition that LULCC has partly contributed to the observed increase in surface temperature ~~from 1951 until 2005~~.

4.4 Physical mechanisms

After analyzing ~~changes in local~~ the changes in surface variables and the large-scale in the ~~model set of model experiments~~, one pertinent question arises. How does LULCC lead to ~~a~~ reduction in moderate rainfall events? ~~Calculations based on regression show that evapotranspiration over land controls about 10–20% of the interannual variability in rainfall over CI (Saha et al., 2012). In another study (Halder et al., 2015) we found showed~~ that surface ET can strongly modulate the terrestrial segment of land–atmosphere coupling strength (Dirmeyer, 2011) ~~for and the chances of triggering of convection and~~ precipitation during the Indian summer monsoon. ~~We note, that in the PLC experiment, decrease in~~ From comparison of the PLC and HLC experiments, we note a decrease of about 3 cm in the total evapotranspiration over CI is around 3 (when LHF in Wis converted to mm day⁻¹) ~~that constitutes about, that is~~ 40–60 % of the maximum decrease in ~~total rainfall~~ the total

rainfall magnitude. Although an increase ~~(decrease) in crops (forests) in crop cover and decrease in forest~~ increases the temperature near the surface and within the boundary layer, the associated decrease in local moisture flux could possibly also lower the large-scale convective instability. ~~Vertically~~ To better understand that, we analyze changes in vertically integrated moist static energy (VIMSE) which is a good measure of instability and precipitation in the Tropics (Srinivasan et al., 1996). VIMSE from surface to 500 hPa during JJAS in PLC depicts high values over those areas of land that show maximum seasonal mean rainfall (Fig. 12a). ~~There~~ Differences show that there is a large-scale reduction in VIMSE in the PLC experiment, with significant decrease over a major part of CI and the north (Fig. 12b). ~~Difference~~ Additionally, the difference (PLC-HLC) in the vertical profile of dry static energy (DSE, blue line) ~~depicts~~ suggests an increase in temperature ~~at in~~ the lower levels of ~~atmosphere the troposphere~~ over CI. Despite that, the effect of ~~decrease in moisture~~ decreasing moisture in the lower levels effectively reduces the MSE (red line, Fig. 12c) thereby increasing atmospheric stability and hence ~~lowers~~ lowering the chances of triggering of moist convection over land in the PLC experiment. ~~Lesser~~ Reduced large-scale low-level moisture convergence over the land part in the PLC than HLC, on account of a reduction in surface roughness length (Fig. 6b) also contributes to the ~~reduced~~ reduction in convective instability. These two factors together reduce rainfall in the moderate category.

5 Discussions

~~LULGC in the form of extensive cropland and pasture expansion and deforestation have taken place over India ever since the inception of the Green revolution in the 1960's; associated with urbanization and industrialization.~~ This study explores the hypothesis how LULCC over India has contributed to the observed decrease in moderate rainfall events over CI and increase in extreme daily surface temperatures during the monsoon season: ~~A RCM, on account of its dynamical downscaling capability and better representation of land-atmosphere feedbacks over the monsoon region has been used to prove the hypothesis, from 1951 through 2005, using a RCM. The climatological mean features of~~

Indian summer monsoon are very well captured by the RCM RegCM4. The statistics of the differences between the long simulations with fixed present day and historical land cover (2005) and historical (1950) PFT distributions, LAIs and SAIs demonstrate the impact of LULCC on daily surface temperature and precipitation variability during the monsoon season (JJAS). Another two similar experiments are also made conducted, but with SSTs de-trended over the ocean within the RCM domain in order to eliminate the effect of warming trend in the positive trend in Indian Ocean SSTs on temperature and precipitation changes over land. The climatological mean features of Indian summer monsoon are very well captured by the RCM.

Differences show that seasonal rainfall and large-scale moisture convergence is are significantly decreased in the PLC and PLCS experiments when compared to the HLC and HLCS experiments, respectively. The decrease is enhanced in the case of the PLCS experiment compared to PLC. That decrease in seasonal rainfall is mostly contributed by a significant decrease in moderate rainfall events and amount over CI. Changes in extreme rainfall events are not significant. We demonstrate that a significant increase in surface wind speed over land on account of reduction in surface roughness length is responsible for the decrease in moisture convergence. This is the dynamical response to LULCC. Decrease The increase in surface wind speed is attributed to an increase in crop cover at the expense of forests and hence a reduction in surface roughness length. This way, the dynamical response of regional climate over India to LULCC is demonstrated. Decreases in forest cover also reduces and increases in crops between 1950 and 2005 also lead to reductions in the regional moisture flux emanating from the surface significantly. Therefore, despite significant increase increases in surface and low-level temperature, boundary layer temperature, a decrease of moisture reduces the large-scale convective instability and chances of triggering of convection and hence precipitation over central and north India. This effect mechanism constitutes the thermodynamic response of the regional climate to LULCC. A decrease in total cloud cover increases the surface Net Radiation that net radiation, which together with a decrease in surface moisture results in an increase in the surface sensible heat flux and the Bowen-bowen ratio. Together, they

these changes contribute to the increase in surface temperature mean surface temperature and its extremes. It is ~~important to note~~ noteworthy that the order of increase in surface temperature extremes over India during the summer monsoon season is comparable to that of the observed changes when de-trended SSTs within the model domain are used. Likewise, the order of decrease in moderate rainfall events over CI also become comparable to ~~observed changes in~~ the observed changes during the period 1951–2005.

It is important to note ~~choice of the model parameterization schemes, accuracy of the lateral boundary conditions and the criteria for calculating thresholds for daily moderate and extreme rainfall events have an influence on the results.~~ Apart from that, the model's grid resolution is about half of that of IMD rainfall data. Hence area of a single grid box in IMD data (which represents an event) is about four times of that of the model. This could also be one of the reasons for disparity in the number of moderate rainfall events between observations and the model that the deficiency in the RCM in terms of capturing the frequency distribution of daily very heavy rainfall events over CI realistically could have a bearing on our inferences. Hence, our results are partly dependent on the choice of model parameterization schemes. However, this is a well-known problem related to climate models (Frei et al., 2003; Kang et al., 2014) and similar studies when repeated with other RCMs is expected to give us further evidence on the role of LULCC in affecting the frequency distribution of daily rainfall events over India. Apart from that, the criteria used for calculating thresholds for daily moderate and extreme rainfall events in the RCM may also have influence on the results. There is a cold temperature bias over land in the model RegCM4, and positive rainfall bias over the ocean (figure not shown). ~~that is also evident from earlier studies~~ (Saha et al., 2011, 2012; Halder et al., 2015). Apparently, in all these experiments the global warming signal is also present in the large-scale LBCs used from NCEP/NCAR reanalysis that force the model in one way only. A part of the change in simulated surface temperature and rainfall in the model may also be attributed to non-linear interactions (internal variability) that is model dependent. However, we expect the differences between two simulations with the same model to reduce the effect of these factors and demonstrate the impact of LULCC on regional climate over India. Use of a high-

resolution RCM is more advantageous in excluding large-scale remote feedbacks that take place in a coarse-resolution GCM and ~~helps therefore helps to~~ better resolve regional ~~land surface processes~~land-atmosphere feedbacks. Apart from that, we believe that the LULC data prepared from multiple sources and used as fixed lower boundary condition ~~in this study~~ is much improved compared to other historical reconstructed data ~~(such as that of Ramankutty et al., 2008), although there may be certain level of uncertainty in their estimates, particularly during pre-satellite era. However there is no ambiguity about a continuous decrease in forest cover over the last many decades over the south Asian region- utilized in earlier studies.~~ Nonetheless, our experiments ~~show demonstrate~~ that the decrease in moderate rainfall events over central India is partly attributed to changes in LULC from 1950 to 2005.

6 Conclusions

Apart from an accelerated warming trend in the global mean surface temperature in the later half of the 20th century, the number of extreme events in terms of temperature as well as precipitation has been reported to increase. As regional or local changes in these extremes in different seasons can have different signatures due to complex regional feedbacks associated with the GHGs, clouds, aerosols and other anthropogenic activities such as LULCC, they need greater attention and proper attribution. Regional land-atmosphere feedbacks associated with LULCC are one of the potential drivers of climate change. LULC data shows significant decrease ~~(increase)~~ in the forest ~~(crop)~~ and increase in crop cover over central, south and northwest part of India between ~~1951-1950~~ and 2005. From 1951 ~~till until~~ 2005, the observed mean (extreme) surface temperature over India has increased by a maximum of 1.11 °C (1.48 °C) during the summer monsoon season. There have also been significant changes in the rainfall distribution during ~~last those~~ 55 years ~~over central India. Observed extreme. While observed heavy and very heavy~~ precipitation events have ~~been increasing increased~~ over central India ~~during the monsoon season and have been potentially associated with the significant warming trend over the Indian Ocean. However,~~

due to a significant decrease in moderate rainfall events, the overall seasonal rainfall has reportedly remained stable ~~over India during the 55-year~~ during that period. In this study, we ~~demonstrate that observed LULG cannot reject the hypothesis that LULCC over India~~ has partly contributed to the observed decrease in moderate rainfall events and increase in extreme surface temperature during the summer monsoon season.

~~In order to examine whether changes in surface temperature and rainfall distribution pattern are linked with changes in LULG, two sets of long simulations (27 years) using PFTs of 1950 and 2005 as fixed lower boundary condition are carried using the regional climate model RegCM4.0 coupled with the CLM3.5 land surface model. Model simulations with PFT distribution of 1950 and 2005 are referred to as HLC and PLC experiments respectively. Another two sensitivity experiments with similar model set-up are conducted, but with de-trended SSTs over the ocean in order to remove the effect of warming. They are respectively named as PLGS (with 2005 PFTs) and HLGS (with 1950 PFTs) experiments. It is found that, increase in mean and extreme surface temperatures by 1–1.2 °C over CI in PLC the present land cover experiment coincides with the region of decrease (increase) in forest (crop) in forest and increase in crop type of PFTs. Furthermore, that increase is found to be even higher and more widespread over the Indian land mass in the PLGS experiment.~~

~~Reduction in local evapotranspiration followed by an increase in surface sensible heat flux and Bowen ratio in the PLC experiment is consistent with the decrease in forest cover. Furthermore, wind speed at the lower atmospheric levels increase significantly in the PLC experiment that suggests a decrease in region when the positive trend in the Indian Ocean SSTs is removed. There is a reduction in large-scale convective instability and moisture convergence over land. Therefore, there is a net decrease in moisture supply over land in PLC as compared to the HLC experiment. Apart from that, decrease in local moisture supply also significantly reduces the large-scale convective instability over land and formation of clouds. Net radiation is increased and that contributes to the increase in surface sensible heat flux and temperature extremes. that leads to decrease in seasonal precipitation in the PLC experiment compared to HLC. As the major part of monsoon portion~~

of monsoon seasonal rainfall occurs through moderate ~~events (about 85)~~ rainfall events (Goswami et al. (2006)), it is expected that ~~decrease in total~~ the decrease in moisture flux and large-scale convective instability over land would also ~~decrease lead to a decrease~~ in the moderate rainfall ~~events~~ category. The model results indeed support ~~that theory~~ your hypothesis, and show that regions with a decrease in forest cover ~~depict decrease also depict a decrease in~~ the number of moderate rainfall events as well as the ~~quantity of total amount of~~ rainfall in that category. Changes in heavy rainfall events ~~that are more determined by large-scale moisture availability and triggered by local instabilities, are not~~ are not found to be significant. These results are further supported by the two additional sensitivity experiments PLCS and HLCS. ~~Availability of local surface moisture flux that is associated with the type of LULC is~~ We conclude that changes in local/regional moisture flux and surface roughness length that are associated with this type of LULCC are crucial in determining the changes in large-scale instability and moisture convergence over land and ~~hence chances of convective triggering of rainfall~~ the frequency distribution of daily rainfall events over the Indian monsoon region. Therefore, this study demonstrates that LULC changes-LULCC in the last 55 years have contributed partly to the observed decrease in moderate rainfall events over India as well as increase in extreme surface temperatures.

Understanding the mechanisms responsible for observed changes in daily rainfall distribution and extreme surface temperature ~~and rainfall distribution~~ in the monsoon regions is important for the scientific community and policy makers as well. It is conceivable that, as the global mean temperature becomes warmer and the ~~climate~~ regional climates possibly more unpredictable, LULC change due to population growth, deforestation/afforestation, agricultural expansion and urbanization would add more uncertainties through its dynamic (~~e.g. strength of~~ changes in large-scale circulation) and thermodynamic effects (albedo, evaporation and instability changes). ~~Effect of urbanization is expected to increase warming, reduce local evaporation and also possibly affect rainfall distribution.~~ However, this study does not include urbanization effects. Apart from that, impact of aerosols, GHGs and irrigation activity ~~has~~ have also not been considered here which would introduce competing influences. Therefore, part of the regional warming over

India seen in observations could not be explained only through ~~changes in LULC~~ LULCC that we have isolated here. ~~It would be interesting to study in future the effect of LULC change~~ Investigation of the impact of LULCC in a high-resolution ~~global coupled global climate~~ model where the ~~effect of other anthropogenic forcing are included, and also where the LULC land cover~~ changes with time or dynamic vegetation is used, would make another interesting study. Furthermore, studies similar as this with a suite of climate models would further augment our understanding about the role of LULCC in Indian monsoon climate. Nevertheless, this study shows that it is highly important to include projected anthropogenic changes in regional LULC in IPCC future climate change scenarios.

The Supplement related to this article is available online at [doi:10.5194/hessd-0-1-2015-supplement](https://doi.org/10.5194/hessd-0-1-2015-supplement).

Acknowledgements. Authors thank the Abdus Salam International Centre for Theoretical Physics in Trieste, Italy for making the model codes of RegCM4.0 available for this study from <http://gforge.ictp.it/gf/>. Global time varying forcing datasets for the RCM and fixed datasets for the CLM3.5 land surface model are obtained from <http://users.ictp.it/pubregcm/RegCM3RegCM4/globedat.htm>. The PFT based global vegetation datasets were prepared at the Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, USA. Daily high-resolution gridded ~~rainfall (Product No. 1/2005) and~~ temperature data (Product No. 3/2008) ~~which are~~ prepared from quality controlled station observations can be purchased for a fee from the National Climate Centre (NCC), IMD, Pune, India. Details about the procedure of purchase can be searched in the link “NCC Gridded Data Product”, present in the webpage http://www.imdpune.gov.in/publication/pub_index.html. The figures have been prepared using the freely available software package GrADS.

Authors gratefully acknowledge the use of data and resources at the Indian Institute of Tropical Meteorology (IITM), Pune, India. Fellowship of S. Halder was funded by the CSIR, New Delhi, India during completion of this study for his PhD thesis while at IITM. S. Halder and P. Dirmeyer are supported by a grant from the Ministry of Earth Sciences (~~India~~), Govt. of India to the Center for Ocean-Land-Atmosphere Studies of George Mason University. Support for the Twentieth

Century Reanalysis Project dataset is provided by the U.S. Department of Energy, Office of Science
Innovative and Novel Computational Impact on Theory and Experiment (DOE INCITE) program,
and Office of Biological and Environmental Research (BER), and by the National Oceanic and
Atmospheric Administration Climate Program Office.

References

- Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P., Janowiak, J., Rudolf, B., Schneider, U.,
Curtis, S., Bolvin, D., Gruber, A., Susskind, J., and Arkin, P.: The version 2 Global Precipitation
Climatology Project (GPCP) monthly precipitation analysis (1979–present), *J. Hydrometeorol.*, 4,
1147–1167, doi:10.1175/1525-7541(2003)004<1147:TVGPCP>2.0.CO;2, 2003.
- Alexander, L. V., Zhang, X., Peterson, T. C., Caesar, J., Gleason, B., Tank, A. M. G. K., Haylock, M.,
Collins, D., Trewin, B., Rahimzadeh, F., Tagipour, A., Kumar, K. R., Revadekar, J., Griffiths, G.,
Vincent, L., Stephenson, D. B., Burn, J., Aguilar, E., Brunet, M., Taylor, M., New, M., Zhai, P.,
Rusticucci, M., and Vazquez-Aguirre, J. L.: Global observed changes in daily climate extremes
of temperature and precipitation, *J. Geophys. Res.*, 111, D05109, doi:10.1029/2005JD006290,
2006.
- Ali, H., Misra, V., and Pai, D. S.: Observed and projected urban extreme rainfall events in India, *J.*
Geophys. Res., 19, 12621–12642, doi:10.1002/2014JD022264, 2014.
- Allen, M. R. and Ingram, W. J.: Constraints on future changes in climate and the hydrologic cycle,
Nature, 419, 224–231, doi:10.1038/nature01092, 2002.
- Avila, F. B., Pitman, A. J., Donat, M. G., Alexander, L. V., and Abramowitz, G.: Climate model
simulated changes in temperature extremes due to land cover change, *J. Geophys. Res.*, 117,
D04108, doi:10.1029/2011JD016382, 2012.
- Behera, S. K., Krishnan, R., and Yamagata T.: Unusual ocean atmosphere conditions in the tropical
Indian ocean during 1994, *Geophys. Res. Lett.*, 26, 3001–3004, doi:10.1029/1999GL010434,
1999.
- Bonan, G. B., Oleson, K. W., Vertenstein, M., Levis, S., Zeng, X., Dai, Y., Dickin-
son, R. E., and Yang, Z.-L.: The land surface climatology of the community land
model coupled to the NCAR community climate model, *J. Climate*, 15, 3123–3149,
doi:10.1175/1520-0442(2002)015<3123:TLSCOT>2.0.CO;2, 2002.

- Bounoua, L., DeFries, R., Collatz, G. J., Sellers, P., and Khan, H.: Effects of land cover conversion on surface climate, *Climatic Change*, 52, 29–64, 2002.
- 1095 Chow, K. C., Cha, J. C. L., Pal, J. S., and Giorgi, F.: Convection suppression criteria applied to the MIT cumulus parameterization scheme for simulating the Asian summer monsoon, *Geophys. Res. Lett.*, 32, L24709, doi:10.1029/2006GL028026, 2006.
- Chowdary, J. S., John, N., and Gnanaseelan, C.: Interannual variability of surface air-temperature over India: impact of ENSO and Indian Ocean Sea surface temperature, *Int. J. Climatol.*, 34, 2, 416–429, doi:10.1002/joc.3695, 2013.
- 1100 Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Matsui, N., Allan, R. J., Yin, X., Gleason, B. E., Vose, R. S., Rutledge, G., Bessemoulin, P., Bronnimann, S., Brunet, M., Crouthamel, R. I., Grant, A. N., Groisman, P. Y., Jones, P. D., Kruk, M. C., Kruger, A. C., Marshall, G. J., Maugeri, M., Mok, H. Y., Nordli Ø., Ross, T. F., Trigo, R. M., Wang, X. L., Woodruff, S. D., and Worley, S. J.: Review article: the Twentieth Century Reanalysis Project, *Q. J. Roy. Meteor. Soc.*, 137, 1–28, doi:10.1002/qj.776, 2011.
- Davin, E. L. and De Noblet-Ducoudrè, N.: Climatic impact of global-scale deforestation: radiative versus nonradiative processes, *J. Climate*, 23, 97–112, 2010.
- Dirmeyer, P. A.: The terrestrial segment of soil moisture-climate coupling, *Geophys. Res. Lett.*, 38, 16, doi:10.1029/2011GL048268, 2011.
- 1110 Dirmeyer, P. A., Niyogi, D., de Noblet-Ducoudrè, N., Dickinson, R. E., and Snyder, P. K.: Editorial: impacts of land use change on climate, *Int. J. Climatol.*, 30, 1905–1907, 2010.
- Douglas, E., [Beltrán-Przekurat, A., Niyogi, D., Pielke Sr., R. A., and Vörösmarty, C. J.: The impact of agricultural intensification and irrigation on land-atmosphere interactions and Indian monsoon precipitation — A mesoscale modeling perspective, *Global Planet. Chang.*, 67, 117–128, doi:10.1016/j.gloplacha.2008.12.007, 2009.](#)
- 1115 [Douglas, E. M., Niyogi, D., Frolking, S., Yeluripati, J. B., Pielke Sr., R. A., Vorosmarty, C. J., and Mohanty, U. C.: Change in moisture and energy fluxes due to agricultural land use and irrigation in the Indian monsoon belt, *Geophys. Res. Lett.*, 33, L14403, doi:10.1029/2006GL026550, 2006.](#)
- 1120 Dutta, S. K., Das, S., Kar, S. C., Mohanty, U. C., and Joshi, P. C.: Impact of vegetation on the simulation of seasonal monsoon rainfall over the Indian subcontinent using a regional model, *J. Earth Syst. Sci.*, 118, 5, 413–440, 2009.
- Elguindi, N., Bi, X., Giorgi, F., Nagarajan, B., Pal, J., Solmon, F., Rauscher, S., and Zakey, A.: RegCM version 4.0 User's Guide, ICTP, Trieste, Italy, available at:

- gforge.ictp.it/gf/download/docmanfileversion/6/253/UserGuide.pdf, last access: June 2015, 2010.
- Feddema, J., Oleson, K., Bonan, G., Mearns, L., Washington, W., Meehl, G., and Nychka, D.: A comparison of a GCM response to historical anthropogenic land cover change and model sensitivity to uncertainty in present-day land cover representations, *Clim. Dynam.*, 25, 581–609, doi:10.1007/s00382-005-0038-z, 2005.
- Feser, F. and Barcikowska, M.: The influence of spectral nudging on typhoon formation in regional climate models, *Environ. Res. Lett.*, 7, 014024, doi:10.1088/1748-9326/7/1/014024, 2012.
- Feser, F., Rockel, B., von Storch, H., Winterfeldt, J., and Zahn, M.: Regional climate models add value to global model data: a review and selected examples, *B. Am. Meteorol. Soc.*, 92, 1181–1192, doi:10.1175/2011BAMS3061.1, 2011.
- Frei, C., Christensen, J. H., De'que, M., Jacob, D., Jones, R. G., and Vidale, P. L.: Daily precipitation statistics in regional climate models: evaluation and intercomparison for the European Alps, *J. Geophys. Res.*, 108, 4124, doi:10.1029/2002JD002287, 2003.
- Fritsch, J. M. and Chappell, C. F.: Numerical prediction of convectively driven mesoscale pressure systems. Part I: Convective parameterization, *J. Atmos. Sci.*, 37, 1722–1733, doi:10.1175/1520-0469(1980)037<1722:NPOCDM>2.0.CO;2, 1980.
- Garratt, J. R.: Sensitivity of climate simulations to land-surface and atmospheric boundary layer treatments – a review, *J. Climate*, 6, 419–449, doi:10.1175/1520-0442(1993)006<0419:SOCSTL>2.0.CO;2, 1993.
- Gao, X., Shi, Y., Zhang, D., Wu, J., Giorgi, F., Ji, Z., and Wang, Y.: ~~Uncertainties in monsoon precipitation projections over China: results from two high-resolution RCM simulations~~, *Clim. Res.*, 52, 213–226, 2012.
- Giorgi, F. and Bates, G. T.: The climatological skill of a regional model over complex terrain, *Mon. Weather Rev.*, 117, 2325–2347, 1989.
- Giorgi, F. and Mearns, L. O.: Introduction to special section: Regional climate modeling revisited, *J. Geophys. Res.*, 104, D6, 6335–6352, 1999.
- Giorgi, F., Coppola, E., Solmon, F., Mariotti, L., Sylla, M. B., Bi, X., Elguindi, N., Diro, G. T., Nair, V., Giuliani, G., Turuncoglu, U. U., Cozzini, S., Guttler, I., O'Brien, T. A., Tawfik, A. B., Shalaby, A., Zakey, A. S., Steiner, A. L., Stordal, F., Sloan, L. C., and Brankovic, C.: RegCM4: model description and preliminary tests over multiple CORDEX domains, *Clim. Res.*, 52, 7–29, doi:10.3354/cr01018, 2012.

- Goldewijk, K. K.: Estimating global land use change over the past 300 years: the HYDE database, *Global Biogeochem. Cy.*, 15, 417–433, 2001.
- Goswami, B. N., Venugopal, V., Sengupta, D., Madhusoodanan, M. S., and Xavier, P. K.: Increasing trend of extreme rain events over India in a warming environment, *Science*, 314, 1442–1445, doi:10.1126/science.1132027, 2006.
- Grell, G. A.: Prognostic evaluation of assumptions used by cumulus parameterization, *Mon. Weather. Rev.*, 121, 764–787, doi:10.1175/1520-0493(1993)121<0764:PEOAUB>2.0.CO;2, 1997.
- Halder, S., Dirmeyer, P., and Saha, S. K.: Uncertainty in the mean and variability of Indian summer monsoon due to land atmosphere feedback in RegCM4, *J. Geophys. Res.*, ~~in-review~~, [120](#), [18](#), [9437–9458](#), doi:10.1002/2015JD023101, 2015.
- Haerter, J. O. and Berg, P.: Unexpected rise in extreme precipitation caused by a shift in rain type?, *Nat. Geosci.*, 2, 372–373, doi:10.1038/ngeo523, 2001.
- Holtzlag, A. A. M., Bruijn, E. I. F. D., and Pan, H. L.: A high resolution air mass transformation model for short-range weather forecasting, *Mon. Weather. Rev.*, 118, 1561–1575, doi:10.1175/1520-0493(1990)118<1561:AHRAMT>2.0.CO;2, 1999.
- Hurtt, G. C., Frolking, S., Fearon, M., Moore, B., Shevliakova, E., Malyshev, S., Pacala, S., and Houghton, R.: The underpinnings of land-use history: three centuries of global gridded land-use transitions, wood-harvest activity, and resulting secondary lands, *Glob. Change Biol.*, 12, 1208–1229, doi:10.1111/j.1365-2486.2006.01150.x, 2006.
- Jaswal, A., Rao, G. P., and De, U.: Spatial and temporal characteristics of evaporation trends over India during 1971–2000, *Mausam*, 59, 149–158, 2008.
- Jaswal, A. K., Rao, P. C. S., and Singh, V.: Climatology and trends of summer high temperature days in India during 1969–2013, *J. Earth Syst. Sci.*, 124, 1–15, 2015.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–470, 1996.
- Kang, I.-S., Yang, Y. M., and Tao, W. K.: GCMs with implicit and explicit representation of cloud microphysics for simulation of extreme precipitation frequency, *Clim. Dynam.*, 45, 325–335, doi:10.1007/s00382-014-2376-1, 2014.

- Karl, T. R., Jones, P. D., Knight, R. W., Kukla, G., Mummer, N., Razuvayev, V., Gallo, K. P., Lindseay, J., Charlson, R. J., and Peterson, T. C.: Asymmetric trends of daily maximum and minimum temperature, *B. Am. Meteorol. Soc.*, 74, 6, 1007–1023, 1993.
- Kiehl, J. T., Hack, J. J., Bonan, G. B., Boville, B. A., Briegleb, B. P., Williamson, D. L., and Rasch, P. J.: Description of the NCAR Community Climate Model (CCM3), NCAR, Boulder, Colorado, USA, Tech. Rep., NCAR, NCAR/TN-420+STR, 1996.
- Kishtawal, C., Niyogi, D., Tewari, M., Pielke Sr., R. A., and Shepherd, M.: Urbanization signature in the observed heavy rainfall climatology over India, *Int. J. Climatol.*, 30, 1908–1916, doi:10.1002/joc.2044, ~~2010~~-2009.
- Koster, R. D., Suarez, M. J., and Heiser, M.: Variance and predictability of precipitation at seasonal-to-interannual timescales, *J. Hydrometeorol.*, 1, 26–46, doi:10.1175/1525-7541(2000)001<0026:VAPOPA>2.0.CO;2, 2001.
- Kothawale, D. R. and Rupa Kumar, K.: On the recent changes in surface temperature trends over India, *Geophys. Res. Lett.*, 32, L18714, doi:10.1029/2005GL023528, 2005.
- Kothawale, D. R., Munot, A. A., and Kumar, K. K.: Surface air temperature variability over India during 1901–2007, and its association with ENSO, *Clim. Res.*, 42, 89–104, doi:10.3354/cr00857, 2010.
- Krishnan, R. and Ramanathan, V.: Evidence of surface cooling from absorbing aerosols, *Geophys. Res. Lett.*, 29, 9, 1340, 89–104, doi:10.1029/2002GL014687, 2002.
- Krishnan, R., Sabin, T. P., Vellore, R., Mujumdar, M., Sanjay, J., Goswami, B. N., Hourdin, F., Dufresne, J. -L., and Terray, P.: Deciphering the desiccation trend of the South Asian monsoon hydroclimate in a warming world, *Clim. Dynam.*, doi:10.1007/s00382-015-2886-5, 2015.
- Laprise, R., de Elía, R., Caya, D., Biner, S., Lucas-Picher, P., Diaconescu, E., Leduc, M., Alexandru, A., and Separovic, L.: Challenging some tenets of regional climate modeling, *Meteorol. Atmos. Phys.*, 100, 3–22, 2008.
- Lawrence, P. J. and Chase, T. N.: Investigating the climate impacts of global land cover change in the community climate system model, *Int. J. Climatol.*, 30, 2066–2087, doi:10.1002/joc.2061, 2010.
- Leduc, M. and Laprise, R.: Regional climate model sensitivity to domain size, *Clim. Dynam.*, 32, 833–854, 2009.
- Lee, E., Chase, T. N., Rajagopalan, B., Barry, R. G., Biggs, T. W., and Lawrence, P. J.: Effects of irrigation and vegetation activity on early Indian summer monsoon variability, *Int. J. Climatol.*, 29, 573–581, doi:10.1002/joc.1721, 2009.

1220 [Lei, M., Niyogi, D., Kishtawal, C., Pielke Sr., R. A., Beltrán-Przekurat, A., Nobis, T., and Vaidya, S.: Effect of explicit urban land surface representation on the simulation of the 26 July 2005 heavy rain event over Mumbai, India, Atmos. Chem. Phys. Discussions, 8, 8773–8816, doi:10.5194/acp-8-5975-2008, 2008.](#)

1225 Lohar, D. and Pal, B.: The effect of irrigation on premonsoon season over South West Bengal, India, J. Climate, 8, 2567–2570, doi:10.1175/1520-0442(1995)008<2567:TEOIOIP>2.0.CO;2, 1995.

Mahmood, R., Pielke Sr., R. A., Hubbard, K., Niyogi, D., Dirmeyer, P., McAlpine, C., Carleton, A., Hale, R., Gameda, S., Beltran-Przekurat, A., Baker, B., McNider, R., Legates, D. R., Shepherd, M., Du, J., Blanken, P., Frauenfeld, O. W., Nair, U. S., and Fall, S.: Land cover changes and their biogeophysical effects on climate, Int. J. Climatol., 34, 929–953, 2010.

1230 Nayak, S. and Mandal, M.: Impact of land-use and land-cover changes on temperature trends over Western India, Curr. Sci. India, 102, 1166–1173, 2012.

Niranjan, Kumar, K., Rajeevan, M., Pai, D. S., Srivastava, A. K., and Preethi, B.: On the observed variability of monsoon droughts over India, Weather and Climate Extremes, 1, 42–50, doi:10.1016/j.wace.2013.07.006, 2013.

1235 Niyogi, D., [Chang, H. -I., Chen, F., Gu, L., Kumar, A., Menon, S., and Pielke Sr., R., A.: Potential impacts of aerosol-land-atmosphere interactions on the Indian monsoonal rainfall characteristics, Natural Hazards, Special Issue on Monsoons, Invited Contribution, 42, 2, 345–359, DOI 10.1007/s11069-006-9085-y, 2007.](#)

1240 [Niyogi, D.,](#) Kishtawal, C. M., Tripathi, S., and Govindaraju, R. S.: Observational evidence that agricultural intensification and land use change may be reducing the Indian summer monsoon rainfall, Water Resour. Res., 46, W03533, doi:10.1029/2008WR007082, 2010.

Oleson, K. W., Niu, G. Y., Yang, Z. L., Lawrence, D. M., Thornton, P. E., Lawrence, P. J., Stockli, R., Dickinson, R. E., Bonan, G. B., Levis, S., Dai, A., and Qian, T.: Improvements to the Community Land Model and their impact on the hydrological cycle, J. Geophys. Res., 113, G01021, doi:10.1029/2007JG000563, 2008.

1245 Oleson, K. W., Lawrence, D. M., Bonan, G. B., Flanner, M. G., Kluzek, E., Lawrence, P. J., Levis, S., Swenson, S. C., Thornton, P. E., Dai, A., Decker, M., Dickinson, R., Feddema, J., Heald, C. L., Hoffman, F., Lamarque, J. F., Mahowald, N., Niu, G. Y., Qian, T., Randerson, J., Running, S., Sakaguchi, K., Slater, A., Stockli, R., Wang, A., Yang, Z. L., Zeng, X., and Zeng, X.: Technical description of version 4.0 of the Community Land Model (CLM), NCAR, Boulder, Colorado, USA, NCAR Technical Note NCAR/TN-387+STR, NCAR, 257 pp., 2010.

- Pai, D. S. and Sridhar, L.: Long term trends in the extreme rainfall events over India, in: High-Impact Weather Events over the SAARC Region, edited by: Ray, K., Mohapatra, M., Bandyopadhyay, B. K., and Rathore, L. S., Springer International Publishing, Cham, Switzerland with Capital Publishing Company, New Delhi, India, 229–240, 2015.
- Pai, D. S., Nair, S. A., and Ramanathan, A. N.: Long term climatology and trends of heat waves over India during the recent 50 years (1961–2010), *Mausam*, 64, 585–604, 2013.
- Pal, J. S., Giorgi, F., Bi, X., Elguindi, N., Solmon, F., Gao, X., Rauscher, S. A., Francisco, R., Zakey, A., Winter, J., Ashfaq, M., Syed, F. S., Bell, J. L., Diffenbaugh, N. S., Karmacharya, J., Konar, A., Martinez, D., da Rocha, R. P., Sloan, L. C., and Steiner, A. L.: Regional climate modeling for the developing world: the ICTP RegCM3 and RegCNET, *B. Am. Meteorol. Soc.*, 88, 9, 1395–1409, doi:10.1175/BAMS-88-9-1395, 2007.
- Panda, D. K. and Kumar, A.: The changing characteristics of monsoon rainfall in India during 1971–2005 and links with large scale circulation, *Int. J. Climate*, 3881–3899, doi:10.1002/joc.3948, 2014.
- Pielke Sr., R. A.: Influence of the spatial distribution of vegetation and soils on the prediction of cumulus convective rainfall, *Rev. Geophys.*, 39, 151–177, doi:10.1029/1999RG000072, 2001.
- Pielke Sr., R. A., Pitman, A., Niyogi, D., Mahmood, R., McAlpine, C., Hossain, F., Goldewijk, K. K., Nair, U., Betts, R., Fall, S., Reichstein, M., Kabat, P., and de Noblet, N.: Land use/land cover changes and climate: modeling analysis and observational evidence, *WIREs Climate Change*, 2, 828–850, doi:10.1002/wcc.144, 2011.
- Pitman, A. J., de Noblet-Ducoudré, N., Avila, F. B., Alexander, L. V., Boisier, J.-P., Brovkin, V., Delire, C., Cruz, F., Donat, M. G., Gayler, V., van den Hurk, B., Reick, C., and Voldoire, A.: Effects of land cover change on temperature and rainfall extremes in multi-model ensemble simulations, *Earth Syst. Dynam.*, 3, 213–231, doi:10.5194/esd-3-213-2012, 2012.
- Rajeevan, M., Bhate, J., Kale, J. D., and Lal, B.: High resolution daily gridded rainfall data for Indian region: analysis of break and active monsoon spells, *Curr. Sci. India*, 9, 296–306, 2006.
- Rajeevan, M., Bhate, J., and Jaswal, A. K.: Analysis of variability and trends of extreme rainfall events over India using 104 years of gridded daily rainfall data, *Geophys. Res. Lett.*, 35, L18707, doi:10.1029/2008GL035143, 2008.
- Ramankutty, N. and Foley, J. A.: Estimating historical changes in global land cover: croplands from 1700 to 1992, *Global Biogeochem. Cy.*, 13, 997–1027, doi:10.1029/1999GB900046, 1999.

- 1285 Ramankutty, N., Evan, A. T., Monfreda, C., and Foley, J. A.: Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000, *Global Biogeochem. Cy.*, 22, GB1003, doi:10.1029/2007GB002952, 2008.
- Ratnam, J. V., Giorgi, F., Kaginalkar, A., and Cozzini, S.: Simulation of the Indian monsoon using the RegCM3-ROMS regional coupled model, *Clim. Dynam.*, 33, 119–139, doi:10.1007/s00382-008-0433-3, 2009.
- 1290 Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., and Wang, W.: An improved in situ and satellite SST analysis for climate, *J. Climate*, 15, 1609–1625, doi:10.1175/1520-0442(2002)015<1609:AIISAS>2.0.CO;2, 2002.
- Roxy, K. M., Kapoor, R., Terray, P., and Masson, S.: The curious case of Indian Ocean warming, *J. Climate*, 27, 8501–8509, doi:10.1175/JCLI-D-14-00471.1, 2014.
- 1295 [Roxy, K. M., Kapoor, R., Terray, P., Murtugudde, R., Ashok, K., and Goswami, B. N.: Drying of Indian subcontinent by rapid Indian Ocean warming and a weakening land-sea thermal gradient, *Nat. Commun.*, 6, doi:10.1038/ncomms8423, 2015.](#)
- Saeed, F., Hagemann, S., and Jacob, D.: Impact of irrigation on the South Asian summer monsoon, *Geophys. Res. Lett.*, 36, L20711, doi:10.1029/2009GL040625, 2009.
- 1300 Saha, S. K., Halder, S., Kumar, K. K., and Goswami, B. N.: Pre-onset land surface processes and “internal” interannual variabilities of the Indian summer monsoon, *Clim. Dynam.*, 36, 2077–2089, doi:10.1029/2011JD017291, 2011.
- Saha, S. K., Halder, S., Rao, A. S., and Goswami, B. N.: Modulation of ISOs by land–atmosphere feedback and contribution to the interannual variability of Indian summer monsoon, *J. Geophys. Res.*, 117, D1301, doi:10.1029/2011JD017291, 2012.
- 1305 Sampaio, G., Nobre, C. A., Costa, M. H., Satyamurty, P., Soares-Filho, B. S., and Cardoso, M.: Regional climate change over eastern Amazonia caused by pasture and soybean cropland expansion, *Geophys. Res. Lett.*, 34, L17709, doi:10.1029/2007GL030612, 2007.
- Sen Roy, S., Mahmood, R., Niyogi, D., Lei, M., Foster, S. A., Hubbard, K. G., Douglas, E., and Pielke Sr., R. A.: Impacts of the agricultural Green Revolution induced land use changes on air temperatures in India, *J. Geophys. Res.*, 112, D21108, doi:10.1029/2007JD008834, 2007.
- 1310 Sen Roy, S., Mahmood, R., Quintanar, A. I., and Gonzalez, A.: Impacts of irrigation on dry season precipitation in India, *Theor. Appl. Climatol.*, 104, 193–207, doi:10.1007/s00704-010-0338-z, 2011.

- Shkol'nik, I., Meleshko, V., Efimov, S., and Stafeeva, E.: Changes in climate extremes on the territory of Siberia by the middle of the 21st century: an ensemble forecast based on the MGO Regional Climate Model, *Russ. Meteorol. Hydrol.*, 37, 71–84, 2012.
- Seneviratne, S., Nicholls, N., Easterling, D., Goodess, C. M., Kanae, S., Kossin, J., Luo, Y., Marengo, J., McInnes, K., Rahimi, M., Reichstein, M., Sorteberg, A., Vera, C., and Zhang, X.: Changes in climate extremes and their impacts on the natural physical environment, in: *SREX: Special Report on Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*, edited by: Field, C. B., Barros, V., Stocker, T. F., Qin, D., Dokken, D. J., Ebi, K. L., Mastrandrea, M. D., Mach, K. J., Plattner, G.-K., Allen, S. K., Tignor, M., and Midgley, P. M., A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC), Cambridge University Press, Cambridge, UK and New York, NY, USA, 109–230, 2012.
- Sheikh, M. M., Manzoor, N., Ashraf, J., Adnan, M., Collins, D., Hameed, S., Manton, M. J., Ahmed, A. U., Baidya, S. K., Borgaonkar, H. P., Islam, N., Jayasinghearachchi, D., Kothawale, D. R., Premalal, K. H. M. S., Revadekar, J. V., and Shrestha, M. L.: Trends in extreme daily rainfall and temperature indices over South Asia, *Int. J. Climate*, 35, 1625–1637, doi:10.1002/joc.4081, 2014.
- Singh, D., Tsiang, M., Rajaratnam, B., and Diffenbaugh, N. S.: Observed changes in extreme wet and dry spells during the South Asian summer monsoon season, *Nature Climate Change*, 4, 456–461, doi:10.1038/nclimate2208, 2014.
- Snyder, P. K.: The influence of tropical deforestation on the Northern Hemisphere climate by atmospheric teleconnections, *Earth Interact.*, 14, 1–32, doi:10.1175/2010EI280.1, 2010.
- Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L. (Eds.): *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, New York, 996 pp., 2007.
- Srinivasan, J. and Smith, G. L.: The role of heat fluxes and moist static energy in tropical convergence zones, *Mon. Weather Rev.*, 124, 2089–2099, doi:10.1175/1520-0493(1996)124<2089:TROHFA>2.0.CO;2, 1996.
- Srivastava, A. K., Rajeevan, M., and Kshirsagar, S. R.: Development of a high resolution daily gridded temperature data set (1969–2005) for the Indian region, *Atmos. Sci. Lett.*, 10, 249–254, doi:10.1002/asl.232, 2009.
- Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M. (Eds.): *IPCC, 2013: Climate Change 2013: The Physical Science*

Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK and New York, NY, USA, 1535 pp., 2013.

Stockli, R., Lawrence, D. M., Niu, G. Y., Oleson, K. W., Thornton, P. E., Yang, Z. L., Bonan, G. B., Denning, A. S., and Running, S. W.: Use of FLUXNET in the Community Land Model development, *J. Geophys. Res.*, 113, G01025, doi:10.1029/2007JG000562, 2008.

Sud, Y., Shukla, J., and Mintz, Y.: Influence of land surface roughness on atmospheric circulation and precipitation: a sensitivity study with a general circulation model, *J. Appl. Meteorol.*, 27, 1036–1054, doi:10.1175/1520-0450(1988)027<1036:IOLSRO>2.0.CO;2, 1998.

Takata, K., Saito, K., and Yasunari, T.: Changes in the Asian monsoon climate during 1700–1850 induced by preindustrial cultivation, *P. Natl. Acad. Sci. USA*, 106, 9586–9589, doi:10.1073/pnas.0807346106, 2009.

~~Tawfik, A. and Steiner, A. L.: The role of soil ice in land-atmosphere coupling over the United States: a soil moisture-precipitation-winter feedback mechanism, *J. Geophys. Res.*, 116, D02113, doi:, 2011.~~

Tian, H., Banger, K., Bo, T., and Dadhwal, V. K.: History of land use in India during 1880–2010: large-scale land transformations reconstructed from satellite data and historical archives, *Global Planet. Change*, 121, 76–88, doi:10.1016/j.gloplacha.2014.07.005, 2014.

Trenberth, K. E.: Atmospheric moisture residence times and cycling: implications for rainfall rates and climate change, *Climatic Change*, 39, 667–694, 1998.

Trenberth, K. E.: Changes in precipitation with climate change, *Clim. Res.*, 47, 123–138, doi:10.3354/cr00953, 2011.

Trenberth, K. E., Dai, A., Rasmussen, R. M., and Parsons, D. B.: The changing character of precipitation, *B. Am. Meteorol. Soc.*, 84, 1205–1217, doi:10.1175/BAMS-84-9-1205, 2003.

Tuinenburg, O. A., Hutjes, R. W. A., Jacobs, C. M. J., and Kabat, P.: Diagnosis of local land-atmosphere feedbacks in India, *J. Climate*, 24, 251–266, doi:10.1175/2010JCLI3779.1, 2011.

Wang, B., Ding, Q., and Joseph, P. V.: Objective definition of the Indian Summer Monsoon onset, *J. Climate*, 22, 3303–3316, doi:10.1175/2008JCLI2675.1, 2009.

Willett, K. M., Jones, P. D., Thorne, P. W., and Gillett, N. P.: A comparison of large scale changes in surface humidity over land in observations and CMIP3 general circulation models, *Environ. Res. Lett.*, 5, 025210, doi:10.1088/1748-9326/5/2/025210, 2010.

- Xavier, P. K., Marzin, C., and Goswami, B. N.: An objective definition of the Indian summer monsoon season and a new perspective on the ENSO-monsoon relationship, *Q. J. Roy. Meteor. Soc.*, 133, 749–764, doi:10.1002/qj.45, 2007.
- Xue, Y., Janjic, Z., Dudhia, J., Vasic, R., and De Sales, F.: A review on regional dynamical downscaling on intraseasonal to seasonal simulation/prediction and major factors that affect downscaling ability, *Atmos. Res.*, 147–148, 68–85, 2014.
- Yatagai, A., Kamiguchi, K., Arakawa, O., Hamada, A., Yasutomi, N., and Kito, A.: APHRODITE: constructing a long-term daily gridded precipitation dataset for ~~asia~~ Asia based on a dense network of rain gauges, *B. Am. Meteorol. Soc.*, 93, 1401–1415, doi:10.1175/BAMS-D-11-00122.1, 2012.
- Zeng, X. and Beljaars, A.: A prognostic scheme of sea surface skin temperature for modeling and data assimilation, *Geophys. Res. Lett.*, 32, L14605, doi:10.1029/2005GL023030, 2005.
- Zeng, X., Shaikh, M., Dai, Y., Dickinson, R. E., and Myneni, R.: Coupling of the Common Land Model to the NCAR Community Climate Model, *J. Climate*, 15, 1832–1854, 2002.

Table 1. Experimental set-up for LULCC based simulations with RegCM4.

Experiment name	Lateral boundary conditions	Sea surface temperature	Year of fixed LULC	Period of simulation
PLC	01 Nov 1981–31 Dec 2008	Observed (1981–2008)	2005	27 years
HLC	01 Nov 1981–31 Dec 2008	Observed (1981–2008)	1950	27 years
PLCS	01 Nov 1981–31 Dec 2008	De-trended (1981–2008)	2005	27 years
HLCS	01 Nov 1981–31 Dec 2008	De-trended (1981–2008)	1950	27 years




Figure 1. PFT distribution of ~~crop and~~ forest and crop cover (in %) used as fixed lower boundary condition in the model experiments. (a) Forest and (b) crop of the year 1950 (HLC). (c) Forest and (d) crop of the year 2005 (PLC). Differences (PLC-HLC) in (e) forest and (f) crop cover.




Figure 2. Time series of number of observed moderate rainfall events over CI and total rainfall in JJAS (in mm; 1951–2007) from ~~moderate events in IMD~~ APHRODITE rainfall data. **(a)** Moderate rainfall events and **(b)** total amount of rainfall in moderate category. Black dotted line represents the linear trend.




Figure 3. Observed trend (1969–2005) in seasonal (JAS) 2 m air temperature from IMD (in $^{\circ}\text{C decade}^{-1}$). Trend in **(a)** seasonal average of daily mean, **(b)** 90th percentile of daily mean, **(c)** seasonal average of daily maximum, and **(d)** 90th percentile of daily maximum temperature. Observed trend (1951–2005) in **(e)** seasonal average of daily mean and **(f)** 90th percentile of daily mean 2 m air temperature from 20CR reanalysis data (in $^{\circ}\text{C decade}^{-1}$). Green contour encloses the region where trends are significant at 90 % confidence level.




Figure 4. (a) Seasonal (JJAS) averaged rainfall from GPCP (in mm day⁻¹, ~~1979–2008~~1982–2008). (b) Seasonal averaged rainfall (in mm day⁻¹) in PLC experiment with RegCM4 (1982–2008). (c) Seasonal averaged rainfall in PLCS experiment (in mm day⁻¹, 1982–2008). (d) Seasonal averaged rainfall based on ~~IMD–APHRODITE~~ data (in mm day⁻¹, ~~1979–2007~~1982–2007). (e) Seasonal averaged rainfall ~~based on APHRODITE data (in mm day⁻¹, 1979–2007).~~ (f) ~~Seasonal averaged rainfall~~ only over land in PLC experiment (in mm day⁻¹). (f) ~~Representation of orography in RegCM4.~~ Units are in m.




Figure 5. Seasonal (JJAS) averaged wind at 10 m (in m s^{-1} , 1982–2008). **(a)** Climatological mean (PLC experiment) and **(b)** difference (PLC-HLC). The shaded color depicts magnitude and arrows show the direction. Green contour shows differences significant at 90 % confidence level.




Figure 6. (a) Difference (PLC-HLC experiment) in climatological mean seasonal rainfall (in cm, 1982–2008) shown in shaded color. (b) Same as in (a) but for PLCS-HLCS experiments. Dashed (solid) black contours show the decrease (increase) in velocity potential analog (or the divergent component) of vertically integrated moisture flux qV (from surface–300 hPa). The contour interval is in $1 \times 10^6 \text{ kg s}^{-1}$. Green contour shows differences significant at the 90 % confidence level.




Figure 7. Difference (PLC-HLC) in (a) number of moderate rainfall events during JJAS and (b) total amount of moderate rainfall (in mm day⁻¹, 1982–2008). Green contour shows differences significant at the 90 % confidence level.




Figure 8. Difference (PLC-HLC) in JJAS averaged 2 m air temperature (in °C, 1982–2008), for (a) daily mean, and (b) daily maximum temperature. Green contour shows significance at 90 % confidence levels.




Figure 9. Difference (PLC-HLC) in daily variance of 2 m air temperature in JJAS (in $^{\circ}\text{C}^2$, 1982–2008) for (a) daily mean, and (b) daily maximum temperature. Green contour shows significance at 90% confidence level. Difference (PLC-HLC) in the 90th percentile of daily 2 m air temperature in JJAS (in $^{\circ}\text{C}$, 1982–2008), for (c) daily mean and (d) daily maximum temperature.

Figure 10. (a) Seasonal (JJAS) averaged surface pressure in PLC experiment (in black contours) and its difference (PLC-HLC) in shaded color (in hPa, 1982–2008). Green contour shows differences significant at 90 % confidence level. **(b–i)** Show similar differences as in **(a)** but for surface soil moisture (0–10 cm, in mm), 2 m specific humidity (in g kg^{-1}), surface albedo (unitless), total cloud cover (in %), surface net radiation (in W m^{-2}), surface sensible heat flux (in W m^{-2}), surface latent heat flux (in W m^{-2}), respectively. **(i)** Shows only the difference (PLC-HLC) in sum of transpiration and evaporation of canopy-intercepted water (in mm day^{-1}). In **(d)** and **(i)**, only the differences significant at 90 % confidence level are shaded.




Figure 11. Difference (PLCS-HLCS) in seasonal (JJAS) averaged 2m air temperatures (in °C, 1982–2008) for (a) daily mean, and (b) daily maximum temperature. Green contour shows differences significant at 90 % confidence level. Difference (PLCS-HLCS) in the 90th percentile of daily 2m air temperature in JJAS (in °C, 1982–2008), for (c) daily mean, and (d) daily maximum temperature.




Figure 12. (a) Seasonal average (JJAS) of vertically integrated moist static energy (VIMSE, surface–500 hPa) in PLC experiment (in $1 \times 10^4 \text{ kJ kg}^{-1}$, 1982–2008). (b) Difference (PLC–HLC) in VIMSE (in kJ kg^{-1} , 1982–2008). Green contour shows differences significant at 90 % confidence level. (c) Difference (PLC–HLC) in CI averaged moist static energy (MSE, in red) and dry static energy (DSE, in blue) in units of kJ kg^{-1} .