

**Rainfall erosivity in catchments contaminated with fallout from the Fukushima Dai-ichi nuclear power plant accident**

**J.P. Laceby<sup>1</sup>, C. Chartin<sup>2</sup>, O.Evrard<sup>1</sup>, Y.Onda<sup>3</sup>, L. Garcia-Sanchez<sup>4</sup>, O. Cerdan<sup>5</sup>**

<sup>1</sup>Laboratoire des Sciences du Climat et de l'Environnement (LSCE), Unité Mixte de Recherche 8212 (CEA-CNRS-UVSQ/IPSL), Université Paris-Saclay, Gif-sur-Yvette, France

<sup>2</sup> Georges Lemaître Centre for Earth and Climate Research, Earth and Life Institute, Université catholique de Louvain, Belgium

<sup>3</sup>Graduate School of Life and Environmental Sciences, Center for Research in Isotopes and Environmental Dynamics (CRIED), University of Tsukuba, Tsukuba, Japan

<sup>4</sup>Laboratoire de Biogéochimie, Biodisponibilité et Transferts de Radionucléides, IRSN/PRP-ENV/SERIS/L2BT, Cadarache, France

<sup>5</sup>Bureau de Recherches Géologiques et Minières, Orléans, France

Correspondence to: J.P. Laceby (laceby@lsce.ipsl.fr)

## **Abstract:**

The Fukushima Dai-ichi nuclear power plant (FDNPP) accident in March 2011 resulted in the fallout of significant quantities of radiocesium over the Fukushima region. After reaching the soil surface, radiocesium is quickly bound to fine soil particles. Thereafter, rainfall and snowmelt runoff events transfer particle-bound radiocesium downstream. Characterizing the precipitation regime of the fallout-impacted region is thus important for understanding post-deposition radiocesium dynamics. Accordingly, 10-minute (1995-2015) and daily precipitation data (1977-2015) from 42 meteorological stations within a 100 km radius of the FDNPP were analysed. Monthly rainfall erosivity maps were developed to depict the spatial heterogeneity of rainfall erosivity for catchments entirely contained within this radius. The mean average precipitation in the region surrounding the FDNPP is 1420 mm yr<sup>-1</sup> (SD 235) with a mean rainfall erosivity of 3696 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> (SD 1327). Tropical cyclones contribute 22% of the precipitation (422 mm yr<sup>-1</sup>) and 40% of the rainfall erosivity (1462 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> (SD 637)). The majority of precipitation (60%) and rainfall erosivity (82%) occurs between June and October. At a regional scale, rainfall erosivity increases from the North to the South during July and August, the most erosive months. For the remainder of the year, this gradient occurs mostly from northwest to southeast. Relief features strongly influence the spatial distribution of rainfall erosivity at a smaller scale, with the coastal plains and coastal mountain range having greater rainfall erosivity than the inland, Abukuma river valley. Understanding these patterns, particularly their spatial and temporal (both inter and intra annual) variation, is important for contextualizing soil and particle-bound radiocesium transfers in the Fukushima region. Moreover, understanding the impact of tropical cyclones will be important for managing sediment and sediment-bound contaminant transfers in regions impacted by these events.

**Key Words:** R-Factor, FDNPP, EI<sub>30</sub>, Erosion, Cesium, Tropical Cyclones

**Author Contributions:** J.P.L & L.G-S downloaded and prepared the data, J.P.L and C.C. performed the analyses, all authors were fundamental to drafting and editing the manuscript.

## 1 Introduction

In March 2011, the Great Tohoku earthquake triggered a giant tsunami that resulted in the Fukushima Dai-ichi nuclear power plant (FDNPP) release of the largest amount of radioactive material since Chernobyl (Chino et al., 2011;Thakur et al., 2013). After the decay of short-lived radionuclides (e.g.  $^{131}\text{I}$ ,  $^{110\text{m}}\text{Ag}$ ,  $^{129}\text{Te}$ ), the radionuclide with the most serious short and long term human health concerns is radiocesium ( $t_{1/2}$   $^{134}\text{Cs}$  – 2 yr,  $^{137}\text{Cs}$  – 30yr) (Kitamura et al., 2014;Saito et al., 2015).

Radiocesium is rapidly bound to fine soil particles and remains stored within the top 0-5cm of the soil profile (Tamura, 1964;Sawhiney, 1972;Kato et al., 2012). As radiocesium is adsorbed to soil particles in the upper layer of the soil profile, soil erosion and particle transport processes are considered to be the main processes responsible for transferring radiocesium downstream (Saito and Onda, 2015;Yamashiki et al., 2014;Evrard et al., 2015). Rainfall drives erosion by detaching particles from the soil surface, primarily through the kinetic energy of raindrops (Wischmeier and Smith, 1958;Renard and Freimund, 1994). The potential of rainfall to erode soils is referred to as rainfall erosivity. Quantifying rainfall erosivity is important to determine the climatic vulnerability of a region, like Fukushima, to rainfall-driven soil erosion (Mello et al., 2013).

The Universal Soil Loss Equation (USLE) (Wischmeier and Smith, 1978) and its revised edition (RUSLE) (Renard et al., 1997) are some of the most broadly adopted soil erosion modelling frameworks (Oliveira et al., 2013). The USLE predicts average annual soil loss with six factors pertaining to soil, topography, vegetation, management and rainfall erosivity, known as the R-factor. The R-factor estimates rainfall erosivity as a combined function of event frequency, intensity, and rainfall amount (Wischmeier and Smith, 1958). It sums event-based erosivity individually calculated through multiplying each storm's total kinetic energy with its maximum measured 30-minute rainfall intensity. To incorporate variations in inter-annual rainfall, the R-factor is averaged over long temporal periods (~20 years) (Renard and Freimund, 1994;Wischmeier and Smith, 1978).

When all other factors in the (R)USLE are held constant, there is a proportional relationship between soil loss and the R-factor (Lu and Yu, 2002;Wischmeier and Smith, 1958;Wischmeier and Smith, 1978). The utility of the R-factor is reflected in its global application (McFarlane et al., 1986;Ma et al., 2014;Oliveira et al., 2013;Moore, 1979;Panagos et al., 2015) including Asia

in general (Lee and Heo, 2011;Shamshad et al., 2008;Yu et al., 2001) and Japan in particular (Shiono et al., 2013).

There are limitations with the application of the R-factor and the USLE (Kinnell, 2010). Although the USLE and R-factor were developed from plot scale soil erosion research (Wischmeier and Smith, 1978), they have been applied at the catchment scale (Wilkinson et al., 2009;Kitamura et al., 2014;Belyaev et al., 2005). A debate on the applicability of the USLE within catchment scale modelling frameworks and the universality of the R-factor are beyond the scope of this current research. What is important is that the R-factor and a thorough characterization of the precipitation regime are useful to contextualize local soil erosion, which represents the first step of potential downstream particle-bound contaminant transfers.

Research in the Fukushima region has clearly indicated that precipitation drives downstream radiocesium migration (Evrard et al., 2015). In fact, the majority of soil erosion and concomitant radiocesium migration was reported to coincide with extreme rainfall events (Mouri et al., 2014;Nagao et al., 2013;Onishi et al., 2014). For example, Yamaguchi et al. (2014) reported that most sediment-bound radiocesium migration occurs in flood events that may only occur 1 to 2 times per year. **Tropical cyclones** in this region likely transfer significant quantities of radiocesium from hillslopes to the Pacific Ocean (Evrard et al., 2015).

As improved estimates of rainfall erosivity result in more accurate modelling results (Renard et al., 1991;Lee and Heo, 2011), a comprehensive examination of rainfall erosivity, including the impact of **tropical cyclones** on erosivity, may improve our understanding of sediment and radiocesium dynamics in the Fukushima region. Therefore, the objective of this research is to evaluate the spatial and temporal (monthly and inter-annual) precipitation and rainfall erosivity in general, and from **tropical cyclones** in particular, for catchments contained entirely within a 100 km radius of the FDNPP accident. Rainfall erosivity maps (monthly and annual) are provided for the Fukushima region along with precipitation and rainfall erosivity data for all events selected by the RUSLE criteria, and information summarizing **tropical cyclone** events impacting the area since 2011 as supplementary information.

## 2 Methods

### 2.1 Study area

Most of the terrestrial fallout from the FDNPP accident occurred over the coastal catchments of the Fukushima Prefecture (Fig. 1). Soil contamination  $>25 \text{ kBq kg}^{-1}$  of  $^{134+137}\text{Cs}$  occurred within small catchments ( $< 700 \text{ km}^2$ ) draining to the Pacific Ocean, with the highest contamination levels in the coastal mountain range northwest of the FDNPP. The coastal mountain range is separated from the larger westernmost mountain range by the Abukuma river catchment ( $\sim 5200 \text{ km}^2$ ) extending on a SSW-NNE direction and characterised by a  $^{134+137}\text{C}$  soil contamination  $<25 \text{ kBq kg}^{-1}$ .

The region has a humid, subtropical climate along the Pacific coast (Cwa in the Köppen-Geiger climate classification (Peel et al., 2007)), transitioning to a humid continental climate (Dfa) to the west. The westward continental region has large differences in seasonal temperatures, with hot and humid summers and cold winters. The subtropical coastal region has slightly warmer winters, along with hot humid summers and more precipitation in the summer months compared to relatively dry winters. There is **tropical cyclone** driven precipitation in the Fukushima region that results in high volumes and intensities of precipitation occurring during extreme events (Evrard et al., 2014;Chartin et al., 2013). The Japanese **tropical cyclone** season occurs between May and October with the peak activity occurring between July and September. The mean annual temperature in the Fukushima region is  $11.3^\circ\text{C}$  (**SD 1.7**) ranging from a mean of  $0.0$  (**SD 2.1**) in January to  $23.4$  (**SD 1.4**) in August.

### 2.2 Meteorological Stations

All precipitation monitoring stations within a 100 km radius of the FDNPP were analyzed (Fig. 1). Fourteen stations within this 100 km **radius** were omitted due to an operation period of less than 7 years ( $n=5$ ) or operation pause during winter months ( $n=9$ ). Up to 21 years were available for the 10-minute (1995–2015) and 39 years for daily precipitation data (1977–2015) for 42 stations. Daily precipitation was also analyzed from three additional long-term stations (including one located within a 115 km radius of the FDNPP) as they had **similar** data available from 1890–2015. The long-term stations were excluded from the **R-factor and daily rainfall calculations**

owing to high levels of missing data (mean 33% daily) and short temporal periods (7 years – 10-minute). All data were downloaded from the Japan Meteorological Agency's (JMA) website (<http://www.jma.go.jp/jma/index.html>).

Daily precipitation data were examined for **breakpoints** and homogeneity with the RHTest program from the Meteorological Service of Canada (Wang et al., 2010). No significant break points required correction. Duan et al. (2015) similarly did not report significant break points for the JMA daily precipitation data. No homogenization analyses were performed on the 10-minute precipitation data owing the difficulties in assessing heterogeneity in 10-minute precipitation data and the potential to overcorrect the data. Shiono et al. (2013) similarly did not correct 10-minute JMA data. For the daily data, there was an average of only 0.23% of missing data (**SD** 0.20%, maximum 0.94%) compared to 0.30% (**SD** 0.20%, maximum 0.78%) for the 10-minute data.

### 2.3 R-factor calculations

The Rainfall Intensity Summarisation Tool (RIST) software (USDA, 2013) was used to calculate the R-factor, which is a product of the kinetic energy of an event (E) and its maximum 30-minute intensity ( $I_{30}$ ) (Renard and Freimund, 1994):

$$R = \frac{1}{n} \sum_{j=1}^n \sum_{k=1}^{m_j} (EI_{30})_k \quad (1)$$

where R equals the annual average rainfall erosivity in  $\text{MJ mm ha}^{-1} \text{ hr}^{-1} \text{ yr}^{-1}$ ,  $n$  represents the number of years of data utilized,  $m_j$  is the number of erosive events in year  $j$ , and  $EI_{30}$  was calculated as:

$$EI_{30} = \left( \sum_{r=1}^o e_r v_r \right) I_{30} \quad (2)$$

with  $e_r$  representing **rainfall energy per unit depth of rainfall** in  $\text{MJ ha}^{-1} \text{ mm}^{-1}$ ,  $v_r$  is the volume of rainfall (mm) during a given time interval ( $r$ ), and  $I_{30}$  is the maximum rainfall intensity over a 30 minute period of the rainfall event ( $\text{mm h}^{-1}$ ). For each time interval,  $e_r$  was calculated as:

$$e_r = 0.29[1 - 0.72^{(-0.05i_r)}] \quad (3)$$

with  $i_r$  being rainfall intensity ( $\text{mm h}^{-1}$ )(Renard and Freimund, 1994).

To calculate the annual and monthly R-factors, erosive rainfall events were summed for each monitoring station for the considered period (months or year). Two criteria within the RIST software were used to define an erosive event: 1) cumulative rainfall of a given event is  $>12.7\text{mm}$ ; and 2) accumulations  $<1.27\text{mm}$  over a 6 hour period separates events into two different periods.

To consider the impact of snowfall, precipitation was removed from R-factor calculations when the temperature dropped below  $0^{\circ}\text{C}$  for 31 stations (Meusburger et al., 2012). When removing precipitation that may have occurred as snowfall, there was not a significant difference in the R-factor (t-test  $p$ : 0.469) with a mean difference of  $26\text{ MJ mm ha}^{-1}\text{ hr}^{-1}\text{ yr}^{-1}$  (SD 35). Owing to the limited impact of potential snowfall on R-factor calculations, 11 stations that did not record temperature were included in the analyses.

To examine the impact of tropical cyclones on precipitation and rainfall erosivity, the best track data of all tropical cyclones from the JMA Tokyo-Typhoon Center (JMA, 2016) were cross-referenced with regional events that occurred from 1995-2015. Each cyclone path and summary information was also reviewed in the Wikipedia Pacific Typhoon database (Wikipedia, 2016). Furthermore, news reports on tropical cyclones in the region (from Google news searches) were also examined to provide complimentary information. A databased was then created that allowed for the quantification of the precipitation and rainfall erosivity that occurred during tropical cyclone related events. In the Northwest Pacific Ocean, tropical cyclones are referred to as tropical depressions, tropical storms and typhoons depending on the sustained wind speed. Tropical cyclones are non-frontal synoptic scale systems that have organized convection along with cyclonic surface wind circulation. Typhoons, tropical storms and tropical depressions are all referred to as tropical cyclones in the remainder of this manuscript.

## 2.4 R-factor spatial interpolation

Annual and monthly R-factor data were used to derive maps of annual and monthly rainfall erosivity at a resolution of 250 m over the eastern part of the Fukushima Prefecture. A regression approach, based on relationships established between the R-factor and spatially distributed covariates, was used to produce these maps rather than interpolation by kriging in order to identify the environmental covariates that are driving the spatial distribution of rainfall erosivity.

The covariates used for the spatial interpolation of annual and monthly R-factors over the study area are described below:

- Climatic data (i.e. monthly mean precipitation and mean temperature) from the WorldClim database reported for the period 1950–2000 and with a 30 arc-seconds resolution (equivalent to 1 km resolution) (Hijmans et al., 2005) were used for the monthly models (Panagos et al., 2015). From the monthly data, annual mean precipitation and mean temperature were derived to be included as covariates in the annual R-factor model. These covariates are expected to impact the spatial distribution of rainfall erosivity.
- Based on their known influence on spatial patterns and intensities of rainfall, different morphometric attributes have been included as covariates in the mapping procedure, as proposed by Daly et al. (2002). Precipitation varies strongly with elevation (Barry and Chorley, 2009). Conceptually, relief features can induce different precipitation-elevation gradients depending partially on their ability to block and uplift moisture-bearing air (Daly, 2002). In mountainous areas, the steepest features oriented perpendicular to the air flow are more likely to produce a greater elevation-precipitation gradient than gentle sloping areas parallel to the air flux, such as coastal plains. Moreover, hillslopes experience different climate regimes depending on their orientation, relative to air currents at large scales (i.e., leeward and windward sides of mountains), and solar radiation. Areas located at similar elevations may thus have different precipitation intensities.

Accordingly, a 90m-resolution Digital Elevation Model (DEM) was obtained from Shuttle Radar Topography Mission (SRTM) of the NASA (Jarvis et al., 2008). Considering the local relief in the fallout affected region (Fig. 1), the DEM was smoothed with a Gaussian filter to two wavelengths of 5 and 20 km. From the 20 km wavelength smoothed DEM, slope and aspect were derived. The slope layer was used in the modelling procedure to account for potential impact of large relief features on moisture-bearing air. The aspect layer derived from the same 20 km wavelength smoothed DEM was used to delineate the opposite sides of mountains and represent leeward and windward sides. Then, this aspect layer was inserted in the model as a categorical covariate (i.e., a 2 values layer, one for



each leeward and windward ‘facets’). From the 5 km wavelength smoothed DEM, the easting and northing were computed to represent the degree to which aspect is close to the east and the north, respectively (Zar, 2010). These two layers were used as covariates in the modelling procedure to account for slope orientation at a smaller scale than entire side of a mountains range at the coarser scale.

- Distance to the Pacific Ocean coast (DC = distance to the coast) was calculated for each pixel by using the Euclidean distance tool from ArcGis 10 (ESRI, 2011) as the proximity of the ocean, a major source of moisture, may influence the spatial distribution of precipitation.

Before the fitting of the models, the continuous covariates presented above were rescaled at a 250 m resolution in ArcGIS10 (ESRI, 2011) with bilinear interpolation, that determined the output value of a cell based on a weighted distance average of the four nearest input cell centres. Then, the spatial variation of rainfall erosivity in the study region was modelled with Generalized Additive Models (GAM; Hastie and Tibshirani (1986)), implemented in the mgcv package in R (Wood, 2001). The GAM regression technique is a generalization of linear regression models (GLM) in which the coefficients may be a set of smoothing functions. GAMs consider the non-linearity that may exist between the target variable (Y) and covariates (X) providing more flexibility to the model fitting than a GLM.

As for GLMs, GAMs specify a distribution for the response variable Y and use a link function g relating the conditional mean  $\mu(Y)$  of the response variable to an additive function of the selected predictors:

$$g[\mu(Y)] = \alpha + f_1 X_1 + f_2 X_2 + \dots + f_p X_p \quad (4)$$

where Y is the response variable,  $X_1, X_2, \dots, X_p$  represent the covariates and the  $f_i$ 's the smooth (non-parametric) functions.

Based on monthly and annual mean values for each of the 42 stations, 13 GAMs were fitted to spatially model the R-factor over the study area (i.e. one model per month and one model for the year). A Gaussian distribution model incorporated the conditional mean  $\mu(Y)$  and, due to the predominant logarithmic distribution of the monthly and annual data, a log-linear link function  $g(\mu) = \log(\mu)$  was implemented. The smoothing functions of the models were built using

regression splines fitted by penalized Maximum Likelihood to avoid over-fitting (Wood, 2001). An extra penalty was added to each smoothing term so that each could potentially be set to zero during the fitting process, especially in case of multi-collinearity or multi-concurvity. The interaction of geographical coordinates was then added to each model (as a two-dimensional spline on latitude and longitude) to incorporate spatial trends of the target variable at the regional scale.

A model including all the covariates was first developed (the ‘full’ model), based on each of the total datasets (13 datasets of  $n=42$ ). Next, a backward stepwise procedure was applied to select the appropriate covariates where each of the covariates were sequentially removed from the full model (Poggio et al., 2013; de Brogniez et al., 2015). The difference ( $\Delta AIC$ ) between AIC (Akaike’s Information Criterion; Akaike (1974)) was calculated for the ‘full model’ and compared with AIC obtained when removing sequentially each of the covariates from the model in order to evaluate the influence of each covariate on the overall capabilities of model prediction.

Explained variance was calculated during the procedure to assess the evolution of ‘fitting performance’ and to support the decision to keep or remove a covariate into the final model. Once the final model was obtained, a Bayesian approach was used to compute standard errors for the predictions as proposed in the mgcv package. A leave-one-out cross-validation (LOOCV) procedure was applied to each fitted model. Predicted and observed values were compared with the calculation of the Root Mean Square Errors (RMSE) and the coefficient of determination ( $R^2$ ). Finally, maps were delineated according to the limits of the target area, i.e. entire hydrological basins contained within the 100 km radius of around the FDNPP.

### 3 Results

#### 3.1 Precipitation

Mean annual precipitation ranged from 1098 mm yr<sup>-1</sup> at station 281 to 2201 mm yr<sup>-1</sup> at station 1116 (Table 1; Fig 2A). Mean annual precipitation was 1420 mm yr<sup>-1</sup> (SD 235 mm yr<sup>-1</sup>)(Fig. 2A) with a coefficient of variation of 17%. Between 1977 and 2015, regional mean annual precipitation ranged from 869 mm in 1984 to 1844 mm in 2006 (Fig. 3A) with a coefficient of variation of 15%. This inter-annual variation in precipitation was maintained over a longer period

(Fig. 4) with a mean annual precipitation for the long-term stations (126 yr) being 1159 mm yr<sup>-1</sup> (SD 187mm) with a 16% coefficient of variation. The 5 year running average for these long-term stations demonstrates that the region experiences prolonged wet and dry periods that last approximately 5 – 10 years. Of note, these long-term stations plot near or below one standard deviation on the annual precipitation mean for all of the stations, and therefore they may not be indicative of all precipitation trends in the region.

The majority of precipitation (60%) occurs between June and October (Fig. 5A) coinciding with a wet season that includes tropical cyclones. In fact, 39% of the precipitation occurred between July and September alone, compared to only 22% between March and May, and 17% between November and February. The monthly precipitation also varied spatially. For example, the mean monthly maximum precipitation was 318 mm m<sup>-1</sup> at site 1116 in July compared to 256 mm m<sup>-1</sup> for site 281 in July.

The highest daily maximum precipitation recorded was 607 mm in 1998 at Site 326, compared to the lowest daily maximum precipitation of 38 mm in 1984 at site 294 (Fig. 6A). The mean daily maximum precipitation for all the stations and every year was 115 mm (SD 27 mm). The highest mean daily maximum precipitation for all the stations (169 mm) was recorded in 2011 compared to a minimum of 62 mm in 1984. Although the maximum daily precipitation is not an optimal indicator of precipitation depth owing to the probability of events occurring over multiple days, the year of the FDNPP accident (2011), had the highest daily maximum precipitation as a result of Typhoon Roke.

### 3.2 Rainfall Erosivity

The annual R-factor ranged from 1972 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> at station 290 to 8274 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> at station 326 (Table 1 - Fig. 2B). The mean R-factor for the 42 stations was 3696 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> (SD 1327) with a coefficient of variation of 36%. The coefficient of variation was double the corresponding coefficient of variation for annual average precipitation indicative of more spatial variability.

The regional mean annual R-factor ranged from a minimum of 1866 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> in 1996 to a maximum of 7159 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> in 1998 (Fig. 3C). The annual R-factor standard

deviation was 1252 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> with a coefficient of variation of 34%. The coefficient of variation again was more than double **that of** the annual precipitation.

The highest monthly R-factors occurred in the summer months and early fall (Fig. 5B) coinciding with the wet summer and the peak **tropical cyclone** season (July-September). In fact, 82% of the rainfall erosivity occurred between June and October in the Fukushima region, an increase of 22% compared to the amount of precipitation during this period. In particular, 64% of the rainfall erosivity occurs between July and September. Winter (November-February) only contributes 6% of the total annual R-factor with spring (March – May) contributing the remaining 12%.

### **3.3 Spatial Distribution of Rainfall Erosivity**

The **variance** explained by the final erosivity models produced by the calibration procedure (from the total dataset of n=42) varied from 82.5% for September to 97.7% for April, and 95.5% for the annual R-factor model (Table 2). Predicted values produced with the LOOCV procedure for the 42 stations showed similar distributions as the observed values (Table 2) for the year and for the months between October and June. For the months between July and September, which includes the **tropical cyclone** season, the standard deviations of predicted values were smaller than for the observed values. For mean observed values ranging from 778.7 to 799.7 MJ mm ha<sup>-1</sup> h<sup>-1</sup> m<sup>-1</sup> between July and September, the RMSE varied between 165.3 and 313.6 MJ mm ha<sup>-1</sup> h<sup>-1</sup> m<sup>-1</sup> with a R<sup>2</sup> between 0.56 and 0.67. The model fitted for August demonstrated a bias by underestimating R-factor (mean observed values = 778.7 MJ mm ha<sup>-1</sup> h<sup>-1</sup> m<sup>-1</sup> and mean predicted values = 755.2 MJ mm ha<sup>-1</sup> h<sup>-1</sup> m<sup>-1</sup>).

For the months between October and June, and for the year, the differences between mean observed and predicted values remained relatively low relative to corresponding observed values and their variability. For these periods, the R<sup>2</sup> produced by the validation procedure varied between 0.62 for April to 0.86 in October, with a R<sup>2</sup> of 0.77 for the annual model. **In general, the annual map and the months with higher erosivity had lower R<sup>2</sup> values. Contrarily, months with lower erosivity often had higher R<sup>2</sup>. This is possibly due to localized impacts of high precipitation events.**

For all of the monthly models, the categorical variable ‘facets’, was selected as one of the most influential explanatory covariates, whereas facets were less influential in the annual model (Table

3). The selection of ‘facets’ highlights how opposite sides of the mountain ranges in this region have different rainfall erosivity regimes.

For 10 of the 13 models, mean precipitation, for the considered period, was selected as a significant explanatory covariate. Mean temperature was selected by the procedure for the three models for which mean precipitation was not retained as significant covariates (i.e., February, November and December). Mean temperature and mean precipitation was selected for the annual R-factor model. According to the  $\Delta AIC$ , mean precipitation and mean temperature capture the most R-factor spatial variability within the final models (Table 3). Easting and/or northing were selected in all the other models, except July and September, accounting for the influence of hillslope aspects at smaller scales than entire mountain sides. The selection of elevation, slope and/or the distance to the coast as significant explanatory variables in some of the final models varied.

Although elevation, slope and distance to the coast may influence precipitation patterns and intensities, their non-consistent selection in the final models does not mean that they potentially do not influence rainfall erosivity. Covariates have the potential not to be retained in the final model when there may be multi-collinearity or multi-concurvity between covariates. The non-selection of covariates signifies that other covariates likely have a similar, though stronger, mathematical relationships with the target variable. To achieve parsimony, only select variables from the original set of covariates are included in the final model.

The annual R-factor has a predominantly positive gradient from North North West to South South East (Fig. 7A) and is impacted by the relief of the region. The coastal plain receives lower rainfall erosivity than the adjacent coastal mountain range, both of which have greater rainfall erosivity than the inland Abukuma river valley. The South South East oriented side of the westernmost mountain range bordering the Abukuma valley has lower rainfall erosivity than the easternmost side of this valley. The coefficient of variation (CV) of modelling errors is rather low, predominantly under 10% in the study area (Fig 7B).

In July and August, the most erosive months of the year, rainfall erosivity increases mainly from the North to the South of the region, while during the rest of the year, this gradient occurs mostly from North West to South East. The spatial patterns relative to the relief features are also observed in the monthly rainfall erosivity distributions with the coastal plains and coastal

mountain range receiving higher rainfall erosivity than the parallel Abukuma river valley. The maps of errors associated with each monthly R-factor models are available as supplementary information (Fig. S1). The coastal plain and coastal mountainous ranges, including the FDNPP and a large part of the contamination plume with high levels of radiocesium contamination, have greater rainfall erosivity than the less contaminated Abukuma River valley.

### 3.4 Tropical Cyclones

In the Fukushima region, tropical cyclones contributed 22% ( $331 \text{ mm y}^{-1}$ ) of the total annual precipitation (Fig 3A). This contribution varies from only 5% ( $68 \text{ mm y}^{-1}$ ) in 2003 to 35% ( $639 \text{ mm y}^{-1}$ ) in 1998. In September, 60% of the precipitation is derived from tropical cyclone related events ( $119 \text{ mm m}^{-1}$ ) compared to 43% in October ( $64 \text{ mm m}^{-1}$ ), 28% in August ( $49 \text{ mm m}^{-1}$ ) and 25% in July ( $47 \text{ mm m}^{-1}$ ). In the remainder of year there is less than a  $15 \text{ mm m}^{-1}$  contribution from tropical cyclones (<11% of the month precipitation totals)(Fig 5A).

Although tropical cyclones only contribute 22% of the annual precipitation in the region, they contribute 40% of the annual rainfall erosivity ( $1462 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ yr}^{-1}$ ). This contribution varies from 6% ( $142 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ yr}^{-1}$ ) in 1995 to 64% in 1998 ( $4547 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ yr}^{-1}$ )(Fig 3B). In September, 75% of the rainfall erosivity ( $589 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ m}^{-1}$ ) is derived from tropical cyclone activity, followed by 57% in October ( $208 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ m}^{-1}$ ), 41% in August ( $321 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ m}^{-1}$ ), 32% in July ( $257 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ m}^{-1}$ ), and 19% in June ( $59 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ m}^{-1}$ )(Fig 5B). Tropical cyclones, which only occurred on average 3.9 times per year (SD 2.0) contributed 40% of the rainfall erosivity in the Fukushima region.

## 4 Discussion

### 4.1 Precipitation

The location and intensity of annual tropical cyclones influence the spatial and temporal (intra- and inter-annual) variation in precipitation in the Fukushima region. The majority of precipitation occurs in the summer months, the result of the high humidity and seasonal tropical cyclone activity. Long-term climatic oscillations (e.g., El Niño) also influence the precipitation patterns, illustrated by the 5–10 year running average reported in the long-term analyses (Fig. 4). Duan et

al. (2015) reported periodic variations in extreme precipitation though at frequencies of 2-3 years and 5-13 years.

The average precipitation in the Fukushima region ( $1420 \text{ mm yr}^{-1}$ ) is  $\sim 250 \text{ mm}$  less than the mean annual precipitation in Japan ( $1684 \text{ mm yr}^{-1}$ ) (Kim et al., 2010). Along the western and southeast coast of Japan, annual precipitation may be greater than  $3000 \text{ mm yr}^{-1}$  (Kim et al., 2010). In the southeast of Japan, the elevated annual precipitation is related to an increasingly tropical climate progressing southwards and the increased occurrence of tropical cyclones in these warmer climates with higher precipitation intensities during summer months.

## 4.2 Rainfall Erosivity

Rainfall erosivity varied spatially and temporally (monthly and inter-annually) in the Fukushima region. Rainfall erosivity has similarly varied in other regions of the world, (e.g. the Mediterranean region (Capolongo et al., 2008) and the Middle East (Eltaif et al., 2010)). Although researchers have demonstrated relationships between precipitation and rainfall erosivity (Ma et al., 2014), in the Fukushima region the relationship was positive, though not significant ( $r^2$  0.39). The lack of significant relationship was likely the result of rainfall erosivity varying more than precipitation. There are also possibly decadal variations in rainfall erosivity, similarly to precipitation, that require long-term analyses to investigate.

There have been a range of R-factors reported for Japan and the Fukushima region. In the Fukushima region, between July 17, 2011 and November 18, 2012, Yoshimura et al. (2015) calculated an R-factor of  $3677 \text{ MJ mm ha}^{-1} \text{ h}^{-1}$  (SD 463) for 3 sites near Kawamata town,  $\sim 13 \text{ km}$  from station 1130. Although station 1130 recorded a similar R-factor for this time period ( $3853 \text{ MJ mm ha}^{-1} \text{ h}^{-1}$ ), the R-factor recorded at this station was 40% lower than the regional average for this period ( $6559 \text{ MJ mm ha}^{-1} \text{ h}^{-1}$  (SD 3232)). This demonstrates the potential impact of the spatial heterogeneity on rainfall erosivity and limitations of extrapolating data from one sampling point to the entire region.

Yamaguchi et al. (2014) quantified an average annual R-factor of  $3366 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ yr}^{-1}$  for 23 JMA stations in this region for a 10 year period between 2001 and 2011. For a sensitivity analysis, these authors also modelled a maximum case of  $4217 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ yr}^{-1}$  and a minimum case of  $1986 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ yr}^{-1}$ . The average annual R-factor reported by Yamaguchi

et al. (2014) was 9% lower than that reported for our current research, likely the result of the longer temporal period and the high levels of rainfall erosivity in the late 1990s not being included in the calculations of Yamaguchi et al. (2014). Further, there was only a 120 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> difference between our minimum cases, though our maximum case was 41% higher (7159 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup>) in 1998 than Yamaguchi et al. (2014), and again 33% higher (6288 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup>) than their maximum case in 1999.

Around Japan a variety of R-factors have been reported, for example 5880 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> in the Onga River basin (Fukuoka prefecture - South Japan)(Tran et al., 2011), 4568 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> in the Shiraska River catchment (Aichi prefecture - Central Japan) (Karki and Shibano, 2006), and 2890 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> in the Hirose River catchment (Miyagi prefecture - Northern Japan)(Takeuchi and Ishidaira, 2001). Shiono et al. (2013) found that rainfall erosivity varied from 721 to 35299 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> across Japan with a mean of 5130 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup>. In our current analyses, rainfall erosivity in the Fukushima region was found to be similar to the results of Shiono et al. (2013) who spatially depicted rainfall erosivity in the Fukushima region being between 2500 and 5000 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup>.

Shiono et al. (2013) also demonstrated that rainfall erosivity in the Fukushima region is relatively low compared to the rest of Japan. Similarly to annual precipitation, there is a rainfall erosivity gradient in Japan from north to south, particularly along the Pacific coast with rainfall erosivity increasing southwards from the colder temperate/subarctic regions in the north towards the tropical regions in the south. Importantly, the rainfall erosivity of the Fukushima region is projected, by Shiono et al. (2013), to increase significantly as a result of climate change.

The difference between Shiono et al. (2013) and our current research is the comprehensive analysis of precipitation, and rainfall erosivity at a scale relevant to the fallout from the FDNPP accident. In particular, our current research highlights that the coastal mountain range, which received a significant amount of radiocesium fallout, has higher rainfall erosivity relative to the remainder of the fallout-impacted region, with a large proportion of this rainfall erosivity being derived for **tropical cyclones**.



### 4.3 Tropical cyclones

Regions subject to tropical cyclones generally have much high rainfall erosivity. In Mexico, the mean annual rainfall erosivity was  $6525 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ yr}^{-1}$  (García-Oliva et al., 1995). In Peninsular Malaysia, the R-factor was shown to range between 9000 and 14000  $\text{MJ mm ha}^{-1} \text{ h}^{-1} \text{ yr}^{-1}$  (Shamshad et al., 2008) in one study and between 1360 and 21600  $\text{MJ mm ha}^{-1} \text{ h}^{-1} \text{ yr}^{-1}$  in another (Yu et al., 2001). In these regions, the majority of the annual precipitation and rainfall erosivity may be derived from only a few storms (García-Oliva et al., 1995). In the Fukushima region, this current research demonstrated that 40% of the rainfall erosivity is derived from tropical cyclones. Accordingly, to model erosion and contaminant transfers in the Fukushima region, it may be important to focus on the spatial distribution of rainfall erosivity from major tropical cyclone events (Chartin et al., in revision).

### 4.4 Implications for radiocesium dynamics

Precipitation characteristics, such as rainfall erosivity, are fundamental inputs into a variety of approaches to modelling downstream contaminant transfers (Karydas et al., 2015; Yamaguchi et al., 2014; Rosa et al., 1996). Although precipitation and rainfall erosivity alone are insufficient to directly model downstream radiocesium migration, their characterization is important to improve our understanding of radiocesium dynamics. In the Fukushima region, rainfall drives three major contaminant transfer processes: i) wet fallout of particle from the atmosphere to the surface controls the spatial distribution of deposited contaminants, ii) rainfall splash effects trigger the detachment of particle-bound contaminants, and iii) detached particles are potentially transported by overland flow generated by rainfall when infiltration capacity is exceeded or when the soil is saturated with water. In addition, the timing and intensity of events in general, and tropical cyclones in particular, are likely of particular importance for understanding these and other post-fallout radiocesium dynamics in the Fukushima region (Fig 9).

The Fukushima region has been demonstrated to be highly reactive to major precipitation events (Evrard et al., 2013) with the climate driving a massive and episodic transfer of radiocesium downstream (Chartin et al., 2013). In fact, Yamaguchi et al. (2014) state that the majority of sediment and radiocesium is transferred downstream in 1–2 major events each year. In Figure 9, the potential impact of major events is quickly evident. In each year, the rainfall erosivity from

**tropical cyclones** is greater than the mean monthly rainfall erosivity (individual **tropical cyclone** and tropical storm data for events since 2011 are presented in Table S1 in the supplementary information).

Although the Chernobyl and Fukushima regions have similar humid continental climates (Dfb), the **tropical cyclones** and other significant summer rainfall events result in a higher precipitation (~2 times) and rainfall erosivity in the Fukushima region (~3–4 times). In the Kharkiv region of Ukraine, the mean annual precipitation is 460 mm yr<sup>-1</sup> (Nazarok et al., 2014), in Odessa it is <500 mm yr<sup>-1</sup> (Svetlitchnyi, 2009) compared to 1420 mm yr<sup>-1</sup> for the Fukushima region. In Ukraine, the region within a 100 km radius from Chernobyl has a rainfall erosivity that ranges from 830 to 1250 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> (Larionov, 1993). Within 100 km from the FDNPP accident, the R factor is ~3 to 4 times higher (3696 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup>). As there is positive relationship between rainfall erosivity and soil loss, where other factors are held equal, higher erosion rates and therefore a more rapid radiocesium export is anticipated in the Fukushima region compared to Chernobyl, based on these differences in their climatic contexts.

**R-factor maps focusing on the region impacted by fallout from the FDNPP accident that depict the annual and monthly variation in the R-factor are important for research and modelling of radiocesium dynamics in this region. These R-factor maps are thus provided (Dataset 1).** Further, all the event data (total precipitation, rainfall erosivity and kinetic energy) are included in the supplementary information (Dataset 2). More research is nevertheless required to examine the impact of different rain drop size distributions and resultant variations in kinetic energy in this region.

## **5 Conclusions**

**Rainfall erosivity is an important indicator of a region's climatic vulnerability to rainfall-driven soil erosion and also the potential downstream transfer of sediment and sediment-bound contaminants.** Rainfall erosivity in the Fukushima region is 3696 MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup> and the region receives 1420 mm yr<sup>-1</sup> of precipitation. Although these are below average values for Japan, the rainfall erosivity is more than 3–4 times higher than in Chernobyl. The majority of precipitation (60%) and rainfall erosivity (86%) occurs between June and October. Major **tropical cyclone** events are responsible for 22% of the precipitation (422 mm yr<sup>-1</sup>) and 40% of the rainfall

erosivity ( $1462 \text{ MJ mm ha}^{-1} \text{ h}^{-1} \text{ yr}^{-1}$  (SD 637)). The highest areas of contamination in the coastal mountain range correspond to the contaminated area that receives high rainfall erosivity.

The R-factor maps demonstrate that areas most contaminated with radiocesium with the highest rainfall erosivity are located along the coastal mountain range of the Pacific Ocean west of the FDNPP. Catchments in these areas will have the highest sensitivity to rainfall induced soil erosion and thus potential concomitant radiocesium transfers to downstream catchments and ultimately to the Pacific Ocean. The highest rainfall erosivity in these areas occurs during the tropical cyclone season (July-September). Monitoring radiocesium transfers after major events will be important. In other catchments that may be impacted by future radiocesium contamination it will be important to similarly investigate the climatic context, as precipitation strongly influences post-fallout radiocesium dynamics.

## Acknowledgements

We would like to thank the reviewers for their constructive comments that helped improve this manuscript. This research was funded by the ANR (*French National Research Agency*) in the framework of the AMORAD project (ANR-11-RSNR-0002).

## References

- Akaike, H.: A new look at the statistical model identification, *Automatic Control, IEEE Transactions on*, 19, 716-723, 1974.
- Barry, R. G., and Chorley, R. J.: *Atmosphere, weather and climate*, Routledge, 2009.
- Belyaev, V. R., Golosov, V. N., Ivanova, N. N., Markelov, M. V., and Tishkina, E. V.: Human-accelerated soil redistribution within an intensively cultivated dry valley catchment in southern European Russia, *Iahs-Aish P*, 291, 11-20, 2005.
- Capolongo, D., Diodato, N., Mannaerts, C. M., Piccarreta, M., and Strobl, R. O.: Analyzing temporal changes in climate erosivity using a simplified rainfall erosivity model in Basilicata (southern Italy), *Journal of Hydrology*, 356, 119-130, <http://dx.doi.org/10.1016/j.jhydrol.2008.04.002>, 2008.
- Chartin, C., Evrard, O., Onda, Y., Patin, J., Lefèvre, I., Ottlé, C., Ayrault, S., Lepage, H., and Bonté, P.: Tracking the early dispersion of contaminated sediment along rivers draining the Fukushima radioactive pollution plume, *Anthropocene*, 1, 23-34, 2013.
- Chartin, C., Evrard, O., Onda, Y., Lacey, P., Cerdan, O., Lepage, H., Ottlé, C., Lefèvre, I., and Bonté, P.: The impact of typhoons on sediment connectivity: Lessons learnt from contaminated coastal catchments of Fukushima Prefecture (Japan), *Earth Surface Processes and Landforms*, in revision.
- Chino, M., Nakayama, H., Nagai, H., Terada, H., Katata, G., and Yamazawa, H.: Preliminary estimation of release amounts of  $^{131}\text{I}$  and  $^{137}\text{Cs}$  accidentally discharged from the Fukushima Daiichi nuclear power plant into the atmosphere, *Journal of nuclear science and technology*, 48, 1129-1134, 2011.
- Daly, C.: Variable influence of terrain on precipitation patterns: Delineation and use of effective terrain height in PRISM, World Wide Web document. <http://www.ocs.orst.edu/prism/effter.pdf>, 2002.
- Daly, C., Gibson, W. P., Taylor, G. H., Johnson, G. L., and Pasteris, P.: A knowledge-based approach to the statistical mapping of climate, *Climate research*, 22, 99-113, 2002.
- de Brogniez, D., Ballabio, C., Stevens, A., Jones, R., Montanarella, L., and van Wesemael, B.: A map of the topsoil organic carbon content of Europe generated by a generalized additive model, *European Journal of Soil Science*, 66, 121-134, 2015.
- Duan, W., He, B., Takara, K., Luo, P., Hu, M., Alias, N. E., and Nover, D.: Changes of precipitation amounts and extremes over Japan between 1901 and 2012 and their connection to climate indices, *Climate Dynamics*, 45, 2273-2292, 2015.
- Eltaiif, N. I., Gharaibeh, M. A., Al-Zaitawi, F., and Alhamad, M. N.: Approximation of Rainfall Erosivity Factors in North Jordan, *Pedosphere*, 20, 711-717, [http://dx.doi.org/10.1016/S1002-0160\(10\)60061-6](http://dx.doi.org/10.1016/S1002-0160(10)60061-6), 2010.
- ESRI: ArcGIS Desktop: Release 10, in, Redlands. Environmental Systems Research Institute, CA, 2011.
- Evrard, O., Chartin, C., Onda, Y., Patin, J., Lepage, H., Lefèvre, I., Ayrault, S., Ottlé, C., and Bonté, P.: Evolution of radioactive dose rates in fresh sediment deposits along coastal rivers draining Fukushima contamination plume, *Scientific reports*, 3, 2013.
- Evrard, O., Chartin, C., Onda, Y., Lepage, H., Cerdan, O., Lefevre, I., and Ayrault, S.: Renewed soil erosion and remobilisation of radioactive sediment in Fukushima coastal rivers after the 2013 typhoons, *Sci. Rep.*, 4, 10.1038/srep04574, 2014.

551 Evrard, O., Laceby, J. P., Lepage, H., Onda, Y., Cerdan, O., and Ayrault, S.: Radiocesium transfer from  
 552 hillslopes to the Pacific Ocean after the Fukushima Nuclear Power Plant accident: A review, *Journal of*  
 553 *Environmental Radioactivity*, 148, 92-110, <http://dx.doi.org/10.1016/j.jenvrad.2015.06.018>, 2015.

554 García-Oliva, F., Maass, J. M., and Galicia, L.: Rainstorm Analysis and Rainfall Erosivity of a Seasonal  
 555 Tropical Region with a Strong Cyclonic Influence on the Pacific Coast of Mexico, *Journal of Applied*  
 556 *Meteorology*, 34, 2491-2498, 10.1175/1520-0450(1995)034<2491:RAAREO>2.0.CO;2, 1995.

557 Hastie, T., and Tibshirani, R.: Generalized additive models, *Statistical science*, 297-310, 1986.

558 Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., and Jarvis, A.: Very high resolution interpolated  
 559 climate surfaces for global land areas, *International journal of climatology*, 25, 1965-1978, 2005.

560 Jarvis, A., Reuter, H. I., Nelson, A., and Guevara, E.: Hole-filled SRTM for the globe Version 4, available  
 561 from the CGIAR-CSI SRTM 90m Database (<http://srtm.csi.cgiar.org>), 2008.

562 JMA (Japanese Meteorological Agency) RSMC Tokyo - Typhoon Center: [http://www.jma.go.jp/jma/jma-](http://www.jma.go.jp/jma/jma-eng/jma-center/rsmc-hp-pub-eg/RSMC_HP.htm)  
 563 [eng/jma-center/rsmc-hp-pub-eg/RSMC\\_HP.htm](http://www.jma.go.jp/jma/jma-eng/jma-center/rsmc-hp-pub-eg/RSMC_HP.htm), access: Feb 12, 2016.

564 Karki, K. B., and Shibano, H.: Soil loss in a forested watershed underlain by deeply weathered granite:  
 565 comparison of observations to predictions of a GIS-based USLE, *Bulletin of the Tokyo University Forests*,  
 566 2006.

567 Karydas, C. G., Tzoraki, O., and Panagos, P.: A New Spatiotemporal Risk Index for Heavy Metals:  
 568 Application in Cyprus, *Water*, 7, 4323-4342, 2015.

569 Kato, H., Onda, Y., and Teramage, M.: Depth distribution of <sup>137</sup>Cs, <sup>134</sup>Cs, and <sup>131</sup>I in soil profile after  
 570 Fukushima Dai-ichi Nuclear Power Plant Accident, *Journal of Environmental Radioactivity*, 111, 59-64,  
 571 <http://dx.doi.org/10.1016/j.jenvrad.2011.10.003>, 2012.

572 Kim, S., Nakakita, E., Tachikawa, Y., and Takara, K.: Precipitation changes in Japan under the A1B climate  
 573 change scenario, *Annual Journal of Hydraulic Engineering, JSCE*, 54, 127-132, 2010.

574 Kinnell, P. I. A.: Event soil loss, runoff and the Universal Soil Loss Equation family of models: A review,  
 575 *Journal of Hydrology*, 385, 384-397, <http://dx.doi.org/10.1016/j.jhydrol.2010.01.024>, 2010.

576 Kitamura, A., Yamaguchi, M., Kurikami, H., Yui, M., and Onishi, Y.: Predicting sediment and cesium-137  
 577 discharge from catchments in eastern Fukushima, *Anthropocene*, 5, 22-31,  
 578 <http://dx.doi.org/10.1016/j.ancene.2014.07.001>, 2014.

579 Larionov, G.: Erosion and deflation of soils, in, Moscow University Press. Moscow, Russia (in Russian),  
 580 pp200, 1993.

581 Lee, J.-H., and Heo, J.-H.: Evaluation of estimation methods for rainfall erosivity based on annual  
 582 precipitation in Korea, *Journal of Hydrology*, 409, 30-48,  
 583 <http://dx.doi.org/10.1016/j.jhydrol.2011.07.031>, 2011.

584 Lu, H., and Yu, B.: Spatial and seasonal distribution of rainfall erosivity in Australia, *Soil Research*, 40, 887-  
 585 901, 2002.

586 Ma, X., He, Y., Xu, J., van Noordwijk, M., and Lu, X.: Spatial and temporal variation in rainfall erosivity in a  
 587 Himalayan watershed, *CATENA*, 121, 248-259, <http://dx.doi.org/10.1016/j.catena.2014.05.017>, 2014.

588 McFarlane, D., Westcott, T., and Davies, J.: Rainfall erosivity in Western Australia, *Hydrology and Water*  
 589 *Resources Symposium 1986: River Basin Management; Preprints of Papers*, 1986, 350,

590 Mello, C. R., Viola, M. R., Beskow, S., and Norton, L. D.: Multivariate models for annual rainfall erosivity in  
591 Brazil, *Geoderma*, 202–203, 88–102, <http://dx.doi.org/10.1016/j.geoderma.2013.03.009>, 2013.

592 Meusburger, K., Steel, A., Panagos, P., Montanarella, L., and Alewell, C.: Spatial and temporal variability  
593 of rainfall erosivity factor for Switzerland, *Hydrol. Earth Syst. Sci.*, 16, 167–177, 10.5194/hess-16-167-  
594 2012, 2012.

595 Moore, T.: Rainfall erosivity in East Africa, *Geografiska Annaler. Series A. Physical Geography*, 147–156,  
596 1979.

597 Mouri, G., Golosov, V., Shiiba, M., and Hori, T.: Assessment of the caesium-137 flux adsorbed to  
598 suspended sediment in a reservoir in the contaminated Fukushima region in Japan, *Environmental*  
599 *Pollution*, 187, 31–41, <http://dx.doi.org/10.1016/j.envpol.2013.12.018>, 2014.

600 Nagao, S., Kanamori, M., Ochiai, S., Tomihara, S., Fukushi, K., and Yamamoto, M.: Export of <sup>134</sup>Cs and  
601 <sup>137</sup>Cs in the Fukushima river systems at heavy rains by Typhoon Roke in September 2011, *Biogeosciences*  
602 *Discuss.*, 10, 2767–2790, 10.5194/bgd-10-2767-2013, 2013.

603 Nazarok, P., Kruglov, O., Menshov, O., Kutsenko, M., and Sukhorada, A.: Mapping soil erosion using  
604 magnetic susceptibility. A case study in Ukraine, *Solid Earth Discussions*, 6, 831–848, 2014.

605 Oliveira, P. T. S., Wendland, E., and Nearing, M. A.: Rainfall erosivity in Brazil: A review, *CATENA*, 100,  
606 139–147, <http://dx.doi.org/10.1016/j.catena.2012.08.006>, 2013.

607 Onishi, Y., Kurikami, H., and Yokuda, S.: Simulation of Sediment and Cesium Transport in the Ukedo River  
608 and the Ogi Dam Reservoir during a Rainfall Event using the TODAM Code, *Pacific Northwest National*  
609 *Laboratory (PNNL), Richland, WA (US)*, 2014.

610 Panagos, P., Ballabio, C., Borrelli, P., Meusburger, K., Klik, A., Rousseva, S., Tadić, M. P., Michaelides, S.,  
611 Hrabalíková, M., and Olsen, P.: Rainfall erosivity in Europe, *Science of the Total Environment*, 511, 801–  
612 814, 2015.

613 Peel, M. C., Finlayson, B. L., and McMahon, T. A.: Updated world map of the Köppen-Geiger climate  
614 classification, *Hydrology and earth system sciences discussions*, 4, 439–473, 2007.

615 Poggio, L., Gimona, A., and Brewer, M. J.: Regional scale mapping of soil properties and their uncertainty  
616 with a large number of satellite-derived covariates, *Geoderma*, 209, 1–14, 2013.

617 Renard, K. G., Foster, G. R., Weesies, G. A., and Porter, J. P.: RUSLE: Revised universal soil loss equation,  
618 *Journal of soil and Water Conservation*, 46, 30–33, 1991.

619 Renard, K. G., and Freimund, J. R.: Using monthly precipitation data to estimate the R-factor in the  
620 revised USLE, *Journal of Hydrology*, 157, 287–306, [http://dx.doi.org/10.1016/0022-1694\(94\)90110-4](http://dx.doi.org/10.1016/0022-1694(94)90110-4),  
621 1994.

622 Renard, K. G., Foster, G. R., Weesies, G. A., McCool, D., and Yoder, D.: Predicting soil erosion by water: a  
623 guide to conservation planning with the revised universal soil loss equation (RUSLE), *Agriculture*  
624 *Handbook* (Washington), 1997.

625 Rosa, D. d. I., Cromptvoets, J., Mayol Rodríguez, F., and Moreno Arce, J. A.: Land vulnerability evaluation  
626 and climate change impacts in Andalusia, Spain: soil erosion and contamination, 1996.

627 Saito, K., and Onda, Y.: Outline of the national mapping projects implemented after the Fukushima  
628 accident, *Journal of environmental radioactivity*, 139, 240–249, 2015.

629 Saito, K., Tanihata, I., Fujiwara, M., Saito, T., Shimoura, S., Otsuka, T., Onda, Y., Hoshi, M., Ikeuchi, Y., and  
630 Takahashi, F.: Detailed deposition density maps constructed by large-scale soil sampling for gamma-ray  
631 emitting radioactive nuclides from the Fukushima Dai-ichi Nuclear Power Plant accident, *Journal of*  
632 *environmental radioactivity*, 139, 308-319, 2015.

633 Sawhney, B.: Selective sorption and fixation of cations by clay minerals: a review, *Clays Clay Miner*, 20,  
634 93-100, 1972.

635 Shamshad, A., Azhari, M. N., Isa, M. H., Hussin, W. M. A. W., and Parida, B. P.: Development of an  
636 appropriate procedure for estimation of RUSLE EI30 index and preparation of erosivity maps for Pulau  
637 Penang in Peninsular Malaysia, *CATENA*, 72, 423-432, <http://dx.doi.org/10.1016/j.catena.2007.08.002>,  
638 2008.

639 Shiono, T., Ogawa, S., Miyamoto, T., and Kameyama, K.: Expected impacts of climate change on rainfall  
640 erosivity of farmlands in Japan, *Ecological Engineering*, 61, 678-689, 2013.

641 Svetlitchnyi, A.: Soil Erosion Induced Degradation of Agrolandscapes in Ukraine: Modeling, Computation  
642 and Prediction in Conditions of the Climate Changes, in: *Regional Aspects of Climate-Terrestrial-*  
643 *Hydrologic Interactions in Non-boreal Eastern Europe*, edited by: Groisman, P., and Ivanov, S., NATO  
644 Science for Peace and Security Series C: Environmental Security, Springer Netherlands, 191-199, 2009.

645 Takeuchi, B. K., and Ishidaira, H.: The examination of sediment production estimation by the USLE  
646 method in mountainous basins of Japan, *Hydraulic Engineering*, 45, 85-90, 2001.

647 Tamura, T.: Consequences of activity release: selective sorption reactions of cesium with soil minerals,  
648 *Nuclear Safety* 5, 262-268, 1964.

649 Thakur, P., Ballard, S., and Nelson, R.: An overview of Fukushima radionuclides measured in the northern  
650 hemisphere, *Science of The Total Environment*, 458-460, 577-613,  
651 <http://dx.doi.org/10.1016/j.scitotenv.2013.03.105>, 2013.

652 Tran, T. A., Mitani, Y., Ikemi, H., and Matsuki, H.: Human Impacts on Erosion and Deposition in Onga  
653 River Basin, Kyushu, Japan, *Memoirs of the Faculty of Engineering, Kyushu University*, 71, 47-65, 2011.

654 USDA: Rainfall Intensity Summarization Tool (RIST), United States Department of Agriculture (USDA),  
655 Agriculture Research Service, National Sedimentation Laboratory, Oxford, Mississippi, Version 3.88, in,  
656 2013.

657 Wang, X. L., Chen, H., Wu, Y., Feng, Y., and Pu, Q.: New Techniques for the Detection and Adjustment of  
658 Shifts in Daily Precipitation Data Series, *Journal of Applied Meteorology and Climatology*, 49, 2416-2436,  
659 10.1175/2010JAMC2376.1, 2010.

660 Wikipedia Pacific typhoon climatology: [https://en.wikipedia.org/wiki/Pacific\\_typhoon\\_climatology](https://en.wikipedia.org/wiki/Pacific_typhoon_climatology),  
661 access: Feb 13, 2016.

662 Wilkinson, S. N., Prosser, I. P., Rustomji, P., and Read, A. M.: Modelling and testing spatially distributed  
663 sediment budgets to relate erosion processes to sediment yields, *Environment Modelling and Software*,  
664 24, 489-501, 2009.

665 Wischmeier, W. H., and Smith, D. D.: Rainfall energy and its relationship to soil loss, *Transactions*  
666 *American Geophysical Union*, 39, 285-291, 10.1029/TR039i002p00285, 1958.

667 Wischmeier, W. H., and Smith, D. D.: Predicting rainfall erosion losses-A guide to conservation planning,  
668 *Predicting rainfall erosion losses-A guide to conservation planning*, 1978.

669 Wood, S. N.: mgcv: GAMs and generalized ridge regression for r. *R news*, 1(2), 2-25, 2001.



670 Yamaguchi, M., Kitamura, A., Oda, Y., and Onishi, Y.: Predicting the long-term <sup>137</sup>Cs distribution in  
671 Fukushima after the Fukushima Dai-ichi nuclear power plant accident: a parameter sensitivity analysis,  
672 Journal of Environmental Radioactivity, 135, 135-146, <http://dx.doi.org/10.1016/j.jenvrad.2014.04.011>,  
673 2014.

674 Yamashiki, Y., Onda, Y., Smith, H. G., Blake, W. H., Wakahara, T., Igarashi, Y., Matsuura, Y., and  
675 Yoshimura, K.: Initial flux of sediment-associated radiocesium to the ocean from the largest river  
676 impacted by Fukushima Daiichi Nuclear Power Plant, Scientific Reports, 4, 10.1038/srep03714, 2014.

677 Yoshimura, K., Onda, Y., and Kato, H.: Evaluation of radiocaesium wash-off by soil erosion from various  
678 land uses using USLE plots, Journal of Environmental Radioactivity, 139, 362–369,  
679 <http://dx.doi.org/10.1016/j.jenvrad.2014.07.019>, 2015.

680 Yu, B., Hashim, G., and Eusof, Z.: Estimating the R-factor with limited rainfall data: a case study from  
681 peninsular Malaysia, Journal of Soil and Water Conservation, 56, 101-105, 2001.

682 Zar, J. H.: Biostatistical Analysis, in, 5th Edition ed., Prentice Hall, New Jersey, 944, 2010.

683

684

685



**Table 1.** Station coordinates along with the annual precipitation, R-Factor and kinetic energy (KE) mean and standard deviation (SD).

Station	Latitude	Longitude	Annual Precipitation (mm)		R-Factor (MJ mm ha <sup>-1</sup> h <sup>-1</sup> yr <sup>-1</sup> )		KE (MJ ha <sup>-1</sup> yr <sup>-1</sup> )	
			Mean	SD	Mean	SD	Mean	SD
254 <sup>A-4</sup>	38.178	140.635	1429	302	3295	3045	167	61
256	38.015	140.612	1292	243	2497	1296	149	42
257	38.025	140.858	1256	259	2922	1313	148	38
279	37.912	140.143	1343	221	2049	572	142	28
281	37.852	140.588	1098	172	2045	792	121	26
285	37.783	140.925	1350	266	3859	1928	171	48
288 <sup>A</sup>	37.638	140.983	1333	272	3819	1527	173	39
290	37.552	140.108	1265	242	1972	1029	124	35
291	37.583	140.430	1176	219	2337	1206	136	33
294	37.435	140.577	1168	210	2718	1355	141	36
295	37.492	140.965	1493	299	4327	1694	196	43
299	37.368	140.330	1154	176	2465	982	130	30
300 <sup>A-D</sup>	37.347	141.017	1532	295	4249	1517	203	40
304	37.287	140.625	1231	200	2984	1361	145	31
307	37.147	140.460	1280	205	3405	1530	157	34
310 <sup>A</sup>	37.065	140.877	1426	281	4425	2105	193	49
312	36.938	140.408	1399	234	4342	2145	182	46
314 <sup>A</sup>	36.868	140.637	2001	380	7177	2405	274	55
315	36.833	140.772	1472	247	4126	1629	185	42
316 <sup>10</sup>	36.778	140.345	1419	204	5075	2370	193	39
326	37.123	140.035	1954	373	8274	8491	279	78
1011	36.580	140.645	1474	232	4406	2002	191	37
1034	37.233	141.000	1568	280	4530	1545	205	45
1116 <sup>3</sup>	37.668	140.260	2201	355	5056	2385	262	57
1127 <sup>C-6</sup>	37.008	140.737	1435	285	3636	1263	178	46
1129	37.337	140.808	1444	250	4193	2115	185	42
1130	37.695	140.747	1307	245	2989	1563	157	41
1132 <sup>10</sup>	38.003	140.207	1254	199	2350	1380	127	34
1150 <sup>A10</sup>	37.560	140.753	1341	239	3470	1369	165	35
1173 <sup>A10</sup>	36.778	140.482	1437	234	4432	1981	191	42
1220 <sup>10</sup>	37.932	140.778	1262	245	2878	1234	149	40
1268 <sup>A8</sup>	37.288	140.218	1366	256	3680	2595	170	50
1274 <sup>A8</sup>	37.207	140.750	1440	272	4156	1812	181	43
1282 <sup>9</sup>	37.722	140.058	1762	316	3351	1764	200	49
1293 <sup>9</sup>	37.892	140.437	1363	178	2361	916	144	29
1294 <sup>9</sup>	37.277	140.063	1568	281	3914	2567	190	51
1298 <sup>A8</sup>	37.827	140.728	1462	270	3765	2209	175	46
1386 <sup>A7</sup>	37.388	140.090	1311	231	2726	1569	142	39
1421 <sup>A5</sup>	36.743	140.593	1851	284	6448	2422	249	50
1464	38.138	140.917	1129	174	2742	1522	133	27
1466 <sup>B-2</sup>	37.227	140.427	1232	213	3150	1055	146	33
1564	38.127	140.680	1356	170	2647	1030	154	26

<sup>A</sup>No temperature data available

<sup>A</sup> For R-factor, 10 years of data available, <sup>B</sup> 13y, <sup>C</sup> 14y, <sup>D</sup> 18y

<sup>1</sup>For precipitation analyses, 10y available, <sup>2</sup> 13y, <sup>3</sup> 21y, <sup>4</sup> 28y, <sup>5</sup> 30y, <sup>6</sup> 32y, <sup>7</sup> 33y, <sup>8</sup> 36 y, <sup>9</sup> 37 y, <sup>10</sup> 38y


**Table 2.** Observed and predicted mean and standard deviation (**SD**) for monthly (in MJ mm ha<sup>-1</sup> h<sup>-1</sup> m<sup>-1</sup>) and annual R-factors (in MJ mm ha<sup>-1</sup> h<sup>-1</sup> yr<sup>-1</sup>) with GAM performance parameters (RMSE: Root Mean Square Error; R<sup>2</sup>: coefficient of determination).

Month or Year	Observed R		Predicted R		Variance explained (%)	Model performance (LOOCV)	
	Mean	SD	Mean	SD		RMSE	R <sup>2</sup>
Year	3696	1327	3696	1293	96	640	0.77
January	43	26	44	25	90	16	0.77
February	24	16	23	15	90	9	0.68
March	64	31	64	30	93	13	0.83
April	160	104	165	97	92	66	0.62
May	206	160	214	163	98	75	0.80
June	312	124	312	116	88	66	0.72
July	800	283	800	244	83	167	0.65
August	779	469	755	305	92	314	0.56
September	785	287	781	263	83	165	0.67
October	368	247	374	237	95	92	0.86
November	73	28	72	25	90	12	0.80
December	83	66	84	62	93	33	0.76


**Table 3.** Results from the backward stepwise procedure used to select covariates to include in final GAMs for annual and monthly R-factor maps computation. The values ( $\Delta AIC$ ) correspond to the difference in AIC scores obtained when the particular covariate is dropped from the ‘full’ model, i.e., the model containing all the covariates. Scores in bold indicate which covariates were selected in the final models.

Model	$\Delta AIC$							
	Elevation	Slope	DC*	Easting	Northing	Rainfall	Temperature	Facets
Annual	<b>8.7</b>	1.0	<b>10.8</b>	<b>12.1</b>	2.8	<b>13.6</b>	<b>10.1</b>	<b>5.6</b>
January	0.0	<b>6.1</b>	<b>2.1</b>	<b>0.7</b>	0.0	<b>8.3</b>	0.0	<b>4.0</b>
February	<b>2.2</b>	0.0	0.1	<b>2.7</b>	<b>2.6</b>	0.2	<b>4.6</b>	<b>8.6</b>
March	0.0	<b>9.9</b>	0.0	<b>1.3</b>	0.1	<b>5.3</b>	0.0	<b>3.2</b>
April	<b>3.7</b>	0.0	<b>17.6</b>	0.0	<b>0.5</b>	<b>17.3</b>	0.0	<b>14.0</b>
May	0.1	<b>8.5</b>	0.0	<b>0.7</b>	<b>4.5</b>	<b>8.4</b>	0.0	<b>10.9</b>
June	0.4	0.3	0.0	0.2	<b>1.4</b>	<b>8.6</b>	0.0	<b>12.6</b>
July	0.0	0.3	<b>5.5</b>	0.0	0.0	<b>13.2</b>	0.0	<b>13.8</b>
August	<b>0.7</b>	<b>1.6</b>	0.0	0.0	<b>8.6</b>	<b>8.8</b>	0.1	<b>7.8</b>
September	0.0	0.4	<b>5.5</b>	0.0	0.0	<b>13.2</b>	0.0	<b>13.8</b>
October	0.4	<b>4.8</b>	<b>1.2</b>	0.0	<b>5.7</b>	<b>16.3</b>	0.0	<b>19.8</b>
November	0.0	<b>26.3</b>	<b>2.7</b>	<b>3.0</b>	0.2	0.2	<b>11.2</b>	<b>8.8</b>
December	0.3	0.2	0.0	<b>3.2</b>	0.4	0.6	<b>11.7</b>	<b>5.5</b>


\* Distance to Coast




**Figure 1.** Standard, omitted and long-term stations in the Fukushima region, along with catchments (grey lines) contained entirely within a 100km buffer around the FDNPP (dashed line)  $^{134}\text{+}^{137}\text{Cs}$  activities are from Chartin et al. (2013)




**Figure 2.** Total (grey) and typhoon-related (red) mean annual precipitation (A-Top) and R-factor (B-Bottom) with the solid line representing the mean and the dashed lines being one standard deviation on the mean. The error bars indicate one standard deviation on the mean for individual stations.



**Figure 3.** Annual (grey) and typhoon-related (red) mean annual precipitation (A) and corresponding standardized anomaly index (SAI – B) and the **annual** (grey) and typhoon related (red) R-factor (C) and the R-factor's SAI (D). The grey solid line represent the regional mean (including typhoons) and the grey dashed lines represent one standard deviation on the total mean whereas the red dashed line representing the mean contribution from typhoon events. The blue (total) and red (typhoon) solid lines depict the 5-year running average.




**Figure 4.** Mean annual precipitation (A) and the standardized **anomaly** index (SAI - B) for three long-term rainfall stations including mean annual precipitation (solid line), the standard deviation (dashed line) for the entire period and with the 5-year running average (blue line).



**Figure 5.** Total (grey) and typhoon related (red) mean monthly rainfall (A-top) and R-factor (B-bottom) with error bars indicating one standard deviation on the mean and the solid line representing the monthly mean and dashed lines representing on standard deviation on the mean. The mean monthly temperature is plotted (red line) along with mean monthly rainfall (A-top).






**Figure 6.** Mean maximum daily precipitation (A) and corresponding SAI with error bars indicating one standard deviation on the mean for each station, along with the mean (solid) and one standard deviation on the mean (dashed line) for the daily rainfall analyses period and the blue solid line depicting the 5-year running average.


697

698


699



**Figure 7:** Annual R-factor (A) and model errors depicted as the coefficient of variation (i.e.  $CV = (\text{standard error} / \text{mean}) * 100$ ) (B) for catchments entirely contained within a 100 km radius of the FDNPP. <sup>134+137</sup>Cs activities are from Chartin et al. (2013).



**Figure 8.** Monthly R-factor distribution for catchments contained entirely within a 100 km radius of the FDNPP. <sup>134+137</sup>Cs activities are from Chartin et al. (2013).



**Figure 9.** Total (grey) and typhoon related (red) monthly rainfall erosivity from 2011–2015 with the mean line representing mean monthly rainfall erosivity and the dashed line being one standard deviation on the mean for the period of 1995-2015 for the 42 rainfall stations analyzed.