

Review of “Does drought alter hydrological functions in forest soils? An infiltration experiment” hess-2015-255 by K. Gimbel, H. Puhmann, and M. Weiler

## 1 Response to Reviewers and Summary of Changes

First, we would like to thank the editor and the referees for the review and the helpful comments to improve the paper. We have addressed all the comments as explained below.

### Editor comments:

-Abstract: “Climate change is expected to change the water cycle”. I would suggest replacing the second “change” with “impact” or “modify”.

**Answer:** We changed the sentence, the sentence now reads as follows:

“Climate change is expected to impact the water cycle and severely affect precipitation patterns across central Europe and in other parts of the world, leading to more frequent and severe droughts.”

-Pag. 26, line 6: “often in addition to elevated” – remove “often”

**Answer:** We removed “often”

-Pag. 26, line 29: “First” should be “first”

**Answer:** We corrected it.

-Pag. 26, line 12-13: “...only few studies focus on forest ecosystems or take a closer look at drought impacts on soils where often only soil moisture is observed (Ozolinčius 12 et al., 2009,; Albert et al., 2011,; Glaser et al., 2013)”. According to the response to reviewers, this should be “...only few studies focus on forest ecosystems or take a closer look at drought impacts on soils where often only soil moisture is observed to change and no variations in other soil properties are monitored (Ozolinčius et 34 al., 2009; Albert et al., 2011; Glaser et al., 2013).”

**Answer:** We changed the sentence as asked above.

-Pag. 33, lines 4-6: “the reverse moisture...in the data”. This is a bit confusing, please provide other plausible explanations, rather than hydraulic redistribution (since you say you did not find any sign of hydraulic redistribution).

**Answer:** We agree and changed the sentence to:

“The reverse moisture effect might be caused by effects associated with the installation and placement of the probes (e.g. preferential flow along probe cables, concentration of throughfall – although installations were performed with great care), or root effects, for example hydraulic redistribution. However,...”

-Fig. 4: for completeness, add in the caption the time period over which the cumulative soil moisture

**Answer:** We changed the caption and added following sentence:

“The computations begin with the start of the rainfall reduction and end at the date of the dye tracer experiment 2013.”

# 1 Does drought alter hydrological functions in forest soils?

## 2 An infiltration experiment

3  
4 K. F. Gimbel<sup>1</sup>, H. Puhlmann<sup>2</sup>, M. Weiler<sup>1</sup>

5  
6 [1]{ Hydrology, Faculty of Environment and Natural Resources, University of Freiburg,  
7 Germany }

8 [2]{ Forest Research Institute Baden-Württemberg, Freiburg, Germany }

9 Correspondence to: K. F. Gimbel (katharina.gimbel@hydrology.uni-freiburg.de)

### 10 11 **Abstract**

12 Climate change is expected to ~~change~~impact the water cycle and severely affect precipitation  
13 patterns across central Europe and in other parts of the world, leading to more frequent and  
14 severe droughts. Usually when projecting drought impacts on hydrological systems, it is  
15 assumed that system properties, like soil properties, remain stable and will not be affected by  
16 drought events. To study if this assumption is appropriate, we address the effects of drought  
17 on the infiltration behavior of forest soils using dye tracer experiments on six sites in three  
18 regions across Germany, which were forced into drought conditions. The sites cover clayey,  
19 loamy and sandy textured soils. In each region, we compared a deciduous and a coniferous  
20 forest stand to address differences between the main tree species. The results of the dye tracer  
21 experiments show clear evidence for changes in infiltration behavior at the sites. The  
22 infiltration changed at the clayey plots from regular and homogeneous flow to fast  
23 preferential flow. Similar behavior was observed at the loamy plots, where large areas in the  
24 upper layers remained dry, displaying signs of strong water repellency. This was confirmed  
25 by water drop penetration times (WDPT) tests, which revealed, in all except one plot,  
26 moderate to severe water repellency. Water repellency was also accountable for the change of  
27 regular infiltration to fingered flow in the sandy soils. The results of this study suggest that  
28 the “drought-history” or generally the climatic conditions in the past of a soil are more  
29 important than the actual antecedent soil moisture status regarding hydrophobicity and

1 infiltration behavior; and also, that drought effects on infiltration need to be considered in  
2 hydrological models to obtain realistic predictions concerning water quality and quantity in  
3 runoff and groundwater recharge.

4

## 5 **1 Introduction**

6 Soils moderate how water moves through the vadose zone and govern the percolation of water  
7 to groundwater and stream flow. Soils not only store water for plant growth, function as a  
8 habitat for different biota and as transition zone to groundwater, but are also important –  
9 especially the top layers – for sorption and degradation of contaminants and (agri-)chemicals  
10 (Hendrickx and Flury 2001). The efficiency of this important ecosystem service for  
11 groundwater and surface water protection depends on the behavior of pollutants in the soil  
12 and the hydrological transport processes (Keesstra et al. 2012). How fast water passes the  
13 vadose zone depends on its hydraulic soil properties and distribution such as pore volume  
14 distribution, soil aggregation, water repellency and rooting pattern.

15 Due to climate change and increasing human intervention, the global water cycle is expected  
16 to change with probably increasing summer dryness and winter wetness in many regions  
17 across the world including Western and Central Europe (IPCC 2012, Prudhomme et al. 2014).  
18 In addition, droughts are expected to be more frequent and severe in the future (Prudhomme  
19 et al. 2014, Seneviratne et al. 2006). Drought conditions can alter the hydrological functions  
20 of soils, and soil structure is responding to drought by shrinkage and fracturing of soil  
21 aggregates. These soil shrinkage cracks channel the infiltrating water and by that foster the  
22 bypassing of the soil matrix (Hendrickx and Flury 2001, Ritsema et al. 1997) and therefore  
23 alter the infiltration patterns in soil. Thus, the infiltration and redistribution of water within  
24 the soil changes and hence also the proportion of water reaching the groundwater (Hendrickx  
25 and Flury 2001).

26 Soils under drought conditions are prone to become water repellent, depending on soil  
27 properties and organic matter content (DeBano 1981 and 2000). Due to modifications of the  
28 three-dimensional distribution and dynamics of soil moisture, water repellency has far  
29 reaching consequences for infiltration processes (Doerr and Ritsema 2006). Water repellency  
30 hinders infiltration and thus either increases overland flow (Doerr and Ritsema 2006) or  
31 redirects the water into preferential flow paths and creates instable wetting fronts (fingered  
32 preferential flow; Ritsema et al. 1993 and 2000, Dekker and Ritsema 2000).

1 To assess the impacts of drought, rainfall exclusion experiments are valuable and often  
2 applied tools (e.g. English et al. 2005, Phillips et al. 2009, Da Costa et al. 2010, Kopittke et al.  
3 2014), ~~often~~ in addition to elevated CO<sub>2</sub> concentrations (e.g. Dermody et al. 2007), and night-  
4 time warming (e.g. Albert et al. 2011, Selsted et al. 2012). While many studies focus on single  
5 aspects of drought effects like plant growth and seedling activity (Meijer et al., 2011; Wu and  
6 Chen, 2013) or on particular ecosystems like grassland (Suttle and Thomson, 2007; Bütof 5 et  
7 al., 2012) and heather ecosystems (Albert et al., 2011; Selsted et al., 2012), only few studies  
8 focus on forest ecosystems or take a closer look at drought impacts on soils where often only  
9 soil moisture is observed to change and no variations in other soil properties are monitored  
10 (Ozolinčius et al., 2009; Albert et al., 2011; Glaser et al., 2013). ~~only few studies focus on~~  
11 ~~forest ecosystems or take a closer look at drought impacts on soils where often only soil~~  
12 ~~moisture is observed (Ozolinčius et al., 2009; Albert et al., 2011; Glaser et al., 2013).~~

13 To study drought effects, often extreme short-time events equivalent to droughts with  
14 occurrence probabilities of up to 100 or even 1000 years have been introduced to the  
15 examined soils (e.g. Glaser et al. 2013). By introducing these extreme events, the question of  
16 transferability of the results to natural systems in respect to the expected behavior under  
17 predicted future drought conditions arises. Therefore, this study employs a moderate rainfall  
18 reduction equivalent to an annual drought with a 40-year return period, in accordance to  
19 climate predictions, thereby avoiding tentativeness due to an overreaction to an unnatural  
20 extreme drought (Gimbel et al. 2015).

21 To monitor changes in soil hydraulic properties, the changes in infiltration patterns in the soil  
22 after two years of prolonged drought were observed in three regions across Germany.  
23 Infiltration patterns were chosen because they reflect the integrated changes of soil  
24 hydrological functions and directly show how water moves in the soil under altered  
25 conditions. In this paper, we present results of several dye tracer infiltration experiments  
26 before and after two years of prolonged artificial drought. The objectives of this study are:  
27 fFirst, to investigate whether droughts predicted by climate projections affect the infiltration  
28 behavior of forest soils, and second, whether changes in infiltration patterns can be attributed  
29 to changes in the hydrologic properties of the soils. Three hypotheses will be tested: (1)  
30 Induced drought alters infiltration patterns due to changes in soil hydraulic properties; e.g.  
31 soil water repellency and forming of shrinkage cracks, leading to preferential flow paths and  
32 faster infiltration. (2) The main tree species have an effect on the magnitude of the observed

1 response. (3) The drought will increase water repellency depending on tree species and soil  
2 properties.

3

## 4 **2 Material and Methods**

### 5 **2.1 Study sites**

6 To identify the influence of drought on infiltration patterns of forest soils, six plots in three  
7 different regions across Germany were selected. The plots were located in Schwäbische Alb  
8 (South-West Germany), Hainich-Dün (Central Germany) and Schorfheide-Chorin (North-East  
9 Germany) (Figure 1). All plots are part of the Biodiversity Exploratories framework that  
10 incorporates, in total, 150 sites on grassland and 150 sites in forest (for more information on  
11 the Biodiversity Exploratories, refer to Fischer et al. 2010). In each of the Exploratories, two  
12 forest plots were selected, which are – within each Exploratory – similar with respect to  
13 topography and soil texture type (Figure 2) but differ in tree species composition. In each site,  
14 one plot with a coniferous and one with a deciduous main tree species was selected. At the  
15 Schwäbische Alb and Hainich-Dün sites, beech (*Fagus sylvatica*) and spruce (*Picea abies*)  
16 were chosen, in Schorfheide-Chorin beech and pine (*Pinus sylvestris*).

17 The Schwäbische Alb soils are shallow (25 to 35 cm) Leptosols on Jurassic shell limestone  
18 with a high stone content (Figure 2, top). The mean annual temperature at this site is 6.5° C  
19 and the mean annual precipitation amounts to 940 mm. The underlying geology of the  
20 Hainich-Dün is Triassic limestone. The soils at this site are loamy Stagnosols with depths  
21 between 45 and 65 cm. At the Hainich-Dün site the mean annual temperature is 7.2° C and  
22 the mean annual precipitation is 533 mm. The Schorfheide-Chorin plots are located in a  
23 young glacial landscape where the dominant geological substrate is glacial till covered by  
24 glacio-fluvial and aeolian sands. The soils at this site are deep, sandy Cambisols. At the  
25 Schorfheide-Chorin site, mean annual temperature is 8.5° C and the mean annual precipitation  
26 amounts to 589 mm. All climate data are taken from nearby stations of the German weather  
27 service (DWD, years 1950–2010).

28 The experiments of this study are part of the interdisciplinary project ‘Global Change Effects  
29 on Forest Understorey: Interactions between Drought and Land-use Intensity’ (Gimbel et al.  
30 2015). The artificial imposed drought was created by a 10 m x 10 m roofed subplot, covered  
31 with transparent panels. In addition, a control plot with the same technical equipment, but

1 without the roofing was installed. The distance between the roofed and the control plots range  
2 between 15 m and 30 m. The control and roofed plots include a central adult overstorey tree,  
3 which are similar in age, size, and canopy structure. To provide sufficient exchange with  
4 ambient air (avoiding a ‘greenhouse effect’), all four sides of the roof are open. To collect  
5 water from the roof, rain gutters are mounted alongside the timber construction. The roof is  
6 designed to reduce precipitation between 11 and 100 % - 11 % already intercepted by the  
7 roofing construction and rain gutters itself. The incoming precipitation was reduced between  
8 March and November to the level equivalent to an annual drought with a return period of 40  
9 years. The resulting annual target precipitation inputs under the roofs were 700 mm (26 %  
10 reduction) for Schwäbische Alb, 355 mm (33 % reduction) at the Hainich-Dün, and 395 mm  
11 (27 % reduction) at the Schorfheide-Chorin site. For a more detailed description of the whole  
12 experimental drought setting and of the study plots see Gimbel et al. (2015).

## 13 **2.2 Soil moisture measurements**

14 To observe the impact of reduced precipitation input on soil moisture, soil moisture probes  
15 were installed on the drought and on the control subplots of every site. The probes (5TM and  
16 5TE, Decagon Devices Inc.) were inserted in 5 cm, 15 cm and 30 cm depths in three  
17 replicates on the plots at 2, 3, and 4 m distance from the central trees. The accuracy according  
18 to the technical data sheets of the 5TE and 5TM probes is  $\pm 15\%$  of the measured value for  
19 the volumetric water content. The readings of every probe are logged at 15 min intervals. For  
20 better comparability among the sites, the mean values of the three replicates of every depth  
21 per control and drought plot were cumulated and normalized to the maximum cumulated  
22 value of the control plot.

## 23 **2.3 Soil water repellency**

24 Hydrophobicity in soil was measured with the water drop penetration time (WDPT) test (e.g.,  
25 Bisdom et al. 1993). This test determines how long water repellency persists on a porous  
26 surface. The tests were performed immediately before the dye tracer experiments in 2013, in  
27 the drought and control profiles of the deciduous plots and in the drought profiles of the  
28 coniferous plots. For the WDPT tests, a water droplet is placed on a planar soil surface with a  
29 pipette and the time is taken until the water drop is completely taken up by the soil. The  
30 observation was stopped after exceeding a time of 3600 seconds. Depending on the profile  
31 depths, WDPT tests were performed in several depths of the profile, covering the main soil

1 horizontation. In each depth, five sampling locations were used to traverse the profile, and the  
2 tests were repeated three times per location, resulting in 15 WDPTs per depth (Figure 3b).  
3 The mean and maximum values of the WDPT test were classified after Bisdom et al. (1993)  
4 (Table 1).

## 5 **2.4 Dye tracer experiments**

6 The dye tracer experiments were conducted in August 2011 before installation of the roofs  
7 and in August 2013 after two years of drought. For each experiment, an area of 80 x 120 cm  
8 was prepared by cutting smaller vegetation (grasses, herbaceous plants, and small tree  
9 offshoots), covering the surroundings with a thin plastic sheet, and dividing the area into three  
10 sub-areas with a size of 80 x 40 cm each (Figure 3 a). The experimental area was kept shaded  
11 and sheltered from rain in all weather conditions to minimize evaporation and uncontrolled  
12 water input during the experiments. Brilliant Blue FCF was diluted in water of local origin to  
13 a concentration of 4 g/l and was sprayed with a backpack nozzle sprayer for even distribution  
14 (Bachmair et al, 2009). For an overall application amount of 20, 40 and 60 mm, each sub-area  
15 was sprinkled with an intensity of 20 mm/h. The applied rainfall intensity of 20 mm/h reflects  
16 a heavy rainfall event in all regions, therefore the sprinkling amounts simulate one, two, and  
17 three hours of heavy rainfall. After sprinkling, the experimental area was covered with plastic  
18 sheets to prevent evaporation and further water input through eventual rain.

19 The next day (after waiting at least 12 hours), three vertical soil profiles per sub-area were  
20 prepared. Keeping a 10 cm buffer stripe at the beginning and between the individual sub-  
21 areas, every sub-area was divided in three sections, spaced 10 cm from each other (Figure  
22 3 a). To obtain the dye pattern, the surface of the excavated soil profiles was smoothed with a  
23 spatula and loose particles were removed with a brush, avoiding smearing. Stones were left in  
24 place and shaped into relief when needed. Roots were trimmed. Pictures were taken from each  
25 profile with a standard digital compact camera with a resolution of 10 megapixels  
26 (3648 x 2736 pixel). The single profiles were photographed with a ruler frame and a grey  
27 scale under even illumination and different illumination settings (Weiler and Flühler, 2004).  
28 The picture with the best image quality from each profile was used for further processing.

## 1 **2.5 Image processing and data analysis**

2 To objectively compare the flow pattern of the different profiles, we used the image analysis  
3 procedure developed by Weiler and Flühler (2004). We provide a short description of the  
4 process here, for more detailed information, refer to Weiler (2001) and Weiler and Flühler  
5 (2004). The image processing consists of three main steps. In the first step, geometric  
6 distortion of the image is corrected by establishing a relationship between the image pixel  
7 location and the true location on the soil profile. During this step, the image is also scaled  
8 such that, one pixel corresponds to a square of  $1 \times 1 \text{ mm}^2$ . In the second step, the spectral  
9 composition changes in daylight are balanced to ensure inter-picture comparability. This is  
10 done by a color adjustment of the image using the photographed grey scale. In the third step,  
11 the images are classified into stained and unstained areas. Applying a semi-supervised  
12 classification technique, a binary image of stained versus unstained areas is obtained. In  
13 contrast to the work of Weiler and Flühler (2004), we did not use the information of different  
14 dye tracer concentrations, due to the high heterogeneity of the background color. In this step,  
15 objects like stones and vegetation are manually digitized, too. All calculations were done with  
16 the programming language IDL (Interactive Data Language, Exelis Inc.).

## 17 **2.6 Dye pattern analysis**

18 To obtain objective measures to compare the dye patterns of the different profiles and sites,  
19 we derived three depth related variables of the binary images: (1) volume density, (2) surface  
20 density and (3) stained path width as basis for further delineation of flow processes. The  
21 volume density (VD) is similar to the frequently used dye coverage. It is defined as stained  
22 volume divided by the reference space and is originating from the methods of stereology,  
23 which relates a three-dimensional parameter to two-dimensional measurements (Weibel  
24 1979). Surface density (SD) is defined as surface area of an object divided by the volume of  
25 the reference space. Surface density provides information on the size and number of features:  
26 a high SD is caused by a large number of small objects, whereas a low SD indicates less but  
27 larger objects (Weiler 2001). The stained path width (SPW) is derived by measuring the width  
28 of every stained object at a certain depth. The SPW of every depth were classified into three  
29 classes of  $< 20 \text{ mm}$ ,  $20 - 200 \text{ mm}$ , and  $> 200 \text{ mm}$  (Weiler and Flühler 2004). The sum of the  
30 three SPW classes per depth corresponds to the VD of the regarding depth. Using the  
31 frequency distribution of the SPW of every depth, the dye pattern can be related to distinct  
32 flow processes. For example, macropore flow with low interaction can be identified by long



1 and narrow stains, whereas macropore flow with mixed interaction shows a broader  
2 distribution of shapes (Weiler and Flühler 2004). The classification introduced by Weiler and  
3 Flühler (2004) was used to distinguish five flow processes, depending on the proportion of  
4 stains in each SPW class: two types of matrix flow ((1) homogeneous and (2) heterogeneous)  
5 and three types of macropore flow ((3) low, (4) mixed and (5) high interaction with matrix),  
6 where interaction is understood as the lateral water flow from macropores into the  
7 surrounding soil matrix (Weiler and Naef 2003). To assess the differences in the VD values  
8 between the treatments (pre-drought, control, drought), the Kruskal-Wallis test and the  
9 Nemenyi post-hoc test were applied, using R (version 3.2.3, The R Foundation for  
10 Statistical Computing, 2015) and the package “PMCMR” (version 4.1 by Thorsten Pohlert)  
11 within. Differences between treatments were supposed significant, when p-values are  $\leq 0.01$ .

12

### 13 **3 Results**

#### 14 **3.1 Soil moisture**

15 Figure 4 shows the normalized cumulated sums of the soil moisture measurements of the  
16 control and the drought plots over the course of two years. All plots developed a soil moisture  
17 deficit compared to the control plots in the upper 5 cm of the soil, as shown by the black line  
18 below the 1:1 line. The water deficit is also transduced to the 15 cm and 30 cm depths in both  
19 Schwäbische Alb plots and in the coniferous plot of Hainich-Dün, but is generally less  
20 pronounced. The plots at the Schorfheide-Chorin site show no deficit (deciduous plot) or even  
21 a small plus in soil moisture (coniferous plot) compared to the control plot. The sandy soils of  
22 Schorfheide-Chorin are already very dry without drought treatment. The reverse moisture  
23 effect might be caused by [effects associated with the installation and placement of the probes](#)  
24 [\(e.g. preferential flow along probe cables, concentration of throughfall – although installations](#)  
25 [were performed with great care\), or](#) root effects, for example hydraulic redistribution.  
26 However, we did not find any signs for hydraulic redistribution in the data. The deciduous  
27 plot of the Hainich-Dün site experienced major probe failures due to animal damage during  
28 the summer month of 2012 and again in 2013. Therefore, only the data taken during the  
29 winter month could be used for the comparison. For this reason, the data do not cover the  
30 months with the highest expected soil moisture deficits.

## 1 **3.2 Soil water repellency**

2 Figure 5 shows the results of the WDPT test in 2013. All drought treated plots at all sites –  
3 coniferous and deciduous – exhibit water repellency (WDPT data from control plot under  
4 coniferous not available). All control plot soils are wettable (WDPT class 1) or feature at least  
5 lower water repellency than the drought treated plots. The coniferous plots under drought of  
6 Hainich-Dün and Schwäbische Alb showed higher WDPTs than the deciduous plots. This is  
7 valid for both mean and maximum values. The Schorfheide-Chorin deciduous plot showed  
8 higher water repellency than the coniferous plot. In all soil profiles, water repellency is  
9 highest in the topsoil and diminishes at a depth of about 20 cm. However, in the Schorfheide-  
10 Chorin deciduous plot, water repellency is present up to a depth of 50 – 60 cm. When present,  
11 strong to severe water repellency is dominant in the measured drought treated plots. Only the  
12 Hainich-Dün deciduous plot soil is classified as wettable in average and the Schorfheide-  
13 Chorin coniferous plot as slightly water repellent. Highest values in mean and maximum  
14 water repellency were found in the coniferous plots Hainich-Dün (mean 941 s; max 3600 s) in  
15 about 10 – 15 cm depth and in the Schwäbische Alb (mean 990 s; max 2340 s) in the topsoil  
16 (Figure 5).

## 17 **3.3 Dye tracer experiments and dye pattern analysis**

### 18 **3.3.1 Comparison between pre-drought pattern and control pattern**

19 Differences between pre-drought and control plots (without drought treatment) reflect  
20 differences in soil structure, texture and moisture due to a distance of 20 - 40 m between the  
21 drought and reference plot, but may also include time dependent changes of the soil  
22 characteristics, which are independent from the drought treatment. To ensure validity of the  
23 dye pattern analyses, it is necessary to assure comparability among the plots. To exclude time  
24 dependent changes as reasons for differences in pre-drought and drought treated dye patterns,  
25 the pre-drought pattern were checked against the pattern of the control plots. Figure 6  
26 compares the pre-drought pattern and the control pattern of the deciduous plots. In addition,  
27 Figure 7 provides boxplots of VD for different depths of the pre-drought and the control  
28 profiles for direct comparison.

29 The Schwäbische Alb pre-drought plot (Figure 6, top left) shows high VD in the top 10 cm in  
30 all profiles. The 40 mm and 60 mm sprinkling volume profiles show a high SPW in the top  
31 5 to 10 cm. On the control plot (Figure 6, top right) also large areas of the profiles top 10 cm

1 are stained, but the VD and SPW are not as high as the pre-drought profiles. This is especially  
2 evident in the VD boxplots of the upper 0 – 10 cm (Figure 7, top). All Schwäbische Alb  
3 profiles have high stone contents, in some cases exceeding 50 % of the profile width (Figure  
4 6). Below 10 cm depth, the control plot profiles are almost completely stained. This pattern is  
5 similar to the pre-drought profiles. In general, the patterns of the control profiles are similar in  
6 VD, SPW values, and distribution to the 20 mm pre-drought profile. The 60 mm control  
7 profile is reflecting the high VD and SPW values in top layers, which are characteristic of the  
8 40 mm and 60 mm pre-drought profiles.

9 The Hainich-Dün pre-drought profiles (Figure 6, center left) show low to medium SPW in all  
10 depths. VD values are high in the top 5 cm in all profiles and between 10 cm to 30 cm in the  
11 40 mm and 60 mm sprinkling amount profiles. The 20 mm profile displays only small VD  
12 values below 10 cm depth. All profiles have a medium to high stone content (30 – 60 %)  
13 below 30 cm depth. The control plot profiles (Figure 6, center right) are very similar in VD  
14 and SPW to the pre-drought profile pattern, but with generally lower VD in the 60 mm  
15 sprinkling amount profile (Figure 7, center right). Except for the 20 mm profile, which  
16 displays no stones, the control plot profiles have a medium to high stone content below 25 cm  
17 depth. In all profiles, large areas of the profile stayed unstained. However, although having a  
18 low VD in top layer, the 60 mm control plot profile is not following the pronounced drop in  
19 VD between 5 cm and 10 cm depths and the subsequent rise between 15 cm and 25 cm, which  
20 is characteristic for all other profiles (pre-drought and control). These distinct differences are  
21 apparent in the boxplots (Figure 7)

22 In the Schorfheide-Chorin pre-drought profiles (Figure 6, bottom left), high VD and SPW  
23 values are present. The highest VD and SPW values can be found in the 60 mm sprinkling  
24 amount profile. Below 10 cm depth, the 20 mm pre-drought profile displays only small to  
25 medium SPW and – in comparison to the 40 mm and 60 mm profiles – small VD values. The  
26 control plot profiles (Figure 6, bottom right), show in general high VD and SPW values, but  
27 have lower values in the top 10 cm than the pre-drought profiles (Figure 6 bottom right and  
28 Figure 7, bottom). This is more apparent in the 20 mm and 40 mm profiles (Figure 7). In the  
29 pre-drought and control plot, infiltration reached down to depths over 70 cm and no stones are  
30 present.

31 To summarize, the comparison between the pre-drought and control plots showed a broad  
32 agreement. Differences, that need to be accounted for, are the lower VD in the profile top

1 layers, especially at the Schwäbische Alb and Schorfheide-Chorin site. These differences  
2 might be due to spatial heterogeneities, e.g. slight differing in soil layer boundary depths,  
3 given the distance between the control and the pre-drought plots (15 m to 30 m). In addition,  
4 the initial conditions (soil moisture) were also slightly different possibly resulting in the  
5 observed differences. Choosing 10 cm steps for statistical comparison of the VD may in  
6 addition introduce differences, if soil layer boundary depths differ. Therefore, not only the  
7 VD, but also the SPW and the determined flow processes need to be taken into account for  
8 comparison. However, the pre-drought and drought experiment were performed in close  
9 vicinity (1 m). In the Hainich-Dün, the drop and rise of VD in all profiles points to a soil layer  
10 boundary effect on infiltration. This is not time dependent and present in both pre-drought and  
11 control profiles, therefore the comparability between the pre-drought and drought pattern is  
12 not affected.

### 13 **3.3.2 Comparison between pre-drought pattern and drought pattern**

14 As can be seen in Figure 8, all plots show marked differences between pre- and after-drought  
15 infiltration patterns. The clayey and loamy sites (Schwäbische Alb and Hainich-Dün) develop  
16 unstained (=unwetted) areas in the topsoil layers. This is more pronounced in the coniferous  
17 plots, where unstained areas are already visible in the pre-drought infiltration pattern. Figure 9  
18 compares VD in boxplot for different depths of the drought and pre-drought profiles including  
19 the statistical significance.

20

#### 21 *Schwäbische Alb coniferous plot*

22 At the Schwäbische Alb site, medium to low volume densities (VD) were found on the pre-  
23 drought coniferous plot throughout the whole profile for the 20 mm and 40 mm sprinkling  
24 depth and high VD for 60 mm sprinkling depth (Figure 8, top left). The drought 40 mm and  
25 60 mm profiles are lower in VD in the top layers (0 – 10 cm), than the pre-drought profiles  
26 (Figure 8 and 9, top left); the 40 mm profile is displaying even unstained areas (no VD). The  
27 20 mm pre-drought profile is already very low in VD, therefore the differences to the after  
28 drought profile is not distinct (Figure 8, top left). The drought coniferous plot shows a rise of  
29 VD culminating around 20 cm depth (Figure 8, top) for all sprinkling amounts (20 mm,  
30 40 mm and 60 mm). Below 20 cm depth, the 20 mm and 40 mm profiles show (Figure 9, top

1 left) higher VD in the after drought profiles than in the pre-drought profiles, whereas the  
2 60 mm profile show the same extent of VD in the drought and in the pre-drought profile.

3 The stained path ways (SPW) of the Schwäbische Alb coniferous pre-drought profiles are  
4 small to medium in the 20 mm and 40 mm profiles and high in the 60 mm profile (Figure 8,  
5 top left). After drought, low to medium SPW are dominant in the 20 mm and 60 mm profiles;  
6 high SPW values are occurring in the 40 mm profile below 20 cm. The flow processes  
7 identified in this depth as matrix flow, are caused by local saturation due to low  $K_s$  (Figure 8,  
8 top left). The dominating flow types in the pre-drought profiles are identified as macropore  
9 flow, with low, mixed and high interaction depending on soil layer and infiltration volume.  
10 Dominating flow types in the drought plot are macropore flow with low, medium and high  
11 interaction.

12

### 13 *Schwäbische Alb deciduous plot*

14 The Schwäbische Alb deciduous plot shows in the 40 mm and 60 mm pre-drought profiles  
15 high SPW and in all infiltrating volumes high VD in the top layer (0 – 10 cm; Figure 8 and  
16 Figure 9, top right). Medium to high VD are maintained throughout the whole 40 mm and  
17 60 mm profiles, and to lesser extend in the 20 mm profile. The drought profiles show lower  
18 VD in the top 10 cm, compared to the pre-drought profiles (Figure 9, top right). Below 20 –  
19 25 cm depths, the 20 cm and 40 cm drought profiles show higher VD than the pre-drought  
20 profiles. However, the drought profiles are more similar in shape to the VD pattern of the  
21 control than to the pre-drought profiles (Figure 6, top). Also, the stone contents in the three  
22 pre-drought profiles are higher than in the drought profiles (Figure 8, top right).

23 The dominating flow types in the Schwäbische Alb deciduous pre-drought profiles are  
24 identified as macropore flow with low, mixed, and high interaction, and as matrix flow,  
25 depending on soil layer and infiltration volume (Figure 8, top right). The flow processes  
26 identified in the top layers of the 40 mm and 60 mm pre-drought profiles as matrix flow are  
27 caused by local saturation due to low  $K_s$ . The dominating flow types of the drought deciduous  
28 profiles are identified as macropore flow with low, mixed, and high interaction (Figure 8, top  
29 right).

30

### 31 *Hainich-Dün coniferous plot*

1 The Hainich-Dün coniferous pre-drought profiles show low VD for all sprinkling amounts,  
2 especially in the topsoil between 4 cm and 22 cm (Figure 8, center left). The 20 mm and  
3 40 mm pre-drought profiles show unstained areas (no VD). The small VD values are even  
4 more pronounced in the drought profiles (Figure 8 and Figure 9, center left), in which all  
5 profiles exhibit unstained areas. Below the unstained layer, the VD rises to a maximum in 15  
6 to 20 cm depth and drops again around 30 cm depth. The 20 mm and 60 mm drought profiles  
7 show throughout all depths low VD (Figure 8, center left).

8 In all Hainich-Dün coniferous pre-drought profiles, no large SPW occur and flow types are  
9 classified as macropore flow with low, mixed and high interaction. This applies also for the  
10 20 mm and 60 mm drought profiles. In contrast, the 40 mm drought profile exhibits high SPW  
11 between 15 cm and 25 cm depth. Therefore, the flow types in this depths are identified as  
12 matrix flow (Figure 8, center left). The main flow types in the coniferous drought profiles are  
13 macropore flow with low, mixed and high interaction. The pre-drought profiles are dominated  
14 by macropore flow with low and mixed interaction. In both, pre-drought and drought profiles,  
15 the stone content is comparable (Figure 8, center left).

16

#### 17 *Hainich-Dün deciduous plot*

18 The Hainich-Dün deciduous drought profiles exhibit smaller VD in the top 5 cm compared to  
19 the pre-drought profiles (Figure 8 and Figure 9, center right). Unstained areas are present in  
20 the top 5 – 10 cm of the 20 mm drought profile. The 40 mm and 60 mm pre-drought profiles  
21 show high VD values between 10 cm and 25 cm. High VD values are also present in the  
22 drought profiles, maintaining high values throughout the whole profile. While no high SPW  
23 values are found in the pre-drought profiles, high SPW values can be found in the 40 mm  
24 drought profile between 10 cm and 30 cm and in the 60 mm drought profile between 10 cm  
25 and 40 cm (Figure 8 and Figure 9, center right). The flow types of the deciduous pre-drought  
26 profiles are classified as macropore flow with low, mixed, and high interaction. The drought  
27 profiles are also classified as macropore flow with low, mixed, and high interaction and,  
28 where high SPW values occur, as matrix flow (homogeneous and heterogeneous) (Figure 8,  
29 center right). The stone contents of the pre-drought and drought profiles are increasing with  
30 depth below 25 – 30 cm; the drought profiles are exhibiting a slightly higher stone content  
31 than the pre-drought profiles (Figure 8).

32

1 *Schorfheide-Chorin coniferous plot*

2 The pre-drought pattern of the Schorfheide-Chorin site show high SPW and VD in the top  
3 layers (0 - 10 cm depth) decreasing with depth (Figure 8, bottom left). While the 20 mm and  
4 40 mm pre-drought profiles show a maximum infiltration depth of about 45 cm and 30 cm,  
5 respectively, the 60 mm pre-drought profile is stained below 70 cm, exhibiting medium VD  
6 values (Figure 8, bottom left). High SPW values are found in the 20 mm pre-drought profile  
7 up to a depth of 15 cm and in the 40 mm and 60 mm profiles up to 10 cm and 30 cm,  
8 respectively. The drought profiles of the coniferous plots show far lower VD values in the top  
9 layers compared to the pre-drought profiles (Figure 8 and Figure 9, bottom left). The 40 mm  
10 drought profile is exhibiting even an unstained layer in about 5 cm depth. High SPW values  
11 can be found in the 40 mm and 60 mm drought profile, not in the top layers, but between  
12 20 cm and 25 cm depth (40 mm profile), and between 10 cm and 25 cm depth (60 mm profile)  
13 (Figure 8, bottom left). This is reflected in the flow type classification. Whereas matrix flow  
14 is dominating the top layers in pre-drought profiles (at least the top 10 cm), matrix flow is  
15 occurring below 10 cm depth in the 40 mm and 60 mm drought profiles (Figure 8, bottom  
16 left).

17

18 *Schorfheide-Chorin deciduous plot*

19 The Schorfheide-Chorin deciduous pre-drought and drought patterns do not exhibit much  
20 differences in shape and in VD values in the 20 mm and 40 mm profiles (Figure 8, bottom  
21 right). The largest differences in VD can be found in the top 10 cm of the 20 mm profiles and  
22 in the 60 mm profile (Figure 9, bottom right). In addition, the 20 mm drought profile exhibits  
23 an unstained layer around 40 cm depth (Figure 8, bottom right). The difference between pre-  
24 drought and drought is more evident in the SPW values: Whereas high SPW values are found  
25 in the 40 mm and 60 mm pre-drought profiles in the top and bottom half of the profile, high  
26 SPW values are found in the drought profile in the bottom half, plus a small layer of two  
27 centimeter of high SPW around 10 cm depth in the 40 mm profile (Figure 8, bottom right). In  
28 the Schorfheide-Chorin deciduous pre-drought profiles, flow types of the 40 mm and 60 mm  
29 are dominated by matrix flow (Figure 8, bottom right). However, all profiles in the pre-  
30 drought plots have, a proportion of macropore flow. In the drought profiles, matrix flow is  
31 only occurring in bottom half of the 40 mm and 60 mm profiles.

32

1 To summarize, compared with pre-drought infiltration pattern, the drought pattern of all plots  
2 reveal differences in infiltration processes. For example, over 90% of the depths ranges show  
3 significant differences in VD between the drought and pre-drought site. Clayey and loamy  
4 soils behave similarly, developing unwatered soil layers. High SPW values in 20 to 30 cm  
5 depth of the drought pattern indicate local saturation. In sandy soils, the change from high  
6 SPW values of the pre-drought pattern to medium and low in the drought pattern exhibit a  
7 change from front-like to a more scattered infiltration. In general, the effects were more  
8 pronounced at the coniferous plots. These findings correspond well with the results of the  
9 WDPT tests: In the clayey and loamy soils (except Hainich-Dün deciduous plot), the  
10 unstained topsoil layers are coinciding with the high WDPTs (Figure 5). Coniferous plot  
11 Hainich-Dün stays unwatered up to a depth of about 15 to 20 cm and Schwäbische Alb plots to  
12 a depth of about 10 cm, which is corresponding to the depths where the highest WDPT values  
13 were observed (Hainich-Dün: WDPT class 4; Schwäbische Alb: WDPT classes 4 and 3,  
14 respectively). In the sandy soils of the Schorfheide-Chorin profiles, low SPW values  
15 correspond to high WDPTs (class 2 and 3). Below the water repellent zone, SPW values are  
16 increasing again (Figure 5, bottom).

17

## 18 **4 Discussion**

### 19 **4.1 Infiltration patterns and influence of main tree species**

20 The comparison of pre-drought infiltration patterns of the drought plots with patterns of the  
21 control plots (without drought treatment) showed broad agreements. All control plot profiles  
22 are comparable to the pre-drought plot profiles, including differences that can be addressed to  
23 small scale heterogeneities of soil properties. When interpreting the patterns, the differences  
24 in VD in the top layers of all plots need to be taken into account. When doing this, at all sites,  
25 the dye experiments before and during drought conditions can be directly compared.

26 In this study, it was hypothesized that the induced drought alters infiltration patterns due to  
27 changes in soil hydraulic properties (e.g., soil water repellency and forming of shrinkage  
28 cracks) and the main tree species is having an effect on the magnitude of the response. The  
29 results of the infiltration experiment show a clear evidence for changes in infiltration pattern  
30 as well as the importance of tree species on infiltration pattern: Schwäbische Alb plots have  
31 clayey soils with a high stone content, and show, in pre-drought and control plots, a slow and



1 even infiltration. The drought-treated plots developed large areas with small volume densities  
2 and SPWs in the topsoil, while for deeper layers, broad stains (large SPW) were observed  
3 which cover the profiles for the most part (high VD). This is typical for preferential flow that  
4 follows the shrinkage cracks of clayey soils or biopores of roots or soil fauna (Dekker and  
5 Ritsema 2000, Hendrickx and Flury 2001, Hardie et al. 2011). Water infiltrates quickly to  
6 deeper layers, bypassing a large proportion of the soil matrix. In deeper soil layers where the  
7 cracks or biopores end, local saturation occurs, and lateral redistribution into the soil matrix  
8 due to the now lower infiltration capacity and velocity can be observed. This also explains the  
9 similar pattern in the loamy Hainich-Dün soils.

10 A trend to more preferential flow was also observed in the Hainich-Dün plots, where the  
11 dense and loamy soils are also prone to shrinkage. Furthermore, in the Hainich-Dün drought  
12 profiles unstained (i.e., unwetted) areas in the topsoil layers were observed. This is more  
13 pronounced in the coniferous plot, where unstained areas were already visible in the pre-  
14 drought experiment. Preferential flow does not only originate from cracks and biopores, but  
15 also from textural boundaries and instable wetting fronts (Doerr and Ritsema. 2006, Hendrickx  
16 and Flury 2001). Unstable wetting fronts can occur due to air entrapment or hydrophobicity,  
17 which effectively hinders infiltration and redirects the water to structural and textural  
18 preferential flow paths (Doerr and Ritsema 2006). The unwetted topsoil layers of the Hainich-  
19 Dün coniferous plots can be explained by the combination of severe water repellency and  
20 shrinkage cracks acting as effective bypasses.

21 In contrast to the other sites, Schorfheide-Chorin soils are sandy and highly permeable with  
22 low stone content. In both Schorfheide-Chorin plots, the infiltration patterns changed from a  
23 regular front-like stable infiltration to unstable, more scattered and fingered infiltration  
24 patterns. Following the conceptualization of unstable flow in water repellent soils by Ritsema  
25 et al. (1993, 2000), water flows, after entering the soil, through preferential pathways through  
26 the water repellent layer and distributes laterally in the divergence layer underneath. In fact,  
27 such flow patterns were pronounced in the Schorfheide-Chorin deciduous drought plot:  
28 medium VD and SPW up to a depth of 50 - 60 cm and larger SPW in the layer beneath. This  
29 fits with the results of the WDPT tests, which show a slight to strong water repellency in the  
30 top 50 - 60 cm of the profile.

31 In general, drought induced major changes on the infiltration behavior of the examined soils.  
32 Clayey and loamy soils developed preferential flow. In these soils, the bypassing of the top

1 10 cm – 20 cm is fostered by water repellency, leading to unwetted topsoil layers. Sandy soils  
2 developed fingered infiltration patterns, due to the forming of a water repellent layer. In all  
3 three sites, the effects of the drought treatment were more pronounced in soils with coniferous  
4 main tree species than with deciduous main tree species.

5

## 6 **4.2 Water repellency**

7 In this study, it was hypothesized that the artificial induced drought will increase soil water  
8 repellency depending on the main tree species and soil properties. The highest water  
9 repellency was found in the coniferous plots of Schwäbische Alb and Hainich-Dün. Soils  
10 under coniferous trees often feature acidic soil conditions, which promote water repellency  
11 (Orfánus et al. 2014). In a study by Orfánus et al. (2014) liming practices and associated rise  
12 of pH-values significantly reduced water repellency of former pine forest soils. Jost et al.  
13 (2004) explained the difference in recharge under a beech and a spruce forest stand, with the  
14 higher hydrophobicity, and therefore the hindering of infiltration, combined with higher  
15 surface runoff of the spruce stand. This is in contrast to the findings of Buczko et al. (2006),  
16 who found the highest proportion of water repellent soils in pure beech stands compared to  
17 pure pine and mixed stands on sandy soils. However, in our study the sandy Schorfheide-  
18 Chorin plots showed higher mean WDPTs in the deciduous (beech) plot, than on the  
19 coniferous (pine) plot.

20 The soil texture can also influence the water repellency: A study of Gonzalez-Penalozza et al.  
21 (2013) suggests that water repellency is related to soil particle size. They induced water  
22 repellency by using different concentrations of stearic acid on samples of fine, medium, and  
23 coarse sand. Water repellency was extreme in coarse textured samples. The authors explained  
24 that by the lower specific surface compared to fine textured samples and therefore smaller  
25 area that has to be covered by water repellent agents. We could not observe this effect in our  
26 sandy soils. The coarser textured coniferous plot was less water repellent than the finer  
27 textured deciduous Schorfheide-Chorin plot. However, water repellency can originate from a  
28 broad range of factors. The degree of water repellency of a soil also depends on the amount  
29 and type of organic matter that is incorporated in it (DeBano 1981, Bisdom et al. 1993,  
30 Buczko et al. 2006, Vogelmann et al. 2013), the age and type of forest and litter type (Neris et  
31 al. 2013).

1 Several studies detected a significant impact of spruce litter on infiltration processes, either by  
2 hydrophobicity (Schume et al. 2004) or interception (Neris et al. 2013). Schume et al. (2004)  
3 found that spruce litter can intercept up to 5 mm of precipitation and Neris et al. (2013) found  
4 infiltration rates of 20 mm/h compared to that of 50mm/h of deciduous stands, doubling the  
5 runoff of the sites. In this study, we did not record the interception of the litter layer, which  
6 may have altered the total amount of water infiltrating into the soil. However, a natural litter  
7 layer is always present and intercepts precipitation (e.g. Gerrits et al. 2010). By keeping the  
8 natural litter layer in our experimental setup, our test results include the two influencing  
9 factors of the systems natural response in the infiltration pattern: The redistribution of  
10 incoming precipitation by the litter layer, leading to more spatial heterogeneous water input in  
11 the soil, compared to a soil with removed or hydrophilic litter layer. The measured infiltration  
12 pattern is a result of both factors, giving a more natural representation than a separate  
13 observation of litter layer and soil response.

14 Furthermore, the plants of the forest understory can also influence hydrophobicity of the soil;  
15 plants are covered with a cuticle composed of hydrophobic liquids, embedded in a polyester  
16 matrix and wax crystalloids (Holloway 1994, Barthlott and Neinhuis 1997). Water repellent  
17 plant coatings can be found in all plant life forms with a clear dominance in among herbs  
18 (Neinhuis and Barthlott 1997, Dekker and Ritsema 2000). It is even discussed that  
19 hydrophobic exudates might be a strategy for plants, microorganisms and fungi, to suppress  
20 germination and growth of competing vegetation by reducing evaporation and nutrient  
21 leaching (Doerr et al. 2007).

22 Hydrophobicity is dependent on the moisture status of the soil, which is defined by Doerr and  
23 Thomas (2003) as critical moisture or transition zone. Vogelmann et al. (2013) found a  
24 critical water threshold of 0.36 to 0.57 cm<sup>3</sup> cm<sup>-3</sup> beyond which hydrophobic soils become  
25 hydrophilic, varying as a function of soil organic matter content. In contrast to the findings of  
26 Doerr and Thomas (2003), we found very similar water contents in drought treated and  
27 control soils, but very different hydrophobicity conditions. This indicates that the “drought-  
28 history” or generally the climatic condition in the past of a soil is more important than the  
29 actual antecedent soil moisture status regarding hydrophobicity and infiltration behavior.

30 In our rainfall exclusion experiment drought stress was not intense enough to induce mortality  
31 or strong changes in above-ground biomass of a particular species (Gimbel et al. 2015).  
32 Nevertheless, drought and water repellency may promote the die-off of fine roots, which

1 thereupon contribute to the total organic matter in the soil. The amount of soil organic matter  
2 and its composition has a strong influence on the strength of water repellency (e.g.  
3 Vogelmann et al. 2013, Bisdorf et al. 1993, DeBano 1981). Therefore, the die-off of fine  
4 roots may lead to a self-reinforcing circle of water repellency.

5 Soil organic matter can form micro- and macro-aggregates by acting as binding agent between  
6 soil components (e.g. Tisdall and Oades 1982, Annabi et al. 2011) or by covering soil  
7 particles (e.g., Vogelmann et al. 2013 a). Vogelmann et al. (2013 b) concluded in their study,  
8 that water repellency leads to slower wetting of soil aggregates. Therefore, cohesive forces  
9 hold up longer, which increases the resistance to disaggregation and thus, indirectly aiding in  
10 maintaining soil structure. Terrestrial fungi are also in the focus of research concerning soil  
11 water repellency and aggregation (e.g. Tisdall and Oades 1982, Rillig and Mummey 2006,  
12 Chau et al. 2012). Zheng et al. (2014) found in three of nine species of ectomycorrhizal fungi  
13 associated with *Pinus sylvestris* seedlings increased soil water repellency and in six of nine  
14 species an increase of water stable aggregation. In our study, only the coniferous plot in  
15 Schorfheide-Chorin has *Pinus sylvestris* as main tree species. In fact, the plot showed slight  
16 (mean values) to strong (maximum values) water repellency in the top 20 cm. Nevertheless,  
17 the WDPT values of the deciduous plot in this area indicated stronger water repellency (in  
18 mean and maximum values).

19 All of our experimental plots showed clear response to the drought treatment, irrespective of  
20 their soil type and vegetation cover. Especially the fast bypassing of the topsoil layer and the  
21 developing of unstained and hence not wetted areas may bear consequences in the upcoming  
22 climate change. Sorption and degradation of contaminants is strongest in the topsoil and  
23 decrease with soil depth (Hendrickx and Flury 2001). Thus, bypassing of the topsoil soil  
24 matrix foster early arrival times and high concentrations of contaminants in the groundwater,  
25 which was shown by several tracer field studies (e.g. Hendrickx and Flury 2001, Ritsema et  
26 al. 1997, Hardie et al. 2011). Once formed, dry zones persist further wetting and additional  
27 water infiltrates through already existing preferential pathways, further stabilizing established  
28 flow paths (Dekker and Ritsema 2000; Hagedorn and Bundt 2002). Under present climate  
29 conditions, soil water repellency is already a widespread phenomenon (Buczko et al. 2006).  
30 For the predicted climate conditions, where droughts will be more common, an even higher  
31 level of hydrophobicity is to be expected, according to the findings of our rainfall reduction  
32 experiments.

1

## 2 **5 Conclusions**

3 Two years of rainfall reduction equivalent to an annual drought with a 40 year return interval  
4 was sufficient to change the soil properties and hence the infiltration pathways of six forest  
5 soils independent of soil type and tree species. All drought treated soils, except one,  
6 developed slight to severe water repellency. Main tree species had a particular effect on  
7 hydrophobicity, but is only accounting for minor differences in infiltration pattern. The  
8 “drought-history” or generally the climatic condition in the past had more effect on the  
9 observed hydrophobicity and infiltration behavior than the actual antecedent soil moisture  
10 conditions of the soils. The results of this study suggest that drought effects on infiltration  
11 processes need to be considered in hydrological models to obtain realistic predictions  
12 regarding water quality and quantity in runoff and groundwater recharge.

13

## 14 **Acknowledgements**

15 The authors thank Emil Blattmann, Delon Wagner and Lukas Neuhaus for their technical  
16 assistance. Field support by Stefan Seeger, Frank Lehmann, Christoph Ries and Ruth Magh is  
17 gratefully acknowledged. The work has been funded by the DFG Priority Program 1374  
18 "Infrastructure-Biodiversity-Exploratories" (DFG-Refno. WE4598/3-1, GE1090/11-1, and  
19 BR1698/16-1). We thank the managers of the three exploratories, Swen Renner, Sonja  
20 Gockel, Andreas Hemp, Martin Gorke and Simone Pfeiffer for their work in maintaining the  
21 plot and project infrastructure, and Markus Fischer, the late Elisabeth Kalko, Eduard  
22 Linsenmair, Dominik Hessenmöller, Jens Nieschulze, Daniel Prati, Ingo Schöning, François  
23 Buscot, Ernst-Detlef Schulze and Wolfgang W. Weisser for their role in setting up the  
24 Biodiversity Exploratories project. Field work permits were issued by the responsible state  
25 environmental offices of Baden-Württemberg, Thüringen, and Brandenburg (according to §  
26 72 BbgNatSchG). The article processing charge was funded by the German Research  
27 Foundation (DFG) and the Albert Ludwigs University Freiburg in the funding programme  
28 Open Access Publishing.

29

## 1 **References**

- 2 Albert, K. R., Ro-Poulsen, H., Mikkelsen, T. N., Michelsen, A., van der Linden, L., and  
3 Beier, C.: Effects of elevated CO<sub>2</sub>, warming and drought episodes on plant carbon uptake in a  
4 temperate heath ecosystem are controlled by soil water status, *Plant, Cell & Environment*, 34,  
5 1207–1222, doi:10.1111/j.1365-3040.2011.02320.x, 2011.
- 6 Annabi, M., Y. Le Bissonnais, M. Le Villio-Poitrenaud, and S. Houot. Improvement of soil  
7 aggregate stability by repeated applications of organic amendments to a cultivated silty loam  
8 soil. *Agriculture, Ecosystems & Environment* 144(1):382–389.  
9 doi:10.1016/j.agee.2011.07.005, 2011.
- 10 Bachmair, S., Weiler, M., and Nützmann, G.: Controls of land use and soil structure on water  
11 movement: Lessons for pollutant transfer through the unsaturated zone, *Journal of Hydrology*,  
12 369, 241–252, doi:10.1016/j.jhydrol.2009.02.031, 2009.
- 13 Barthlott, W. and Neinhuis, C.: Purity of the sacred lotus, or escape from contamination in  
14 biological surfaces, *Planta*, 202, 1–8, doi:10.1007/s004250050096, 1997
- 15 Bisdorf, E., Dekker, L. W., and Schoute, J.: Water repellency of sieve fractions from sandy  
16 soils and relationships with organic material and soil structure, *Geoderma*, 56, 105–118,  
17 doi:10.1016/0016-7061(93)90103-R, 1993.
- 18 Buczko, U., Bens, O., and Hüttel, R. F.: Water infiltration and hydrophobicity in forest soils of  
19 a pine–beech transformation chronosequence, *Journal of Hydrology*, 331, 383–395,  
20 doi:10.1016/j.jhydrol.2006.05.023, 2006.
- 21 Bütof, A., von Riedmatten, Lars R., Dormann, C. F., Scherer-Lorenzen, M., Welk, E., and  
22 Brühlheide, H.: The responses of grassland plants to experimentally simulated climate change  
23 depend on land use and region, *Glob Change Biol*, 18, 127–137, doi:10.1111/j.1365-  
24 2486.2011.02539.x, 2012.
- 25 Chau, H.W., Y.K. Goh, V. Vujanovic, and B.C. Si: Wetting properties of fungi mycelium  
26 alter soil infiltration and soil water repellency in a  $\gamma$ -sterilized wettable and repellent soil.  
27 *Fungal Biology* 116(12):1212–1218. doi:10.1016/j.funbio.2012.10.004, 2012
- 28 da Costa, A. C. L., Galbraith, D., Almeida, S., Portela, Bruno Takeshi Tanaka, da Costa, M.,  
29 de Athaydes Silva Junior, João, Braga, A. P., de Gonçalves, Paulo H. L., de Oliveira, Alex  
30 AR, Fisher, R., Phillips, O. L., Metcalfe, D. B., Levy, P., and Meir, P.: Effect of 7 yr of

1 experimental drought on vegetation dynamics and biomass storage of an eastern Amazonian  
2 rainforest, *New Phytologist*, 187, 579–591, doi:10.1111/j.1469-8137.2010.03309.x, 2010.

3 DeBano, L.F.: Water repellent soils: a state-of-the-art. USDA For. Serv. Gen.Tech. Rep.  
4 PSW-46, pp. 21, 1981

5 DeBano, L. 2000. Water repellency in soils: a historical overview. *Journal of Hydrology* 231-  
6 232:4–32. doi:10.1016/S0022-1694(00)00180-3.

7 Dekker, L. and Ritsema, C. J.: Wetting patterns and moisture variability in water repellent  
8 Dutch soils, *Journal of Hydrology*, 231-232, 148–164, doi:10.1016/S0022-1694(00)00191-8,  
9 2000.

10 Dermody, O., Weltzin, J. F., Engel, E. C., Allen, P., and Norby, R. J.: How do elevated  
11 [CO<sub>2</sub>], warming, and reduced precipitation interact to affect soil moisture and LAI in an old  
12 field ecosystem?, *Plant Soil*, 301, 255–266, doi:10.1007/s11104-007-9443-x, 2007.

13 Doerr, S.H., Thomas, A.D., Soil moisture: a controlling factor in water repellency? In:  
14 Ritsema, C.J., Dekker, L.W. (Eds.), *Soil Water Repellency: Occurrence, Consequences and*  
15 *Amelioration*. Elsevier Science, pp. 137–149., 2003

16 Doerr, S. H. and Ritsema, C. J.: Water Movement in Hydrophobic Soils, in: *Encyclopedia of*  
17 *Hydrological Sciences*, John Wiley & Sons, Ltd, 2006.

18 Doerr, S. H., Ritsema, C. J., Dekker, L. W., Scott, D. F., and Carter, D.: Water repellence of  
19 soils: new insights and emerging research needs, *Hydrol. Process.*, 21, 2223–2228,  
20 doi:10.1002/hyp.6762, 2007.

21 English, N.B., Weltzin, J.F., Fravolini, A., Thomas, L., and Williams, D.G.: The influence of  
22 soil texture and vegetation on soil moisture under rainout shelters in a semi-desert grassland.  
23 *Journal of arid environments*, 63(1), 324-343, doi:10.1016/j.jaridenv. 2005.03.013, 2005.

24 FAO, 2006. *Guidelines for Soil Description*, 4th edition. Food and Agriculture Organization  
25 of the United Nations, Rome.

26 Federer, C. A., Vörösmarty, C., and Fekete, B.: Sensitivity of Annual Evaporation to Soil and  
27 Root Properties in Two Models of Contrasting Complexity, *J. Hydrometeor*, 4, 1276–1290,  
28 doi:10.1175/1525-7541(2003)004<1276:SOAETS>2.0.CO;2, 2003.

29 Fischer, Markus.; Bossdorf, O.; Gockel, S.; Hänsel, F.; Hemp, A.; Hessenmöller, D.; Korte,  
30 G.; Nieschulze, J.; Pfeiffer, S.; Prati, D.R.S.; Schöning, I.; Schumacher, U.; Wells, K.;

1 Buscot, F.; Kalko, E.K.V.; Linsenmair, K.E.; Schulze, E.-D.; Weisser, W.W.: Implementing  
2 large-scale and long-term functional biodiversity research: The Biodiversity Exploratories.  
3 *Basic and Applied Ecology*, 11, 6, 473–485, doi:10.1016/j.baae.2010.07.009, 2010

4 Gerrits, A. M. J., L. Pfister, and Savenije, H. H. G. 2010. Spatial and temporal variability of  
5 canopy and forest floor interception in a beech forest. *Hydrol. Process.* 24(21):3011–3025.  
6 doi:10.1002/hyp.7712.

7 Gimbel, K. F.; Felsmann, K.; Baudis, M.; Puhlmann, H.; Gessler, A.; Bruelheide, H.;  
8 Kayler, Z.; Ellerbrock, R. H.; Ulrich, A.; Welk, E.; Weiler, M.: Drought in forest understory  
9 ecosystems – a novel rainfall reduction experiment. *Biogeosciences*, 12, 961–975,  
10 doi:10.5194/bg-12-961-2015, 2015

11 Glaser, B., Jentsch, A., Kreyling, J., and Beierkuhnlein, C.: Soil-moisture change caused by  
12 experimental extreme summer drought is similar to natural inter-annual variation in a loamy  
13 sand in Central Europe, *Z. Pflanzenernähr. Bodenk.*, 176, 27–34,  
14 doi:10.1002/jpln.201200188, 2013.

15 González-Peñaloza, F. A., Zavala, L. M., Jordán, A., Bellinfante, N., Bárcenas-Moreno, G.,  
16 Mataix-Solera, J., Granged, A. J., Granja-Martins, F. M., and Neto-Paixão, H. M.: Water  
17 repellency as conditioned by particle size and drying in hydrophobized sand, *Geoderma*, 209-  
18 210, 31–40, doi:10.1016/j.geoderma.2013.05.022, 2013.

19 Hagedorn, F., and M. Bundt. 2002. The age of preferential flow paths. *Geoderma* 108(1-  
20 2):119–132. doi:10.1016/S0016-7061(02)00129-5

21 Hammel, K., Kennel, M., 2001. Charakterisierung und Analyse der Wasserverfügbarkeit und  
22 des Wasserhaushalts von Waldstandorten in Bayern mit dem Simulationsmodell BROOK90,  
23 *Forstliche Forschungsberichte München* 185.

24 Hardie, M. A., Cotching, W. E., Doyle, R. B., Holz, G., Lisson, S., and Mattern, K.: Effect of  
25 antecedent soil moisture on preferential flow in a texture-contrast soil, *Journal of Hydrology*,  
26 398, 191–201, doi:10.1016/j.jhydrol.2010.12.008, 2011.

27 Hendrickx, J. M. H. and Flury, M.: Uniform and preferential flow mechanisms in the vadose  
28 zone, *Conceptual models of flow and transport in the fractured vadose zone*, 149–187, 2001.

29 Holloway, P.J., 1994. Plant cuticles: physiochemical characteristics and biosynthesis. In:  
30 Percy, K.E. (Ed.). *Air Pollution and the Leaf Cuticle*, Springer, Berlin, pp. 1–13.



1 Jost, G., Schume, H., and Hager, H.: Factors controlling soil water-recharge in a mixed  
2 European beech (*Fagus sylvatica* L.)?Norway spruce [*Picea abies* (L.) Karst.] stand, *Eur J*  
3 *Forest Res*, doi:10.1007/s10342-004-0033-7, 2004.

4 Keesstra, S. D., Geissen, V., Mosse, K., Piirainen, S., Scudiero, E., Leistra, M., and van  
5 Schaik, L.: Soil as a filter for groundwater quality, *Current Opinion in Environmental*  
6 *Sustainability*, 4, 507–516, doi:10.1016/j.cosust.2012.10.007, 2012.

7 Kopittke, G. R., Tietema, A., Loon, E. E., and Asscheman, D.: Fourteen Annually Repeated  
8 Droughts Suppressed Autotrophic Soil Respiration and Resulted in an Ecosystem Change,  
9 *Ecosystems*, 17, 242–257, doi:10.1007/s10021-013-9720-x, 2014.

10 Meijer, S. S., Holmgren, M., and Van der Putten, W. H.: Effects of plant-soil feedback on tree  
11 seedling growth under arid conditions, *Journal of Plant Ecology*, 4, 193–200,  
12 doi:10.1093/jpe/rtr011, 2011.

13 Neinhuis, C. and Barthlott, W.: Characterization and Distribution of Water-repellent, Self-  
14 cleaning Plant Surfaces, *Annals of Botany*, 79, 667–677, doi:10.1006/anbo.1997.0400, 1997.

15 Neris, J., Tejedor, M., Rodríguez, M., Fuentes, J., and Jiménez, C.: Effect of forest floor  
16 characteristics on water repellency, infiltration, runoff and soil loss in Andisols of Tenerife  
17 (Canary Islands, Spain), *CATENA*, 108, 50–57, doi:10.1016/j.catena.2012.04.011, 2013.

18 Orfánus, T., Dlapa, P., Fodor, N., Rajkai, K., Sándor, R., and Nováková, K.: How severe and  
19 subcritical water repellency determines the seasonal infiltration in natural and cultivated  
20 sandy soils, *Soil and Tillage Research*, 135, 49–59, doi:10.1016/j.still.2013.09.005, 2014.

21 Ozolinčius, R., Stakėnas, V., Varnagirytė-Kabašinskienė, I., and Buožytė, R.: Artificial  
22 Drought in Scots Pine Stands: Effects on Soil, Ground Vegetation and Tree Condition,  
23 *Annales Botanici Fennici*, 46, 299–307, doi:10.5735/085.046.0406, 2009.

24 Phillips, O. L., Aragao, L. E. O. C., Lewis, S. L., Fisher, J. B., Lloyd, J., Lopez-Gonzalez, G.,  
25 Malhi, Y., Monteagudo, A., Peacock, J., Quesada, C. A., van der Heijden, G., Almeida, S.,  
26 Amaral, I., Arroyo, L., Aymard, G., Baker, T. R., Banki, O., Blanc, L., Bonal, D., Brando, P.,  
27 Chave, J., de Oliveira, A. C. A., Cardozo, N. D., Czimczik, C. I., Feldpausch, T. R., Freitas,  
28 M. A., Gloor, E., Higuchi, N., Jimenez, E., Lloyd, G., Meir, P., Mendoza, C., Morel, A.,  
29 Neill, D. A., Nepstad, D., Patino, S., Penuela, M. C., Prieto, A., Ramirez, F., Schwarz, M.,  
30 Silva, J., Silveira, M., Thomas, A. S., Steege, H. T., Stropp, J., Vasquez, R., Zelazowski, P.,  
31 Davila, E. A., Andelman, S., Andrade, A., Chao, K.-J., Erwin, T., Di Fiore, A., C., E. H.,

1 Keeling, H., Killeen, T. J., Laurance, W. F., Cruz, A. P., Pitman, N. C. A., Vargas, P. N.,  
2 Ramirez-Angulo, H., Rudas, A., Salamao, R., Silva, N., Terborgh, J., and Torres-Lezama, A.:  
3 Drought Sensitivity of the Amazon Rainforest, *Science*, 323, 1344–1347,  
4 doi:10.1126/science.1164033, 2009.

5 Prudhomme, C., Giuntoli, I., Robinson, E. L., Clark, D. B., Arnell, N. W., Dankers, R.,  
6 Fekete, B. M., Franssen, W., Gerten, D., Gosling, S. N., Hagemann, S., Hannah, D. M., Kim,  
7 H., Masaki, Y., Satoh, Y., Stacke, T., Wada, Y., and Wisser, D.: Hydrological droughts in the  
8 21st century, hotspots and uncertainties from a global multimodel ensemble experiment,  
9 *Proceedings of the National Academy of Sciences*, 111, 3262–3267,  
10 doi:10.1073/pnas.1222473110, 2014.

11 Puhlmann H, von Wilpert K. Test und Entwicklung von Pedotransferfunktionen für  
12 Wasserretention und hydraulische Leitfähigkeit von Waldböden. *Waldökol Landschaftsforsch*  
13 *Naturschutz*. 12: 61–71.

14 Rillig, M. C. and Mummey, D. L.: Mycorrhizas and soil structure. *New Phytologist* 171,  
15 41e53, 2006

16 Ritsema, C. J., Dekker, L. W., Hendrickx, J. M. H., and Hamminga, W.: Preferential flow  
17 mechanism in a water repellent sandy soil, *Water Resour. Res.*, 29, 2183–2193,  
18 doi:10.1029/93WR00394, 1993.

19 Ritsema, C. J., Dekker, L. W., van den Elsen, E. G. M., Oostindiel, K., Steenhuis, T. S., and  
20 Nieber, J. L.: Recurring fingered flow pathways in a water repellent sandy field soil, *Hydrol.*  
21 *Earth Syst. Sci.*, 1, 777–786, doi:10.5194/hess-1-777-1997, 1997.

22 Ritsema, C. and Dekker, L.: Preferential flow in water repellent sandy soils: principles and  
23 modeling implications, *Journal of Hydrology*, 231-232, 308–319, doi:10.1016/S0022-  
24 1694(00)00203-1, 2000.

25 Schume, H., Jost, G., and Hager, H.: Soil water depletion and recharge patterns in mixed and  
26 pure forest stands of European beech and Norway spruce, *Journal of Hydrology*, 289, 258–  
27 274, doi:10.1016/j.jhydrol.2003.11.036, 2004.

28 Selsted, M. B., Linden, L., Ibrom, A., Michelsen, A., Larsen, K. S., Pedersen, J. K.,  
29 Mikkelsen, T. N., Pilegaard, K., Beier, C., and Ambus, P.: Soil respiration is stimulated by  
30 elevated CO<sub>2</sub> and reduced by summer drought: three years of measurements in a multifactor

1 ecosystem manipulation experiment in a temperate heathland (CLIMAITE), *Glob Change*  
2 *Biol*, 18, 1216–1230, doi:10.1111/j.1365-2486.2011.02634.x, 2012.

3 Seneviratne, S.I., Luethi, D., Litschi, M., and Schaer, C.: Land-atmosphere coupling and  
4 climate change in Europe, *Nature*, 443, 205-209, doi:10.1038/nature05095, 2006

5 Suttle, B. K. and Thomsen, M. A.: Climate change and grassland restoration in California:  
6 Lessons from six years of rainfall manipulation in a North Coast grassland, *Madroño*, 54,  
7 225–233, doi:10.3120/0024-9637(2007)54[225:CCAGRI]2.0.CO;2, 2007.

8 Tisdall, J. M. and Oades, J. M.: Organic matter and water-stable aggregates in soils. *Journal*  
9 *of Soil Science* 33, 141e163, 1982

10 Vogelmann, E. S., Reichert, J. M., Prevedello, J., Consensa, C., Oliveira, A., Awe, G. O., and  
11 Mataix-Solera, J.: Threshold water content beyond which hydrophobic soils become  
12 hydrophilic: The role of soil texture and organic matter content, *Geoderma*, 209-210, 177–  
13 187, doi:10.1016/j.geoderma.2013.06.019, 2013 a.

14 Vogelmann, E.S., J.M. Reichert, J. Prevedello, G.O. Awe, and J. Mataix-Solera.:Can  
15 occurrence of soil hydrophobicity promote the increase of aggregates stability? *CATENA*  
16 110:24–31. doi:10.1016/j.catena.2013.06.009, 2013 b.

17 Weibel, E. R.: In: *Stereological Methods. Practical Methods for Biological Morphometry*,  
18 vol.1 Academia Press, London, 1979.

19 Weiler, M: *Mechanics Controlling Macropore flow during Infiltration – dye Tracer*  
20 *Experiments and Simulations*. ETH, Zürich, Switzerland. 151 pp, 2001.

21 Weiler, M., Naef, F.: An experimental tracer study of the role of macropores in infiltration in  
22 grassland soils. *Hydrological Processes* 17, 477–493, 2003

23 Weiler, M. and Flühler, H.: Inferring flow types from dye patterns in macroporous soils,  
24 *Geoderma*, 120, 137–153, doi:10.1016/j.geoderma.2003.08.014, 2004.

25 World reference base for soil resources, 2006: A framework for international classification,  
26 correlation and communication. 2006 ed. World soil resources reports 103. Food and  
27 Agriculture Organization of the United Nations, Rome.

28 Wu, C. and Chen, J. M.: Diverse responses of vegetation production to interannual summer  
29 drought in North America, *International Journal of Applied Earth Observation and*  
30 *Geoinformation*, 21, 1–6, doi:10.1016/j.jag.2012.08.001, 2013.

1 Zheng, W., E.K. Morris, and M.C. Rillig.: Ectomycorrhizal fungi in association with *Pinus*  
2 *sylvestris* seedlings promote soil aggregation and soil water repellency. *Soil Biology and*  
3 *Biochemistry* 78:326–331. doi:10.1016/j.soilbio.2014.07.015, 2014.  
4

1 *Table 1: Classification of water repellency by WDPT time, after Bisdorn et al. (1993)*


---

WDPT in s	Classification	Class
< 5	wettable	1
5 – 60	slightly water repellent	2
60 – 600	strongly water repellent	3
600 – 3600	severely water repellent	4
> 3600	extremely water repellent	5

---


2

3



1  
2  
3  
4  
5  
6

Figure 1: Location of the study sites (red squares) within Germany (light blue); South-West: Schwäbische Alb; center: Hainich-Dün; North-East: Schorfheide-Chorin.



1


2

3 Figure 2: Soil horizons, texture, and rock fractions of the six experimental plots. Soil type  
 4 classification according to the World reference base for soil (FAO 2006).

5

6

1



2


3 Figure 3: Scheme for profile excavation (a) and WDPT experiment (b). The 20 mm, 40 mm,  
4 and 60 mm in (a) denote the applied sprinkling volumes. For the WDPT experiment (b), five  
5 sampling locations (boxes) were used traversing the profile. On every sampling location, the  
6 tests were repeated three times.

7

8



1




2

3 Figure 4: Normalized cumulated sums of soil moisture of the drought versus the control  
4 subplots of the investigated soils. [The computations begin with the start of the rainfall](#)  
5 [reduction and end at the date of the dye tracer experiment 2013.](#)

6

7



1

2

3 Figure 5: Mean and maximum water drop penetration times (WDPTs) of the control (green)


4 and drought (red) plots. Orange lines and numbers refer to the WDPT classes after Bisdorf et

5 al. (1993) (see Table 2).


6



1 Figure 6: Comparison between stained path width (SPW) of pre-drought (2011) and control  
2 (2013) plot. Blue shades indicate the SPW classes. The sum of SPW is the volume density  
3 (VD) per depth. Grey and black indicate the VD of stones.  
4



- 1 Figure 7: VD boxplots of the drought and the pre-drought pattern. Depth ranges are omitted,
- 2 where one of the profile is shorter than the other. Statistically significant ( $p\text{-value} \leq 0.01$ )
- 3 differences between the treatments are marked with an asterisk.
- 4




1

2

1 Figure 8: Comparison between before drought (2011) and after drought (2013) stained path  
2 widths (SPW) and flow processes for coniferous and deciduous stand plots. The sum of SPW  
3 is the volume density (VD) per depth. Grey and black indicate the VD of stones.  
4





1

2

3 Figure 9: VD boxplots of the drought and the pre-drought pattern. Depth ranges are omitted,  
 4 where one of the profile is shorter than the other. Statistically significant ( $p\text{-value} \leq 0.01$ )  
 5 differences between the treatments are marked with an asterisk.