

Comments to the Author:

The paper investigates the role of drainage and soil storage properties on modulating climate-evapotranspiration relations for three mountainous catchments in western U.S. As mentioned by the reviewers, the study results are of interest for both field researchers and modellers. The paper is clearly written though the conclusions section will benefit from changes suggested by the reviewers.

The reviewers have raised a number of additional interesting questions that have been carefully responded by the authors, who have also clearly identified the strategy to address these issues in the revised paper. The number of revisions is minor but I would like to see a revised version. I am therefore recommending minor revisions in line the reviewer's suggestions and with my own assessment of the manuscript.

**We thank the editor for both the positive response to the original manuscript and for the opportunity to share our revisions to the full manuscript text.**

**Our revised manuscript includes all revisions mentioned in our response to the referees, including the abbreviated conclusion and focusing of the results and discussion. There are additional minor edits throughout the text that modify our use of the word 'soil' to 'subsurface storage' or 'geologic'. These edits are in direct response to referee #1's comment on the use of terminology "soil AWC" and our agreement that storage, in the physical world and in our model, includes more than the soil. Our edited manuscript even suggests a change to the paper's title to change 'Soil storage...' to 'Subsurface storage capacity', the full title would be:**

Subsurface storage capacity influences climate-evapotranspiration interactions in three western United States catchments

**REFEREE #1 COMMENTS**

Garcia and Tague present an interesting comparison of hydrologic partitioning in three catchments, reaching the conclusion that differences in landscape characteristics, specifically subsurface water storage, attenuates the role of climate in controlling ET. The paper is a nice example of how the timing and amount of precipitation interact with variable storage to control the fate of precipitation in mountain catchments and should be of interest to the range of observationists and modelers.

Although the authors focus on the implications for these three catchments, the take home messages potentially are applicable to a wide range of systems where the spatially and temporally explicit interplay between climate and landscape has the potential to result in different hydrologic responses in locations with similar mean climate.

Although I am generally supportive of the work, I do have a number of concerns and suggestions that I hope will focus the presentation.

**Reply: We thank the referee for the supportive comments and have addressed the concerns and incorporated many of the suggestions below.**

Shorten and focus conclusions to highlight key implications. The current take home points are somewhat buried, including both modeling issues (e.g. error introduced by absence of soil calibration) and broader science take homes (e.g. precipitation timing vs. storage interactions)

**Reply: We appreciate this suggestion and have removed/moved text that is better suited to the discussion and restructured the conclusions to focus on the bigger take-home points. The conclusion now reads as follows:**

We demonstrate how subsurface storage and drainage properties (AWC and parameters that control lateral redistribution) interact with climate-related drivers to influence ET in three western U.S. mountain watersheds with distinctive precipitation regimes. These watersheds reflect conditions found in many other western U.S. snow-dominated systems, where summer water availability is influenced by the magnitude of precipitation, timing of soil moisture recharge and spring temperature and its effect on snowmelt. We found that, for our three watersheds, estimates of longer-term average (15-year) watershed-scale ET vary across a range of physically realistic storage/drainage parameters. For all watersheds, the range in long term mean ET estimates across AWC estimates (e.g., mean ET at a high AWC versus mean ET at a low AWC) may be as large as inter-annual variation in ET, suggesting that the influence of AWC and drainage can be substantial.

Our results also point to the importance of lateral redistribution as a control on ET, particularly for CA-SIER. Only a few studies have emphasized the role of lateral redistribution in plot to watershed scale climate responses in the Western U.S. (Barnard et al., 2010; Tague and Peng, 2013). For the CA-SIER site, our model results suggest that there can also be interactions between AWC and hillslope to watershed scale redistribution as controls on ET. Lateral redistribution was less important for the CO-ROC, where summer precipitation was a more important contributor to annual ET values and the least important for the wetter OR-CAS site. Results emphasize that the role of subsurface properties, including both storage and drainage, will be different for different climate regimes.

These results have important implications both for predicting ET in basins where data is not available for calibration and for understanding and predicting the spatial variability of ET within a basin. AWC also affects the sensitivity of annual ET to climate drivers, particularly in the two more seasonally water-limited basins. Although the three watersheds show different responses of annual ET to these climate drivers, there are values of AWC that would eliminate these cross-basin differences. These sensitivities highlight the need for improved information on spatial patterns of subsurface properties to contribute to the development of science-based information on forest vulnerabilities to climate change. Improved accounting for plant accessibility to moisture has improved model-data ET comparisons in previous modeling studies at regional and global scales (Hwang et al., 2009; Tang et al., 2013; Thompson et al., 2011). With expected decreases in fractional precipitation received as snow with climate change (Diffenbaugh et al., 2013; Knowles et al., 2006), we might expect soil storage to play a more important role

in providing water for forests in the future. Improved understanding of how climate and subsurface storage/drainage combine to control ET can enhance our understanding of forest water stress related to increased mortality (van Mantgem et al., 2009). Western U.S. forests show substantial vulnerability to drought, with declines in productivity and increases in mortality and disturbance in drought years (Allen et al., 2010; Hicke et al., 2012; Williams et al., 2013). Understanding these ecosystems' responses to primary climate drivers is of particular concern given recent warming trends (Sterl et al., 2008) and multi-year droughts (Cook et al., 2004; Dai et al., 2004). Identifying the physical conditions in which our ability to estimate ET is most sensitive or limited by knowledge of subsurface geologic properties helps to prioritize regional data acquisition agendas. Integrating results from recent advances in geophysical measurements and models such as those emerging from Critical Zone Observatories in the U.S. and elsewhere (Anderson et al., 2008) will be essential for analysis of climate ET interactions.

#### **REFeree #1 COMMENTS CONTINUE BELOW**

The paper could and should be improved by explicitly addressing alternative explanations for the differences between the three catchments. For example, they vary significantly in size, elevation, and total precipitation and the differences between catchment responses plausibly could be explained by these factors.

Similarly, how do the specifics of climate across the three sites influence results? For example, what does PET look like across time and space for your study catchments? Presumably, higher elevations in CO are always energy limited, while lower elevations switch are water limited. In contrast, CA and OR experience the seasonal pattern in energy vs. water limitation that is your focus.

**Reply: Though text throughout the paper addresses how cross-site differences in physical characteristics are influencing the response of ET to climate, we have included an additional paragraph to the discussion (now third paragraph) to be explicit about how the range of responses we observe in model results are related to these characteristics:**

**The range of sensitivities of ET to climate in this study is a direct function of climatic and physical characteristics of the catchments presented in this study. For example, OR-CAS receives twice as much precipitation and spans a much lower elevation range than either CA-SIER or CO-ROC (Table 2). Because OR-CAS is considerably wetter, its sensitivity of ET to magnitude of annual  $P$  is lessened considerably. OR-CAS' lower elevations, and related mean winter temperatures, also result in smaller average snowpacks reducing the strength of spring temperature as an explanatory variable for ET. Differences between CA-SIER and CO-ROC largely reflect seasonal distribution of precipitation, and reflect the importance of summer precipitation in CO-ROC. While climate is the dominant factor, topographic differences are also important. As discussed above,**

**topographically driven flowpath convergence in CA-SIER tends to increase sensitivity of ET to parameters that influence lateral drainage. This effect is less evident in the other two watersheds. We also note that CO-ROC is considerably larger than our other two study sites and, as such, includes significant fractions of other land cover including rock and meadow. We expect the different vegetation types to influence the response of ET to climate.**

Abstract begins with winter-wet summer dry but CO-ROC receives 46% precipitation in growing season while other sites are much less. This is an important part of your paper but suggests using a more objective metric perhaps AET: PET to describe differences between supply and demand.

**Reply: P and PET are averages of annual sums. The aridity index, P:PET, is a helpful summary metric for normalizing how water-limited the catchments are. We added these values to Table 2.**

	CO-ROC	OR-CAS	CA-SIER
P:PET	0.9	2.3	1.2

On a related note, the introduction begins with Mediterranean climates, but CO is a cold continental climate; I'm not certain that OR is technically Mediterranean either.

**Reply: We appreciate this point and have removed the two occurrences of 'Mediterranean' as a description in the text --in the first line of the introduction, and the last paragraph of the introduction.**

The results section as written reads too much like a discussion with numerous references and comparisons other work, making it difficult to focus on the key points of this effort.

**Reply: We have edited the results section to move some of the discussion of results to the discussion section, and in some cases remove text that is in the discussion already:**

**Moved to discussion:**

Among the predicted consequences of increased temperatures are an earlier start to the vegetation growing season (Cayan et al., 2001), and an increase in vapor pressure deficits and water demand (Isaac & van Wijngaarden, 2012).

CA-SIER does not show a significant relationship between  $T_{AMJ}$  and ET because the effect of temperature is strongly dependent on the amount of snowpack the basin receives in a year (Tague & Peng, 2013), which is more variable than the amount of snowpack received in CO-ROC or OR-CAS. These results suggest that the dominant effect of warmer spring temperatures is earlier meltout of snowpack, which leads to more snowmelt lost as runoff and results in less net recharge. A mechanism we suggest for this loss of runoff is that soils are more likely to be saturated in spring months. Later into the growing season, increased ET demands will have depleted soil

stores and throughfall/snowmelt will enter the soil matrix and be available for plant water use.

**Removed: Thus warmer spring temperatures could potentially increase total annual ET through lengthening of the early growing season**

I suggest you either changing the term “soil AWC” or more clearly define it to include other potential water sources. There is a growing body of literature that suggests that soil storage alone is often not sufficient to represent available water in mountain catchments. There this is rock water, groundwater, mobile vs. immobile water, etc. You have an opportunity to broaden the discussion and awareness among the land surface/hydroclimate modeling community of these distinctions with this work.

**Reply: We agree with the reviewer that plants often access water beneath what is typically defined as soil. In RHESSys plant available water storage is not restricted to “soil” but can include sapprolite, and rock water – and water from groundwater flow. We agree that the terminology “soil AWC” is indeed misleading. We have revised this terminology throughout the text and included the following text to emphasize that storage occurs not only in the soil:**

Previous studies have shown that plants access to stored water is a substantial contributor to summer evapotranspiration in semi-arid regions (Bales et al., 2011). Plant accessible storage includes both water stored in soil and in sapprolite and bedrock layers that can be accessed by plant roots (McNamara et al., 2011).

Addressing the above issues should not require large amount of work, but should help focus the paper on important take home messages by addressing and removing distracting aspects of the current presentation likely to distract a critical reader.

#### References

- Bales, R., Hopmans, J., O’Green, A., Meadows, M., Hartsough, P., Kirchner, P., ... Beaudette, D. (2011). Soil Moisture Response to Snowmelt and Rainfall in a Sierra Nevada Mixed-Conifer Forest. *Vadose Zone Journal*, 10(3), 786–799. doi:10.2136/vzj2011.0001
- Cayan, D. R., Dettinger, M. D., Kammerdiener, S. a., Caprio, J. M., & Peterson, D. H. (2001). Changes in the Onset of Spring in the Western United States. *Bulletin of the American Meteorological Society*, 82(3), 399–415. doi:10.1175/1520-0477(2001)082<0399:CITOOS>2.3.CO;2
- Isaac, V., & van Wijngaarden, W. a. (2012). Surface Water Vapor Pressure and Temperature Trends in North America during 1948–2010. *Journal of Climate*, 25(10), 3599–3609. doi:10.1175/JCLI-D-11-00003.1
- McNamara, J. P., Tetzlaff, D., Bishop, K., Soulsby, C., Seyfried, M., Peters, N. E., ... Hooper, R. (2011). Storage as a metric of catchment comparison. *Hydrological Processes*, 25, 3364–3371. doi:10.1002/hyp.8113

Tague, C., & Peng, H. (2013). The sensitivity of forest water use to the timing of precipitation and snowmelt recharge in the California Sierra: Implications for a warming climate. *Journal of Geophysical Research: Biogeosciences*, 118, 1–13. doi:10.1002/jgrg.20073

## **RESPONSE TO REFEREE #2**

Authors have performed an interesting study to assess the role of soil storage on climate-evapotranspiration (ET) interactions in three mountainous catchments using a distributed ecohydrologic model. In particular, the role of soil storage is considered by incorporating uncertainty of soil storage parameters in deriving precipitation, recharge and temperature relationships with ET. The manuscript is very well written and discussion of the results is very clear. However, the readers can benefit from a more focused conclusion summarizing main take home messages of the paper and its broader impact.

**Reply: We thank the reviewer for the supportive comments and address the detailed comments below. Both referees suggest a more focused conclusion. We have edited the conclusion with this suggestion in mind -- some points have been moved to the discussion and other text has been removed. The edited conclusion now reads as follows:**

[Edited conclusion removed for brevity. It is pasted into response to Reviewer #1 above.]

## **REVIEWER #2 COMMENTS CONTINUE BELOW**

### **Detailed comments:**

1) Authors have used a spatially distributed model to perform simulations across three catchments but the final results are aggregated at the catchment scale. It will be very interesting to see how these climate sensitivities change across the catchment? Are they observing differences between uplands and lowland areas?

**REPLY: We agree that the spatial patterns of these climate sensitivities would be interesting to observe across the catchment, however a thorough analysis is beyond the scope of this paper and we will explore more spatial patterns in future work. It would afford less room in the manuscript length to address how these climate sensitivities interact with subsurface properties, which we believe to be the novel contribution of our work.**

2) Does the sensitivity of ET change for different land cover types in a given catchment?

**REPLY: We expect that ET estimates would vary with different land cover types in each catchment. Two of our catchments, CA-SIER and OR-CAS, are uniformly covered in conifers. CO-ROC, which is significantly larger than the other two catchments, is comprised of other land types including meadows and rock. We**

expect that these land cover types are responding to climate drivers differently than the conifers. We have added text to acknowledge this important point:

We also note that CO-ROC is considerably larger than our other two study sites and, as such, includes significant fractions of other land cover including rock and meadow. We expect the different vegetation types to confound the response of ET to climate.

3) What about sensitivities of plant transpiration and NPP to AWC and precipitation?

**REPLY:** We explored the response of transpiration and NPP to the climate drivers presented in our study and found similar patterns in response across watersheds. We chose to focus on ET because we were able to validate our model estimates of annual values to field based observations in each catchment, which we believe strengthen our paper's results and discussion. We were unable to find similar transpiration observations at an appropriate scale/temporal resolution for model validation. We used annual NPP estimates to validate our carbon cycle, but chose to focus our presentation on the sensitivity of our hydrologic metric (ET) because it is more directly related to availability of soil moisture (AWC). Adding discussions of NPP/plant transpiration would also substantially lengthen an already long paper but we will consider this in future work.

4) Can authors specify which of the soil parameters generate most of the scatter in their results like in Figure 5 or 6? In other words, what is the most sensitive parameter? Is the most sensitive parameter different among the catchments?

**REPLY:**

We examined the influence of individual soil parameters to the sensitivity of ET estimates (not shown) and found that the sensitivity often varied with combinations for parameters rather than a single parameter value. For all catchments, streamflow estimates were most sensitivity to the 'm' parameter that controls the decay of conductivity with depth and defines an effective soil depth.

5) Can authors specify which metric they used for annual NPP during calibration (page 7899)?

**REPLY:** We used estimates of annual NPP that we found in peer-reviewed literature to define a minimum and maximum range of NPP values then selected parameters that fell within this range. That range of values is provided in Table 2. For calibration we selected parameters that fell within this NPP range and also provided reasonable estimates of streamflow based on the NSE and the daily bias.

6) Since R75 is not the actual recharge, I suggest authors rephrase it to timing of potential recharge.

**REPLY:** We agree that timing of potential recharge is more appropriate and have rephrased as follows:

To assess the impact of timing of potential recharge (as influenced either by year to year variation in precipitation timing, snowmelt or rain-snow partitioning) we calculate  $R_{75}$ , the day of water year by which 75% of the total potential annual recharge has occurred.

7) Can authors briefly describe patch elements in RHESSys (page 7898)?

**REPLY: Patch units are not necessarily grid shaped, but instead are delineated based on landscape characteristics including elevation, land cover classification, and aspect. Average patch sizes range from 90 to 8100m<sup>2</sup> with average patch size of 3600m<sup>2</sup>. Soil, vegetation and climate processes are calculated at the scale of the patch.**

8) A brief description of snow module will be helpful. How the results are impacted by the snow parameters?

**REPLY:**

**We have added the following text to the methods section that describes the RHESSys model:**

RHESSys partitions rain to snow at a daily timestep based on each patch's air temperature. Snowmelt is estimated using a combination of an energy budget approach for radiation-driven melt and a temperature index-based approach for latent heat-drive melt processes.

**We agree with the reviewer that results are potentially sensitive to snow parameters that control the rate of accumulation and melt. However we assume that in order to achieve reasonable rates of model performance relative to daily streamflow observations, the snow parameters used are reasonable and provide a basis for assessing the sensitivity to subsurface characteristics, which is the central focus of this paper.**



1 | [Subsurface storage capacity](#) influences climate-evapotranspiration interactions in three  
2 | western United States catchments

3

4 | Garcia, Elizabeth S., Department of Atmospheric Sciences, University of Washington,  
5 | Seattle, WA, USA.

6

7 | Tague, Christina L., Bren School of Environmental Science and Management, University  
8 | of California, Santa Barbara, California, USA.

9

10 | Corresponding author: E. S. Garcia, Department of Atmospheric Sciences, University of  
11 | Washington, Seattle, WA 98195 (esgarcia@uw.edu)

12

13

Naomi Tague 11/1/2015 6:09 PM

Deleted: Soil storage

15 ABSTRACT

16

17 In the winter-wet, summer-dry forests of the western United States, total annual  
18 evapotranspiration (ET) varies with precipitation and temperature. Geologically mediated  
19 drainage and storage properties, however, may strongly influence these relationships  
20 between climate and ET. We use a physically based process model to evaluate how plant  
21 accessible water storage capacity (AWC) and rates of drainage influence model estimates  
22 of ET-climate relationships for three snow-dominated, mountainous catchments with  
23 differing precipitation regimes. Model estimates show that total annual precipitation is a  
24 primary control on inter-annual variation in ET across all catchments and that the timing  
25 of recharge is a second order control. Low AWC, however, increases the sensitivity of  
26 annual ET to these climate drivers by three to five times in our two study basins with  
27 drier summers. ET – climate relationships in our Colorado basin receiving summer  
28 precipitation are more stable across subsurface drainage and storage characteristics.  
29 Climate driver-ET relationships are most sensitive to subsurface storage (AWC) and  
30 drainage parameters related to lateral redistribution in the relatively dry Sierra site that  
31 receives little summer precipitation. Our results demonstrate that uncertainty in  
32 geophysically mediated storage and drainage properties can strongly influence model  
33 estimates of watershed scale ET responses to climate variation and climate change. This  
34 sensitivity to uncertainty in geophysical properties is particularly true for sites receiving  
35 little summer precipitation. A parallel interpretation of this parameter sensitivity is that  
36 spatial variation in storage and drainage properties are likely to lead to substantial within-  
37 watershed plot scale differences in forest water use and drought stress.

38

39 1. INTRODUCTION

40

41 In high-elevation forested ecosystems in the western U.S., the majority of precipitation  
42 falls during the winter there is often a disconnect between seasonal water availability and  
43 growing season water demand. Consequently forests in these regions are frequently water  
44 limited, even when annual precipitation totals are high (Boisvenue and Running, 2006;  
45 Hanson and Weltzin, 2000). This disconnect between water inputs and energy demands

46 also highlights the importance of storage of winter recharge by both snowpack and by  
47 soils. The importance of snowpack storage in these systems for hydrologic fluxes has  
48 received significant attention, particularly given their vulnerability to climate warming.  
49 Warmer temperatures are already shifting seasonal water availability in the western U.S.  
50 through reductions in snowpack accumulation (Knowles et al., 2006) and earlier  
51 occurrence of peak snowpack (Mote et al., 2005) and shifts in streamflow timing (Stewart  
52 et al., 2005). Recently, field and modeling studies have shown that the years with greater  
53 snowpack accumulation can be a strong predictor of vegetation water use and  
54 productivity for sites in the California Sierra (Tague and Peng, 2013; Trujillo et al., 2012).  
55

56 Less attention, however, has been paid to the role of subsurface storage and drainage that  
57 can influence whether or not winter precipitation or snowmelt is available for plant water  
58 use during the summer months. Previous studies have shown that plant access to stored  
59 water is a substantial contributor to summer evapotranspiration in semi-arid regions  
60 (Bales et al., 2011). Plant accessible storage includes both water stored in soil and in  
61 sapprolite and bedrock layers that can be accessed by plant roots (McNamara et al., 2011).

62 Like snowpack, the storage of water in the subsurface has the potential to act as a water  
63 reservoir, storing winter precipitation for use later in the growing season (Geroy et al.,  
64 2011). The amount of water that can be stored varies substantially in space with  
65 topography, geologic properties, and antecedent moisture conditions (Famiglietti et al.,  
66 2008; McNamara et al., 2005). If the rate of snowmelt allows for subsurface moisture,  
67 stores to be replenished later into the growing season, more of the winter precipitation is  
68 made available for plant water use. If, storage capacity is too shallow to capture a  
69 significant amount of runoff or if the rate of rain or snowmelt inputs exceeds the rate of  
70 infiltration, then subsurface storage will not be physically able to extend water  
71 availability. While field studies in the Western US have shown that shallow soils can  
72 limit how much snowmelt is available for ecological use during the summer (Kampf et al.,  
73 2014; Smith et al., 2011), these studies cannot fully characterize the relative impact of  
74 subsurface storage on ET given inter-annual and cross-site variation in climate drivers.

75

Elizabeth Garcia 11/20/2015 7:28 AM

Comment [1]: Reference added

Naomi Tague 11/1/2015 6:14 PM

Deleted: by a soil

Naomi Tague 11/1/2015 6:14 PM

Deleted: soil

Naomi Tague 11/1/2015 6:24 PM

Deleted: soi

Elizabeth Garcia 11/20/2015 7:28 AM

Deleted: l

Naomi Tague 11/1/2015 6:24 PM

Deleted: oils

Naomi Tague 11/1/2015 6:24 PM

Deleted: are

Naomi Tague 11/1/2015 6:24 PM

Deleted: soil

Naomi Tague 11/1/2015 6:24 PM

Deleted: soil

84 | In this paper, we focus on the potential for plant accessible subsurface water storage to  
85 | mediate the sensitivity of ET to inter-annual variation in climate drivers, precipitation and  
86 | temperature. Understanding how ET varies with climate drivers is important, both from  
87 | the perspective of how ET influences downstream water supply and water availability for  
88 | forests and other vegetation (Grant et al., 2013). Western U.S. forests show substantial  
89 | vulnerability to drought, with declines in productivity and increases in mortality and  
90 | disturbance in drought years (Allen et al., 2010; Hicke et al., 2012; Williams et al., 2013).  
91 | Understanding these ecosystems' responses to primary climate drivers is of particular  
92 | concern given recent warming trends (Sterl et al., 2008) and multi-year droughts (Cook et  
93 | al., 2004; Dai et al., 2004) and that these changes in water and energy demands are  
94 | expected to intensify (Ashfaq et al., 2013). Increased temperatures also effect plant  
95 | phenology, leading to earlier spring onset of plant water use and productivity (Cayan et  
96 | al., 2001) and thus can influence water requirements and water use. However, increases  
97 | in early season water use, combined with higher atmospheric moisture demand, may lead  
98 | to increased soil water deficit later in the season.

99 |  
100 | Forest evapotranspiration is also a substantial component of the water budget (Post and  
101 | Jones, 2001) and thus any change in forest water use will potentially have significant  
102 | impacts on downstream water use. *Goulden et al.* [2012], for example, use flux tower and  
103 | remote sensing data to argue that warming may result in an increase of up to 60% in  
104 | vegetation water use at high elevations in the Upper Kings River watershed in  
105 | California's Southern Sierra watershed. We note however that these projected increases  
106 | depend on how subsurface storage capacity interacts with snowpack at high elevations.

107 |  
108 | This manuscript's primary research objective is to quantify the interaction between  
109 | subsurface storage characteristics and key climate-related metrics that influence forest  
110 | water availability and use in snow-dominated environments receiving a range of summer  
111 | precipitation. Heterogeneity in subsurface properties in soil, saprolite and bedrock layers  
112 | make the characterization of subsurface storage difficult at the watershed scale. Here we  
113 | use a spatially distributed process-based model, the Regional Hydro-Ecologic Simulation  
114 | System (RHESSys), to quantify how uncertainty or spatial variation in subsurface storage,

Naomi Tague 11/1/2015 6:25 PM

Deleted: soil

Naomi Tague 11/1/2015 6:25 PM

Deleted: o

Naomi Tague 11/1/2015 6:25 PM

Deleted: il

Naomi Tague 11/1/2015 6:25 PM

Deleted: oil

Naomi Tague 11/1/2015 6:26 PM

Deleted: soil

Elizabeth Garcia 10/30/2015 10:26 AM

Deleted: Mediterranean

Naomi Tague 11/1/2015 6:26 PM

Deleted: oil

Naomi Tague 11/1/2015 6:26 PM

Deleted: s soil

Naomi Tague 11/1/2015 6:27 PM

Deleted: oil

124 properties might be expected to influence watershed response to these climate-related  
125 drivers. We apply RHESSys in three case study watersheds of differing precipitation  
126 regimes to investigate how climate and [subsurface storage](#) combine to control inter-  
127 annual variation in ET.

128

## 129 2. METHODS

130

131 We apply our model at a daily time step to three watersheds located in the western  
132 Oregon Cascades (OR-CAS), central Colorado Rocky Mountains (CO-ROC) and central  
133 California Sierras (CA-SIER). All three watersheds receive a substantial fraction of  
134 precipitation as snowfall, but vary in their precipitation and temperature regimes and  
135 amount of precipitation that falls as snow (Figure 1). We compare a humid, seasonally  
136 dry watershed (OR-CAS) to two catchments that receive half as much precipitation  
137 annually. The more water-limited catchments differ in that CO-ROC receives a  
138 significant amount of its precipitation budget during the summer growing season. We use  
139 these case studies to estimate ET sensitivity to [storage and drainage properties](#) for  
140 several different precipitation and temperature regimes common in western U.S.

141 mountain watersheds. For each watershed, we quantify how subsurface storage and  
142 drainage properties interact with a combination of inter-annual variation in precipitation  
143 timing and magnitude, and shifts in snowpack storage. We first establish how inter-  
144 annual variation in three primary climate-related metrics (precipitation, average spring  
145 temperature, and timing of soil moisture recharge) influence annual ET with average  
146 [subsurface storage](#) properties. We then explore how these relationships change across  
147 physically plausible [storage values](#).

148

### 149 2.1 RHESSys MODEL DESCRIPTION

150

151 We use a physically based model (RHESSys v.5.15) to calculate vertical water, energy,  
152 and carbon fluxes in our three watersheds (Tague and Band, 2004). RHESSys is a  
153 spatially explicit model that partitions the landscape into units representative of the  
154 different hydro-ecological processes modeled (Band et al., 2000). RHESSys has been

Naomi Tague 11/1/2015 6:27 PM

Deleted: soil

Naomi Tague 11/1/2015 6:27 PM

Deleted: soil

Naomi Tague 11/1/2015 6:27 PM

Deleted: soil

Naomi Tague 11/1/2015 6:28 PM

Deleted: soil

Naomi Tague 11/1/2015 6:28 PM

Deleted: properties

160 used to address diverse eco-hydrologic questions across many watersheds (Baron et al.,  
161 2000; Shields and Tague, 2012; Tague and Peng, 2013). Key model processes are  
162 described below and a full account is provided in *Tague and Band* [2004].

163

164 RHESSys requires data describing spatial landscape characteristics and climate forcing; a  
165 digital elevation model (DEM), geologic, and vegetation maps are used to represent the  
166 topographic, geologic, carbon and nitrogen characteristics within a watershed. RHESSys  
167 accounts for variability of climate processes within the catchment using algorithms  
168 developed for extrapolation of climate processes from point station measurements over  
169 spatially variable terrain (Running and Nemani, 1987). Hydrologic processes modeled in  
170 RHESSys include interception, evapotranspiration, infiltration, vertical and lateral  
171 subsurface drainage, and snow accumulation and melt. The Penman-Monteith formula  
172 (Monteith, 1965) is used to calculate evaporation of canopy interception, snow

173 sublimation, evaporation from subsurface, and litter stores, and transpiration by leaves. A  
174 model of stomatal conductance allows transpiration to vary with soil water availability,  
175 vapor pressure deficit, atmospheric CO<sub>2</sub> concentration, and radiation and temperature  
176 (Jarvis, 1976). A radiation transfer scheme that accounts for canopy overstory and  
177 understory, as well as sunlit and shaded leaves, controls energy available for transpiration.  
178 RHESSys accounts for changes in vapor pressure deficit for fractions of days that rain  
179 occurs (wet versus dry periods). Plant canopy interception and ET are also a function of  
180 leaf area index (LAI) and gappiness of the canopy such that as LAI increases and gap size  
181 decreases, plant interception capacity and transpiration potential increases. RHESSys

182 partitions rain to snow at a daily timestep based on each patch's air temperature.  
183 Snowmelt is estimated using a combination of an energy budget approach for  
184 radiation-driven melt and a temperature index-based approach for latent heat-drive  
185 melt processes. Subsurface water availability varies as a function of infiltration and  
186 water loss through transpiration, evaporation and drainage. RHESSys also routes water  
187 laterally and thus patches can receive additional moisture inputs as either re-infiltration of  
188 surface flow or through shallow subsurface flow from upslope contributing areas. Lateral  
189 subsurface drainage routes subsurface and surface water between spatial units and it is a

Naomi Tague 11/1/2015 6:28 PM

Deleted: soil

Naomi Tague 11/1/2015 6:28 PM

Deleted: soil

Naomi Tague 11/1/2015 6:29 PM

Deleted: oil

193 function of topography and soil and saprolite drainage characteristics. Deep groundwater  
194 stores are drained to the stream using a simple linear reservoir representation.

195

196 Carbon and nitrogen cycling in RHESSys was modified from BIOME-BGC (Thornton,  
197 1998) to account for dynamic rooting depth, sunlit and shaded leaves, multiple canopy  
198 layers, variable carbon allocation strategies, and drought stress mortality. The Farquhar  
199 equation is used to calculate gross primary productivity (GPP) (Farquhar et al., 1980).  
200 Plant respiration costs include both growth and maintenance respiration and are  
201 influenced by temperature following *Ryan* [1991]. Net primary productivity (NPP) is  
202 calculated by subtracting total respiration costs from GPP.

203

204 In our three study sites, RHESSys is driven with daily records of precipitation and  
205 maximum and minimum temperature. Each basin is calibrated for seven parameters that  
206 characterize subsurface storage and drainage properties. Drainage rates are controlled by  
207 saturated hydraulic conductivity ( $K$ ) and its decay with depth ( $m$ ). Air-entry pressure  
208 ( $\varphi_{ae}$ ), pore size index ( $b$ ), and rooting depth ( $Z_r$ ) control subsurface water holding  
209 capacity (Brooks and Corey, 1964). In all basins, we assume that geologic properties  
210 allow for deeper groundwater stores that are inaccessible to vegetation (Table 2).

211 Vegetation however can access more shallow groundwater flow. These deep groundwater  
212 stores are controlled by two parameters representing the percentage of water that passes  
213 to the store ( $g_{w1}$ ) and the rate of its release to streamflow ( $g_{w2}$ ). Calibration is conducted  
214 with a Monte-Carlo based approach, the generalized likelihood uncertainty estimation  
215 (GLUE) method (Beven and Binley, 1992). Parameter sets (1000 total) are generated by  
216 random sampling from uniform distributions of literature-constrained estimates for the  
217 individual parameters; all calibration parameter sets are physically viable representations  
218 of soils within each basin. In other words, though a single parameter set may not meet  
219 streamflow and annual NPP calibration metrics, that particular subsurface storage  
220 capacity may still exist within the basin.

221

222 Model validation and drainage/storage parameter calibration were performed using two  
223 measures: daily streamflow statistics and annual measures of NPP. Streamflow statistics

Elizabeth Garcia 10/30/2015 6:44 PM  
Formatted: Font:(Default) Times, Bold

Naomi Tague 11/1/2015 6:29 PM  
Deleted: soil

Naomi Tague 11/1/2015 6:29 PM  
Deleted: Soil d

Naomi Tague 11/1/2015 6:29 PM  
Deleted: Soil

Naomi Tague 11/1/2015 6:29 PM  
Deleted: a

Naomi Tague 11/1/2015 6:30 PM  
Deleted: s

Naomi Tague 11/1/2015 6:29 PM  
Deleted: oil

Naomi Tague 11/1/2015 6:31 PM  
Deleted: soil type

Naomi Tague 11/1/2015 6:31 PM  
Deleted: soil

232 | were set such that good parameters resulted in daily flow magnitude errors less than 15%,  
233 | Nash-Sutcliffe efficiencies (NSE, a measure of hydrograph shape) greater than 0.65, and  
234 | logged NSE values greater than 0.7 (a test of peak and low flows) (Nash and Sutcliffe,  
235 | 1970). We select all parameter sets from these acceptable values; the total number of  
236 | parameters equals 87, 246, and 47 for CA-SIER, CO-ROC, and OR-CAS, respectively.  
237 | Daily hydrologic fluxes are calculated over 15 years for each soil parameter set in order  
238 | to account for variability due to parameters in establishing relationships with our climate  
239 | related indices, the results of which are presented in Figs. 2-4. We verify our annual ET  
240 | estimates against limited field estimates published in literature for subwatersheds of CO-  
241 | ROC and OR-CAS (Baron and Denning, 1992; Webb et al., 1978). The average of our  
242 | model estimated annual ET matches these limited field-based measurements and also fall  
243 | within the bounds of annual ET estimated through water balance by subtracting annual  
244 | streamflow from our records of annual precipitation. We assess the performance of the  
245 | carbon-cycling model by comparing with published forest field measurements of annual  
246 | NPP (values reported in Table 2). In our fully coupled eco-hydrologic model, accurate  
247 | estimates of NPP also suggest that ET estimates are reasonable. Finally we note that  
248 | RHESSys estimates of ET and NPP have been evaluated in a number of previous studies  
249 | by comparison with flux tower and tree ring data and these studies confirm that RHESSys  
250 | provides reasonable estimates of ET and its sensitivity to climate drivers (Vicente-  
251 | Serrano et al., 2015; Zierl et al., 2007) . We quantify the sensitivity of ET-climate  
252 | relationships to geologic properties by varying subsurface storage parameters (Figs. 5-6).

Naomi Tague 11/1/2015 6:31 PM

Deleted: soil

Naomi Tague 11/1/2015 6:32 PM

Deleted: soil

Naomi Tague 11/1/2015 6:32 PM

Deleted: soil

Naomi Tague 11/1/2015 6:32 PM

Deleted: soil

Naomi Tague 11/1/2015 6:32 PM

Deleted: soil

## 254 2.2 STUDY SITES

255  
256 | These analyses are conducted in three western U.S. mountain catchments: Big Thompson  
257 | in Colorado's Rocky Mountains (CO-ROC), Lookout Creek in Oregon's Western  
258 | Cascades (OR-CAS), and Sagehen Creek Experimental Forest in California's Northern  
259 | Sierra Nevada (CA-SIER). Basin characteristics pertinent to modeling annual ET are  
260 | listed in Table 2 and we highlight important similarities and differences here. All sites are  
261 | located on steep, mountainous slopes and are dominated by forest cover. All basins have  
262 | climates typical of the western U.S., on average receiving 54% - 81% of their annual



268 precipitation during the winter, 29% - 64% of the annual  $P$  falls as snow, and they do not  
269 meet potential evaporative demand during the growing season (Fig. 1, Table 2). On  
270 average, OR-CAS is a much wetter basin and receives more than twice as much annual  
271 precipitation than CO-ROC and CA-SIER. Despite OR-CAS receiving more precipitation,  
272 a much lower fraction of that winter precipitation is received as snow. On average OR-  
273 CAS's peak streamflow occurs in December, four to five months earlier than CO-ROC  
274 and CA-SIER (Fig. 1). The drier watersheds, CO-ROC and CA-SIER, receive more than  
275 half of their annual precipitation as snow (Table 2). CO-ROC also experiences a summer  
276 monsoonal season and on average receives 46% its annual precipitation from April –  
277 September. Landscape carbon (C) and nitrogen (N) stores in general vary with total  
278 annual  $P$  across basins. For example, OR-CAS receives the most precipitation and also  
279 supports stands of large, old-growth forests; its LAI is more than twice that of either CO-  
280 ROC or CA-SIER. As presented in the model description (Sect. 2.1), we use a stable,  
281 climatic optimum for vegetation biomass for all analyses in this paper. *Garcia et al.*  
282 [2013] and *Tague and Peng* [2013] provide detailed descriptions of the [geology](#), and  
283 climate data, model vegetation, and [organic](#) soil carbon store spin-up and calibration used  
284 for model implementations of OR-CAS and CA-SIER, respectively. We note that all  
285 precipitation and temperature data were derived from daily measurements made at  
286 climate stations located within the basins and extrapolated across the terrain using MT-  
287 CLM algorithms (Running and Nemani, 1987) and 30-m resolution DEMs. Though  
288 RHESSys has previously been used in CO-ROC (Baron et al., 2000), we have made  
289 significant updates in RHESSys since that time, so we re-implemented the model as  
290 described in the next section.

291

### 292 2.2.1 RHESSys MODEL DEVELOPMENT FOR CO-ROC

293

294 In CO-ROC, landscape topographic characteristics including elevation, slope and aspect  
295 were derived from a digital elevation model (DEM) downloaded from the U.S. Geologic  
296 Survey (USGS) National Elevation Data set at 1/3 arc second resolution  
297 (<http://datagateway.nrcs.usda.gov/>). A stream network was then derived to accumulate  
298 surface and subsurface flow at USGS gage #06733000. Sub-catchments were delineated

Naomi Tague 11/1/2015 6:33 PM

Deleted: soil

300 using GRASS GIS's watershed basin analysis program, *r.watershed*. Terrestrial data was  
301 aggregated such that the average size of the patch units, the smallest spatial units for  
302 calculation of vertical model processes, was 3600 m<sup>2</sup>. Soil classification data was  
303 downloaded from the Soil Survey Geographic database (SSURGO) and aggregated to  
304 four primary soil types: gravelly loam, sandy loam, loamy sandy, and rock  
305 (<http://datagateway.nrcs.usda.gov/>). Parameter values associated with these soil types are  
306 based on literature values (Dingman, 1994; Flock, 1978) and adjusted using model  
307 calibration, as described above. We note that these initial values are approximate and  
308 calibration permits storage values that reflect plant access to water stored in both organic  
309 soil layers and in sapprolite and rock. Vegetation land cover from the National Land  
310 Cover Database (NLCD) was aggregated to four primary vegetation types: subalpine  
311 conifer, aspen, shrubland, and meadow (Homer et al., 2007). Because a shift in  
312 precipitation patterns occurs at approximately 2700 meters, we use daily records of  
313 precipitation,  $T_{\max}$ , and  $T_{\min}$  from two points within the watershed. RHESys then  
314 interpolates data from these points based on MTN-CLM (Running and Nemani, 1987) to  
315 provide spatial estimates of temperature, precipitation and other meteorologic drivers for  
316 each patch. Climate data from 1980-2008 was downloaded from the DAYMET system  
317 for two locations – one at elevation 2460 m (latitude 40.35389, longitude -105.58361)  
318 and the second at 3448 m (latitude 40.33769, longitude -105.70315) (Thornton et al.,  
319 2012).

320

321 Plant C and N stores were initialized by converting remote-sensing derived LAI to leaf,  
322 stem and woody carbon and nitrogen values using allometric equations appropriate to the  
323 vegetation type (<http://daac.ornl.gov/MODIS/>; MOD15A2 Collection 5). In order to  
324 stabilize organic soil C and N stores relative to the LAI-derived plant C and N, we run the  
325 model repeatedly over the basin's climate record until the change in stores stabilizes  
326 (Thornton and Rosenbloom, 2005). After stabilizing soil biogeochemical processes, we  
327 remove vegetation C and N stores and then dynamically 'regrow' them using daily  
328 allocation equations (Landsberg and Waring, 1997) for 160 years in order to stabilize  
329 plant and soil C and N stores with model climate drivers. For all three basins, an optimum

330 maximum size for each vegetation type was determined using published, field-derived  
331 estimates of LAI and aboveground and total annual NPP.

332

333 2.3 FRAMEWORK for PRIMARY CONTROLS on *ET*

334

335 In these seasonally water-limited basins, we use total annual precipitation ( $P$ ) as a metric  
336 of gross climatic water input. Annual precipitation  $P$  is summed over a water year (Oct. 1  
337 to Sep. 30 of the following calendar year) and summer season  $P$  is summed over July,  
338 August, and September. For all climate metrics we use spatially averaged watershed  
339 values. To assess the impact of timing of soil moisture recharge (as influenced either by  
340 year to year variation in precipitation timing, snowmelt or rain-snow partitioning) we  
341 calculate  $R_{75}$ , the day of water year by which 75% of the total annual recharge has  
342 occurred. Recharge is defined as liquid water (e.g. rain throughfall or snowmelt) that  
343 reaches the soil surface. For this metric, we do not differentiate between water that, upon  
344 reaching the soil surface becomes runoff, and water that infiltrates into the soil. We treat  
345 this variable as a temporal marker of potential water availability that denotes the timing  
346 within the water year that either rain throughfall or snowmelt may potentially infiltrate  
347 the soil. To examine energy inputs, we identify a season when temperature most strongly  
348 influences estimates of annual ET modeled using historic climate. We performed linear  
349 regressions between model estimate of total annual ET and one and three-month averages  
350 of daily maximum ( $T_{\max}$ ), minimum ( $T_{\min}$ ) and average temperatures ( $T_{\text{avg}} = (T_{\max} +$   
351  $T_{\min})/2$ ) for all watersheds and for all months of the year. We test the correlation  
352 significance with a  $p$ -value and set a significance threshold at 0.05, i.e., a  $p$ -value greater  
353 than 0.05 is not significant. Our analysis found a three-month average of daily  $T_{\text{avg}}$  in  
354 April, May and June ( $T_{\text{AMJ}}$ ) to have the greatest explanatory power as a temperature  
355 variable for estimating inter-annual variation in annual ET under historic climate  
356 variability across our three study watersheds (results not shown). We note that the  $p$ -  
357 value for  $T_{\text{AMJ}}$  in CA-SIER was greater than 0.05 so it is not reported as a significant  
358 result. The growing season is assumed to extend from May 1 to September 30 in all  
359 watersheds. For all climate metrics we use spatially-averaged watershed values.

360

361 We examine the role of storage through AWC. As noted above, plants access water  
 362 organic soils as well as water stored in sapprolite and rock (Schwinning et al., 2010). We  
 363 consider an aggregate storage and do not distinguish between these layers. AWC  
 364 represents the water stored after gravity drainage (field capacity) that can be extracted by  
 365 plant root suction (wilting point), and is thus still viable for plant water use [Dingman,  
 366 1994, p. 236]. We calculate AWC as:

$$368 \text{ AWC} = (\theta_{fc} - \theta_{wp}) Z_r \quad (2)$$

370 Where  $\theta_{fc}$  represents the average field capacity per unit depth,  $\theta_{wp}$  the average  
 371 characteristic wilting point also per unit depth, and AWC is scaled by vegetation rooting  
 372 depth,  $Z_r$ , a model calibration parameter. The field capacity and wilting point are  
 373 calculated, respectively, as

$$375 \theta_{fc} = \phi (\varphi_{ae} / 0.033)^b \quad (3)$$

$$376 \theta_{wp} = \phi (\varphi_{ae} / \psi_v)^{1/b} \quad (4)$$

378 Where  $\phi$  is average subsurface porosity,  $\varphi_{ae}$  represents the air-entry pressure (in meters),  
 379  $b$  is a pore size distribution index that describes the moisture-characteristic curve, and  $\psi_v$   
 380 describes the pressure at which the plants' stomata close. Variables  $\varphi_{ae}$  and  $b$  are also  
 381 model calibration parameters.

383 Larger AWC indicates that more water can be held in the subsurface, and potentially  
 384 interacts with climate to extend plant water availability by capturing snowmelt, one of the  
 385 primary sources of water for forest ET. Our results present each watershed's average  
 386 AWC; watersheds are represented by one (OR-CAS), two (CA-SIER), and five (CO-  
 387 ROC) soil types and their characterizations are described in Table 2. All values of AWC  
 388 calculated in calibration represent physically feasible values for each watershed.

390 We use RHESSys to calculate total annual ET over the entire available climate record in  
 391 each basin (28-50 years; Table 2) and use linear regression to quantify how much of the

Naomi Tague 11/1/2015 6:36 PM  
**Comment [2]:**

Naomi Tague 11/1/2015 6:37 PM  
**Comment [3]:** Schwinning, S. The  
 ecohydrology of roots in rocks.  
*Ecohydrology* **3**, 238–245 (2010).

Naomi Tague 11/1/2015 6:36 PM  
**Deleted:** in the soil

Naomi Tague 11/1/2015 6:37 PM  
**Deleted:** soil's

Naomi Tague 11/1/2015 6:37 PM  
**Deleted:** soil's

Naomi Tague 11/1/2015 6:37 PM  
**Deleted:** soil

Naomi Tague 11/1/2015 6:38 PM  
**Deleted:** soil

Naomi Tague 11/1/2015 6:38 PM  
**Deleted:** soil

Naomi Tague 11/1/2015 6:38 PM  
**Deleted:** oil

Naomi Tague 11/1/2015 6:39 PM  
**Deleted:** We note that we use the term soil  
 in a hydrologic sense to denote subsurface  
 media that stores water. This can include both  
 organic and mineral soil as well as fractured  
 bedrock and sapprolite that would be accessible  
 by deeper plant roots

Naomi Tague 11/1/2015 6:40 PM  
**Deleted:** The range in soil parameter values  
 is bounded by literature values specific to each  
 site's soil properties (Dingman, 1994). This  
 means a

Naomi Tague 11/1/2015 6:39 PM  
**Deleted:** soils

410 inter-annual variation in ET is related to each of the three climate metrics— $P$ ,  $T_{AMJ}$ , and  
411  $R_{75}$ . We set a limit of less than 0.05 for p-values to determine significance. We then  
412 investigate how long-term mean ET and its relationship with these climate-related  
413 indicators are influenced by AWC.

414

415 To examine how subsurface storage capacity may influence long term average ET, we  
416 calculate average annual ET over a 15-year period (1985-2000) for a range of 1000 AWC  
417 values and linearly regress the long-term averaged ET values against AWC. We then  
418 characterize the interacting influences of AWC and each climate driver. For the 1000  
419 values of AWC, we calculate the slope of annual ET estimates to each climate predictor  
420 ( $P$ ,  $T_{AMJ}$ ,  $R_{75}$ ).

421

### 422 3. RESULTS

423

#### 424 3.1 ANNUAL $P$ vs. ET

425

426 In all watersheds higher  $P$  results in greater total annual ET (Fig. 2). This is a statistically  
427 significant relationship in all watersheds (CO-ROC and CA-SIER, correlations and p-  
428 values reported in Table 3) where the years of highest annual  $P$  are correlated with the  
429 years of greatest annual ET. Of the three basins, CO-ROC's annual ET shows the greatest  
430 sensitivity to  $P$ , having the steepest slope. Annual  $P$  is the strongest explanatory variable  
431 of annual ET in both CO-ROC ( $r^2 = 0.9$ ) and CA-SIER ( $r^2 = 0.75$ ) (Table 3). For CO-ROC,  
432 annual  $P$  has a greater influence (steeper slope) in the drier years when  $P$  is less than  
433 1000 mm (Fig. 2). OR-CAS has the least significant relationship between  $P$  and ET on an  
434 annual scale. OR-CAS is a relatively wet basin and on average receives more than twice  
435 the amount of winter (Jan-Mar) precipitation than CA-SIER or CO-ROC receives. High  
436 annual  $P$  in OR-CAS in most years likely diminishes the sensitivity of ET to the  
437 magnitude of  $P$ .

438

#### 439 3.2 TIMING OF RECHARGE vs ET

440

Naomi Tague 11/1/2015 6:40 PM

Deleted: soil

Naomi Tague 11/1/2015 6:40 PM

Deleted: soil

Naomi Tague 11/1/2015 6:40 PM

Deleted: soil

Naomi Tague 11/1/2015 6:40 PM

Deleted: soil

445 For all three catchments, later  $R_{75}$  has a significant positive correlation with ET (Fig. 3).  
446 In OR-CAS and CA-SIER,  $R_{75}$  occurs between February and May. There is more scatter  
447 in the predictive power of  $R_{75}$  for annual ET when  $R_{75}$  is earlier in the water year. The  
448 earliest  $R_{75}$  are in OR-CAS, where a greater fraction of winter precipitation falls as rain.  
449 CA-SIER and CO-ROC are more sensitive to the timing of recharge than OR-CAS.  
450 Summer monsoonal pulses in CO-ROC push  $R_{75}$  to later in the water year as compared to  
451 OR-CAS or CA-SIER. The explanatory power of  $R_{75}$  for ET is greatest in CA-SIER  
452 where greater accumulation of snowpack and warmer spring temperatures can interact to  
453 increase forest water use earlier in the growing season.

454

### 455 3.3 SPRING TEMPERATURE vs. ET

456

457 Warmer spring temperature ( $T_{AMJ}$ ) in all basins generally reduces annual ET (Fig. 4a) and  
458 is significantly correlated with lower ET in CO-ROC and OR-CAS. CA-SIER does not  
459 show a significant relationship between  $T_{AMJ}$  and ET. In CO-ROC and OR-CAS  
460 increasing  $T_{AMJ}$  leads to a reduction in water availability and a decline in later season ET.  
461 The relationship between spring air temperature and snowmelt timing is demonstrated by  
462 significant correlations between  $T_{AMJ}$  and  $R_{75}$  for CO-ROC (Fig. 4b). The colder  
463 temperatures and more persistent snowpack in the CO-ROC basin is more sensitive,  
464 relative to OR-CAS, in ET response to earlier snowmelt due to temperature increases.

465

### 466 3.4 AWC vs. ET

467

468 Increased AWC increases the long-term average ET in all basins. Figure 5 shows a  
469 nonlinear relationship between long-term mean ET and AWC suggesting that the effect  
470 of increasing storage diminishes for higher AWC values. Each basin reaches an  
471 approximate storage capacity above which a further increase in storage (AWC) is less  
472 important and climate (i.e.,  $P$  and energy) variables limit ET. Following *Muggeo* [2003],  
473 for each basin, we calculate that breakpoint value of AWC where ET is less sensitive to  
474 AWC. We find that the threshold value of AWC varies across basins and is substantially  
475 higher in CO-ROC (265 mm) as compared to CA-SIER (195 mm) and OR-CAS (190

Elizabeth Garcia 10/30/2015 10:33 AM

**Deleted:** , which is somewhat counter-intuitive. Among the predicted consequences of increased temperatures are an earlier start to the vegetation growing season (Cayan et al., 2001), and an increase in vapor pressure deficits and water demand (Isaac and van Wijnngaarden, 2012). Thus warmer spring temperatures could potentially increase total annual ET through lengthening of the early growing season. However, warmer spring temperatures are

Elizabeth Garcia 10/30/2015 5:14 PM

**Deleted:** because the effect of temperature is strongly dependent on the amount of snowpack the basin receives in a year (Tague and Peng, 2013), which is more variable than the amount of snowpack received in CO-ROC or OR-CAS. These results suggest that the dominant effect of warmer spring temperatures is earlier meltout of snowpack, which leads to more snowmelt lost as runoff and results in less net recharge. A mechanism we suggest for this loss of runoff is that soils are more likely to be saturated in spring months. Later into the growing season, increased ET demands will have depleted soil stores and throughfall/snowmelt will enter the soil matrix and be available for plant water use. So instead of higher  $T_{AMJ}$  leading to increased water demand and early growing season ET, i

Naomi Tague 11/1/2015 6:45 PM

**Deleted:** SOIL

506 | mm) (Fig. 5). Regression of AWC against annual ET show that a significant relationship  
507 | exists in OR-CAS and CO-ROC (Table 3).

508

509 | The effect of varying lateral redistribution or lateral drainage parameters can be seen in  
510 | the range of slopes for a given AWC (e.g., the scatter in the slope-AWC relationship).  
511 | All three watersheds show some sensitivity of climate-ET relationships to lateral  
512 | redistribution parameters for a given AWC. CA-SIER shows the greatest sensitivity,  
513 | followed by OR-CAS and CO-ROC. The greater sensitivity of CA-SIER to lateral  
514 | drainage parameters may reflect the strong contribution of snowmelt recharge in its drier  
515 | and winter precipitation dominated climate. The topography of CA-SIER is also  
516 | distinctive and includes many swale-like features that concentrate drainage from upslope  
517 | areas. We calculate the topographic wetness index (TWI) using a 30m resolution DEM  
518 | for each watershed (Moore et al., 1991) (Table 2). The TWI reflects the propensity of a  
519 | location to develop saturated conditions under the assumption that topography controls  
520 | water flow. Higher TWI values represent flatter, converging terrain and lower values  
521 | reflect steep topography. The mean TWI for CA-SIER is greater than, and significantly  
522 | different from (Welch's t-test) the mean TWI for CO-ROC and OR-CAS. Particularly for  
523 | CA-SIER, changing storage parameters associated with drainage rates can alter the  
524 | timing of flow into areas that concentrate flow and subsequently alter their ET rates.

525

526 | 3.5 SENSITIVITY OF ET to CLIMATE DRIVERS with AWC

527

528 | We analyze the sensitivity of ET relationships with climate drivers to subsurface storage  
529 | properties by plotting the slope of linear regressions between ET and  $P$ ,  $R_{75}$ , and  $T_{AMJ}$ ,  
530 | across all storage parameter sets in Fig. 6. We note that the slope of the relationships  
531 | between climate drivers and ET has been normalized by the watersheds' mean AWC in  
532 | these plots to facilitate cross-site comparison.

533

534 | 3.5.1 SENSITIVITY to  $P$  with AWC

535

Elizabeth Garcia 10/30/2015 7:18 AM

Deleted: logged values of

Naomi Tague 11/1/2015 6:45 PM

Deleted: soil

Naomi Tague 11/1/2015 6:45 PM

Deleted: SOIL

Naomi Tague 11/1/2015 6:45 PM

Deleted: oil

Naomi Tague 11/1/2015 6:45 PM

Deleted: soil

541 Of the climate drivers explored, ET relationships with annual precipitation  $P$  have the  
542 greatest robustness across subsurface storage parameter sets, as suggested by number of  
543 sets that show a statistically significant relationship between annual  $P$  and annual ET (Fig.  
544 6A). As expected, slopes are positive between  $P$  and ET across all basins. Only the drier  
545 basins CO-ROC and CA-SIER have p-values less than 0.001, highlighting the strength of  
546  $P$  as a climatic driver in these drier basins, as discussed above. The response in slope  
547 sensitivity across AWC is similar in OR-CAS and CA-SIER where ET's sensitivity to  $P$   
548 is highest at low AWC and decreases with increased AWC. OR-CAS has a much smaller  
549 range in sensitivities (slope varies from 0.2-0.6) compared to CA-SIER (slope varies  
550 from 0.0-0.8). Thus in CA-SIER for low values of AWC, year-to-year variation in  $P$   
551 becomes a greater control on year-to-year variation in ET. For both OR-CAS and CA-  
552 SIER, increasing AWC becomes less important at higher values of AWC. Higher scatter  
553 in slope of annual  $P$  versus ET relationship for CA-SIER also reflects the greater  
554 sensitivity of ET to subsurface parameters that influence lateral drainage as discussed  
555 above (Sect. 3.4).

556

557 The variation of ET response to  $P$  across AWC in CO-ROC is noteworthy for two  
558 reasons. First, CO-ROC has the highest slope values (0.6-0.8), which again reflects the  
559 consistency of annual  $P$  as a control on inter-annual variation in ET in this basin. Second,  
560 unlike OR-CAS and CA-SIER, increasing AWC does not substantially reduce that  
561 sensitivity (i.e., slope) to  $P$ . Though CO-ROC's sensitivity to  $P$  does not change with  
562 AWC, the scatter in slopes (0.6-0.8) suggests that lateral drainage has a strong effect on  
563 this climate-ET relationship. We note that CO-ROC has a seasonal precipitation regime  
564 where a significant fraction of its annual precipitation is received later in the growing  
565 season as summer monsoonal pulses. When precipitation occurs during the growing  
566 season, the water available for ET is less likely to be limited by storage capacity. Instead  
567 ET is limited by the amount or intensity of precipitation. Water that does recharge the  
568 system is used relatively quickly, making variation in storage (or AWC) less important as  
569 a control on how much  $P$  can be used in CO-ROC.

570

571 3.5.2 SENSITIVITY to  $R_{75}$  with AWC

Naomi Tague 11/1/2015 6:46 PM

Deleted: soil

Naomi Tague 11/1/2015 6:46 PM

Deleted: soil



574

575 After precipitation, the timing of recharge ( $R_{75}$ ) most significantly correlates with  
576 increased ET across all AWC and all basins (Fig. 6B). There are several similarities in  
577 the response of ET's sensitivity to  $R_{75}$  across AWC when compared to sensitivity to  $P$   
578 (Fig. 6A). For example, the dry basins CO-ROC and CA-SIER have the highest degree of  
579 sensitivity (significant slopes  $> 1.0$ ) as compared to OR-CAS (slopes  $< 1.0$ ) and CA-  
580 SIER has the greatest variability in its sensitivity to AWC with slopes ranging from 1.0-  
581 3.0 across variation in [storage](#) parameters. CO-ROC once again has the least variability in  
582 the ET versus  $R_{75}$  relationship, with consistently high (2.0-2.5) slopes unaffected by  
583 AWC.

Naomi Tague 11/1/2015 6:46 PM

Deleted: oil

584

### 585 3.5.3 SENSITIVITY to $T_{AMJ}$ with AWC

586

587 Finally,  $T_{AMJ}$  has the fewest [subsurface storage/drainage](#) parameter sets with significant  
588 correlation with ET. None of the linear regressions of ET on  $T_{AMJ}$  have statistical  
589 significance less than 0.001 (Fig. 6C). The slopes are always negative because earlier  
590 occurrence of snowmelt results in less ET. For all basins, the sensitivity of ET to  $T_{AMJ}$  is  
591 greatest at the lowest values of AWC, though CO-ROC once again demonstrates the least  
592 variability in slopes across the entire range of AWC (-0.2 – -0.3). At OR-CAS,  $T_{AMJ}$  is  
593 only significant for the lower AWC values. We suggest this is in part due to the small  
594 fraction of  $P$  that falls as snow. Because  $T_{AMJ}$ 's largest effect is through timing of  
595 snowmelt (Fig. 4), AWC interacts with  $T_{AMJ}$  to modulate the melt response. With  
596 relatively less snowmelt in OR-CAS, only the [systems](#) with the smallest capacities will  
597 have a significant negative interaction effect with AWC.

Naomi Tague 11/1/2015 6:46 PM

Deleted: soil

598

## 599 4. DISCUSSION

600

601 Our model estimates show differences in the response of ET to climate-related drivers  
602 across the three watersheds, primarily due to differences in their precipitation regimes.

603 Spatial heterogeneity in [soil and geology](#), both within and between watersheds

604 substantially alter these relationships. [Our model-based study provides a simplified](#)

Naomi Tague 11/1/2015 6:47 PM

Deleted: soils

Naomi Tague 11/1/2015 6:47 PM

Deleted: soils

Elizabeth Garcia 10/30/2015 10:49 AM

Moved (insertion) [1]

609 representation of these interactions, ignoring many additional complexities. In particular,  
610 we assume no adaptation of the ecosystem structure and composition that would  
611 influence productivity, evapotranspiration and their relationship with climate  
612 (Loudermilk et al., 2013). Future work will investigate these coupled carbon cycling-  
613 hydrology interactions. In this study we focus on the energy and moisture drivers of ET  
614 and how subsurface properties influence their interaction.

615  
616 The degree to which climate drivers affect ET varies with the magnitude and seasonality  
617 of basin precipitation. Total annual  $P$  is the first order control of ET in the two drier  
618 watersheds, CO-ROC and CA-SIER. In OR-CAS, most of the inter-annual variation in  
619 precipitation is reflected in inter-annual variation in runoff rather than ET. In most years,  
620 subsurface storage is filled by this annual precipitation during the winter and spring,  
621 asynchronously to late growing season demands (Fig. 1). Our results extend findings by  
622 previous studies demonstrating that vegetation productivity and water use relates to the  
623 fraction of regional precipitation available to plants (Brooks et al., 2011; Thompson et al.,  
624 2011). The fraction of water available to plants tends to decrease with larger rainfall  
625 (given saturated soil stores a greater proportion is lost) and with synchronicity between  
626 the timing of recharge and growing season water demands.

627  
628 Our analysis highlights the timing of water availability ( $R_{75}$ ) as a key predictor of total  
629 annual ET; annual ET increases when recharge occurs later in the water year, during the  
630 growing season and period of highest water demand. Previous research has shown how  
631 delayed soil moisture recharge (Tague and Peng, 2013) and snowpack dynamics (Tague  
632 and Heyn, 2009; Trujillo et al., 2012) are able to increase ET in the Sierra Nevada. In  
633 these mountain basins, the sensitivity of ET to timing of recharge is related to the fraction  
634 of precipitation received as snow. The climate metrics related to snowmelt,  $R_{75}$  and  $T_{AMJ}$ ,  
635 are important secondary controls of ET, especially in the colder, snow-dominated  
636 watersheds, CA-SIER and CO-ROC. We note that CA-SIER does not show a significant  
637 relationship between  $T_{AMJ}$  and ET because the effect of temperature is strongly dependent  
638 on the amount of snowpack the basin receives in a year (Tague and Peng, 2013), which is  
639 more variable than the amount of snowpack received in CO-ROC or OR-CAS. In OR-

Elizabeth Garcia 10/30/2015 10:49 AM

Deleted: -

Naomi Tague 11/1/2015 6:47 PM

Deleted: soil

Elizabeth Garcia 10/30/2015 5:25 PM

Moved (insertion) [4]

642 CAS and CO-ROC, spring temperature  $T_{AMJ}$  is more strongly related to ET through its  
 643 effect on snowmelt and correlates negatively with ET. These results suggest that the  
 644 dominant effect of warmer spring temperatures is earlier meltout of snowpack, which  
 645 leads to more snowmelt lost as runoff and results in less net recharge. This greater loss of  
 646 runoff occurs when storage capacity is exceeded. Later into the growing season,  
 647 increased ET demands will have depleted subsurface stores and throughfall/snowmelt  
 648 will enter the soil matrix and be available for plant water use. Previous work has shown  
 649 seasonal increases in spring ET with warmer spring temperatures (Hamlet et al., 2007)  
 650 which may be related to an earlier start to the vegetation growing season (Cayan et al.,  
 651 2001), and an increase in vapor pressure deficits and water demand (Isaac and van  
 652 Wijngaarden, 2012). Our work suggests that though early season ET may increase with  
 653 warming temperatures, warmer spring temperatures may in some cases decrease total  
 654 annual ET by melting the snowpack stores earlier in the water year and reducing soil  
 655 moisture recharge later in the spring when energy demand is high.

656

657 The range of sensitivities of ET to climate in this study is a direct function of climatic and  
 658 physical characteristics of the catchments presented in this study. For example, OR-CAS  
 659 receives twice as much precipitation and spans a much lower elevation range than either  
 660 CA-SIER or CO-ROC (Table 2). Because OR-CAS is considerably wetter, its sensitivity  
 661 of ET to magnitude of annual  $P$  is lessened considerably. OR-CAS' lower elevations, and  
 662 related mean winter temperatures, also result in smaller average snowpacks reducing the  
 663 strength of spring temperature as an explanatory variable for ET. Differences between  
 664 CA-SIER and CO-ROC largely reflect seasonal distribution of precipitation, and reflect  
 665 the importance of summer precipitation in CO-ROC. While climate is the dominant  
 666 factor, topographic differences are also important. As discussed above, topographically  
 667 driven flowpath convergence in CA-SIER tends to increase sensitivity of ET to  
 668 parameters that influence lateral drainage. This effect is less evident in the other two  
 669 watersheds.

670

671 Over a range of physically realistic storage characteristics, long-term averages of ET  
 672 increase with greater storage (AWC) in all basins. Our analysis found the greatest

- Naomi Tague 11/1/2015 6:56 PM  
**Deleted:** A mechanism we suggest for this
- Naomi Tague 11/1/2015 6:56 PM  
**Deleted:** is that
- Naomi Tague 11/1/2015 6:48 PM  
**Deleted:** soils
- Naomi Tague 11/1/2015 6:48 PM  
**Deleted:** are more likely to be saturated
- Naomi Tague 11/1/2015 6:56 PM  
**Deleted:** in spring months
- Naomi Tague 11/1/2015 6:48 PM  
**Deleted:** oil
- Naomi Tague 11/1/2015 6:57 PM  
**Deleted:** the lateness of
- Elizabeth Garcia 10/30/2015 5:25 PM  
**Moved up [4]:** In OR-CAS and CO-ROC, spring temperature  $T_{AMJ}$  is more strongly related to ET through its effect on snowmelt and correlates negatively with ET.
- Naomi Tague 11/1/2015 6:58 PM  
**Formatted:** Font:Not Bold
- Naomi Tague 11/1/2015 6:58 PM  
**Formatted:** Font:Not Bold
- Naomi Tague 11/1/2015 6:58 PM  
**Deleted:** The range of sensitivities of ET to climate in this study is a direct function of the physical characteristics of the catchments presented in this study. For example, OR-CAS receives twice as much precipitation and spans a much lower elevation range than either CA-SIER or CO-ROC (Table 2). Because OR-CAS is considerably wetter, its sensitivity of ET to magnitude of annual  $P$  is lessened considerably. Similarly, OR-CAS' lower elevations, and related mean winter temperatures, result in smaller average snowpacks reducing the strength of spring temperature as an explanatory variable for ET. We also note that CO-ROC is considerably larger than our other two study sites and, as such, includes ... [1]
- Naomi Tague 11/1/2015 6:58 PM  
**Formatted:** Font:Not Bold
- Elizabeth Garcia 10/30/2015 9:51 AM  
**Formatted:** Font:Not Bold
- Elizabeth Garcia 10/30/2015 9:56 AM  
**Formatted:** Font:Not Bold
- Elizabeth Garcia 10/30/2015 10:56 AM  
**Deleted:** Increasing AWC generally increases the magnitude of long-term a... [2]
- Naomi Tague 11/1/2015 6:49 PM  
**Deleted:** soil
- Naomi Tague 11/1/2015 6:49 PM  
**Deleted:** soil

735 sensitivity of long-term average annual ET to variation in AWC in OR-CAS (Table 3). In  
736 CO-ROC, ET ranges from 380-600 mm across annual  $P$  variation, and across all  
737 calibrated subsurface parameters long-term average ET ranges from 450-600 mm. This  
738 variation in CO-ROC's ET associated with subsurface storage characteristics is on the  
739 same order of magnitude as inter-annual variation in ET with  $P$ . Similarly, in CA-SIER,  
740 ET ranges from 400-800 mm across the  $P$  record and across all storage parameters, and  
741 ranges from 700-1000 mm long-term. There is a nonlinear relationship between ET and  
742 AWC in each basin. We suggest that below a threshold point in each basin (195 - 265mm  
743 of AWC), long-term average ET is more sensitive to AWC and above these threshold  
744 values the effect of climate on ET is greater than an increase in subsurface storage.

746 The sensitivity of ET to year-to-year variability of climate drivers is also influenced by  
747 AWC. The sensitivity of ET estimates to climate drivers varies by two to five magnitudes  
748 in CA-SIER and OR-CAS across the range of plausible storage parameters. These basins  
749 receive the smallest fraction of annual  $P$  in the summer and their annual ET estimates are  
750 most sensitive to  $P$ ,  $R_{75}$ , and  $T_{AMJ}$  at low water capacity (AWC). CO-ROC has a high  
751 sensitivity to climate drivers but this sensitivity does not change with AWC. We suggest  
752 that a strong summer  $P$  signal in CO-ROC explains the negligible change in ET's  
753 sensitivity to climate drivers across values of AWC, similar to other studies that show  
754 that summer  $P$  can offset the dependence of ET on soil replenishment or winter snowpack  
755 (Hamlet et al., 2007; Litaor et al., 2008). The relative importance of AWC to regional  
756 climate differences is apparent if we consider that a similar sensitivity to  $P$  and  $T_{AMJ}$  can  
757 be achieved for all basins by varying AWC. For example, ET at the smallest AWC values  
758 in OR-CAS are similarly sensitive (slope of 0.6) to inter-annual variation precipitation as  
759 stands in CO-ROC (Fig. 6A).

761 The two more water-limited basins demonstrate similarly high sensitivities of ET to  
762 climate drivers, but differ in the response of their sensitivity to climate across AWCs.  
763 Despite CO-ROC and CA-SIER showing similarly strong sensitivities to climate, their  
764 response across AWC differs considerably. CA-SIER's sensitivity to climate drivers is  
765 highly variable across all AWC but still demonstrates slightly higher sensitivity at lower

Naomi Tague 11/1/2015 6:49 PM

Deleted: soil

Naomi Tague 11/1/2015 6:59 PM

Deleted: soil

Naomi Tague 11/1/2015 6:49 PM

Deleted: soil

Naomi Tague 11/1/2015 6:49 PM

Deleted: oil

Naomi Tague 11/1/2015 6:49 PM

Deleted: soil

Naomi Tague 11/1/2015 6:50 PM

Deleted: oil

Naomi Tague 11/1/2015 6:50 PM

Deleted: soil

Naomi Tague 11/1/2015 6:50 PM

Deleted: content

Naomi Tague 11/1/2015 6:50 PM

Deleted: soil

Naomi Tague 11/1/2015 6:50 PM

Deleted: soil

Naomi Tague 11/1/2015 6:50 PM

Deleted: soil

777 AWC values. Its lack of summer precipitation, like OR-CAS, gives water storage a more  
778 significant role in mediating late summer water stress. With lower AWC values there is  
779 less potential for water storage and ET becomes more sensitive to climate drivers.

780

781 In addition to the sensitivity to AWC, our results show that lateral redistribution strongly  
782 influences the sensitivity of ET to climate drivers in the drier basins; in CA-SIER and  
783 CO-ROC there is considerable scatter in the slopes for  $P$  and  $R_{75}$  across a single AWC  
784 (e.g., for an AWC of 400 mm, the  $P:ET$  ranges from 0.6 to 0.8 and 0.2 to 0.7 for CO-  
785 ROC and CA-SIER, respectively in Fig. 6A). We note that this additional sensitivity of  
786 ET-climate relationships to drainage rates, even given similar AWC or storage conditions,  
787 emphasizes the role played by lateral connections. In other words, results suggest that for  
788 the two more water limited sites, the timing of upslope contributions to downslope areas  
789 can mediate the sensitivity of watershed scale vegetation water use.

790

791 Our results have general implications for model based estimates of ET in this region.  
792 Because there is substantial heterogeneity in subsurface storage characteristics within  
793 each basin (Dahlgren et al., 1997; Denning et al., 1991; McGuire et al., 2007) we might  
794 expect that the full range of AWCs can be observed when we look across individual  
795 forest stands within a basin. Thus, our estimates that show substantial changes in climate-  
796 ET relationships across subsurface parameters suggest that there may be substantial  
797 within-basin spatial heterogeneity in vegetation responses to climate variation and change.  
798 Even if model estimates are focused on basin aggregate responses such as streamflow,  
799 our results point to the importance of calibration data for defining subsurface storage and  
800 drainage properties. Estimates of subsurface parameters are often derived from readily  
801 available products such as STATSGO and SSURGO [Natural Resources Conservation  
802 Service] that provide relatively coarse scale and imperfect information about hydrologic  
803 properties. Consequently, hydrologic models are typically calibrated to obtain estimates  
804 of storage and drainage parameters (Beven, 2011). Our results suggest that in areas where  
805 streamflow data is not available for calibration, watershed scale estimates of ET  
806 responses to climate drivers may have substantial errors.

807

Naomi Tague 11/1/2015 6:50 PM

Deleted: soil

Elizabeth Garcia 10/30/2015 5:54 PM

Moved (insertion) [2]

Naomi Tague 11/1/2015 6:51 PM

Deleted: soil

Naomi Tague 11/1/2015 6:51 PM

Deleted:

Naomi Tague 11/1/2015 6:51 PM

Deleted: at soils

Naomi Tague 11/1/2015 6:51 PM

Deleted: oil

Naomi Tague 11/1/2015 7:06 PM

Deleted: -

814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844

## 5. CONCLUSIONS

We demonstrate how subsurface storage and drainage properties (AWC and parameters that control lateral redistribution) interact with climate-related drivers to influence ET in three western U.S. mountain watersheds with distinctive precipitation regimes. These watersheds reflect conditions found in many other western U.S. snow-dominated systems, where summer water availability is influenced by the magnitude of precipitation, timing of soil moisture recharge and spring temperature and its effect on snowmelt. We found that, for our three watersheds, estimates of longer-term average (15-year) watershed-scale ET vary across a range of physically realistic storage/drainage parameters. For all watersheds, the range in long term mean ET estimates across AWC estimates (e.g., mean ET at a high AWC versus mean ET at a low AWC) may be as large as inter-annual variation in ET, suggesting that the influence of AWC and drainage can be substantial.

Our results also point to the importance of lateral redistribution as a control on ET, particularly for CA-SIER. Only a few studies have emphasized the role of lateral redistribution in plot to watershed scale climate responses in the Western U.S. (Barnard et al., 2010; Tague and Peng, 2013). For the CA-SIER site, our model results suggest that there can also be interactions between AWC and hillslope to watershed scale redistribution as controls on ET. Lateral redistribution was less important for the CO-ROC, where summer precipitation was a more important contributor to annual ET values and the least important for the wetter OR-CAS site. Results emphasize that the role of subsurface properties, including both storage and drainage, will be different for different climate regimes.

These results have important implications both for predicting ET in basins where data is not available for calibration and for understanding and predicting the spatial variability of ET within a basin. AWC also affects the sensitivity of annual ET to climate drivers.

Naomi Tague 11/1/2015 7:06 PM

**Deleted:** Lateral redistribution strongly influences the sensitivity of ET to climate drivers in the drier basins; in CA-SIER and CO-ROC there is considerable scatter in the slopes for  $P$  and  $R_{75}$  across a single soil AWC (e.g., for an AWC of 400 mm, the P:ET ranges from 0.6 to 0.8 and 0.2 to 0.7 for CO-ROC and CA-SIER, respectively in Fig. 6A). We note that this additional sensitivity of ET-climate relationships to drainage rates, even given similar AWC or storage conditions, emphasizes the role played by lateral connections. In other words, results suggest that for the two more water limited sites, the timing of upslope contributions to downslope areas can mediate the sensitivity of watershed scale vegetation water use.

Elizabeth Garcia 10/30/2015 10:49 AM

**Moved up [1]:** Our model-based study provides a simplified representation of these interactions, ignoring many additional complexities. In particular, we assume no adaptation of the ecosystem structure and composition that would influence productivity, evapotranspiration and their relationship with climate (Loudermilk et al., 2013). Future work will investigate these coupled carbon cycling-hydrology interactions. In this study we focus on the energy and moisture drivers of E... [3]

Elizabeth Garcia 10/30/2015 8:41 AM

**Deleted:** - ... [4]

Naomi Tague 11/1/2015 6:52 PM

**Deleted:** oil

Naomi Tague 11/1/2015 6:52 PM

**Deleted:** soil parameters

Elizabeth Garcia 10/30/2015 7:43 AM

**Deleted:** e.g

Naomi Tague 11/1/2015 6:52 PM

**Deleted:** soil

Elizabeth Garcia 10/30/2015 9:04 AM

**Moved (insertion) [3]**

Naomi Tague 11/1/2015 7:13 PM

**Deleted:** These results have important implications both for predicting ET in basins where soil data is not available for calib... [5]

Elizabeth Garcia 10/30/2015 9:04 AM

**Moved up [3]:** These results have important for predicting ET in basins where soil data is not available for calibration and for ... [6]

Elizabeth Garcia 10/30/2015 5:54 PM

**Moved up [2]:** Because there is substantial heterogeneity in soil characteristics within each basin (Dahlgren et al., 1997; Denn... [7]

Naomi Tague 11/1/2015 6:53 PM

**Deleted:** soil



962 particularly in the two more seasonally water-limited basins. Although the three  
 963 watersheds show different responses of annual ET to these climate drivers, there are  
 964 values of AWC that would eliminate these cross-basin differences. These sensitivities  
 965 highlight the need for improved information on spatial patterns of subsurface properties  
 966 to contribute to the development of science-based information on forest vulnerabilities to  
 967 climate change. Improved accounting for plant accessibility to moisture has improved  
 968 model-data ET comparisons in previous modeling studies at regional and global scales  
 969 (Hwang et al., 2009; Tang et al., 2013; Thompson et al., 2011). With expected decreases  
 970 in fractional precipitation received as snow with climate change (Diffenbaugh et al.,  
 971 2013; Knowles et al., 2006), we might expect soil storage to play a more important role  
 972 in providing water for forests in the future. Improved understanding of how climate and  
 973 subsurface storage/drainage combine to control ET can enhance our understanding of  
 974 forest water stress related to increased mortality (van Mantgem et al., 2009). Western U.S.  
 975 forests show substantial vulnerability to drought, with declines in productivity and  
 976 increases in mortality and disturbance in drought years (Allen et al., 2010; Hicke et al.,  
 977 2012; Williams et al., 2013). Understanding these ecosystems' responses to primary  
 978 climate drivers is of particular concern given recent warming trends (Sterl et al., 2008)  
 979 and multi-year droughts (Cook et al., 2004; Dai et al., 2004). Identifying the physical  
 980 conditions in which our ability to estimate ET is most sensitive or limited by knowledge  
 981 of subsurface geologic properties helps to prioritize regional data acquisition agendas.  
 982 Integrating results from recent advances in geophysical measurements and models such  
 983 as those emerging from Critical Zone Observatories in the U.S. and elsewhere (Anderson  
 984 et al., 2008) will be essential for analysis of climate ET interactions.

985  
 986 ACKNOWLEDGEMENTS

987  
 988 Data is available upon request from the author. This work was supported by funding from  
 989 the U.S. Geological Survey through the Western Mountain Initiative (Award Number  
 990 G09AC00337) and the U.S. National Science Foundation through the Willamette  
 991 Watershed 2100 Project (EAR-1039192). We also acknowledge support from the  
 992 Southern Sierra Critical Zone Observatory (EAR-0725097) and the Center for Scientific

- Naomi Tague 11/1/2015 7:14 PM  
**Deleted:** Our study
- Naomi Tague 11/1/2015 7:14 PM  
**Deleted:** suggests that
- Naomi Tague 11/1/2015 7:14 PM  
**Deleted:** is needed in
- Naomi Tague 11/1/2015 7:15 PM  
**Deleted:** developing
- Naomi Tague 11/1/2015 7:15 PM  
**Deleted:** these
- Naomi Tague 11/1/2015 6:53 PM  
**Deleted:** soil

- Naomi Tague 11/1/2015 6:53 PM  
**Deleted:** soils
- Elizabeth Garcia 10/30/2015 8:59 AM  
**Deleted:** These findings may also extend to understanding of forest disturbances, contributing to mechanistic explanations for previous work showing that the area burned by wildfire can be correlated to ecosystems and their climates (Littell et al., 2009).
- Elizabeth Garcia 10/30/2015 9:11 AM  
**Deleted:**
- Elizabeth Garcia 10/30/2015 9:10 AM  
**Deleted:** - ... [8]

- Naomi Tague 11/1/2015 6:54 PM  
**Deleted:** oil characterizations
- Elizabeth Garcia 10/30/2015 9:01 AM  
**Deleted:** can also

- 1011 Computing from the CNSI, MRL: an NSF MRSEC (DMR-1121053) and NSF CNS-  
1012 0960316  
1013
- 1014 LITERATURE CITED
- 1015 Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Venetier,  
1016 M., Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E. H. (Ted), Gonzalez, P.,  
1017 Fensham, R., Zhang, Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S. W.,  
1018 Semerci, A. and Cobb, N.: A global overview of drought and heat-induced tree mortality  
1019 reveals emerging climate change risks for forests, *For. Ecol. Manage.*, 259(4), 660–684,  
1020 doi:10.1016/j.foreco.2009.09.001, 2010.
- 1021 Anderson, S. P., Bales, R. C. and Duffy, C. J.: Critical Zone Observatories: Building a  
1022 network to advance interdisciplinary study of Earth surface processes, *Mineral. Mag.*,  
1023 72(1), 7–10, doi:10.1180/minmag.2008.072.1.7, 2008.
- 1024 Arthur, M. and Fahey, T.: Biomass and nutrients in an Engelmann spruce - subalpine fir  
1025 forest in north central Colorado: pools, annual production, and internal cycling, *Can. J.*  
1026 *For. Res.*, 22(3), 315–325, doi:10.1139/x92-041, 1992.
- 1027 Ashfaq, M., Ghosh, S., Kao, S.-C., Bowling, L. C., Mote, P., Touma, D., Rauscher, S. a.  
1028 and Diffenbaugh, N. S.: Near-term acceleration of hydroclimatic change in the western  
1029 U.S., *J. Geophys. Res. Atmos.*, 118(January), 1–18, doi:10.1002/jgrd.50816, 2013.
- 1030 Bales, R., Hopmans, J., O’Green, A., Meadows, M., Hartsough, P., Kirchner, P.,  
1031 Hunsaker, C. and Beaudette, D.: Soil Moisture Response to Snowmelt and Rainfall in a  
1032 Sierra Nevada Mixed-Conifer Forest, *Vadose Zo. J.*, 10(3), 786–799,  
1033 doi:10.2136/vzj2011.0001, 2011.
- 1034 Band, L. E., Tague, C. L., Brun, S. E., Tenenbaum, D. E. and Fernandes, R. A.:  
1035 Modelling Watersheds as Spatial Object Hierarchies: Structure and Dynamics, *Trans.*  
1036 *GIS*, 4(3), 181–196, doi:10.1111/1467-9671.00048, 2000.
- 1037 Barnard, H., Graham, C., van Verseveld, W., Brooks, J. R., Bond, B. J. and McDonnell, J.  
1038 J.: Mechanistic assessment of hillslope transpiration controls of diel subsurface flow: a  
1039 steady-state irrigation approach, *Ecohydrology*, 3, 133–142, doi:10.1002/eco.114, 2010.
- 1040 Baron, J., Hartman, M. and Band, L.: Sensitivity of a high-elevation Rocky Mountain  
1041 watershed to altered climate and CO<sub>2</sub>, *Water Resour. Res.*, 36(1), 89–99 [online]  
1042 Available from: <http://www.agu.org/pubs/crossref/2000/1999WR900263.shtml>  
1043 (Accessed 26 January 2012), 2000.
- 1044 Baron, J. S. and Denning, A.: Hydrologic budget estimates, in *Biogeochemistry of a*



- 1045 Subalpine Ecosystem, edited by J. Baron, pp. 28–47, Springer-Verlag, New York., 1992.
- 1046 Beven, K. J.: Rainfall-runoff modelling: the primer, John Wiley & Sons., 2011.
- 1047 Beven, K. J. and Binley, A.: THE FUTURE OF DISTRIBUTED MODELS: MODEL  
1048 CALIBRATION AND UNCERTAINTY PREDICTION, *Hydrol. Process.*, 6(3), 279–  
1049 298, doi:10.1002/hyp.3360060305, 1992.
- 1050 Boisvenue, C. and Running, S. W.: Impacts of climate change on natural forest  
1051 productivity - evidence since the middle of the 20th century, *Glob. Chang. Biol.*, 12(5),  
1052 862–882, doi:10.1111/j.1365-2486.2006.01134.x, 2006.
- 1053 Bradford, J. B., Birdsey, R. a., Joyce, L. a. and Ryan, M. G.: Tree age, disturbance  
1054 history, and carbon stocks and fluxes in subalpine Rocky Mountain forests, *Glob. Chang.*  
1055 *Biol.*, 14(12), 2882–2897, doi:10.1111/j.1365-2486.2008.01686.x, 2008.
- 1056 Brooks, P. D., Troch, P. a., Durcik, M., Gallo, E. and Schlegel, M.: Quantifying regional  
1057 scale ecosystem response to changes in precipitation: Not all rain is created equal, *Water*  
1058 *Resour. Res.*, 47(10), W00J08, doi:10.1029/2010WR009762, 2011.
- 1059 Brooks, R. and Corey, A.: Hydraulic properties of porous media, in *Hydrology Paper 3*, p.  
1060 27, Colorado State University, Fort Collins., 1964.
- 1061 Cayan, D. R., Dettinger, M. D., Kammerdiener, S. a., Caprio, J. M. and Peterson, D. H.:  
1062 Changes in the Onset of Spring in the Western United States, *Bull. Am. Meteorol. Soc.*,  
1063 82(3), 399–415, doi:10.1175/1520-0477(2001)082<0399:CITOOS>2.3.CO;2, 2001.
- 1064 Cook, E. R., Woodhouse, C. a, Eakin, C. M., Meko, D. M. and Stahle, D. W.: Long-term  
1065 aridity changes in the western United States., *Science*, 306(5698), 1015–8,  
1066 doi:10.1126/science.1102586, 2004.
- 1067 Dahlgren, R. a., Boettinger, J. L., Huntington, G. L. and Amundson, R. G.: Soil  
1068 development along an elevational transect in the western Sierra Nevada, California,  
1069 *Geoderma*, 78(3-4), 207–236, doi:10.1016/S0016-7061(97)00034-7, 1997.
- 1070 Dai, A., Trenberth, K. E. and Qian, T.: A global dataset of Palmer Drought Severity  
1071 Index for 1870-2002: Relationship with soil moisture and effects of surface warming, *J.*  
1072 *Hydrometeorol.*, 5(6), 1117–1130 [online] Available from:  
1073 <http://journals.ametsoc.org/doi/pdf/10.1175/JHM-386.1> (Accessed 29 January 2012),  
1074 2004.
- 1075 Denning, A., Baron, J., Mast, M. and Arthur, M.: Hydrologic pathways and chemical  
1076 composition of runoff during snowmelt in Loch Vale watershed, Rocky Mountain  
1077 National Park, Colorado, USA, *Water. Air. Soil Pollut.*, 59(1-2), 107–123 [online]  
1078 Available from: <http://link.springer.com/article/10.1007/BF00283175> (Accessed 3

- 1079 October 2013), 1991.
- 1080 Diffenbaugh, N. S., Scherer, M. and Ashfaq, M.: Response of snow-dependent  
1081 hydrologic extremes to continued global warming., *Nat. Clim. Chang.*, 3(4), 379–384,  
1082 doi:10.1038/nclimate1732, 2013.
- 1083 Dingman, S. L.: *Physical Hydrology*, 2nd ed., Prentice Hall, Englewood Cliffs, NJ., 1994.
- 1084 Famiglietti, J. S., Ryu, D., Berg, A. a., Rodell, M. and Jackson, T. J.: Field observations  
1085 of soil moisture variability across scales, *Water Resour. Res.*, 44(1), W01423,  
1086 doi:10.1029/2006WR005804, 2008.
- 1087 Farquhar, G. D., Caemmerer, S. Von and Berry, J. A.: A biochemical model of  
1088 photosynthetic CO<sub>2</sub> assimilation in leaves of C<sub>3</sub> species, *Planta*, 149, 78–90 [online]  
1089 Available from: <http://link.springer.com/article/10.1007/BF00386231> (Accessed 30 April  
1090 2013), 1980.
- 1091 Flock, J.: Lichen-Bryophyte Distribution along a Snow-Cover-Soil-Moisture Gradient,  
1092 Niwot Ridge, Colorado, *Arct. Alp. Res.*, 10(1), 31–47 [online] Available from:  
1093 <http://www.jstor.org/stable/10.2307/1550655> (Accessed 7 October 2013), 1978.
- 1094 Garcia, E. S., Tague, C. L. and Choate, J. S.: Method of spatial temperature estimation  
1095 influences ecohydrologic modeling in the Western Oregon cascades, *Water Resour. Res.*,  
1096 49, 1611–1624, doi:10.1002/wrcr.20140, 2013.
- 1097 Geroy, I. J., Gribb, M. M., Marshall, H. P., Chandler, D. G., Benner, S. G. and  
1098 McNamara, J. P.: Aspect influences on soil water retention and storage, *Hydrol. Process.*,  
1099 25(25), 3836–3842, doi:10.1002/hyp.8281, 2011.
- 1100 Gholz, H. L.: Environmental limits on aboveground net primary production, leaf area,  
1101 and biomass in vegetation zones of the Pacific Northwest, *Ecology*, 63(2), 469–481  
1102 [online] Available from: <http://www.esajournals.org/doi/abs/10.2307/1938964> (Accessed  
1103 30 November 2011), 1982.
- 1104 Goulden, M. L., Anderson, R. G., Bales, R. C., Kelly, a. E., Meadows, M. and Winston,  
1105 G. C.: Evapotranspiration along an elevation gradient in California’s Sierra Nevada, *J.*  
1106 *Geophys. Res.*, 117(G3), G03028, doi:10.1029/2012JG002027, 2012.
- 1107 Grant, G. E., Tague, C. L. and Allen, C. D.: Watering the forest for the trees: an emerging  
1108 priority for managing water in forest landscapes, *Front. Ecol. Environ.*, 11(6), 314–321,  
1109 doi:10.1890/120209, 2013.
- 1110 Grier, C. C. and Logan, R. S.: Old-growth *Pseudotsuga menziesii* communities of a  
1111 western Oregon watershed: biomass distribution and production budgets, *Ecol. Monogr.*,  
1112 47(4), 373–400 [online] Available from: <http://www.jstor.org/stable/10.2307/1942174>

- 1113 (Accessed 18 August 2013), 1977.
- 1114 Hamlet, A. F., Mote, P. W., Clark, M. P. and Lettenmaier, D. P.: Twentieth-Century  
1115 Trends in Runoff, Evapotranspiration, and Soil Moisture in the Western United States\*, J.  
1116 Clim., 20(8), 1468–1486, doi:10.1175/JCLI4051.1, 2007.
- 1117 Hanson, P. and Weltzin, J.: Drought disturbance from climate change: response of United  
1118 States forests, *Sci. Total Environ.*, 262(3), 205–220, doi:10.1016/S0048-9697(00)00523-  
1119 4, 2000.
- 1120 Hicke, J. a., Allen, C. D., Desai, A. R., Dietze, M. C., Hall, R. J., Ted Hogg, E. H.,  
1121 Kashian, D. M., Moore, D., Raffa, K. F., Sturrock, R. N. and Vogelmann, J.: Effects of  
1122 biotic disturbances on forest carbon cycling in the United States and Canada, *Glob.*  
1123 *Chang. Biol.*, 18(1), 7–34, doi:10.1111/j.1365-2486.2011.02543.x, 2012.
- 1124 Homer, C., Dewitz, J., Fry, J., Coan, M., Hossain, N., Larson, C., Herold, N., Mckerrow,  
1125 A., Vandriel, J. N. and Wickham, J.: Completion of the 2001 National Land Cover  
1126 Database for the Conterminous United States, *Photogramm. Eng. Remote Sensing*, 73(4),  
1127 337–341, 2007.
- 1128 Hudiburg, T., Law, B., Turner, D. P., Campbell, J., Donato, D. and Duane, M.: Carbon  
1129 dynamics of Oregon and Northern California forests and potential land-based carbon  
1130 storage., *Ecol. Appl.*, 19(1), 163–180, doi:10.1890/07-2006.1, 2009.
- 1131 Hwang, T., Band, L. and Hales, T. C.: Ecosystem processes at the watershed scale:  
1132 Extending optimality theory from plot to catchment, *Water Resour. Res.*, 45(11),  
1133 W11425, doi:10.1029/2009WR007775, 2009.
- 1134 Isaac, V. and van Wijngaarden, W. a.: Surface Water Vapor Pressure and Temperature  
1135 Trends in North America during 1948–2010, *J. Clim.*, 25(10), 3599–3609,  
1136 doi:10.1175/JCLI-D-11-00003.1, 2012.
- 1137 Jarvis, P.: The interpretation of the variations in leaf water potential and stomatal  
1138 conductance found in canopies in the field, *Philos. Trans. R. Soc. London. B, Biol. Sci.*,  
1139 273(927), 593–610, doi:10.1098/rstb.1976.0035, 1976.
- 1140 Kampf, S., Markus, J., Heath, J. and Moore, C.: Snowmelt runoff and soil moisture  
1141 dynamics on steep subalpine hillslopes, *Hydrol. Process.*, 29(5), 712–723,  
1142 doi:10.1002/hyp.10179, 2014.
- 1143 Knowles, N., Dettinger, M. D. and Cayan, D. R.: Trends in snowfall versus rainfall in the  
1144 western United States, *J. Clim.*, 19(18), 4545–4559, doi:10.1175/JCLI3850.1, 2006.
- 1145 Landsberg, J. and Waring, R.: A generalised model of forest productivity using simplified  
1146 concepts of radiation-use efficiency, carbon balance and partitioning, *For. Ecol. Manage.*,

- 1147 95(3), 209–228 [online] Available from:  
1148 <http://www.sciencedirect.com/science/article/pii/S0378112797000261> (Accessed 30  
1149 November 2011), 1997.
- 1150 Litaor, M. I., Williams, M. and Seastedt, T. R.: Topographic controls on snow  
1151 distribution, soil moisture, and species diversity of herbaceous alpine vegetation, Niwot  
1152 Ridge, Colorado, *J. Geophys. Res.*, 113(G2), G02008, doi:10.1029/2007JG000419, 2008.
- 1153 Loudermilk, E. L., Scheller, R. M., Weisberg, P. J., Yang, J., Dilts, T. E., Karam, S. L.  
1154 and Skinner, C.: Carbon dynamics in the future forest: the importance of long-term  
1155 successional legacy and climate-fire interactions., *Glob. Chang. Biol.*, 19(11), 3502–3515,  
1156 doi:10.1111/gcb.12310, 2013.
- 1157 van Mantgem, P. J., Stephenson, N. L., Byrne, J. C., Daniels, L. D., Franklin, J. F., Fulé,  
1158 P. Z., Harmon, M. E., Larson, A. J., Smith, J. M., Taylor, A. H. and Veblen, T. T.:  
1159 Widespread increase of tree mortality rates in the western United States., *Science*,  
1160 323(5913), 521–4, doi:10.1126/science.1165000, 2009.
- 1161 McGuire, K. J., Weiler, M. and McDonnell, J. J.: Integrating tracer experiments with  
1162 modeling to assess runoff processes and water transit times, *Adv. Water Resour.*, 30(4),  
1163 824–837, doi:10.1016/j.advwatres.2006.07.004, 2007.
- 1164 McNamara, J. P., Chandler, D., Seyfried, M. and Achet, S.: Soil moisture states, lateral  
1165 flow, and streamflow generation in a semi-arid, snowmelt-driven catchment, *Hydrol.*  
1166 *Process.*, 19(20), 4023–4038, doi:10.1002/hyp.5869, 2005.
- 1167 McNamara, J. P., Tetzlaff, D., Bishop, K., Soulsby, C., Seyfried, M., Peters, N. E.,  
1168 Aulenbach, B. T. and Hooper, R.: Storage as a metric of catchment comparison, *Hydrol.*  
1169 *Process.*, 25, 3364–3371, doi:10.1002/hyp.8113, 2011.
- 1170 Monteith, J.: Evaporation and Environment, in *Proceedings of the 19th Symposium of the*  
1171 *Society for Experimental Biology*, vol. 19, pp. 205–234, Cambridge University Press,  
1172 New York. [online] Available from:  
1173 <http://www.unc.edu/courses/2007fall/geog/801/001/www/ET/Monteith65.pdf> (Accessed  
1174 30 November 2011), 1965.
- 1175 Moore, I. D., Grayson, R. B. and Ladson, A. R.: DIGITAL TERRAIN MODELLING : A  
1176 REVIEW OF HYDROLOGICAL, GEOMORPHOLOGICAL, AND BIOLOGICAL  
1177 APPLICATIONS, *Hydrol. Process.*, 5(1), 3–30, 1991.
- 1178 Mote, P. W., Hamlet, A. F., Clark, M. P. and Lettenmaier, D. P.: Declining Mountain  
1179 Snowpack in Western North America\*, *Bull. Am. Meteorol. Soc.*, 86(1), 39–49,  
1180 doi:10.1175/BAMS-86-1-39, 2005.
- 1181 Muggeo, V. M. R.: Estimating regression models with unknown break-points., *Stat. Med.*,

- 1182 22(19), 3055–71, doi:10.1002/sim.1545, 2003.
- 1183 Nash, J. E. and Sutcliffe, J.: River flow forecasting through conceptual models part I —  
1184 A discussion of principles, *J. Hydrol.*, 10(3), 282–290 [online] Available from:  
1185 <http://www.sciencedirect.com/science/article/pii/0022169470902556> (Accessed 28  
1186 November 2011), 1970.
- 1187 Natural Resources Conservation Service, S. S. S.: Soil Survey Geographic (SSURGO)  
1188 Database. [online] Available from: <http://sdmdataaccess.nrcs.usda.gov/>, n.d.
- 1189 Post, D. A. and Jones, J. A.: Hydrologic regimes of forested, mountainous, headwater  
1190 basins in New Hampshire, North Carolina, Oregon, and Puerto Rico, *Adv. Water Resour.*,  
1191 24(9), 1195–1210, doi:10.1016/S0309-1708(01)00036-7, 2001.
- 1192 Running, S. and Nemani, R.: Extrapolation of synoptic meteorological data in  
1193 mountainous terrain and its use for simulating forest evapotranspiration and  
1194 photosynthesis, *Can. J. For. Res.*, 17(6), 472–483 [online] Available from:  
1195 <http://www.nrcresearchpress.com/doi/abs/10.1139/x87-081#.VctfPmRViko> (Accessed 30  
1196 November 2011), 1987.
- 1197 Ryan, M. G.: Effects of climate change on plant respiration, *Ecol. Appl.*, 1(2), 157–167  
1198 [online] Available from: <http://www.esajournals.org/doi/abs/10.2307/1941808> (Accessed  
1199 1 December 2011), 1991.
- 1200 Shields, C. A. and Tague, C. L.: Assessing the Role of Parameter and Input Uncertainty  
1201 in Ecohydrologic Modeling: Implications for a Semi-arid and Urbanizing Coastal  
1202 California Catchment, *Ecosystems*, 15(5), 775–791, doi:10.1007/s10021-012-9545-z,  
1203 2012.
- 1204 Smith, T. J., McNamara, J. P., Flores, A. N., Gribb, M. M., Aishlin, P. S. and Benner, S.  
1205 G.: Small soil storage capacity limits benefit of winter snowpack to upland vegetation,  
1206 *Hydrol. Process.*, 25(25), 3858–3865, doi:10.1002/hyp.8340, 2011.
- 1207 Sterl, A., Severijns, C., Dijkstra, H., Hazeleger, W., Jan van Oldenborgh, G., van den  
1208 Broeke, M., Burgers, G., van den Hurk, B., Jan van Leeuwen, P. and van Velthoven, P.:  
1209 When can we expect extremely high surface temperatures?, *Geophys. Res. Lett.*, 35(14),  
1210 L14703, doi:10.1029/2008GL034071, 2008.
- 1211 Stewart, I. T., Cayan, D. R. and Dettinger, M. D.: Changes toward Earlier Streamflow  
1212 Timing across Western North America, *J. Clim.*, 18(8), 1136–1155,  
1213 doi:10.1175/JCLI3321.1, 2005.
- 1214 Tague, C. and Band, L.: RHESSys: regional hydro-ecologic simulation system-an object-  
1215 oriented approach to spatially distributed modeling of carbon, water, and nutrient cycling,  
1216 *Earth Interact.*, 8(19), 1–42, doi:10.1175/1087-3562(2004)8<1:RRHSSO>2.0.CO;2, 2004.

- 1217 Tague, C. and Heyn, K.: Topographic controls on spatial patterns of conifer transpiration  
1218 and net primary productivity under climate warming in mountain ecosystems,  
1219 *Ecohydrology*, 554(October), 541– 554, doi:10.1002/eco.88, 2009.
- 1220 Tague, C. and Peng, H.: The sensitivity of forest water use to the timing of precipitation  
1221 and snowmelt recharge in the California Sierra: Implications for a warming climate, *J.*  
1222 *Geophys. Res. Biogeosciences*, 118, 1–13, doi:10.1002/jgrg.20073, 2013.
- 1223 Tang, J., Pilesjö, P., Miller, P. a., Persson, A., Yang, Z., Hanna, E. and Callaghan, T. V.:  
1224 Incorporating topographic indices into dynamic ecosystem modelling using LPJ-GUESS,  
1225 *Ecohydrology*, 7(4), 1147–1162, doi:10.1002/eco.1446, 2013.
- 1226 Thompson, S., Harman, C., Konings, A., Sivapalan, M., Neal, A. and Troch, P.:  
1227 Comparative hydrology across AmeriFlux sites: The variable roles of climate, vegetation,  
1228 and groundwater, *Water Resour. Res.*, 47(null), W00J07, doi:10.1029/2010WR009797,  
1229 2011.
- 1230 Thornton, P. E.: Description of a numerical simulation model for predicting the dynamics  
1231 of energy, water, carbon, and nitrogen in a terrestrial ecosystem, University of Montana,  
1232 Missoula, MT., 1998.
- 1233 Thornton, P. E. and Rosenbloom, N. A.: Ecosystem model spin-up: Estimating steady  
1234 state conditions in a coupled terrestrial carbon and nitrogen cycle model, *Ecol. Modell.*,  
1235 189(1-2), 25–48, doi:10.1016/j.ecolmodel.2005.04.008, 2005.
- 1236 Thornton, P., Thornton, M., Mayer, B., Wilhelmi, N., Wei, Y. and Cook, R.: Daymet:  
1237 Daily surface weather on a 1 km grid for North America, 1980-2012, Oak Ridge Natl.  
1238 Lab. Distrib. Act. Arch. Cent., doi:10.3334/ORNLDAAC/Daymet\_V2, 2012.
- 1239 Trujillo, E., Molotch, N. P., Goulden, M. L., Kelly, A. E. and Bales, R. C.: Elevation-  
1240 dependent influence of snow accumulation on forest greening, *Nat. Geosci.*, 5(10), 705–  
1241 709, doi:10.1038/ngeo1571, 2012.
- 1242 Van Tuyl, S., Law, B. E., Turner, D. P. and Gitelman, a. I.: Variability in net primary  
1243 production and carbon storage in biomass across Oregon forests—an assessment  
1244 integrating data from forest inventories, intensive sites, and remote sensing, *For. Ecol.*  
1245 *Manage.*, 209(3), 273–291, doi:10.1016/j.foreco.2005.02.002, 2005.
- 1246 Vicente-Serrano, S. M., Camarero, J. J., Zabalza, J., Sangüesa-Barreda, G., López-  
1247 Moreno, J. I. and Tague, C. L.: Evapotranspiration deficit controls net primary production  
1248 and growth of silver fir: Implications for Circum-Mediterranean forests under forecasted  
1249 warmer and drier conditions, *Agric. For. Meteorol.*, 206, 45–54,  
1250 doi:10.1016/j.agrformet.2015.02.017, 2015.
- 1251 Webb, W., Szarek, S., Lauenroth, W., Kinerson, R. and Smith, M.: Primary Productivity

- 1252 and Water Use in Native Forest, Grassland, and Desert Ecosystems, *Ecology*, 59(6),  
1253 1239–1247, doi:10.2307/1938237, 1978.
- 1254 Williams, A., Allen, C., Macalady, A., Griffin, D., Woodhouse, C., Meko, D., Swetnam,  
1255 T. W., Rauscher, S. a. and Seager, R.: Temperature as a potent driver of regional forest  
1256 drought stress and tree mortality, *Nat. Clim. Chang.*, 3(September), 292–297,  
1257 doi:10.1038/NCLIMATE1693, 2013.
- 1258 Zierl, B., Bugmann, H. and Tague, C.: Water and carbon fluxes of European ecosystems:  
1259 An evaluation of the ecohydrological model RHESSys, *Hydrol. Process.*, 21(24), 3328–  
1260 3339, doi:10.1002/hyp.6540, 2007.
- 1261
- 1262

1263 Table 1. Explanatory variables

Abbreviation	Definition
P	Total annual precipitation
T <sub>AMJ</sub>	Average daily temperature for April, May, June
R <sub>75</sub>	Day of water year that 75% of soil water recharge occurs
AWC	Available water capacity of soil (field capacity-wilting point)

1264

1265



1266 Table 2. Basin topography, geology, vegetation and climate characteristics. Climate  
 1267 descriptions are averaged over total available climate record (duration noted in table).

Watershed	CO-ROC	OR-CAS	CA-SIER
Location	Colorado	Oregon	California
U.S. Geological Survey gage number	06733000	14161500	10343500
Geology	Holocene glacial till, rock; Precambrian gneiss, granite	Western Cascade basalt	Sierra granite, with Miocene andesite cap
Elevation range (m)	1470-4345	410-1630	1800-2650
Drainage Area (km <sup>2</sup> )	350	64	26
Topographic Wetness Index- Mean (Std Dev)	7.0 (1.9)	6.6 (1.7)	7.9 (1.8)
Climate record	1980 – 2008	1958-2008	1960-2000
Mean Annual Precipitation (mm)	1000	2250	850
Annual Precipitation as snow (%)	64	29	55
Precipitation received in Growing Season (%)	46	21	19
Min/Max winter <i>T</i> (JFM) (oC)	-12.1/-0.02	-0.9 / 5.2	-9.5/3.7
Min/Max spring <i>T</i> (AMJ) (oC)	-2.7/10.9	4.0/14.0	-2.5/13.8
<a href="#">P:PET</a>	<a href="#">0.9</a>	<a href="#">2.3</a>	<a href="#">1.2</a>
Vegetation	Subalpine fir, aspen, meadows, shrub	Douglas-fir, Western Hemlock	Mixed Conifer, Jeffrey and Lodgepole Pine
Mean basin LAI	3.5	9.0	4.1
Annual NPP range for calibration (gC m <sup>-2</sup> yr <sup>-1</sup> )	280-520	620-1100	450-800
Literature sources used to bound annual NPP range	Arthur and Fahey [1992] Bradford et al. [2008]	Grier and Logan [1977] Gholz [1982]	Hudiburg et al. [2009] Goulden et al. [2012] <sup>a</sup>

1268 <sup>a</sup>Values reported as gross primary productivity, converted to NPP using RHESSys  
 1269 calculated values of respiration.

1270

1271 Table 3. Statistics for ET predictors based on linear regression models.

Watershed		CO-ROC	OR-CAS	CA-SIER
Precipitation (P)	p-value	< 0.001	< 0.05	< 0.001
	r <sup>2</sup>	0.9	0.1	0.75
	slope	0.4	0.1	0.2
Timing (R <sub>75</sub> )	p-value	<0.001	< 0.01	<0.001
	r <sup>2</sup>	0.2	0.2	0.4
	slope	3.8	1.2	4.6
Temperature T <sub>AMJ</sub>	p-value	<0.001	<0.05	>0.1
	r <sup>2</sup>	0.4	0.1	-0.01
	slope	-26.3	-25.7	15
Soil Capacity (AWC)	p-value	0.001	0.001	0.001
	r <sup>2</sup>	0.43	0.53	0.11
	slope	0.1	0.2	0.1

1272

1273

Elizabeth Garcia 10/30/2015 7:17 AM  
Formatted: Superscript

1274 **Figure Captions**

1275

1276 Figure 1. Locations and average daily water fluxes averaged from 1980-2000 for three  
1277 case study watersheds located in (A) the western Oregon Cascades (OR-CAS), (B)  
1278 Colorado Rockies (CO-ROC), and (C) California Sierra Nevada (CA-SIER).

1279

1280 Figure 2. (A) Total annual ET increases with total annual precipitation. Lines indicate  
1281 statistically significant relationships ( $p\text{-value} < 0.05$ ).

1282

1283 Figure 3. Later occurrence of soil moisture recharge ( $R_{75}$ ) is significantly correlated with  
1284 increased annual ET in all study watersheds.

1285

1286 Figure 4. (A) Warmer spring temperatures are correlated with lower total annual ET in  
1287 the two snow-dominated watersheds. (B) An earlier occurrence of soil moisture recharge  
1288 is correlated with warmer temperatures in CO-ROC.


1289

1290 Figure 5. Each point represents the 15-year average annual ET from WY 1985-2000 for a  
1291 physically viable mean basin soil available water capacity (AWC). Vertical lines  
1292 represent the calculated breakpoint in the nonlinear relationship between long-term ET  
1293 and AWC for each basin.

1294

1295 Figure 6. The impact of soil AWC on the slope a linear regression model of annual ET as  
1296 a function of climate predictors: (A) precipitation, (B)  $R_{75}$ , and (C)  $T_{AMJ}$ . The slope of  
1297 ET:climate predictor is plotted across a physically viable range of mean basin soil AWC  
1298 for each climate predictor and for each study basin: OR-CAS (left column), CO-ROC  
1299 (middle column), and CA-SIER (right column). The slopes are normalized to facilitate  
1300 inter-basin comparison.

1301




1302

1303 Figure 1. Locations and average daily water fluxes averaged from 1980-2000 for three

1304 case study watersheds located in (A) the western Oregon Cascades (OR-CAS), (B)

1305 Colorado Rockies (CO-ROC), and (C) California Sierra Nevada (CA-SIER).


1306




1307

1308 Figure 2. (A) Total annual ET increases with total annual precipitation. Lines indicate  
 1309 statistically significant relationships ( $p$ -value < 0.05).


1310



1311 Figure 3. Later occurrence of soil moisture recharge ( $R_{75}$ ) is significantly correlated with  
 1312 increased annual ET in all study watersheds.




1313 Figure 4. (A) Warmer spring temperatures are correlated with lower total annual ET in  
 1314 the two snow-dominated watersheds. (B) An earlier occurrence of soil moisture recharge  
 1315 is correlated with warmer temperatures in CO-ROC.  
 1316



1317  
 1318 Figure 5. Each point represents the 15-year average annual ET from WY 1985-2000 for a  
 1319 physically viable mean basin soil available water capacity (AWC). Vertical lines  
 1320 represent the calculated breakpoint in the nonlinear relationship between long-term ET  
 1321 and AWC for each basin.





1322

1323 Figure 6. The impact of soil AWC on the slope a linear regression model of annual ET as  
 1324 a function of climate predictors: (A) precipitation, (B)  $R_{75}$ , and (C)  $T_{AMJ}$ . The slope of  
 1325 ET:climate predictor is plotted across a physically viable range of mean basin soil AWC  
 1326 for each climate predictor and for each study basin: OR-CAS (left column), CO-ROC

1327 (middle column), and CA-SIER (right column).The slopes are normalized to facilitate  
1328 inter-basin comparison.

