

## Responses to minor comments.

We are grateful of both reviewers help in improving our manuscript. Below we provide detailed responses to their last round of comments.

### **Report 1**

I just have one small comment to the Fig.2. :

Comment	Response
-to change hidro-climatological in the caption	Done
-to clarify what is the meaning of total SWE	We added explanatory text in the caption
-and maybe to revise the runoff/basin area caption by using specific runoff	We modified the title in the Y-axis following the reviewer's suggestion

### **Report 2**

Comment	Response
Lines 36-38: this statement is difficult to be understood here, I suggest removing or rephrasing;	We modified the wording in this section.
Line 82: I think it would be useful here to specify that SWE reconstruction focuses on the depletion season;	Added a sentence to this effect
Line 662: it may be useful citing Figure S4 here?	We agree, and added a citation to Figure S4
Line 744 - 746: if I understand results properly, in POR model estimates are usually closer to quartiles than to median values, whereas during 2013 in LAG model estimate is close to minimum observed SWE during manual operations. This is not a problem for your results, but it may be useful to add some additional specifications here on this;	We added a more specific description int he text
Line 780 - 786: do these R2 refer to a 1:1 line? Please specify this. If not, providing regression equations may be useful;	R2 values refer to the best fit line. We added a sentence to this effect, and provide slope and intercept values in table 4

Line 816: unimpaired flow is already mentioned at line 705 so it may be useful moving "(no human extractions)" there;	Done
Line 825, 829, 831, 846: please consider comment 5 here again;	In this case R2 values also refer to the best fit line. However, application of our results to streamflow forecasting is the subject of future work and we prefer to limit the discussion here to correlation values only.
Line 1142: please specify if 0.80 is an R2;	Done.

1 Spatio-temporal variability of snow water equivalent in the extra-tropical Andes  
2 cordillera from distributed energy balance modeling and remotely sensed snow  
3 cover.

4

5 Edward Cornwell<sup>1</sup>, Noah P. Molotch<sup>2,3</sup> and James McPhee<sup>\*1,4</sup>

6 *1: Advanced Mining Technology Center, Facultad de Ciencias Físicas y Matemáticas,  
7 Universidad de Chile*

8 *2: Department of Geography and Institute of Arctic and Alpine Research, University of  
9 Colorado, Boulder*

10 *3: Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California,  
11 USA.*

12 *4: Departamento de Ingeniería Civil, Facultad de Ciencias Físicas y Matemáticas,  
13 Universidad de Chile*

14

15 *\*: corresponding author: jmcphee@u.uchile.cl*

16

## 17 Abstract

18 Seasonal snow cover is the primary water source for human use and ecosystems along  
19 the extratropical Andes Cordillera. Despite its importance, relatively little research has  
20 been devoted to understanding the properties, distribution and variability of this natural  
21 resource. This research provides high-resolution (500-meter), daily distributed estimates  
22 of end-of-winter and spring snow water equivalent over a 152,000-km<sup>2</sup> domain that  
23 includes the mountainous reaches of central Chile and Argentina. Remotely sensed  
24 fractional snow covered area and other relevant forcings are combined with extrapolated  
25 data from meteorological stations and a simplified physically-based energy balance  
26 model in order to obtain melt-season melt fluxes which are then aggregated to estimate  
27 end-of-winter (or peak) snow water equivalent (SWE). Peak SWE estimates show an  
28 overall coefficient of determination R<sup>2</sup> of 0.68 and RMSE of 274 mm compared to  
29 observations at 12 automatic snow water equivalent sensors distributed across the model  
30 domain, with R<sup>2</sup> values between 0.32 and 0.88. Regional estimates of peak SWE

31 accumulation show differential patterns strongly modulated by elevation, latitude and  
32 position relative to the continental divide. The spatial distribution of peak SWE shows  
33 that the 4000 - 5000 m.a.s.l. elevation band is significant for snow accumulation,  
34 despite having a smaller surface area than the 3000 – 4000 m.a.s.l. band. On average,  
35 maximum snow accumulation is observed in early September in the western Andes, and  
36 in early October on the eastern side of the continental divide. The results presented here  
37 have the potential of informing applications such as seasonal forecast model assessment  
38 and improvement, regional climate model validation, as well as evaluation of  
39 observational networks and water resource infrastructure development.

40

## 41 1 Introduction

42 Accurately predicting the spatial and temporal distribution of snow water equivalent  
43 (SWE) in mountain environments remains a significant challenge for the scientific  
44 community and water resource practitioners around the world. The Andes Cordillera, a  
45 formidable mountain range that constitutes the backbone of the South American  
46 continent, remains one of the relatively least studied mountain environments due to its  
47 generally low accessibility and complex topography. The extratropical stretch of the  
48 Andes, extending south from approximately latitude 27 ° S, is a snow-dominated  
49 hydrological environment that provides key water resources for a majority of the  
50 population in Chile and Argentina. Until now, a very sparse network of snow courses  
51 and automated snow measuring stations (snow pillows) has been the only source of  
52 information about this key resource. In a context of sustained climate change  
53 characterized by warming trends and likely future precipitation reductions (Vera et al.,  
54 2006; Vicuña et al., 2011), it becomes ever more relevant to understand the past  
55 dynamics of the seasonal snowpack in order to validate predictive models of future

James McPhee 1/1/2016 11:37
<b>Eliminado:</b> two
James McPhee 1/1/2016 11:37
<b>Eliminado:</b> snow-volume storage zones characterized by a larger areal extent (3000-4000 m.a.s.l.) and greater
James McPhee 1/1/2016 11:37
<b>Eliminado:</b> (
James McPhee 1/1/2016 11:37
<b>Eliminado:</b> - 5000
James McPhee 1/1/2016 11:37
<b>Eliminado:</b> ..) respectively
James McPhee 1/1/2016 11:37
<b>Eliminado:</b> peaks
James McPhee 1/1/2016 11:37
<b>Eliminado:</b> whereas maximum accumulation occurs

66 snow-water resources. This research presents the first spatially and temporally explicit  
67 high-resolution SWE reconstruction over the snow-dominated extratropical Andes of  
68 central Chile and Argentina based on a physical representation of the snowpack energy  
69 balance (Kustas et al., 1994) and remotely sensed snow extent (Dietz et al., 2012)  
70 between years 2001 and 2014. A key advantage of the presented product is its  
71 independence from notoriously scarce and unreliable precipitation measurements at  
72 high elevations. Estimates of maximum SWE accumulation and depletion curves are  
73 obtained at 500-m resolution, coincident with the MODIS Fractional Snow Cover  
74 product MOD10A1 (Hall et al., 2002).

75 Patterns of hydroclimatic spatiotemporal variability in the extratropical Andes have  
76 been studied with increased intensity over the last couple of decades, as pressure for  
77 water resources has mounted while at the same time rapid changes in land use and  
78 climate have highlighted the societal need for increased understanding of water resource  
79 variability and trends under present and future climates. The vast majority of studies  
80 have relied on statistical analyses of instrumental records and regional climate models  
81 to present synoptic-scale summaries of precipitation (e.g. Aravena and Luckman, 2009;  
82 Falvey and Garreaud, 2007; Garreaud, 2009), temperature (Falvey and Garreaud, 2009),  
83 snow accumulation (Masiokas et al., 2006) and streamflow variability (Cortés et al.,  
84 2011; Núñez et al., 2013). Currently, no high-resolution, large-scale distributed  
85 assessments of snow water equivalent are available for the Andes region.

86 | The SWE reconstruction method seeks to estimate end-of-winter accumulation by back  
87 | accumulating melt energy fluxes during the depletion season. The methods and  
88 assumptions required for SWE reconstruction have been tested and refined since initial  
89 development (Cline et al., 1998). Applications across a variety of scales have been  
90 presented in recent years. In the Sierra Nevada, Jepsen et al. (2012) compared SWE

reconstructions to distributed snow surveys in a 19.1 km<sup>2</sup> basin ( $R^2 = 0.79$ ), while Guan et al. (2013) obtained good correlation with SWE observations from an operational snow sensor network across the entire Sierra Nevada ( $R^2 = 0.74$ ). In the Rocky Mountains, Jepsen et al. (2012) obtained an  $R^2$  value of 0.61 when comparing reconstructed SWE to spatial regression from snow surveys, and Molotch (2009) estimated SWE with a mean absolute error (MAE) of 23% compared to intensive study areas. A useful discussion on the uncertainties of the SWE reconstruction method -albeit one based on temperature-index melt equations- was presented by Slater et al. (2013), who demonstrated that errors in forcing data are at least, if not more, important than snow covered area data availability. The vast majority, if not all, of SWE reconstruction exercises have been developed in the northern hemisphere, under environmental conditions quite different from those predominant in the extratropical Andes Cordillera. Here, snow distribution and properties have been analyzed in a few local studies (e.g. Ayala et al., 2014; Cortés et al., 2014; Gascoin et al., 2013), but no large-scale estimations at a relevant temporal and spatial resolution for hydrologic applications have been presented. In fact, the Andes of Chile and Argentina display near-ideal conditions for the SWE reconstruction approach due to (1) the near absence of forest cover over a large fraction of the domain where snow accumulation is hydrologically significant; (2) the sharp climatological distinction between wet (winter: June through August) and dry (spring/summer: September through March) seasons, with most of annual precipitation falling during the former; and (3) the low prevalence of cloudy conditions during spring and summer months over the mountains, which afford a high availability of remotely sensed snow cover information. Conversely, the SWE reconstruction presented here is certainly subject to a series of uncertainty sources, such

115 as the sparseness of the hydrometeorological observational network, which limits both  
116 the availability of forcing and validation data.

117 However, this is the first estimation of peak SWE and snow depletion distribution at  
118 this scale and spatial resolution for the extratropical Andes, and the information shown  
119 here can be useful for several applications such as understanding year-to year  
120 differential accumulation patterns that may impact the performance of seasonal  
121 streamflow forecast models that rely on point-scale data only. Also, the SWE  
122 reconstruction can be used to validate output from global or regional climate models  
123 and reanalysis, which are being increasingly employed to estimate hydrological states  
124 and fluxes in ungauged regions. By analyzing the spatial correlation of snow  
125 accumulation and hydrometeorological variables, distributed SWE estimates can inform  
126 the design of improved climate observation networks. Likewise, from analyzing the  
127 obtained SWE estimates in light of the necessary modeling assumptions and data  
128 availability we are able to highlight future research directions aimed at quantifying and  
129 reducing these uncertainties.

130 The objectives of this research include: 1) To assess the dominant patterns of spatio-  
131 temporal variability in snow water equivalent of the snow-dominated extratropical  
132 Andes cordillera; and, 2) to explicitly evaluate the strengths and weaknesses of the  
133 SWE reconstruction approach in different sub-regions of the extratropical Andes using  
134 snow sensors and distributed snow surveys.

## 135 2 Study Area

136 | [Figure 1](#) shows the study area, which includes headwater basins in the Andes Mountains  
137 of central Chile and Argentina, between 27 ° S and 38 ° S. These basins supply fresh  
138 water to low valleys located on both sides of the Cordillera, a topographic barrier more

James McPhee 1/1/2016 11:37  
**Eliminado:** Figure 1

140 than 5 km high which strongly controls the spatial variability in atmospheric processes  
141 (Garreaud, 2009; Montgomery et al., 2001). In Chile, runoff from the Andes Mountains  
142 benefits 75% of the population (<http://www.ine.cl>) as well as most of the country's  
143 agricultural output, hydropower and industrial activities. In the case of Argentina, 7% of  
144 the population is located in the provinces of La Rioja, San Juan, Mendoza and northern  
145 Neuquén (<http://www.indec.gov.ar/>), with primary water uses in agriculture and  
146 hydropower. The selected watersheds have unimpeded streamflow observations and a  
147 snow-dominated hydrologic regime (Figure 2). River basins included in this study have  
148 been grouped in eight clusters, or hydrologic response units, based on the seasonality of  
149 river flow; numbered C1 to C8 in Figure 1b. Due to differences in topography and  
150 locations of stream gages, the number of headwater basins contained within clusters  
151 differs markedly on both sides of the Cordillera, with larger watersheds on the  
152 Argentinean side.

153 The hydro-climate is mostly controlled by orographic effects on precipitation (Falvey  
154 and Garreaud, 2007) and inter-annual variability associated with the Pacific Ocean  
155 through the El Niño-Southern Oscillation and Pacific Decadal Oscillation (Masiokas et  
156 al., 2006; Newman et al., 2003; Rubio-Álvarez and McPhee, 2010). Precipitation is  
157 concentrated in winter months on the western slope (Aceituno, 1988) and sporadic  
158 spring and summer storms occur on the mountain front plains of the eastern slope. The  
159 vegetation cover presents a steppe type condition on the west slope up to 33 ° S,  
160 transitioning to the south into tall bushes and sparse mountain forest. On the eastern  
161 slope the steppe vegetation prevails until 37 ° S with an intermittent presence of  
162 mountain forests in the Patagonian plains (Eva et al., 2004).

163 | [Figure 2](#) summarizes the dominant climatology and associated hydrological regime of  
164 rivers in the study region. The temperature seasonality (upper left) is typical of a

James McPhee 1/1/2016 11:37  
**Eliminado:** Figure 2

166 temperate, Mediterranean climate, and precipitation is strongly concentrated in the fall-  
167 winter months of May through August (upper right). The hydrological regime is  
168 markedly snow-dominated in the northern part of the domain, which can be seen from  
169 the sharp increase in river flow from October and into the summer months of Dec, Jan  
170 and Feb (lower right) that follows the seasonal melt of snow (lower left). Only rivers in  
171 the southern subregion display a significant rainfall-dominated seasonal hydrograph.

172 The importance of SWE for the region is demonstrated by the fact that for the studied  
173 basins, ablation-season (September - March) river flow accounts for two-thirds of  
174 average annual streamflow. Maximum SWE accumulation is reached between the  
175 months of August and September on the western side and between late September and  
176 early October on the eastern side.[\(Figure S4\)](#) Scattered snow showers in mid spring  
177 (September through November) affect the study area, but they do not affect significantly  
178 the decreasing trend of snow-covered area during the melt season (see timing of peak  
179 SWE and fractional Snow Covered Area (fSCA) analysis in online supplementary  
180 material). This feature is essential for choosing the SWE reconstruction methodology  
181 used in this work, which is most applicable to snow regimes with distinct snow  
182 accumulation and snow ablation seasons.

183 By and large, the existing network of high-elevation meteorological stations does not  
184 include appropriately shielded solid precipitation sensors. Some climate reanalysis  
185 products exist, but their representation of Andean topography is crude, and their spatial  
186 resolution is not readily amenable to hydrological applications without significant bias  
187 correction (Krogh et al., 2015; Scheel et al., 2011). Previous attempts at estimating  
188 precipitation amounts at high elevation reaches in the Andes suggest uncertainties on  
189 the order of 50% (Castro et al., 2014; Falvey and Garreaud, 2007; Favier et al., 2009).  
190 In some basins, runoff is partially dictated by glacier contributions, which occur in

James McPhee 1/1/2016 11:37  
Eliminado:

192 summer. According to the Randolph Glacier Inventory (<http://www.glims.org/RGI/>) the  
193 central Andes cordillera has a glacier area of 2,245 km<sup>2</sup> between 27 ° S and 38 ° S,  
194 which is equivalent to 1.5% of the modeling domain surface area (~152,000 km<sup>2</sup>).

195

### 196 3 Methods

#### 197 3.1 SWE reconstruction model

198 A retrospective SWE reconstruction model based on the convolution of the fSCA  
199 depletion curve and time-variant energy inputs for each domain pixel is implemented.  
200 For each year, the model is run at a daily time step between Aug 15 (end of winter) and  
201 Jan 15 (mid-summer). This time window ensures capturing the most likely time at  
202 which peak SWE occurs –which itself is variable from year to year- and the almost  
203 complete depletion of the seasonal snowpack. Isolated pixels with non-negative fSCA  
204 values may remain after Jan 15 at glacier and perennial snowpack sites. However, the  
205 relative area that these pixels represent with respect to the entire model domain is very  
206 low (< 1.5%), and can be neglected in the context of this work.

207 The energy balance model adopted here derives from the formulation proposed by  
208 Brubaker et al. (1996), which considers explicit net shortwave and longwave radiation  
209 terms and a conceptual, pseudo-physically based formulation for turbulent fluxes that  
210 depends only on the degree-day air temperature:

$$M_p = \max\{(Q_{nsw} + Q_{nlw}) f_B + T_d a_r, 0\} \quad [1]$$

211 Were  $M_p$  is potential melt;  $Q_{nsw}$  is the net shortwave energy flux;  $Q_{nlw}$  is the net  
212 longwave energy flux;  $T_d$  is the degree-day temperature,  $a_r$  [mm °C<sup>-1</sup> day<sup>-1</sup>] is the  
213 restricted degree-day factor, and  $f_B$  is the energy-to-mass conversion factor with a value  
214 of 0.26 [mm W<sup>-1</sup> m<sup>2</sup> day<sup>-1</sup>]. Actual melt is obtained by multiplying potential melt by  
215 fractional snow cover area:

$$M = M_p fSCA^{fc} \quad [2]$$

216 where  $fSCA^{fc}$  is the fSCA MOD10A1 estimate adjusted to forest cover correction by a  
 217 vegetation fractional  $f_{veg}$  (0 to 1) from the MOD44B product (Hansen et al., 2003):

$$fSCA^{fc} = \frac{fSCA^{obs}}{(1 - f_{veg})} \quad [3]$$

218 The SWE for each pixel is computed for each year by accumulating the melt fluxes  
 219 back in time during the melt season, starting from the day in which fSCA reaches a  
 220 minimum value, and up to a date such that winter fSCA has plateaued, according to the  
 221 relations:

$$SWE_t = SWE_0 - \sum_1^t M = M_{t+1} + SWE_{t+1} \quad [4]$$

$$SWE_0 = \sum_{t=1}^n M_t ; SWE_n = 0 \quad [5]$$

222 where  $SWE_0$  is end-of-winter or initial maximum SWE accumulation,  $SWE_n$  is a  
 223 minimum or threshold value. The model was run retrospectively until Aug 15, an  
 224 adequate date before which little melt can be expected for most of the winter seasons  
 225 within the modeling period in this region (please see Fig. S5 in the online  
 226 supplementary material).

### 227 3.2 Fractional Snow Covered Area and land use data

228 Spatio-temporal evolution of snow covered area was estimated using the fSCA product  
 229 from the Moderate Resolution Imaging Spectroradiometer (MODIS) on-board the Terra  
 230 satellite (MOD10A1 C5 Level 3). The MOD10A1 product provides daily fSCA  
 231 estimates at 500-m resolution. Percentages of snow extent (i.e. 0% to 100%) are derived  
 232 from an empirical linearization of the Normalized Difference Snow Index (NDSI),  
 233 considering the total MODIS reflectance in the visible range (0.545 - 0.565  $\mu\text{m}$ ; band 4)

234 and shortwave infrared (1.628 - 1.652  $\mu\text{m}$ ; band 6) (Hall et al., 2002; Hall and Riggs,  
235 2007).

236 Binary and fractional MODIS fSCA estimates are limited by the use of an empirical  
237 NDSI-based method. These errors are notoriously sensitive to surface features such as  
238 fractional vegetation and surface temperature (Rittger et al., 2013). Arsenault et al.  
239 (2014) reviewed MODIS fSCA accuracy estimates from several studies under different  
240 climatic conditions, and report a range between 1.5% and 33% in terms of absolute  
241 error with respect to ground observations and operational snow cover datasets. Errors  
242 stem mainly from cloud masking and detection of very thin snow (<10 mm depth),  
243 forest cover and terrain complexity. In general, commission and omission errors are  
244 greatest in the early and late portions of the snow cover season (Hall and Riggs, 2007)  
245 and decrease with increasing elevation (Arsenault et al., 2014). Molotch and Margulis  
246 (2008) compared MODIS and Landsat Enhanced Thematic Mapper performance in the  
247 context of SWE reconstruction, showing that significant differences in SWE estimates  
248 were a result of SCA estimation accuracy and less so of model spatial resolution. The  
249 latter conclusion supports the feasibility of using the snow covered area products at a  
250 500 m spatial resolution for regional scale studies. In order to minimize the effect of  
251 cloud cover on the temporal continuity and extent of the fSCA estimates, the  
252 MOD10A1 fSCA product was post-processed by a modified algorithm for non-binary  
253 products, based on the algorithm proposed by Gafurov and Bárdossy (2009). Their  
254 method is adapted here to the fractional snow cover product, applying a three-step  
255 correction consisting of: (1) a pixel-specific linear temporal interpolation over 1, 2 or 3  
256 days prior and posterior to a cloudy pixel; (2) a spatial interpolation over the eight-pixel  
257 kernel surrounding the cloudy pixel, retaining information from lower-elevation pixels  
258 only; and (3) assigning the 2001-2014 fSCA pixel specific average when steps (1) and

259 (2) where not feasible. This step minimized the effect of cloud cover on data availability  
260 over the spatial domain, yielding cloud cover percentages ranging from 21% in  
261 September to 8% in December.

262 The Normalized Difference Vegetation Index (NDVI) (Huete et al., 2002) derived from  
263 the product MOD13Q1 v5 MODIS Level 3 (16 days - 250 m) is used to classify forest  
264 presence for each model pixel. For pixels classified as forested, both fSCA and energy  
265 fluxes were corrected: fractional SCA was modified on the basis of percentage forest  
266 cover (Molotch, 2009; Rittger et al., 2013), using the average of the forest percentage  
267 product from MOD44B V51. Forest attenuation (below canopy) of energy fluxes at the  
268 snow surface was estimated from forest cover following the method from Ahl et al.  
269 (2006) assuming invariant LAI over each melt season. The selected LAI pattern is  
270 obtained by averaging the four LAI scenes available between December - January time  
271 window through 14 study years. This time window displays the average state of  
272 evergreen forest with the maximum amount of data.

273

### 274 **3.3 Model forcings**

275 Spatially distributed forcings are required at each grid element in order to run the SWE  
276 reconstruction model. In order to ensure the tractability of the extrapolation process, we  
277 divided the model domain into sub-regions or clusters, composed by one or more river  
278 basins. The river basins were grouped using a clusterization algorithm (please see  
279 section S2 on the online supplementary material) based on melt season river flow  
280 volume as described in Rubio-Álvarez and McPhee (2010). Then, spatially distributed  
281 variables (surface temperature, fSCA, global irradiance) are combined with  
282 homogeneous variables for each cluster (e.g. cloud cover index) and point data from  
283 meteorological stations in order to obtain a distributed product as described below. A

284 further benefit of the clustering process is that it allows us to analyze distinct regional  
285 features of the SWE reconstruction parameters, input variables and output estimates.

286 Net shortwave radiation,  $Q_{nsw}$  is estimated as a function of incoming solar radiation  
287 based on the equation:

$$Q_{nsw} = (1 - \alpha_s)(G_{\downarrow})\tau_a \quad [6]$$

288 where  $\alpha_s$  is snow surface broadband albedo;  $G_{\downarrow}$  is incoming solar radiation (global  
289 irradiance);  $\tau_a$  is the shortwave transmissivity as a function of LAI for mixed forest  
290 cover (Pontailler et al., 2003; Sicart et al., 2004), which in turn is estimated as:

$$\tau_a = e^{(-\kappa LAI)} ; \quad LAI = -1.323 \ln \left( \frac{0.88 - NDVI}{0.72} \right) \quad [7]$$

291 with  $\kappa = 0.52$  for mixed forest species (Dewalle and Rango, 2008). Equation 7 is valid  
292 for NDVI values between 0.16 and 0.87. Global irradiance under cloudy sky conditions  
293 is estimated considering a daily distributed spatial pattern of clear sky irradiance  
294  $G_{c\downarrow}$  derived by the *r.sun* GRASS GIS module (Hofierka et al., 2002; Neteler et al., 2012)  
295 and the clear sky index  $K_c$  derived from the insolation incident on a horizontal surface  
296 from the "Climatology Resource for Agroclimatology" project in the NASA Prediction  
297 Worldwide Energy Resource "POWER" (<http://power.larc.nasa.gov/>) 1°x1° gridded  
298 product.

$$G_{\downarrow} = K_c G_{c\downarrow} ; \quad K_c = (\overline{G_{r\downarrow}} / \overline{G_{c\downarrow}}) \quad [8]$$

299 In Equation 8,  $\overline{G_{r\downarrow}}$  and  $\overline{G_{c\downarrow}}$  are spatial averages over each hydrologic response unit  
300 (cluster) of the POWER and *r.sun*-derived products, respectively.

301 A snow-age decay function based on snowfall detection is implemented to estimate  
302 daily snow surface albedo (Molotch and Bales, 2006) constrained between values of  
303 0.85 and 0.40 (Army Corps of Engineers - USACE, 1960). Snowfall events were

304 diagnosed using a unique minimum threshold for fSCA increments of 2.5% for each  
305 hydrologic unit area.

306 Net long wave radiation estimates are derived using:

$$Q_{nlw} = L_{\downarrow} f_{sv} \varepsilon_s + \sigma T_a^4 (1 - f_{sv}) \varepsilon_{sf} - \sigma T_s^4 \varepsilon_s \quad [9]$$

$$L_{\downarrow} = 0.575 e_a^{1/7} \sigma T_a^4 (1 + a_c C^2) \quad [10]$$

307 Where  $T_a$  is air temperature,  $T_s$  is the snow surface temperature,  $\varepsilon_s$  is the snow  
308 emissivity (i.e. 0.97),  $\varepsilon_{sf}$  is the canopy emissivity (i.e. 0.97),  $f_{sv}$  is the sky-view factor  
309 (i.e. assumed equal to shortwave transmissivity; Pomeroy et al., 2009; Sicart et al.,  
310 2004),  $\sigma$  is the Stefan-Boltzmann constant, and  $L_{\downarrow}$  is the incoming long wave radiation.

311 Air vapor pressure ( $e_a$ ) required for long wave radiation estimates was derived from air  
312 temperature and relative humidity, which in turn was assumed constant throughout the  
313 melt period and equal to 40% based on observations at selected high-elevation  
314 meteorological stations. The multiplying factor  $(1 + a_c C^2)$  represents an increase in  
315 energy input relative to clear sky conditions due to cloud cover, where  $a_c$  equals 0.17  
316 and  $C = 1 - K_c$  is an estimate of the cloud cover fraction (DeWalle and Rango, 2008).

317 Spatially distributed air temperature is generated by combining daily air temperature  
318 recorded at index meteorological stations and a weekly spatial pattern of skin  
319 temperature derived from the MODIS Land Surface Temperature product  
320 (MOD11A1.V5) (Wan et al., 2004; Wan et al., 2002). The product MOD11A1 V5  
321 Level 3 estimates surface temperature from thermal infrared brightness temperatures  
322 under clear sky conditions using daytime and nighttime scenes and has been shown to  
323 adequately represent measurements at meteorological stations ( $R^2 \geq 0.7$ ), displaying  
324 moderate overestimation in spring and underestimation in fall (Neteler, 2010). Other  
325 studies have reported similar accuracies, with RSME values around 4.5 °K in cold

James McPhee 1/1/2016 11:37  
Eliminado:

327 mountain environments (Williamson et al., 2014). Taking into account the high  
328 correlation between air temperature and LST (Benali et al., 2012; Colombi et al., 2007;  
329 Williamson et al., 2014), we define:

$$T_a = T_{a\ base} + \Delta T_a = T_{a\ base} + \mu (LST - LST_{base}) + v \quad [11]$$

330 were  $T_{a\ base}$  is daily air temperature at an index station for each cluster and  $\Delta T_a$  is the  
331 difference in air temperature between any pixel and the pixel where the index station is  
332 located. To determine  $\Delta T_a$  we use a linear regression between MODIS LST data and  
333  $\Delta T_a$  considering pairs of stations located at high altitude and valley (base) sites, taking  
334 into account the melt season average values over the 2001-2014 period. In equation 11,  
335  $LST - LST_{base}$  denotes the difference between skin temperatures from any pixel and the  
336 index station pixel. The linear regression between skin temperature and air temperature  
337 differences has a slope  $\mu$  of 0.65, an intersect  $v$  of -0.5 and  $R^2$  of 0.93 (Figure S3 in  
338 online supplementary material). Estimation of LST during cloudy conditions is done as  
339 follows: (1) a pixel-specific linear temporal interpolation is performed over 1 and 2 days  
340 prior and posterior to the cloudy pixel; and (2) estimation of remaining null values by  
341 an LST-elevation linear regression (Rhee and Im, 2014).

342 This spatial extrapolation method was preferred over more traditional methods -for  
343 example based on vertical lapse-rates (Minder et al., 2010; Molotch and Margulis,  
344 2008)- after initial tests showed that the combined effect of the relatively low elevation  
345 of index stations and the large vertical range of the study domain resulted in  
346 unreasonably low air temperatures at pixels with the highest elevations. Likewise, the  
347 scarcity of high-elevation meteorological stations and the large spatial extent of the  
348 model domain precluded us from adopting more sophisticated temperature estimation  
349 methods (e.g. Ragettli et al., 2014).

350 Snow surface temperature and degree-day temperature are estimated (Brubaker et al.,  
351 1996) as:

$$T_d = \max(T_a, 0) ; T_s = \min(T_a - \Delta_T, 0) \quad [12]$$

352 where  $\Delta_T$  is the difference between air and snow surface temperature. To the best of our  
353 knowledge, no direct, systematic values of snow surface temperature exist in this  
354 region, so for the purposes of this paper we adopt an average value  $\Delta_T = 2.5$  [°C],  
355 following the suggestion in Brubaker et al., (1996). Slightly higher values ranging from  
356 3 to 6°C are shown for continental and alpine snow types (Raleigh et al., 2013)  
357 indicating an additional source of uncertainty over net long wave radiation  
358 computations. More sophisticated parametrizations for  $T_s$ , for example based on heat  
359 flow through the snowpack, have been proposed (e.g. Rankinen et al., 2004; Tarboton  
360 and Luce, 1996) but those require explicit knowledge about the snowpack temperature  
361 profile and/or more complex model formulations to estimate the internal snowpack heat  
362 and mass budgets simultaneously.

363 The  $a_r$  coefficient in the restricted degree-day energy balance equation was computed  
364 using a combination of station and reanalysis data, and assumed spatially homogeneous  
365 within each of the clusters that subdivide the model domain. Brubaker et al. (1996)  
366 propose a scheme in which this parameter can be explicitly computed from air and snow  
367 surface temperature, air relative humidity, and atmospheric pressure and wind speed.  
368 Wind speed was obtained from the NASA POWER reanalysis described previously. A  
369 correction for atmospheric stability is applied on the bulk transfer coefficient  $C_h$   
370 according to the formulation presented by Kustas et al. (1994), assuming a surface  
371 roughness of 0.0005 m:

$$C_h = \begin{cases} (1 - 58R_i)^{0.25} \text{ for } R_i < 0 \\ (1 + 7R_i)^{-0.1} \text{ for } R_i > 0 \end{cases} ; \quad R_i = \frac{gz(T_a - T_s)}{u^2 T_a} \quad [13]$$

372 Where  $R_i$  is the Richardson number,  $g$  is the gravity acceleration (9.8 [m s<sup>-2</sup>]),  $z$  is the  
 373 standard air temperature measurement height (2 m) and  $u$  is wind speed. The  
 374 calculation of  $R_i$  and  $a_r$  is based on the standard assumptions of  $T_s$  at the freezing point  
 375 and a water vapor saturated snow surface over all high-elevation meteorological stations  
 376 with available air temperature and relative humidity records (Molotch and Margulis,  
 377 2008). Further in the text, we discuss some implications of these assumptions and of the  
 378 input data used on the ability of the model of simulating relevant components of the  
 379 snowpack energy exchange.

James McPhee 1/1/2016 11:37

Eliminado: ... [1]

380 [Table 1](#) shows the main cluster characteristics and regionalized model parameters. It can  
 381 be seen that for those clusters located in the southern and middle reaches of the model  
 382 domain, the  $a_r$  parameter values range from 0.10 to 0.23 [cm °C<sup>-1</sup> day<sup>-1</sup>], which is  
 383 similar to values reported in previous studies performed in other mountain ranges in the  
 384 Northern Hemisphere (0.20 – 0.25 in Martinec (1989), 0.17 in Kustas et al., (1994),  
 385 0.20 in Brubaker et al., (1996), 0.15 in Molotch and Margulis (2008)). However, values  
 386 associated to the northernmost clusters of our study area are quite low, reaching under  
 387 0.02 for the C1 cluster in northern Chile.

388 Clear sky index ( $K_c$ ) values range between 0.78 and 0.89 which is similar to values  
 389 reported by Salazar and Raaijijk (2014) who estimate  $K_c$  values on the order of 0.90 for  
 390 a single location at 1200 m.a.s.l. in northern Argentina. A 5 to 6 °C difference can be  
 391 observed in mean air temperature at index stations between the northern and southern  
 392 edge of the domain. Temperatures for the C4 cluster are subject to greater uncertainty,

393 because no high-elevation climate station data was available for this study ([Figure S4](#)).  
 394 Forest cover values are lower than 6% throughout the model domain, with the exception

James McPhee 1/1/2016 11:37

Eliminado: .

398 of cluster C3, with a value of 13.8%. The difference in forest cover between clusters C3  
399 and C8 can be attributed to the precipitation shadow effect induced by the Andes ridge.  
400 Forest corrections applied to MODIS fSCA resulted in a 17% increase with respect to  
401 the original values over the southern sub-domain (C3).

402 **3.4 Evaluation data: SWE, snow depth and river flow observations**

403 Operational daily snow-pillow data from stations maintained by government agencies in  
404 Chile and Argentina were available for this study (Table 2). Only stations with ten or  
405 more years of record were included and manual snow course data were neglected  
406 because of their discontinuous nature. Approximately 10% of observed maximum SWE  
407 accumulation values were discarded due to obvious measurement errors and data gaps.  
408 An analysis of the seasonal variability of snow-pillow records on the western and  
409 eastern slopes of the Andes suggests that peak-SWE date is somewhat delayed on the  
410 latter, by approximately one month. Therefore, peak-SWE estimates for Chilean and  
411 Argentinean stations are evaluated on September 1 and October 1, respectively,  
412 although in the results section we show values for September 15 in order to use a unique  
413 date for the entire domain. Manual snow depth observations were taken in the vicinity  
414 of selected snow-pillow locations in order to evaluate the representativeness of these  
415 measurements at the MODIS grid scale during the peak-SWE time window. These  
416 depth observations were obtained in regular grid patterns within an area the  
417 approximate size of a MODIS pixel (500 m), centered about the snow-pillow location.  
418 On average, 120 depth observations spaced at approximately 50-m increments were  
419 obtained at each snow pillow site. Snow density was estimated by a depth-weighted  
420 average of snow densities measured in snow pits with a 1000-cc snow cutter. Samples  
421 where obtained either at regular 10-cm depth intervals along the snow pit face, or at the  
422 approximate mid depth of identifiable snow strata for very shallow snow pack

423 conditions. Weights were computed as the fraction of total depth represented by each  
424 snow sample.

425 Distributed snow depth observations were available from snow surveys carried out  
426 during late winter between 2010 and 2014 at seven study catchments in the western side  
427 of the Andes, between latitudes 30 ° S and 37 ° S ([Figure1](#), [Table 3](#)). Snow depths were  
428 recorded with 3 m graduated avalanche probes inserted vertically into the snow pack.

429 Depending on the terrain conditions, between three and five individual point snow depth  
430 measurements were obtained at each location, from which a mean snow depth and  
431 standard error are calculated; i.e. three-point observations are made forming a line with  
432 a spacing of one meter and five-point observations are made forming a cross with an  
433 angle of 90 degrees and a spacing of one meter. Pixel-scale SWE estimates are obtained  
434 by averaging all depth observations within the limits of MODIS pixels and multiplying  
435 them by density observations from snow pits excavated at the time of each snow survey;  
436 i.e. two or three snow pits per field campaign. After this, individual depth observations  
437 are converted into SWE for model validation. Modeled SWE values are averaged at all  
438 MODIS pixels where manual depth observations are available, and their summary  
439 statistics are compared to those of SWE estimated from manual depth observations at  
440 the same pixels, multiplied by average density from snow pits.

441 Spring and summer season (September to March) total river flow volume (SSRV) for  
442 the 2001-2014 period are obtained from unimpaired ([no human extractions](#)) streamflow  
443 records at river gauges located in the mountain front along the model domain. Data  
444 were pre-selected leaving out series that showed too many missing values, and verified  
445 through the double mass curve method (Searcy and Hardison, 1960) in order to discard  
446 anomalous values and to ensure homogeneity throughout the period of study. Regional  
447 consistency was verified through regression analysis, only including streamflow records

James McPhee 1/1/2016 11:37

**Eliminado:** ([Figure 1](#), [Table 3](#)).

449 with  $R^2$  values greater than 0.5 among neighboring catchments. Missing values  
450 constituted about 3.7% of the entire period and were filled through linear regression.

451

## 452 4 Results

### 453 4.1 Model validation

454 | [Figure 3](#) compares reconstructed peak SWE (gray circles) to observed values at three  
455 snow-pillow locations (black diamonds) where additional validation sampling at the  
456 MODIS pixel scale was conducted (box plots). At the Cerro Vega Negra site (CVN),  
457 located in cluster C1, the model overestimates peak SWE (September 1) with respect to  
458 the snow-pillow value by 97% in 2013 and by 198% in 2014. At the Portillo site (POR,  
459 cluster C2), reconstructed SWE underestimates recorded values by 51% in 2013 and  
460 72% 2014. At the Laguna Negra site (LAG, also C2), reconstructed peak SWE slightly  
461 overestimates recorded values (8%) (Table 4). However, reconstructed SWE compares  
462 favorably to distributed manual SWE observations obtained in the vicinity of the snow  
463 pillows at the POR and LAG sites. [At POR, model estimates approach upper \(2012\) and](#)  
464 [lower \(2013 and 2104\) quartiles, while at LAG the model estimates are closer to the](#)  
465 [minimum value observed in 2013 and very similar to the observed mean in 2014.](#)

James McPhee 1/1/2016 11:37

Eliminado: Figure 3

466 | [Figure 4](#) depicts the comparison between reconstructed SWE and snow surveys carried  
467 out at pilot basins throughout the model domain. From left to right, it can be seen that  
468 the model slightly overestimates SWE with respect to observations at CVN (i.e. 18%  
469 overestimation). Further south, there is a very good agreement at ODA - MAR (i.e. 4%  
470 underestimation), with less favorable results at MOR - LVD (i.e. 39% underestimation)  
471 and OB - RBL (i.e. 36% underestimation). At CHI the model significantly  
472 underestimates SWE (i.e. by 67%); note this site is heavily forested. For the 2013a and  
473 2014a boxes (Figure 4) –which correspond to clearing sites-, there is still

James McPhee 1/1/2016 11:37

Eliminado: Figure 4

476 underestimation, but of lesser magnitude (20%). Summarizing, we detect model  
477 overestimation respect to snow survey medians in four cases and underestimation in  
478 fifteen cases. In 11 out of 19 cases, reconstructed SWE lies within the snow survey data  
479 uncertainty bounds (standard deviation).

480 Figure 5 shows a comparison between model estimates of peak (Sep 15<sup>th</sup>) SWE and  
481 corresponding observations at snow pillow sites. In general, directly contrasting pixel-  
482 based estimates with sensor observations should be attempted with caution. In areas  
483 with complex topography, slight variations in the position of the sensor with respect to  
484 the model grid, combined with high spatial variability in snow accumulation could lead  
485 to large differences between model estimates and observations. Also, small-scale  
486 variations in snow accumulation near the sensor, for example induced by protective  
487 fences, could introduce bias to the results (e.g. Meromy et al., 2013; Molotch and Bales,  
488 2006; Rice and Bales, 2010). Taking the above into consideration, [Figure 5](#) suggests  
489 that the model tends to overestimate observed peak SWE at the two northernmost sites  
490 on the Chilean side (QUE and CVN); the equivalent cluster on the Argentinean side  
491 (C4) lacks SWE observations. [The R<sup>2</sup> values indicated below refer to the best linear fit;](#)  
492 [regression line slope and intercept coefficients are provided in Table 4.](#) Overall, we find  
493 a better agreement at the eastern slope sites (i.e. R<sup>2</sup> = 0.74) than at their western  
494 counterparts (i.e. R<sup>2</sup> = 0.43), with a combined R<sup>2</sup> value of 0.61. Individually, the worst  
495 and best linear agreements are obtained at POR (R<sup>2</sup>=0.32) and LOA (R<sup>2</sup>=0.88),  
496 respectively. Time series of observed SWE and model estimates for these two extreme  
497 cases are shown in the supplementary online material, and indicate a significant degree  
498 of inter-annual variability in model discrepancies in terms of peak SWE, but less in  
499 terms of, for instance, snow cover duration. Average standard error, SE<sub>x̄</sub> is 284 mm  
500 (SE<sub>x̄</sub> =242 mm at the west slope; SE<sub>x̄</sub>=302 mm at the east slope), with a range between

James McPhee 1/1/2016 11:37  
**Eliminado:** Figure 5

502 72 mm (TOS) and 378 mm (ATU) (Table 4). Relative errors display some variability,  
503 with overestimation higher than 30% at the two northernmost (QUE and CVN) and at  
504 the southernmost (PEH) snow pillows. For all other snow pillows, the model estimates  
505 are lower than the sensor observation; the range of relative errors for those sites with  
506 underestimation goes from -52% to -5%.

507 **4.2 Correlation with melt-season river flows**

508 Under the assumption of unimpaired flows, peak SWE and seasonal flow volume  
509 should show some degree of correlation, even though no assumptions can be made here  
510 about other relevant hydrologic processes, such as flow contributions from glaciated  
511 areas, subsurface storage carryover at the basin scale and influence of spring and  
512 summer precipitation. Differences can be expected due to losses to evapotranspiration  
513 and sublimation affecting the snowpack and soil water throughout the melt season.  
514 Hence, basin-averaged peak SWE should always be higher than melt season river  
515 volume. A clear regional pattern emerges when inspecting the results of this comparison  
516 in [Figure 6](#). Correlation between peak SWE and melt season river flow is higher in  
517 clusters C1 and C4 with  $R^2$  values of 0.84 and 0.86, respectively. The result for Cluster  
518 C4 indicates that liquid precipitation during the melt season (Figure 2) does not result in  
519 decreased correlation between peak SWE and river flow. Clusters C2, C5, C6 and C7  
520 display a somewhat lower correlation, with some individual years departing more  
521 significantly from the overall linear trend.  $R^2$  values range between 0.46 and 0.78 in  
522 these cases. Finally, not only are correlation coefficients lower for the southern clusters  
523 C3 ( $R^2 = 0.56$ ) and C8 ( $R^2 = 0.48$ ), but also estimated peak SWE is always lower than  
524 river flow, which indicates the importance of spring and summer precipitation in  
525 determining streamflow variability. In fact, Castro et al. (2014) analyze patterns of daily  
526 precipitation in this area and document average spring and summer rainfall amounts of

James McPhee 1/1/2016 11:37

Eliminado: (no human extractions).

James McPhee 1/1/2016 11:37

Eliminado: Figure 6.

529 approximately 520 mm in C3 and 85 mm in C8. A promising avenue for further  
530 research in this region emerges when comparing the correlation between melt-season  
531 river flow and the spatially distributed reconstructed product versus that of river flows  
532 and snow pillow data. Table 5 shows values of  $R^2$  for the linear regression between  
533 these variables. It can be seen that for two of the three clusters on the western side of the  
534 continental divide, the end-of-winter distributed reconstruction has more predictive  
535 power than observed SWE. Only for central Chile the *Laguna Negra* (LAG, with a  
536 value of 0.82) site has a better correlation with river flows, but the reconstructed  
537 product has a value of 0.78, which lies in between those found for LAG and for *Portillo*  
538 (POR, with a value of 0.68). For the eastern side of the continental divide, the  
539 distributed product shows similar skill than that of snow pillows except for Atuel, which  
540 has a very high correlation ( $R^2$  of 0.87) with cluster C6 river flows, and for cluster C7,  
541 in which the reconstruction shows higher predictive power ( $R^2$  of 0.89) than the  
542 available SWE observations (VAL and PEH).

543

#### 544 4.3 Regional SWE estimates

545 | [Figure 7](#) shows the Sep 15 SWE average over the 2001 – 2014 period obtained from the  
546 reconstruction model, and the percent annual deviations (anomalies) from that average.  
547 Steep elevation gradients can be inferred from the climatology, as well as the latitudinal  
548 variation expected from precipitation spatial patterns. For the northern clusters (C1 and  
549 C4), the peak SWE averaged over snow covered areas is on the order of 300 mm while  
550 in the middle of the domain (C2, C5, C6), it averages approximately 750 mm. The  
551 southern clusters (C3, C7, C8) do show high accumulation averages (approximately 650  
552 mm), despite the sharp decrease in the Andes elevation south of latitude 34 ° S. The  
553 anomaly maps convey the important degree of inter-annual variability, as well as

James McPhee 1/1/2016 11:37  
Eliminado: Figure 7

555 distinct spatial patterns associated with it. Between 2001 and 2014, years 2002 and  
556 2005 stand out for displaying large positive anomalies throughout the entire  
557 mountainous region of the model domain, with values 2000 mm and more above the  
558 simulation period average. Other years prior to 2010 show differential accumulation  
559 patterns, where either the northern or southern parts of the domain are more strongly  
560 affected by positive or negative anomalies. Overall, the northern clusters (C1 and C4)  
561 show above-average accumulation in only three (2002, 2005 and 2007) of the 14  
562 simulated years, whereas the other clusters show above-average accumulation for six  
563 years (2001, 2002, 2005, 2006, 2008 and 2009). In particular, years 2007 and 2009  
564 show a bimodal spatial structure, with excess accumulation (deficit) in the northern  
565 (southern) clusters during the former, and the inverse pattern in the latter year.

566 A longitudinal pattern in the distribution of negative anomalies can be discerned from  
567 Figure 7, whereby drought conditions tend to be more acute on one side of the divide  
568 versus the other. Conversely, during positive anomaly years, both sides of the Andes  
569 seem to show similar behavior. Further research on the mechanisms of moisture  
570 transport during below-average precipitation years may shed light on this result.

571 | [Figure 8](#) provides a different perspective on the region's peak SWE climatology by  
572 presenting our results aggregated into elevation bands for each hydrologic unit.  
573 Elevation bands are defined at 1000-m increments starting from 1000 m.a.s.l. Crosses  
574 indicate average peak SWE for each band (mm), and circle areas are proportional to the  
575 surface area covered by each elevation band. From north to south, hydrologic unit C4  
576 shows slightly higher SWE than C1 between 3000 and 5000 m.a.s.l., but much larger  
577 surface areas (~32,000 vs. ~17,000 km<sup>2</sup>), indicating a larger water resource potential.  
578 C2 stands out as having the greatest area-weighted cluster SWE and the greatest SWE  
579 for each elevation band. Compared to its counterpart on the eastern side of the Andes

James McPhee 1/1/2016 11:37  
**Eliminado:** Figure 8

range (C5), C2 shows higher accumulations (up to ~1800 mm) at all elevations. The area included between 2000 and 4000 m.a.s.l. (~13,000 km<sup>2</sup>), which shows an estimated peak SWE accumulation on the order of 600 mm, represents the most predominant snow volume accumulation zone. Although the 4000-5000 m.a.s.l. elevation band contributes approximately half the 2000 - 4000 band surface area in C2, its average peak SWE is roughly twice that of the 3000-4000 band (~6,000 km<sup>2</sup>). This makes this subregion interesting for future research, because most snow observations in the area are obtained below 4000 m.a.s.l.; the same is true for unit C5. Further to the south, the barrier effect of the Andes is also suggested by the displacement of the SWE-elevation distribution in C6 and C7 when compared to C3. On the eastern side of the model domain, it is interesting to see a steepening of the average peak SWE elevation profile between C6 and C8, suggesting that C8 is less affected by Andes blockage than its northern counterparts.

Estimated net energy inputs ([Figure 9](#)) shows a decrease from the northern (C1 and C4) into the mid-range clusters (C2, C5 and C6), with increases again in the southern reaches of the domain (C3, C7 and C8). This is a result of a combination of an increasing trend in net shortwave radiation in the south-north direction and a reverse spatial trend in net long wave radiation exchange, which increases (approaches less-negative values) in the north-south direction. Modeled turbulent energy fluxes (Equation 1) are negligible in the northern clusters, but their contribution to the net energy exchange increases with latitude as a result of the spatial variation in the  $a_r$  parameter.

[Figure 10](#) shows the temporal (seasonal) variation in average fSCA and SWE for each cluster, and [Table 6](#) shows peak SWE at the watershed scale, averaged both over the entire basin and over the snow covered area. Maximum fSCA increases in the north-

James McPhee 1/1/2016 11:37

**Eliminado:** (Figure 9)

James McPhee 1/1/2016 11:37

**Eliminado:** ... [2]

James McPhee 1/1/2016 11:37

**Con formato:** EspacioPosterior: 10 pto

James McPhee 1/1/2016 11:37

**Eliminado:** Table 5

610 south direction, consistently with the climatological increase in winter precipitation and  
611 decrease in temperature. A dramatic increase in snow coverage is observed between the  
612 northern (i.e. C1 and C4) and adjacent southern clusters (i.e. C2 and C5), with average  
613 peak fSCA increasing from 20% to 50%. The highest average snow coverage is  
614 observed for cluster C8, with more than 60%. Snow water equivalent displays a similar  
615 regional variability with lower seasonal variability than snow cover for all clusters  
616 except for C2, where fSCA and SWE variability throughout the melt season are  
617 identical. Mean peak SWE in northern Chile is the lowest among the eight clusters, with  
618 approximately 100 mm SWE over the 2001-2014 period. The largest estimate is for  
619 cluster C2, central Chile, where mean peak SWE exceeds 500 mm. The rain shadow  
620 effect of the Andes range is apparent in the comparison of SWE and fSCA in C2 and  
621 C5-C6-C7. Fractional snow covered area is lower on the east side because of the larger  
622 basin sizes which increases the proportional area of lower elevation terrain. In addition,  
623 peak SWE is approximately 25% lower on the east side, with less than 400 mm SWE  
624 for the eastern clusters. Cluster C4 is not affected by this phenomenon, showing higher  
625 snow coverage and water equivalent accumulation than its counterpart, C1. Cluster C8  
626 represents an interesting exception in that its average fSCA is the largest within the  
627 model domain, but peak SWE is not significantly higher than the estimates in the other  
628 clusters on the Argentinean side of the Andes.

629

## 630 **5 Discussion**

### 631 **5.1 Sensitivity analysis**

632 The Andes cordillera, on one hand, displays ideal conditions for SWE reconstruction,  
633 including low cloud cover, infrequent snowfall during spring and summer, and very low  
634 forest cover. On the other hand, the scarcity of basic climate data poses challenges that

would affect any modeling exercise. A local sensitivity analysis is implemented in order to gain insights regarding the influence of some of the assumptions required for SWE modeling ([Figure 11](#)). The influence of the clear sky factor ( $K_c$ ), snow surface albedo ( $\alpha_s$ ), the slope of the  $\Delta_{LST}$  vs.  $\Delta_{Ta}$  relationship ( $\mu$ ), the  $a_r$  parameter, and the difference between air and snow surface temperature are explored. Results are shown for the model pixels corresponding to two of the snow pillow sites, each located at the northern and southern sub-regions of the model domain respectively. The clear sky factor, snow albedo and  $\Delta_{LST}$  vs.  $\Delta_{Ta}$  slope are the most sensitive parameters at the northern (CVN) site. Increasing the slope in the  $\Delta_{LST}$  vs.  $\Delta_{Ta}$  relationship results in decreasing temperature at pixels with higher elevations than the index station, thus lowering long wave cooling and resulting in higher SWE estimates. The impact of increasing slope values decreases progressively, because an increasing slope results in increased pixel air temperature, but snow surface temperature cannot exceed 0° C. The influence of snow albedo is analyzed by perturbing the entire albedo time series for each season from the values predicated by the USACE model. Increasing albedo values restricts the energy available for melt therefore decreasing peak SWE estimates. Again, a nonlinear effect is observed, constrained by a minimum albedo value of 0.4. The sensitivity of the clear sky factor, on the other hand, is monotonic, with increasing values generating more available solar energy, resulting in higher SWE estimates. At the southern site (ALT), the shape of the sensitivity functions is the same as at CVN, but the magnitude of SWE variations as a function of parameter perturbations is smaller. This is likely related to the fact that turbulent fluxes constitute a larger fraction of the simulated overall energy balance at the southern sites;  $a_r$  parameter values are greater in the southern portions of the domain. Therefore, perturbations of the other terms account for a smaller fraction of the energy exchange at the southern sites.

James McPhee 1/1/2016 11:37

**Eliminado:** ([Figure 11](#)).

661    **5.2 Model performance and conceptual energy balance representation**

662    Among the many factors that influence model performance, the sub-region delineation  
663    involves the selection of index meteorological stations for extrapolating input data at the  
664    domain level. Thus, for example, two adjacent pixels that are part of different sub-  
665    regions may be assigned input data derived from two different meteorological stations  
666    that are many kilometers apart. It would be preferable to use distributed inputs only, but  
667    these were not available for this domain. Future research is needed to explore  
668    alternative strategies for domain clustering.

669    Overall, the model performance, evaluated against SWE observations, is comparable to  
670    that achieved in other mountain regions of the world. Our average coefficient of  
671    determination  $R^2$  of 0.68 is lower than that obtained by Guan et al. (2013) in the Sierra  
672    Nevada (0.74) when comparing operational snow pillow observations, although this  
673    value is affected by three stations with much lower agreement (POR, LAG, ATU); the  
674    median  $R^2$  in our study, on the other hand, is 0.73, which we consider satisfactory in  
675    light of the scarcity of forcing data and direct snow properties observations available in  
676    this region. The overall relative error is -2% for observations from snow pillows within  
677    our study region, but this value is strongly affected by two stations where we observed  
678    significant overestimation (QUE and CVN). When including the remaining ten snow  
679    pillows only, relative error increases to -16%. Given that forest cover is minimal in our  
680    modeling domain, we can attribute this bias to either weaknesses in the simplified  
681    energy balance model formulation or to errors in the MOD10A1 fSCA product.  
682    Previous work in the northern hemisphere (Rittger et al., 2013) has shown that MODIS  
683    can underestimate fractional snow cover during the snowmelt season. On one hand, land  
684    cover heterogeneity at spatial resolutions lower than the MODIS scale (i.e. 500 m)  
685    result in mixed-pixel detection problems. On the other hand, spectral unmixing based on

686 the NDSI approach tends to underestimate fSCA under patchy snow distributions. In  
687 addition, surface temperatures greater than 10 °C –more likely to exist during late  
688 spring- induce MODIS fSCA underestimation. Molotch and Margulis (2008) tested the  
689 SWE reconstruction model using Landsat ETM and MOD10A1 and found that  
690 maximum basin-wide mean SWE estimates were significantly lower when using  
691 MOD10A1. More recently, Cortés et al. (2014) showed that a similar pattern can be  
692 seen for the extratropical Andes, whereby MODIS fSCA consistently underestimated  
693 LANDSAT TM fSCA retrievals. MODIS fSCA underestimation during spring  
694 combined with increased net energy fluxes over the snowpack can result in a marked  
695 underestimation (~20%) for available energy flux for snowpack melting and  
696 consequently (~45%) for maximum SWE (Molotch and Margulis, 2008).

697 Comparisons against spatial interpolations from intensive-study areas in the Sierra  
698 Nevada or Rocky Mountains (e.g. Erxleben et al., 2002; Jepsen et al., 2012) are not  
699 directly applicable, because in this study we do not employ interpolation methods to  
700 derive our manual snow survey SWE estimates. However, the average overestimation  
701 found with respect to snow survey data could be explained by the fact that manual  
702 surveys are limited by site accessibility and sampling procedures. For example, snow  
703 probes utilized are only 3.0 m long, which precludes observation of deeper snowpack;  
704 likewise, deep snow is expected in sites exposed to avalanching, which were generally  
705 avoided in snow survey design due to safety considerations. On the other hand, manual  
706 snow surveys do not visit steep snow-free areas where snow depth is expected to be  
707 lower than the 500-m pixel reconstruction. The combined effect of these two contrasting  
708 effects is the subject of further research in this region.

709 Another possible explanation for model errors is the simplified formulation of the  
710 | energy balance equation<sup>2</sup>, which may be problematic when applied over a large,

711 climatically variable model domain. To explore the implications of the simplified  
712 energy balance with respect to model errors, we focus on the representation of turbulent  
713 energy fluxes, represented here through a linear temperature-dependent term. [Figure 12](#)  
714 describes the spatial distribution of the  $a_r$  parameter, and its dependence on air  
715 temperature and relative humidity observed at index meteorological stations. The  
716 implication for energy balance modeling is that turbulent fluxes would account for a  
717 very small portion of the snowpack energy and mass balance in the northern area (C1  
718 and C4), which is characterized by low air temperatures and relative humidity, which  
719 yield very low  $a_r$  values. The reader must recall that  $a_r$  values were computed based on  
720 index station data and assumed spatially homogeneous over each cluster. The simplified  
721 model formulation used in this research, however, although pseudo-physically based -  
722 compared to degree-day or fully calibrated models- allows only for positive net  
723 turbulent fluxes, because both the  $a_r$  and the degree-day temperature index are positive  
724 values. However, previous studies in this region (Corripio and Purves, 2005; Favier et  
725 al., 2009) have suggested that latent heat fluxes have a relevant role because of high  
726 sublimation rates favored by high winds and low relative humidity conditions  
727 predominant in the area.

728 In order to diagnose differential performance of the model across the hydrologic units  
729 defined in this study, we compute the Bowen ratio ( $\beta$ ) at the point scale from data  
730 available only at the few high elevation meteorological stations in the region with  
731 recorded relative humidity. The calculations show that at stations located within cluster  
732 C1, latent heat fluxes are opposite in sign and larger in magnitude than sensible heat  
733 fluxes (Figure S6 in supplementary material). While this results in net turbulent cooling  
734 of the snowpack, this energy loss is not considered in our simplified energy balance  
735 approach. Note that for the clusters C5, C6, C7 and C8, all located on the eastern

James McPhee 1/1/2016 11:37

Con formato: español

James McPhee 1/1/2016 11:37

Eliminado: Figure 12

James McPhee 1/1/2016 11:37

Con formato: español

737 (Argentinean) slope of the Andes, sensible and latent heat fluxes are positive, compared  
738 to negative latent heat fluxes for all the index stations within clusters C2 and C3 on the  
739 Chilean side. This result is consistent with Insel et al. (2010), who applied a Regional  
740 Circulation Model (RegCM3) in the area and showed a significant difference in relative  
741 humidity (~70% east side vs. ~40% west side). The fact that we extrapolate the  $a_r$   
742 parameter value based on relatively low elevation meteorological observations  
743 throughout the southern Argentinean hydrologic units may result in a yet not quantified  
744 overestimation of seasonal energy inputs and peak SWE for those clusters.

745

## 746 **6 Conclusions**

747 Snow water equivalent is the foremost water source for the extratropical Andes region  
748 in South America. This paper presents the first high-resolution distributed assessment of  
749 this critical resource, combining instrumental records with remotely sensed snow  
750 covered area and a physically-based snow energy balance model. Overall errors in  
751 estimated peak SWE, when compared with operational station data, amount to -2.2%,  
752 and correlation with observed melt-season river flows is high, with ~~an R<sup>2</sup>~~ value of 0.80.  
753 MODIS Fractional SCA data proved adequate for the goals of this study, affording high  
754 temporal resolution observations and an appropriate spatial resolution given the extent  
755 of the study region. These results have implications for evaluating seasonal water  
756 supply forecasts, analyzing synoptic-scale drivers of snow accumulation, and validating  
757 precipitation estimates from regional climate models. In addition, the strong correlation  
758 between peak SWE and seasonal river flow indicates that our results could be useful for  
759 the evaluation of alternative water resource projects as part of development and climate  
760 change adaptation initiatives. Finally, the regional SWE and anomaly estimates  
761 illustrate the dramatic spatial and temporal variability of water resources in the

James McPhee 1/1/2016 11:37

Eliminado: a

763 extratropical Andes, and provide a striking visual assessment of the progression of the  
764 drought that has affected the region since 2009. These results should motivate further  
765 research looking into the climatic drivers of this spatially distributed phenomenon.

766 **Acknowledgements**

767 This research was conducted with support from CONICYT, under grants FONDECYT  
768 1121184, SER-03, FONDEF CA13I10277 and CHILE-USA2013. The authors wish to  
769 thank everybody involved in field data collection, included brothers Santiago and  
770 Gonzalo Montserrat, Mauricio Cartes, Alvaro Ayala, and many others. Gonzalo Cortés  
771 provided insightful comments to working drafts of this manuscript.

772 **Bibliography**

773 Aceituno, P., 1988. On the functioning of the Southern Oscillation in the South  
774 American sector. Part I: Surface climate. Mon. Weather Rev. 116, 505–524.

775 Ahl, D.E., Gower, S.T., Burrows, S.N., Shabanov, N.V., Myneni, R.B., Knyazikhin, Y.,  
776 2006. Monitoring spring canopy phenology of a deciduous broadleaf forest using  
777 MODIS. Remote Sens. Environ. 104, 88–95.

778 Aravena, J.-C., Luckman, B.H., 2009. Spatio-temporal rainfall patterns in southern  
779 South America. Int. J. Climatol. 29, 2106–2120.

780 Army Corps of Engineers, U., 1960. Engineering and design: runoff from snowmelt.  
781 [publisher not identified], [Washington.

782 Arsenault, K.R., Houser, P.R., De Lannoy, G.J.M., 2014. Evaluation of the MODIS  
783 snow cover fraction product. Hydrol. Process. 28, 980–998. doi:10.1002/hyp.9636

784 Ayala, A., McPhee, J., Vargas, X., 2014. Altitudinal gradients, midwinter melt, and

- 785 wind effects on snow accumulation in semiarid midlatitude Andes under La Niña  
786 conditions. *Water Resour. Res.* 50, 3589–3594. doi:10.1002/2013WR014960
- 787 Benali, A., Carvalho, A.C., Nunes, J.P., Carvalhais, N., Santos, A., 2012. Estimating air  
788 surface temperature in Portugal using MODIS LST data. *Remote Sens. Environ.* 124,  
789 108–121.
- 790 Brubaker, K., Rango, A., Kustas, W., 1996. Incorporating Radiation Inputs into the  
791 Snowmelt Runoff Model. *Hydrol. Process.* 10, 1329–1343. doi:10.1002/(SICI)1099-  
792 1085(199610)10:10<1329::AID-HYP464>3.0.CO;2-W
- 793 Castro, L.M., Gironás, J., Fernández, B., 2014. Spatial estimation of daily precipitation  
794 in regions with complex relief and scarce data using terrain orientation. *J. Hydrol.* 517,  
795 481–492.
- 796 Cline, D.W., Bales, R.C., Dozier, J., 1998. Estimating the spatial distribution of snow in  
797 mountain basins using remote sensing and energy balance modeling. *Water Resour.*  
798 *Res.* 34, 1275–1285.
- 799 Colombi, A., De Michele, C., Pepe, M., Rampini, A., 2007. Estimation of daily mean  
800 air temperature from MODIS LST in Alpine areas. *EARSeL EProceedings* 6, 38–46.
- 801 Corripio, J.G., Purves, R.S., 2005. Surface energy balance of high altitude glaciers in  
802 the central Andes: The effect of snow penitentes. *Clim. Hydrol. Mt. Areas* 15–27.
- 803 Cortés, G., Cornwell, E., McPhee, J.P., Margulis, S.A., 2014. Snow Cover  
804 Quantification in the Central Andes Derived from Multi-Sensor Data, in: AGU Fall  
805 Meeting Abstracts. p. 0410.
- 806 Cortés, G., Girotto, M., Margulis, S.A., 2014. Analysis of sub-pixel snow and ice extent

807 over the extratropical Andes using spectral unmixing of historical Landsat imagery.  
808 Remote Sens. Environ. 141, 64–78. doi:10.1016/j.rse.2013.10.023

809 Cortés, G., Vargas, X., McPhee, J., 2011. Climatic sensitivity of streamflow timing in  
810 the extratropical western Andes Cordillera. J. Hydrol. 405, 93–109.  
811 doi:10.1016/j.jhydrol.2011.05.013

812 DeWalle, D., Rango, A., 2008. Principles of snow hydrology. Cambridge University  
813 Press, New York.

814 Dietz, A.J., Kuenzer, C., Gessner, U., Dech, S., 2012. Remote sensing of snow—a  
815 review of available methods. Int. J. Remote Sens. 33, 4094–4134.

816 Erxleben, J., Elder, K., Davis, R., 2002. Comparison of spatial interpolation methods for  
817 estimating snow distribution in the Colorado Rocky Mountains. Hydrol. Process. 16,  
818 3627–3649.

819 Eva, H.D., Belward, A.S., De Miranda, E.E., Di Bella, C.M., Gond, V., Huber, O.,  
820 Jones, S., Sgrenzaroli, M., Fritz, S., 2004. A land cover map of South America. Glob.  
821 Change Biol. 10, 731–744.

822 Falvey, M., Garreaud, R., 2007. Wintertime precipitation episodes in central Chile:  
823 Associated meteorological conditions and orographic influences. J. Hydrometeorol. 8,  
824 171–193.

825 Falvey, M., Garreaud, R.D., 2009. Regional cooling in a warming world: Recent  
826 temperature trends in the southeast Pacific and along the west coast of subtropical South  
827 America (1979–2006). J. Geophys. Res. Atmospheres 114, D04102.  
828 doi:10.1029/2008JD010519

- 829 Favier, V., Falvey, M., Rabatel, A., Praderio, E., López, D., 2009. Interpreting  
830 discrepancies between discharge and precipitation in high-altitude area of Chile's Norte  
831 Chico region (26–32°S). *Water Resour. Res.* 45, W02424. doi:10.1029/2008WR006802
- 832 Gafurov, A., Bárdossy, A., 2009. Cloud removal methodology from MODIS snow  
833 cover product. *Hydrol. Earth Syst. Sci.* 13, 1361–1373.
- 834 Garreaud, R.D., 2009. The Andes climate and weather. *Adv. Geosci.* 22, 3–11.
- 835 Gascoin, S., Lhermitte, S., Kinnard, C., Bortels, K., Liston, G.E., 2013. Wind effects on  
836 snow cover in Pascua-Lama, Dry Andes of Chile. *Adv. Water Resour.* 55, 25–39.  
837 doi:10.1016/j.advwatres.2012.11.013
- 838 Guan, B., Molotch, N.P., Waliser, D.E., Jepsen, S.M., Painter, T.H., Dozier, J., 2013.  
839 Snow water equivalent in the Sierra Nevada: Blending snow sensor observations with  
840 snowmelt model simulations. *Water Resour. Res.* 49, 5029–5046.  
841 doi:10.1002/wrcr.20387
- 842 Hall, D.K., Riggs, G.A., 2007. Accuracy assessment of the MODIS snow products.  
843 *Hydrol. Process.* 21, 1534–1547.
- 844 Hall, D.K., Riggs, G.A., Salomonson, V.V., DiGirolamo, N.E., Bayr, K.J., 2002.  
845 MODIS snow-cover products. *Remote Sens. Environ.* 83, 181–194.
- 846 Hansen, M.C., DeFries, R.S., Townshend, J.R.G., Carroll, M., Dimiceli, C., Sohlberg,  
847 R.A., 2003. Global percent tree cover at a spatial resolution of 500 meters: First results  
848 of the MODIS vegetation continuous fields algorithm. *Earth Interact.* 7, 1–15.
- 849 Hofierka, J., Suri, M., others, 2002. The solar radiation model for Open source GIS:  
850 implementation and applications, in: *Proceedings of the Open Source GIS-GRASS*

- 851 Users Conference. pp. 1–19.
- 852 Huete, A., Didan, K., Miura, T., Rodriguez, E.P., Gao, X., Ferreira, L.G., 2002.  
853 Overview of the radiometric and biophysical performance of the MODIS vegetation  
854 indices. *Remote Sens. Environ.* 83, 195–213.
- 855 Insel, N., Poulsen, C.J., Ehlers, T.A., 2010. Influence of the Andes Mountains on South  
856 American moisture transport, convection, and precipitation. *Clim. Dyn.* 35, 1477–1492.
- 857 Jepsen, S.M., Molotch, N.P., Williams, M.W., Rittger, K.E., Sickman, J.O., 2012.  
858 Interannual variability of snowmelt in the Sierra Nevada and Rocky Mountains, United  
859 States: Examples from two alpine watersheds. *Water Resour. Res.* 48.
- 860 Krogh, S.A., Pomeroy, J.W., McPhee, J., 2015. Physically Based Mountain  
861 Hydrological Modeling Using Reanalysis Data in Patagonia. *J. Hydrometeorol.* 16,  
862 172–193. doi:10.1175/JHM-D-13-0178.1
- 863 Kustas, W.P., Rango, A., Uijlenhoet, R., 1994. A simple energy budget algorithm for  
864 the snowmelt runoff model. *Water Resour. Res.* 30, 1515–1527.
- 865 Martinec, J., 1989. Hour-to-hour snowmelt rates and lysimeter outflow during an entire  
866 ablation period. *Snow Cover Glacier Var.* 19–28.
- 867 Masiokas, M.H., Villalba, R., Luckman, B.H., Le Quesne, C., Aravena, J.C., 2006.  
868 Snowpack variations in the central Andes of Argentina and Chile, 1951–2005: Large-  
869 scale atmospheric influences and implications for water resources in the region. *J. Clim.*  
870 19, 6334–6352.
- 871 Meromy, L., Molotch, N.P., Link, T.E., Fassnacht, S.R., Rice, R., 2013. Subgrid  
872 variability of snow water equivalent at operational snow stations in the western USA.

- 873 Hydrol. Process. 27, 2383–2400.
- 874 Minder, J.R., Mote, P.W., Lundquist, J.D., 2010. Surface temperature lapse rates over  
875 complex terrain: Lessons from the Cascade Mountains. *J. Geophys. Res. Atmospheres*  
876 1984–2012 115.
- 877 Molotch, N.P., 2009. Reconstructing snow water equivalent in the Rio Grande  
878 headwaters using remotely sensed snow cover data and a spatially distributed snowmelt  
879 model. *Hydrol. Process.* 23, 1076–1089. doi:10.1002/hyp.7206
- 880 Molotch, N.P., Bales, R.C., 2006. Comparison of ground-based and airborne snow  
881 surface albedo parameterizations in an alpine watershed: Impact on snowpack mass  
882 balance. *Water Resour. Res.* 42.
- 883 Molotch, N.P., Margulis, S.A., 2008. Estimating the distribution of snow water  
884 equivalent using remotely sensed snow cover data and a spatially distributed snowmelt  
885 model: A multi-resolution, multi-sensor comparison. *Adv. Water Resour., Hydrologic*  
886 *Remote Sensing* 31, 1503–1514. doi:10.1016/j.advwatres.2008.07.017
- 887 Montgomery, D.R., Balco, G., Willett, S.D., 2001. Climate, tectonics, and the  
888 morphology of the Andes. *Geology* 29, 579–582.
- 889 Neteler, M., 2010. Estimating Daily Land Surface Temperatures in Mountainous  
890 Environments by Reconstructed MODIS LST Data. *Remote Sens.* 2, 333–351.  
891 doi:10.3390/rs1020333
- 892 Neteler, M., Bowman, M.H., Landa, M., Metz, M., 2012. GRASS GIS: A multi-purpose  
893 open source GIS. *Environ. Model. Softw.* 31, 124–130.
- 894 Newman, M., Compo, G.P., Alexander, M.A., 2003. ENSO-forced variability of the

- 895 Pacific decadal oscillation. *J. Clim.* 16, 3853–3857.
- 896 Núñez, J., Rivera, D., Oyarzún, R., Arumí, J.L., 2013. Influence of Pacific Ocean  
897 multidecadal variability on the distributional properties of hydrological variables in  
898 north-central Chile. *J. Hydrol.* 501, 227–240.
- 899 Pomeroy, J.W., Marks, D., Link, T., Ellis, C., Hardy, J., Rowlands, A., Granger, R.,  
900 2009. The impact of coniferous forest temperature on incoming longwave radiation to  
901 melting snow. *Hydrol. Process.* 23, 2513.
- 902 Pontailler, J.-Y., Hymus, G.J., Drake, B.G., 2003. Estimation of leaf area index using  
903 ground-based remote sensed NDVI measurements: validation and comparison with two  
904 indirect techniques. *Can. J. Remote Sens.* 29, 381–387.
- 905 Ragettli, S., Cortés, G., McPhee, J., Pellicciotti, F., 2014. An evaluation of approaches  
906 for modelling hydrological processes in high-elevation, glacierized Andean watersheds.  
907 *Hydrol. Process.* 28, 5674–5695. doi:10.1002/hyp.10055
- 908 Raleigh, M.S., Landry, C.C., Hayashi, M., Quinton, W.L., Lundquist, J.D., 2013.  
909 Approximating snow surface temperature from standard temperature and humidity data:  
910 New possibilities for snow model and remote sensing evaluation. *Water Resour. Res.*  
911 49, 8053–8069.
- 912 Rankinen, K., Karvonen, T., Butterfield, D., 2004. A simple model for predicting soil  
913 temperature in snow-covered and seasonally frozen soil: model description and testing.  
914 *Hydrol. Earth Syst. Sci. Discuss.* 8, 706–716.
- 915 Rhee, J., Im, J., 2014. Estimating high spatial resolution air temperature for regions with  
916 limited in situ data using MODIS products. *Remote Sens.* 6, 7360–7378.

- 917 Rice, R., Bales, R.C., 2010. Embedded-sensor network design for snow cover  
918 measurements around snow pillow and snow course sites in the Sierra Nevada of  
919 California. *Water Resour. Res.* 46.
- 920 Rittger, K., Painter, T.H., Dozier, J., 2013. Assessment of methods for mapping snow  
921 cover from MODIS. *Adv. Water Resour.* 51, 367–380.
- 922 Rubio-Álvarez, E., McPhee, J., 2010. Patterns of spatial and temporal variability in  
923 streamflow records in south central Chile in the period 1952–2003. *Water Resour. Res.*  
924 46, W05514. doi:10.1029/2009WR007982
- 925 Salazar, G., Raaijijk, C., 2014. Evaluation of clear-sky conditions in high altitude sites.  
926 *Renew. Energy* 64, 197–202.
- 927 Scheel, M.L.M., Rohrer, M., Huggel, C., Santos Villar, D., Silvestre, E., Huffman, G.J.,  
928 2011. Evaluation of TRMM Multi-satellite Precipitation Analysis (TMPA) performance  
929 in the Central Andes region and its dependency on spatial and temporal resolution.  
930 *Hydrol Earth Syst Sci* 15, 2649–2663. doi:10.5194/hess-15-2649-2011
- 931 Searcy, J.K., Hardison, C.H., 1960. Double-mass curves.
- 932 Sicart, J.E., Essery, R.L., Pomeroy, J.W., Hardy, J., Link, T., Marks, D., 2004. A  
933 sensitivity study of daytime net radiation during snowmelt to forest canopy and  
934 atmospheric conditions. *J. Hydrometeorol.* 5, 774–784.
- 935 Slater, A.G., Barrett, A.P., Clark, M.P., Lundquist, J.D., Raleigh, M.S., 2013.  
936 Uncertainty in seasonal snow reconstruction: Relative impacts of model forcing and  
937 image availability. *Adv. Water Resour., Snow–Atmosphere Interactions and*  
938 *Hydrological Consequences* 55, 165–177. doi:10.1016/j.advwatres.2012.07.006

939 Tarboton, D.G., Luce, C.H., 1996. Utah energy balance snow accumulation and melt  
940 model (UEB). Citeseer.

941 Vera, C., Silvestri, G., Liebmann, B., González, P., 2006. Climate change scenarios for  
942 seasonal precipitation in South America from IPCC-AR4 models. Geophys. Res. Lett.  
943 33.

944 Vicuña, S., Garreaud, R.D., McPhee, J., 2011. Climate change impacts on the  
945 hydrology of a snowmelt driven basin in semiarid Chile. Clim. Change 105, 469–488.  
946 doi:10.1007/s10584-010-9888-4

947 Wan, Z., Zhang, Y., Zhang, Q., Li, Z., 2002. Validation of the land-surface temperature  
948 products retrieved from Terra Moderate Resolution Imaging Spectroradiometer data.  
949 Remote Sens. Environ. 83, 163–180.

950 Wan, Z., Zhang, Y., Zhang, Q., Li, Z.-L., 2004. Quality assessment and validation of  
951 the MODIS global land surface temperature. Int. J. Remote Sens. 25, 261–274.

952 Williamson, S.N., Hik, D.S., Gamon, J.A., Kavanagh, J.L., Flowers, G.E., 2014.  
953 Estimating temperature fields from MODIS land surface temperature and air  
954 temperature observations in a sub-Arctic Alpine environment. Remote Sens. 6, 946–  
955 963.

956

957 List of Tables

- 958 Table 1. Study area subdivision, relevant characteristics and model parameters
- 959 Table 2. Snow pillow measurements available within the study domain
- 960 Table 3. Summary of snow depth and density intensive study campaigns
- 961 Table 4. Model validation statistics against intensive study area observations around  
962 snow pillows and at catchment scale
- 963 Table 5. Coefficient of determination R<sup>2</sup> between river melt season flows (SSRV),  
964 estimated and observed SWE (end-of-winter)
- 965 Table 6. Peak SWE 2001 - 2014 climatology for river basins within the study region.  
966 Basin-wide averages, SCA-wide averages and basin-wide water volumes shown
- 967

James McPhee 1/1/2016 11:37
<b>Eliminado:</b> Table 1. Study area subdivision, relevant characteristics and model parameters... [3]
James McPhee 1/1/2016 11:37
<b>Con formato:</b> Interlineado: sencillo
James McPhee 1/1/2016 11:37
<b>Con formato:</b> Inglés (americano)
James McPhee 1/1/2016 11:37
<b>Con formato:</b> Inglés (americano)
James McPhee 1/1/2016 11:37
<b>Con formato:</b> Inglés (americano)
James McPhee 1/1/2016 11:37
<b>Con formato:</b> Inglés (americano)
James McPhee 1/1/2016 11:37
<b>Eliminado:</b> ... [4]
James McPhee 1/1/2016 11:37
<b>Con formato:</b> español
James McPhee 1/1/2016 11:37
<b>Con formato:</b> Interlineado: sencillo

973

974

**Table 1.** Study area subdivision, relevant characteristics and model parameters

Cluster	Area x10 <sup>3</sup> [km <sup>2</sup> ]	Average elevation [m.a.s.l.]	Average cluster latitude [°]	Clear sky Index (Kc)	Avg. $\alpha_r$ [cm/°C/day]	$T_a$ [°C]	Forest Cover [%]
C1	26.5	3300	-29.4	0.78	0.02	18.3	2.0
C2	17.9	2760	-33.7	0.89	0.11	16.1	5.5
C3	9.20	1890	-36.4	0.83	0.18	12.2	13.8
C4	49.3	3520	-30.1	0.8	0.04	20.4	1.4
C5	18.5	2855	-33.4	0.83	0.15	15.6	3.0
C6	7.60	2807	-34.8	0.83	0.21	13.9	2.3
C7	14.8	2167	-36.1	0.85	0.20	16.7	2.5
C8	8.30	1840	-37.0	0.82	0.23	15.7	4.9
Total / Average	152.1	2320	***	0.83	0.14	***	3.3

975

976

James McPhee 1/1/2016 11:37

**Con formato:** Fuente: 12 pt, Sin Negrita

James McPhee 1/1/2016 11:37

**Con formato:** Interlineado: sencillo

977 **Table 2. Snow pillow measurements available within the study domain**

978

ID	SWE data	Symbol	Lat. (S)	Long. (W)	Elevation [m.a.s.l.]	Reference cluster
<b>CHILE</b>						
1	Quebrada Larga	QUE	30° 43'	70° 16'	3500	C1
2	Cerro Vega Negra	CVN	30° 54'	70° 30'	3600	C1
3	El Soldado	SOL	32° 00'	70° 19'	3290	C2
4	Portillo	POR	32° 50'	70° 06'	3000	C2
5	Laguna Negra	LAG	33° 39'	70° 06'	2780	C2
6	Lo Aguirre	LOA	35° 58'	70° 34'	2000	C3
7	Alto Mallines	ALT	37° 09'	70° 14'	1770	C3
<b>ARGENTINA</b>						
8	Toscas	TOS	33° 09'	69° 53'	3000	C5
9	Laguna Diamante	DIA	34° 11'	69° 41'	3300	C6
10	Laguna Atuel	ATU	34° 30'	70° 02'	3420	C6
11	Valle Hermoso	VAL	35° 08'	70° 12'	2250	C7
12	Paso Pehuenches	PEH	35° 08'	70° 23'	2545	C7

979

980

981 **Table 3. Summary of snow depth and density intensive study campaigns**

982

Year	ID (Figure 1)	Symbol	Field site	Date	Snow-pit density [kg/m <sup>3</sup> ]	SWE average [mm]	SWE std. dev. [mm]	SWE range [mm]	Sample size
2010	2	ODA	Ojos de Agua	25-sep	352	450	163	848 - 0	134
2011	2	ODA	Ojos de Agua	30-agosto	341	705	199	1194 - 136	374
	5	MOR	Morales	01-sept	367	642	282	1101 - 0	171
	8	OBL	Olla Blanca DET	31-agosto	333	539	217	1032 - 79	289
2012	1	CVN	Cerro Vega Negra	28-agosto	308	296	115	700 - 40	166
	3	MAR	Juncal - Mardones	30-agosto	373	530	230	1120 - 40	163
	5	MOR	Morales	12-sept	412	590	360	1240 - 150	152
	8	OBL	Olla Blanca DET	03-sept	411	590	260	1230 - 0	309
	4	POR	Portillo	15-sept	410	170	180	1230 - 0	181
2013	1	CVN	Cerro Vega Negra	21-agosto	356	405	165	1040 - 10	282
	2	ODA	Ojos de Agua	23-agosto	355	540	220	1310 - 100	300
	10	CHI	Nevados Chillán <sup>a</sup>	27-agosto	416	980	240	1270 - 30	104
	10	CHI	Nevados Chillán <sup>b</sup>	27-agosto	416	600	240	1230 - 70	216
	4	POR	Portillo	23-agosto	392	340	210	1120 - 0	91
	6	LAG	Laguna Negra	30-agosto	455	480	250	1770 - 0	32
2014	1	CVN	Cerro Vega Negra	05-agosto	321	163	85	620 - 0	326
	5	MOR	Morales	12-agosto	401	510	250	1190 - 0	329
	7	LVD	Lo Valdez	13-agosto	365	710	290	1260 - 0	186
	8	OBL	Olla Blanca DET	12-sept	363	420	240	1210 - 0	334
	9	RBL	Río Blanco DET	06-sept	354	620	290	1210 - 0	99
	10	CHI	Nevados Chillán <sup>a</sup>	26-sept	504	830	400	380 - 1510	18
	10	CHI	Nevados Chillán <sup>b</sup>	26-sept	504	980	250	530 - 1500	87
	4	POR	Portillo	19-agosto	436	170	140	850 - 0	73
	6	LAG	Laguna Negra	30-agosto	365	300	110	540 - 0	117

(a) without forest cover (upper part of basin).

(b) with forest cover (lower part of basin).

983

984

985

**Table 4.** Model validation statistics against intensive study area observations around snow pillows and at catchment scale

Reconstructed SWE vs. MODIS pixel (grid) sampling (selected snow-pillows)							
	Avg. Sampling [mm] (1)	Std. Dev. Sampling [mm] (2)	Avg. Model [mm] (3)	SP (sensor) [mm] (4)	RE% (avg.)	RE% (avg.)	RE% (avg.)
CVN	223	110	334	200	49%	-10%	98%
POR	227	177	170	353	-25%	-35%	51%
LAG	395	180	283	280	-28%	-30%	8%
Reconstructed SWE vs. snow surveys (pilot-basins)							
	avg. Sampling [mm] (1)	std. dev. Sampling [mm] (2)	avg. Model [mm] (3)	std. dev. Model [mm]	RE% (1) vs. (3)	RE% (2) vs. (4)	RE% (std. dev.)
CVN	253	133	298	63 (4)	18%	-53%	
ODA-MAR	556	203	535	128	-4%	-37%	
MOR-LVD	613	295	375	115	-39%	-61%	
OBL-RBL	497	252	317	89	-36%	-65%	
CHI (forest)	790	245	257	46	-67%	-81%	
CHI (clear)	905	320	724	170	-20%	-47%	
Reconstructed SWE vs. snow-pillows (Sep 1 – Chile & Oct 1 – Argentina)							
	R <sup>2</sup>	Slope	Intercept	SE <sub>s</sub> [mm]	RE%	RMSE [mm]	Mod. SWE average [mm]
QUE	0.71	1.39	131	208	79	335	529
CVN	0.78	0.92	247	140	56	251	609
SOL	0.68	0.85	-16	112	19	127	401
POR	0.32	0.52	87	277	-36	398	437
LAG	0.42	0.76	16	217	-21	230	424
LOA	0.88	0.79	101	123	-5	171	734
ALT	0.83	0.56	5	89	-41	332	489
TOS	0.78	0.41	26	72	-52	251	120
DIA	0.76	0.85	38	141	-4	137	455
ATU	0.56	1.04	77	378	9	349	1263
VAL	0.72	0.74	11	211	-24	273	457
PEH	0.74	1.01	303	334	32	436	1302
Average	0.68			192	-2	274	602
							330

- Con formato ... [5] James McPhee 1/1/2016 11:37 Celdas insertadas ... [8] James McPhee 1/1/2016 11:37 Con formato ... [6] James McPhee 1/1/2016 11:37 Tabla con formato ... [7] James McPhee 1/1/2016 11:37 Con formato ... [9] James McPhee 1/1/2016 11:37 Con formato ... [10] James McPhee 1/1/2016 11:37 Con formato ... [11] James McPhee 1/1/2016 11:37 Con formato ... [12] James McPhee 1/1/2016 11:37 Con formato ... [13] James McPhee 1/1/2016 11:37 Con formato ... [14] James McPhee 1/1/2016 11:37 Con formato ... [15] James McPhee 1/1/2016 11:37 Con formato ... [16] James McPhee 1/1/2016 11:37 Con formato ... [17] James McPhee 1/1/2016 11:37 Con formato ... [18] James McPhee 1/1/2016 11:37 Con formato ... [19] James McPhee 1/1/2016 11:37 Con formato ... [20] James McPhee 1/1/2016 11:37 Con formato ... [21] James McPhee 1/1/2016 11:37 Con formato ... [22] James McPhee 1/1/2016 11:37 Con formato ... [23] James McPhee 1/1/2016 11:37 Con formato ... [24] James McPhee 1/1/2016 11:37 Con formato ... [25] James McPhee 1/1/2016 11:37 Con formato ... [26] James McPhee 1/1/2016 11:37 Con formato ... [27] James McPhee 1/1/2016 11:37 Con formato ... [28] James McPhee 1/1/2016 11:37 Con formato ... [29] James McPhee 1/1/2016 11:37 Con formato ... [30] James McPhee 1/1/2016 11:37 Con formato ... [31] James McPhee 1/1/2016 11:37 Con formato ... [32] James McPhee 1/1/2016 11:37 Con formato ... [33] James McPhee 1/1/2016 11:37 Con formato ... [34] James McPhee 1/1/2016 11:37 Con formato ... [35] James McPhee 1/1/2016 11:37 Con formato ... [36] James McPhee 1/1/2016 11:37 Con formato ... [37] James McPhee 1/1/2016 11:37 Con formato ... [38] James McPhee 1/1/2016 11:37 Con formato ... [39] James McPhee 1/1/2016 11:37 Con formato ... [40] James McPhee 1/1/2016 11:37 Con formato ... [41] James McPhee 1/1/2016 11:37 Con formato ... [42] James McPhee 1/1/2016 11:37 Con formato ... [43] James McPhee 1/1/2016 11:37 Con formato ... [44]

991      **Table 5. Coefficient of determination R<sup>2</sup> between river melt season flows (SSRV), estimated and**  
 992      **observed SWE (end-of-winter).**

993

R2 values specific SSRV vs. Estimated SWE per cluster		R2 values specific SSRV vs. SWE at snow pillows (2001 – 2013)*		
	2001 – 2014	Neglecting 2009 at Argentinean clusters**	Best	2 <sup>nd</sup> best
c1	0.84	***	0.74 (CVN)	0.69 (QUE)
c2	0.78	***	0.82 (LAG)	0.68 (POR)
c3	0.57	***	0.17 (LOA)	0.16 (ALT)
c4	0.87	***	***	-
c5	0.66	0.82	0.81 (TOS)	-
c6	0.45	0.76	0.87 (ATU)	0.77 (DIA)
c7	0.64	0.89	0.77 (VAL)	0.41 (PEH)
c8	0.48	0.64	***	-

\* 2014 flows in Argentina unavailable to us at the moment of writing.

\*\* 2009 is considered an outlier year for the reconstruction at Argentinean sites.

994

995

996

997

**Table 6.** Peak SWE 2001 - 2014 climatology for river basins within the study region. Basin-wide averages, SCA-wide averages and basin-wide water volumes shown

998

ID	Basin - gauge station	Lat. S	Long. W	Outlet Elev. [m. a.s.l.]	Area [km <sup>2</sup> ]	SWE		
						Basin-wide [mm]	Over- SCA [mm]	Basin-wide [m <sup>3</sup> x10 <sup>6</sup> ]
<b>CHILE</b>								
1	Copiapó en Pastillo	27° 59'	69° 58'	1300	7470	45	120	336
2	Huasco en Algodones	28° 43'	70° 30'	750	7180	68	161	488
3	Elqui en Algarrobal	29° 59'	70° 35'	760	5710	151	269	862
4	Hurtado en San Agustín	30° 27'	70° 32'	2050	676	302	325	204
5	Grande en Puntilla San Juan	30° 41'	70° 55'	2140	3545	137	306	486
6	Cogotí en La Fraguita	31° 06'	70° 53'	1021	491	182	335	89
7	Illapel en Huinitil	31° 33'	70° 57'	650	1046	180	305	188
8	Chalina en San Agustín	31° 41'	70° 43'	920	437	142	332	62
9	Choapa en Salamanca	31° 48'	70° 55'	560	2212	214	356	473
10	Sobrante en Piñadero	32° 12'	70° 42'	2057	126	172	198	22
11	Alicahue en Colliguay	32° 18'	70° 44'	852	344	92	184	32
12	Putaendo en Resg. Los Patos	32° 30'	70° 34'	1218	890	273	346	243
13	Aconcagua en Chacabuco	32° 51'	70° 30'	950	2110	609	692	1285
14	Mapocho en Los Almendros	33° 22'	70° 27'	970	640	269	342	172
15	Maipo en El Manzano	33° 35'	70° 22'	850	4840	692	760	3349
16	Cachapoal en Puente Termas	34° 15'	70° 34'	700	2455	700	814	1719
17	Tinguiririca en Los Briones	34° 43'	70° 49'	560	1785	532	677	950
18	Teno en Claro	34° 59'	70° 49'	650	1210	438	524	530
19	Lontué en Colorado - Palos	35° 15'	71° 02'	600	1330	656	759	872
20	Maule en Armerillo	35° 42'	70° 10'	470	5465	525	554	2869
21	Nuble en San Fabián	36° 34'	71° 33'	410	1660	376	430	624
22	Polcura en Laja	37° 19'	77° 32'	675	2088	358	378	748
<b>ARGENTINA</b>								
23	Jachal en Pachimoco	30° 12'	68° 49'	1563	24266	79	175	1917
24	San Juan en km 101	31° 15'	69° 10'	1129	23860	308	569	7349
25	Mendoza en Guido	32° 54'	69° 14'	1479	7304	460	672	3360
26	Tunuyán en Zapata	33° 46'	69° 16'	852	11230	289	592	3245
27	Diamante en La Jaula	34° 40'	69° 18'	1451	19332	395	489	7636
28	Atuel en Loma Negra	35° 15'	69° 14'	1353	3696	338	525	1249
39	Malargüe en La Barda	35° 33'	69° 40'	1568	1055	171	284	180
30	Colorado en Buta Ranquil	37° 04'	69° 44'	817	14896	288	495	4290
31	Neuquén en Rahueco	37° 21'	70° 27'	870	8266	356	446	2943

1000

1001

James McPhee 1/1/2016 11:37

**Con formato:** español

James McPhee 1/1/2016 11:37

**Eliminado:** 6.

James McPhee 1/1/2016 11:37

**Con formato:** español

James McPhee 1/1/2016 11:37

**Con formato:** Normal

1003 List of Figures

1004 Figure 1. Study area and model domain: (a) river basins, stream gages (red circles) and  
1005 sites where snow survey data are available (green circles), (b) hydrologic units (C1 to  
1006 C8) and snow-pillow stations (white circles).

1007 Figure 2. Summarized hydro-climatology of the model domain. Data from  
1008 meteorological stations located within zones C1, C4, C3 and C8 summarized the hidro-  
1009 climatological regime of northern-west, northern-east, southern-west and southern-east  
1010 zones respectively. Total SWE is SWE measured at selected snow pillow stations.

James McPhee 1/1/2016 11:37

Eliminado: climatology

1011 Figure 3. Reconstructed SWE validation at selected snow-pillow sites. Black diamonds  
1012 are instrumental records, gray circles are model estimates, and box-plots summarize  
1013 manual verification dataset around the pillow site. Upper and lower box limits are the  
1014 75% and 25% quartiles, horizontal line is the median, white box is the mean, upper and  
1015 lower dashes represent plus and minus 2.5 standard deviations from the mean, and  
1016 crosses are outlying values.

1017 Figure 4. Reconstructed SWE validation at pixels with snow survey data. Box plots  
1018 summarize all individual measurements at pixels co-located with SWE reconstruction.  
1019 Symbology analogous to Figure 3.

1020 Figure 5. Comparison between peak reconstructed and observed SWE at snow-pillow  
1021 sites. Solid line represents the 1:1 line.

1022 Figure 6. Area-specific spring - summer runoff volume (SSRV) versus peak SWE.  
1023 Clusters 1 through 3 include rivers on the Chilean (western) slope of the Andes range;  
1024 clusters 4 through 8 correspond to Argentinean (eastern) rivers. Solid line represents 1:1  
1025 line. C4 and C8 SSRV were estimated by area-transpose method.

1026 Figure 7. Regional peak (Sep 15th) SWE Climatology for the 2001 - 2014 period (upper  
1027 left panel), and annual peak SWE anomalies.

1028 Figure 8. Maximum SWE through 1000 m elevation bands (EB). Crosses are mean  
1029 values within EB, lines are estimated SWE-elevation profile. Circle radius indicate EB  
1030 area [km<sup>2</sup>] scaled by 0.05 and takes values from SWE axis.

1031 Figure 9. Time series of energy fluxes over snow surface (average over 14 years) and  
1032 global average per cluster. Unique axes scale for all plots.

1033 Figure 10. Average seasonal evolution of fSCA and SWE in the study region. Lower  
1034 right panel shows the spatial correlation between time-averaged fSCA, SWE and  
1035 Specific melt-season river discharge.

1036 Figure 11. Sensitivity of peak SWE estimates to model forcings and parameters.  
1037 Average over the 2001 - 2014 period at selected snow pillow sites.  $\Delta_x$  represents the  
1038 percentage change over each parameter studied respect to the base case.

1040 Figure 12. Restricted Degree Day factor as a function of space (basin cluster) and  
1041 climatological properties. Bowen ( $\beta$ ) coefficient shown between parenthesis in legend.

1042

Página 16: [1] Eliminado James McPhee 1/1/16 11:37

Table 1

Página 24: [2] Eliminado James McPhee 1/1/16 11:37

Figure 10

Página 40: [3] Eliminado James McPhee 1/1/16 11:37

Table 1. Study area subdivision, relevant characteristics and model parameters

Table 2. Snow pillow measurements available within the study domain

Table 3. Summary of snow depth and density intensive study campaigns

Página 40: [4] Eliminado James McPhee 1/1/16 11:37

Table 6. Peak SWE 2001 - 2014 climatology for river basins within the study region. Basin-wide averages, SCA-wide averages and basin-wide water volumes shown

Página 44: [5] Con formato James McPhee 1/1/16 11:37

Fuente: 10 pt

Página 44: [6] Con formato James McPhee 1/1/16 11:37

Fuente: 8 pt

Página 44: [7] Tabla con formato James McPhee 1/1/16 11:37

Tabla con formato

Página 44: [8] Celdas insertadas James McPhee 1/1/16 11:37

Celdas insertadas

Página 44: [9] Con formato James McPhee 1/1/16 11:37

Fuente: 8 pt

Página 44: [10] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [11] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [12] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [13] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [14] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [15] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [16] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [17] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [18] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [19] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

**Página 44: [20] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [21] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [22] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [23] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [24] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [25] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [26] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [27] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [28] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [29] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [30] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [31] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [32] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [33] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [34] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [35] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [36] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [37] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [38] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [39] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [40] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

Página 44: [41] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [42] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [43] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [44] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [45] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [46] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [47] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [48] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [49] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [50] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

**Página 44: [51] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [52] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [53] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [54] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [55] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [56] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [57] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [58] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [59] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [60] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [61] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [62] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [63] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [64] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [65] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [66] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [67] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [68] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [69] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [70] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [71] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

Página 44: [72] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [73] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [74] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [75] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [76] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [77] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [78] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [79] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [80] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [81] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

**Página 44: [82] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [83] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [84] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [85] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [86] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [87] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [88] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [89] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [90] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [91] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [92] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [93] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [94] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [95] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [96] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [97] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [98] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [99] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [100] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [101] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [102] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

Página 44: [103] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [104] Celdas insertadas

James McPhee

1/1/16 11:37

Celdas insertadas

Página 44: [105] Celdas insertadas

James McPhee

1/1/16 11:37

Celdas insertadas

Página 44: [106] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [107] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [108] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [109] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [110] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [111] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [112] Celdas eliminadas

James McPhee

1/1/16 11:37

Celdas eliminadas

**Página 44: [113] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [114] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [115] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [116] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [117] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [118] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [119] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [120] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [121] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [122] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [123] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [124] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [125] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [126] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [127] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [128] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [129] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [130] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [131] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [132] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [133] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

Página 44: [134] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [135] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [136] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [137] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [138] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [139] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [140] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [141] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [142] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [143] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

**Página 44: [144] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [145] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [146] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [147] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [148] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [149] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [150] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [151] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [152] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [153] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [154] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [155] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [156] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [157] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [158] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [159] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [160] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [161] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [162] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [163] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [164] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

Página 44: [165] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [166] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [167] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [168] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [169] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [170] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [171] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [172] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [173] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

Página 44: [174] Con formato

James McPhee

1/1/16 11:37

Fuente: 8 pt

**Página 44: [175] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [176] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [177] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [178] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [179] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [180] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [181] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [182] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [183] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [184] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [185] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [186] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [187] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [188] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [189] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [190] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [191] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [192] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [193] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [194] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [195] Con formato** James McPhee **1/1/16 11:37**

Fuente: 8 pt

**Página 44: [196] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [197] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [198] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [199] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [200] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [201] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [202] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt

**Página 44: [203] Con formato**

**James McPhee**

**1/1/16 11:37**

Fuente: 8 pt